Thermal cycling-hyperthermia in combination with polyphenols, epigallocatechin gallate and chlorogenic acid, exerts synergistic anticancer effect against human pancreatic cancer PANC-1 cells

Chueh-Hsuan Lu[1,2] , Wei-Ting Chen[1,2] , Chih-Hsiung Hsieh[1,2], Yu-Yi Kuo[1,2], and Chih-Yu Chao[1,2,3,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Biomedical & Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Institute of Applied Physics, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author
E-mail: cychao@phys.ntu.edu.tw (CYC)

Keywords: combination treatment, polyphenolic compound, pancreatic cancer, hyperthermia, synergistic effect

32 **Abstract**

Hyperthermia (HT) has shown feasibility and potency as an anticancer therapy.

Administration of HT in the chemotherapy has previously enhanced the cytotoxicity

of drugs against pancreatic cancer. However, the drugs used when conducting these

studies are substantially conventional chemotherapeutic agents that may cause

unwanted side effects. Additionally, the thermal dosage in the treatment of cancer

cells could also probably harm the healthy cells. The purpose of this work was to

investigate the potential of the two natural polyphenolic compounds, epigallocatechin

gallate (EGCG) and chlorogenic acid (CGA), as heat synergizers in the thermal

treatment of the PANC-1 cells. Furthermore, we have introduced a unique strategy

entitled the thermal cycling-hyperthermia (TC-HT) that is capable of providing a

maximum synergy and minimal side effect with the anticancer compounds. Our

results demonstrate that the combination of the TC-HT and the CGA or EGCG

markedly exerts the anticancer effect against the PANC-1 cells, while none of the

single treatment induced such changes. The synergistic activity was attributed to the

cell cycle arrest at the G2/M phase and the induction of the ROS-dependent

mitochondria-mediated apoptosis. These findings not only represent the first *in vitro*

thermal synergistic study of natural compounds in the treatment of pancreatic cancer,

but also highlight the potential of the TC-HT as an alternative strategy in thermal

treatment.

Introduction

Pancreatic cancer is one of the leading causes in cancer death and remains one of

the deadliest solid human malignancies worldwide [1]. Patients with pancreatic cancer

are commonly diagnosed at the unresectable stage, and in most cases, patients with

advanced pancreatic cancer have a poor response to chemotherapy or radiotherapy. In

spite of the fact that therapeutic methods have been improved, the prognosis for

pancreatic cancer patients still remains poor with a low five-year survival rate [2].

Therefore, there is a need for continued research in novel agents or alternative

therapeutic strategies for treating pancreatic cancers, thereby making an improvement

for the patients' quality of life.

Hyperthermia (HT) has emerged as a promising method for treating cancer over

the past decades [3]. It is a procedure exposing the tumor tissue to high temperatures

that cause cancer cell damage and death. Researches have shown that HT exhibits

therapeutic potential against cancer cells through multiple cellular changes, such as

protein denaturation and aggregation, inhibition of DNA synthesis, cytoskeleton

disruption, and alteration in the calcium homeostasis [4-6]. In addition, HT can

directly activate the immune response against the tumors, increase the tumor

oxygenation, and improve the drug delivery [7-9]. Although these encouraging results

have expanded our understanding of the cytotoxic effects of HT on the cancer cells, in

the case of HT as single treatment, it has been shown not to be sufficient to kill cancer

cells [10]. To strengthen the effectiveness of HT, several investigations have explored

combinations of HT and other cancer therapies, such as radiotherapy and

chemotherapy [11]. It has been demonstrated to be effective against various types of

cancer, including pancreatic cancer, in that HT enhanced the cytotoxicity of

gemcitabine through the inhibition of nuclear factor kappa B (NF-κB) [12-14]. There

have also been reports of gemcitabine and other drugs, such as cisplatin and

carbonplatin, combined with HT, that demonstrated the clinical efficacy in patients

with pancreatic cancer [15, 16]. These data suggest that HT could modify the

cytotoxicity of the anticancer drugs, thereby yielding better outcomes in treating

pancreatic cancer. However, the drugs used in these combined treatments are

conventional chemotherapeutic drugs, which have been known to cause unpleasant

and even dangerous side effects.

Nowadays, there has been an increasing interest in natural compounds research

due to their lower toxicity and diverse biological properties. Phenolic compounds are

among the most studied in cancer prevention and cure, and also the largest group of

phytochemicals, as well as being widely distributed in our diet. Particularly, regular

intakes of dietary polyphenols have been linked to lower risks of many cancers [17].

Tea and coffee are two of the most consumed beverages worldwide, and the natural

phenolic compounds, epigallocatechin gallate (EGCG) and chlorogenic acid (CGA),

are the major components in both drinks, respectively. It has been shown that EGCG

and CGA have healthy benefits such as antioxidative, anti-inflammatory, and

anticancer activities [18-20]. Recently, published findings from animal experimental

[21, 22] and clinical studies [23, 24] have demonstrated the ability of EGCG and

CGA to suppress tumor cell growth such as breast, lung and bladder cancer. On the

other hand, there are a number of studies indicating that natural compounds, including

polyphenols, can act as heat synergizers, and thereby improve the anticancer effect

[25-29]. Therefore, this has prompted us to conduct the first combined experiment of

HT and natural phenolic compounds in pancreatic cancer. Furthermore, we propose a

novel approach named thermal cycling-hyperthermia (TC-HT), which allows cells to

receive the equivalent thermal dosage through the repeated heat-and-cold cycle. As a

matter of fact, there is some evidence indicating that HT lacks tumor selectivity and

could also cause harm to normal cells [30, 31]. In this thermal cycled strategy, the

cells could avoid prolonged continuous heating under thermal treatment, thereby

reducing the toxicity of HT to highlight the synergistic anticancer efficiency of the

combined therapy. Therefore, the aim of this study was to investigate the synergistic

anti-pancreatic cancer effect of the TC-HT with the CGA and the TC-HT with the

EGCG.

In this paper, we examined the effects of the EGCG or CGA combined with the

TC-HT on the growth inhibition of PANC-1 cells and evaluated the cell cycle

regulation, apoptosis, and the expression of associated proteins to elucidate their

underlying mechanisms. Our results demonstrate that the co-administration with the

TC-HT and the EGCG or CGA significantly inhibited the cell proliferation and

increased the cell death by inducing the cell cycle arrest and mitochondrial apoptotic

pathway. These findings first indicate that both the EGCG and CGA might be

effective heat synergizers and that the maximum synergistic effects on the PANC-1

cells could be obtained when the TC-HT was administered. Furthermore, the TC-HT

as a thermal treatment could be much gentler and feasible, mainly because of the view

that the TC-HT itself is relatively harmless to the cells. We believe that this study

provides the interesting concept of cyclic thermal application in treatment of cancer *in*

vitro, and highlights the potential of the TC-HT as an alternative thermal treatment.

Materials and methods

Cell culture

Human pancreatic cancer cell line PANC-1 was obtained from the Bioresource

Collection and Research Center of the Food Industry Research and Development

Institute (Hsinchu, Taiwan). Cells were plated in 75 cm^3 cell culture flasks and grown

in high-glucose Dulbecco's modified Eagle's medium (DMEM) (Hyclone)

supplemented with 10% fetal bovine serum (FBS) (Hyclone) and 1%

penicillin-streptomycin (Gibco) in a humidified 5% CO_2 incubator at 37°C.

Drug treatment and TC-HT exposure

The EGCG was dissolved in distilled water at a concentration of 20 mM, and the

CGA was dissolved in dimethyl sulfoxide (DMSO)(Sigma). Subsequently, the two

samples were stored at -20°C. The stocks were diluted with a culture medium to the

indicated concentration for treatment before usage, and the final concentration of

DMSO in each well was 0.05% (v/v). After overnight incubation at 37^{o}C, cells were

treated with various concentrations of EGCG or CGA, as well as the solvent vehicle

(0.05% DMSO) for CGA. For the TC-HT, the cells were subject to a high and low

temperature water bath using a temperature controller modified by a PCR machine

(**Fig 1A**). The cells were exposed to HT (1-cycle TC-HT) at 46°C for 30 min without

a break or treated with 46-30°C TC-HT through 3, 6, and 10 cycles of 10, 5, and 3

min, respectively, to receive the equivalent thermal dosages (**Fig 1B**). In the

combined treatment, cells in a medium containing the EGCG or CGA were exposed

to the TC-HT. The temperatures of the cancer cells actually sensed were monitored by

placing a needle thermocouple at the bottom of the well (**Fig 1C**). During heat

exposure experiment, the non-heating groups (non-treated and drug-treated groups)

were exposed to a similar ambient environment as the heating groups over the

experimental period of time. After the treatment was completed, cells were stored in a

cell culture incubator until the time of analysis.

Fig 1. Exposure to the TC-HT via a temperature controller. (A)(B) Schematic of

experiment setup and the TC-HT programs setting. (C) The actual temperature

recorded every 20 sec in the PANC-1 cells throughout the exposure period.

MTT assay

Cell viability was accessed by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium

bromide (MTT) (Sigma) assay. Cells were seeded in 24-well plates and incubated

overnight at 37°C. After a single-agent treatment or combined treatment of the

TC-HT with the 0.05% DMSO (vehicle control), EGCG or CGA for 24 h, the

medium was removed and the cells were washed with phosphate buffered saline

(PBS). Cells were then incubated in DMEM containing 0.5 mg/ml MTT for an

additional 4 h at 37°C. Then the medium was removed and DMSO was added to

dissolve the formazan crystals. The supernatant from each sample was transferred into

96-well plates, and the absorbance was read at 570 nm using an ELISA microplate

reader. The calculation of synergism quotient (SQ) was dividing the combined effect

by the sum of individual effects. The given treatment shows a synergy when SQ is

greater than 1.0.

Clonogenic survival assay

PANC-1 cells were seeded at 1000 cells/dish in 35 mm Petri dishes for 24 h and

treated with 0.05% DMSO (vehicle control), CGA, EGCG and TC-HT alone or in

combination. Cell medium was replaced after the treatment, and the dishes were

cultured in a humidified 5% CO_2 incubator at 37°C for additional 14 days. At last, the

cells were fixed with 4% paraformaldehyde (PFA) (Sigma) for 10 min and stained

with 0.1% crystal violet (Sigma). The colonies containing more than 50 cells were

counted, and the number of colonies in each treatment group was normalized to

control group.

Cell cycle analysis

Cells were seeded in 35 mm Petri dishes and incubated overnight at 37°C. After 24 h

treatment with DMSO (vehicle control), CGA, EGCG and TC-HT alone or in

combination, the cells were harvested, washed with PBS, and fixed with 70% ethanol

at 4°C for 30 min. The cells were then stained with propidium iodide (PI) (BD

Bioscience) and RNase A (Thermal Scientific) for 30 min in the dark. The stained

cells were subject to the cell cycle analysis by using a flow cytometer (FACSCanto II;

BD Biosciences), and data were analyzed with ModFit LT software.

186   DAPI staining assay

DAPI staining was used to detect morphological characteristics of the nucleus. Cells

were cultured on glass coverslips in 35 mm Petri dishes. At the end of each 24 h

treatment, cells were washed twice with PBS and fixed with 4% PFA for 10 min at

room temperature. After washing twice with PBS, the glass coverslips with attached

cells were mounted using Fluoroshield mounting medium with DAPI (Abcam) and

examined with a fluorescence microscope (Axio Imager A1; ZEISS).

193   Annexin-V/PI double staining assay

Apoptosis was determined by using the Annexin V-FITC/PI apoptosis detection kit

(BD Biosciences). Briefly, PANC-1 cells were treated with the TC-HT combined with

the DMSO (vehicle control), CGA or EGCG, and then the cells were harvested with

trypsin-EDTA (Gibco) and collected by centrifugation at 2,000 × g for 5 min, washed

twice with cold PBS, and resuspended in binding buffer containing Annexin V-FITC

and PI. The cell suspensions were incubated for 15 min at room temperature in the

dark and analyzed by a FACS Calibur flow cytometer.

Mitochondria membrane potential (MMP) measurement

202 The cells treated with 0.05% DMSO (vehicle control), CGA, EGCG and TC-HT

203 alone or in combination for 24h were harvested, and resuspended with PBS followed

204 by staining with 20 nM $DiOC_6(3)$ (Enzo Life Sciences International Inc.) for 30 min

205 at 37°C in the dark. The fraction of cells showing low MMP was then measured by a

206 flow cytometer.

ROS detection

Cellular Reactive Oxygen Species (ROS) levels of superoxide anion ($O_2^{\bullet-}$) were

detected using the fluorescent dye dihydroethidium (DHE) (Sigma). Cells were

treated with the indicated treatments, washed with PBS, and then incubated with 5

μM DHE for 30 min at 37°C in the dark. The fluorescence intensities were measured

by flow cytometry and ROS levels were expressed as mean fluorescence intensity

(MFI).

Western blot analysis

After treatment with CGA, EGCG, and TC-HT for 24 h alone or in combination, cells

were harvested, washed with cold PBS, and lysed on ice for 30 min in lysis buffer (50

mM Tris-HCl, pH 7.4, 150 mM NaCl, 0.25% deoxycholic acid, 1% NP-40, 1.0%

Triton X-100, 0.1% SDS, 1 mM EDTA, 1% phosphate and protease inhibitor cocktail)

(Millipore). Cell lysates were clarified by centrifugation at 23,000 × g for 30 min at

4°C, and the protein concentration in the supernatant fraction was quantified using the

Bradford protein assay (Bioshop). Proteins were resolved by 10% SDS-PAGE and

electrotransferred onto polyvinylidene fluoride membrane (PVDF) (Millipore) in

transfer buffer (10 mM CAPS, pH 11.0, 10% methanol). The membranes were

blocked with 5% nonfat dry milk/TBST (blocking buffer) for 1 h at room temperature

and then incubated overnight at 4°C with diluted primary antibodies in blocking

buffer. The specific primary antibodies against Bcl-2, cleaved caspase-8, cleaved

caspase-9, cleaved caspase-3 (Cell Signaling), Bax (Santa Cruz), Cdc2, cyclin B1,

cleaved PARP and β-actin (GeneTex) were used. After washing with TBST, the

membranes were incubated with HRP-conjugated anti-goat (GeneTex) or anti-rabbit

(Jackson Immunoresearch) secondary antibody. Chemiluminescence was detected

using WesternBright ECL western blotting reagent (Advansta).

Statistical analysis

The results were presented as mean ± standard deviation (SD). Statistical analysis

using one-way analysis of variance (ANOVA) performed with SigmaPlot software.

The results were considered to be statistically significant when p-values were less

than 0.05. Each experiment was done in triplicate.

Results

TC-HT in combination with polyphenols synergistically inhibits proliferation of PANC-1 cells

The TC-HT was performed by a modified PCR machine to raise the temperature to a

desired level followed by a rapid return to normothermic temperature. The actual

temperature in the cells measured by a needle thermocouple could be elevated from

$36^{o}C$ to 43.5, 44.2 and $44.9^{o}C$ within 3, 5 and 10 min, respectively, and rapidly

returned to the physiological temperature (**Fig 1C**). The effect of the TC-HT and the

CGA or EGCG on cell growth was first explored using MTT assay. As shown in **Figs**

2A and **2B**, cells were treated with various concentrations of the CGA or EGCG for

24 h, and both of the compounds only slightly affected the viability of PANC-1 cells.

DMSO (0.05%) treated cells with or without the TC-HT did not show a significant

difference in viability compared to untreated control cells. Moreover, the viability of

PANC-1 cells in response to treatment with the TC-HT decreased in a

cycle-dependent manner. When the TC-HT was combined with either of the two

compounds, a significant dose-dependent decrease in viability was observed at all

four cycling parameters. Although the 1-cycle TC-HT and either the compound

cooperatively reduced the proliferation of PANC-1 cells, the heat alone resulted in

significant cytotoxicity to the cells. Notably, there was no difference in the cell

viability under the condition of the 10-cycles TC-HT when compared to that in the

control group, while it still was capable of working synergistically with either the

CGA or the EGCG to exert the anti-proliferative activity. The SQ values for the

corresponding treatments are as shown in **Table 1**. This suggests that both the CGA

and the EGCG could exhibit a synergistic cytotoxic effect when co-administered with

the TC-HT, particularly in the treatment of 10 cycles. We next performed a

clonogenic survival assay to confirm the effect of the TC-HT combined with CGA or

EGCG on cell proliferation. As shown in **Figs 2C** and **2D**, colony formation in

PANC-1 cells was dramatically reduced following both combined treatments. Based

on these data, the 10-cycles TC-HT (43.5-36 °C) in combination with the

concentration of 200 μM CGA, or the concentration of 20 μM EGCG, was used for

all the following experiments.

Fig 2. Effects of CGA or EGCG combined with the TC-HT on proliferation in

PANC-1 cells. (A) The cells were treated with different cycle numbers of the

TC-HT, various concentrations of CGA and 0.05% DMSO (vehicle control) alone or

in combination for 24 h. (B) The cells were treated with different cycle numbers of

the TC-HT and various concentrations of EGCG alone or in combination for 24 h. (C)

Representative images of clonogenic survival assay. (D) Analysis of colony formation

rate. Data are presented as mean ± S.D. in triplicate.

Table 1. Synergy quotient for CGA or EGCG in combination with the TC-HT.

		TC-HT			
		1-cycle	**3-cycles**	**6-cycles**	**10-cycles**
CGA (μM)	**100**	0.84	0.86	1.08	1.51
	200	1.10	1.32	1.51	2.82

	300	1.11	1.34	1.38	2.40
	10	1.18	1.11	1.27	1.95
EGCG (μM)	**20**	1.28	1.35	1.76	3.09
	30	1.23	1.40	1.70	2.69

The combination of CGA or EGCG with TC-HT caused G2/M cell cycle arrest in PANC-1 cells

282  To evaluate the mode of the anti-proliferative effects of the TC-HT, CGA and

283 EGCG, the cell cycle progression in the PANC-1 cells was examined by flow

284 cytometric analysis. As shown in **Figs 3A** and **3B**, treatment with the CGA, EGCG,

285 and TC-HT had no obvious effect on the cell progression when compared with the

286 group of untreated cells. Also, 0.05% DMSO used as the solvent vehicle of CGA

287 either alone or in combination with TC-HT had no effect on the cell progression.

288 Interestingly, the TC-HT combined with either of the two polyphenols resulted in a

289 significant accumulation of cells in the G2/M phase with a concomitant reduction of

290 cells in the G1 phase. To investigate the molecular mechanism of the results of the

291 combined effect on the cell cycle distribution, we examined the relevant proteins

involved in the G2/M progression. As shown in **Fig 3C**, the expression of Cdc2 and

cyclin B1 were markedly reduced in response to the combined treatment with TC-HT

and CGA or the combined treatment with TC-HT and EGCG. These results indicate

that either the CGA or EGCG combined with the TC-HT synergistically induced cell

cycle arrest in the G2/M phase by decreasing cyclin B1 and Cdc2 in the cells, and

eventually inhibited the cell growth of PANC-1 cells.

Fig 3. Effects of CGA or EGCG combined with the TC-HT on the G2/M cell

cycle arrest in PANC-1 cells. Cells were treated with the TC-HT (10 cycles), 0.05%

DMSO (vehicle control), CGA (200 μM), and EGCG (20 μM) alone or in

combination (TC-HT + DMSO, TC-HT + CGA, TC-HT + EGCG) for 24 h and

stained with propidium iodide (PI) for cell cycle analysis. (A) Representative DNA

content profiles. (B) The percentage of cell population in each phase of the cell cycle.

(C) Western bolt analysis of the expression of cell cycle regulator proteins cyclin B1

and Cdc2. β-actin was used as an internal control. Data are presented as mean ± S.D.

in triplicate. ($^{**}p < 0.01$ and $^{***}p < 0.001$ compared with the untreated control)

TC-HT combined with CGA or EGCG induces apoptosis in PANC-1 cells

We then investigated as to whether the synergistic cytotoxic effect of the TC-HT

with the CGA or EGCG was associated with the induction of apoptosis, and the flow

cytometric analysis was performed using Annexin V-FITC/PI staining. As shown in

Figs 4A and **4B**, the fraction of apoptotic cells after the treatment of PANC-1 cells

with either the single agent CGA, or the EGCG or TC-HT were barely increased as

compared with untreated control cells. There was no significant difference in

apoptosis between the 0.05% DMSO (vehicle control) combined with or without

TC-HT and the untreated control. Interestingly, the TC-HT in combination with the

CGA significantly increased the early apoptotic cell death in PANC-1 cells

($18.6\pm5.2\%$). A similar result was also obtained in the combination of the TC-HT and

the EGCG, but to a greater extent ($24.3\pm0.8\%$). These results also showed that the late

apoptotic cell death was increased following the exposure of TC-HT with CGA

($15.7\pm4.6\%$) or EGCG ($18.4\pm4.6\%$). Furthermore, in contrast to control cells, the

DAPI staining demonstrated that the cells treated with either of the two combined

treatments had more condensed and fragmented nuclei, which are typical

morphological alterations for the apoptosis (**Fig 4C**). These results suggest that the

TC-HT has a potential synergistic effect with the CGA or EGCG on the apoptosis in

the PANC-1 cells.

Fig 4. Combination of the TC-HT and the CGA or EGCG induces apoptosis in

PANC-1 cells. The apoptosis analysis for the cells following the treatment with the

TC-HT (10 cycles), 0.05% DMSO (vehicle control), CGA (200 μM), and EGCG (20

μM) alone or in combination (TC-HT + DMSO, TC-HT + CGA, TC-HT + EGCG)

for 24 h. (A) Flow cytometric detection of the apoptosis with Annexin V-FITC/PI

double staining. (B) Histogram quantifying the percentage of PANC-1 cells in early

and late apoptosis. (C) The nuclei morphology alterations (arrow) were examined

using DAPI staining. Data are presented as mean ± S.D. in triplicate. ($^{**}p < 0.01$ and

$^{***}p < 0.001$ compared with the untreated control)

Combination of the TC-HT with CGA or EGCG triggers

significant loss in the mitochondrial membrane potential in

PANC-1 cells

Collapse of the mitochondrial integrity is a critical event in the cells undergoing

apoptosis. To examine whether the combination of the TC-HT and CGA- or

EGCG-induced apoptosis involved mitochondrial disruption, the mitochondrial

membrane potential (MMP) was assessed using $DiOC_6(3)$ fluorescence staining by

flow cytometric analysis [32]. As shown in **Figs 5A** and **5B**, the treatment with the

EGCG, the CGA and its corresponding vehicle control (0.05% DMSO) did not

change the MMP level in comparison with the untreated control. In response to

treatment with the TC-HT alone or together with vehicle control, there was no

significant change in MMP within cells; however, the cells treated with TC-HT

showed a stronger effect on the MMP depolarization after the administration of CGA

or EGCG (44±6.4% and 75.7±7.6%), which was consistent with the results of the

apoptosis analysis as described in **Fig 4**. These results indicate that the TC-HT

combined with the CGA- or EGCG-induced apoptosis in the PANC-1 cells is

mediated by mitochondrial dysfunction.

Fig 5. Combination of the TC-HT and the CGA or EGCG induces mitochondrial

dysfunction. (A) Flow cytometric analysis of MMP using $DiOC_6(3)$ after treatment

with the TC-HT (10 cycles), 0.05% DMSO (vehicle control), CGA (200 μM), and

EGCG (20 μM) alone or in combination (TC-HT + DMSO, TC-HT + CGA, TC-HT +

EGCG) for 24 h. The M1 regions indicate the cells with the loss of MMP. (B)

Histogram represents the percentage of cells in the M1 region. Data are presented as

mean ± S.D. in triplicate. ([**]$p < 0.01$ and [***]$p < 0.001$ compared with the untreated

control)

Combination of the TC-HT with CGA or EGCG induces apoptosis through the mitochondrial pathway

It is well known that Bcl-2 family proteins and caspases, along with PARP, play

important roles in the mitochondria-mediated apoptosis. To further explore the

mechanism by which the TC-HT combined with the CGA or EGCG triggered

apoptosis in the PANC-1 cells, we evaluated the expression of the apoptosis-related

proteins using western blot analysis. As shown in **Fig 6**, when compared with control

cells, the cleaved caspase-9, -3, and the cleaved PARP were markedly increased after

the co-administration with the TC-HT and either the CGA or EGCG. In addition, both

of the combined treatments down-regulated the expression of Bcl-2 and up-regulated

the expression of Bax, and thus decreased the ratio of Bcl-2 to Bax. These results

reveal that the TC-HT combined with the CGA or EGCG promotes apoptosis in the

PANC-1 cells via activation of the mitochondrial pathway.

Fig 6. Effects of the TC-HT, CGA and EGCG on the expression of the

apoptosis-related proteins in PANC-1 cells. (A)(B) The protein levels of cleaved

caspase-9, -3, cleaved PARP, Bcl-2 and Bax of the PANC-1 cells treated with the

TC-HT (10 cycles), CGA (200 μM) and EGCG (20 μM) alone or in combination

treatment (TC-HT + CGA, TC-HT + EGCG) for 24 h were examined by western blot

analysis. GAPDH and β-actin were used as internal controls.

The role of ROS in the TC-HT combined with the CGA- or

EGCG-induced apoptosis in PANC-1 cells

As ROS generation is a critical event in the induction of the apoptosis, we next

examined the role of ROS in apoptosis induced by treatment with the TC-HT in

combination with either the CGA or EGCG. The intracellular ROS was measured by

flow cytometry using a fluorescence probe, DHE, which reacts with $O_2^{\bullet-}$ [33]. As

shown in **Fig 7A**, the treatment of PANC-1 cells with 0.05% DMSO (vehicle control),

CGA or EGCG alone did not alter the level of ROS when compared with the

untreated control cells. Also, the level of ROS induced by TC-HT and its combination

with DMSO were found to be comparable to the untreated control in the MFI. It is

worth mentioning that the TC-HT combined with CGA and the TC-HT combined

with EGCG significantly increased the level of ROS in MFI by approximately 2.3-

and 3.6-fold, respectively (**Fig 7B**). These results suggest that the TC-HT combined

with the CGA or EGCG may induce apoptosis through the excessive ROS production

in the PANC-1 cells.

Fig 7. Combined effects of the TC-HT and the CGA or EGCG on ROS

generation in PANC-1 cells. (A) The DHE ($O_2^{\bullet-}$) levels were measured by flow

cytometry after treatment with the TC-HT (10 cycles), 0.05% DMSO (vehicle

control), CGA (200 μM), and EGCG (20 μM) alone or in combination (TC-HT +

DMSO, TC-HT + CGA, TC-HT + EGCG) for 24 h. (B) Graph shows the fold change

in MFI of DHE relative to control. Data are presented as mean ± S.D. in triplicate.

($^{**}p < 0.01$ and $^{***}p < 0.001$ compared with the untreated control)

Discussion

Despite previous studies having shown that HT or polyphenols, the EGCG and CGA, possessed anticancer activities against pancreatic cancer cells [34-37], the effect of these two agents as a combined therapy on pancreatic cancer has not been reported. In this study, we have explored as to whether the herbal compound, CGA or EGCG, could cooperate with heat against the pancreatic cancer cells, and elucidated the cellular mechanism underlying the biological effects. Our results have demonstrated that the exposure to heat using the TC-HT in combination with the CGA or EGCG synergistically inhibited the growth and induced the apoptosis in the PANC-1 cells. Previous studies have shown the benefits of HT as an adjuvant in combination with chemotherapy against many various cancers, including pancreatic cancer [13, 38, 39]. However, there is a major concern that HT can cause unavoidable thermal injury to normal cells. Therefore, it is important to refine the method of heat administration to achieve the desired thermal dosage with the minimal cytotoxicity. To the best of our knowledge, this is the first report indicating a novel approach TC-HT synergizes with the CGA or EGCG on pancreatic cancer PANC-1 cells.

Previous research has reported the response of PANC-1 cells to mild HT at

different temperatures [40]. Particularly, the temperature in a range of about 42 to

46°C has been shown capable of inducing the cell death of the pancreatic cell line

PANC-1. Our survival analysis revealed a consistent result that the cell growth of

PANC-1 in the 1-cycle group was significantly inhibited. Although HT has a

cooperative cytotoxicity with the CGA or EGCG, its severe thermal toxicity to a cell

is what we want to avoid. The usage of the repeated cycles of heat exposure could

provide a means to synergize with the anticancer compounds in the heating process,

and avoid the thermal damage accumulation in the following non-heating process.

Earlier studies have shown the time-dependent modification of cancer cells during

exposure to HT for which the analysis of cellular growth indicated that the survival

rate decreased with increasing HT exposure time [41, 42]. Namely, the short exposure

of cancer cells to HT may induce cellular stress without affecting their proliferation,

while the prolonged exposure may lead to cytotoxicity. It has also been reported that

short HT treatment of PANC-1 cells at the temperature 46°C for 5 min had no effect

on cell viability [43]. Therefore, we suggest that the 10-cycles TC-HT induced stress

rather than damage in cells during each heating cycle. As expected, the cells treated

with the TC-HT of different cycling parameters exhibited a decreased cytotoxicity

with the increased cycles. It is worth noting that 10 cycles of the TC-HT (43.5-36 °C),

or either compound, did not cause a significant alteration in the viability, however, the

combination of 10 cycles of the TC-HT with the CGA or EGCG strongly resulted in a

dose-dependent decline of the cell viability. Similar results from colony formation

were also obtained with the cells treated with the combination of TC-HT and CGA or

the combination of TC-HT and EGCG. These results suggested that the either CGA or

EGCG may act as heat synergizer and co-administered with the use of the TC-HT

could have significantly combined effects against pancreatic cancer PANC-1 cells.

Furthermore, we have also tested one liver cancer (HepG2) and one non-cancerous

(HEK293) cell lines to examine the combined action. Results shown in **S1 Fig**

revealed that the 10-cycles TC-HT did not cause cytotoxicity in both cancer and

non-cancerous cells. Interestingly, such substantial cell death in the PANC-1 cells

treated with the TC-HT in combination with CGA or EGCG was not observed in the

HepG2 and HEK293 cells, which indicated cell specificity of the combination

treatments.

HT has been demonstrated to inhibit the growth of human cancer cells through

interfering with the cell cycle progression [27, 29, 44]. In our results, we found that

the TC-HT did not cause an obvious accumulation of cells in the G2/M phase. When

cells undergoing the TC-HT were given the CGA or EGCG concurrently, the

proportion of cells in the G2/M phase was significantly increased. Additionally, this

observation was supported by the marked downregulation of Cdc2 and cyclin B1

proteins in the combination of the TC-HT and the CGA or EGCG. The Cdc2 (cell

division cycle 2), also known as the CDK1 (cyclin-dependent kinase 1), is a core

regulator that drives the cells through G2 and into mitosis. Several studies have

reported that the binding of Cdc2 to cyclin B1 complex plays an important role in the

G2/M progression [45, 46]. Collectively, these results indicated that the synergistic

cytotoxicity of CGA or EGCG under the exposure of TC-HT, at least in part, was

associated with the inhibition of the Cdc2/cyclin B1 kinase activity.

Apoptosis, also the best known form of programmed cell death, plays a pivotal

role in defending against cancer. The conventional HT has been demonstrated to

affect apoptotic pathways in various types of cancer cells [47]. It has also been

reported that polyphenols exhibit anticancer activities against different human cancer

cells through activating the apoptotic pathway [48, 49]. However, in our study, the

results of apoptosis analysis performed using FITC Annexin V and PI double staining

indicated that neither the TC-HT nor these two compounds induced significant

apoptotic cells (**Fig 4**). Only when the TC-HT in combination with the CGA or

EGCG was conducted, both of the combined treatments significantly elevated the

percentage of the early and late apoptotic cells. This finding was further confirmed by

the nuclear morphological alterations of the apoptosis in the TC-HT-treated cells in

combination with the CGA or EGCG. The apoptotic process can be divided into the

death receptor and mitochondrial pathways, and the mitochondrial pathway of cell

death is thought to be the major mechanism of apoptosis in mammals. The Bcl-2

family proteins are key regulators of the mitochondrial apoptotic pathway, which

comprise of both the anti-apoptotic and pro-apoptotic members. Unbalanced

Bcl-2/Bax ratio within the cells induces the disruption of the mitochondrial membrane,

release of cytochrome c, activation of caspases, and the subsequent cleavage of PARP

[50, 51]. Results from **Figs 5** and **6** indicated that the combined treatment with the

TC-HT and the CGA or EGCG markedly decreased the level of MMP and Bcl-2

expression, and increased Bax expression. We also found that the activation of

caspase-9, -3, and the cleavage of PARP were promoted with both of the combined

treatments **(Fig 6)**. Collectively, these data indicated that the mitochondria-dependent

pathway is involved in the synergistic apoptosis following the combined treatments.

Mitochondria are widely believed to be the main cellular source of ROS. The

excessive production of ROS could result in mitochondrial dysfunction, which in turn

triggers the apoptosis [52]. We then confirmed if the observation of mitochondrial

dysfunction in cells treated with the TC-HT in combination with the CGA or EGCG

was promoted by ROS generation. Our result showed that the cellular ROS generation

was elevated significantly following the administration of combination of the TC-HT

with the CGA or EGCG, indicating that the high levels of ROS production played an

important role in the apoptosis induced by the combination of the TC-HT with CGA

and the TC-HT with EGCG. HT has been known to act as an adjuvant treatment

modality to improve the cytotoxicity of several anticancer agents such as

chemotherapeutic drugs [53-55] and herbal compounds [56, 57]. It is thought that HT

can enhance the sensitivity of cancer cells to treatment with drugs [58], thereby

presenting synergistic anticancer activity. Therefore, these observations of this study

tempted us to suggest that the heat stress during temperature cycles may contribute to

the potentiation of the CGA or EGCG-triggered cytotoxic responses. Most

importantly, the TC-HT could serve as a mild approach to synergize with the natural

compound, CGA or EGCG. Furthermore, we also investigate the feasibility of the

TC-HT by using standard clinical compound gemcitabine shown to exhibit thermal

enhancement of cytotoxicity when combined with standard HT [53, 59]. Similar result

as the combination of 1-cycle TC-HT and gemcitabine was observed from analysis of

cell viability in the combined treatment with the 10-cycles TC-HT and gemcitabine,

whereas the 10-cycles TC-HT alone had no effect on cell viability (**S2 Fig**). This

finding suggests that this approach could be potentially extended to other anticancer

agent with heat synergy in cancer research.

In conclusion, a novel method for producing a desired thermal dosage is

proposed in which it applies repeated thermal treatments of short exposure to prevent

the toxic effects from prolonged exposure. Our findings have demonstrated the

TC-HT shows capability of synergizing with the natural compound, CGA or EGCG,

and minimizing the thermal damage resulting from HT. The synergistic activity

against the PANC-1 cells was performed primarily via the G2/M arrest and the

apoptosis induced through the ROS-mediated mitochondrial pathway leading to an

imbalance between Bcl-2 and Bax, activation of caspase-9 and -3, and the cleavage of

PARP. This study represents an interesting concept of cyclic thermal application in

treatment of cancer *in vitro*, which may potentially demonstrate an alternative to heat

treatment. Gaining an optimal combined effect for a variety of cancers via the

examination of the relation between the TC-HT parameters and other anticancer drugs

is an attractive subject that warrants further investigation.

Funding

This work was supported by grants from Ministry of Science and Technology

(MOST 105-2112-M-002-006-MY3 to CYC) and Ministry of Education (MOE

106R880708 to CYC) of the Republic of China. The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the

manuscript.

Competing Interests

The authors have declared that no competing interests exist.

Acknowledgments

We would like to thank Technology Commons in College of Life Science, National Taiwan University for use of flow cytometry system, and the staff of the imaging core at the First Core Labs, National Taiwan University Hospital for technical assistance.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Chueh-Hsuan Lu.

Formal analysis: Chueh-Hsuan Lu, Wei-Ting Chen, Chih-Hsiung Hsieh, Yu-Yi Kuo, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Chueh-Hsuan Lu, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Resources: Chih-Hsiung Hsieh, Chih-Yu Chao.

Supervision: Chih-Yu Chao.

563   **Validation:** Chueh-Hsuan Lu.

564   **Writing – original draft:** Chueh-Hsuan Lu, Wei-Ting Chen, Chih-Yu Chao.

565   **Writing – review & editing:** Chih-Yu Chao.

567 Submission

573 References

1.   Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. CA Cancer J Clin.
2016;66(1):7-30. doi: 10.3322/caac.21332. PubMed PMID: 26742998.
2.   Fryer RA, Galustian C, Dalgleish AG. Recent advances and developments in
treatment strategies against pancreatic cancer. Curr Clin Pharmacol. 2009;4(2):102-12.
PubMed PMID: 19442075.
3.   Overgaard J. The Current and Potential Role of Hyperthermia in Radiotherapy.
Int J Radiat Oncol. 1989;16(3):535-49.
4.   Roti JLR. Cellular responses to hyperthermia (40-46 degrees C): Cell killing and
molecular events. Int J Hyperther. 2008;24(1):3-15. doi:
10.1080/02656730701769841. PubMed PMID: WOS:000252581400002.
5.   Oei AL, Vriend LEM, Crezee J, Franken NAP, Krawczyk PM. Effects of
hyperthermia on DNA repair pathways: one treatment to inhibit them all. Radiat
Oncol. 2015;10. doi: ARTN 165
10.1186/s13014-015-0462-0. PubMed PMID: WOS:000358983800001.
6.   Huang SH, Yang KJ, Wu JC, Chang KJ, Wang SM. Effects of hyperthermia on
the cytoskeleton and focal adhesion proteins in a human thy carcinoma cell line. J

Cell Biochem. 1999;75(2):327-37. doi: Doi

10.1002/(Sici)1097-4644(19991101)75:2<327::Aid-Jcb14>3.0.Co;2-G. PubMed

PMID: WOS:000082783600014.

7. Calderwood SK, Theriault JR, Gong JL. How is the immune response affected

by hyperthermia and heat shock proteins? Int J Hyperther. 2005;21(8):713-6. doi:

10.1080/02656730500340794. PubMed PMID: WOS:000233906600007.

8. Ponce AM, Vujaskovic Z, Yuan F, Needham D, Dewhirst MW. Hyperthermia

mediated liposomal drug delivery. Int J Hyperther. 2006;22(3):205-13. doi:

10.1080/02656730600582956. PubMed PMID: WOS:000238093900004.

9. Song CW, Shakil A, Osborn JL, Iwata K. Tumour oxygenation is increased by

hyperthermia at mild temperatures. Int J Hyperther. 2009;25(2):91-5. doi: Pii

910057702

10.1080/02656730902744171. PubMed PMID: WOS:000264750400001.

10. Hildebrandt B, Wust P, Ahlers O, Dieing A, Sreenivasa G, Kerner T, et al. The

cellular and molecular basis of hyperthermia. Crit Rev Oncol Hematol.

2002;43(1):33-56. PubMed PMID: 12098606.

11. Kaur P, Hurwitz MD, Krishnan S, Asea A. Combined hyperthermia and

radiotherapy for the treatment of cancer. Cancers (Basel). 2011;3(4):3799-823. doi:

10.3390/cancers3043799. PubMed PMID: 24213112; PubMed Central PMCID:

PMCPMC3763397.

12. Au JLS, Wientjes MG. Combination Intravesical Hyperthermia and

Chemotherapy for Bladder Cancer The Rampersaud/Vujaskovic/Inman Article

Reviewed. Oncology-Ny. 2010;24(12):1155-60. PubMed PMID:

WOS:000293341400009.

13. Adachi S, Kokura S, Okayama T, Ishikawa T, Takagi T, Handa O, et al. Effect

of hyperthermia combined with gemcitabine on apoptotic cell death in cultured

human pancreatic cancer cell lines. Int J Hyperthermia. 2009;25(3):210-9.

14. Refaat T, Sachdev S, Sathiaseelan V, Helenowski I, Abdelmoneim S, Pierce MC,

et al. Hyperthermia and radiation therapy for locally advanced or recurrent breast

cancer. Breast. 2015;24(4):418-25. doi: 10.1016/j.breast.2015.03.008. PubMed PMID:

25900383; PubMed Central PMCID: PMCPMC4791303.

15. Bakshandeh-Bath A, Stoltz AS, Homann N, Wagner T, Stolting S, Peters SO.

Preclinical and Clinical Aspects of Carboplatin and Gemcitabine Combined with

Whole-body Hyperthermia for Pancreatic Adenocarcinoma. Anticancer Res.

2009;29(8):3069-77. PubMed PMID: WOS:000268846800029.

16.   Tschoep-Lechner KE, Milani V, Berger F, Dieterle N, Abdel-Rahman S, Salat C,

et al. Gemcitabine and cisplatin combined with regional hyperthermia as second-line

treatment in patients with gemcitabine-refractory advanced pancreatic cancer. Int J

Hyperther. 2013;29(1):8-16. doi: 10.3109/02656736.2012.740764. PubMed PMID:

WOS:000313672700002.

17.   Scalbert A, Williamson G. Dietary intake and bioavailability of polyphenols. J

Nutr. 2000;130(8):2073s-85s. PubMed PMID: WOS:000088760200003.

18.   Richelle M, Tavazzi I, Offord E. Comparison of the antioxidant activity of

commonly consumed polyphenolic beverages (coffee, cocoa, and tea) prepared per

cup serving. J Agr Food Chem. 2001;49(7):3438-42. doi: DOI 10.1021/jf0101410.

PubMed PMID: WOS:000170043400047.

19.   Min KJ, Kwon TK. Anticancer effects and molecular mechanisms of

epigallocatechin-3-gallate. Integr Med Res. 2014;3(1):16-24. doi:

10.1016/j.imr.2013.12.001. PubMed PMID: 28664074; PubMed Central PMCID:

PMCPMC5481703.

20.   Belkaid A, Currie JC, Desgagnes J, Annabi B. The chemopreventive properties

of chlorogenic acid reveal a potential new role for the microsomal

glucose-6-phosphate translocase in brain tumor progression. Cancer Cell Int. 2006;6:7.

doi: 10.1186/1475-2867-6-7. PubMed PMID: 16566826; PubMed Central PMCID:

PMCPMC1440869.

21.   Xu RS, Kang QM, Ren J, Li ZK, Xu XP. Antitumor Molecular Mechanism of

Chlorogenic Acid on Inducting Genes GSK-3 beta and APC and Inhibiting Gene

beta-Catenin (vol 2013, 951319, 2013). J Anal Methods Chem. 2015. doi: Artn

189328

10.1155/2015/189328. PubMed PMID: WOS:000352383000001.

22.   Feng RT, Lu YJ, Bowman LL, Qian Y, Castranova V, Ding M. Inhibition of

activator protein-1, NF-kappa B, and MAPKs and induction of phase 2 detoxifying

enzyme activity by chlorogenic acid. J Biol Chem. 2005;280(30):27888-95. doi:

10.1074/jbc.M503347200. PubMed PMID: WOS:000230678600051.

23.   Zhao H, Zhu W, Jia L, Sun X, Chen G, Zhao X, et al. Phase I study of topical

epigallocatechin-3-gallate (EGCG) in patients with breast cancer receiving adjuvant

radiotherapy. Br J Radiol. 2016;89(1058):20150665. Epub 2015/11/27. doi:

10.1259/bjr.20150665. PubMed PMID: 26607642; PubMed Central PMCID:

PMCPMC4985212.

24.   Gee JR, Saltzstein DR, Kim K, Kolesar J, Huang W, Havighurst TC, et al. A

Phase II Randomized, Double-blind, Presurgical Trial of Polyphenon E in Bladder

Cancer Patients to Evaluate Pharmacodynamics and Bladder Tissue Biomarkers.

Cancer Prev Res. 2017;10(5):298-307. doi: 10.1158/1940-6207.Capr-16-0167.

PubMed PMID: WOS:000400625500005.

25.   Kim JH, Kim SH, Alfieri AA, Young CW. Quercetin, an Inhibitor of Lactate

Transport and a Hyperthermic Sensitizer of Hela-Cells. Cancer Res. 1984;44(1):102-6.

PubMed PMID: WOS:A1984RX69400019.

26.   Cui ZG, Piao JL, Kondo T, Ogawa R, Tsuneyama K, Zhao QL, et al. Molecular

mechanisms of hyperthermia-induced apoptosis enhanced by docosahexaenoic acid:

Implication for cancer therapy. Chem-Biol Interact. 2014;215:46-53. doi:

10.1016/j.cbi.2014.03.005. PubMed PMID: WOS:000336357600007.

27.   Zhao PJ, Jiang H, Su D, Feng JG, Ma SL, Zhu XH. Inhibition of cell

proliferation by mild hyperthermia at 43 degrees C with Paris Saponin I in the lung

adenocarcinoma cell line PC-9. Mol Med Rep. 2015;11(1):327-32. doi:

10.3892/mmr.2014.2655. PubMed PMID: WOS:000346651100045.

28.   Cui ZG, Piao JL, Rehman MUR, Ogawa R, Li P, Zhao QL, et al. Molecular

mechanisms of hyperthermia-induced apoptosis enhanced by withaferin A. Eur J

Pharmacol. 2014;723:99-107. doi: 10.1016/j.ejphar.2013.11.031. PubMed PMID:

WOS:000330574600014.

29.   Hatashita M, Taniguchi M, Baba K, Koshiba K, Sato T, Jujo Y, et al. Sinodielide

A exerts thermosensitizing effects and induces apoptosis and G2/M cell cycle arrest in

DU145 human prostate cancer cells via the Ras/Raf/MAPK and PI3K/Akt signaling

pathways. Int J Mol Med. 2014;33(2):406-14. doi: 10.3892/ijmm.2013.1568. PubMed

PMID: 24285252.

30.   Dewey WC. Arrhenius Relationships from the Molecule and Cell to the Clinic.

Int J Hyperther. 1994;10(4):457-83. doi: Doi 10.3109/02656739409009351. PubMed

PMID: WOS:A1994NX92300001.

31.   Sminia P, Vanderzee J, Wondergem J, Haveman J. Effect of Hyperthermia on

the Central-Nervous-System - a Review. Int J Hyperther. 1994;10(1):1-30.

32.   Korchak HM, Rich AM, Wilkenfeld C, Rutherford LE, Weissmann G. A

carbocyanine dye, DiOC6(3), acts as a mitochondrial probe in human neutrophils.

Biochem Biophys Res Commun. 1982;108(4):1495-501. PubMed PMID: 7181903.

33.   Gomes A, Fernandes E, Lima JL. Fluorescence probes used for detection of

reactive oxygen species. Journal of biochemical and biophysical methods.

2005;65(2-3):45-80.

34.   Xiaoping L, Leizhen Z. Advances of high intensity focused ultrasound (HIFU)

for pancreatic cancer. Int J Hyperthermia. 2013;29(7):678-82. doi:

10.3109/02656736.2013.837199. PubMed PMID: 24102396.

35.   Wang K, Chen L, Meng Z, Lin J, Zhou Z, Wang P, et al. High intensity focused

ultrasound treatment for patients with advanced pancreatic cancer: a preliminary

dosimetric analysis. Int J Hyperthermia. 2012;28(7):645-52. doi:

10.3109/02656736.2012.713541. PubMed PMID: 22946544.

36.   Bimonte S, Cascella M, Leongito M, Palaia R, Caliendo D, Izzo F, et al. An

overview of pre-clinical studies on the effects of (-)-epigallocatechin-3-gallate, a

catechin found in green tea, in treatment of pancreatic cancer. Recenti Prog Med.

2017;108(6):282-7. Epub 2017/06/21. doi: 10.1701/2715.27715. PubMed PMID:

28631776.

37.   Dong J, Zou J, Yu XF. Coffee drinking and pancreatic cancer risk: a

meta-analysis of cohort studies. World J Gastroenterol. 2011;17(9):1204-10. Epub

2011/03/31. doi: 10.3748/wjg.v17.i9.1204. PubMed PMID: 21448427; PubMed

Central PMCID: PMCPMC3063915.

38.   Miyamoto R, Oda T, Hashimoto S, Kurokawa T, Inagaki Y, Shimomura O, et al.

Cetuximab delivery and antitumor effects are enhanced by mild hyperthermia in a

xenograft mouse model of pancreatic cancer. Cancer Sci. 2016;107(4):514-20.

39.   Issels RD. Hyperthermia adds to chemotherapy. Eur J Cancer.

2008;44(17):2546-54. doi: 10.1016/j.ejca.2008.07.038. PubMed PMID: 18789678.

40.   Ware MJ, Nguyen LP, Law JJ, Krzykawska-Serda M, Taylor KM, Cao HST, et

al. A new mild hyperthermia device to treat vascular involvement in cancer surgery.

Sci Rep. 2017;7(1):11299. doi: 10.1038/s41598-017-10508-6. PubMed PMID:

28900126; PubMed Central PMCID: PMCPMC5595878.

41.   Dewey WC, Hopwood LE, Sapareto SA, Gerweck LE. Cellular responses to

combinations of hyperthermia and radiation. Radiology. 1977;123(2):463-74. Epub

1977/05/01. doi: 10.1148/123.2.463. PubMed PMID: 322205.

42.   Raaphorst GP, Freeman ML, Dewey WC. Radiosensitivity and recovery from

radiation damage in cultured CHO cells exposed to hyperthermia at 42.5 or 45.5

degrees C. Radiat Res. 1979;79(2):390-402. PubMed PMID: 482603.

43.   Curley SA, Palalon F, Sanders KE, Koshkina NV. The Effects of Non-Invasive

Radiofrequency Treatment and Hyperthermia on Malignant and Nonmalignant Cells.

Int J Env Res Pub He. 2014;11(9):9142-53. doi: 10.3390/ijerph110909142. PubMed

PMID: WOS:000342027500031.

44. Lim CU, Zhang Y, Fox MH. Cell cycle dependent apoptosis and cell cycle

blocks induced by hyperthermia in HL-60 cells. Int J Hyperthermia. 2006;22(1):77-91.

doi: 10.1080/02656730500430538. PubMed PMID: 16423754.

45. Jessus C, Beach D. Oscillation of Mpf Is Accompanied by Periodic Association

between Cdc25 and Cdc2-Cyclin-B. Cell. 1992;68(2):323-32. doi: Doi

10.1016/0092-8674(92)90473-P. PubMed PMID: WOS:A1992HB35300011.

46. Fisher D, Krasinska L, Coudreuse D, Novak B. Phosphorylation network

dynamics in the control of cell cycle transitions. J Cell Sci. 2012;125(Pt 20):4703-11.

doi: 10.1242/jcs.106351. PubMed PMID: 23223895.

47. Ahmed K, Tabuchi Y, Kondo T. Hyperthermia: an effective strategy to induce

apoptosis in cancer cells. Apoptosis. 2015;20(11):1411-9. doi:

10.1007/s10495-015-1168-3. PubMed PMID: 26354715.

48. Rodríguez M, Estrela J, Ortega Á. Natural polyphenols and apoptosis induction

in cancer therapy. J Carcinog Mutag S. 2013;6.

49. Yamagata K, Izawa Y, Onodera D, Tagami M. Chlorogenic acid regulates

apoptosis and stem cell marker-related gene expression in A549 human lung cancer

cells. Mol Cell Biochem. 2018;441(1-2):9-19. doi: 10.1007/s11010-017-3171-1.

PubMed PMID: 28875417.

50. Tait SWG, Green DR. Mitochondrial Regulation of Cell Death. Csh Perspect

Biol. 2013;5(9). doi: ARTN a008706

10.1101/cshperspect.a008706. PubMed PMID: WOS:000325168700011.

51. Gross A, McDonnell JM, Korsmeyer SJ. BCL-2 family members and the

mitochondria in apoptosis. Genes Dev. 1999;13(15):1899-911. PubMed PMID:

10444588.

52. Orrenius S. Reactive oxygen species in mitochondria-mediated cell death. Drug

Metab Rev. 2007;39(2-3):443-55. doi: 10.1080/03602530701468516. PubMed PMID:

17786631.

53. Jin H, Zhao Y, Yang J, Zhang X, Ma S. Hyperthermia enhances the sensitivity of

pancreatic cancer SW1990 cells to gemcitabine through ROS/JNK signaling.

Oncology letters. 2018;16(5):6742-8.

54. Cesna V, Sukovas A, Jasukaitiene A, Naginiene R, Barauskas G, Dambrauskas Z,

et al. Narrow line between benefit and harm: Additivity of hyperthermia to cisplatin

cytotoxicity in different gastrointestinal cancer cells. World journal of

gastroenterology. 2018;24(10):1072.

55. Mohamed F, Marchettini P, Stuart OA, Urano M, Sugarbaker PH. Thermal

enhancement of new chemotherapeutic agents at moderate hyperthermia. Annals of

surgical oncology. 2003;10(4):463-8.

56. Zhao P, Jiang H, Su D, Feng J, Ma S, Zhu X. Inhibition of cell proliferation by

mild hyperthermia at 43 C with Paris Saponin I in the lung adenocarcinoma cell line

PC 9. Mol Med Rep. 2015;11(1):327-32.

57. Feng Q, Sun L, Teng B, Yanping X. Growth inhibitory effect of triptolide

combined with hyperthermia on Hep-2 cells. Journal of Jilin University (Medicine

Edition). 2006;(05).

58. Kampinga HH. Cell biological effects of hyperthermia alone or combined with

radiation or drugs: a short introduction to newcomers in the field. Int J Hyperther.

2006;22(3):191-6.

59. Adachi S, Kokura S, Okayama T, Ishikawa T, Takagi T, Handa O, et al. Effect

of hyperthermia combined with gemcitabine on apoptotic cell death in cultured

human pancreatic cancer cell lines. Int J Hyperther. 2009;25(3):210-9. doi: Pii

911094556

10.1080/02656730802657036. PubMed PMID: WOS:000265991500006.

783 Supporting information

S1 Fig. (A) HepG2 and (B) HEK293 cell viability results from MTT assay after

the treatments of CGA, EGCG, and TC-HT (10 cycles) alone or in combination

(TC-HT + CGA, TC-HT + EGCG) for 24 h.

S2 Fig. PANC-1 cell viability results from MTT assay after the treatments of

gemcitabine (5 μM) and TC-HT (1 cycle and 10 cycles) alone or in combination

for 24 h.

A

| TC-HT | − | − | + | + |
| CGA | − | + | − | + |

Cleaved caspase-9

Cleaved caspase-3

Cleaved PARP

GAPDH

Bcl-2

Bax

β-actin

B

| TC-HT | − | − | + | + |
| EGCG | − | + | − | + |

Cleaved caspase-9

Cleaved caspase-3

Cleaved PARP

GAPDH

Bcl-2

Bax

β-actin