Thermal cycling-hyperthermia in combination with polyphenols, epigallocatechin gallate and chlorogenic acid, exerts synergistic anticancer effect against human pancreatic cancer PANC-1 cells

Chueh-Hsuan Lu[1,2], Wei-Ting Chen[1,2], Chih-Hsiung Hsieh[1,2], Yu-Yi Kuo[1,2], and Chih-Yu Chao[1,2,3,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Biomedical & Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Institute of Applied Physics, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author
E-mail: cychao@phys.ntu.edu.tw (CYC)

Keywords: combination treatment, polyphenolic compound, pancreatic cancer, hyperthermia, synergistic effect

Abstract

Hyperthermia (HT) has shown feasibility and potency as an anticancer therapy. Administration of HT in the chemotherapy has previously enhanced the cytotoxicity of drugs against pancreatic cancer. However, the drugs used when conducting these studies are substantially conventional chemotherapeutic agents that may cause unwanted side effects. Additionally, the thermal dosage in the treatment of cancer cells could also probably harm the healthy cells. The purpose of this work was to investigate the potential of the two natural polyphenolic compounds, epigallocatechin gallate (EGCG) and chlorogenic acid (CGA), as heat synergizers in the thermal treatment of the PANC-1 cells. Furthermore, we have introduced a novel strategy entitled the thermal cycling-hyperthermia (TC-HT) that is capable of providing a maximum synergy and minimal side effect with the anticancer compounds. Our results demonstrate that the combination of the TC-HT and the CGA or EGCG markedly exerts the anticancer effect against the PANC-1 cells, while none of the single treatment induced such changes. The synergistic activity was attributed to the cell cycle arrest at the G2/M phase and the induction of the ROS-dependent mitochondria-mediated apoptosis. These findings not only represent the first thermal

49   synergistic study of natural compounds in the treatment of pancreatic cancer, but also

50   highlight the potential application of the TC-HT as an alternative strategy in anticancer

51   treatment.

Introduction

Pancreatic cancer is one of the leading causes in cancer death and remains one of

the deadliest solid human malignancies worldwide [1]. Patients with pancreatic cancer

are commonly diagnosed at the unresectable stage, and in most cases, patients with

advanced pancreatic cancer have a poor response to chemotherapy or radiotherapy. In

spite of the fact that therapeutic methods have been improved, the prognosis for

pancreatic cancer patients still remains poor with a low five-year survival rate [2].

Therefore, there is a need for continued research in novel agents or alternative

therapeutic strategies for treating pancreatic cancers, thereby making an improvement

for the patients' quality of life.

Hyperthermia (HT) has emerged as a promising method for treating cancer over

the past decades [3]. It is a procedure exposing the tumor tissue to high temperatures

that cause cancer cell damage and death. Researches have shown that HT exhibits

therapeutic potential against cancer cells through multiple cellular changes, such as

protein denaturation and aggregation, inhibition of DNA synthesis, cytoskeleton

disruption, and alteration in the calcium homeostasis [4-6]. In addition, HT can directly

activate the immune response against the tumors, increase the tumor oxygenation, and

improve the drug delivery [7-9]. Although these encouraging results have expanded our

understanding of the cytotoxic effects of HT on the cancer cells, the HT alone has a

limited role in cancer treatment. To strengthen the effectiveness of HT, several

investigations have explored combinations of HT and other cancer therapies, such as

radiotherapy and chemotherapy [10]. It has been demonstrated to be effective against

various types of cancer, including pancreatic cancer, in that HT enhanced the

cytotoxicity of gemcitabine through the inhibition of nuclear factor kappa B (NF-κB)

[11-13]. There have also been reports of gemcitabine and other drugs, such as cisplatin

and carbonplatin, combined with HT, that demonstrated the clinical efficacy in patients

with pancreatic cancer [14, 15]. These data suggest that HT could modify the

cytotoxicity of the anticancer drugs, thereby yielding better outcomes in treating

pancreatic cancer. However, the drugs used in these combined treatments are

conventional chemotherapeutic drugs, which have been known to cause unpleasant side

effects.

Nowadays, there has been an increasing interest in natural compounds research

due to their lower toxicity and diverse biological properties. Phenolic compounds are

among the most studied in cancer prevention and cure, and also the largest group of

phytochemicals, as well as being widely distributed in our diet. Particularly, regular

intakes of dietary polyphenols have been linked to lower risks of many cancers [16].

Tea and coffee are two of the most consumed beverages worldwide, and the natural

phenolic compounds, epigallocatechin gallate (EGCG) and chlorogenic acid (CGA),

are the major components in both drinks, respectively. It has been shown that EGCG

and CGA have healthy benefits such as antioxidative, anti-inflammatory, and

anticancer activities [17-19]. Recently, published findings from animal experimental

and clinical studies have demonstrated the ability of EGCG and CGA to suppress tumor

cell growth such as breast, lung and bladder cancer [20-23]. On the other hand, there are

a number of studies indicating that natural compounds, including polyphenols, can act

as heat synergizers, and thereby improve the anticancer effect [24-28]. Therefore, this

has prompted us to conduct the first combined experiment of HT and natural phenolic

compounds in pancreatic cancer. Furthermore, we propose a refined method for HT

that we have termed thermal cycling-hyperthermia (TC-HT), which allows cells to

receive the equivalent thermal dosage through the repeated heat-and-cold cycle. As a

matter of fact, there is some evidence indicating that HT lacks tumor selectivity and

could also cause harm to normal cells [29, 30]. In this thermal cycled strategy, the cells

could avoid prolonged continuous heating under thermal treatment, thereby reducing

the toxicity of HT to highlight the synergistic anticancer efficiency of the combined

therapy. Therefore, the aim of this study was to investigate the synergistic

anti-pancreatic cancer effect of the TC-HT with the CGA and the TC-HT with the

EGCG.

In this paper, we examined the effects of the EGCG or CGA combined with the

TC-HT on the growth inhibition of PANC-1 cells and evaluated the cell cycle

regulation, apoptosis, and the expression of associated proteins to elucidate their

underlying mechanisms. Our results demonstrate that the co-administration with the

TC-HT and the EGCG or CGA significantly inhibited the cell proliferation and

increased the cell death by inducing the cell cycle arrest and mitochondrial apoptotic

pathway. These findings first indicate that the EGCG and CGA might be effective heat

synergizers and that the maximum synergistic effects on the PANC-1 cells could be

obtained when the TC-HT was administered. Furthermore, the TC-HT as an alternative

strategy for the thermal therapy could be much safer and feasible, mainly because of the

view that the TC-HT itself is relatively harmless to the cells. We believe that the

TC-HT as an alternative thermal treatment may present a promising strategy for the

anti-pancreatic cancer combined therapy.

Materials and methods

Cell culture

Human pancreatic cancer cell line PANC-1 was obtained from the Bioresource

Collection and Research Center of the Food Industry Research and Development

Institute (Hsinchu, Taiwan). Cells were plated in 75 cm^3 cell culture flasks and grown

in high-glucose Dulbecco's modified Eagle's medium (DMEM) (Hyclone)

supplemented with 10% fetal bovine serum (FBS) (Hyclone) and 1%

penicillin-streptomycin (Gibco) in a humidified 5% CO_2 incubator at 37°C.

Drug treatment and TC-HT exposure

The EGCG was dissolved in distilled water at a concentration of 20 mM, and the CGA

was dissolved in dimethyl sulfoxide (DMSO)(Sigma). Subsequently, the two samples

were stored at -20°C. The stocks were diluted with a culture medium to the indicated

concentration for treatment before usage, and the final concentration of DMSO in each

well was 0.05% (v/v). After overnight incubation at 37^{o}C, cells were treated with

various concentrations of EGCG and CGA, as well as the solvent vehicle (0.05%

DMSO) for CGA. For the TC-HT, the cells were subject to a high and low temperature

water bath using a temperature controller modified by a PCR machine (**Fig 1A**). The

cells were exposed to HT (1-cycle TC-HT) at 46°C for 30 min without a break or

treated with 46-30°C TC-HT through 3, 6, and 10 cycles of 10, 5, and 3 min,

respectively, to receive the equivalent thermal dosages (**Fig 1B**). In the combined

treatment, cells in a medium containing the EGCG or CGA were exposed to the TC-HT.

The temperatures of the cancer cells actually sensed were monitored by placing a

needle thermocouple at the bottom of the well (**Fig 1C**). During heat exposure

experiment, the controls (non-treated and DMSO groups) and the drug-treated groups

were exposed to a similar hypoxic environment as the heating groups over the

experimental period of time. After the treatment was completed, cells were stored in a

cell culture incubator until the time of analysis.

Fig 1. Exposure to the TC-HT via a temperature controller. (A)(B) Schematic of

experiment setup and the TC-HT programs setting. (C) The actual temperature

recorded every 20 sec in the PANC-1 cells throughout the exposure period.

MTT assay

Cell viability was accessed by 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium

bromide (MTT) (Sigma) assay. Cells were seeded in 24-well plates and incubated

overnight at 37°C. After a single-agent treatment or combined treatment of the TC-HT

with the 0.05% DMSO (vehicle control), EGCG or CGA for 24 h, the medium was

removed and the cells were washed with phosphate buffered saline (PBS). Cells were

then incubated in DMEM containing 0.5 mg/ml MTT for an additional 4 h at 37°C.

Then the medium was removed and DMSO was added to dissolve the formazan

crystals. The supernatant from each sample was transferred into 96-well plates, and the

absorbance was read at 570 nm using an ELISA microplate reader. The calculation of

synergism quotient (SQ) was dividing the combined effect by the sum of individual

effects. The given treatment shows a synergy when SQ is greater than 1.0.

Clonogenic survival assay

PANC-1 cells were seeded at 1000 cells/dish in 35 mm Petri dishes for 24 h and treated

with 0.05% DMSO (vehicle control), CGA, EGCG and TC-HT alone or in combination.

Cell medium was replaced after the treatment, and the dishes were cultured in a

humidified 5% CO_2 incubator at 37°C for additional 14 days. At last, the cells were

fixed with 4% paraformaldehyde (PFA) (Sigma) for 10 min and stained with 0.1%

crystal violet (Sigma). The colonies containing more than 50 cells were counted, and

the number of colonies in each treatment group was normalized to control group.

Cell cycle analysis

Cells were seeded in 35 mm Petri dishes and incubated overnight at 37°C. After 24 h

treatment with DMSO (vehicle control), CGA, EGCG and TC-HT alone or in

combination, the cells were harvested, washed with PBS, and fixed with 70% ethanol at

4°C for 30 min. The cells were then stained with propidium iodide (PI) (BD Bioscience)

and RNase A (Thermal Scientific) for 30 min in the dark. The stained cells were subject

to the cell cycle analysis by using a flow cytometer (FACSCanto II; BD Biosciences),

and data were analyzed with ModFit LT software.

DAPI staining assay

DAPI staining was used to detect morphological characteristics of the nucleus. Cells

were cultured on glass coverslips in 35 mm Petri dishes. At the end of each 24 h

treatment, cells were washed twice with PBS and fixed with 4% PFA for 10 min at

room temperature. After washing twice with PBS, the glass coverslips with attached

cells were mounted using Fluoroshield mounting medium with DAPI (Abcam) and

examined with a fluorescence microscope (Axio Imager A1; ZEISS).

Annexin-V/PI double staining assay

Apoptosis was determined by using the Annexin V-FITC/PI apoptosis detection kit

(BD Biosciences). Briefly, PANC-1 cells were treated with the TC-HT combined with

the DMSO (vehicle control), CGA or EGCG, and then the cells were harvested with

trypsin-EDTA (Gibco) and collected by centrifugation at $2,000 \times g$ for 5 min, washed

twice with cold PBS, and resuspended in binding buffer containing Annexin V-FITC

and PI. The cell suspensions were incubated for 15 min at room temperature in the dark

and analyzed by a FACS Calibur flow cytometer.

Mitochondria membrane potential (MMP) measurement

The cells treated with 0.05% DMSO (vehicle control), CGA, EGCG and TC-HT for

24h alone or in combination were harvested, and resuspended with PBS followed by

staining with 20 nm $DiOC_6(3)$ (Enzo Life Sciences International Inc.) for 30 min at

37°C in the dark. The fraction of cells showing low MMP was then measured by a flow

cytometer.

ROS detection

Cellular Reactive Oxygen Species (ROS) levels of superoxide anion ($O_2^{\bullet-}$) were

detected using the fluorescent dye dihydroethidium (DHE) (Sigma). Cells were treated

with the indicated treatments, washed with PBS, and then incubated with 5 μM DHE

for 30 min at 37°C in the dark. The fluorescence intensities were measured by flow

cytometry and ROS levels were expressed as mean fluorescence intensity (MFI).

Western blot analysis

After treatment with CGA, EGCG, and TC-HT for 24 h alone or in combination, cells

were harvested, washed with cold PBS, and lysed on ice for 30 min in lysis buffer (50

mM Tris-HCl, pH 7.4, 150 mM NaCl, 0.25% deoxycholic acid, 1% NP-40, 1.0% Triton

X-100, 0.1% SDS, 1 mM EDTA, 1% phosphate and protease inhibitor cocktail)

(Millipore). Cell lysates were clarified by centrifugation at $23,000 \times g$ for 30 min at 4°C,

and the protein concentration in the supernatant fraction was quantified using the

Bradford protein assay (Bioshop). Proteins were resolved by 10% SDS-PAGE and

electrotransferred onto polyvinylidene fluoride membrane (PVDF) (Millipore) in

transfer buffer (10 mM CAPS, pH 11.0, 10% methanol). The membranes were blocked

with 5% nonfat dry milk/TBST (blocking buffer) for 1 h at room temperature and then

incubated overnight at 4°C with diluted primary antibodies in blocking buffer. The

specific primary antibodies against Bcl-2, cleaved caspase-8, cleaved caspase-9,

cleaved caspase-3 (Cell Signaling), Bax (Santa Cruz), Cdc2, cyclin B1, cleaved PARP

and β-actin (GeneTex) were used. After washing with TBST, the membranes were

incubated with HRP-conjugated anti-goat (GeneTex) or anti-rabbit (Jackson

Immunoresearch) secondary antibody. Chemiluminescence was detected using

WesternBright ECL western blotting reagent (Advansta).

Statistical analysis

The results were presented as mean ± standard deviation (SD). Statistical analysis using

one-way analysis of variance (ANOVA) performed with SigmaPlot software. The

results were considered to be statistically significant when *p*-values were less than 0.05.

Each experiment was done in triplicate.

Results

TC-HT in combination with polyphenols synergistically

inhibits proliferation of PANC-1 cells

The TC-HT was performed by a modified PCR machine to raise the temperature to a

desired level followed by a rapid return to normothermic temperature. The actual

temperature in the cells measured by a needle thermocouple could be elevated from

$36^{o}C$ to 43.5, 44.2 and $44.9^{o}C$ within 3, 5 and 10 min, respectively, and rapidly returned

to the physiological temperature (**Fig 1C**). The effect of the TC-HT and the CGA or

EGCG on cell growth was first explored using MTT assay. As shown in **Figs 2A** and

2B, cells were treated with various concentrations of the CGA or EGCG for 24 h, and

both of the compounds only slightly affected the viability of PANC-1 cells. DMSO

(0.05%) treated cells with or without the TC-HT did not show a significant difference

in viability compared to untreated control cells. Moreover, the viability of PANC-1

cells in response to treatment with the TC-HT decreased in a cycle-dependent manner.

When the TC-HT was combined with either of the two compounds, a significant

dose-dependent decrease in viability was observed at all four cycling parameters.

Although the 1-cycle TC-HT and either the compound cooperatively reduced the

proliferation of PANC-1 cells, the heat alone resulted in significant cytotoxicity to the

cells. Notably, there was no difference in the cell viability under the condition of the

10-cycles TC-HT when compared to that in the control group, while it still was capable

of working synergistically with either the CGA or the EGCG to exert the

anti-proliferative activity. The SQ values for the corresponding treatments are as shown

in **Table 1**. This suggests that the CGA and the EGCG could exhibit a synergistic

cytotoxic effect when co-administered with the TC-HT, particularly in the treatment of

10 cycles. We next performed a clonogenic survival assay to confirm the effect of the

TC-HT combined with each of CGA and EGCG on cell proliferation. As shown in **Figs**

2C and **2D**, colony formation in PANC-1 cells was dramatically reduced following

both combined treatments. Based on these data, the 10-cycles TC-HT (43.5-36 °C) in

combination with the concentration of 200 μM CGA, or the concentration of 20 μM

EGCG, was used for all the following experiments.

Fig 2. Effects of CGA or EGCG combined with the TC-HT on proliferation in

PANC-1 cells. (A) The cells were treated with different cycle numbers of the TC-HT,

various concentrations of CGA and 0.05% DMSO (vehicle control) alone or in

combination for 24 h. (B) The cells were treated with different cycle numbers of the

TC-HT and various concentrations of EGCG alone or in combination for 24 h. (C)

Representative images of clonogenic survival assay. (D) Analysis of colony formation

rate. Data are presented as mean ± S.D. in triplicate.

Table 1. Synergy quotient for polyphenols in combination with the TC-HT.

		TC-HT			
		1-cycle	**3-cycles**	**6-cycles**	**10-cycles**
	100	0.84	0.86	1.08	1.51
CGA (μM)	**200**	1.10	1.32	1.51	2.82
	300	1.11	1.34	1.38	2.40
	10	1.18	1.11	1.27	1.95
EGCG (μM)	**20**	1.28	1.35	1.76	3.09
	30	1.23	1.40	1.70	2.69

The combination of polyphenols and TC-HT caused G2/M cell cycle arrest in PANC-1 cells

To evaluate the mode of the anti-proliferative effects of the TC-HT, CGA and

EGCG, the cell cycle progression in the PANC-1 cells was examined by flow

cytometric analysis. As shown in **Figs 3A** and **3B**, treatment with the CGA, EGCG, and

TC-HT had no obvious effect on the cell progression when compared with the group of

untreated cells. Also, 0.05% DMSO used as the solvent vehicle of CGA either alone or

in combination with TC-HT had no effect on the cell progression. Interestingly, the

TC-HT combined with either of the two polyphenols resulted in a significant

accumulation of cells in the G2/M phase with a concomitant reduction of cells in the G1

phase. To investigate the molecular mechanism of the results of the combined effect on

the cell cycle distribution, we examined the relevant proteins involved in the G2/M

progression. As shown in **Fig 3C**, the expression of Cdc2 and cyclin B1 were markedly

reduced in response to both of the combined treatments. These results indicate that the

CGA or EGCG together with the TC-HT synergistically induced cell cycle arrest in the

G2/M phase by decreasing cyclin B1 and Cdc2 in the cells, and eventually inhibited the

cell growth of PANC-1 cells.

Fig 3. Effects of CGA or EGCG combined with the TC-HT on the G2/M cell cycle

arrest in PANC-1 cells. Cells were treated with the TC-HT (10 cycles), 0.05% DMSO

(vehicle control), CGA (200 µM), and EGCG (20 µM) alone or in combination for 24 h

and stained with propidium iodide (PI) for cell cycle analysis. (A) Representative DNA

content profiles. (B) The percentage of cell population in each phase of the cell cycle.

(C) Western bolt analysis of the expression of cell cycle regulator proteins cyclin B1

and Cdc2. β-actin was used as an internal control. Data are presented as mean ± S.D. in

triplicate. ($^{**}p < 0.01$ and $^{***}p < 0.001$ compared with the untreated control)

TC-HT combined with polyphenols induces apoptosis in PANC-1 cells

We then investigated as to whether the synergistic cytotoxic effect of the TC-HT

with the CGA or EGCG was associated with the induction of apoptosis, and the flow

cytometric analysis was performed using Annexin V-FITC/PI staining. As shown in

Figs 4A and **4B**, the fraction of apoptotic cells after the treatment of PANC-1 cells with

either the single agent CGA, or the EGCG or TC-HT were barely increased as

compared with untreated control cells. There was no significant difference in apoptosis

between the 0.05% DMSO (vehicle control) combined with or without TC-HT and the

untreated control. Interestingly, the TC-HT in combination with the CGA significantly

increased the early apoptotic cell death in PANC-1 cells (18.6±5.2%). A similar result

was also obtained in the combination of the TC-HT and the EGCG, but to a greater

extent (24.3±0.8%). These results also showed that the late apoptotic cell death was

increased following the exposure of TC-HT with CGA (15.7±4.6%) or EGCG

(18.4±4.6%). Furthermore, in contrast to control cells, the DAPI staining demonstrated

that the cells treated with either of the two combined treatments had more condensed

and fragmented nuclei, which are typical morphological alterations for the apoptosis

(**Fig 4C**). These results suggest that the TC-HT has a potential synergistic effect with

the CGA or EGCG on the apoptosis in the PANC-1 cells.

Fig 4. Combination of the TC-HT and the CGA or EGCG induces apoptosis in

PANC-1 cells. The apoptosis analysis for the cells following the treatment with the

TC-HT (10 cycles), 0.05% DMSO (vehicle control), CGA (200 μM), and EGCG (20

μM) alone or in combination for 24 h. (A) Flow cytometric detection of the apoptosis

with Annexin V-FITC/PI double staining. (B) Histogram quantifying the percentage of

PANC-1 cells in early and late apoptosis. (C) The nuclei morphology alterations (arrow)

were examined using DAPI staining. Data are presented as mean ± S.D. in triplicate.

($^{**}p < 0.01$ and $^{***}p < 0.001$ compared with the untreated control)

Combination of the TC-HT and polyphenols triggers significant loss in the mitochondrial membrane potential in PANC-1 cells

Collapse of the mitochondrial integrity is a critical event in the cells undergoing

apoptosis. To examine whether the combination of the TC-HT and CGA- or

EGCG-induced apoptosis involved mitochondrial disruption, the mitochondrial

membrane potential (MMP) was assessed using $DiOC_6(3)$ fluorescence staining by

flow cytometric analysis [31]. As shown in **Figs 5A** and **5B**, the treatment with the

EGCG, the CGA and its corresponding vehicle control (0.05% DMSO) did not change

the MMP level in comparison with the untreated control. In response to treatment with

the TC-HT alone or together with vehicle control, there was no significant change in

MMP within cells; however, the cells treated with TC-HT showed a stronger effect on

the MMP depolarization after the administration of CGA or EGCG (44±6.4% and

75.7±7.6%), which was consistent with the results of the apoptosis analysis as

described in **Fig 4**. These results indicate that the TC-HT combined with the CGA- or

EGCG-induced apoptosis in the PANC-1 cells is mediated by mitochondrial

dysfunction.

Fig 5. Combination of the TC-HT and the CGA or EGCG induces mitochondrial

dysfunction. (A) Flow cytometric analysis of MMP using $DiOC_6(3)$ after treatment

with the TC-HT (10 cycles), 0.05% DMSO (vehicle control), CGA (200 μM), and

EGCG (20 μM) alone or in combination for 24 h. The M1 regions indicate the cells

with the loss of MMP. (B) Histogram represents the percentage of cells in the M1

region. Data are presented as mean ± S.D. in triplicate. ($^{**}p < 0.01$ and $^{***}p < 0.001$

compared with the untreated control)

Combination of the TC-HT and polyphenols induces apoptosis through the mitochondrial pathway

It is well known that Bcl-2 family proteins and caspases, along with PARP, play important roles in the mitochondria-mediated apoptosis. To further explore the mechanism by which the TC-HT combined with the CGA or EGCG triggered apoptosis in the PANC-1 cells, we evaluated the expression of the apoptosis-related proteins using western blot analysis. As shown in **Fig 6**, when compared with control cells, the cleaved caspase-9, -3, and the cleaved PARP were markedly increased after the co-administration with the TC-HT and either the CGA or EGCG. In addition, both of the combined treatments down-regulated the expression of Bcl-2 and up-regulated the expression of Bax, and thus decreased the ratio of Bcl-2 to Bax. These results reveal that the TC-HT combined with the CGA or EGCG promotes apoptosis in the PANC-1 cells via activation of the mitochondrial pathway.

Fig 6. Effects of the TC-HT, CGA and EGCG on the expression of the apoptosis-related proteins in PANC-1 cells. (A)(B) The protein levels of cleaved caspase-9, -3, cleaved PARP, Bcl-2 and Bax of the PANC-1 cells treated with the

374   TC-HT (10 cycles), CGA (200 μM) and EGCG (20 μM) alone or in combination

375   treatment for 24 h were examined by western blot analysis. GAPDH and β-actin were

376   used as internal controls.

The role of ROS in the TC-HT combined with the CGA- or EGCG-induced apoptosis in PANC-1 cells

As ROS generation is a critical event in the induction of the apoptosis, we next

examined the role of ROS in apoptosis induced by treatment with the TC-HT in

combination with either the CGA or EGCG. The intracellular ROS was measured by

flow cytometry using a fluorescence probe, DHE, which reacts with $O_2^{\bullet-}$ [32]. As

shown in **Fig 7A**, the treatment of PANC-1 cells with 0.05% DMSO (vehicle control),

CGA or EGCG alone did not alter the level of ROS when compared with the untreated

control cells. Also, the level of ROS induced by TC-HT and its combination with

DMSO were found to be comparable to the untreated control in the MFI. It is worth

mentioning that the TC-HT combined with CGA and the TC-HT combined with EGCG

significantly increased the level of ROS in MFI by approximately 2.3- and 3.6-fold,

respectively (**Fig 7B**). These results suggest that the TC-HT combined with the CGA or

EGCG may induce apoptosis through the excessive ROS production in the PANC-1

cells.

Fig 7. Combined effects of the TC-HT and the CGA or EGCG on ROS generation

in PANC-1 cells. (A) The DHE ($O_2^{\bullet-}$) levels were measured by flow cytometry after

treatment with the TC-HT (10 cycles), 0.05% DMSO (vehicle control), CGA (200 μM),

and EGCG (20 μM) alone or in combination for 24 h. (B) Graph shows the fold change

in MFI of DHE relative to control. Data are presented as mean ± S.D. in triplicate. ([**]p

< 0.01 and [***]p < 0.001 compared with the untreated control)

Discussion

Despite previous studies having shown that HT or polyphenols, the EGCG and

CGA, possessed anticancer activities against pancreatic cancer cells [33-36], the effect

of these two agents as a combined therapy on pancreatic cancer has not been reported.

In this study, we have explored as to whether the herbal compounds of the CGA and

EGCG could sensitize the pancreatic cancer cells to heat, and elucidated the cellular

mechanism underlying the biological effects. Our results have demonstrated that the

exposure to heat using the TC-HT in combination with the CGA or EGCG

synergistically inhibited the growth and induced the apoptosis in the PANC-1 cells.

Previous studies have shown the benefits of HT as an adjuvant in combination with

chemotherapy against many various cancers, including pancreatic cancer [12, 37, 38].

However, there is a major concern that HT can cause unavoidable thermal injury to

normal cells. Therefore, it is important to refine the method of heat administration to

achieve the desired thermal dosage with the minimal cytotoxicity. To the best of our

knowledge, this is the first report indicating an effective and safe combination of the

TC-HT and the CGA or EGCG on pancreatic cancer cells.

Previous research has reported the response of PANC-1 cells to mild HT at

different temperatures [39]. Particularly, the temperature in a range of about 42 to 46°C

has been shown capable of inducing the cell death of the pancreatic cell line PANC-1.

Our survival analysis revealed a consistent result that the cell growth of PANC-1 in the

1-cycle group was significantly inhibited. Although HT has a cooperative cytotoxicity

with the CGA and EGCG, its severe thermal toxicity to a cell is what we want to avoid.

The usage of the repeated cycles of heat exposure could provide a means to synergize

with the anticancer compounds in the heating process, and avoid the thermal damage

accumulation in the following non-heating process. Earlier studies have shown the

time-dependent modification of cancer cells during exposure to HT for which the

analysis of cellular growth indicated that the survival rate decreased with increasing HT

exposure time [40, 41]. Namely, the short exposure of cancer cells to HT may induce

cellular stress without affecting their proliferation, while the prolonged exposure may

lead to cytotoxicity. It has also been reported that short HT treatment of PANC-1 cells

at the temperature 46°C for 5 min had no effect on cell viability [42]. Therefore, we

suggest that the 10-cycles TC-HT induced stress rather than damage in cells during

each heating cycle. As expected, the cells treated with the TC-HT of different cycling

parameters exhibited a decreased cytotoxicity with the increased cycles. It is worth

noting that 10 cycles of the TC-HT (43.5-36 °C), or either compound, did not cause a

significant alteration in the viability, however, the combination of 10 cycles of the

TC-HT with the CGA or EGCG strongly resulted in a dose-dependent decline of the

cell viability. Similar results from colony formation were also obtained with the cells

treated with both combinations. These results suggested that the CGA and EGCG may

act as heat synergizers and co-administered with the use of the TC-HT could have

significantly combined effects against pancreatic cancer cells.

Cancer is characterized by the loss of ability in the cell division arising from the

aberrant activity of various cell cycle regulators. HT has been demonstrated to inhibit

the growth of human cancer cells through interfering with the cell cycle progression [26,

28, 43]. In our results, we found that the TC-HT did not cause an obvious accumulation

of cells in the G2/M phase. When cells undergoing the TC-HT were given the CGA or

EGCG concurrently, the proportion of cells in the G2/M phase was significantly

increased. Additionally, this observation was supported by the marked downregulation

of Cdc2 and cyclin B1 proteins in the combination of the TC-HT and the CGA or

EGCG. The Cdc2 (cell division cycle 2), also known as the CDK1 (cyclin-dependent

kinase 1), is a core regulator that drives the cells through G2 and into mitosis. Several

studies have reported that the binding of Cdc2 to cyclin B1 complex plays an important

role in the G2/M progression [44, 45]. Collectively, these results indicated that the

synergistic cytotoxicity of CGA and EGCG under the exposure of TC-HT, at least in

part, was associated with the inhibition of the Cdc2/cyclin B1 kinase activity.

Apoptosis, also the best known form of programmed cell death, plays a pivotal

role in defending against cancer. The conventional HT has been demonstrated to affect

apoptotic pathways in various types of cancer cells [46]. It has also been reported that

polyphenols exhibit anticancer activities against different human cancer cells through

activating the apoptotic pathway [47, 48]. However, in our study, the results of

apoptosis analysis performed using FITC Annexin V and PI double staining indicated

that neither the TC-HT nor these two compounds induced significant apoptotic cells

(**Fig 4**). Only when the TC-HT in combination with the CGA or EGCG was conducted,

both of the combined treatments significantly elevated the percentage of the early and

late apoptotic cells. This finding was further confirmed by the nuclear morphological

alterations of the apoptosis in the TC-HT-treated cells in combination with the CGA or

EGCG. The apoptotic process can be divided into the death receptor and mitochondrial

pathways, and the mitochondrial pathway of cell death is thought to be the major

mechanism of apoptosis in mammals. The Bcl-2 family proteins are key regulators of

the mitochondrial apoptotic pathway, which comprise of both the anti-apoptotic and

pro-apoptotic members. Unbalanced Bcl-2/Bax ratio within the cells induces the

disruption of the mitochondrial membrane, release of cytochrome c, activation of

caspases, and the subsequent cleavage of PARP [49, 50]. Results from **Figs 5** and **6**

indicated that the combined treatment with the TC-HT and the CGA or EGCG

markedly decreased the level of MMP and Bcl-2 expression, and increased Bax

expression. We also found that the activation of caspase-9, -3, and the cleavage of

PARP were promoted with both of the combined treatments (**Fig 6**). Collectively, these

data indicated that the mitochondria-dependent pathway is involved in the synergistic

apoptosis following the combined treatments.

Mitochondria are widely believed to be the main cellular source of ROS. The

excessive production of ROS could result in mitochondrial dysfunction, which in turn

triggers the apoptosis [51]. HT-mediated oxidative stress has been suggested to be

responsible for the HT-induced apoptosis [52]. Several reports have provided evidence

that HT increased cellular level of ROS including $O_2^{\bullet-}$, which is the precursor of most

ROS and a mediator in oxidative chain reactions [25, 27, 53]. We then confirmed if the

observation of mitochondrial dysfunction in cells treated with the TC-HT in

combination with the CGA or EGCG was promoted by ROS generation. Our result

showed that the cellular ROS generation was elevated significantly following the

administration of combination of the TC-HT with the CGA or EGCG, while no

significant change was observed in the level of ROS when the TC-HT was applied

alone. Taken together, these results indicated that the high levels of ROS production

played an important role in the apoptosis induced by the combination of the TC-HT

with CGA and the TC-HT with EGCG. Most importantly, based on our results, the

TC-HT could serve as a harmless approach to synergize with the natural compounds,

CGA and EGCG. We also investigate the feasibility of the TC-HT by using standard

clinical compound gemcitabine known to synergize with standard HT [54, 55]. Similar

result as the combination of 1-cycle TC-HT and gemcitabine was observed from

analysis of cell viability in the combined treatment with the 10-cycles TC-HT and

gemcitabine, whereas the 10-cycles TC-HT alone had no effect on cell viability (**S1**

Fig.). This finding suggests that this approach can be extended to other anticancer agent

with heat synergy in treatment of cancer.

High-intensity focused ultrasound (HIFU) as a hyperthermal technique has been

extensively applied in cancer treatment, such as cancers of liver, prostate and pancreas

[56]. Besides its use for thermal ablation, it has also been shown to be adopted to

produce mild HT in tumor [57, 58]. Particularly, the use of multi-focal HIFU has

demonstrated the improvements in controlling temperature accurately and uniformly

when performing mild HT with multi-focal HIFU in a scanning manner [59]. On the

other hand, the technique of multiple focal points induced simultaneously by the

electronically focussed beam allows the size of focal region to be increased [59, 60].

The TC-HT introduced in this study represents an approach for mild thermal treatment

performed by using a series of short period of heat exposure within the desired time. In

view of a fast elevation of temperature in local region achievable by HIFU exposure,

we suggest that the HIFU can be applied to reach the target temperature during each

heating cycle of the TC-HT. Furthermore, the concept of pre-cooling [61] is also

needed to be involved for rapidly returning to physiological temperature following the

heating cycle, through which by cooling the local area surrounding tumor cells to a

temperature lower than 37 °C to quickly dissipate the heat away from tumor. Therefore,

we believe that this work opens the way for development of the TC-HT as a cancer

treatment strategy.

In conclusion, a novel method for producing a desired thermal dosage is proposed

in which it applies repeated thermal treatments of short exposure to prevent the toxic

effects from prolonged exposure. Our findings have demonstrated the TC-HT shows

capability of synergizing with the natural compounds of CGA and EGCG and

minimizing the thermal damage resulting from HT. The synergistic activity against the

PANC-1 cells was performed primarily via the G2/M arrest and the apoptosis induced

through the ROS-mediated mitochondrial pathway leading to an imbalance between

Bcl-2 and Bax, activation of caspase-9 and -3, and the cleavage of PARP. We believe

this strategy potentially provides an improvement to the therapeutic window of HT in

treating cancer. Gaining an optimal therapy for a variety of cancers via the examination

of the relation between the TC-HT parameters and other anticancer drugs is an

attractive subject that warrants further investigation.

Funding

534   This work was supported by grants from Ministry of Science and Technology

535 (MOST 105-2112-M-002-006-MY3 to CYC) and Ministry of Education (MOE

536 106R880708 to CYC) of the Republic of China. The funders had no role in study

537 design, data collection and analysis, decision to publish, or preparation of the

538 manuscript.

Competing Interests

541   The authors have declared that no competing interests exist.

Acknowledgments

544   We would like to thank Technology Commons in College of Life Science,

545   National Taiwan University for use of flow cytometry system, and the staff of the

546   imaging core at the First Core Labs, National Taiwan University hospital for technical

547   assistance.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Chueh-Hsuan Lu.

Formal analysis: Chueh-Hsuan Lu, Wei-Ting Chen, Chih-Hsiung Hsieh, Yu-Yi Kuo,

Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Chueh-Hsuan Lu, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Resources: Chih-Hsiung Hsieh, Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Chueh-Hsuan Lu.

Writing – original draft: Chueh-Hsuan Lu, Wei-Ting Chen, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

Submission

567
568

References

1.   Siegel RL, Miller KD, Jemal A. Cancer statistics, 2016. CA Cancer J Clin.
2016;66(1):7-30. doi: 10.3322/caac.21332. PubMed PMID: 26742998.

2.   Fryer RA, Galustian C, Dalgleish AG. Recent advances and developments in
treatment strategies against pancreatic cancer. Curr Clin Pharmacol.
2009;4(2):102-12. PubMed PMID: 19442075.

3.   Overgaard J. The Current and Potential Role of Hyperthermia in
Radiotherapy. Int J Radiat Oncol. 1989;16(3):535-49.

4.   Roti JLR. Cellular responses to hyperthermia (40-46 degrees C): Cell killing
and molecular events. Int J Hyperther. 2008;24(1):3-15. doi:
10.1080/02656730701769841. PubMed PMID: WOS:000252581400002.

5.   Oei AL, Vriend LEM, Crezee J, Franken NAP, Krawczyk PM. Effects of
hyperthermia on DNA repair pathways: one treatment to inhibit them all. Radiat
Oncol. 2015;10. doi: ARTN 165
10.1186/s13014-015-0462-0. PubMed PMID: WOS:000358983800001.

6.   Huang SH, Yang KJ, Wu JC, Chang KJ, Wang SM. Effects of hyperthermia on
the cytoskeleton and focal adhesion proteins in a human thy carcinoma cell line. J
Cell Biochem. 1999;75(2):327-37. doi: Doi
10.1002/(Sici)1097-4644(19991101)75:2<327::Aid-Jcb14>3.0.Co;2-G. PubMed
PMID: WOS:000082783600014.

7.   Calderwood SK, Theriault JR, Gong JL. How is the immune response affected
by hyperthermia and heat shock proteins? Int J Hyperther. 2005;21(8):713-6. doi:

10.1080/02656730500340794. PubMed PMID: WOS:000233906600007.

8.   Ponce AM, Vujaskovic Z, Yuan F, Needham D, Dewhirst MW. Hyperthermia

mediated liposomal drug delivery. Int J Hyperther. 2006;22(3):205-13. doi:

10.1080/02656730600582956. PubMed PMID: WOS:000238093900004.

9.   Song CW, Shakil A, Osborn JL, Iwata K. Tumour oxygenation is increased by

hyperthermia at mild temperatures. Int J Hyperther. 2009;25(2):91-5. doi: Pii

910057702

10.1080/02656730902744171. PubMed PMID: WOS:000264750400001.

10.   Kaur P, Hurwitz MD, Krishnan S, Asea A. Combined hyperthermia and

radiotherapy for the treatment of cancer. Cancers (Basel). 2011;3(4):3799-823.

doi: 10.3390/cancers3043799. PubMed PMID: 24213112; PubMed Central

PMCID: PMCPMC3763397.

11.   Au JLS, Wientjes MG. Combination Intravesical Hyperthermia and

Chemotherapy for Bladder Cancer The Rampersaud/Vujaskovic/Inman Article

Reviewed. Oncology-Ny. 2010;24(12):1155-60. PubMed PMID:

WOS:000293341400009.

12.   Adachi S, Kokura S, Okayama T, Ishikawa T, Takagi T, Handa O, et al. Effect of

hyperthermia combined with gemcitabine on apoptotic cell death in cultured

human pancreatic cancer cell lines. Int J Hyperthermia. 2009;25(3):210-9.

13.   Refaat T, Sachdev S, Sathiaseelan V, Helenowski I, Abdelmoneim S, Pierce MC,

et al. Hyperthermia and radiation therapy for locally advanced or recurrent

breast cancer. Breast. 2015;24(4):418-25. doi: 10.1016/j.breast.2015.03.008.

PubMed PMID: 25900383; PubMed Central PMCID: PMCPMC4791303.

14.   Bakshandeh-Bath A, Stoltz AS, Homann N, Wagner T, Stolting S, Peters SO.

Preclinical and Clinical Aspects of Carboplatin and Gemcitabine Combined with

Whole-body Hyperthermia for Pancreatic Adenocarcinoma. Anticancer Res.

2009;29(8):3069-77. PubMed PMID: WOS:000268846800029.

15.   Tschoep-Lechner KE, Milani V, Berger F, Dieterle N, Abdel-Rahman S, Salat C,

et al. Gemcitabine and cisplatin combined with regional hyperthermia as

second-line treatment in patients with gemcitabine-refractory advanced

pancreatic cancer. Int J Hyperther. 2013;29(1):8-16. doi:

10.3109/02656736.2012.740764. PubMed PMID: WOS:000313672700002.

16.   Scalbert A, Williamson G. Dietary intake and bioavailability of polyphenols. J

Nutr. 2000;130(8):2073s-85s. PubMed PMID: WOS:000088760200003.

17.   Richelle M, Tavazzi I, Offord E. Comparison of the antioxidant activity of

commonly consumed polyphenolic beverages (coffee, cocoa, and tea) prepared

per cup serving. J Agr Food Chem. 2001;49(7):3438-42. doi: DOI

10.1021/jf0101410. PubMed PMID: WOS:000170043400047.

18. Min KJ, Kwon TK. Anticancer effects and molecular mechanisms of

epigallocatechin-3-gallate. Integr Med Res. 2014;3(1):16-24. doi:

10.1016/j.imr.2013.12.001. PubMed PMID: 28664074; PubMed Central PMCID:

PMCPMC5481703.

19. Belkaid A, Currie JC, Desgagnes J, Annabi B. The chemopreventive properties

of chlorogenic acid reveal a potential new role for the microsomal

glucose-6-phosphate translocase in brain tumor progression. Cancer Cell Int.

2006;6:7. doi: 10.1186/1475-2867-6-7. PubMed PMID: 16566826; PubMed

Central PMCID: PMCPMC1440869.

20. Zhao H, Zhu W, Jia L, Sun X, Chen G, Zhao X, et al. Phase I study of topical

epigallocatechin-3-gallate (EGCG) in patients with breast cancer receiving

adjuvant radiotherapy. Br J Radiol. 2016;89(1058):20150665. Epub 2015/11/27.

doi: 10.1259/bjr.20150665. PubMed PMID: 26607642; PubMed Central PMCID:

PMCPMC4985212.

21. Gee JR, Saltzstein DR, Kim K, Kolesar J, Huang W, Havighurst TC, et al. A

Phase II Randomized, Double-blind, Presurgical Trial of Polyphenon E in Bladder

Cancer Patients to Evaluate Pharmacodynamics and Bladder Tissue Biomarkers.

Cancer Prev Res. 2017;10(5):298-307. doi: 10.1158/1940-6207.Capr-16-0167.

PubMed PMID: WOS:000400625500005.

22. Xu RS, Kang QM, Ren J, Li ZK, Xu XP. Antitumor Molecular Mechanism of

Chlorogenic Acid on Inducting Genes GSK-3 beta and APC and Inhibiting Gene

beta-Catenin (vol 2013, 951319, 2013). J Anal Methods Chem. 2015. doi: Artn

189328

10.1155/2015/189328. PubMed PMID: WOS:000352383000001.

23. Feng RT, Lu YJ, Bowman LL, Qian Y, Castranova V, Ding M. Inhibition of

activator protein-1, NF-kappa B, and MAPKs and induction of phase 2 detoxifying

enzyme activity by chlorogenic acid. J Biol Chem. 2005;280(30):27888-95. doi:

10.1074/jbc.M503347200. PubMed PMID: WOS:000230678600051.

24. Kim JH, Kim SH, Alfieri AA, Young CW. Quercetin, an Inhibitor of Lactate

Transport and a Hyperthermic Sensitizer of Hela-Cells. Cancer Res.

1984;44(1):102-6. PubMed PMID: WOS:A1984RX69400019.

25. Cui ZG, Piao JL, Kondo T, Ogawa R, Tsuneyama K, Zhao QL, et al. Molecular

mechanisms of hyperthermia-induced apoptosis enhanced by docosahexaenoic acid: Implication for cancer therapy. Chem-Biol Interact. 2014;215:46-53. doi: 10.1016/j.cbi.2014.03.005. PubMed PMID: WOS:000336357600007.

26. Zhao PJ, Jiang H, Su D, Feng JG, Ma SL, Zhu XH. Inhibition of cell proliferation by mild hyperthermia at 43 degrees C with Paris Saponin I in the lung adenocarcinoma cell line PC-9. Mol Med Rep. 2015;11(1):327-32. doi: 10.3892/mmr.2014.2655. PubMed PMID: WOS:000346651100045.

27. Cui ZG, Piao JL, Rehman MUR, Ogawa R, Li P, Zhao QL, et al. Molecular mechanisms of hyperthermia-induced apoptosis enhanced by withaferin A. Eur J Pharmacol. 2014;723:99-107. doi: 10.1016/j.ejphar.2013.11.031. PubMed PMID: WOS:000330574600014.

28. Hatashita M, Taniguchi M, Baba K, Koshiba K, Sato T, Jujo Y, et al. Sinodielide A exerts thermosensitizing effects and induces apoptosis and G2/M cell cycle arrest in DU145 human prostate cancer cells via the Ras/Raf/MAPK and PI3K/Akt signaling pathways. Int J Mol Med. 2014;33(2):406-14. doi: 10.3892/ijmm.2013.1568. PubMed PMID: 24285252.

29. Dewey WC. Arrhenius Relationships from the Molecule and Cell to the Clinic. Int J Hyperther. 1994;10(4):457-83. doi: Doi 10.3109/02656739409009351. PubMed PMID: WOS:A1994NX92300001.

30. Sminia P, Vanderzee J, Wondergem J, Haveman J. Effect of Hyperthermia on the Central-Nervous-System - a Review. Int J Hyperther. 1994;10(1):1-30.

31. Korchak HM, Rich AM, Wilkenfeld C, Rutherford LE, Weissmann G. A carbocyanine dye, DiOC6(3), acts as a mitochondrial probe in human neutrophils. Biochem Biophys Res Commun. 1982;108(4):1495-501. PubMed PMID: 7181903.

32. Gomes A, Fernandes E, Lima JL. Fluorescence probes used for detection of reactive oxygen species. Journal of biochemical and biophysical methods. 2005;65(2-3):45-80.

33. Xiaoping L, Leizhen Z. Advances of high intensity focused ultrasound (HIFU) for pancreatic cancer. Int J Hyperthermia. 2013;29(7):678-82. doi: 10.3109/02656736.2013.837199. PubMed PMID: 24102396.

34. Wang K, Chen L, Meng Z, Lin J, Zhou Z, Wang P, et al. High intensity focused ultrasound treatment for patients with advanced pancreatic cancer: a preliminary dosimetric analysis. Int J Hyperthermia. 2012;28(7):645-52. doi: 10.3109/02656736.2012.713541. PubMed PMID: 22946544.

35.   Bimonte S, Cascella M, Leongito M, Palaia R, Caliendo D, Izzo F, et al. An
overview of pre-clinical studies on the effects of (-)-epigallocatechin-3-gallate, a
catechin found in green tea, in treatment of pancreatic cancer. Recenti Prog Med.
2017;108(6):282-7. Epub 2017/06/21. doi: 10.1701/2715.27715. PubMed PMID:
28631776.

36.   Dong J, Zou J, Yu XF. Coffee drinking and pancreatic cancer risk: a
meta-analysis of cohort studies. World J Gastroenterol. 2011;17(9):1204-10.
Epub 2011/03/31. doi: 10.3748/wjg.v17.i9.1204. PubMed PMID: 21448427;
PubMed Central PMCID: PMCPMC3063915.

37.   Miyamoto R, Oda T, Hashimoto S, Kurokawa T, Inagaki Y, Shimomura O, et al.
Cetuximab delivery and antitumor effects are enhanced by mild hyperthermia in
a xenograft mouse model of pancreatic cancer. Cancer Sci. 2016;107(4):514-20.

38.   Issels RD. Hyperthermia adds to chemotherapy. Eur J Cancer.
2008;44(17):2546-54. doi: 10.1016/j.ejca.2008.07.038. PubMed PMID:
18789678.

39.   Ware MJ, Nguyen LP, Law JJ, Krzykawska-Serda M, Taylor KM, Cao HST, et al.
A new mild hyperthermia device to treat vascular involvement in cancer surgery.
Sci Rep. 2017;7(1):11299. doi: 10.1038/s41598-017-10508-6. PubMed PMID:
28900126; PubMed Central PMCID: PMCPMC5595878.

40.   Dewey WC, Hopwood LE, Sapareto SA, Gerweck LE. Cellular responses to
combinations of hyperthermia and radiation. Radiology. 1977;123(2):463-74.
Epub 1977/05/01. doi: 10.1148/123.2.463. PubMed PMID: 322205.

41.   Raaphorst GP, Freeman ML, Dewey WC. Radiosensitivity and recovery from
radiation damage in cultured CHO cells exposed to hyperthermia at 42.5 or 45.5
degrees C. Radiat Res. 1979;79(2):390-402. PubMed PMID: 482603.

42.   Curley SA, Palalon F, Sanders KE, Koshkina NV. The Effects of Non-Invasive
Radiofrequency Treatment and Hyperthermia on Malignant and Nonmalignant
Cells. Int J Env Res Pub He. 2014;11(9):9142-53. doi: 10.3390/ijerph110909142.
PubMed PMID: WOS:000342027500031.

43.   Lim CU, Zhang Y, Fox MH. Cell cycle dependent apoptosis and cell cycle
blocks induced by hyperthermia in HL-60 cells. Int J Hyperthermia.
2006;22(1):77-91. doi: 10.1080/02656730500430538. PubMed PMID:
16423754.

44.   Jessus C, Beach D. Oscillation of Mpf Is Accompanied by Periodic Association
between Cdc25 and Cdc2-Cyclin-B. Cell. 1992;68(2):323-32. doi: Doi

10.1016/0092-8674(92)90473-P. PubMed PMID: WOS:A1992HB35300011.

45.   Fisher D, Krasinska L, Coudreuse D, Novak B. Phosphorylation network

dynamics in the control of cell cycle transitions. J Cell Sci. 2012;125(Pt

20):4703-11. doi: 10.1242/jcs.106351. PubMed PMID: 23223895.

46.   Ahmed K, Tabuchi Y, Kondo T. Hyperthermia: an effective strategy to induce

apoptosis in cancer cells. Apoptosis. 2015;20(11):1411-9. doi:

10.1007/s10495-015-1168-3. PubMed PMID: 26354715.

47.   Rodríguez M, Estrela J, Ortega Á. Natural polyphenols and apoptosis

induction in cancer therapy. J Carcinog Mutag S. 2013;6.

48.   Yamagata K, Izawa Y, Onodera D, Tagami M. Chlorogenic acid regulates

apoptosis and stem cell marker-related gene expression in A549 human lung

cancer cells. Mol Cell Biochem. 2018;441(1-2):9-19. doi:

10.1007/s11010-017-3171-1. PubMed PMID: 28875417.

49.   Tait SWG, Green DR. Mitochondrial Regulation of Cell Death. Csh Perspect

Biol. 2013;5(9). doi: ARTN a008706

10.1101/cshperspect.a008706. PubMed PMID: WOS:000325168700011.

50.   Gross A, McDonnell JM, Korsmeyer SJ. BCL-2 family members and the

mitochondria in apoptosis. Genes Dev. 1999;13(15):1899-911. PubMed PMID:

10444588.

51.   Orrenius S. Reactive oxygen species in mitochondria-mediated cell death.

Drug Metab Rev. 2007;39(2-3):443-55. doi: 10.1080/03602530701468516.

PubMed PMID: 17786631.

52.   Katschinski DM, Boos K, Schindler SG, Fandrey J. Pivotal role of reactive

oxygen species as intracellular mediators of hyperthermia-induced apoptosis. J

Biol Chem. 2000;275(28):21094-8. doi: DOI 10.1074/jbc.M001629200. PubMed

PMID: WOS:000088230600020.

53.   Davidson JF, Whyte B, Bissinger PH, Schiestl RH. Oxidative stress is involved

in heat-induced cell death in Saccharomyces cerevisiae. P Natl Acad Sci USA.

1996;93(10):5116-21. doi: DOI 10.1073/pnas.93.10.5116. PubMed PMID:

WOS:A1996UL25500109.

54.   Jin HB, Zhao YY, Zhang SR, Yang JF, Zhang XF, Ma SL. Hyperthermia inhibits

the motility of gemcitabine-resistant pancreatic cancer PANC-1 cells through the

inhibition of epithelial-mesenchymal transition. Mol Med Rep.

2018;17(5):7274-80. doi: 10.3892/mmr.2018.8763. PubMed PMID:

WOS:000430556800132.

55. Adachi S, Kokura S, Okayama T, Ishikawa T, Takagi T, Handa O, et al. Effect of

hyperthermia combined with gemcitabine on apoptotic cell death in cultured

human pancreatic cancer cell lines. Int J Hyperther. 2009;25(3):210-9. doi: Pii

911094556

10.1080/02656730802657036. PubMed PMID: WOS:000265991500006.

56. Hsiao YH, Kuo SJ, Tsai HD, Chou MC, Yeh GP. Clinical Application of

High-intensity Focused Ultrasound in Cancer Therapy. J Cancer.

2016;7(3):225-31. Epub 2016/02/27. doi: 10.7150/jca.13906. PubMed PMID:

26918034; PubMed Central PMCID: PMCPMC4747875.

57. Partanen A, Yarmolenko PS, Viitala A, Appanaboyina S, Haemmerich D,

Ranjan A, et al. Mild hyperthermia with magnetic resonance-guided

high-intensity focused ultrasound for applications in drug delivery. Int J

Hyperther. 2012;28(4):320-36. doi: 10.3109/02656736.2012.680173. PubMed

PMID: WOS:000304352400006.

58. Frazier N, Payne A, de Bever J, Dillon C, Panda A, Subrahmanyam N, et al.

High intensity focused ultrasound hyperthermia for enhanced macromolecular

delivery. J Control Release. 2016;241:186-93. doi: 10.1016/j.jconrel.2016.09.030.

PubMed PMID: WOS:000386251700016.

59. Partanen A, Tillander M, Yarmolenko PS, Wood BJ, Dreher MR, Kohler MO.

Reduction of peak acoustic pressure and shaping of heated region by use of

multifoci sonications in MR-guided high-intensity focused ultrasound mediated

mild hyperthermia. Medical Physics. 2013;40(1). doi: Artn 013301

10.1118/1.4769116. PubMed PMID: WOS:000313033200054.

60. Payne A, Vyas U, Todd N, de Bever J, Christensen DA, Parker DL. The effect of

electronically steering a phased array ultrasound transducer on near-field tissue

heating. Med Phys. 2011;38(9):4971-81. doi: 10.1118/1.3618729. PubMed PMID:

21978041; PubMed Central PMCID: PMCPMC3166338.

61. Vaity C, Al-Subaie N, Cecconi M. Cooling techniques for targeted temperature

management post-cardiac arrest. Crit Care. 2015;19. doi: ARTN 103

10.1186/s13054-015-0804-1. PubMed PMID: WOS:000352009500002.

Supporting information

S1 Fig. PANC-1 cell viability results from MTT assay after the treatments of

gemcitabine (5 μM) and TC-HT (1 cycle and 10 cycles) alone or in combination

for 24 h.

A

TC-HT	−	−	+	+
CGA	−	+	−	+

Cleaved caspase-9

Cleaved caspase-3

Cleaved PARP

GAPDH

Bcl-2

Bax

β-actin

B

TC-HT	−	−	+	+
EGCG	−	+	−	+

Cleaved caspase-9

Cleaved caspase-3

Cleaved PARP

GAPDH

Bcl-2

Bax

β-actin