Resolving multisensory and attentional influences across cortical depth in sensory cortices

Authors/Affiliations

Remi Gau[a,e*], Pierre-Louis Bazin[b,c], Robert Trampel[b], Robert Turner[b,d], Uta Noppeney[a,f]

a Computational Neuroscience and Cognitive Robotics Centre, University of Birmingham, B15 2TT Birmingham, UK

b Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

c Integrative Model-based Cognitive Neuroscience research unit, University of Amsterdam, Amsterdam, The Netherlands

d Sir Peter Mansfield Imaging Centre, University of Nottingham, NG7 2RD, Nottingham, UK

e Institute of Psychology and Institute of Neuroscience, Université Catholique de Louvain, 1348 Louvain-la-Neuve, Belgium

f Donders Institute for Brain, Cognition, Behaviour, Radboud University, Nijmegen, The Netherlands

* Corresponding author: Remi Gau <remi.gau@gmail.com>

Classification

Social sciences/Psychological and cognitive sciences

Biological sciences/Neuroscience

Keywords: multisensory integration, audiovisual, primary sensory cortices, deactivations, crossmodal attention, cortical layers, layer-dependent fMRI, ultra-high-resolution fMRI, 7 Tesla

ABSTRACT

In our environment our senses are bombarded with a myriad of signals, only a subset of which is

relevant for our goals. Using sub-millimeter-resolution fMRI at 7T we resolved BOLD-response

and activation patterns across cortical depth in early sensory cortices to auditory, visual and

audiovisual stimuli under auditory or visual attention. In visual cortices, auditory stimulation

induced widespread inhibition irrespective of attention, whereas auditory relative to visual attention

suppressed mainly central visual field representations. In auditory cortices, visual stimulation

suppressed activations, but amplified responses to concurrent auditory stimuli, in a patchy

topography. Critically, multisensory interactions in auditory cortices were stronger in deeper

laminae, while attentional influences were greatest at the surface. These distinct depth-dependent

profiles suggest that multisensory and attentional mechanisms regulate sensory processing via

partly distinct circuitries. Our findings are crucial for understanding how the brain regulates

information flow across senses to interact with our complex multisensory world.

INTRODUCTION

In our natural environment our senses are exposed to a constant influx of sensory signals that arise from many different sources. How the brain flexibly regulates information flow across the senses to enable effective interactions with the world remains unclear.

Mounting evidence from neuroimaging (Beauchamp et al., 2004; Noesselt et al., 2007; Rohe and Noppeney, 2016), neurophysiology (Atilgan et al., 2018; Kayser et al., 2010, 2008; Lakatos et al., 2007) and neuroanatomy (Falchier et al., 2002; Rockland and Ojima, 2003) suggests that interactions across the senses are pervasive in neocortex, arising even in primary cortices (Driver and Noesselt, 2008; Ghazanfar and Schroeder, 2006; Liang et al., 2013; Schroeder and Foxe, 2002). Visual stimuli can directly drive as well as modulate responses in cortices that are dedicated to other sensory modalities. Most prominently, functional magnetic resonance imaging (fMRI) in humans has shown that visual stimuli can induce crossmodal deactivations in primary and secondary auditory cortices (Laurienti et al., 2002; Leitão et al., 2013; Mozolic et al., 2008), yet enhance the response to a concurrent auditory stimulus (Werner and Noppeney, 2011, 2010). Further, neurophysiological research has suggested that multisensory influences emerge early in sensory cortices and are to some extent preserved in anaesthetized animals (Butler et al., 2012; Ibrahim et al., 2016; Iurilli et al., 2012; Kayser et al., 2007; Mercier et al., 2013). Yet, the ability to extrapolate from neurophysiological findings in animals to human fMRI studies is limited by the nature of the BOLD response, which pools neural activity over time and across a vast number of neurons (Logothetis, 2008).

Information flow is regulated not only by multisensory but also by attentional mechanisms that are guided by our current goals (Fairhall and Macaluso, 2009; Talsma et al., 2010). Critically, multisensory and attentional mechanisms are closely intertwined. Both enhance perceptual sensitivity (Leo et al., 2011) and precision of sensory representations (Ernst and Bülthoff, 2004; Fetsch et al., 2012; Meijer et al., 2019; Rohe and Noppeney, 2015). Most importantly, the co-occurrence of two congruent sensory stimuli boosts the salience of an event (Lewis and Noppeney,

2010; Van der Burg et al., 2008), which may thereby attract greater attention. Conversely, a

stimulus presented in one sensory modality alone may withdraw attentional resources from other

sensory modalities. Behavioural and functional imaging studies have shown that shifting attention

endogenously to one sensory modality reduces processing and activations in the unattended sensory

systems (Ciaramitaro et al., 2007; Johnson and Zatorre, 2005; Mozolic et al., 2008; Rohe and

Noppeney, 2018, 2015; Shomstein and Yantis, 2004). As a consequence, attentional mechanisms

may contribute to competitive and cooperative interactions across the senses, for instance by

amplifying responses for congruent audiovisual stimuli, and generating crossmodal deactivations

for unisensory stimuli. Inter-sensory attention can also profoundly modulate multisensory

interactions (Talsma et al., 2006). Most prominently, the influence of visual stimuli on auditory

cortices was shown to be enhanced when attention was focused on the visual sense (Lakatos et al.,

2009).

While previous neurophysiological studies have revealed influences of modality-specific attention

predominantly in superficial laminae in non-human primates (Lakatos et al., 2009), visual

influences on auditory cortices have recently been shown to be most prominent in deep layer 6 of

auditory cortex in rodents (Morrill and Hasenstaub, 2018). In other words, combined

neurophysiological evidence from primates and rodents suggests a double disassociation of

attentional and multisensory influences.

To investigate whether this double dissociation can be found in human neocortex we exploited

recent advances in submillimeter-resolution fMRI at 7T that allow the characterization of depth-

dependent activation profiles (De Martino et al., 2015a; Duong et al., 2003; Harel et al., 2006; Kok

et al., 2016; Koopmans et al., 2010; Muckli et al., 2015; Polimeni et al., 2010; Trampel et al., 2012).

While gradient-echo echo-planar-imaging (GE-EPI) BOLD fMRI is not yet able to attribute

activations unequivocally to specific cortical layers (Duong et al., 2003; Goense et al., 2012; Harel

et al., 2006; Huber et al., 2017; Markuerkiaga et al., 2016; Trampel et al., 2017), the observation in

the same cortical territories of distinct laminar activation profiles induced by multisensory and

attentional influences would strongly imply distinct neural mechanisms.

The current study investigated the processing of auditory, visual or audiovisual looming stimuli

under auditory and visual attention in the human brain. Combining submillimeter-resolution fMRI

at 7T with laminar and multivariate pattern analyses we show distinct depth-dependent activation

profiles and/or patterns for multisensory and attentional influences in early auditory and visual

cortices. These results suggest that multisensory and attentional mechanisms regulate sensory

processing in early sensory cortices via partly distinct neural circuitries.

RESULTS

In this fMRI study, participants were presented with blocks of auditory (A), visual (V) and audiovisual (AV) looming stimuli interleaved with fixation (Figure 1A). We used looming motion as a biologically relevant and highly salient stimulus that reliably evokes crossmodal influences in sensory cortices in human neuroimaging and animal neurophysiology (Cappe et al., 2012; Maier et al., 2008; Tyll et al., 2013). Modality-specific attention was manipulated by requiring participants to detect and respond selectively to weak auditory or visual targets, which were adjusted prior to the main study to threshold performance in sound amplitude or visual size for each participant. The targets were interspersed throughout all auditory, visual and audiovisual looming blocks (e.g. visual targets were presented during both visual and auditory looming blocks; see Figure 1B).

For the behavioural analysis, the percentages of visual and auditory targets that gained a response (see Supplementary Figure 1 and Supplementary Table 1 for '% target responses') were entered into a 2 (target modality: auditory vs. visual) X 2 (attended modality: auditory vs. visual) X 3 (stimulus block modality: auditory, visual, audio-visual) repeated measures ANOVA. We observed significant main effects of attended modality ($F_{(1, 10)}=291.263$, $p<0.001$, $\eta_p^2=0.967$), target modality ($F_{(1, 10)}=13.289$, $p=0.004$, $\eta_p^2=0.571$), and stimulus modality ($F_{(1.986, 19.864)}=5.304$, $p=0.014$, $\eta_p^2=0.347$), together with a significant interaction between target modality and attended modality ($F_{(1, 10)}=14.786$, $p=0.003$, $\eta_p^2=0.597$). This interaction confirmed that participants had successfully maintained auditory or visual attention as instructed.

Further, we observed significant interactions between target modality and stimulus block modality ($F_{(1.813, 18.126)}=8.149$, $p=0.004$, $\eta_p^2=0.449$) and between attended modality and stimulus block modality ($F_{(1.436, 14.360)}=24.034$, $p<0.001$, $\eta_p^2=0.706$). These interactions resulted from greater hit rates for auditory targets given auditory attention than for visual targets given visual attention during both auditory ($t_{(10)}=2.845$, $p=0.017$, Hedges $g_{av}=0.804$) and visual blocks ($t_{(10)}=4.432$, $p=0.001$, Hedges $g_{av}=2.037$), but not for audio-visual blocks ($t_{(10)}=0.276$, $p=0.788$, Hedges $g_{av}=0.081$) suggesting that observers' performance was not completely matched across all

conditions. Specifically, the presentation of auditory and visual stimuli in the audiovisual blocks

interfered with the detection of auditory targets. For completeness, there was no significant 3-way

interaction (F(1.783, 17.826)=2.467, p=0.118, η_p^2=0.198).

Using sub-millimeter resolution fMRI, we characterized the laminar profiles in auditory (primary

auditory cortex, A1; planum temporale, PT) and visual (primary, V1; higher order, V2/3) regions

for the following effects: 1. sensory deactivations in unisensory contexts for non-preferred stimuli

(i.e. crossmodal, e.g. [V-Fix] in auditory cortices), 2. crossmodal modulation (e.g. [AV-A] in

auditory cortices, [AV-V] in visual cortices) in audiovisual context and 3. direct and modulatory

effects of modality-specific attention.

Briefly, we used the following methodological approach (see Material and methods): in each ROI

and participant, we estimated the regional BOLD response (i.e. 'B parameter estimate') (e.g. Figure

2A row 1, left) and the multivariate pattern decoding accuracy for each of the six laminae (e.g.

Figure 3A row 1, right). We then characterized the laminar profiles of BOLD response and

decoding accuracy in terms of a constant and a linear shape parameter (i.e. 'S parameter estimates')

and show their across-subjects' mean and distribution in violin plots (e.g. Figure 2A row 2). Finally,

we characterized the spatial topography of those S parameter estimates by projecting their group

mean onto a normalized group cortical surface (e.g. Figure 2B).

All statistical results are presented in Table 1, Table 2, Table 3 and Supplementary Table 3 and 4.

Additional descriptive statistics as well as effect sizes can be found here: https://osf.io/tbh37/.

Auditory cortices

Auditory stimuli evoked a positive BOLD response in primary auditory cortex and especially in

anterior portions of the planum temporale. As expected from the typical physiological point spread

function of the GE-EPI BOLD signal, the positive BOLD signal increased roughly linearly towards

the cortical surface (Markuerkiaga et al., 2016) (Supplementary Figure 1 and Supplementary table

1).

Deactivations induced by crossmodal visual stimuli (i.e. a negative BOLD response for [V-Fix])

were observed in both A1 and PT with a constant response profile and based on visual inspection

even a trend towards stronger deactivations in deeper laminae (Figure 2A left, Table 1). These

visually-induced deactivations were generally observed for both auditory and visual attention

conditions, with no significant difference between them ([Att_A-Att_V] for visual stimuli in A1 and

PT: $F(2,40)= 0.644$, $p = 0.530$).

We did not observe a significant crossmodulatory effect of visual stimuli on the BOLD-response in

A1 and PT in the context of concurrent auditory stimuli (Figure 3A left, Table 2A). However,

pattern classifiers succeeded in discriminating between patterns elicited by [AV vs. A] conditions

across all laminae in both PT and A1 (Figure 3A right, Table 2B).Thus, even when the mean BOLD

response across vertices averaged across A1 and PT did not differ significantly for AV and A

stimuli at a particular depth, the visual stimulus changed the activation pattern elicited by a

concurrent auditory stimulus. This provides evidence that sub-regions with crossmodal

enhancement and suppression co-exist in A1 and PT. The visual induced changes in activation

patterns were again not significantly affected by modality-specific attention (for the classification

[AV-V]$_{att A}$ VS [AV-V]$_{att V}$, $F(2,40)= 0.185$, $p = 0.832$).

Taken together these results suggest that salient visual looming stimuli affected auditory cortices

irrespective of the direction of modality-specific attention in both unisensory and audiovisual

contexts. Further, the surface projections of the shape parameters of the BOLD response profile at

the group level revealed a patchy topography both for visually-induced deactivations in a

unisensory context (Figure 2B left) and for visual-induced modulations of the auditory response

(Figure 3B left i and ii).

We next explored whether visual stimuli influenced activation patterns in auditory cortices in a

similar patchy topography during unisensory visual and audiovisual contexts. For this, we defined a

generarl linear model for each subject that used the 'constant' or 'linear' shape parameters from the

visual deactivations (i.e. [V-Fix]) for each vertex to predict the 'constant', and respectively 'linear',

shape parameters for the visually-induced crossmodal modulation (i.e. [AV-A]) over vertices (again

in a cross-validated fashion). We visualized the results of this regression in the form of raster plots

(Figure 3B right i and ii). These raster plots show the laminar BOLD response profile for the

difference [AV-A] across vertices, sorted according to their BOLD response profile for [V-fix].

In A1 and PT, the shape profile of a vertex for visual deactivations significantly predicted its profile

for cross-modal modulation suggesting similar patchy topographies for visual deactivations and

crossmodal modulation. In PT, the constant parameter for [V-Fix] predicted the constant for [AV-

A]: the less a vertex is deactivated in PT by visual stimuli in a unisensory context, the more it

shows a visually-induced enhancement of the response to a concurrent auditory stimulus (constant:

$t(10)=3.460$, P=0.006, linear: $t(10)=1.853$, p=0.094, see Figure 3Bi right). More interestingly, in A1

the linear shape parameter for [V-Fix] predicted the linear shape parameter of [AV-A]: vertices with

less deactivations in deeper relative to superficial laminae in unisensory visual context showed a

robust crossmodal enhancement that was most pronounced in deeper laminae (linear: $t(10)=3.121$,

p=0.011, constant: $t(10)=0.021$, p=0.983, Figure 3Bii right). These results strongly suggest that

visual stimuli generate a BOLD response in primary auditory cortex with a patchy topography

similar for unisensory and audiovisual contexts.

Because these commonalities in topography across two contrasts could in principle arise from

spurious factors (such as registration errors, curvature-dependent segmentation errors,

heterogeneous occurrence of principal veins, curvature dependent occurrence of veins, orientation

dependence to B0, signal leakage of kissing gyri) that could affect BOLD-response magnitude

similarly in different contrasts, we repeated the statistical analysis and raster plots for the constant

shape parameters of the [V-Fix] and [A-Fix] contrasts in auditory areas. However, as shown in the

raster plots in Supplementary Figure 7 this control analysis did not reveal any significant

relationship between the two contrasts. The absence of a significant effect in this control analysis

thus suggests that the similarity in topography between visually induced deactivations and cross-

modal modulations reflects similarities in neural organization rather than non-specific effects.

In contrast to these multisensory influences, attentional modulation was greatest at the cortical

surface in both A1 and PT for the regional BOLD response (i.e. significant positive linear effect in

A1 and PT, Figure 4Ai left, Table 3A). Averaged across A1 and PT, pattern classifiers also

discriminated between the auditory and visual attention conditions (Figure 4Ai right, Table 3B).

The decoding accuracy profiles were characterized by a significant constant term and a non-

significant trend for the linear term. Surprisingly, even though the mean BOLD-response profiles

revealed a profound effect of attention, the discrimination between auditory and visual attention

conditions was rather limited. Discriminating between activation patterns may be limited, if

modality-specific attention predominantly amplifies and scales the BOLD response for A, V or AV

stimuli whilst preserving the activation patterns, whereas the crossmodal influences impacts the

activation patterns.

To investigate this explanation further, we quantified and compared the similarity in A1 between

activation patterns of auditory vs. visual attention conditions (averaging over i. A_Att_V vs. A_Att_A;

and ii. AV_Att_V vs. AV_Att_A) and of auditory vs. audiovisual conditions (averaging over iii.

A_Att_A vs. AV_Att_A; and iv. A_Att_V vs. AV_Att_V) using the Spearman correlation coefficient

computed over vertices. The average similarity between auditory and visual attention conditions

was indeed greater than the average similarity between A and AV conditions (two-sided exact sign

permutation test on Fisher transformed correlation coefficients, $p = 0.008$). These results suggest

that auditory attention mainly scales the BOLD response in A1 whilst preserving the activation

pattern, whereas visual stimuli alter the activation pattern in A1.

In summary, we observed distinct laminar BOLD response profiles and activation patterns for

multisensory and attentional influences in auditory cortices. Visual stimuli induced deactivations

with a constant profile both in A1 and PT. Likewise, they influenced the activation pattern in

auditory cortices similarly across all laminae (i.e. constant effect for decoding accuracy) in

audiovisual context leading to successful pattern decoding accuracy for [AV vs. A] even in deeper

laminae. In fact, in A1 and PT visual inputs altered the activation pattern with a similar patchy

topography when presented alone (i.e. visual-induced deactivation) or together with an auditory

stimulus (i.e. crossmodal modulation).

Modality-specific attention amplified the mean BOLD-signal within A1 and PT mainly at the

cortical surface (i.e. significant linear parameter). Likewise the decoding accuracy was better than

chance with a non-significant increase towards the surface.

Visual cortices

In V1 and V2/3 visual stimuli induced activations that increased towards the cortical surface as

expected for GE-EPI BOLD (Markuerkiaga et al., 2016) (Supplementary Figure 1 and

Supplementary table 1). Contrary to previous research (Chen et al., 2013; Koopmans et al., 2010)

we did not observe a more selective increase in activations for layer 4. Potentially, this selective

increase in BOLD-response may have been smoothed across multiple layers, because the relative

cortical depth of layer four varies between foveal and lateral projections in V1 and our region of

interest is larger than those used in previous work (for similar findings see figure 6B in (Polimeni et

al., 2010) for fMRI and Figure 6E in (Waehnert et al., 2016) for structural anatomy). Consistent

with previous research (Shmuel et al., 2002; Silver et al., 2008; Tootell et al., 1998), the centrally

presented visual looming stimuli activated predominantly areas representing the centre of the visual

field with response suppressions (i.e. negative BOLD) in adjacent areas (see dashed white lines in

Supplementary Figure 3B right and also Supplementary Figure 5 left). By contrast, auditory

deactivations were not confined to the central field that was activated by visual stimuli. Instead,

they were particularly pronounced in the rostral part of the calcarine sulcus representing more

eccentric positions of the visual field (Figure 2B right and Supplementary Figure 5 left right).

Furthermore, auditory looming stimuli induced deactivations that increased in absolute magnitude

towards the cortical surface (significant constant and linear effect, Table 1). Again as in auditory

areas, these auditory deactivations were not significantly modulated by modality-specific attention

([Att$_V$-Att$_A$] for auditory stimuli in V1 and V2/3: $F_{(2,40)}= 1.651$, $p = 0.205$).

Despite the pronounced deactivations elicited by unisensory auditory stimuli, we did not observe

any significant cross-modal modulation of the regional BOLD response in V1 and V2/3 (see

Supplementary Figure 4 left and Supplementary Table 4A). Likewise, the support vector classifiers

were not able to discriminate between activation patterns for [AV vs. V] stimuli better than chance.

In summary, the effect of auditory stimuli on visual cortices was abolished under concurrent visual

stimulation both in terms of regional BOLD response and activation patterns (see Supplementary

Figure 4 right and Supplementary Table 4B).

With respect to attentional modulation, we observed no significant differences in mean BOLD

responses in V1 or V2/3 for visual vs. auditory attention conditions (see Table 3A). As shown in the

violin plots of Figure 4Aii left, only few participants showed a substantial attentional modulation

effect when averaged across the entire regions of interest. Yet, the classifier was able to

discriminate between patterns for visual and auditory attention conditions successfully across all

cortical depth surfaces in V1 and V2/3 (i.e. significant constant term in V1 and V2/3 and linear term

in V2/3, see Table 3B, Figure 4Aii right). The surface projection of shape parameters of the BOLD

response profile at the group level explains this discrepancy between the attentional effects on

regional BOLD response and activation patterns (Figure 4Bii). Consistent with mechanisms of

contrast enhancement (Bressler et al., 2013; Müller and Kleinschmidt, 2004; Smith et al., 2000;

Tootell et al., 1998), attending to vision relative to audition increased responses in regions that were

activated by visual looming stimuli, yet suppressed the deactivations in the surrounding regions that

were deactivated by visual stimuli thereby cancelling out global attentional effects for the regional

BOLD response (Figure 4Bii, note the white lines indicate the border between visual activation and

deactivation from Supplementary Figure 3B right).

In summary, auditory looming induced widespread deactivations in visual cortices that were

maximal at the cortical surface and extended into regions that represent more peripheral visual

fields. By contrast, attention to vision predominantly increased BOLD response in central parts that

were activated by visual stimuli and suppressed BOLD responses in the periphery that were

deactivated   by   visual   stimuli.

DISCUSSION

The current high-resolution 7T fMRI study revealed distinct depth-dependent BOLD response profiles and patterns for multisensory and attentional influences in early sensory cortices.

Distinct laminar profiles and activation patterns for multisensory and attentional influences in auditory cortices

Visual looming suppressed activations in auditory cortices (see also (Laurienti et al., 2002; Leitão et al., 2013; Mozolic et al., 2008), but enhanced the response to concurrent auditory looming in posterior auditory cortices (Werner and Noppeney, 2011, 2010). In other words, intersensory competition for purely visual stimuli turned into cooperative interactions for audiovisual stimuli. While the visual-induced response suppression was constant across cortical depth in A1 and PT, the crossmodal BOLD-response enhancement was observed mainly in the caudal parts of PT, i.e. in parts of PT where visual influences have previously been shown (Kayser et al., 2007). At a finer spatial resolution, multivariate analyses revealed significant differences between audiovisual and auditory activation patterns across all laminae in both A1 and PT. As shown in the surface projections (see Figure 3B), visual looming enhanced and suppressed auditory responses in adjacent patches consistent with neurophysiological results in non-human primates (Kayser et al., 2008). Critically, as illustrated in the raster plots, the response profile of a vertex to visual stimuli significantly predicted its laminar profile for crossmodal modulation (see Figure 3B right). In A1, vertices whose activations were only weakly suppressed by visual looming in deeper laminae showed a greater visual-induced response enhancement for audiovisual looming again in deeper laminae. These results suggest that visual looming influences activations in A1 mainly in deeper layers with similar patchy topographies in unisensory and audiovisual contexts.

By contrast, modality-specific attention affected auditory and visual evoked responses predominantly at the cortical surface (see significant linear term in Table 3). The large attentional BOLD-response effects in superficial laminae dovetail nicely with previous neurophysiological

research, which located effects of modality-specific attention in supragranular layers of auditory cortices (De Martino et al., 2015a; Lakatos et al., 2009). Yet, because the GE-EPI BOLD response measured at superficial laminae includes contributions from deeper laminae (Harel et al., 2006; Markuerkiaga et al., 2016), we cannot unequivocally attribute the attentional influences at the cortical surface solely to superficial layers as neural origin. It is also possible that the increase in attentional BOLD-response effects towards the cortical surface may arise from neural effects across all cortical layers. Consistent with this conjecture, the BOLD-response profiles to auditory induced activations (Supplementary Figure 2A left) and attentional modulation (Figure 4Ai left) in A1 and PT are quite similar.

Critically, however, the laminar profiles for attentional effects were distinct from those for multisensory (i.e. crossmodal) effects in auditory cortices. This dissociation in laminar profiles in the same territories cannot be explained by vascular effects that limit the laminar specificity of the BOLD response, but strongly implies that multisensory and attentional mechanisms regulate information flow in auditory cortices via partly distinct neural circuitries (De Martino et al., 2015a; Lakatos et al., 2009). Further, the constant laminar profiles for crossmodal influences in auditory cortices suggest that visual signals influence auditory cortices mainly via infragranular layers – a finding that converges with a recent neurophysiological study in mouse that likewise highlighted deep layer 6 as the key locus for visual influences on auditory cortex (Morrill and Hasenstaub, 2018). Anatomical studies in monkeys have previously suggested that visual inputs can influence auditory cortices via three routes (Ghazanfar and Schroeder, 2006; Musacchia and Schroeder, 2009; Schroeder et al., 2003; Smiley and Falchier, 2009): i. thalamic afferents, ii. direct connectivity between sensory areas (Falchier et al., 2002; Rockland and Ojima, 2003) and iii. connections from higher order association cortices.

Common laminar profiles, but distinct activation patterns for multisensory and attentional influences in visual cortices

In visual cortices, visual and auditory stimuli evoked responses that were maximal in absolute

amplitude at the cortical surface, yet differed in their activation pattern. Visual looming stimuli

induced activations in areas representing the centre of the visual field and deactivations in

surrounding areas representing the periphery (Shmuel et al., 2002). In contrast to this visual 'centre-

surround' pattern, auditory stimuli induced widespread deactivations (Laurienti et al., 2002; Leitão

et al., 2013; Mozolic et al., 2008) particularly in peripheral visual field representations known to

have denser direct fibre connections to auditory cortices (Falchier et al., 2002; Rockland and Ojima,

2003).

Thus, our study revealed three types of deactivations. In visual cortices, both visual and auditory

stimuli induced deactivations that were most evident at the cortical surface, as previously reported

for deactivations to ipsilateral visual stimuli (Fracasso et al., 2018), though see (Goense et al.,

2012). In auditory cortices, the visual induced deactivations were constant across cortical depth.

These differences in laminar profiles or BOLD patterns between visual and auditory cortices may

reflect differences in neural mechanisms. For instance, neurophysiological studies in rodents have

shown asymmetries in multisensory influences between auditory and visual cortices, reporting

robust auditory induced inhibition in layers 2/3 in visual cortices but not visual induced inhibition in

auditory cortices (Iurilli et al., 2012).

While we observed no statistically significant crossmodal nor attentional effects on the regional

BOLD response profiles in visual cortices, multivariate pattern analyses indicated that modality-

specific attention altered the activation patterns at a sub-regional spatial resolution (Figure 4Aii):

attention to vision amplified activations in areas representing the central visual field and suppressed

activations in areas representing peripheral visual fields (see Figure 4Bii).

The differences in activation patterns for multisensory and attentional influences indicate that the

impact of purely auditory looming (Maier et al., 2008, 2004) on visual cortices cannot solely be

attributed to a withdrawal of attentional resources from vision. Instead, purely auditory looming

may inhibit activations in visual cortices via direct connectivity between auditory and visual areas

(Bizley et al., 2007; Budinger et al., 2006; Campi et al., 2010; Falchier et al., 2002; Ibrahim et al.,

2016; Rockland and Ojima, 2003). In line with this conjecture, research in rodents has shown that

auditory stimuli modulates activity in supragranular layers in primary visual areas via direct

connectivity from auditory cortices and translaminar inhibitory circuits – though the studies

disagreed in whether layer 5 (Iurilli et al., 2012) or layer 1 (Deneux et al., 2019; Ibrahim et al.,

2016) were the primary targets of auditory inputs.

Multisensory effects in visual and auditory cortices – a comparison

Our results reveal distinct laminar profiles for multisensory deactivations in auditory and visual

cortices. In auditory cortices the visual induced deactivations were constant across laminae, in

visual cortices the deactivations linear increased (in absolute magnitude) towards cortical surface.

These distinct laminar profiles converge with previous research in rodents that also reveal visual

influences in auditory cortices predominantly in infragranular layer (Morrill and Hasenstaub, 2018),

but auditory influences in visual cortices mainly in supragranular layers (Deneux et al., 2019;

Ibrahim et al., 2016; Iurilli et al., 2012). Potentially, these distinct laminar profiles in auditory and

visual cortices may suggest that multisensory influences are mediated by different neural circuitries

in auditory and visual cortices. However, the distinct laminar profiles across cortical areas may not

only reflect differences in neural circuitries but also in vascular organization. Moreover, a wealth of

research has previously shown that multisensory responses profoundly depend on stimulus salience

and other input characteristics (Deneux et al., 2019; Meijer et al., 2017; Stein et al., 2019; Werner

and Noppeney, 2010). It is currently unclear how these factors influence laminar BOLD-response

profiles. Finally, interpreting laminar BOLD-response profiles in relation to previous findings in

rodents remains tentative because of cross-species and methodological differences.

CONCLUSIONS

Using submillimetre resolution fMRI at 7T, we resolved BOLD response and activation patterns of

multisensory and attentional influences across cortical depth in early sensory cortices. In visual

cortices auditory stimulation induced widespread inhibition, while attention to vision enhanced

central but suppressed peripheral field representations. In auditory cortices competitive and

cooperative multisensory interactions were stronger deep within the cortex, while attentional

influences were greatest at the cortical surface. The distinctiveness of these depth-dependent

activation profiles and patterns suggests that multisensory and attentional mechanisms control

information flow via partly distinct neural circuitries.

MATERIAL AND METHODS

All procedures were approved by the Ethics Committee of the University of Leipzig.

Participants

Thirteen healthy German native speakers (6 females; 7 males; mean age: 24.8 years, standard deviation: 1.5 years, range: 22-27 years) gave written informed consent to participate in this fMRI study. Participants reported no history of psychiatric or neurological disorders, and no current use of any psychoactive medications. All had normal or corrected to normal vision and reported normal hearing. Two of those subjects did not complete all fMRI sessions and were therefore not included in the analysis.

Looming stimuli and targets

The visual looming stimulus (500 ms duration) was a sequence of 1 to 4 radially expanding white annuli (Figure 1B). Each of the four annuli started in the centre of the screen with a diameter of 0.25º visual angle and expanded at a radial speed of 15.52º visual angle per second to a maximum radius of 3.88º visual angle which covered the entire screen visible in the scanner bore along the vertical dimension. During the 500 ms sequence, an additional annulus started in the centre every 125 ms. Given the retinotopic organization of early visual cortices (Wandell et al., 2007) these parameters ensured that visual looming elicited relatively widespread activations in visual cortices.

The auditory looming stimulus (500 ms duration) was composed of several pure tones (55, 110, 150, 220, 330, 380, 440, 660, 880, 990, 1110, 1500, 2500, 3000 Hz) that increased in frequency and amplitude over the entire 500 ms duration. The amplitude of the sound was exponentially modulated with the amplitude at time t given by $A_t = A_0 \, e^{0.68t}\text{-}1$ and A_0 being the initial baseline amplitude for each pure tone. The frequency of each pure tone component was increased by two thirds of its original base frequency over the 500 ms. Given the tonotopic organization of primary

auditory cortices (Formisano et al., 2003) this mixture of sound frequencies ensured widespread

activations in auditory cortices.

Audiovisual stimuli were generated by presenting the auditory and visual stimuli in synchrony

together.

The visual target was a grey dot presented for 50 ms within a circular area that increased over the

500 ms duration consistent with the auditory, visual or audiovisual looming stimulus (Figure 1B).

Auditory targets were a single 440 Hz pure tone that lasted for 100 ms. The size of the visual target

and the sound amplitude of the auditory target were adjusted in a subject-specific fashion to match

the approximate target detection performance across sensory modalities and then held constant

across the entire experiment. This was done using the method of constant stimuli in a brief

'psychometric function experiment', performed prior to the fMRI study inside the scanner and with

scanner noise present.

Experimental design

The experiment conformed to a 3 (stimulus modality: auditory, visual, audiovisual) X 2 (attended

modality: auditory, visual) factorial design (Figure 1A). In an intersensory selective attention

paradigm, participants were presented with blocks of audio (A), visual (V) or audiovisual (AV)

looming stimuli as highly salient stimuli that drive bottom-up attention and elicit crossmodal

deactivations (Leitão et al., 2013). Brief auditory (i.e. weak and brief beep) and visual (i.e. small

and brief grey dot) targets were presented interspersed in all stimulation blocks irrespective of the

sensory modality of the looming stimuli or the focus of modality-specific attention (e.g. A and V

targets were presented during a unisensory auditory looming block and auditory attention). The

visual grey dot could be presented at any time at any location within the area defined by the radially

expanding outermost annulus until they reach a radius of maximal 3.88º visual angle. Likewise, the

auditory target is presented at random time interspersed in the looming sound. Consistent with

previous research on modality-specific attention (Lakatos et al., 2009; Werner and Noppeney, 2011,

2010) these additional targets were included to ensure that the effects of attention could not be

attributed to top-down effects associated with explicit responses.

During auditory attention (Att$_A$), participants were instructed to respond selectively to all auditory

targets and ignore visual ones. During visual attention (Att$_V$) participants were instructed to respond

selectively to all visual targets, i.e. brief grey dots and ignore auditory targets. In this way, we

matched the auditory (i.e. attention to frequency) and visual attention (i.e. attention to targets in

spatial field) tasks to the tonotopic and retinotopic organizational principles of primary auditory and

visual areas as well as the specifics of auditory and visual looming bottom-up salient stimuli.

Experimental procedures

Auditory, visual or audiovisual looming stimuli were presented in 33 seconds blocks of 50 stimuli

(stimulus duration: 500 ms, inter-stimuli interval: 160 ms giving a fixed stimulus onset asynchrony

of 660 ms with no jitter) (Figure 1B). Activation blocks were preceded by 16 seconds fixation

baseline periods. We opted for a block design as they have the highest design efficiency despite

their potential drawbacks from a cognitive perspective (Friston et al., 1999). Throughout the entire

experiment, participants fixated a cross (0.16º visual angle) in the centre of the black screen.

Therefore, processes related to fixation are present in all conditions (i.e. all stimulation and fixation

conditions). The fixation cross was white during the activation conditions and blue during the

fixation baseline periods. A blue letter ('V' or 'A') presented 3 seconds prior to the activation

blocks instructed participants to direct their attention to the visual or auditory modality. During

visual attention blocks, participants were instructed to attend generally to the visual modality and

respond selectively to visual targets. Conversely, during auditory attention blocks, participants had

to attend generally to the auditory modality and respond selectively to auditory targets. Sixteen

targets of each modality were presented for each condition. To maintain observer's attention and

minimize the effect of target stimuli on measured stimulus-induced activations, targets were more

frequently presented at the end of the block (i.e. two thirds of the blocks included one auditory

target and one visual target, the last third of blocks included 2 targets of each type). The final block

of each run was followed by a fixation baseline period of 20 seconds. The order of blocks was

pseudo randomized across participants. Each fMRI run included 18 blocks (3 blocks for each

condition in our 2 X 3 design) and lasted 15 minutes. There were 4 runs per participant yielding 12

blocks per condition per participant.

Experimental set up

Stimuli were presented using Presentation (v17.6, Neurobehavioral system, www.neurobs.com) on

a PC desktop (Windows XP). Visual stimuli were back-projected onto a plexiglas screen using an

LCD projector (60 Hz) visible to the participant through a mirror mounted on the MR head coil.

The MR head coil enabled a visual field with a width of 10.35° and height of 7.76° visual angle.

Auditory stimuli were presented using MR-compatible headphones (MR-Confon; HP-SI01 adapted

for the head-coil). Behavioral responses were collected using a MR-compatible button device

connected to the stimulus computer. The code to run the fMRI experiment is available at

https://doi.org/10.5281/zenodo.3581316.

Behavioral data analysis

The response window extended from the target onset and for either 2.5 seconds or until the

beginning of the next target. Any recorded response outside of that window was coded as an extra-

response (i.e. to be modelled in the fMRI design matrix). The percentage of auditory (or visual)

targets that participants responded to in a particular condition were entered as dependent variables

into a 2 (target modality: auditory vs. visual) X 2 (attended modality: auditory vs. visual) X 3

(stimulus block modality: auditory, visual, audio-visual) repeated measures ANOVA (results were

corrected for non-sphericity using Greenhouse-Geiser correction). Alpha was set to 0.05. This

analysis was done with IBM SPSS Statistics, version 17.0 (IBM, Armonk, NY, USA)

MRI Data acquisition

All experiments were performed on a Siemens 7T MAGNETOM MRI scanner (Siemens Healthineers, Erlangen, Germany) equipped with an SC72 gradient coil (Siemens Healthineers). For radio-frequency transmission and reception, a single-channel-transmit/32-channel-receive phased array head coil (Nova Medical, Wilmington, MA, USA) was used. The experiment was run over 2 days with 50 minutes MRI scanning in total per day per participant.

On the first day a whole-brain MP2RAGE sequence (Marques et al., 2010) was used to acquire a quantitative T1 map and a weighted T1 image and a second inversion image to enable high-resolution segmentation and cortical laminae definition [repetition time (TR) / echo time (TE) = 5000/2.45 ms, inversion time 1 = 900 ms, flip angle 1 = 5°, inversion time 2 = 2750 ms, flip angle 2 = 3°, 240 slices, matrix = 320×320, acquisition time = 600 s, spatial resolution = 0.7×0.7×0.7 mm^3 voxels, partial Fourier factor = 6/8, parallel imaging using GRAPPA with an acceleration factor of 4, field of view = 168 X 224 mm]. On the second day, we used a shorter version of that sequence with a coarser resolution only in order to position our EPI acquisition slab.

Two functional runs were acquired every day using a 2D gradient echo single-shot echo planar imaging (EPI) readout [TR/TE = 3000/25 ms, flip angle = 90º, 48 axial slices acquired in descending direction, matrix size = 256×240, phase encoding along the second dimension (posterior to anterior), slice thickness = 0.75 mm, pixel bandwidth: 1085 Hz, brain coverage = 36 mm, spatial resolution = 0.75×0.75×0.75 mm^3 voxels, partial Fourier factor = 6/8, parallel imaging using GRAPPA with an acceleration factor of 4, number of volumes = 302, acquisition time = 906 s, field of view = 192 X 180 mm. In order to avoid spin history effects due to imperfect pulse profiles a slice gap of 0.05 mm was introduced. For every session, a new and optimum set of shim values was used.]. A corresponding vendor-provided homodyne online image reconstruction algorithm was used. The first six volumes in each run were automatically discarded to avoid for T1 saturation effects.

The acquisition slab was inclined and positioned to include Heschl's gyri, the posterior portions of

superior temporal gyri and the calcarine sulci. Both hemispheres were acquired. On the second day

the slab was auto-aligned on the acquisition slab of the first day. This provided us with nearly full

coverage over all our four regions of interest (see Supplementary Table 2).

One field-map was acquired on each day with a coverage and orientation matching that of the EPI

slab [TR/TE1/TE2 = 1500/6/7.02 ms, flip angle = 72º; 18 axial slices, matrix = 96 X 90, slice

thickness = 2 mm, spatial resolution = 2.00 X 2.00 X 2.00 mm^3 voxels, field of view: FOV= 192 X

180 mm].

MRI structural analysis

The MRI structural data were processed using the CBS toolbox (v3.0.8, Max Planck Institute for

human cognitive and brain science, Leipzig, Germany, www.nitrc.org/projects/cbs-tools/) based on

the MIPAV software package (v7.0.1, NIH, Bethesda, USA, www.mipav.cit.nih.gov, (McAuliffe et

al., 2001)) and the JIST pipeline environment(v2.0, (Landman et al., 2013; Lucas et al., 2010)). The

T1 map was coregistered to the Montreal's Neurological Institute brain space (rigid body

registration and normalized mutual information as a cost function) and resampled to an isometric

resolution of 0.4 mm. The image was segmented fully automatically, and the cortical boundary

surfaces were reconstructed (Bazin et al., 2014). Between these surfaces, we computed 6

intracortical surfaces that defined cortical laminae with the equivolume approach (Waehnert et al.,

2014) (Figure 1C and Supplementary Figure 2). It is important to note that the term 'lamina' used in

this communication does not directly refer to cytoarchitectonically defined cortical layers: for

example, the variation across cortical regions of the relative thickness between layers was not

modelled here (Nieuwenhuys, 2013). However, it should be mentioned that this method provides

biologically plausible cortical contours (Waehnert et al., 2014).

Definition of Region of Interest

We defined four regions of interest (ROI), each combining left and right hemispheres, as follows

(see Table S2 for ROI effective sizes):

A) Several studies have shown that the core of primary auditory cortex (A1) because of its

characteristic myelination profile can be localized using its low T1 intensity (De Martino et

al., 2015b; Dick et al., 2012; Hackett et al., 2001). We therefore defined it by manual

delineation (RG, UN) that was guided by anatomical structure (Heschl's gyrus) and

informed by the quantitative T1 map that was sampled at mid-depth of the cortex and

projected on the corresponding inflated surface. The final bilateral A1 region was defined as

the union of the two delineations (RG, UN). As a result the sub-region of Heschl's gyrus

that we refer to as primary auditory cortex may differ from primary auditory cortex defined

based on cytoarchitecture.

B) The planum temporale (PT) was defined bilaterally using the area 41-42 of the brainnetome

atlas (Fan et al., 2016) (http://atlas.brainnetome.org/). The full probability maps of this ROI

and its anatomical neighbours were inverse-normalized into native space using the

deformation field given by the SPM segmentation of the T1 weighted structural scan. The

probability map was then re-gridded to an isometric resolution of 0.4 mm, sampled at mid-

depth of the cortex and projected on the corresponding surface. Vertices were included as

part of PT if their probability exceeded 40% unless i. they were already defined as being

part of A1 or ii. the sum of the probabilities of the neighbouring regions exceeded that of the

area 41-42.

C) Primary Visual area (V1) and high order visual areas (V2 and V3) were defined bilaterally

in a similar fashion as PT. We used the probabilistic retinotopic maps of the ROIs for V1,

V2/3 (Wang et al., 2015) (http://scholar.princeton.edu/napl/resources). In this case vertices

were included if their probability exceeded 10% unless the sum of the probabilities of the

neighbouring regions exceeded that of the area of interest.

We obtained > 95% coverage for A1, PT and V1 and > 82% coverage for V2/3. (For the number of

vertices and percentage coverage across subjects, Supplementary Table 2).

fMRI preprocessing

The fMRI data were pre-processed and analyzed using statistical parametric mapping (SPM12 –

v6685; Wellcome Center for Neuroimaging, London, UK; www.fil.ion.ucl.ac.uk/spm) running on

matlab (Mathworks). The fieldmaps were co-registered to the first functional scan of the first run

(i.e. rigid-body transformations optimized using normalized mutual information as cost function)

and a voxel displacement map was then created. Functional scans from each participant were

realigned and unwarped using the first scan as a reference (interpolation: 4^{th} degree b-spline,

unwarping done using the voxel displacement map of the corresponding day for each run).

Functional to anatomical coregistration

We co-registered the mean EPI image to the pre-processed T1 map (i.e. rigid-body transformations

optimized using normalised mutual information as cost function). We assessed the accuracy of co-

registration using FSLview (FSL 5.0 - Analysis Group, FMRIB, Oxford, UK,

https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSL) by flipping back and forth between the mean EPI and the

T1 map images in all the ROIs (see Supplementary Figure 2). If the co-registration was not

sufficiently precise, the mean EPI was initially co-registered manually and the co-registration

repeated until anatomical structures of the mean EPI and the T1 map were precisely realigned. The

transformation matrix of the co-registration was later applied to the beta images from the fMRI

general linear model (see).

fMRI statistical analysis

At the first (i.e subject) level, we performed a mass univariate analysis with a linear regression at

each voxel, using generalized least squares with a global approximate AR(1) autocorrelation model

and a drift fit with discrete cosine transform basis (128 s cut-off). We modelled the fMRI

experiment with a mixed block-event related model with regressors entered into the run-specific

design matrix after convolving the onsets of each block or event with a canonical hemodynamic

response function (HRF) and its temporal derivative. More specifically, we modelled each of the

three looming blocks in each run for each of the 6 conditions in our 3 (visual, auditory, audiovisual)

X 2 (auditory vs. visual attention) factorial design separately. Hence for each run the statistical

model included the following regressors: 3 block regressors for each of the 6 conditions (A_Att$_A$,

A_Att$_V$, V_Att$_A$, V_Att$_V$, AV_Att$_A$, AV_Att$_V$), 4 event-related regressors for each target type under

each attention condition (i.e. TargetA_Att$_A$, TargetA_Att$_V$, TargetV_Att$_A$, TargetV_Att$_v$) and one

event-related regressor modelling any additional responses that participants produced in the absence

of targets. We modelled the targets and additional responses as independent regressors in our first

(i.e. subject) level general linear model (GLM) to minimize confounding effects of perceptual

decision making, response selection and motor preparation on the activations reported for the A, V,

AV looming activation blocks that are the focus of this communication. Nuisance covariates

included the realignment parameters to account for residual motion artefacts.

All subsequent analyses included only the beta images pertaining to the HRF of the activation

blocks in our 2 (modality specific attention) X 3 (stimulation modality) design: 3 blocks per

condition per run X 6 conditions X 4 runs = 72 beta images per subject.

To minimize the possibility that attentional lapses reduced the sensitivity of our analysis we also

repeated our neuroimaging analysis selectively for blocks that included neither misses (i.e. missed

responses to targets of the attended modality) nor false alarms (i.e. responses to targets from the

unattended modality). This control analysis basically showed similar results similar to those from

the main analysis that is reported in this manuscript.

Sampling of the BOLD-response along cortical depth

The 72 beta images (6 conditions X 3 blocks per condition per run X 4 runs) were co-registered to

the MRI structural by applying the transformation matrix obtained from the mean EPI image co-

registration and up-sampled to a 0.4 mm isometric resolution (4th degree b-spline interpolation).

Finally, we sampled each beta image along vertices defined by the normal to the mid-cortical

surface at the depths defined by the 6 intra-cortical surfaces from the MRI structural analysis (tri-

linear interpolation) (see 'MRI structural analysis' above, Figure 1C).

In total the fMRI data were therefore resampled 3 times for: 1. realignment+unwarping, 2.

upsampling the beta images to 0.4 mm and 3. sampling the beta images along the surfaces. The

smoothness estimation using AFNI 3dFWHMx gave the following results (FWHM mean and

standard deviation across subjects in X/Y/Z): smoothness of the raw data: 1.70 (0.08), 2.05 (0.09),

1.65 (0.08) mm; smoothness after upsampling at 0.4 mm: 2.22 (0.14), 2.88 (0.31), 2.11 (0.21) mm.

We were not able to estimate the smoothness of the data after sampling along the surface*s*. Note that

no additional smoothing was applied beyond the one due to interpolation during pre-processing.

The activity values at the vertices of these 6 intra-cortical surfaces in our four ROIs (i.e. primary

auditory, planum temporale, V1, V2/3) pooled over both hemispheres form the basis for our

univariate ROI analysis of the laminar BOLD-response profiles and laminar decoding accuracy

profiles (based on multivariate pattern classification).

Laminar BOLD-response profiles for contrasts of interest

The laminar profiles of each ROI were obtained for each block per run by collapsing data over

vertices. For each of our 72 beta images (6 conditions X 3 blocks per condition per run X 4 runs)

we summarized the BOLD response at each of the six cortical depth levels in terms of the median of

the parameter estimates across all vertices at this cortical depth within a particular ROI (n.b. each

ROI pools over both hemispheres). The median was used as a summary index because the

parameter estimates distribution was skewed especially for the most superficial laminae.

For each ROI and lamina, we computed the following contrasts of interest independently for the i^{th}

block of a given condition of the j^{th} run:

1. the deactivation induced

a. by auditory stimuli relative to fixation baseline irrespective of modality-specific

658   attention:

$$[\text{A-Fix}] = [\text{A\_Att}_A + \text{A\_Att}_V]_{i,j} / 2,$$

b. by visual stimuli relative to fixation baseline irrespective of modality-specific attention:

$$[\text{V-Fix}] = [\text{V\_Att}_A + \text{V\_Att}_V]_{i,j} / 2,$$

2. the crossmodal enhancement or suppression irrespective of attention, i.e. specifically for

auditory and visual regions:

a. in A1 and PT: the visual-induced modulation of auditory activations irrespective of

665   modality-specific attention:

666    $$[\text{AV - A}] = [(\text{AV\_Att}_A - \text{A\_Att}_A) + (\text{AV\_Att}_V - \text{A\_Att}_V)]_{i,j} / 2,$$

667   b. In V1 and V2/3: the auditory-induced modulation of visual activations irrespective of

668   modality-specific attention:

$$[\text{AV - V}] = [(\text{AV\_Att}_A - \text{V\_Att}_A) + (\text{AV\_Att}_V - \text{V\_Att}_V)]_{i,j} / 2,$$

3. attentional modulation irrespective of stimulus modality, i.e. specifically for auditory and visual

regions

a. A1 and PT: Modulation of stimulus responses by auditory relative to visual attention

$$[\text{Att}_A - \text{Att}_V]_{A,V,AV} = [(\text{Att}_A - \text{Att}_V)_A + (\text{Att}_A - \text{Att}_V)_V + (\text{Att}_A - \text{Att}_V)_{AV}]_{i,j} / 3.$$

b. V1 and V2/3: Modulation of stimulus responses by visual relative to auditory attention

$$[\text{Att}_V - \text{Att}_A]_{A,V,AV} = [(\text{Att}_V - \text{Att}_A)_A + (\text{Att}_V - \text{Att}_A)_V + (\text{Att}_V - \text{Att}_A)_{AV}]_{i,j} / 3$$

For completeness, we also assessed whether the differences in BOLD-response for the contrasts

listed 1 and 2 – in cases when they were significantly different from zero - depended on modality-

specific attention (i.e. by testing for the interaction of sensory evoked responses or crossmodal

enhancement with modality-specific attention). Moreover, we also show the activations > fixation

for auditory or visual stimuli in the Supplementary Figure 3 to confirm that our experimental

manipulations were effective.

Shape parameters for characterization of laminar BOLD response profiles

The laminar profile of each ROI was then summarized by parameters by collapsing across blocks

per runs. For each statistical comparison, lamina and ROI, we thus obtained 12 (i.e. 3 blocks X 4

runs) contrast estimates (as the median over vertices within a bilateral ROI) per subject i.e. 12

laminar profiles. We refer to those parameter estimates that quantify the BOLD response for a

particular lamina as BOLD response parameter estimates. For instance, Figure 2A (first row) shows

the. B parameter estimates (across participants mean $\pm$ SEM) as a line plot for each of the six

laminae as a function of cortical depth along the x-axis.

To characterize the overall shape of the laminar profile, we estimated for each subject a 2^{nd} level

general linear model (i.e. a laminar GLM) that modelled the activation in each lamina in a given

ROI as dependent variable by two predictors: 1. a constant term (i.e. activation mean across

laminae) and 2. a linear term characterising a linear increase across laminae (mean-centered and

orthogonalized with respect to the constant term). The parameter estimates of this 2^{nd} or laminar

GLM are referred to as shape parameters. For instance, Figure 2A (2^{nd} row) shows the S parameter

estimates as violin plots for the constant and the linear term. The S-parameters enable us to make

inferences about the shape of a laminar BOLD-response profile rather than using an omnibus F-test

that assesses whether the BOLD-response differs across any of the six laminae followed by

numerous post hoc pairwise comparisons between layers. However, we acknowledge that this

characterization makes our inference less specific about locating even the BOLD-response to a

particular lamina. Moreover, because the laminar GLM in the current report did not include any

higher order (e.g quadratic) terms they would not be to model U-shaped laminar profiles.

Statistical analysis for the shape parameters at the between subject i.e. group level

To enable generalization to the population, we entered these shape parameters into linear mixed

effects models at the group level separately for each contrast. To limit the number of statistical tests

we used the following step-down approach:

1. We formed 2 (shape parameter: constant, linear) x 2 (ROI: primary, non-primary) linear mixed

effects models separately for the six contrasts, i.e. 2 (sensory cortices: visual, auditory) x 3

(contrasts: crossmodal deactivation, crossmodal modulation, attentional modulation). In each linear

mixed effects model we tested whether the constant or the linear parameter, each averaged across

primary and higher order ROIs, was significantly different from zero using a two dimensional F-

contrast. Hence, we computed three F-tests for visual cortices and three F-tests for auditory cortices.

2. If a two dimensional F-test was significant, we computed follow-up one dimensional F-tests

separately for the constant and the linear parameters (again averaged across primary and higher

order sensory cortices).

3. If a one dimensional F-test was significant, we computed follow-up t-tests separately for the

primary and the higher order sensory cortices.

Based on our a-priori predictions (Leitão et al., 2013) that auditory stimuli induce deactivations in

visual areas and that visual stimuli induce deactivations in auditory areas, we employed one-sided t-

tests in step 2 and 3 (i.e. negative constant term for contrasts). In all other cases unidirectional F-

tests were used. Unless otherwise stated, we report statistical results as significant at $p < 0.05$.

Multivariate analysis of pattern across vertices and decoding accuracy profiles

Multivariate pattern analyses were performed using a linear support vector classifier (SVC) that was

trained in a leave-one-run-out cross-validation scheme (LIBSVM 3.21,

https://www.csie.ntu.edu.tw/~cjlin/libsvm/, with C = 1 and mean-centred activations for each

feature separately for training and test sets).

We performed multivariate pattern analysis directly on the BOLD response patterns pooled over the

two hemispheres, i.e. the patterns of B-parameters, independently at each of the 6 cortical depths to

generate laminar profiles of decoding accuracy for the following comparisons:

1. $[AV \text{ vs } A]_{AttA, AttV}$ (pooled over attended modality) for A1 and PT,

2. $[AV \text{ vs } V]_{AttA, AttV}$ (pooled over attended modality) for V1 and V2-3,

3. $[Att_A \text{ vs } Att_V]_{A,V,AV}$ (pooled over stimulation modality) for A1, PT, V1 and V2/3.

For completeness, we also assessed whether the classification performance for the comparisons

listed above – in cases when they were significantly different from zero - depended on modality-

specific attention by running the comparison for the equivalent contrast between the the 2 modality

specific attention conditions (e.g the comparison $[AV-V]_{att\ A}$ vs $[AV-V]_{att\ V}$ for the 1 in the list

above).

In line with our analysis of laminar BOLD response profiles, we modelled the laminar profiles of

decoding accuracy using a laminar GLM with a constant term and a linear term as linear increase

across laminae as predictors.

Again as in our analysis of laminar BOLD-response profiles, the 'constant' and 'linear' shape

parameters characterizing the decoding accuracy profiles were entered into 2 (shape parameter:

constant, linear) x 2 (ROI: primary, non-primary) linear mixed effects models separately for each of

the four decoding comparisons, i.e. 2 (sensory cortices: visual, auditory) x 2 (contrasts: crossmodal

modulation, attentional modulation). We then applied the step down procedure exactly as described

in detail for the BOLD-response profiles. Please note that the laminar profiles of decoding accuracy

need to be interpreted cautiously. First, decoding accuracy depends on BOLD-signal and noise

characteristics that can both vary across laminae. Second, the laminar profile of decoding accuracy

is more difficult to interpret, because accuracy is bounded between 0 and 1.

Raster plots: statistical relationship of BOLD response profiles between different conditions

Next we explored whether the response profile in a vertex as characterized by their 'constant' and

'linear' shape parameters in one condition is statistically predictive of this vertex laminar profile in

another condition or contrast (see figure 5 from (Fracasso et al., 2018). For instance, we asked

whether the magnitude of visual induced deactivations (e.g. [V-Fix] averaged over attention

conditions) in a vertex predicts its crossmodal enhancement (e.g. [AV-A] averaged over attention

conditions). For this, we sorted and averaged the vertices in percentile-like bins based on the value

of a shape parameter (e.g. 'constant' or 'linear') for a particular sorting contrast (e.g. [V–Fix]).

This bin order was then used to sort and average the values of the 2^{nd} contrast (e.g. [AV-A]). We

entered the shape parameter value (e.g. 'constant' or 'linear') for the sorting contrast (e.g. [V–Fix])

for each bin as a linear regressor (+ a constant) to predict the corresponding shape parameter in each

bin in the sorted contrast (e.g. [AV-A]) in a subject-specific general linear model.

To allow for generalization to the population, we entered the parameter for the linear term (i.e.

slope of regression line) of this regression model into one sample t-tests at the group level.

We illustrated the statistical relationships of the laminar profiles between different contrasts at the

group level in raster plots by averaging the laminar profiles for each bin across subjects of the

sorting (i.e. predicting) and the sorted (i.e. predicted) values. For instance, in Figure 3B i (right) we

sorted the vertices according to [V-fix] in PT such that the constant parameter (i.e. average BOLD

response across laminae) increases from bottom to top (i.e. predicting contrast). Unsurprisingly, the

raster plot for the [V-Fix] panel as the predicting contrast thus goes from a negative BOLD response

profile (i.e. coded in blue) in the bottom rows of the panel to a positive BOLD response profile (i.e.

coded in red) in the top rows of the panel. Under the null-hypothesis the laminar BOLD response

profile for the predicting contrast (e.g. [V-fix]) in a vertex is unrelated to its laminar BOLD

response profile for the predicted contrast (e.g. [AV-A]) and we would not expect any structure in

the raster plots for the predicted contract (e.g. [AV-A], n.b. we accounted for spurious correlation

by performing all these analyses in a crossvalidated fashion). Conversely, if the shape parameter for

[V-fix] in a vertex significantly predicts its shape parameters for [AV-A], we would expect a

structured raster plot also for [AV-A].

Hence, these raster plots can reveal additional BOLD signal structure that is averaged out by the

group surface projections: while group surface projections (see next section, or Figure 3B left)

reveal only spatial topographies which are consistent across participants, the raster plots reveal

whether the similarity between patterns of laminar profiles from different conditions is consistent

across subjects, even when the activation patterns themselves are not similar across subjects. In

other words, raster plots illustrate the similarity or covariance between patterns that is consistent

across subjects even when the patterns themselves vary across subjects.

Please note that both the regression model and the raster plots were computed in a leave one day out

cross-validation to avoid biases and spurious correlations between sorting and sorted contrast. The

number of bins was set to the smallest number of vertices found within an ROI (pooled over

hemispheres) across subjects. For visualization purposes all depicted raster plots were smoothed

along the vertical axis (FWHM = 1% of the number of data bins).

We also note that the statistical results were basically equivalent without any binning (i.e. directly

entering the shape parameter values of vertices into the GLM), the binning was applied only to

enable illustration of the results in raster plots.

These regression models over vertices and raster plots were used to evaluate whether crossmodal

modulation for auditory or visual stimuli in A1, PT, V1 or V2/3 depended on the unisensory visual

or auditory response.

Intersubject registration for surface projection of group results

To visualize patterns of the laminar profile shape parameters (i.e. 'constant' or 'linear') that are

consistent across subjects on the cortical surface, we transformed the individual surface projections

into a standardized study group space using a multimodal, multi-contrast surface registration

(MMSR) approach (Tardif et al., 2015). The transformation matrix for group normalization was

computed for the level-set corresponding to the mid-cortical surface and the high-resolution T1

maps. The T1 map was sampled radially along the cortical profile, averaged between 20% and 80%

cortical depth and smoothed tangentially (FWHM = 1.5 mm). To reduce computation time, the

level-set and the T1 map data were down-sampled to a resolution of 0.8 mm isotropic. We used a

two-stage registration. In the first step, we computed an intermediate mean surface by using the

median subject in the study group as the initial target. In the second step, MMSR was repeated

using this intermediate mean surface as the new target.

To apply this registration to our fMRI data, we computed the mean beta images for each of our 6

conditions, sampled them at the depths defined by the 6 intra-cortical surfaces (trilinear

interpolation) and then down-sampled to a resolution of 0.8 mm isotropic. We then used the

deformation field resulting from the second stage MMSR to transform those 36 images per subject

(6 conditions X 6 depths) into the group surface template space (trilinear interpolation). Mass

univariate laminar GLM were then performed in normalized group space and the parameters

corresponding to the 'constant' and 'linear' terms were averaged across subjects for each vertex.

Materials and Data Availability

The raw data of the results presented here are available in a BIDS format upon request

(remi.gau@gmail.com). Beta values extracted from our layers / ROIs for each participant as well as

the summary data necessary to reproduce our figures have been uploaded as CSV or mat files on the

open-science framework (https://osf.io/63dba/). Group average statistical maps are available in an

NIDM format from neurovault (https://neurovault.org/collections/5209/).

The results of the quality control MRIQC pipeline (https://mriqc.readthedocs.io/en/stable/) on the

BOLD data as well as information about motion and framewise displacement during scanning is

also available from the repository.

The code for the analysis is available at https://doi.org/10.5281/zenodo.3581319. The code to run

the fMRI experiment is available at https://doi.org/10.5281/zenodo.3581316.

ACKNOWLEDGMENTS

This project was funded by the ERC starter grant (mult-sens) and the Max Planck Society

AUTHOR CONTRIBUTIONS

Conceptualization: U.N., R.Tu.; Investigation: R.G.; Formal Analysis and Validation: R.G., U.N.;

Methodology: R.G., U.N., R.Tr., R.Tu.; Software: R.G., PL.B.; Resources: PL.B, R.Tr., R.Tu.;

Writing – Original Draft: R.G., UN; Writing – Review and Editing: R.G., U.N., PL.B., R.Tr., R.Tu.;

Visualization: R.G., Supervision: U.N.; Project administration: U.N, R.Tu., Funding Acquisition:

U.N., R.Tu.

REFERENCES

Atilgan H, Town SM, Wood KC, Jones GP, Maddox RK, Lee AKC, Bizley JK. 2018. Integration of Visual Information in Auditory Cortex Promotes Auditory Scene Analysis through Multisensory Binding. *Neuron* **97**:640-655.e4. doi:10.1016/j.neuron.2017.12.034

Bazin P-L, Weiss M, Dinse J, Schäfer A, Trampel R, Turner R. 2014. A computational framework for ultra-high resolution cortical segmentation at 7Tesla. *NeuroImage* **93**:201–209. doi:10.1016/j.neuroimage.2013.03.077

Beauchamp MS, Lee KE, Argall BD, Martin A. 2004. Integration of Auditory and Visual Information about Objects in Superior Temporal Sulcus. *Neuron* **41**:809–823. doi:10.1016/S0896-6273(04)00070-4

Bizley JK, Nodal FR, Bajo VM, Nelken I, King AJ. 2007. Physiological and Anatomical Evidence for Multisensory Interactions in Auditory Cortex. *Cereb Cortex* **17**:2172–2189. doi:10.1093/cercor/bhl128

Bressler DW, Fortenbaugh FC, Robertson LC, Silver MA. 2013. Visual spatial attention enhances the amplitude of positive and negative fMRI responses to visual stimulation in an eccentricity-dependent manner. *Vision Res* **85**:104–112. doi:10.1016/j.visres.2013.03.009

Budinger E, Heil P, Hess A, Scheich H. 2006. Multisensory processing via early cortical stages: Connections of the primary auditory cortical field with other sensory systems. *Neuroscience* **143**:1065–1083. doi:10.1016/j.neuroscience.2006.08.035

Butler JS, Foxe JJ, Fiebelkorn IC, Mercier MR, Molholm S. 2012. Multisensory Representation of Frequency across Audition and Touch: High Density Electrical Mapping Reveals Early Sensory-Perceptual Coupling. *J Neurosci* **32**:15338–15344. doi:10.1523/JNEUROSCI.1796-12.2012

Campi KL, Bales KL, Grunewald R, Krubitzer L. 2010. Connections of Auditory and Visual Cortex in the Prairie Vole (Microtus ochrogaster): Evidence for Multisensory Processing in Primary Sensory Areas. *Cereb Cortex* **20**:89–108. doi:10.1093/cercor/bhp082

Cappe C, Thelen A, Romei V, Thut G, Murray MM. 2012. Looming Signals Reveal Synergistic Principles of Multisensory Integration. *J Neurosci* **32**:1171–1182. doi:10.1523/JNEUROSCI.5517-11.2012

Chen G, Wang F, Gore JC, Roe AW. 2013. Layer-specific BOLD activation in awake monkey V1 revealed by ultra-high spatial resolution functional magnetic resonance imaging. *NeuroImage* **64**:147–155. doi:10.1016/j.neuroimage.2012.08.060

Ciaramitaro VM, Buračas GT, Boynton GM. 2007. Spatial and Cross-Modal Attention Alter Responses to Unattended Sensory Information in Early Visual and Auditory Human Cortex. *J Neurophysiol* **98**:2399–2413. doi:10.1152/jn.00580.2007

De Martino F, Moerel M, Ugurbil K, Goebel R, Yacoub E, Formisano E. 2015a. Frequency preference and attention effects across cortical depths in the human primary auditory cortex. *Proc Natl Acad Sci* **112**:16036–16041. doi:10.1073/pnas.1507552112

De Martino F, Moerel M, Xu J, van de Moortele P-F, Ugurbil K, Goebel R, Yacoub E, Formisano E. 2015b. High-Resolution Mapping of Myeloarchitecture In Vivo: Localization of Auditory Areas in the Human Brain. *Cereb Cortex* **25**:3394–3405. doi:10.1093/cercor/bhu150

Deneux T, Harrell ER, Kempf A, Ceballo S, Filipchuk A, Bathellier B. 2019. Context-dependent signaling of coincident auditory and visual events in primary visual cortex. *eLife* **8**:e44006. doi:10.7554/eLife.44006

Dick F, Taylor Tierney A, Lutti A, Josephs O, Sereno MI, Weiskopf N. 2012. In Vivo Functional and Myeloarchitectonic Mapping of Human Primary Auditory Areas. *J Neurosci* **32**:16095–16105. doi:10.1523/JNEUROSCI.1712-12.2012

Driver J, Noesselt T. 2008. Multisensory Interplay Reveals Crossmodal Influences on 'Sensory-Specific' Brain Regions, Neural Responses, and Judgments. *Neuron* **57**:11–23. doi:10.1016/j.neuron.2007.12.013

Duong TQ, Yacoub E, Adriany G, Hu X, U??urbil K, Kim S-G. 2003. Microvascular BOLD contribution at 4 and 7 T in the human brain: Gradient-echo and spin-echo fMRI with suppression of blood effects. *Magn Reson Med* **49**:1019–1027. doi:10.1002/mrm.10472

Ernst MO, Bülthoff HH. 2004. Merging the senses into a robust percept. *Trends Cogn Sci* **8**:162–169. doi:10.1016/j.tics.2004.02.002

Fairhall SL, Macaluso E. 2009. Spatial attention can modulate audiovisual integration at multiple cortical and subcortical sites. *Eur J Neurosci* **29**:1247–1257. doi:10.1111/j.1460-9568.2009.06688.x

Falchier A, Clavagnier S, Barone P, Kennedy H. 2002. Anatomical evidence of multimodal integration in primate striate cortex. *J Neurosci Off J Soc Neurosci* **22**:5749–5759. doi:20026562

Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, Yang Z, Chu C, Xie S, Laird AR, Fox PT, Eickhoff SB, Yu C, Jiang T. 2016. The Human Brainnetome Atlas: A New Brain Atlas Based on Connectional Architecture. *Cereb Cortex* **26**:3508–3526. doi:10.1093/cercor/bhw157

Fetsch CR, Pouget A, DeAngelis GC, Angelaki DE. 2012. Neural correlates of reliability-based cue weighting during multisensory integration. *Nat Neurosci* **15**:146–154. doi:10.1038/nn.2983

Formisano E, Kim DS, Di Salle F, van de Moortele PF, Ugurbil K, Goebel R. 2003. Mirror-symmetric tonotopic maps in human primary auditory cortex. *Neuron* **40**:859–869.

Fracasso A, Luijten PR, Dumoulin SO, Petridou N. 2018. Laminar imaging of positive and negative BOLD in human visual cortex at 7 T. *NeuroImage* **164**:100–111. doi:10.1016/j.neuroimage.2017.02.038

Friston KJ, Zarahn E, Josephs O, Henson RNA, Dale AM. 1999. Stochastic Designs in Event-Related fMRI. *NeuroImage* **10**:607–619. doi:10.1006/nimg.1999.0498

Ghazanfar A, Schroeder C. 2006. Is neocortex essentially multisensory? *Trends Cogn Sci* **10**:278–285. doi:10.1016/j.tics.2006.04.008

Goense J, Merkle H, Logothetis NK. 2012. High-Resolution fMRI Reveals Laminar Differences in Neurovascular Coupling between Positive and Negative BOLD Responses. *Neuron* **76**:629–639. doi:10.1016/j.neuron.2012.09.019

Hackett TA, Preuss TM, Kaas JH. 2001. Architectonic identification of the core region in auditory cortex of macaques, chimpanzees, and humans. *J Comp Neurol* **441**:197–222. doi:10.1002/cne.1407

Harel N, Lin J, Moeller S, Ugurbil K, Yacoub E. 2006. Combined imaging–histological study of cortical laminar specificity of fMRI signals. *NeuroImage* **29**:879–887. doi:10.1016/j.neuroimage.2005.08.016

Huber L, Handwerker DA, Jangraw DC, Chen G, Hall A, Stüber C, Gonzalez-Castillo J, Ivanov D, Marrett S, Guidi M, Goense J, Poser BA, Bandettini PA. 2017. High-Resolution CBV-fMRI Allows Mapping of Laminar Activity and Connectivity of Cortical Input and Output in Human M1. *Neuron* **96**:1253-1263.e7. doi:10.1016/j.neuron.2017.11.005

Ibrahim LA, Mesik L, Ji X, Fang Q, Li H, Li Y, Zingg B, Zhang LI, Tao HW. 2016. Cross-Modality Sharpening of Visual Cortical Processing through Layer-1-Mediated Inhibition and Disinhibition. *Neuron* **89**:1031–1045. doi:10.1016/j.neuron.2016.01.027

Iurilli G, Ghezzi D, Olcese U, Lassi G, Nazzaro C, Tonini R, Tucci V, Benfenati F, Medini P. 2012. Sound-Driven Synaptic Inhibition in Primary Visual Cortex. *Neuron* **73**:814–828. doi:10.1016/j.neuron.2011.12.026

Johnson JA, Zatorre RJ. 2005. Attention to Simultaneous Unrelated Auditory and Visual Events: Behavioral and Neural Correlates. *Cereb Cortex* **15**:1609–1620. doi:10.1093/cercor/bhi039

Kayser C, Logothetis NK, Panzeri S. 2010. Visual Enhancement of the Information Representation in Auditory Cortex. *Curr Biol* **20**:19–24. doi:10.1016/j.cub.2009.10.068

Kayser C, Petkov CI, Augath M, Logothetis NK. 2007. Functional Imaging Reveals Visual Modulation of Specific Fields in Auditory Cortex. *J Neurosci* **27**:1824–1835. doi:10.1523/JNEUROSCI.4737-06.2007

Kayser C, Petkov CI, Logothetis NK. 2008. Visual Modulation of Neurons in Auditory Cortex. *Cereb Cortex* **18**:1560–1574. doi:10.1093/cercor/bhm187

Kok P, Bains LJ, van Mourik T, Norris DG, de Lange FP. 2016. Selective Activation of the Deep Layers of the Human Primary Visual Cortex by Top-Down Feedback. *Curr Biol* **26**:371–376. doi:10.1016/j.cub.2015.12.038

Koopmans PJ, Barth M, Norris DG. 2010. Layer-specific BOLD activation in human V1. *Hum Brain Mapp* **31**:1297–1304. doi:10.1002/hbm.20936

Lakatos P, Chen C-M, O'Connell MN, Mills A, Schroeder CE. 2007. Neuronal Oscillations and Multisensory Interaction in Primary Auditory Cortex. *Neuron* **53**:279–292. doi:10.1016/j.neuron.2006.12.011

Lakatos P, O'Connell MN, Barczak A, Mills A, Javitt DC, Schroeder CE. 2009. The Leading Sense: Supramodal Control of Neurophysiological Context by Attention. *Neuron* **64**:419–430. doi:10.1016/j.neuron.2009.10.014

Landman BA, Bogovic JA, Carass A, Chen M, Roy S, Shiee N, Yang Z, Kishore B, Pham D, Bazin P-L, Resnick SM, Prince JL. 2013. System for Integrated Neuroimaging Analysis and Processing of Structure. *Neuroinformatics* **11**:91–103. doi:10.1007/s12021-012-9159-9

Laurienti PJ, Burdette JH, Wallace MT, Yen Y-F, Field AS, Stein BE. 2002. Deactivation of Sensory-Specific Cortex by Cross-Modal Stimuli. *J Cogn Neurosci* **14**:420–429. doi:10.1162/089892902317361930

Leitão J, Thielscher A, Werner S, Pohmann R, Noppeney U. 2013. Effects of Parietal TMS on Visual and Auditory Processing at the Primary Cortical Level – A Concurrent TMS-fMRI Study. *Cereb Cortex* **23**:873–884. doi:10.1093/cercor/bhs078

Leo F, Romei V, Freeman E, Ladavas E, Driver J. 2011. Looming sounds enhance orientation sensitivity for visual stimuli on the same side as such sounds. *Exp Brain Res* **213**:193–201. doi:10.1007/s00221-011-2742-8

Lewis R, Noppeney U. 2010. Audiovisual Synchrony Improves Motion Discrimination via Enhanced Connectivity between Early Visual and Auditory Areas. *J Neurosci* **30**:12329–12339. doi:10.1523/JNEUROSCI.5745-09.2010

Liang M, Mouraux A, Hu L, Iannetti GD. 2013. Primary sensory cortices contain distinguishable spatial patterns of activity for each sense. *Nat Commun* **4**. doi:10.1038/ncomms2979

Logothetis NK. 2008. What we can do and what we cannot do with fMRI. *Nature* **453**:869–878. doi:10.1038/nature06976

Lucas BC, Bogovic JA, Carass A, Bazin P-L, Prince JL, Pham DL, Landman BA. 2010. The Java Image Science Toolkit (JIST) for Rapid Prototyping and Publishing of Neuroimaging Software. *Neuroinformatics* **8**:5–17. doi:10.1007/s12021-009-9061-2

Maier JX, Chandrasekaran C, Ghazanfar AA. 2008. Integration of Bimodal Looming Signals through Neuronal Coherence in the Temporal Lobe. *Curr Biol* **18**:963–968. doi:10.1016/j.cub.2008.05.043

Maier JX, Neuhoff JG, Logothetis NK, Ghazanfar AA. 2004. Multisensory Integration of Looming Signals by Rhesus Monkeys. *Neuron* **43**:177–181. doi:10.1016/j.neuron.2004.06.027

Markuerkiaga I, Barth M, Norris DG. 2016. A cortical vascular model for examining the specificity of the laminar BOLD signal. *NeuroImage* **132**:491–498. doi:10.1016/j.neuroimage.2016.02.073

Marques JP, Kober T, Krueger G, van der Zwaag W, Van de Moortele P-F, Gruetter R. 2010. MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-mapping at high field. *NeuroImage* **49**:1271–1281. doi:10.1016/j.neuroimage.2009.10.002

McAuliffe MJ, Lalonde FM, McGarry D, Gandler W, Csaky K, Trus BL. 2001. Medical Image Processing, Analysis and Visualization in clinical research. IEEE Comput. Soc. pp. 381–386. doi:10.1109/CBMS.2001.941749

Meijer D, Veselič S, Calafiore C, Noppeney U. 2019. Integration of audiovisual spatial signals is not consistent with maximum likelihood estimation. *Cortex* **119**:74–88. doi:10.1016/j.cortex.2019.03.026

Meijer GT, Montijn JS, Pennartz CMA, Lansink CS. 2017. Audiovisual Modulation in Mouse Primary Visual Cortex Depends on Cross-Modal Stimulus Configuration and Congruency. *J Neurosci* **37**:8783–8796. doi:10.1523/JNEUROSCI.0468-17.2017

Mercier MR, Foxe JJ, Fiebelkorn IC, Butler JS, Schwartz TH, Molholm S. 2013. Auditory-driven phase reset in visual cortex: Human electrocorticography reveals mechanisms of early multisensory integration. *NeuroImage* **79**:19–29. doi:10.1016/j.neuroimage.2013.04.060

Morrill RJ, Hasenstaub AR. 2018. Visual Information Present in Infragranular Layers of Mouse Auditory Cortex. *J Neurosci* **38**:2854–2862. doi:10.1523/JNEUROSCI.3102-17.2018

Mozolic JL, Joyner D, Hugenschmidt CE, Peiffer AM, Kraft RA, Maldjian JA, Laurienti PJ. 2008. Cross-modal deactivations during modality-specific selective attention. *BMC Neurol* **8**. doi:10.1186/1471-2377-8-35

Muckli L, De Martino F, Vizioli L, Petro LS, Smith FW, Ugurbil K, Goebel R, Yacoub E. 2015. Contextual Feedback to Superficial Layers of V1. *Curr Biol* **25**:2690–2695. doi:10.1016/j.cub.2015.08.057

Müller NG, Kleinschmidt A. 2004. The attentional "spotlight's" penumbra: center-surround modulation in striate cortex. *Neuroreport* **15**:977–980.

Musacchia G, Schroeder CE. 2009. Neuronal mechanisms, response dynamics and perceptual functions of multisensory interactions in auditory cortex. *Hear Res* **258**:72–79. doi:10.1016/j.heares.2009.06.018

Nieuwenhuys R. 2013. The myeloarchitectonic studies on the human cerebral cortex of the Vogt–Vogt school, and their significance for the interpretation of functional neuroimaging data. *Brain Struct Funct* **218**:303–352. doi:10.1007/s00429-012-0460-z

Noesselt T, Rieger JW, Schoenfeld MA, Kanowski M, Hinrichs H, Heinze H-J, Driver J. 2007. Audiovisual Temporal Correspondence Modulates Human Multisensory Superior Temporal Sulcus Plus Primary Sensory Cortices. *J Neurosci* **27**:11431–11441. doi:10.1523/JNEUROSCI.2252-07.2007

Polimeni JR, Fischl B, Greve DN, Wald LL. 2010. Laminar analysis of 7T BOLD using an imposed spatial activation pattern in human V1. *NeuroImage* **52**:1334–1346. doi:10.1016/j.neuroimage.2010.05.005

Rockland KS, Ojima H. 2003. Multisensory convergence in calcarine visual areas in macaque monkey. *Int J Psychophysiol Off J Int Organ Psychophysiol* **50**:19–26.

Rohe T, Noppeney U. 2018. Reliability-Weighted Integration of Audiovisual Signals Can Be Modulated by Top-down Attention. *eneuro* **5**:ENEURO.0315-17.2018. doi:10.1523/ENEURO.0315-17.2018

Rohe T, Noppeney U. 2016. Distinct Computational Principles Govern Multisensory Integration in Primary Sensory and Association Cortices. *Curr Biol* **26**:509–514. doi:10.1016/j.cub.2015.12.056

Rohe T, Noppeney U. 2015. Cortical Hierarchies Perform Bayesian Causal Inference in Multisensory Perception. *PLOS Biol* **13**:e1002073. doi:10.1371/journal.pbio.1002073

Schroeder CE, Foxe JJ. 2002. The timing and laminar profile of converging inputs to multisensory areas of the macaque neocortex. *Brain Res Cogn Brain Res* **14**:187–198.

Schroeder CE, Smiley J, Fu KG, McGinnis T, O'Connell MN, Hackett TA. 2003. Anatomical mechanisms and functional implications of multisensory convergence in early cortical processing. *Int J Psychophysiol* **50**:5–17. doi:10.1016/S0167-8760(03)00120-X

Shmuel A, Yacoub E, Pfeuffer J, Van de Moortele PF, Adriany G, Hu X, Ugurbil K. 2002. Sustained negative BOLD, blood flow and oxygen consumption response and its coupling to the positive response in the human brain. *Neuron* **36**:1195–1210.

Shomstein S, Yantis S. 2004. Control of Attention Shifts between Vision and Audition in Human Cortex. *J Neurosci* **24**:10702–10706. doi:10.1523/JNEUROSCI.2939-04.2004

Silver MA, Shenhav A, D'Esposito M. 2008. Cholinergic Enhancement Reduces Spatial Spread of Visual Responses in Human Early Visual Cortex. *Neuron* **60**:904–914. doi:10.1016/j.neuron.2008.09.038

Smiley JF, Falchier A. 2009. Multisensory connections of monkey auditory cerebral cortex. *Hear Res* **258**:37–46. doi:10.1016/j.heares.2009.06.019

Smith AT, Singh KD, Greenlee MW. 2000. Attentional suppression of activity in the human visual cortex. *Neuroreport* **11**:271–277.

Stein BE, Stanford TR, Rowland BA. 2019. Multisensory Integration and the Society for Neuroscience: Then and Now. *J Neurosci* 0737–19. doi:10.1523/JNEUROSCI.0737-19.2019

Talsma D, Doty TJ, Woldorff MG. 2006. Selective Attention and Audiovisual Integration: Is Attending to Both Modalities a Prerequisite for Early Integration? *Cereb Cortex* **17**:679–690. doi:10.1093/cercor/bhk016

Talsma D, Senkowski D, Soto-Faraco S, Woldorff MG. 2010. The multifaceted interplay between attention and multisensory integration. *Trends Cogn Sci* **14**:400–410. doi:10.1016/j.tics.2010.06.008

Tardif CL, Schäfer A, Waehnert M, Dinse J, Turner R, Bazin P-L. 2015. Multi-contrast multi-scale surface registration for improved alignment of cortical areas. *NeuroImage* **111**:107–122. doi:10.1016/j.neuroimage.2015.02.005

Tootell RBH, Hadjikhani N, Hall EK, Marrett S, Vanduffel W, Vaughan JT, Dale AM. 1998. The Retinotopy of Visual Spatial Attention. *Neuron* **21**:1409–1422. doi:10.1016/S0896-6273(00)80659-5

Trampel R, Bazin P-L, Heidemann RM, Schäfer A, Ivanov D, Lohmann G, Geyer S, Turner R. 2012. Laminar-specific fingerprints of different sensorimotor areas obtained during imagined and actual finger tapping. *Proc Int Soc Magn Reson Med* **20**.

Trampel R, Bazin P-L, Pine K, Weiskopf N. 2017. In-vivo magnetic resonance imaging (MRI) of laminae in the human cortex. *NeuroImage*. doi:10.1016/j.neuroimage.2017.09.037

Tyll S, Bonath B, Schoenfeld MA, Heinze H-J, Ohl FW, Noesselt T. 2013. Neural basis of multisensory looming signals. *NeuroImage* **65**:13–22. doi:10.1016/j.neuroimage.2012.09.056

Van der Burg E, Olivers CNL, Bronkhorst AW, Theeuwes J. 2008. Pip and pop: Nonspatial auditory signals improve spatial visual search. *J Exp Psychol Hum Percept Perform* **34**:1053–1065. doi:10.1037/0096-1523.34.5.1053

Waehnert MD, Dinse J, Schäfer A, Geyer S, Bazin P-L, Turner R, Tardif CL. 2016. A subject-specific framework for in vivo myeloarchitectonic analysis using high resolution quantitative MRI. *NeuroImage* **125**:94–107. doi:10.1016/j.neuroimage.2015.10.001

Waehnert MD, Dinse J, Weiss M, Streicher MN, Waehnert P, Geyer S, Turner R, Bazin P-L. 2014. Anatomically motivated modeling of cortical laminae. *NeuroImage* **93**:210–220. doi:10.1016/j.neuroimage.2013.03.078

Wandell BA, Dumoulin SO, Brewer AA. 2007. Visual Field Maps in Human Cortex. *Neuron* **56**:366–383. doi:10.1016/j.neuron.2007.10.012

Wang L, Mruczek REB, Arcaro MJ, Kastner S. 2015. Probabilistic Maps of Visual Topography in Human Cortex. *Cereb Cortex* **25**:3911–3931. doi:10.1093/cercor/bhu277

Werner S, Noppeney U. 2011. The Contributions of Transient and Sustained Response Codes to Audiovisual Integration. *Cereb Cortex* **21**:920–931. doi:10.1093/cercor/bhq161

Werner S, Noppeney U. 2010. Distinct Functional Contributions of Primary Sensory and Association Areas to Audiovisual Integration in Object Categorization. *J Neurosci* **30**:2662–2675. doi:10.1523/JNEUROSCI.5091-09.2010

FIGURES

Figure 1: Experimental design, timeline and cortical layering

A. Experimental design: Participants were presented with auditory, visual and audiovisual

looming stimuli under auditory and visual attention.

B. Example trial and timeline: Participants were presented with brief auditory, visual or

audiovisual looming stimuli in 33 seconds blocks interleaved with 16 seconds fixation. At

the beginning of each block, a cue indicated whether the auditory or visual modality needed

to be attended. Brief visual and auditory targets (grey) were interspersed in the looming

activation blocks. Participants were instructed to respond to the targets in the attended and

ignore the targets in the unattended sensory modality.

C. Cortical layering: **Left**: A parasagittal section of a high resolution T1 map is shown with a

colour coded laminar label for each voxel (voxel size: $(0.4 \text{ mm})^3$). The primary auditory

cortex is circled in yellow. **Right**: The cortical sheet is defined by the pial and white matter

surfaces (thick black solid lines). Six additional surfaces (thin black solid lines) were

determined at different cortical depths. Data were mapped onto those surfaces by sampling

(blue dots) radially along the normal (white dashed line) to the mid-cortical depth surface

(not shown here). WM: white matter, GM: grey matter, CSF: cerebrospinal fluid.

Figure 2: Auditory and visual deactivations

A. BOLD response profiles: **Row 1 and 3:** The BOLD response (i.e. B parameters, across

subjects' mean ± SEM) for visual and auditory looming stimuli averaged over auditory and

visual attention in A1, PT, V1, V2/3 is shown as a function of percentage cortical depth.

WM: white matter; GM: grey matter; CSF: cerebrospinal fluid. Percentage cortical depth is

indicated by the small numbers and colour coded in red. **Rows 2 and 4:** Across subjects'

mean (± SEM) and violin plot of the participants' shape parameter estimates that

characterize the mean (C: constant) and linear increase (L: linear) of the laminar BOLD

response profile. n = 11.

B. Surface projections: Across subject' mean of the 'constant' (row i) and 'linear' increase

(row ii) shape parameter estimates of the laminar BOLD response profile for auditory and

visual looming stimuli (averaged over auditory and visual attention) are projected on an

inflated group mean surface to show auditory and visual regions of the left hemisphere. A1

and V1 are delineated by black solid lines, PT and V2/3 by dashed lines. For visualization

purposes only: i. surface projections were smoothed (FWHM = 1.5 mm); ii. values are also

presented for vertices for which data were not available from all subjects and which were

therefore not included in our formal statistical analysis. Grey areas denote vertices with no

available data for any subject.

Figure 3: Cross-modal modulation in auditory areas

A. Laminar profiles: **Row 1 and 3:** The BOLD response (solid line ; column 1 and 3) and

decoding accuracy (dashed line ; column 2 and 4) (across subjects' mean ± SEM) for [AV-

A] in A1 and PT is shown as a function of percentage cortical depth pooled (i.e. averaged)

over auditory and visual attention. WM: white matter, GM: grey matter, CSF: cerebrospinal

fluid. Percentage cortical depth is indicated by the small numbers and colour coded in red.

Rows 2 and 4: Across subjects' mean (± SEM) and violin plot of participants' shape

parameter estimates that characterize the mean (C: constant) and linear increase (L: linear)

of the laminar BOLD response and decoding accuracy profiles. n = 11.

B. Surface projections and raster plots: **Left**: Across subject' mean of the 'constant' (row i) and

'linear increase' (row ii) shape parameter estimates of the laminar BOLD response profile

for [AV-A] (averaged over auditory and visual attention) are projected on an inflated group

mean surface to show auditory regions of the left hemisphere. A1 and PT are delineated by

black solid and dashed lines. For visualization purposes only: i. surface projections were

893	smoothed (FWHM = 1.5 mm); ii. values are also presented for vertices for which data were
894	not available from all subjects and which were therefore not included in our formal
895	statistical analysis. Grey areas denote vertices with no available data for any subject. **Right:**
896	**Row i**. PT: The raster plot illustrates the statistical relationship between the 'constant' shape
897	parameters for the visual evoked response $[V\text{-}Fix]_{AttA,\ AttV}$ and the crossmodal modulation
898	$[AV\text{-}A]_{AttA,\ AttV}$ in PT. Each raster plot shows the laminar profiles (colour coded along
899	abscissa) of the vertices for the 'predicting contrast' [V-Fix] and of the 'predicted contrast'
900	[AV-A]. The laminar profiles of the vertices were sorted along the ordinate according to the
901	value of the 'constant' shape parameter for [V-Fix]. The raster plot shows that the laminar
902	profile of a vertex for [V-Fix] can predict its laminar profile for [AV-A]: PT vertices with
903	less deactivations across laminae for [V-Fix] are associated with greater crossmodal
904	enhancement [AV-A]. **Row ii**. The raster plot illustrates the statistical relationship between
905	the 'linear slope' shape parameters for the visual evoked response $[V\text{-}Fix]_{AttA,\ AttV}$ and the
906	crossmodal modulation [AV-A] in A1. Each raster plot shows the laminar profiles (colour
907	coded along abscissa) of the vertices for the 'predicting contrast' [V-Fix] and of the
908	'predicted contrast' [AV-A]. The laminar profiles of the vertices are sorted along the
909	ordinate according to the value of the 'linear' shape parameter for [V-Fix]. A1 vertices with
910	less deactivations in deeper laminae for [V-Fix] are associated with greater crossmodal
911	enhancement [AV-A] in deeper laminae.
912	To enable averaging the raster plots across participants, the vertices were binned after
913	sorting (number of bins for A1: 10100, number of bins for PT: 6800). For visualization
914	purposes all raster plots were smoothed along the vertical axis (FWHM = 1% of the number
915	of data bins). The subplot (i.e. black line) to the left of the raster plots shows the across
916	subjects' mean value (+/- STD) of the shape parameters (i.e. i. constant, ii, linear) of the
917	sorting contrast. n = 11.
918	

Figure 4: Attentional modulation

A. Laminar profiles: **Row 1, 3, 5, 7:** The BOLD response (solid line ; column 1 and 3) and

decoding accuracy (dashed line ; column 2 and 4) (across subjects' mean ± SEM) for

attentional modulation (i: top - [Att_A-Att_V] in A1 and PT ; ii: bottom - [Att_V-Att_A] in V1 and

V2/3) is shown as a function of percentage cortical depth pooled (i.e. averaged) over

auditory, visual and audiovisual looming stimuli. WM: white matter, GM: grey matter, CSF:

cerebrospinal fluid. Percentage cortical depth is indicated by the small numbers and colour

coded in red. **Rows 2, 4, 6, 8:** Across subjects' mean (± SEM) and violin plot of the

participants' shape parameter estimates that characterize the mean (C: constant) and linear

increase (L: linear) of the laminar BOLD response and decoding accuracy profiles. n = 11.

B. Surface projections: Across subject' mean of the 'constant' shape parameter estimates for

attentional modulation [Att_A-Att_V]$_{A,V, AV}$ in A1 and PT (i: top) and in V1 and V2/3 (ii:

bottom) are projected on an inflated group mean surface of the left hemisphere. A1 and V1

are delineated by black solid lines. PT and V2/3 are delineated by dashed lines. For

visualization purposes only: i. surface projections were smoothed (FWHM = 1.5 mm); ii.

values are also presented for vertices for which data were not available from all subjects and

which were therefore not included in our formal statistical analysis; iii borders between

visual-induced activations and deactivations (white dashed lines on the right) are reported

here from figure 2B. Grey areas denote vertices with no available data for any subject.

TABLES

Table 1: Auditory and visual deactivations

		linear or constant		constant		linear	
	mean(A1, PT)	F(2,40)=9.280	**p<0.001**	t(10)=-2.460	**p=0.017**a	F(1,20)=2.083	p=0.164
[V-fix]Att_A, Att_V	**A1**			t(10)=-2.077	**p=0.032** a		
	PT			t(10)=-2.042	**p=0.034**a		
	mean(V1, V23)	F(2,40)=58.615	**p<0.001**	t(10)=-5.547	**p<0.001**a	F(1,20)=22.433	**p<0.001**
[A-fix]Att_A, Att_V	**V1**			t(10)=-6.538	**p<0.001** a	t(10)=-5.080	**p<0.001**
	V2-3			t(10)=-4.305	**p<0.001**a	t(10)=-4.142	**p=0.002**

[a] indicates p-values based on a one-sided t-test based on a priori hypotheses. p-values < 0.05 are indicated in bold. n = 11

Using 2 (shape parameter: constant, linear) x 2 (ROI: primary, non-primary) linear mixed effects models, we perfomed the following statistical

comparisons in a 'step down procedure':

1. Two-dimensional F-test assessing whether the constant or linear parameter (e.g. each averaged across ROIs in auditory resp. visual cortices),

    was significantly different from zero (dark grey),

944   2. If this two-dimensional F-test was significant, we computed one dimensional F-tests separately for the constant and the linear parameters (again

945   averaged across auditory resp. visual ROIs) (light grey),

946   3. If the one dimensional F-test was significant, we computed follow-up t-tests separately for each of the two ROIs (white).

Table 2: Effects of the cross-modal modulation on the laminar BOLD response and decoding accuracy profiles in auditory areas

A) BOLD profile

		linear or constant		constant	linear
[AV - A]Att_A, Att_V	mean(A1, PT)	F(2,40)=0.196	p=0.823		

B) Decoding profile

		linear or constant		constant	linear
	mean(A1, PT)	F(2,40)=34.946	**p<0.001**	F(1,20)=21.966 **p<0.001**	F(1,20)=1.850 p=0.189
[AV VS A]att A, att V				**A1** t(10)=3.867 **p=0.003**	
				PT t(10)=4.992 **p<0.001**	

Using 2 (shape parameter: constant, linear) x 2 (ROI: primary, non-primary) linear mixed effects models, we perfomed the following statistical

comparisons in a 'step down procedure':

1. Two-dimensional F-test assessing whether the constant or linear parameter (e.g. each averaged across ROIs in auditory resp. visual cortices),

was significantly different from zero (dark grey),

2. If this two-dimensional F-test was significant, we computed one dimensional F-tests separately for the constant and the linear parameters (again

averaged across auditory resp. visual ROIs) (light grey),

3. If the one dimensional F-test was significant, we computed follow-up t-tests separately for each of the two ROIs (white).

p-values < 0.05 are indicated in bold. n = 11

Table 3: Effects of the attentional modulation (irrespective of stimulus type) on the laminar BOLD response and decoding accuracy profiles

A) BOLD profile

		linear or constant			constant		linear	
[Att_V - Att_A]A, V, AV	**mean(A1, PT)**	F(2,40)=12.602	**p<0.001**		F(1,20)=9.249 **p=0.006**		F(1,20)=12.163	**p=0.002**
				A1	t(10)=1.882 **p=0.089**		t(10)=3.123	**p=0.011**
				PT	t(10)=4.523 **p=0.001**		t(10)=3.361	**p=0.007**
[Att_V - Att_A]A, V, AV	**mean(V1, V23)**	F(2,40)=0.669	p=0.518					

B) Decoding profile

		linear or constant			constant		linear	
[Att_A VS Att_V]A, V, AV	**mean(A1, PT)**	F(2,40)=4.687	**p=0.015**		F(1,20)=4.882 **p=0.039**		F(1,20)=4.028	p=0.058
				A1	t(10)=1.260 p=0.236			

				PT $t(10)=2.031$ $p=0.070$		
[Att_A VS Att_V]A, V, AV	**mean(V1, V23)**	$F(2,40)=20.026$	**p<0.001**	$F(1,20)=13.564$ **p=0.001**	$F(1,20)=9.951$	**p=0.005**
				V1 $t(10)=2.472$ **p=0.033**	$t(10)=1.359$	p=0.204
				V2-3 $t(10)=4.298$ **p=0.002**	$t(10)=3.089$	**p=0.011**

Using 2 (shape parameter: constant, linear) x 2 (ROI: primary, non-primary) linear mixed effects models, we perfomed the following statistical

comparisons in a 'step down procedure':

1. Two-dimensional F-test assessing whether the constant or linear parameter (e.g. each averaged across ROIs in auditory resp. visual cortices),

was significantly different from zero (dark grey),

2. If this two-dimensional F-test was significant, we computed one dimensional F-tests separately for the constant and the linear parameters (again

averaged across auditory resp. visual ROIs) (light grey),

3. If the one dimensional F-test was significant, we computed follow-up t-tests separately for each of the two ROIs (white).

p-values   <   0.05   are   indicated   in   bold.   n   =   11

  SUPPLEMENTARY FIGURES

Supplementary figure 1: Behavioural results

Percentage of responses to auditory (red) and visual (blue) targets in auditory, visual and audiovisual blocks (along the x-axis) and under auditory (top panel) and visual (bottom panel) attention. Thick lines represent group means +/- SEM and each thin line represents a participant. Responses to visual targets under auditory attention and responses to auditory targets under visual attention are false alarms (and vice versa for visual attention).

Supplementary figure 2: Segmentation and coregistration

Sagittal section of T1 structural images (left), the corresponding coregistered mean EPI images of four representative participants. The definition of the six laminae is overlaid in A1

Supplementary figure 3: Auditory and visual activations

A. BOLD response profiles: **Row 1 and 3:** The BOLD response (i.e. B parameters, across subjects' mean ± SEM) for auditory and visual activations induced by looming stimuli averaged over auditory and visual attention in A1, PT, V1, V2/3 is shown as a function of percentage cortical depth. WM: white matter; GM: grey matter; CSF: cerebrospinal fluid. Percentage cortical depth is indicated by the small numbers and colour coded in red. **Rows 2 and 4:** Across subjects' mean (± SEM) and violin plot of the participants' shape parameter estimates that characterize the mean (C: constant) and linear increase (L: linear) of the laminar BOLD response profile. n = 11.

B. Surface projections: Across subject' mean of the 'constant' (row i) and 'linear' increase (row ii) shape parameter estimates of the laminar BOLD response profile for auditory and visual looming stimuli (averaged over auditory and visual attention) are projected on an inflated group mean surface to show auditory and visual regions of the left hemisphere. A1

and V1 are delineated by black solid lines, PT and V2/3 by dashed lines. For visualization

purposes only: i. borders between visual-induced activations and deactivations (white

dashed lines) were defined based on visual inspection; ii. surface projections were smoothed

(FWHM = 1.5 mm); iii. values are also presented for vertices for which data were not

available from all subjects and which were therefore not included in our formal statistical

analysis. Grey areas denote vertices with no available data for any subject.

Supplementary figure 4: Cross-modal modulation in visual areas

Laminar profiles: **Row 1 and 3:** The BOLD response (solid line ; column 1 and 3) and decoding

accuracy (dashed line ; column 2 and 4) (across subjects' mean ± SEM) for [AV-A] in V1 and V2/3

is shown as a function of percentage cortical depth pooled (i.e. averaged) over auditory and visual

attention. WM: white matter, GM: grey matter, CSF: cerebrospinal fluid. Percentage cortical depth

is indicated by the small numbers and colour coded in red. **Rows 2 and 4:** Across subjects' mean (±

SEM) and violin plot of participants' shape parameter estimates that characterize the mean (C:

constant) and linear increase (L: linear) of the laminar BOLD response and decoding accuracy

profiles. n = 11.

Supplementary figure 5: Auditory and visual responses in visual areas

Surface projection for individual subjects: Within subject 'constant' shape parameter estimates of

the laminar BOLD response profile for visual (column i and ii) and auditory (column iii and iv)

stimuli are projected on each subject's inflated surface to show visual regions of the left (column i

and iii) and right (column ii and iv) hemisphere. V1 and V2/3 are delineated by black solid and

dashed lines respectively. For visualization purposes only, surface projections were smoothed

(FWHM = 1.5 mm). Grey areas denote vertices with no available data.

Supplementary figure 6: Auditory and visual responses in auditory areas

<u>Surface projection for individual subjects:</u> Within subject 'constant' shape parameter estimates of

the laminar BOLD response profile for visual (column i and ii) and auditory (column iii and iv) are

projected on each subject's inflated surface to show visual regions of the left (column i and iii) and

right (column ii and iv) hemisphere. A1 and PT are delineated by black solid and dashed lines

respectively. For visualization purposes only, surface projections were smoothed (FWHM = 1.5

mm). Grey areas denote vertices with no available data.

Supplementary figure 7: Raster plots for auditory induced activations and visual induced deactivations in auditory areas

The raster plots illustrate the statistical relationship between the 'constant' shape parameters for the

visual evoked response $[\text{V-Fix}]_{\text{AttA, AttV}}$ and auditory evoked response $[\text{A-Fix}]_{\text{AttA, AttV}}$ in A1 (left)

and PT (right). Each raster plot shows the laminar profiles (colour coded along abscissa) of the

vertices for the 'predicting contrast' [V-Fix] and of the 'predicted contrast' [A-Fix]. The vertex

profiles were sorted along the ordinate according to the value of the 'constant' shape parameter for

[V-Fix]. The raster plot shows that the laminar profile of a vertex for [V-Fix] cannot predict its

laminar profile for [A-Fix]. To enable averaging the raster plots across participants, the vertices

were binned after sorting (number of bins for A1: 10100, number of bins for PT: 6800). For

visualization purposes all raster plots were smoothed along the vertical axis (FWHM = 1% of the

number of data bins). The subplot (i.e. black line) to the left of the raster plots shows the across

subjects' mean value (+/- STD) of the shape parameters (i.e. i. constant, ii, linear) of the sorting

contrast. The violin plot show the distribution across subject of beta values for "a" in the regression

model   *[A-Fix]*   =   *a*   *[V-FIX]*   +   *b*.   n   =   11.

SUPPLEMENTARY TABLES

Supplementary table 1: Behavioural results

Stimuli Targets	Auditory	Visual	Audio-visual
	Attend to auditory		
Auditory	80.11% (13.92%)	93.18% (7.63%)	67.05% (15.08%)
Visual	6.82% (7.63%)	2.27% (4.21%)	11.36% (7.82%)
	Attend to visual		
Auditory	4.55% (11.21%)	6.25% (6.25%)	7.95% (7.95%)
Visual	65.91% (16.14%)	69.32% (14.91%)	65.91% (11.98%)

 Notes: Percentage of target responses (mean and STD across subjects) in the six conditions of our 2 x 3 experimental design. n = 11

 Please note that responses to visual targets under auditory attention and auditory targets under visual attention are false alarms

Supplementary table 2: ROI size and coverage

ROI	Number of vertices (*10^3)			Proportion of ROI covered		
	mean	STD	range	mean	STD	range
A1	19.64	3.60	(15.05- 24.30)	0.96	0.06	(0.84- 1.00)
PT	10.17	2.31	(6.85- 14.36)	0.96	0.06	(0.79- 1.00)
V1	34.54	6.54	(24.64 - 48.44)	0.97	0.02	(0.94 - 1.00)
V2-3	68.74	12.05	(43.59- 87.51)	0.82	0.08	(0.64- 0.90)

028 Notes: 'Number of vertices' refers to the vertices with valid data at all the sampled cortical depths. This vertex count was divided by the total number

029 of vertices included in the initial ROI definition to compute the 'Fraction of the ROI covered'. Note that those numbers are pooled over both

030 hemispheres. n = 11

Supplementary table 3: Auditory and visual activations

		linear or constant		constant		linear	
	mean(A1, PT)	F(2,40)=136.799	**p<0.001**	t(10)=11.537	**p<0.001**a	F(1,20)=52.749	**p<0.001**
[A-fix]Att_A, Att_V	**A1**			t(10)=11.906	**p<0.001** a	t(10)=8.152	**p<0.001**
	PT			t(10)=7.284	**p<0.001**a	t(10)=10.761	**p<0.001**
	mean(V1, V23)	F(2,40)=158.364	**p<0.001**	t(10)=9.864	**p<0.001**a	F(1,20)=49.620	p<0.001
[V-fix]Att_A, Att_V	**V1**			t(10)=8.152	**p<0.001** a	t(10)=6.154	**p<0.001**
	V2-3			t(10)=10.761	**p<0.001**a	t(10)=7.675	**p<0.001**

Using 2 (shape parameter: constant, linear) x 2 (ROI: primary, non-primary) linear mixed effects models, we perfomed the following statistical

comparisons in a 'step down procedure':

1. Two-dimensional F-test assessing whether the constant or linear parameter (e.g. each averaged across ROIs in auditory resp. visual cortices),

was significantly different from zero (dark grey),

2. If this two-dimensional F-test was significant, we computed one dimensional F-tests separately for the constant and the linear parameters (again

averaged across auditory resp. visual ROIs) (light grey),

3. If the one dimensional F-test was significant, we computed follow-up t-tests separately for each of the two ROIs (white).

[a] indicates p-values based on a one-sided t-test based on a priori hypotheses. p-values < 0.05 are indicated in bold. n = 11

Supplementary table 4: Cross-modal modulation in visual areas

A) BOLD profile

		linear or constant		constant	linear
[AV - V]Att_A, Att_V	**mean(V1, V23)**	F(2,40)=0.723	p=0.491		

B) Decoding profile

		linear or constant		constant	linear
[AV VS V]att A, att V	**mean(V1, V23)**	F(2,40)=0.696	p=0.505		

043   Using 2 (shape parameter: constant, linear) x 2 (ROI: primary, non-primary) linear mixed effects models, we perfomed the following statistical

044   comparisons in a 'step down procedure':

1. Two-dimensional F-test assessing whether the constant or linear parameter (e.g. each averaged across ROIs in auditory resp. visual cortices),

   was significantly different from zero (dark grey),

2. If this two-dimensional F-test was significant, we computed one dimensional F-tests separately for the constant and the linear parameters (again

   averaged across auditory resp. visual ROIs) (light grey),

3. If the one dimensional F-test was significant, we computed follow-up t-tests separately for each of the two ROIs (white).

p-values < 0.05 are indicated in bold. n = 11

A   Experimental design

B   Example trial and timeline

100 ms
Auditory target

Visual target
50 ms

stimuli

fixation

50 trials
33 s

16 s

Attention cues

C   Cortical layering

Cortical depth

86 71 57 43 29 14

WM   GM   CSF

Auditory and visual deactivations

A   BOLD-response profiles

A1   Visual deactivations
$[\text{V-fix}]_{\text{Att}_A, \text{Att}_V}$

V1   Auditory deactivations
$[\text{A-fix}]_{\text{Att}_A, \text{Att}_V}$

PT $[\text{V-fix}]_{\text{Att}_A, \text{Att}_V}$

V2-3 $[\text{A-fix}]_{\text{Att}_A, \text{Att}_V}$

B   Surface projections

$[\text{A - fix}]_{\text{Att}_A, \text{Att}_V}$ $[\text{V - fix}]_{\text{Att}_A, \text{Att}_V}$

Constant

Linear

S Param. est. [a u]

constant
-3.0   -1.5   0.0   1.5   3.0
-0.8   -0.4   0.0   0.4   0.8
linear

—— A1   —— V1
······ PT   ······ V2/3