Resolving multisensory and attentional influences across cortical depth in sensory cortices

Authors/Affiliations

Remi Gau[a*], Pierre-Louis Bazin[b,c,], Robert Trampel[b], Robert Turner[b,d], Uta Noppeney[a]

a Computational Neuroscience and Cognitive Robotics Centre, University of Birmingham, B15 2TT

Birmingham, UK

b Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

c Integrative Model-based Cognitive Neuroscience research unit, University of Amsterdam,

Amsterdam, The Netherlands

d Sir Peter Mansfield Imaging Centre, University of Nottingham, NG7 2RD, Nottingham, UK

* Corresponding author: Remi Gau <remi.gau@gmail.com>

Classification

Social sciences/Psychological and cognitive sciences

Biological sciences/Neuroscience

Keywords: multisensory integration, audiovisual, primary sensory cortices, deactivations, crossmodal attention, cortical layers, layer-dependent fMRI, ultra-high-resolution fMRI, 7 Tesla

ABSTRACT

In our natural environment our senses are bombarded with a myriad of diverse signals, only a subset of which is relevant for our current goals. Using sub-millimeter-resolution fMRI at 7T we resolved BOLD response and activation patterns across cortical depth in early sensory cortices to auditory, visual and audiovisual stimuli under auditory and visual attention. Irrespective of attention, auditory stimulation induced widespread inhibition in visual cortices, while auditory relative to visual attention suppressed mainly central visual field representations. Conversely, visual stimulation suppressed activations in auditory cortices, but amplified the response to concurrent auditory stimuli, in a patchy topography. Critically, the competitive and cooperative multisensory interactions in auditory cortices were stronger in relatively deeper cortical laminae, while attentional influences were greatest at the cortical surface. These distinct depth-dependent activation profiles suggest that multisensory and attentional mechanisms regulate sensory processing via partly distinct neural circuitries. Our findings are crucial for understanding how the human brain flexibly regulates information flow across the senses to interact effectively with our complex multisensory world.

INTRODUCTION

In our natural environment our senses are exposed to a constant influx of sensory signals that arise from many different sources. How the brain flexibly regulates information flow across the senses to enable effective interactions with the world remains unclear.

Mounting evidence from neuroimaging (Beauchamp et al., 2004; Noesselt et al., 2007; Rohe and Noppeney, 2016), neurophysiology (Atilgan et al., 2018; Kayser et al., 2010, 2008; Lakatos et al., 2007) and neuroanatomy (Falchier et al., 2002; Rockland and Ojima, 2003) suggests that interactions across the senses are pervasive in neocortex, arising even in primary cortices (Driver and Noesselt, 2008; Ghazanfar and Schroeder, 2006; Liang et al., 2013; Schroeder and Foxe, 2002). Visual stimuli can drive as well as modulate responses in cortices that are dedicated to other sensory modalities. Most prominently, functional magnetic resonance imaging (fMRI) in humans has shown that visual stimuli can induce crossmodal deactivations in primary and secondary auditory cortices (Laurienti et al., 2002; Leitão et al., 2013; Mozolic et al., 2008), yet enhance the response of a concurrent auditory stimulus (Werner and Noppeney, 2011, 2010). Further, neurophysiological research has shown that multisensory influences emerge early in sensory cortices and are to some extent preserved in anaesthetized animals (Atilgan et al., 2018; Butler et al., 2012; Ibrahim et al., 2016; Iurilli et al., 2012; Kayser et al., 2007; Mercier et al., 2013; Morrill and Hasenstaub, 2018). Yet, the ability to extrapolate from neurophysiological findings in animals to human fMRI studies is limited by the nature of the BOLD response, which pools neural activity over time and across a vast number of neurons (Logothetis, 2008).

Information flow is regulated not only by multisensory but also by attentional mechanisms that are guided by our current goals (Fairhall and Macaluso, 2009; Talsma et al., 2010). Critically, multisensory and attentional mechanisms are closely intertwined. Both enhance perceptual sensitivity (Leo et al., 2011) and precision of sensory representations (Ernst and Bülthoff, 2004; Fetsch et al., 2012; Rohe and Noppeney, 2015). Most importantly, the co-occurrence of two congruent sensory stimuli boosts the salience of an event (Van der Burg et al., 2008), which may thereby attract greater

attention. Conversely, a stimulus presented in one sensory modality alone may withdraw attentional resources from other sensory modalities. Behavioural and functional imaging studies have shown that shifting attention endogenously to one sensory modality reduces processing and activations in the unattended sensory systems (Ciaramitaro et al., 2007; Johnson and Zatorre, 2005; Mozolic et al., 2008; Rohe and Noppeney, 2015; Shomstein and Yantis, 2004). As a consequence, attentional mechanisms may contribute to competitive and cooperative interactions across the senses, for instance by amplifying responses for congruent audiovisual stimuli, and generating crossmodal deactivations for unisensory stimuli. Inter-sensory attention can also profoundly modulate multisensory interactions (Talsma et al., 2006). Most prominently, the influence of visual stimuli on auditory cortices was shown to be enhanced or even enabled when attention was focused on the visual sense (Lakatos et al., 2009).

Critically, while previous neurophysiological studies have revealed influences of modality-specific attention predominantly in superficial laminae in non-human primates (Lakatos et al., 2009), visual influences on auditory cortices have recently been shown to be most prominent in deep layer 6 of auditory cortex in rodents (Morrill and Hasenstaub, 2018). In other words, a double dissociation of attentional and multisensory influences was observed in auditory cortices when combining neurophysiological evidence from primates and rodents.

To investigate whether this double dissociation can be found in human neocortex we exploited recent advances in submillimeter-resolution fMRI at 7T that allow the characterization of depth-dependent activation profiles (De Martino et al., 2015a; Kok et al., 2016; Muckli et al., 2015; Trampel et al., 2012). While BOLD fMRI is not yet able to attribute activations unequivocally to specific cortical layers (Goense et al., 2012; Huber et al., 2017; Markuerkiaga et al., 2016; Trampel et al., 2017), the observation in the same cortical territories of distinct laminar activation profiles induced by multisensory and attentional influences would strongly imply distinct neural mechanisms.

The current study investigated the processing of auditory, visual or audiovisual looming stimuli under auditory and visual attention in the human brain. Combining submillimeter-resolution fMRI at 7T

93   with laminar and multivariate pattern analyses we show distinct depth-dependent activation profiles

94   and/or patterns for multisensory and attentional influences in early auditory and visual cortices. These

95   results suggest that multisensory and attentional mechanisms regulate sensory processing in early

96   sensory cortices via partly distinct neural circuitries.

RESULTS

In this fMRI study, participants were presented with blocks of auditory (A), visual (V) and audiovisual (AV) looming stimuli interleaved with fixation (Figure 1A). We used looming motion as a biologically relevant and highly salient stimulus that reliably evokes crossmodal influences in sensory cortices in human neuroimaging and animal neurophysiology (Cappe et al., 2012; Maier et al., 2008; Tyll et al., 2013). Modality-specific attention was manipulated by requiring participants to detect and respond selectively to weak auditory or visual targets, which were adjusted prior to the main study to threshold performance in sound amplitude or visual size for each participant. The targets were interspersed throughout all auditory, visual and audiovisual looming blocks (e.g. visual targets were presented during both visual and auditory looming blocks; see Figure 1B).

For the behavioural analysis, the percentages of visual and auditory targets that gained a response (see supplementary Table 1 for '% target responses') were entered into a 2 (target modality: auditory vs. visual) X 2 (attended modality: auditory vs. visual) X 3 (stimulus block modality: auditory, visual, audio-visual) repeated measures ANOVA. We observed significant main effects of attended modality ($F_{(1, 10)}=291.263$, $p<0.001$, $\eta_p^2=0.967$), target modality ($F_{(1, 10)}=13.289$, $p=0.004$, $\eta_p^2=0.571$), and stimulus modality ($F_{(1.986, 19.864)}=5.304$, $p<0.014$, $\eta_p^2=0.347$), together with a significant interaction between target modality and attended modality ($F_{(1, 10)}=14.786$, $p<0.003$, $\eta_p^2=0.597$). This confirmed that participants had successfully maintained auditory or visual attention as instructed. Further, we observed significant interactions between target modality and stimulus modality ($F_{(1.813, 18.126)}=8.149$, $p<0.004$, $\eta_p^2=0.449$) and between attended modality and stimulus modality ($F_{(1.436, 14.360)}=24.034$, $p<0.001$, $\eta_p^2=0.706$). These interactions resulted from greater hit rates for auditory targets given auditory attention than for visual targets given visual attention during both auditory ($t(10)=2.845$, $p=0.017$, Hedges $g_{av}=0.804$) and visual blocks ($t(10)=4.432$, $p=0.001$, Hedges $g_{av}=2.037$), but not for audio-visual blocks ($t(10)=0.276$, $p=0.788$, Hedges $g_{av}=0.081$) suggesting that

observers' performance was not completely matched across all conditions. For completeness, there

was no significant 3-way interaction (F(1.783, 17.826)=2.467, p<0.118, η_p^2=0.198).

Using sub-millimeter resolution fMRI, we next characterized the laminar profiles in auditory (primary

auditory cortex, A1; planum temporale, PT) and visual (primary, V1; higher order, V2/3) regions for

the following effects: 1. sensory (de)activations in unisensory contexts for the preferred (e.g. [V-Fix]

in visual cortices) and non-preferred stimuli (i.e. crossmodal, e.g. [V-Fix] in auditory cortices), 2.

crossmodal modulation (e.g. [AV-A] in auditory cortices, [AV-V] in visual cortices) in audiovisual

context and 3. direct and modulatory effects of modality-specific attention.

Briefly, we used the following methodological approach (see methods): in each ROI and participant,

we estimated the regional BOLD response (i.e. 'B parameter estimate') and the multivariate pattern

decoding accuracy for each of the six laminae (e.g. figure 2A, row 1). We then characterized the

laminar profiles of BOLD response and decoding accuracy in terms of a constant and a linear shape

parameter (i.e. 'S parameter estimates') and show their across-subjects' mean and distribution in

violin plots (e.g. figure 2A, row 2). Finally, we characterized the spatial topography of those S

parameter estimates by projecting their group mean onto a normalized group cortical surface (e.g. see

figure 2B).

All statistical results (i.e. p-values, effect sizes) are presented in tables 1, 2 and 3.

Figure 1: Experimental design, timeline and cortical layering

A. Experimental design: Participants were presented with auditory, visual and audiovisual looming stimuli under auditory and visual attention.

B. Example trial and timeline: Participants were presented with brief auditory, visual or audiovisual looming stimuli in 33 seconds blocks interleaved with 16 seconds fixation. At the beginning of each block, a cue indicated whether the auditory or visual modality needed to be attended. Brief visual and auditory targets (grey) were interspersed in the looming activation blocks. Participants were instructed to respond to the targets in the attended and ignore the targets in the unattended sensory modality.

C. Cortical layering: **Left**: A parasagittal section of a high resolution T1 map is shown with a colour coded laminar label for each voxel (voxel size: $(0.4 \text{ mm})^3$). The primary auditory cortex is encircled in yellow. **Right**: The cortical sheet is defined by the pial and white matter surfaces (thick black solid lines). Six additional surfaces (thin black solid lines) were determined at different cortical depths. Data were mapped onto those surfaces by sampling (blue dots) radially along the normal (white dashed line) to the mid-cortical depth surface (not shown here). WM: white matter, GM: grey matter, CSF: cerebrospinal fluid.

Auditory cortices

Auditory stimuli evoked a positive BOLD response in primary auditory cortex and especially in

anterior portions of the planum temporale. As expected from the typical physiological point spread

function of the GE-EPI BOLD signal, the positive BOLD signal increased roughly linearly towards

the cortical surface (Markuerkiaga et al., 2016) (Figure 2A and Table 1).

Table 1

Effects of auditory and visual stimuli on the laminar BOLD response

BOLD profile		Constant					Linear				
Contrast	ROI	mean	STD	p value	95% CI effect size		mean	STD	p value	95% CI effect size	
$[\text{A-Fix}]_{\text{att A, att V}}$											
	A1	1.58	(0.44)	**<.001** [a]	[2.38 - 4.82]		0.37	(0.16)	**<.001**	[1.68 - 3.12]	
	PT	1.01	(0.46)	**<.001** [a]	[1.33 - 2.79]		0.24	(0.14)	**<.001**	[1.14 - 2.26]	
	V1	-0.87	(0.44)	**<.001** [a]	[-2.51 - -1.24]		-0.16	(0.11)	**<.001**	[-1.96 - -0.89]	
	V2/3	-0.52	(0.40)	**<.001** [a]	[-1.70 - -0.74]		-0.10	(0.08)	**0.002**	[-1.84 - -0.61]	
$[\text{V-Fix}]_{\text{att A, att V}}$											
	A1	-0.10	(0.15)	**0.034** [a]	[-1.22 - -0.07]		0.01	(0.05)	0.386	[-0.37 - 1.07]	
	PT	-0.15	(0.24)	**0.036** [a]	[-1.68 - 0.06]		0.04	(0.07)	0.110	[-0.06 - 0.87]	
	V1	1.47	(0.60)	**<.001** [a]	[1.42 - 3.54]		0.23	(0.12)	**<.001**	[0.99 - 2.95]	
	V2/3	1.54	(0.47)	**<.001** [a]	[1.69 - 4.34]		0.24	(0.10)	**<.001**	[1.47 - 3.02]	

164 Notes: Unless otherwise stated p-values are based on an exact two-sided sign permutation test. [a] indicates p-values based on an exact one-sided sign

165 permutation test based on a priori hypotheses. p-values < 0.05 are indicated in bold. n = 11

By contrast, deactivations induced by crossmodal visual stimuli (i.e. a negative BOLD response for [V-Fix]) were observed in both A1 and PT, with a non-significant trend in PT towards stronger deactivations in deeper laminae (Figure 2A, B and Table 1). These visually-induced deactivations were generally observed for both auditory and visual attention conditions, with no significant difference between them ([Att_A-Att_V] for visual stimuli in A1: $p = 0.4$ and in PT: $p = 0.6$).

Strikingly, while visual stimuli deactivated auditory cortices when presented alone, they enhanced the BOLD response in PT for an auditory stimulus (i.e. AV > A, Table 2A, Figure 3A). In other words, in PT the suppressive effect of unisensory visual stimuli turned into a crossmodal enhancement, when visual stimuli were presented together with auditory stimuli. Again, this crossmodal enhancement was not significantly modulated by attention. The visually induced amplification of auditory responses was observed predominantly in caudal parts and superficial laminae of PT (i.e. significant positive linear effect in Figure 3A, row 2, and Table 2A).

By contrast, pattern classifiers succeeded in discriminating between patterns elicited by [AV vs. A] conditions across all laminae in both PT and A1, even in those laminae where no crossmodal effect was observed for the regional BOLD response profile (Figure 3A 2^{nd} column and Table 2B). Thus, even when the BOLD response averaged across all vertices in A1 or PT did not differ significantly for AV and A stimuli at a particular depth, the visual stimulus changed the activation pattern elicited by a concurrent auditory stimulus. This provides evidence that sub-regions with crossmodal enhancement and suppression co-exist in A1 and PT. In line with the crossmodal BOLD enhancement in PT, the visual induced changes in activation patterns were not significantly affected by modality-specific attention.

Taken together these results suggest that salient visual looming stimuli affected auditory cortices irrespective of the direction of modality-specific attention in both unisensory and audiovisual contexts. Further, the surface projections of the shape parameters of the BOLD response profile at

the group level revealed a patchy topography both for visually-induced deactivations in a unisensory

context (Figure 2B left) and for visual-induced modulations of the auditory response (Figure 3B left).

Hence, we next investigated whether visual stimuli influenced activation patterns in auditory cortices

in a similar patchy topography during unisensory visual and audiovisual contexts. For this, we defined

a GLM individually for each subject that used the 'constant' or 'linear' shape parameters from the

visual deactivations (i.e. [V-Fix]) for each vertex to predict the 'constant', and respectively 'linear',

shape parameters for the visually-induced crossmodal modulation (i.e. [AV-A]) over vertices (again

in a cross-validated fashion). We visualized the results of a significant regression in the form of raster

plots (Figure 3B right). These raster plots show the laminar BOLD response profile for the difference

[AV-A] across vertices, sorted according to their BOLD response profile for [V-fix].

Auditory and visual responses relative to fixation

A   BOLD-response profiles

B   Surface projections

Figure 2: Auditory and visual responses relative to fixation

A. BOLD response profiles: **Row 1 and 3:** The BOLD response (i.e. B parameters, across subjects' mean ± SEM) for auditory and visual looming stimuli averaged over auditory and visual attention in A1, PT, V1, V2/3 is shown as a function of percentage cortical depth. WM: white matter; GM: grey matter; CSF: cerebrospinal fluid. Percentage cortical depth is

208 indicated by the small numbers and colour coded in red. **Rows 2 and 4:** Across subjects' mean

209 (± SEM) and violin plot of the participants' shape parameter estimates that characterize the

210 mean (C: constant) and linear increase (L: linear) of the laminar BOLD response profile. P

211 values pertain to an exact one-sided sign-permutation test at the group level. Values in bold

212 are significant. n = 11.

213 B. Surface projections: Across subject' mean of the 'constant' (row i) and 'linear' increase (row

214 ii) shape parameter estimates of the laminar BOLD response profile for auditory and visual

215 looming stimuli (averaged over auditory and visual attention) are projected on an inflated

216 group mean surface to show auditory and visual regions of the left hemisphere. A1 and V1

217 are delineated by black solid lines, PT and V2/3 by dashed lines. For visualization purposes

218 only: i. borders between visual-induced activations and deactivations (white dashed lines)

219 were defined based on visual inspection; ii. surface projections were smoothed (FWHM = 1.5

220 mm); iii. values are also presented for vertices for which data were not available from all

221 subjects and which were therefore not included in our formal statistical analysis. Grey areas

222 denote vertices with no available data for any subject.

223 **Table 2**

224 Effects of the cross-modal modulation on the laminar BOLD response and decoding accuracy profiles

A) BOLD profile		Constant				Linear			
Contrast	ROI	mean	STD	p value	95% CI effect size	mean	STD	p value	95% CI effect size
$[AV-A]_{att A, att V}$									
	A1	-0.04	(0.35)	0.691	[-0.75 - 0.50]	-0.01	(0.06)	0.542	[-0.92 - 0.41]
	PT	0.09	(0.34)	0.434	[-0.34 - 0.80]	0.09	(0.10)	**0.014**	[0.30 - 1.40]
$[AV-V]_{att A, att V}$									
	V1	-0.06	(0.38)	0.610	[-0.83 - 0.55]	-0.02	(0.05)	0.179	[-0.85 - 0.18]
	V2/3	-0.09	(0.34)	0.404	[-1.00 - 0.31]	-0.02	(0.06)	0.231	[-0.89 - 0.27]

B) Accuracy profile		Constant				Linear			
Classification	ROI	mean	STD	p value	95% CI effect size	mean	STD	p value	95% CI effect size
$[AV \text{ vs } A]_{att A, att V}$									
	A1	0.12	(0.10)	**0.005**	[0.10 - 2.20]	-0.01	(0.03)	0.387	[-0.84 - 0.34]
	PT	0.17	(0.11)	**<.001**	[0.42 - 2.38]	0.02	(0.02)	**0.007**	[0.05 - 1.70]
$[AV \text{ vs } V]_{att A, att V}$									
	V1	0.00	(0.05)	0.757	[-0.51 - 0.67]	0.00	(0.02)	0.780	[-0.59 - 0.85]
	V2/3	0.02	(0.08)	0.469	[-0.36 - 0.76]	-0.01	(0.02)	0.465	[-0.82 - 0.40]

C) Stimulation dependent crossmodal modulation		Constant				Linear			
	ROI	mean	STD	p value	95% CI effect size	mean	STD	p value	95% CI effect size
$[AV-A]_{att A, att V} = f([V-Fix])$									
	A1	0.00	(0.03)	0.987	[-0.70 - 0.65]	0.02	(0.02)	**0.016**	[0.22 - 1.67]
	PT	0.12	(0.12)	**<.001**	[0.61 - 1.63]	0.06	(0.11)	0.080	[0.02 - 0.96]
$[AV-V]_{att A, att V} = f([A-Fix])$									
	V1	0.01	(0.03)	0.384	[-0.37 - 0.83]	0.00	(0.02)	0.681	[-0.46 - 0.71]
	V2/3	0.01	(0.03)	0.606	[-0.46 - 0.74]	0.00	(0.03)	0.884	[-0.70 - 0.57]

226 Notes: p-values are based on an exact two-sided sign permutation test. p-values < 0.05 are indicated in bold. n = 11

In A1 and PT, a vertex's shape profile for visual deactivations significantly predicted its profile for cross-modal modulation (see Table 2C) suggesting similar patchy topographies for visual deactivations and crossmodal modulation. In PT, the constant parameter for [V-Fix] predicted the constant for [AV-A]: the less a vertex is deactivated in PT by visual stimuli in a unisensory context, the more it shows a visually-induced enhancement of the response to a concurrent auditory stimulus (see Table 2C and Figure 3Bi right). Even more interestingly, in A1 the linear shape parameter for [V-Fix] predicted the linear shape parameter of [AV-A] (see Table 2C): vertices with less deactivations in deeper relative to superficial laminae in unisensory visual context showed a robust crossmodal enhancement that was most pronounced in deeper laminae (Figure 3Bii right and Table 2C). These results strongly suggest that visual stimuli engender a BOLD response in primary auditory cortex with a patchy topography, similar for unisensory and audiovisual contexts.

In contrast to these multisensory influences, attentional modulation was greatest at the cortical surface in both A1 and PT for the regional BOLD response (i.e. significant positive linear effect in A1 and PT, Figure 4A 1^{rst} column and Table 3A). Surprisingly, pattern classifiers did not successfully discriminate between the auditory and visual attention conditions, particularly in deep laminae of A1 (Figure 4A 2^{nd} column and Table 3B). Additional control analyses, with different pre-processing such as various forms of image and feature scaling, or no scaling, replicated this chance classification performance, so that it was not specific to our particular parameter choices. Discriminating between activation patterns may be limited, if modality-specific attention predominantly amplifies and scales the BOLD response for A, V or AV stimuli whilst preserving the activation patterns, whereas the crossmodal influences impacts the activation patterns.

To investigate this explanation further, we quantified and compared the similarity in A1 between activation patterns of auditory vs. visual attention conditions (averaging over i. A_Att_V vs. A_Att_A; and ii. AV_Att_V vs. AV_Att_A) and of auditory vs. audiovisual conditions (averaging over iii. A_Att_A

vs. AV_Att$_A$; and iv. A_Att$_V$ vs. AV_Att$_V$) using the Spearman correlation coefficient computed over vertices. The average similarity between auditory and visual attention conditions was indeed greater than the average similarity between A and AV conditions (two-sided exact sign permutation test on Fisher transformed correlation coefficients, p = 0.008). These results suggest that auditory attention mainly scales the BOLD response in A1 whilst preserving the activation pattern, whereas visual stimuli alter the activation pattern in A1.

In summary, we observed distinct laminar BOLD response profiles and activation patterns for multisensory and attentional influences in auditory cortices. Visual stimuli induced deactivations with a constant profile both in A1 and PT. Likewise, they influenced the activation pattern in auditory cortices similarly across all laminae (i.e. constant effect for decoding accuracy) in audiovisual context leading to successful pattern decoding accuracy for [AV vs. A] even in deeper laminae. In fact, in A1 and PT visual inputs altered the activation pattern with a similar patchy topography when presented alone (i.e. visual-induced deactivation) or together with an auditory stimulus (i.e. crossmodal modulation).

By contrast, modality-specific attention amplified the mean BOLD-signal within A1 and PT mainly at the cortical surface (i.e. significant linear parameter), yet had little impact on the activation pattern across vertices, so that the discrimination between activation patterns for auditory and visual attention was not better than chance.

Visual cortices

In V1 and V2/3 visual stimuli induced activations that increased towards the cortical surface as expected for GE-EPI BOLD (i.e. significant constant and linear effect, see Table 1; (Markuerkiaga et al., 2016). Consistent with previous research (Shmuel et al., 2002; Silver et al., 2008; Tootell et al., 1998), the centrally presented visual looming stimuli activated predominantly areas representing the centre of the visual field with response suppressions (i.e. negative BOLD) in adjacent areas (see

dashed white lines in 3rd column of Figure 2B). By contrast, auditory deactivations were not confined

to the central field that was activated by visual stimuli. Instead, they were particularly pronounced in

the rostral part of the calcarine sulcus representing more eccentric positions of the visual field (Figure

2B 4th column and Figure S1).

Furthermore, auditory looming stimuli induced deactivations that increased in absolute magnitude

towards the cortical surface (i.e. significant negative linear effect, Fig. 2A). Again as in auditory

areas, these auditory deactivations were not significantly modulated by modality-specific attention

(i.e. [Att_V-Att_A] for auditory stimuli: $p > 0.15$ for 'constant' and 'linear' shape parameters in V1 and

V2/3).

Despite the pronounced deactivations elicited by unisensory auditory stimuli, we did not observe any

significant cross-modal modulation of the regional BOLD response in V1 and V2/3 (see Table 2 and

figure 3A). Likewise, the support vector classifiers were not able to discriminate between activation

patterns for [AV vs. V] stimuli better than chance. In summary, the effect of auditory stimuli on visual

cortices was abolished under concurrent visual stimulation both in terms of regional BOLD response

and activation patterns (Figure 3A, 3rd and 4th columns, Table 2).

Crossmodal modulation

A Laminar profiles

B Surface projections and raster plots

Figure 3: Cross-modal modulation

A. Laminar profiles: **Row 1 and 3:** The BOLD response and decoding accuracy (across subjects'

mean ± SEM) for [AV-A] in A1 and PT as well as for [AV-V] in V1 and V2/3 is shown as a

function of percentage cortical depth pooled (i.e. averaged) over auditory and visual attention. WM: white matter, GM: grey matter, CSF: cerebrospinal fluid. Percentage cortical depth is indicated by the small numbers and colour coded in red. **Rows 2 and 4:** Across subjects' mean (± SEM) and violin plot of participants' shape parameter estimates that characterize the mean (C: constant) and linear increase (L: linear) of the laminar BOLD response and decoding accuracy profiles. P values pertain to an exact two-sided sign-permutation test at the group level. Values in bold are significant. n = 11.

B. Surface projections and raster plots: **Lef**t: Across subject' mean of the 'constant' (row i) and 'linear increase' (row ii) shape parameter estimates of the laminar BOLD response profile for [AV-A] in A1 and PT (averaged over auditory and visual attention) are projected on an inflated group mean surface to show auditory regions of the left hemisphere. A1 and PT are delineated by black solid and dashed lines. For visualization purposes only: i. surface projections were smoothed (FWHM = 1.5 mm); ii. values are also presented for vertices for which data were not available from all subjects and which were therefore not included in our formal statistical analysis. Grey areas denote vertices with no available data for any subject. **Right: Row i**. The raster plot shows the statistical relationship between the 'constant' shape parameters for the visual evoked response [V-Fix]$_{AttA, AttV}$ and the crossmodal modulation [AV-A] in PT. Each raster plot shows the laminar profiles (along abscissa) of the 'predicting contrast' [V-Fix]$_{AttA, AttV}$ and of the 'predicted contrast' [AV-A] for the vertices sorted along the ordinate according to the value of the 'constant' shape parameter for [V-Fix]$_{AttA, AttV}$. PT vertices with less deactivations across laminae for [V-Fix]$_{AttA, AttV}$ are associated with greater crossmodal enhancement [AV-A]. **Row ii**. The raster plot shows the statistical relationship between the 'linear slope' shape parameters for the visual evoked response [V-Fix]$_{AttA, AttV}$ and the crossmodal modulation [AV-A] in A1. Each raster plot shows the laminar profiles (along abscissa) of the 'predicting contrast' [V-Fix]$_{AttA, AttV}$ and of the 'predicted contrast' [AV-A] for the vertices sorted along the ordinate according to the value of the linear shape

parameter for [V-Fix]$_{AttA, AttV}$. A1 vertices with less deactivations in deeper laminae for [V-

Fix] are associated with greater crossmodal enhancement [AV-A] in deeper laminae.

For visualization purposes all raster plots were smoothed along the vertical axis (FWHM =

1% of the number of data bins). n = 11.

With respect to attentional modulation, we observed no significant differences in mean BOLD

responses in V1 or V2/3 for visual vs. auditory attention conditions (Table 3A). Yet, the classifier

was able to discriminate between patterns for visual and auditory attention conditions successfully

across all cortical depth surfaces with highest performance in superficial laminae in V1 and V2/3 (i.e.

significant constant term in V1 and V2/3 and linear term in V2/3, Table 3B, Figure 4A 3rd and 4th

columns). The surface projection of shape parameters of the BOLD response profile at the group level

explains this discrepancy between the attentional effects on regional BOLD response and activation

patterns (Figure 4Bii). Consistent with mechanisms of contrast enhancement (Bressler et al., 2013;

Müller and Kleinschmidt, 2004; Smith et al., 2000; Tootell et al., 1998), attending to vision relative

to audition increased responses in regions that were activated by visual looming stimuli, yet

suppressed the deactivations in the surrounding regions that were deactivated by visual stimuli

thereby cancelling out global attentional effects for the regional BOLD response (Figure 4Bii, Note

the white lines indicate the border between visual activation and deactivation from figure 2B).

In summary, auditory looming induced widespread deactivations in visual cortices that were maximal

at the cortical surface and extended into regions that represent more peripheral visual fields (Figure

2B). By contrast, attention to vision predominantly increased BOLD response in central parts that

were activated by visual stimuli and suppressed BOLD responses in the periphery that were

deactivated by visual stimuli.

Table 3

Effects of the attentional modulation (irrespective of stimulus type) on the laminar BOLD response and decoding accuracy profiles

A) BOLD profile		Constant			95% CI effect size			Linear			95% CI effect size		
Contrast	ROI	mean	STD	p value				mean	STD	p value			
$[Att_A - Att_V]_{A, V, AV}$													
	A1	0.14	(0.24)	0.101	[-0.24	-	1.19]	0.04	(0.04)	**0.013**	[-0.05	-	1.81]
	PT	0.21	(0.15)	**<.001**	[0.53	-	2.05]	0.06	(0.06)	**0.008**	[0.06	-	1.82]
$[Att_V - Att_A]_{A,V,AV}$													
	V1	0.11	(0.33)	0.312	[-0.42	-	0.83]	0.02	(0.06)	0.364	[-0.49	-	0.82]
	V2/3	0.00	(0.24)	0.957	[-0.83	-	0.55]	0.00	(0.06)	0.883	[-0.59	-	0.67]

B) Decoding Accuracy profile		Constant			95% CI effect size			Linear			95% CI effect size		
Classification	ROI	mean	STD	p value				mean	STD	p value			
$[Att_A$ vs $Att_V]_{A,V,AV}$													
	A1	0.02	(0.04)	0.228	[-0.32	-	1.08]	0.01	(0.02)	0.196	[-0.19	-	0.93]
	PT	0.04	(0.07)	0.072	[0.11	-	1.27]	0.02	(0.03)	0.064	[-0.18	-	1.11]
	V1	0.06	(0.08)	**0.018**	[0.03	-	1.14]	0.01	(0.01)	0.213	[-0.25	-	0.83]
	V2/3	0.11	(0.09)	**<.001**	[0.70	-	1.80]	0.01	(0.01)	**0.011**	[0.21	-	1.61]

353 Notes: p-values are based on an exact two-sided sign permutation test. p-values < 0.05 are indicated in bold. n = 11

Attentional modulation

A   Laminar profiles

B   Surface projections

Figure 4: Attentional modulation

A. Laminar profiles: **Row 1 and 3:** The BOLD response and decoding accuracy (across subjects'

mean ± SEM) for attentional modulation ([Att$_A$-Att$_V$] in A1 and PT as well as for [Att$_V$-Att$_A$]

in V1 and V2/3) is shown as a function of percentage cortical depth pooled (i.e. averaged)

over auditory, visual and audiovisual looming stimuli. WM: white matter, GM: grey matter,

CSF: cerebrospinal fluid. Percentage cortical depth is indicated by the small numbers and

colour coded in red. **Rows 2 and 4:** Across subjects' mean (± SEM) and violin plot of the

participants' shape parameter estimates that characterize the mean (C: constant) and linear

increase (L: linear) of the laminar BOLD response and decoding accuracy profiles. P values

pertain to an exact two-sided sign-permutation test at the group level. Values in bold are

significant. n = 11.

B. Surface projections : Across subject' mean of the 'constant' shape parameter estimates for

attentional modulation $[Att_A-Att_V]_{A,V,AV}$ in A1 and PT (i: left) and in V1 and V2/3 (ii: right)

are projected on an inflated group mean surface of the left hemisphere. A1 and V1 are

delineated by black solid lines. PT and V2/3 are delineated by dashed lines. For visualization

purposes only: i. surface projections were smoothed (FWHM = 1.5 mm); ii. values are also

presented for vertices for which data were not available from all subjects and which were

therefore not included in our formal statistical analysis; iii borders between visual-induced

activations and deactivations (white dashed lines on the right) are reported here from figure

2B. Grey areas denote vertices with no available data for any subject.

DISCUSSION

The current high-resolution 7T fMRI study revealed distinct depth-dependent BOLD response profiles and patterns for multisensory and attentional influences in early sensory cortices.

Distinct laminar profiles and activation patterns for multisensory and attentional influences in auditory cortices

Visual looming suppressed activations in auditory cortices (Laurienti et al., 2002; Leitão et al., 2013; Mozolic et al., 2008), but enhanced the response to concurrent auditory looming (Werner and Noppeney, 2011, 2010). In other words, intersensory competition for purely visual stimuli turned into cooperative interactions for audiovisual stimuli. While the visual-induced response suppression was constant across cortical depth in A1 and PT, the crossmodal enhancement was observed mainly in the caudal parts of PT at the cortical surface (Kayser et al., 2007). Yet, at a finer spatial resolution, multivariate analyses revealed significant differences between audiovisual and auditory activation patterns across all laminae in both A1 and PT. As shown in the surface projections, visual looming enhanced and suppressed auditory responses in adjacent patches consistent with neurophysiological results in non-human primates (see figure 3 and (Kayser et al., 2008). Critically, a vertex's response profile to purely visual stimuli significantly predicted its laminar profile for crossmodal modulation (see Table 2C, Figure 3Bi, ii). In A1, vertices whose activations were only weakly suppressed by visual looming in deeper laminae showed a greater visual-induced response enhancement for audiovisual looming again in deeper laminae (Figure 3Bii). These results suggest that visual looming influences activations in A1 mainly in deeper layers with similar patchy topographies in unisensory and audiovisual contexts.

By contrast, modality-specific attention affected auditory and visual evoked responses predominantly at the cortical surface (see significant linear term, figure 4A, column 1). Furthermore, we could not discriminate between activation patterns for auditory and visual attention (see figure 4A, column 2).

suggesting that modality-specific attention amplified the mean BOLD response rather than altered

the fine-grained activation pattern.

The large attentional effects in superficial laminae dovetail nicely with previous neurophysiological

research, which located effects of modality-specific attention in supragranular layers of auditory

cortices (De Martino et al., 2015a; Lakatos et al., 2009). Because the BOLD response measured at

superficial laminae includes contributions from deeper laminae (Markuerkiaga et al., 2016), we

cannot unequivocally attribute the attentional influences at the cortical surface solely to superficial

layers as neural origin. Critically, however, the laminar profiles for attentional effects were distinct

from those for multisensory (i.e. crossmodal) effects in auditory cortices. This dissociation in laminar

profiles in the same territories cannot be explained by vascular effects that limit the laminar specificity

of the BOLD response, but strongly implies that multisensory and attentional mechanisms regulate

information flow in auditory cortices via partly distinct neural circuitries (De Martino et al., 2015a;

Lakatos et al., 2009). Further, the constant laminar profiles for crossmodal influences in auditory

cortices suggest that visual signals influence auditory cortices mainly via infragranular cortices – a

finding that converges with a recent neurophysiological study in mouse that likewise highlighted deep

layer 6 as the key locus for visual influences on auditory cortex (Morrill and Hasenstaub, 2018).

Anatomical studies in monkeys have previously suggested that visual inputs can influence auditory

cortices via three routes (Ghazanfar and Schroeder, 2006; Schroeder et al., 2003; Smiley and Falchier,

2009): i. non-specific thalamic afferents terminating mainly in supragranular layers, ii. direct

connectivity between sensory areas terminating in all layers (Falchier et al., 2002; Rockland and

Ojima, 2003) and iii. feed-back connections from higher order association cortices terminating in

infra and supragranular layers (Felleman and Van Essen, 1991). Multisensory influences in our study

may thus rely more on direct connectivity between auditory and visual cortices leading to a greater

involvement of infragranular processes in comparison to attentional influences.

Common laminar profiles, but distinct activation patterns for multisensory and attentional

influences in visual cortices

In visual cortices, visual and auditory stimuli evoked responses that were maximal in absolute amplitude at the cortical surface, yet differed in their activation pattern. Visual looming stimuli induced activations in areas representing the centre of the visual field and deactivations in surrounding areas representing the periphery (Shmuel et al., 2002). In contrast to this visual 'centre-surround' pattern, auditory stimuli induced widespread deactivations (Laurienti et al., 2002; Leitão et al., 2013; Mozolic et al., 2008) particularly in peripheral visual field representations known to have denser direct fibre connections to auditory cortices (Falchier et al., 2002; Rockland and Ojima, 2003)..

Thus, our study revealed three types of deactivations. In visual cortices, both visual and auditory stimuli induced deactivations that were most evident at the cortical surface, as previously reported for deactivations to ipsilateral visual stimuli ((Fracasso et al., 2018), though see (Goense et al., 2012)). In auditory cortices, the visual induced deactivations were constant across cortical depth. These differences in laminar profiles or BOLD patterns between visual and auditory cortices may reflect differences in neural mechanisms. For instance, neurophysiological studies in rodents have shown asymmetries in multisensory influences between auditory and visual cortices, reporting robust auditory induced inhibition in layers 2/3 in visual cortices but not vice versa in auditory cortices (Ibrahim et al., 2016; Iurilli et al., 2012).

While we observed neither crossmodal nor attentional effects on the regional BOLD response profiles in visual cortices, multivariate pattern analyses indicated that modality-specific attention altered the activation patterns at a sub-regional spatial resolution (figure 4A): Attention to vision amplified activations in areas representing the central visual field and suppressed activations in areas representing peripheral visual fields (see figure 4 B).

The differences in activation patterns for multisensory and attentional influences indicate that the

impact of purely auditory looming (Maier et al., 2008, 2004) on visual cortices cannot solely be

attributed to a withdrawal of attentional resources from vision. Instead, purely auditory looming may

inhibit activations in visual cortices via direct connectivity between auditory and visual areas (Bizley

et al., 2007; Budinger et al., 2006; Campi et al., 2010; Falchier et al., 2002; Ibrahim et al., 2016;

Rockland and Ojima, 2003). In line with this conjecture, recent research in rodents has shown that

auditory stimuli elicit strong inhibition in supragranular layers 2/3 in primary visual areas via direct

connectivity from auditory cortices and translaminar inhibitory circuits – though the two studies

disagreed in whether layer 5 or layer 1 (Ibrahim et al., 2016; Iurilli et al., 2012) were the primary

targets of auditory inputs.

CONCLUSIONS

Using submillimetre resolution fMRI at 7T, we resolved BOLD response and activation patterns of

multisensory and attentional influences across cortical depth in early sensory cortices. In visual

cortices auditory stimulation induced widespread inhibition, while attention to vision enhanced

central but suppressed peripheral field representations. In auditory cortices competitive and

cooperative multisensory interactions were stronger deep within the cortex, while attentional

influences were greatest at the cortical surface. The distinctiveness of these depth-dependent

activation profiles and patterns suggests that multisensory and attentional mechanisms control

information flow via partly distinct neural circuitries.

MATERIAL AND METHODS

All procedures were approved by the Ethics Committee of the University of Leipzig.

Participants

Thirteen healthy German native speakers (6 females; 7 males; mean age: 24.8 years, standard

deviation: 1.5 years, range: 22-27 years) gave written informed consent to participate in this fMRI

study. Participants reported no history of psychiatric or neurological disorders, and no current use of

any psychoactive medications. All had normal or corrected to normal vision and reported normal

hearing. Two of those subjects did not complete all fMRI sessions and were therefore not included in

the analysis.

Looming stimuli and targets

The visual looming stimulus (500 ms duration) was a sequence of 1 to 4 radially expanding white

annuli (Figure 1B). Each of the four annuli started in the centre of the screen with a diameter of 0.25°

visual angle and expanded at a radial speed of 15.52° visual angle per second to a maximum radius

of 3.88° visual angle which covered the entire screen visible in the scanner bore along the vertical

dimension. During the 500 ms sequence, an additional annulus started in the centre every 125 ms.

Given the retinotopic organization of early visual cortices (Wandell et al., 2007) these parameters

ensured that visual looming elicited relatively widespread activations in visual cortices.

The auditory looming stimulus (500 ms duration) was composed of several pure tones (55, 110, 150,

220, 330, 380, 440, 660, 880, 990, 1110, 1500, 2500, 3000 Hz) that increased in frequency and

amplitude over the entire 500 ms duration. The amplitude of the sound was exponentially modulated

with the amplitude at time t given by $A_t = A_0 e^{0.68t} - 1$ and A_0 being the initial baseline amplitude for

each pure tone. The frequency of each pure tone component was increased by two thirds of its original

base frequency over the 500 ms. Given the tonotopic organization of primary auditory cortices

(Formisano et al., 2003) this mixture of sound frequencies ensured widespread activations in auditory

cortices.

Audiovisual stimuli were generated by presenting the auditory and visual stimuli in synchrony

together.

The visual target was a grey dot presented for 50 ms within a circular area that increased over the 500

ms duration consistent with the auditory, visual or audiovisual looming stimulus (Figure 1B).

Auditory targets were a single 440 Hz pure tone that lasted for 100 ms. The size of the visual target

and the sound amplitude of the auditory target were adjusted in a subject-specific fashion to match

the approximately target detection performance across sensory modalities and then held constant

across the entire experiment. This was done using the method of constant stimuli in a brief

'psychometric function experiment', performed prior to the fMRI study inside the scanner and with

scanner noise present.

Experimental design

The experiment conformed to a 3 (stimulus modality: auditory, visual, audiovisual) X 2 (attended

modality: auditory, visual) factorial design (Figure 1A). In an intersensory selective attention

paradigm, participants were presented with blocks of audio (A), visual (V) or audiovisual (AV)

looming stimuli as highly salient stimuli that drive bottom-up attention and elicit crossmodal

deactivations (Leitão et al., 2013). Brief auditory (i.e. weak and brief beep) and visual (i.e. small and

brief grey dot) targets were presented interspersed in all stimulation blocks irrespective of the sensory

modality of the looming stimuli or the focus of modality-specific attention (e.g. A and V targets were

presented during a unisensory auditory looming block and auditory attention). The visual grey dot

could be presented at any time at any location covered by the radially expanding white annuli until

they reach a radius of maximal 3.88° visual angle. Likewise, the auditory target is presented at random

time interspersed in the looming sound. Consistent with previous research on modality-specific

attention (Lakatos et al., 2009; Werner and Noppeney, 2011, 2010) these additional targets were

included to ensure that the effects of attention could not be attributed to top-down effects associated

with explicit responses.

During auditory attention (Att$_A$), participants were instructed to respond selectively to all auditory

targets and ignore visual ones. During visual attention (Att$_V$) participants were instructed to respond

selectively to all visual targets, i.e. brief grey dots and ignore auditory targets. In this way, we matched

the auditory (i.e. attention to frequency) and visual attention (i.e. attention to targets in spatial field)

tasks to the tonotopic and retinotopic organizational principles of primary auditory and visual areas

as well as the specifics of auditory and visual looming bottom-up salient stimuli.

Experimental procedures

Auditory, visual or audiovisual looming stimuli were presented in 33 seconds blocks of 50 stimuli

(stimulus duration: 500 ms, inter-stimuli interval: 160 ms, stimulus onset asynchrony: 660 ms)

(Figure 1B). Activation blocks were preceded by 16 seconds fixation baseline periods. We opted for

a block design as they have the highest design efficiency despite their potential drawbacks from a

cognitive perspective (Friston et al., 1999). Throughout the entire experiment, participants fixated a

cross in the centre of the black screen. The fixation cross was white during the activation conditions

and blue during the fixation baseline periods. A blue letter ('V' or 'A') presented 3 seconds prior to

the activation blocks instructed participants to direct their attention to the visual or auditory modality.

During visual attention blocks, participants were instructed to attend generally to the visual modality

and respond selectively to visual targets. Conversely, during auditory attention blocks, participants

had to attend generally to the auditory modality and respond selectively to auditory targets. Sixteen

targets of each modality were presented for each condition. To maintain observer's attention and

minimize the effect of target stimuli on measured stimulus-induced activations, targets were more

frequently presented at the end of the block (i.e. two thirds of the blocks included one auditory target

and one visual target, the last third of blocks included 2 targets of each type). The final block of each

run was followed by a fixation baseline period of 20 seconds. The order of blocks was pseudo

randomized across participants. Each fMRI run included 18 blocks (3 blocks for each condition in

our 2 X 3 design) and lasted 15 minutes. There were 4 runs per participant yielding 12 blocks per

condition per participant.

Experimental set up

Stimuli were presented using Presentation (v17.6, Neurobehavioral system, www.neurobs.com) on a

PC desktop (Windows XP). Visual stimuli were back-projected onto a plexiglas screen using an LCD

projector (60 Hz) visible to the participant through a mirror mounted on the MR head coil. The MR

head coil enabled a visual field with a width of 10.35° and height of 7.76° visual angle. Auditory

stimuli were presented using MR-compatible headphones (MR-Confon; HP-SI01 adapted for the

head-coil). Behavioral responses were collected using a MR-compatible button device connected to

the stimulus computer. The code to run the fMRI experiment is available at https://github.com/Remi-

Gau/AV_Attention-Presentation_code-fMRI/releases/tag/v0.0.1.

Behavioral data analysis

The response window extended from the target onset and for either 2.5 seconds or until the beginning

of the next target. Any recorded response outside of that window was coded as an extra-response (i.e.

to be modelled in the fMRI design matrix). The percentage of auditory (or visual) targets that

participants responded to in a particular condition were entered as dependent variables into a 2 (target

modality: auditory vs. visual) X 2 (attended modality: auditory vs. visual) X 3 (stimulus block

modality: auditory, visual, audio-visual) repeated measures ANOVA (results were corrected for non-

sphericity using Greenhouse-Geiser correction). Alpha was set to 0.05. This analysis was done with

IBM SPSS Statistics, version 17.0 (IBM, Armonk, NY, USA)

MRI Data acquisition

All experiments were performed on a Siemens 7T MAGNETOM MRI scanner (Siemens Healthineers, Erlangen, Germany) equipped with an SC72 gradient coil (Siemens Healthineers). For radio-frequency transmission and reception, a single-channel-transmit/32-channel-receive phased array head coil (Nova Medical, Wilmington, MA, USA) was used. The experiment was run over 2 days with 50 minutes MRI scanning in total per day per participant.

On the first day a whole-brain MP2RAGE sequence (Marques et al., 2010) was used to acquire a quantitative T1 map and a weighted T1 image and a second inversion image to enable high-resolution segmentation and cortical laminae definition [repetition time (TR) / echo time (TE) = 5000/2.45 ms, inversion time 1 = 900 ms, flip angle 1 = 5°, inversion time 2 = 2750 ms, flip angle 2 = 3°, 240 slices, matrix = 320×320, acquisition time = 600 s, spatial resolution = $0.7\times0.7\times0.7$ mm^3 voxels, partial Fourier factor = 6/8, parallel imaging using GRAPPA with an acceleration factor of 4, field of view = 168 X 224 mm]. On the second day, we used a shorter version of that sequence with a coarser resolution only in order to position our EPI acquisition slab.

Two functional runs were acquired every day using a 2D gradient echo single-shot echo planar imaging (EPI) readout [TR/TE = 3000/25 ms, flip angle = 90º, 48 axial slices acquired in descending direction, matrix size = 256×240, slice thickness = 0.75 mm, brain coverage = 3.6 mm, spatial resolution = $0.75\times0.75\times0.75$ mm^3 voxels, partial Fourier factor = 6/8, parallel imaging using GRAPPA with an acceleration factor of 4, number of volumes = 302, acquisition time = 906 s, field of view = 192 X 180 mm] A corresponding vendor-provided homodyne online image reconstruction algorithm was used. The first six volumes in each run were automatically discarded to avoid for T1 saturation effects.

The acquisition slab was inclined and positioned to include Heschl's gyri, the posterior portions of

superior temporal gyri and the calcarine sulci. On the second day the slab was auto-aligned on the

acquisition slab of the first day. This provided us with nearly full coverage over all our four regions

of interest (see table S1).

One field-map was acquired on each day with a coverage and orientation matching that of the EPI

slab [TR/TE1/TE2 = 1500/6/7.02 ms, flip angle = 72º; 18 axial slices, matrix = 96 X 90, slice

thickness = 2 mm, spatial resolution = 2.00 X 2.00 X 2.00 mm^3 voxels, field of view: FOV= 192 X

180 mm].

MRI structural analysis

The MRI structural data were processed using the CBS toolbox (v3.0.8, Max Planck Institute for

human cognitive and brain science, Leipzig, Germany, www.nitrc.org/projects/cbs-tools/) based on

the MIPAV software package (v7.0.1, NIH, Bethesda, USA, www.mipav.cit.nih.gov, (McAuliffe et

al., 2001)) and the JIST pipeline environment(v2.0, (Landman et al., 2013; Lucas et al., 2010)). The

T1 map was coregistered to the Montreal's Neurological Institute brain space (rigid body registration

and normalized mutual information as a cost function) and resampled to an isometric resolution of

0.4 mm. The image was segmented fully automatically, and the cortical boundary surfaces were

reconstructed (Bazin et al., 2014). Between these surfaces, we computed 6 intracortical surfaces that

defined cortical laminae with the equivolume approach (Waehnert et al., 2014) (Figure 1 C). It is

important to note that the term 'lamina' used in this communication does not directly refer to

cytoarchitectonically defined cortical layers: for example, the variation across cortical regions of the

relative thickness between layers was not modelled here (Nieuwenhuys, 2013). However, it should

be mentioned that this method provides biologically plausible cortical contours (Waehnert et al.,

2014).

Definition of Region of Interest

We defined four regions of interest (ROI) as follows (see Table S2 for ROI effective sizes):

A. Several studies have shown that the core of primary auditory cortex (A1) because of its characteristic myelination profile can be localized using its low T1 intensity (De Martino et al., 2015b; Dick et al., 2012; Hackett et al., 2001). We therefore defined it by manual delineation (RG, UN) that was guided by anatomical structure (Heschl's gyrus) and informed by the quantitative T1 map that was sampled at mid-depth of the cortex and projected on the corresponding inflated surface. The final bilateral A1 region was defined as the union of the two delineations (RG, UN).

B. The planum temporale (PT) was defined bilaterally using the area 41-42 of the brainnetome atlas (Fan et al., 2016) (http://atlas.brainnetome.org/). The full probability maps of this ROI and its anatomical neighbours were inverse-normalized into native space using the deformation field given by the SPM segmentation of the T1 weighted structural scan. The probability map was then re-gridded to an isometric resolution of 0.4 mm, sampled at mid-depth of the cortex and projected on the corresponding surface. Vertices were included as part of PT if their probability exceeded 40% unless i. they were already defined as being part of A1 or ii. the sum of the probabilities of the neighbouring regions exceeded that of the area 41-42.

C. Primary Visual area (V1) and high order visual areas (V2 and V3) were defined bilaterally in a similar fashion as PT. We used the probabilistic retinotopic maps of the ROIs for V1, V2/3 (Wang et al., 2015) (http://scholar.princeton.edu/napl/resources). In this case vertices were included if their probability exceeded 10% unless the sum of the probabilities of the neighbouring regions exceeded that of the area of interest.

We obtained > 95% coverage for A1, PT and V1 and > 82% coverage for V2/3. (For the number of vertices and percentage coverage across subjects, see supplementary Table S2).

fMRI preprocessing

The fMRI data were pre-processed and analyzed using statistical parametric mapping (SPM12 – v6685; Wellcome Center for Neuroimaging, London, UK; www.fil.ion.ucl.ac.uk/spm) running on matlab (Mathworks). The fieldmaps were co-registered to the first functional scan of the first run (i.e. rigid-body transformations optimized using normalized mutual information as cost function) and a voxel displacement map was then created. Functional scans from each participant were realigned and unwarped using the first scan as a reference (interpolation: 4^{th} degree b-spline, unwarping done using the voxel displacement map of the corresponding day for each run).

Functional to anatomical coregistration

We co-registered the mean EPI image to the pre-processed T1 map (i.e. rigid-body transformations optimized using normalised mutual information as cost function). We assessed the accuracy of co-registration using FSLview (FSL 5.0 - Analysis Group, FMRIB, Oxford, UK, https://fsl.fmrib.ox.ac.uk/fsl/fslwiki/FSL) by flipping back and forth between the mean EPI and the T1 map images in all the ROIs. If the co-registration was not sufficiently precise, the mean EPI was initially co-registered manually and the co-registration repeated until anatomical structures of the mean EPI and the T1 map were precisely realigned. The transformation matrix of the co-registration was later applied to the beta images from the fMRI general linear model (see 'fMRI statistical analysis').

fMRI statistical analysis

At the first (i.e subject) level, we performed a mass univariate analysis with a linear regression at each voxel, using generalized least squares with a global approximate AR(1) autocorrelation model and a drift fit with discrete cosine transform basis (128 s cut-off). We modelled the fMRI experiment with a mixed block-event related model with regressors entered into the run-specific design matrix after convolving the onsets of each block or event with a canonical hemodynamic response function

(HRF) and its first temporal derivative. More specifically, we modelled each of the three looming

blocks in each run for each of the 6 conditions in our 3 (visual, auditory, audiovisual) X 2 (auditory

vs. visual attention) factorial design separately. Hence for each run the statistical model included the

following regressors: 3 block regressors for each of the 6 conditions (A_Att$_A$, A_Att$_V$, V_Att$_A$,

V_Att$_V$, AV_Att$_A$, AV_Att$_V$), 4 event-related regressors for each target type under each attention

condition (i.e. TargetA_Att$_A$, TargetA_Att$_V$, TargetV_Att$_A$, TargetV_Att$_v$) and one event-related

regressor modelling any additional responses that participants produced in the absence of targets. We

modelled the targets and additional responses as independent regressors in our first (i.e. subject) level

general linear model (GLM) to minimize confounding effects of perceptual decision making,

response selection and motor preparation on the activations reported for the A, V, AV looming

activation blocks that are the focus of this communication.

Nuisance covariates included the realignment parameters to account for residual motion artefacts.

All subsequent analyses included only the beta images pertaining to the HRF of the activation blocks

in our 2 (modality specific attention) X 3 (stimulation modality) design: 3 blocks per condition per

run X 6 conditions X 4 runs = 72 beta images per subject.

To minimize the possibility that attentional lapses reduced the sensitivity of our analysis we also

repeated our neuroimaging analysis selectively for blocks that included neither misses (i.e. missed

responses to targets of the attended modality) nor false alarms (i.e. responses to targets from the

unattended modality). This control analysis basically showed similar results similar to those from the

main analysis that is reported in this manuscript.

The beta images were then co-registered to the MRI structural by applying the transformation matrix

obtained from the mean EPI image co-registration and up-sampled to a 0.4 mm isometric resolution

(4th degree b-spline interpolation). Finally, we sampled each beta image along vertices defined by the

normal to the mid-cortical surface at the depths defined by the 6 intra-cortical surfaces from the MRI

structural analysis (tri-linear interpolation) (see 'MRI structural analysis' above, Figure 1C).

Laminar BOLD response profiles

To obtain laminar BOLD response profiles we summarized the BOLD response at each of the six cortical depth levels in terms of the median of the parameter estimates across all vertices at this cortical depth within an ROI. The median was used as a summary index because the parameter estimates distribution was skewed especially for the most superficial laminae. However, we note that a control analysis using the mean yielded basically equivalent results. We generated laminar BOLD response profiles independently for each of our 72 beta images (6 conditions X 3 blocks per condition per run X 4 runs). For each ROI and lamina, we then computed the following statistical contrasts independently for the ith block of a given condition of the jth run:

1. the activation induced by auditory stimuli relative to fixation baseline irrespective of modality-specific attention:

$$[A\text{-}Fix] = [A\_Att_A + A\_Att_V]_{i,j} / 2,$$

2. the activation induced by visual stimuli relative to fixation baseline irrespective of modality-specific attention:

$$[V\text{-}Fix] = [V\_Att_A + V\_Att_V]_{i,j} / 2,$$

3. the crossmodal enhancement irrespective of attention, i.e. specifically for auditory and visual regions:

   a. in A1 and PT: the visual-induced modulation of auditory activations irrespective of modality-specific attention:

   $$[AV - A] = [(AV\_Att_A - A\_Att_A) + (AV\_Att_V - A\_Att_V)]_{i,j} / 2,$$

   b. In V1 and V2/3: the auditory-induced modulation of visual activations irrespective of modality-specific attention:

   $$[AV - V] = [(AV\_Att_A - V\_Att_A) + (AV\_Att_V - V\_Att_V)]_{i,j} / 2,$$

4. attentional modulation irrespective of stimulus modality, i.e. specifically for auditory and visual regions

   a. A1 and PT: Modulation of stimulus responses by auditory relative to visual attention

$[Att_A - Att_V]_{A,V,AV} = [(Att_A - Att_V)_A + (Att_A - Att_V)_V + (Att - Att_V)_{AV}]_{i,j} / 3.$

b. V1 and V2/3: Modulation of stimulus responses by visual relative to auditory attention

$[Att_V - Att_A]_{A,V,AV} = [(AV\_Att_V - AV\_Att_A) + (A\_Att_V - A\_Att_A) + (V\_Att_V - V\_Att_A)]_{i,j} / 3$

In addition, we also tested for the interaction between stimulus modality and modality-specific

attention by computing the contrasts listed 1, 2 and 3 separately for each modality-specific attention

condition.

For each statistical comparison and lamina, we thus obtained 12 (i.e. 3 blocks X 4 runs) contrast

estimates per subject i.e. 12 laminar profiles. We refer to those parameter estimates that quantify the

BOLD response for a particular lamina as BOLD response parameter estimates (i.e. B parameter

estimates in the figures).

To characterize the overall shape of the laminar profile, we estimated for each subject a 2^{nd} level

general linear model (i.e. a laminar GLM) that modelled the activation in each lamina as dependent

variable by two predictors: 1) a constant term (i.e. activation mean across laminae), 2) a linear

increase across laminae (orthogonalized with respect to the mean). The parameter estimates of this

2^{nd} or laminar GLM are referred to as shape parameters (i.e. S parameter estimates in the figures).

To enable generalization to the population, we entered these shape parameters into independent exact

sign-permutation tests at the group level.

Based on our a-priori predictions (Leitão et al., 2013) that auditory stimuli induce activations in

auditory and deactivations in visual areas and that visual stimuli induce activations in visual and

deactivations in auditory areas, we employed one-sided tests to test for a positive (respective negative)

auditory or visual activations relative to fixation (i.e. the constant term for contrasts: 1,2 from above).

In all other cases two-sided tests were used. Unless otherwise stated, we report statistical results as

significant at $p < 0.05$. The bias-corrected and accelerated bootstrap 95% confidence interval of the

effect size are reported where appropriate. Given our sample size we used Hedges and Olkin's method

to correct for bias in our effect size estimation (Kelley, 2005; Lakens, 2013).

Multivariate analysis of pattern across vertices

Multivariate pattern analyses were performed a linear support vector classifier (SVC) that was trained

in a leave-one-out cross-validation scheme (LIBSVM 3.21,

https://www.csie.ntu.edu.tw/~cjlin/libsvm/, with C = 1 and mean-centred activations for each feature

separately for training and test sets).

We performed multivariate pattern analysis directly on the BOLD response patterns, i.e. the patterns

of B-parameters, independently at each of the 6 cortical depths to generate laminar profiles of

decoding accuracy for the following comparisons:

1. $[AV \text{ vs } A]_{AttA, AttV}$ (pooled over attended modality) for A1 and PT,

2. $[AV \text{ vs } V]_{AttA, AttV}$ (pooled over attended modality) for V1 and V2-3,

3. $[Att_A \text{ vs } Att_V]_{A,V,AV}$ (pooled over stimulation modality) for A1, PT, V1 and V2/3.

In line with our analysis of laminar BOLD response profiles, we modelled the laminar profiles of

decoding accuracy using a laminar GLM with a constant term and a linear increase across laminae as

predictors. These two parameter estimates were then entered into an exact sign-permutation tests at

the group level.

Raster plots: statistical relationship of BOLD response profiles between different conditions

Next we investigated whether the response profile in a vertex as characterized by their 'constant' and

'linear' shape parameters in one condition is statistically predictive of this vertex's laminar profile in

another condition or contrast. For instance, we asked whether the magnitude of visual induced

deactivations (e.g. [V-Fix] averaged over attention conditions) in a vertex predicts its crossmodal

enhancement (e.g. [AV-A] averaged over attention conditions). For this, we sorted and average the

vertices in percentile-like bins based on the value of a shape parameter (e.g. 'constant' or 'linear') for

a particular sorting contrast (e.g. [V–Fix]). This bin order was then used to sort and average the values

of the 2^{nd} contrast (e.g. [AV-A]). We entered the shape parameter value (e.g. 'constant' or 'linear')

for the sorting contrast (e.g. [V–Fix]) for each bin as a linear regressor (+ a constant) to predict the

corresponding shape parameter in each bin in the sorted contrast (e.g. [AV-A]) in a subject-specific

general linear model. To allow for generalization to the population, we entered the parameter for the

linear term into exact sign-permutation tests at the group level.

We illustrated the statistical relationships of the laminar profiles between different contrasts at the

group level in raster plots by averaging the laminar profiles for each bin across subjects of the sorting

(i.e. predicting) and the sorted (i.e. predicted) values (e.g. see Figure 3Bi right). For instance, in figure

3 Bi (right) we sorted the vertices according to [V-fix] in PT such that the constant parameter (i.e.

average BOLD response across laminae) increases from bottom to top (i.e. predicting contrast).

Unsurprisingly, the raster plot for the [V-Fix] panel as the predicting contrast thus goes from a

negative BOLD response profile (i.e. coded in blue) in the bottom rows of the panel to a positive

BOLD response profile (i.e. coded in red) in the top rows of the panel. Under the null-hypothesis the

laminar BOLD response profile for the predicting contrast (e.g. [V-fix]) in a vertex is unrelated to its

laminar BOLD response profile for the predicted contrast (e.g. [AV-A]) and we would not expect any

structure in the raster plots for the predicted contract (e.g. [AV-A], n.b. we accounted for spurious

correlation by performing all these analyses in a crossvalidated fashion). Conversely, if the shape

parameter for [V-fix] in a vertex significantly predicts its shape parameters for [AV-A], we would

expect a structured raster plot also for [AV-A].

Hence, these raster plots can reveal additional BOLD signal structure that is averaged out by the

group surface projections: while group surface projections (see next section, or Fig 3 Bi left) reveal

only spatial topographies which are consistent across participants, the raster plots reveal whether the

similarity between patterns of laminar profiles from different conditions is consistent across subjects,

even when the activation patterns themselves are not similar across subjects. In other words, for raster

plots the patterns themselves can vary across subjects, yet the similarity or covariance between

patterns can still be consistent across subjects.

Please note that both the regression model and the raster plots were computed in a leave one day out

cross-validation to avoid biases and spurious correlations between sorting and sorted contrast.

Further, we also note that the statistical results were basically equivalent without any binning (i.e.

directly entering the shape parameter values of vertices into the GLM), the binning was applied only

to enable illustration of the results in raster plots.

These regression models over vertices and raster plots were used to evaluate whether crossmodal

modulation for auditory or visual stimuli in A1, PT, V1 or V2/3 depended on the unisensory visual

or auditory response.

Intersubject registration for surface projection of group results

To visualize patterns of the laminar profile shape parameters (i.e. 'constant' or 'linear') that are

consistent across subjects on the cortical surface, we transformed the individual surface projections

into a standardized study group space using a multimodal, multi-contrast surface registration

(MMSR) approach (Tardif et al., 2015). The transformation matrix for group normalization was

computed for the level-set corresponding to the mid-cortical surface and the high-resolution T1 maps.

The T1 map was sampled radially along the cortical profile, averaged between 20% and 80% cortical

depth and smoothed tangentially (FWHM = 1.5 mm). To reduce computation time, the level-set and

the T1 map data were down-sampled to a resolution of 0.8 mm isotropic. We used a two-stage

registration. In the first step, we computed an intermediate mean surface by using the median subject

in the study group as the initial target. In the second step, MMSR was repeated using this intermediate

mean surface as the new target.

To apply this registration to our fMRI data, we computed the mean beta images for each of our 6

conditions, sampled them at the depths defined by the 6 intra-cortical surfaces (trilinear interpolation)

and then down-sampled to a resolution of 0.8 mm isotropic. We then used the deformation field

resulting from the second stage MMSR to transform those 36 images per subject (6 conditions X 6

depths) into the group surface template space (trilinear interpolation). Mass univariate laminar GLM

were then performed in normalized group space and the parameters corresponding to the 'constant'

and 'linear' terms were averaged across subjects for each vertex.

Materials and Data Availability

The raw data of the results presented here are available in a BIDS format upon request

(remi.gau@gmail.com). The code for the analysis is available at https://github.com/Remi-Gau/AV-

Attention-7T_code/releases/tag/v0.0.1. The code to run the fMRI experiment is available at

https://github.com/Remi-Gau/AV_Attention-Presentation_code-fMRI/releases/tag/v0.0.1.

ACKNOWLEDGMENTS

This project was funded by the ERC starter grant (mult-sens) and the Max Planck Society

AUTHOR CONTRIBUTIONS

Conceptualization: R.G., U.N., R.Tu.; Investigation: R.G.; Formal Analysis and Validation: R.G., U.N.; Methodology: R.G., U.N., R.Tr., R.Tu.; Software: R.G., PL.B.; Resources: PL.B, R.Tr., R.Tu.; Writing – Original Draft: R.G., UN; Writing – Review and Editing: R.G., U.N., PL.B., R.Tr., R.Tu.; Visualization: R.G., Supervision: U.N.; Project administration: U.N, R.Tu., Funding Acquisition: U.N., R.Tu.

REFERENCES

Atilgan H, Town SM, Wood KC, Jones GP, Maddox RK, Lee AKC, Bizley JK. 2018. Integration of Visual Information in Auditory Cortex Promotes Auditory Scene Analysis through Multisensory Binding. *Neuron* **97**:640-655.e4. doi:10.1016/j.neuron.2017.12.034

Bazin P-L, Weiss M, Dinse J, Schäfer A, Trampel R, Turner R. 2014. A computational framework for ultra-high resolution cortical segmentation at 7Tesla. *NeuroImage* **93**:201–209. doi:10.1016/j.neuroimage.2013.03.077

Beauchamp MS, Lee KE, Argall BD, Martin A. 2004. Integration of Auditory and Visual Information about Objects in Superior Temporal Sulcus. *Neuron* **41**:809–823. doi:10.1016/S0896-6273(04)00070-4

Bizley JK, Nodal FR, Bajo VM, Nelken I, King AJ. 2007. Physiological and Anatomical Evidence for Multisensory Interactions in Auditory Cortex. *Cereb Cortex* **17**:2172–2189. doi:10.1093/cercor/bhl128

Bressler DW, Fortenbaugh FC, Robertson LC, Silver MA. 2013. Visual spatial attention enhances the amplitude of positive and negative fMRI responses to visual stimulation in an eccentricity-dependent manner. *Vision Res* **85**:104–112. doi:10.1016/j.visres.2013.03.009

Budinger E, Heil P, Hess A, Scheich H. 2006. Multisensory processing via early cortical stages: Connections of the primary auditory cortical field with other sensory systems. *Neuroscience* **143**:1065–1083. doi:10.1016/j.neuroscience.2006.08.035

Butler JS, Foxe JJ, Fiebelkorn IC, Mercier MR, Molholm S. 2012. Multisensory Representation of Frequency across Audition and Touch: High Density Electrical Mapping Reveals Early Sensory-Perceptual Coupling. *J Neurosci* **32**:15338–15344. doi:10.1523/JNEUROSCI.1796-12.2012

Campi KL, Bales KL, Grunewald R, Krubitzer L. 2010. Connections of Auditory and Visual Cortex in the Prairie Vole (Microtus ochrogaster): Evidence for Multisensory Processing in Primary Sensory Areas. *Cereb Cortex* **20**:89–108. doi:10.1093/cercor/bhp082

Cappe C, Thelen A, Romei V, Thut G, Murray MM. 2012. Looming Signals Reveal Synergistic Principles of Multisensory Integration. *J Neurosci* **32**:1171–1182. doi:10.1523/JNEUROSCI.5517-11.2012

Ciaramitaro VM, Buračas GT, Boynton GM. 2007. Spatial and Cross-Modal Attention Alter Responses to Unattended Sensory Information in Early Visual and Auditory Human Cortex. *J Neurophysiol* **98**:2399–2413. doi:10.1152/jn.00580.2007

De Martino F, Moerel M, Ugurbil K, Goebel R, Yacoub E, Formisano E. 2015a. Frequency preference and attention effects across cortical depths in the human primary auditory cortex. *Proc Natl Acad Sci* **112**:16036–16041. doi:10.1073/pnas.1507552112

De Martino F, Moerel M, Xu J, van de Moortele P-F, Ugurbil K, Goebel R, Yacoub E, Formisano E. 2015b. High-Resolution Mapping of Myeloarchitecture In Vivo: Localization of Auditory Areas in the Human Brain. *Cereb Cortex* **25**:3394–3405. doi:10.1093/cercor/bhu150

Dick F, Taylor Tierney A, Lutti A, Josephs O, Sereno MI, Weiskopf N. 2012. In Vivo Functional and Myeloarchitectonic Mapping of Human Primary Auditory Areas. *J Neurosci* **32**:16095–16105. doi:10.1523/JNEUROSCI.1712-12.2012

Driver J, Noesselt T. 2008. Multisensory Interplay Reveals Crossmodal Influences on 'Sensory-Specific' Brain Regions, Neural Responses, and Judgments. *Neuron* **57**:11–23. doi:10.1016/j.neuron.2007.12.013

Ernst MO, Bülthoff HH. 2004. Merging the senses into a robust percept. *Trends Cogn Sci* **8**:162–169. doi:10.1016/j.tics.2004.02.002

Fairhall SL, Macaluso E. 2009. Spatial attention can modulate audiovisual integration at multiple cortical and subcortical sites. *Eur J Neurosci* **29**:1247–1257. doi:10.1111/j.1460-9568.2009.06688.x

Falchier A, Clavagnier S, Barone P, Kennedy H. 2002. Anatomical evidence of multimodal
integration in primate striate cortex. *J Neurosci Off J Soc Neurosci* **22**:5749–5759.
doi:20026562
Fan L, Li H, Zhuo J, Zhang Y, Wang J, Chen L, Yang Z, Chu C, Xie S, Laird AR, Fox PT, Eickhoff
SB, Yu C, Jiang T. 2016. The Human Brainnetome Atlas: A New Brain Atlas Based on
Connectional Architecture. *Cereb Cortex* **26**:3508–3526. doi:10.1093/cercor/bhw157
Felleman DJ, Van Essen DC. 1991. Distributed hierarchical processing in the primate cerebral
cortex. *Cereb Cortex N Y N 1991* **1**:1–47.
Fetsch CR, Pouget A, DeAngelis GC, Angelaki DE. 2012. Neural correlates of reliability-based cue
weighting during multisensory integration. *Nat Neurosci* **15**:146–154. doi:10.1038/nn.2983
Formisano E, Kim DS, Di Salle F, van de Moortele PF, Ugurbil K, Goebel R. 2003. Mirror-
symmetric tonotopic maps in human primary auditory cortex. *Neuron* **40**:859–869.
Fracasso A, Luijten PR, Dumoulin SO, Petridou N. 2018. Laminar imaging of positive and negative
BOLD in human visual cortex at 7 T. *NeuroImage* **164**:100–111.
doi:10.1016/j.neuroimage.2017.02.038
Friston KJ, Zarahn E, Josephs O, Henson RNA, Dale AM. 1999. Stochastic Designs in Event-
Related fMRI. *NeuroImage* **10**:607–619. doi:10.1006/nimg.1999.0498
Ghazanfar A, Schroeder C. 2006. Is neocortex essentially multisensory? *Trends Cogn Sci* **10**:278–
285. doi:10.1016/j.tics.2006.04.008
Goense J, Merkle H, Logothetis NK. 2012. High-Resolution fMRI Reveals Laminar Differences in
Neurovascular Coupling between Positive and Negative BOLD Responses. *Neuron* **76**:629–
639. doi:10.1016/j.neuron.2012.09.019
Hackett TA, Preuss TM, Kaas JH. 2001. Architectonic identification of the core region in auditory
cortex of macaques, chimpanzees, and humans. *J Comp Neurol* **441**:197–222.
doi:10.1002/cne.1407
Huber L, Handwerker DA, Jangraw DC, Chen G, Hall A, Stüber C, Gonzalez-Castillo J, Ivanov D,
Marrett S, Guidi M, Goense J, Poser BA, Bandettini PA. 2017. High-Resolution CBV-fMRI
Allows Mapping of Laminar Activity and Connectivity of Cortical Input and Output in
Human M1. *Neuron* **96**:1253-1263.e7. doi:10.1016/j.neuron.2017.11.005
Ibrahim LA, Mesik L, Ji X, Fang Q, Li H, Li Y, Zingg B, Zhang LI, Tao HW. 2016. Cross-
Modality Sharpening of Visual Cortical Processing through Layer-1-Mediated Inhibition
and Disinhibition. *Neuron* **89**:1031–1045. doi:10.1016/j.neuron.2016.01.027
Iurilli G, Ghezzi D, Olcese U, Lassi G, Nazzaro C, Tonini R, Tucci V, Benfenati F, Medini P. 2012.
Sound-Driven Synaptic Inhibition in Primary Visual Cortex. *Neuron* **73**:814–828.
doi:10.1016/j.neuron.2011.12.026
Johnson JA, Zatorre RJ. 2005. Attention to Simultaneous Unrelated Auditory and Visual Events:
Behavioral and Neural Correlates. *Cereb Cortex* **15**:1609–1620. doi:10.1093/cercor/bhi039
Kayser C, Logothetis NK, Panzeri S. 2010. Visual Enhancement of the Information Representation
in Auditory Cortex. *Curr Biol* **20**:19–24. doi:10.1016/j.cub.2009.10.068
Kayser C, Petkov CI, Augath M, Logothetis NK. 2007. Functional Imaging Reveals Visual
Modulation of Specific Fields in Auditory Cortex. *J Neurosci* **27**:1824–1835.
doi:10.1523/JNEUROSCI.4737-06.2007
Kayser C, Petkov CI, Logothetis NK. 2008. Visual Modulation of Neurons in Auditory Cortex.
Cereb Cortex **18**:1560–1574. doi:10.1093/cercor/bhm187
Kelley K. 2005. The Effects of Nonnormal Distributions on Confidence Intervals Around the
Standardized Mean Difference: Bootstrap and Parametric Confidence Intervals. *Educ
Psychol Meas* **65**:51–69. doi:10.1177/0013164404264850
Kok P, Bains LJ, van Mourik T, Norris DG, de Lange FP. 2016. Selective Activation of the Deep
Layers of the Human Primary Visual Cortex by Top-Down Feedback. *Curr Biol* **26**:371–
376. doi:10.1016/j.cub.2015.12.038

Lakatos P, Chen C-M, O'Connell MN, Mills A, Schroeder CE. 2007. Neuronal Oscillations and
954  Multisensory Interaction in Primary Auditory Cortex. *Neuron* **53**:279–292.
955  doi:10.1016/j.neuron.2006.12.011

956 Lakatos P, O'Connell MN, Barczak A, Mills A, Javitt DC, Schroeder CE. 2009. The Leading
957  Sense: Supramodal Control of Neurophysiological Context by Attention. *Neuron* **64**:419–
958  430. doi:10.1016/j.neuron.2009.10.014

959 Lakens D. 2013. Calculating and reporting effect sizes to facilitate cumulative science: a practical
960  primer for t-tests and ANOVAs. *Front Psychol* **4**. doi:10.3389/fpsyg.2013.00863

961 Landman BA, Bogovic JA, Carass A, Chen M, Roy S, Shiee N, Yang Z, Kishore B, Pham D, Bazin
962  P-L, Resnick SM, Prince JL. 2013. System for Integrated Neuroimaging Analysis and
963  Processing of Structure. *Neuroinformatics* **11**:91–103. doi:10.1007/s12021-012-9159-9

964 Laurienti PJ, Burdette JH, Wallace MT, Yen Y-F, Field AS, Stein BE. 2002. Deactivation of
965  Sensory-Specific Cortex by Cross-Modal Stimuli. *J Cogn Neurosci* **14**:420–429.
966  doi:10.1162/089892902317361930

967 Leitão J, Thielscher A, Werner S, Pohmann R, Noppeney U. 2013. Effects of Parietal TMS on
968  Visual and Auditory Processing at the Primary Cortical Level – A Concurrent TMS-fMRI
969  Study. *Cereb Cortex* **23**:873–884. doi:10.1093/cercor/bhs078

970 Leo F, Romei V, Freeman E, Ladavas E, Driver J. 2011. Looming sounds enhance orientation
971  sensitivity for visual stimuli on the same side as such sounds. *Exp Brain Res* **213**:193–201.
972  doi:10.1007/s00221-011-2742-8

973 Liang M, Mouraux A, Hu L, Iannetti GD. 2013. Primary sensory cortices contain distinguishable
974  spatial patterns of activity for each sense. *Nat Commun* **4**. doi:10.1038/ncomms2979

975 Logothetis NK. 2008. What we can do and what we cannot do with fMRI. *Nature* **453**:869–878.
976  doi:10.1038/nature06976

977 Lucas BC, Bogovic JA, Carass A, Bazin P-L, Prince JL, Pham DL, Landman BA. 2010. The Java
978  Image Science Toolkit (JIST) for Rapid Prototyping and Publishing of Neuroimaging
979  Software. *Neuroinformatics* **8**:5–17. doi:10.1007/s12021-009-9061-2

980 Maier JX, Chandrasekaran C, Ghazanfar AA. 2008. Integration of Bimodal Looming Signals
981  through Neuronal Coherence in the Temporal Lobe. *Curr Biol* **18**:963–968.
982  doi:10.1016/j.cub.2008.05.043

983 Maier JX, Neuhoff JG, Logothetis NK, Ghazanfar AA. 2004. Multisensory Integration of Looming
984  Signals by Rhesus Monkeys. *Neuron* **43**:177–181. doi:10.1016/j.neuron.2004.06.027

985 Markuerkiaga I, Barth M, Norris DG. 2016. A cortical vascular model for examining the specificity
986  of the laminar BOLD signal. *NeuroImage* **132**:491–498.
987  doi:10.1016/j.neuroimage.2016.02.073

988 Marques JP, Kober T, Krueger G, van der Zwaag W, Van de Moortele P-F, Gruetter R. 2010.
989  MP2RAGE, a self bias-field corrected sequence for improved segmentation and T1-
990  mapping at high field. *NeuroImage* **49**:1271–1281. doi:10.1016/j.neuroimage.2009.10.002

991 McAuliffe MJ, Lalonde FM, McGarry D, Gandler W, Csaky K, Trus BL. 2001. Medical Image
992  Processing, Analysis and Visualization in clinical research. IEEE Comput. Soc. pp. 381–
993  386. doi:10.1109/CBMS.2001.941749

994 Mercier MR, Foxe JJ, Fiebelkorn IC, Butler JS, Schwartz TH, Molholm S. 2013. Auditory-driven
995  phase reset in visual cortex: Human electrocorticography reveals mechanisms of early
996  multisensory integration. *NeuroImage* **79**:19–29. doi:10.1016/j.neuroimage.2013.04.060

997 Morrill RJ, Hasenstaub AR. 2018. Visual Information Present in Infragranular Layers of Mouse
998  Auditory Cortex. *J Neurosci* **38**:2854–2862. doi:10.1523/JNEUROSCI.3102-17.2018

999 Mozolic JL, Joyner D, Hugenschmidt CE, Peiffer AM, Kraft RA, Maldjian JA, Laurienti PJ. 2008.
1000  Cross-modal deactivations during modality-specific selective attention. *BMC Neurol* **8**.
1001  doi:10.1186/1471-2377-8-35

1002 Muckli L, De Martino F, Vizioli L, Petro LS, Smith FW, Ugurbil K, Goebel R, Yacoub E. 2015.
1003  Contextual Feedback to Superficial Layers of V1. *Curr Biol* **25**:2690–2695.
1004  doi:10.1016/j.cub.2015.08.057

Müller NG, Kleinschmidt A. 2004. The attentional "spotlight's" penumbra: center-surround modulation in striate cortex. *Neuroreport* **15**:977–980.

Nieuwenhuys R. 2013. The myeloarchitectonic studies on the human cerebral cortex of the Vogt–Vogt school, and their significance for the interpretation of functional neuroimaging data. *Brain Struct Funct* **218**:303–352. doi:10.1007/s00429-012-0460-z

Noesselt T, Rieger JW, Schoenfeld MA, Kanowski M, Hinrichs H, Heinze H-J, Driver J. 2007. Audiovisual Temporal Correspondence Modulates Human Multisensory Superior Temporal Sulcus Plus Primary Sensory Cortices. *J Neurosci* **27**:11431–11441. doi:10.1523/JNEUROSCI.2252-07.2007

Rockland KS, Ojima H. 2003. Multisensory convergence in calcarine visual areas in macaque monkey. *Int J Psychophysiol Off J Int Organ Psychophysiol* **50**:19–26.

Rohe T, Noppeney U. 2016. Distinct Computational Principles Govern Multisensory Integration in Primary Sensory and Association Cortices. *Curr Biol* **26**:509–514. doi:10.1016/j.cub.2015.12.056

Rohe T, Noppeney U. 2015. Cortical Hierarchies Perform Bayesian Causal Inference in Multisensory Perception. *PLOS Biol* **13**:e1002073. doi:10.1371/journal.pbio.1002073

Schroeder CE, Foxe JJ. 2002. The timing and laminar profile of converging inputs to multisensory areas of the macaque neocortex. *Brain Res Cogn Brain Res* **14**:187–198.

Schroeder CE, Smiley J, Fu KG, McGinnis T, O'Connell MN, Hackett TA. 2003. Anatomical mechanisms and functional implications of multisensory convergence in early cortical processing. *Int J Psychophysiol* **50**:5–17. doi:10.1016/S0167-8760(03)00120-X

Shmuel A, Yacoub E, Pfeuffer J, Van de Moortele PF, Adriany G, Hu X, Ugurbil K. 2002. Sustained negative BOLD, blood flow and oxygen consumption response and its coupling to the positive response in the human brain. *Neuron* **36**:1195–1210.

Shomstein S, Yantis S. 2004. Control of Attention Shifts between Vision and Audition in Human Cortex. *J Neurosci* **24**:10702–10706. doi:10.1523/JNEUROSCI.2939-04.2004

Silver MA, Shenhav A, D'Esposito M. 2008. Cholinergic Enhancement Reduces Spatial Spread of Visual Responses in Human Early Visual Cortex. *Neuron* **60**:904–914. doi:10.1016/j.neuron.2008.09.038

Smiley JF, Falchier A. 2009. Multisensory connections of monkey auditory cerebral cortex. *Hear Res* **258**:37–46. doi:10.1016/j.heares.2009.06.019

Smith AT, Singh KD, Greenlee MW. 2000. Attentional suppression of activity in the human visual cortex. *Neuroreport* **11**:271–277.

Talsma D, Doty TJ, Woldorff MG. 2006. Selective Attention and Audiovisual Integration: Is Attending to Both Modalities a Prerequisite for Early Integration? *Cereb Cortex* **17**:679–690. doi:10.1093/cercor/bhk016

Talsma D, Senkowski D, Soto-Faraco S, Woldorff MG. 2010. The multifaceted interplay between attention and multisensory integration. *Trends Cogn Sci* **14**:400–410. doi:10.1016/j.tics.2010.06.008

Tardif CL, Schäfer A, Waehnert M, Dinse J, Turner R, Bazin P-L. 2015. Multi-contrast multi-scale surface registration for improved alignment of cortical areas. *NeuroImage* **111**:107–122. doi:10.1016/j.neuroimage.2015.02.005

Tootell RB., Hadjikhani N, Hall EK, Marrett S, Vanduffel W, Vaughan JT, Dale AM. 1998. The Retinotopy of Visual Spatial Attention. *Neuron* **21**:1409–1422. doi:10.1016/S0896-6273(00)80659-5

Trampel R, Bazin P-L, Heidemann RM, Schäfer A, Ivanov D, Lohmann G, Geyer S, Turner R. 2012. Laminar-specific fingerprints of different sensorimotor areas obtained during imagined and actual finger tapping. *Proc Int Soc Magn Reson Med* **20**.

Trampel R, Bazin P-L, Pine K, Weiskopf N. 2017. In-vivo magnetic resonance imaging (MRI) of laminae in the human cortex. *NeuroImage*. doi:10.1016/j.neuroimage.2017.09.037

1055   Tyll S, Bonath B, Schoenfeld MA, Heinze H-J, Ohl FW, Noesselt T. 2013. Neural basis of
1056   multisensory looming signals. *NeuroImage* **65**:13–22.
1057   doi:10.1016/j.neuroimage.2012.09.056
1058   Van der Burg E, Olivers CNL, Bronkhorst AW, Theeuwes J. 2008. Pip and pop: Nonspatial
1059   auditory signals improve spatial visual search. *J Exp Psychol Hum Percept Perform*
1060   **34**:1053–1065. doi:10.1037/0096-1523.34.5.1053
1061   Waehnert MD, Dinse J, Weiss M, Streicher MN, Waehnert P, Geyer S, Turner R, Bazin P-L. 2014.
1062   Anatomically motivated modeling of cortical laminae. *NeuroImage* **93**:210–220.
1063   doi:10.1016/j.neuroimage.2013.03.078
1064   Wandell BA, Dumoulin SO, Brewer AA. 2007. Visual Field Maps in Human Cortex. *Neuron*
1065   **56**:366–383. doi:10.1016/j.neuron.2007.10.012
1066   Wang L, Mruczek REB, Arcaro MJ, Kastner S. 2015. Probabilistic Maps of Visual Topography in
1067   Human Cortex. *Cereb Cortex* **25**:3911–3931. doi:10.1093/cercor/bhu277
1068   Werner S, Noppeney U. 2011. The Contributions of Transient and Sustained Response Codes to
1069   Audiovisual Integration. *Cereb Cortex* **21**:920–931. doi:10.1093/cercor/bhq161
1070   Werner S, Noppeney U. 2010. Distinct Functional Contributions of Primary Sensory and
1071   Association Areas to Audiovisual Integration in Object Categorization. *J Neurosci* **30**:2662–
1072   2675. doi:10.1523/JNEUROSCI.5091-09.2010
1073

1074

SUPPLEMENTARY FIGURES

Auditory and visual responses

A   Surface projections for individual subjects

Figure S1: Auditory and visual responses and their attentional modulation

<u>Surface projection for individual subjects:</u> Within subject 'constant' shape parameter estimates of

the laminar BOLD response profile for visual (column i) and auditory (column ii) are projected on

each subject's inflated surface to show visual regions of the left hemisphere. V1 and V2/3 are

delineated by black solid and dashed lines respectively. For visualization purposes only, surface

projections were smoothed (FWHM = 1.5 mm). Grey areas denote vertices with no available data

SUPPLEMENTARY TABLES

Table 1

Behavioural results

Stimuli / Targets	Auditory			Visual			Audio-visual		
	Attend to auditory								
Auditory	80.11%	(13.92%	)	93.18%	(7.63%	)	67.05%	(15.08%	)
Visual	6.82%	(7.63%	)	2.27%	(4.21%	)	11.36%	(7.82%	)
	Attend to visual								
Auditory	4.55%	(11.21%	)	6.25%	(6.25%	)	7.95%	(7.95%	)
Visual	65.91%	(16.14%	)	69.32%	(14.91%	)	65.91%	(11.98%	)

1087 Notes: Percentage of target responses (mean and STD across subjects) in the six conditions of our 2 x 3 experimental design. n = 11

1088

1089 **Table 2**

1090 ROI size and coverage

ROI	Number of vertices (*10^3)			Proportion of ROI covered		
	mean	STD	range	mean	STD	range
A1	19.33	3.85	(15.05 - 24.28)	0.95	0.07	(0.84 - 1.00)
PT	10.07	2.73	(6.85 - 14.36)	0.98	0.03	(0.91 - 1.00)
V1	33.85	7.50	(24.64 - 48.44)	0.97	0.02	(0.94 - 0.99)
V2-3	67.15	13.91	(43.59 - 87.51)	0.83	0.08	(0.64 - 0.90)

1091 Notes: 'Number of vertices' refers to the vertices with valid data at all the sampled cortical depths. This vertex count was divided by the total number of

1092 vertices included in the initial ROI definition to compute the 'Fraction of the ROI covered'. n = 11

1093

1094