HEXOKINASE 1 Control of Post-Germinative Seedling Growth

Matthew J. Lincoln[1], Ashwin Ganpudi[1], Andrés Romanowski[1,2], Karen J. Halliday[1]

1. Institute for Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh,
Edinburgh EH9 3BF, United Kingdom.

2. Plant-Environment Signalling Group, Department of Biology, Faculty of Science, Utrecht
University, Utrecht 3584 CH, The Netherlands.

Matthew J. Lincoln, Ashwin Ganpudi contributed equally to this manuscript

ORCID IDs: 0000-0003-2822-8501 (M.J.L.); 0000-0002-9515-3899 (A.G.); 0000-0003-0737-2408
(A.R.); 0000-0003-0467-5 104X (K.J.H.)

Corresponding author:

Karen J. Halliday

Institute for Molecular Plant Sciences, School of Biological Sciences, University of Edinburgh,
Edinburgh EH9 3BF, United Kingdom

+44-0131-651-9083

karen.halliday@ed.ac.uk

Summary

- In darkness, PHYTOCHROME INTERACTING FACTOR (PIF)-induced skotomorphogenic

seedling growth, is exemplified by increased hypocotyl elongation. HEXOKINASE1

(HXK1), which is also implicated in seedling establishment, can operate as a glycolytic

enzyme or as a glucose-activated sensor signalling molecule. Under light and nutrient limiting

conditions, the HXK1 sensor-signalling has been shown to control hypocotyl elongation.

Little is known of whether HXK1 glycolytic function, or HXK1 and PIF cross-talk, is

required to control hypocotyl growth.

- We demonstrate HXK1 glycolytic activity is critical for cell expansion, and hypocotyl

growth, post-germination. Notably, application of glucose-6-phosphate, the HXK1 enzymatic

product, can restore short *gin2-1/hxk1-1* mutant hypocotyls to wild-type length. Further,

HXK1 sensor-signalling complex components, VHA-B1 and RPT5B, do not contribute to this

response, for unlike *gin2-1/hxk1-1*, the *vha-B1* and *rpt5b* alleles only disrupt hypocotyl

growth following exogenous glucose application.

- mRNA-seq analysis illustrates that HXK1 and PIF signalling converge at genes with known

roles in light signalling. HXK1 imposes strong regulation on chloroplast and mitochondrial

encoded genes, also branched chain amino acid catabolism pathway genes, which can provide

a source of respiratory substrates in starvation conditions.

- Our study establishes the importance of HXK1 enzymatic function in supporting cell

expansion and hypocotyl growth. We demonstrate a degree of cross-talk between HXK1 and

PIFs through common target gene set.

Key Words: Hexokinase, Phytochrome, Skotomorphogenesis, Hypocotyl, *Arabidopsis thaliana*

Introduction

The first days of life are critical for plant survival; successful seedling establishment relies on the mobilisation of seed reserves and the perception of light through wavelength-specific photoreceptors to support emergence before the switch to photo-autotrophic growth. HEXOKINASE 1 (HXK1), a metabolic enzyme with glucose-sensor signalling properties, and PHYTOCHROME-INTERACTING FACTORS (PIFs), play a pivotal role in coupling carbon availability to growth in the developing seedling (Rolland et al, 2002, Moore et al, 2003, Stewart et al, 2011, Lilley et al, 2012, Sairanen et al, 2013, de Wit et al, 2018, Krahmer et al, 2021).

In Arabidopsis and other oilseed plants, carbon resources are stored primarily in the form of triacylglycerols (TAGs) and seed storage proteins (SSP) which account for approximately 30-40% of the seed's dry weight each (Baud et al, 2008). After germination TAGs are degraded and converted to sucrose. TAG-derived fatty acids undergo B-oxidation, forming acetyl CoA, which fuels the production of sucrose via the glyoxylate cycle and gluconeogenesis. Sucrose is subsequently transported throughout the seedling to support development (Eastmond et al, 2000, Yang and Benning, 2018). In sink tissues, sucrose is cleaved by either SUCROSE SYNTHASE to produce UDP-glucose and fructose, or by INVERTASE to liberate glucose and fructose (Yoon et al, 2021). A vital step in the metabolism of these hexose sugars is their phosphorylation, which is mediated by HXK and FRUCTOKINASE (FRK). HXK can phosphorylate both glucose and fructose, while FRK primarily phosphorylates fructose. That said, FRK has a higher binding affinity for fructose than HXK, suggesting the principal role of HXK is the phosphorylation of glucose to glucose-6-phosphate (G6P) (Harrington and Bush, 2003, Claeyssen and Rivoal, 2007, Granot, 2007). In Arabidopsis, there are six different members of the HXK family: HXK1-3, and HEXOKINASE-LIKE (HKL) 1-3 (Karve et al, 2008). A key difference between the HXK and HKLs is the lack of glucokinase activity in the three HKL proteins, likely attributable to an indel at the adenosine binding site. Across the family, the most well studied member is HXK1.

In addition to its enzymatic role, HXK1 is known to operate as a glucose sensor-signalling molecule, directly coupling nuclear control of gene expression to glucose availability. Here, HXK1 forms a complex with partner proteins VACUOLAR H$^+$ ATPASE SUBUNIT B1 (VHA-B1) and REGULATORY PARTICLE 5B (RPT5B), one of two 19S Proteasome AAA-ATPases. Chromatin immunoprecipitation (ChIP) assays indicate VHA-B1- and RPT5B- dependent HXK1, enrichment at the 5'-272 BP region of the *CHLOROPHYLL A/B BINDING 2 (CAB2)* promoter (which contains DtRE, CUF-1, CCA1, and CGF-1 binding motifs). As none of the complex proteins possess DNA-binding domains, VHA-B1 and RPT5B are proposed to interact with transcription factors to regulate gene expression (Cho et al, 2006). Genetic analysis has shown that HXK1, VHA-B1 or RPT5B mediate glucose-induced repression of the photosynthetic genes, *CAB2* and *CARBONIC*

ANHYDRASE 1 (CAA) expression, and glucose-induced seedling developmental arrest (Cho et al,

2006). The retention of these responses in the HXK1 catalytic mutants G104D (Gly[104] to Asp[104]) and

S177A (Ser[177] to Ala[177]), which retain the glucose-binding site, suggested in these instances, HXK1

operated through glucose-activated signalling, rather than through its glycolytic mode (Moore et al,

2003, Feng et al, 2015).

Interestingly, HXK1 glucose-signalling is not just implicated in *CAB2/CAA* expression and

developmental arrest, which are elicited by high glucose levels, but also in the regulation of hypocotyl

elongation in low light and low nutrient levels (Moore et al, 2003, Cho et al, 2006). The loss of

function HXK1 mutants, glucose-insensitive 2 (*gin2-1)* (Landsberg erecta) and *hxk1-3* (Columbia-0)

exhibit significantly reduced hypocotyl sizes, particularly in limited light conditions (Jang and Sheen,

1997, Jang et al, 1997, Kelly et al, 2021). Additionally, the *vha-B1* and *rpt5b* mutants phenocopy the

gin2-1 short hypocotyl phenotype, while the catalytically inactive alleles *hxk1^{G104D}* and *hxk1^{S177A}*

restore the WT hypocotyl response (Moore et al, 2003; Cho et al, 2006). It would be interesting to

establish whether HXK1 catalytic action assists with the metabolism of seed reserves that fuel

seedling growth prior to the establishment of photosynthetic competence.

The regulation of hypocotyl elongation in darkness and low light conditions is known to be

regulated by PHYTOCHROME INTERACTION FACTORS (PIFs), a family of basic helix-loop-

helix (bHLH) transcription factors. PIFs (PIF1, 2-8) are regulated by the phytochrome (phy) family of

photoreceptors, of which phyB has been seen to be critically important in regulating seedling de-

etiolation (Franklin and Quail, 2010). Excitation of the phyB chromophore with red (600-700 nm)

light, induces a conformational change from the inactive (Pr) to the active (Pfr) form, which

preferentially binds to PIF. This physical interaction results in rapid phosphorylation and subsequent

degradation of PIFs 1, 3, 4, and 5, as well as sequestering them from their target promoters (Leivar et

al, 2008, Park et al, 2012, Park et al, 2018). As a result, these PIFs are most active in darkness and

low light, evident in the short hypocotyl phenotype of the quadruple *pifQ* mutant, which comprises

pif1-1, pif3-1, pif4-2, and *pif5-3* alleles (Leivar et al, 2008). Thus, light-activated phyB repression of

PIFs is critical for the switch from skotomorphogenic to photomorphogenic growth.

Both phy and PIFs have been implicated in carbon resource partitioning, where they have an

important role in adjusting growth allocations to sink tissues in darkness and shaded conditions (Yang

et al, 2016, de Wit et al, 2018, Krahmer et al, 2021). PIFs are involved sucrose regulation of

hypocotyl growth; the *pifQ* mutant is insensitive to sucrose supplementation, and PIF5 protein

stability is moderated by sucrose. (Liu et al, 2011). PIF4 has an important role in coupling carbon

status to temperature-dependent hypocotyl growth (Hwang et al, 2019, Bian et al, 2022). Here,

Trehalose-6-phosphate (Tre6P), a key indicator of sucrose status, stabilises PIF4 by inhibiting SNF1-

RELATED PROTEIN KINASE1 (SnRK1)-mediated PIF4 phosphorylation and 26S proteasome

destruction. Further, phys, together with the nutrient-sensing pathways have been shown to have a

critical role in stimulating seedling shoot apical meristem stem cell division, by activating TARGET

OF RAPAMYCIN (TOR) (Pfeiffer et al, 2016). More recently, a link has been made between guard

cell-located PIF4 and HXK1 and sucrose-mediated hypocotyl elongation in long days. This work

demonstrated that HXK1 overexpression in guard cells enhances *PIF4* expression and hypocotyl

elongation in response to blue light wavelengths (Kelly et al, 2021).

In this study, we assessed seedlings under darkness and low light regimes to determine HXK1

contribution to hypocotyl elongation fuelled by seed reserve utilisation. Genetic, pharmacological,

and molecular analysis point to an important role for HXK1 enzymatic function in supporting

hypocotyl growth in light-limiting conditions. Our mRNA-seq data reveals HXK1 exerts wide-

ranging control of metabolic genes, including a branched chain amino-acid pathway, which provides

an alternative source of carbon in starvation conditions. The mRNA-seq analysis also identifies cross-

talk between HXK1 and PIFs at key light signalling components. Finally, we demonstrate that several

aspects of the HXK1 sensor-signalling pathway do not operate in light-limiting conditions.

Materials and Methods

Plant material, growth conditions, and treatments. The wild-type Arabidopsis thaliana accessions

used in this study are Landsberg erecta (Ler) and Columbia-0 (Col). Seeds for mutants *gin2-1* (Ler),

hxk1-3 (Col), *pfk1-1* (Col), *vha-B1* (Col, SALK_028728), *rpt5b* (Col, SALK_069366), and *king1*

(Col, At3g48530) were obtained from The Nottingham Arabidopsis Stock Centre (NASC), UK. The

hxk1^{S177A} seeds were kindly shared by Professor Ruth Stadler of the University of Erlangen-

Nuernberg. For all experiments, seeds were surface sterilized with bleach and Triton X-100 sown on

0.5X MS plates (0.8% agar, pH 5.7) and stratified in darkness for 2-3 days at 4°C. All plants were

grown at 18°C for 4 days after stratification unless otherwise specified.

Seedling hypocotyl length and cotyledon area measurements. Images of seedlings laid flat on

growth media were used to quantify hypocotyl length and cotyledon area using ImageJ (NIH,

Maryland, USA) and Adobe Photoshop CS6 (Adobe, California, USA), respectively. For glucose-6-

phosphate (G6P) (Sigma G7879), sodium pyruvate (Sigma P2256), and D-mannoheptose (Sigma

M6909-1G) treatments, the required concentrations from filter sterilized stocks were added to

sterilized media and seeds directly sown. Unless otherwise stated, all experiments were performed in

triplicate with 40 seedlings per replicate.

Seedling hypocotyl cell counts and length measurements. For determination of etiolated hypocotyl

epidermal cell lengths/numbers, seedlings were mounted on slides with water and visualized using an

Eclipse E600 Nikon DIC microscope at 20x magnification. Individual cell lengths from each section

of the hypocotyl (basal, middle, upper) were measured using ImageJ and cell number per file was

obtained by manually counting cells from the root emergence point to the apical meristem. Unless

otherwise stated, all experiments were performed in triplicate with at least 20 seedlings per replicate.

Glucose quantification. Seedlings were harvested in liquid nitrogen, finely ground into a powder,

and ethanol extracted thrice. Glucose was then quantified from these ethanol extracts using enzymatic

degradation at 340nm wavelength and normalized to material fresh weight. Unless otherwise stated,

all experiments were performed in triplicate with 50 seedlings per replicate

Gene expression analysis. For qRT-PCR experiments, seedlings harvested in liquid nitrogen were

ground into fine powder. Total RNA was extracted using the RNeasy Plant Mini Kit (Qiagen) with

on-column DNase digestion. cDNA synthesis was performed using the qScript cDNA SuperMix

(Quanta Biosciences) as described by the manufacturer. The qRT-PCR was set up as a 10μL reaction

using SYBR® Green 1 480 Lightcycler® Master (Roche) in a 384-well plate, performed with a

Lightcycler 480 system (Roche). Results were analysed using the Light Cycler 480 software. The

primers used in this study are listed in Supplementary Table S1. Unless otherwise stated, all

experiments were performed in triplicate with 50 seedlings per replicate.

cDNA library preparation and high throughput sequencing. Total RNA was extracted from 4-

day-old etiolated Ler and *gin2-1* seedlings (biological duplicates of 50 seedlings per replicate) as

described above. Samples were then sent to Edinburgh Genomics (University of Edinburgh, UK) for

QC check and sequencing. Briefly, quality check of the samples was performed using Qubit with the

broad range RNA kit (Thermo Fisher Scientific) and Tapestation 4200 with the RNA Screentape for

eukaryotic RNA analysis (Agilent). Libraries were prepared using the TruSeq Stranded mRNA kit

(Illumina) and then validated. Samples were pooled to create 4 multiplexed DNA libraries, which

were paired-end sequenced on an Illumina HiSeq 4000 platform (Machine name K00166, Run

number 346, flowcell AHT2HKBBXX, lanes 5 and 6). On average 26.6 million 150nt PE reads were

obtained for each sample.

Processing of mRNA sequencing reads. Raw sequence reads were trimmed with cutadapt 2.8

(Martin, 2011) with default parameters and --a set to 'AGATCGGAAGAGC', to eliminate adapter

contamination from the PE reads. Trimmed reads were aligned against the *Arabidopsis thaliana*

genome (TAIR10) with HiSat2 v2.2.1 (Kim et al, 2019) with default parameters, except in the case of

the maximum intron length parameter, which was set at 5000. Count tables for the different feature

levels were obtained from bam files using the 'ASpli::readCounts()' function of ASpli package

version 2.4.0 (Mancini et al, 2021) with custom R scripts and considering the AtRTD2 transcriptome

(Zhang et al, 2017). Count tables at the gene level presented a good correlation overall between

replicates and samples (Table S2). Raw sequences (fastq files) used in this paper have been deposited

in the ArrayExpress (Kolesnikov et al, 2015) database at EMBL-EBI (www.ebi.ac.uk/arrayexpress)

under accession number E-MTAB-7654.

Differential gene expression (DGE) analysis. DGE analysis was conducted using custom R scripts

for 21,256 genes whose expression was above a minimum threshold level (10 counts in 0.7 of samples

in the smallest group that express the gene). DGE was estimated using the edgeR package version

3.36.0 (Robinson et al, 2010; Lun et al, 2016), and resulting P values were adjusted using a false

discovery rate (FDR) criterion. Genes with $p < 0.05$, FDR < 0.05 and an absolute log2 fold change >

0.58 were considered differentially expressed. Volcano plots, calculation and plot of chromosomal

distributions, and UpSet plots of differentially expressed genes (DEGs) were generated using R.

GO and KEGG metabolic pathway analysis. Gene set enrichment and KEGG pathway enrichment

analysis were performed using a combination of custom-written R scripts and the clusterProfiler

package (Yu et al, 2012) version 4.2.2 of Bioconductor. The 21,256 expressed genes were used as the

universe gene set. All KEGG pathways of *A. thaliana* were derived from the KEGG Pathway

Database (http://www.kegg.jp; Kanehisa et al, 2012). Only the terms with $p < 0.05$ and FDR <0.1

were further considered. Bubble plots were generated using R.

Statistical analysis. The statistical difference between two populations was tested by two-tailed,

unpaired Student's t-test. To compare three or more populations, a one-way analysis of variance

(ANOVA) followed by Dunnett's test (comparison against a control) was performed. All analyses

were done using GraphPad Prism 7 (GraphPad Software) unless otherwise indicated.

Accession numbers and data availability. Raw sequences (fastq files) used in this paper have been

deposited in the ArrayExpress (Kolesnikov et al, 2015) database at EMBL-EBI

(www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-7654. All custom R scripts are

available at https://github.com/aromanowski/gin2_darkness. Alternatively, they are available upon

request to the authors

Results

HXK1 supports post-germinative hypocotyl growth in a G6P-dependent manner

The *gin2-1* mutant was previously reported to have impaired hypocotyl growth in low nutrient/low

light conditions (Moore et al, 2003), which led us to examine the fluence rate in which HXK1

operates. We established that the *gin2-1* (Ler) and *hxk1-3* (Col-0) hypocotyls were shorter in darkness

and very low irradiance light but were indistinguishable from the wild type at fluence rates greater

than $10\,\mu\mathrm{mol\,m^{-2}\,s^{-1}}$ (Fig. 1a, b). This altered fluence rate response is similar to the quadruple *pifQ*

(*pif1-1, pif3-1, pif4-2, pif5-3)* mutant, though less severe. Interestingly, *gin2-1* and *hxk1-3* seedlings

also have short hypocotyls in 8h light (L), 16h dark (D) short days and in longer photoperiods (16h L,

8h D), but only when the light fluence rate is low (Fig. S1). Thus, the *gin2-1* and *hxk1-3* long

hypocotyl phenotypes are evident in darkness and conditions when light availability is restricted. We

also observe similar reductions in the cotyledon area when light is limited (Fig. 1c).

To establish the cellular basis of the hypocotyl defect, we measured the epidermal cell length in the

basal, middle, and upper regions of the hypocotyl, and recorded the total cell number in epidermal cell

files (Fig. 1d, e). Our results show that *gin2-1* has a lower cell file count and significantly shorter

epidermal cells through the hypocotyl. As hypocotyl cell number is largely determined during embryo

development, this suggests a role for HXK1 at this early stage. As hypocotyl elongation is driven by

hypocotyl cell expansion (Gendreau et al, 1997), our data indicates that HXK1 action may be required

to promote epidermal cell expansion post-germination, particularly when light conditions are limiting.

Previously, HXK1-dependent hypocotyl growth was shown to result from glucose-induced nuclear

signalling, rather than HXK1 enzymatic function (Moore et al, 2003, Cho et al 2006). To explore this

further, we examined the hypocotyl response over a range of glucose concentrations in low-irradiance

light and in darkness. As expected, in WT the glucose response is biphasic, with lower concentrations

promoting, and higher concentrations inhibiting hypocotyl growth (Singh et al, 2017) (Fig. 1f). In

comparison, *gin2-1* and *hxk1-3* mutants exhibited reduced sensitivity to glucose, with a lagging dose-

response curve (Fig. 1f, Fig. S2a, Fig. S2e, S2g). This was not observed in mannitol controls in both

darkness and low light (Fig. 1f, Fig. S2c-d, i-j). The results indicate that HXK1 mutants have an

altered hypocotyl dose response to exogenously applied glucose. Next, we applied a concentration

range of glucose-6-phosphate (G6P) which is the product of HXK1 enzymatic activity. Contrasting

with the glucose response, the WT was largely insensitive to G6P. Further, we found that rising doses

of G6P sequentially increased hypocotyl length in *gin2-1*, and *hxk1-3,* restoring to WT length at 5mM

(Fig. 1g, Fig. S1). Similar results were obtained for seedlings supplied with competitive HXK1

inhibitory molecule, D-mannoheptose, and 5mM G6P (Fig 1h). This suggests that in light-limiting

conditions, and the absence of exogenous glucose, HXK1 enzymatic activity is required to fuel

hypocotyl growth. In support of this proposition, we established that G6P application to *hxk1-3*

successfully restores epidermal cell length in the middle and upper regions of the hypocotyl (Fig. 1i).

As post-germinative hypocotyl cell elongation proceeds in a basal to distal wave, this suggests a

potential window of sensitivity to G6P (Boron and Vissenberg, 2014). Thus, in light-restricted

conditions, HXK1 enzymatic activity appears to play an important role in promoting hypocotyl

elongation post-germination. In comparison, hypocotyl epidermal cells of *pifQ* mutants are much

shorter than WT and *hxk1-3* but are completely insensitive to G6P (Fig. 1i). This data indicates that

PIF control of hypocotyl cell expansion is not due to deficiencies in G6P.

mRNA-seq reveals a role for HXK1 in translation and carbon stress metabolism

To gain a more in-depth understanding of the HXK1 function in seedling establishment, we

performed an mRNA-seq on 4-day-old etiolated Ler and *gin2-1* plants. Briefly, total RNA was

extracted, and 6 stranded libraries were prepared from polyA purified RNA. These were pooled and

sequenced on an Illumina HiSeq 4000 platform. Gene counts were extracted using the ASpli R

package (Mancini et al, 2021) and the AtRTDv2 annotation (Zhang et al, 2017). Raw counts were

filtered to remove lowly expressed genes, normalized to library size and gene expression was

quantified using EdgeR (Robinson et al, 2010). This resulted in 21,256 genes (62% of the 34,212

annotated AtRTD2 genes) being considered for downstream analysis. Using custom R scripts, we

performed differential gene expression (DGE) analysis using EdgeR, followed by gene ontology (GO)

enrichment analysis. We discovered 2,344 genes were significantly misregulated ($|logFC| > 0.58$, $p <$

0.01 and $FDR < 0.05$) in *gin2-1* compared to Ler, with a similar proportion of genes down (1,156) and

upregulated (1,188) (Fig. 2a, Fig. S3). When we evaluated these genes, we found that the

downregulated category of genes included lipid storage, and energy-consuming processes, such as

ribosome biogenesis and cell proliferation (Fig. 2b). In this figure, the bubbles represent each GO

term, the size of the bubble indicating the representation factor (RF) and the colour indicating the *P*-

value score. In contrast, the upregulated category had strong enrichment for genes involved in the

cellular response to starvation, cellular respiration, ATP biosynthesis, amino acid catabolism and

photosynthesis. These patterns are consistent with a starvation-type response and a reduced capacity

to catabolise sugars.

We noted that genes that were strongly upregulated in *gin2-1* included several enzymes involved in

the catabolism of the branched-chain amino acid (*BCAA*) genes: *BRANCHED-CHAIN AMINO ACID*

TRANSAMINASE 2 (*BCAT2), DARK INDUCIBLE 2 (DIN2),* AT1G21400 *(E1A1), ISOVALERYL-*

COA-DEHYDROGENASE (IVD), AT1G03090 *(MCCA),* and *3-METHYLCROTONYL-COA*

CARBOXYLASE (*MCCB*) (Fig. 2c). *BCAA* genes are often suppressed in carbon-rich conditions, and

induced by stress or starvation, as they provide alternative substrates for respiration (Pires et al, 2016,

Binder*,* 2010). Indeed, we also observe strong glucose-induced suppression of the majority of these

genes in both the WT and *gin2-1* (Fig. 2d). In most cases, the efficacy of glucose is greater in *gin2-1*

than in WT, indicating HXK1 has a role in antagonizing the glucose response (Fig. 2c). Our data also

shows that, as for hypocotyl growth (Fig. 1g), G6P application to WT has no significant impact on

BCAA gene expression. However, G6P treatment in *gin2-1* is effective in restoring the BCAA

expression to WT levels. This indicates *BCAA* gene expression is regulated by the HXK1-G6P

pathway.

Another feature of the transcriptomic data analysis is the extent to which HXK1 controls organellar

gene expression, with 23.5% of mitochondrial and 82.8% of chloroplast genes upregulated in *gin2-1*

(Fig 3a). In the mitochondria, we observe enrichment for genes regulating metabolic proton and

electron transport; and in the chloroplast, we see enrichment in all genes except for 23 genes

associated with vesicle formation and chromatin remodelling. Figure 3b shows qPCR verification for

the chloroplast-encoded *RNA POLYMERASE SUBUNIT* genes *RPOA, RPOB, RPOC1* and *RPOC2*,

required for chloroplast gene transcription. Given the dramatic effect on the chloroplast genome, we

wanted to establish whether HXK1 affected seedling greening capacity. This we tested by growing

seedlings in darkness for four days and then exposing them to 3 or 100 μmol m^{-2} s^{-1} white light for 1

hour to promote chlorophyll production. We observed *gin2-1* mutants have significantly higher

chlorophyll (and protochlorophyllide) levels in darkness and following both light treatments,

suggesting chlorophyll production is enhanced in HXK1-deficient seedlings where resources for

growth may be more limited (Fig. 3c).

Transcriptomics data Indicate HXK1 and PIF operate through distinct pathways

As HXK1 and PIFs regulate hypocotyl elongation, albeit through different mechanisms, we wanted to

establish if there was evidence for pathway crosstalk or convergence. By comparing transcriptomic

data from our mRNA-seq with an etiolated, 4-day-old *pifQ* mRNA-seq dataset from Zhang et al

(2013), we discovered that there are 527 genes commonly misregulated between the two mutants: 175

of which are misregulated in the same direction, and 352 which are misregulated in opposite ways

(Fig 3d). Interestingly, for genes that are downregulated in *gin2-1 and pifQ (gin2-1/pifQ down/down)*

we observe enrichment (42 genes, or 24% of genes with synchronous regulation) in lipid storage,

while in the *gin2-1/pifQ* up/up category, there is significant enrichment (18 genes, or 10% of genes

with synchronous regulation) in hypoxia-response genes (Fig. S3). Genes with opposing *gin2-1/pifQ*

down/up regulation are enriched in GO-terms groups for translation, ribosome biogenesis and cell

proliferation. In contrast, opposing *gin2-1/pifQ* up/down category genes include those involved in the

cellular response to starvation and amino acid catabolism. Notably, this latter group includes the

BCAA genes *BCAT2, DIN2, THDP, IVD, MCCA* and *MCCB* that we have identified as HXK1-G6P

regulated (Fig. 2c), all of which show opposing regulation by HXK1 and PIFs (Fig. 3e). We also

noted that the same up/down category included a number of transcription factors such as

PHYTOCHROME INTERACTING FACTOR 5 (PIF5) and *LONG HYPOCOTYL IN FAR-RED*

(HFR1), with pivotal roles in light signalling, *DIG-LIKE 4 (DIL4), ABA 5-BINDING PROTEIN 3*

(AFP3), and *G-BOX BINDING FACTOR 3 (GBF3)*, involved in ABA signalling, *GRETCHEN*

HAGEN 3.12 (GH3.12), the auxin conjugating enzyme, and the *SMALL AUXIN UPREGULATED*

RNA (SAUR) genes *SAUR1 (AT4G34770), SAUR25 (AT4G13790)* and *SAUR42 (AT2G28085)*

(Fig.3f). Thus, HXK1 has a role in repressing the expression of this group of transcription factors that

are promoted by PIFs.

Our mRNA-seq data indicates that in light restricted conditions HXK1 has a role in controlling

nutrient resource management, while our physiological data implicate HXK1-G6P function in

stimulating hypocotyl growth. Previously mutant alleles for the *VHA-B1* and *RPT5B,* which form a

nuclear complex with HXK1, were shown to restrict hypocotyl growth (Cho et al 2006). In

agreement, we found *vha-B1* and *rpt5b* restricted hypocotyl elongation with glucose concentrations as

low as 28mM. However, when grown on media without additional glucose supplement, *vha-B1* and

rpt5b were indistinguishable from WT, while the *hxk1-3* phenotype was retained (Fig. 4a). We also

examined the *king1 (kinγ)* mutant allele for the SNF1-RELATED PROTEIN KINASE

REGULATORY SUBUNIT GAMMA 1, which was recently identified as another interaction partner

for HXK1 (van Dingenen et al, 2019), though neither treatment resulted in a hypocotyl growth defect.

The data suggests HXK1 can regulate hypocotyl elongation through nuclear signaling, but in the

absence of exogenous glucose HXK1 enzymatic function is required.

Earlier reports have implicated HXK1 (VHA-B1 and RPT5B) in glucose repression of *CAB2* and

CAA expression, so we were interested to establish if this regulation was exclusively through the

HXK1 glucose signaling route. In line with these studies, we observe HXK1-dependent repression of

CAB2/CAA when seedlings are grown in the presence of 55mM (1% w/v), but not 28mM (0.5 % w/v)

glucose (Fig. 4b). The 28mM and 55mM application doses lead to 6.5-, and 12.5-fold increases in

internal when glucose levels, while a ~115mM (2% w/v) leads to a 28-fold increase compared to

controls (Fig. 4c). Further, even the lowest concentration (28mM) greatly exceeds glucose levels in

seedlings exposed to the higher light fluence rates of 300 or 600 μmol m^{-2} s^{-1}. We also show that

CAB2 and *CAA* expression is not substantially affected by *gin2-1* following G6P application. Our

data, therefore, lends further support to the notion that *CAB2* and *CAA* are suppressed by HXK1-

VHA-B1-RPT5B in response to exogenous glucose, but they are not targets of the HXK1-G6P

pathway in a physiologically relevant glucose environment.

To further investigate the HXK1 enzymatic role in hypocotyl extension, we next examined the

hxk1^{S177A} substitution allele, which lacks catalytic activity (Moore et al, 2003). The expectation is that

if hypocotyl length (without added glucose) is due to HXK1 enzymatic activity, then *hxk1^{S177A}* would

have a short hypocotyl phenotype. We found that on media without glucose *hxk1^{S177A}* hypocotyls were

indistinguishable from WT, while *hxk1^{S177A}* hypocotyls showed reduced growth compared to WT

plants when glucose is supplemented (Fig. 4d). As this finding is at odds with our other data,

including the *vha-B1, rpt5B* mutant analysis (Fig 4a), we speculated that the effect of *hxk1^{S177A}* may

be masked by a HXK1 glucose-signalling brought about by elevated internal glucose levels. Previous

studies have shown that loss of HXK1 catalytic (in *gin2-1*) activity leads to over-accumulation of

glucose, which we have independently verified for *gin2-1* and *hxk1-3* under our conditions (Fig. 4e).

Further, this is also the case for *hxk1^{S177A}* that has 1.3-fold higher levels than WT (Fig. 4f). Next, we

reasoned that if the *hxk1^{S177A}* hypocotyl elongation results from high internal glucose levels, then

hxk1^{S177A} would exhibit an altered glucose dose-response. In accordance with this hypothesis, our data

shows *hxk1*^{S177A} seedlings have an augmented dose-response that is consistent with elevated internal

glucose levels (Fig. S5).

To consolidate our findings, we analysed the mutant for phosphofructokinase (PFK1), which controls

a rate-limiting step in glycolysis (Mustroph et al, 2007). Our data shows that in low light the *pfk1*

mutant phenocopies the *hxk1-3* short hypocotyl, and as expected *pfk1* is unresponsive to G6P but is

rescued by the application of the glycolytic product, pyruvate (Fig. 4g). These results further reinforce

the important role of the glycolytic pathway in supporting hypocotyl growth in light limiting

conditions.

Discussion

HXK1-mediated hypocotyl elongation is facilitated through its glycolytic role

The seedling hypocotyl is an important model system in *Arabidopsis* for the study of signal

convergence, and growth is an important readout of these signals (Oh et al, 2014, Singh et al, 2017;

Ivakov et al, 2017; Chen et al, 2018). Post-germinative cell expansion requires strong coordination of

these signals and is dependent upon the mobilisation of seed reserves coordinated with signalling

responses (Penfield et al, 2004; Stewart et al, 2011). In this study, we have demonstrated that HXK1

plays an important role in this process via glycolytic action; through seedling establishment, HXK1

operates as a master regulator of the plastome and controls multiple aspects of the metabolic

starvation response. For the most-part HXK1 operates independently of PIF signalling, though a

discrete gene set including HFR1 and PIF5 were identified as possible signal convergence points.

HXK1 has been recognised to have two distinct roles: as a glycolytic enzyme which catalyses the

conversion of glucose to G6P, and as a nuclear-located sugar sensor-signalling molecule (Moore et al,

2003; Cho et al, 2006). Its role in seedling hypocotyl development has previously been ascribed solely

to its nuclear signalling capacity; however, our data shows that when glucose is not supplied

exogenously, HXK1 glycolytic activity is necessary and sufficient to support hypocotyl growth. This

becomes more apparent when light is limited, whether through a shorter light period in diurnal

conditions or with reduced light fluences in both continuous light and diurnal cycles (Fig. 1a-b, Fig.

S2). In contrast to glucose, where dose-response curves are qualitatively similar in WT and *gin2-*

1/*hxk1-3*, G6P selectively restores the *gin2-1*/*hxk1-3* short hypocotyl to wild-type length (Fig. 1f-g,

Fig. S1). These findings emphasise the importance of HXK1 in the breakdown of seed-derived carbon

reserves, which fuel hypocotyl expansion in light-limiting conditions prior to activation of the

photosynthetic machinery. Sucrose production from triacylglycerols and storage proteins feeds into

glycolysis, which in turn allows for the production of energy, carbon skeletons and metabolites that

are required for cell expansion and growth (Penfield et al, 2004, Graham, 2008). Supporting this

notion, we observe HXK1-G6P mediated epidermal cell expansion in seedling hypocotyls (Fig. 1g, 1i,

Fig. S1, Fig. 4g).

Earlier work has implicated PIFs in sucrose-dependent hypocotyl elongation and auxin gene

expression, with PIF5 having a prominent role (Stewart et al, 2011; Lilley et al, 2012). We were

therefore intrigued to establish whether PIFs had a broader role in the integration of light and HXK1

signals to control growth. Our analysis demonstrates that the short hypocotyl *pifQ* mutant is

completely insensitive to G6P, suggesting that G6P is not a limiting factor for growth in *pifQ*.

However, as *pifQ* only represents 4 out of the 8 *PIF* proteins in *Arabidopsis*, there is potential for

other PIF-mediated growth deficiencies operating in a HXK1-G6P responsive way.

HXK1 regulates the carbon stress transcriptome.

When analysing mRNA-seq data we discovered that *gin2-1* has a transcriptome signature that is

reminiscent of sucrose starvation. Earlier studies indicate sucrose starvation leads to strong

upregulation of genes involved in sugar metabolism, photosynthesis, lipid metabolism, cellular

respiration, and downregulation of genes involved lipid storage, cell proliferation and ribosomal

processes (Contento et al, 2004, Nicolai□ et al, 2006). Our study reports a qualitatively similar pattern

in the transcriptome for *gin2-1* (Fig. 2b). Amongst the *gin2-1* upregulated genes are those implicated

in branched-chain amino acid (BCAA) catabolism (Fig. 2b-d), an evolutionary conserved

mitochondrial pathway which can be induced by intense carbon starvation (Binder 2010, Hildebrandt

et al 2015; Heinemann and Hildebrandt, 2021). Activation of the BCAA pathway provides an

alternative source of respiratory substrates that can be utilised during carbohydrate starvation. As G6P

supplementation specifically rescues the expression of these genes, our data indicates *BCAA* genes are

regulated through HXK1-G6P induced signalling. Interestingly, the expression of *BCAA* genes is

known to be directly regulated by SnRK1 kinase-bZIP complexes (Pedrotti et al, 2018). This

evolutionarily conserved protein kinase complex acts to maintain homeostasis under nutrient-stress

conditions. It will therefore be interesting to establish if SnRK1 kinase-bZIP complexes can be

regulated by HXK1-G6P.

In light limiting conditions HXK1 exerts strong control over the plastome.

Our mRNA-seq data reveals a significant effect of HXK1 on the expression of non-nuclear genes. In

dark grown seedlings 82.8% of the chloroplast and 23.5% of the mitochondrial genes were elevated in

gin2-1 (Fig. 3a-c, Table S2). While glucose treatment has been shown to attenuate chlorophyll

degradation in darkness, our data show HXK1 represses chloroplast gene expression in darkness,

which, in turn, suppresses chlorophyll levels following exposure to light (Ueda et al, 2020). Indeed,

the connection between chloroplast biogenesis and metabolism is well established; albino mutants

show high misregulation in genes involved with starvation and mobilisation of storage energy,

consistent with the notion of chloroplast genesis representing a switch of growth strategies to

autotrophism (Grübler et al, 2017). With respect to mitochondria, sucrose starvation has been shown

to cause structural changes, which result in impaired protein production despite unimpeded

transcription activity (Giegé et al, 2005). Additionally, we have shown HXK1 represses the

expression of mitochondrial genes involved in processes including charged particle transport, ATP

synthesis, and oxidative responses. Oxidation in the mitochondria has been linked to metabolism,

feeding back into metabolic processes under both abiotic and biotic stress responses to maintain

signalling capacity (Vanlerberghe, 2013). The impact of *gin2-1* on these organellar transcriptomes

most likely reflects the reality of carbon-resource deprivation, and the need to prime the cellular

machinery to switch to photosynthesis rather than reliance on limited seed reserves to fuel growth.

HXK1 and PIF signalling converges at a small gene subset

Our mRNA-seq analysis identified small group of genes (22% of total *gin2-1* misregulation) that

exhibit similar or opposing regulation by PIF and HXK1 (Fig. 3d-f, Fig. S4). We were interested to

establish that *BCAA* genes, identified as HXK1-repressed genes (Fig 2a-d), were promoted by PIFs.

Likewise, a number of genes implicated in light signalling including *PIF5, HFR1, GH3.12* and *SAUR*

genes, were antagonistically regulated by HXK1 and PIFs. This suggests that while HXK1 and PIF-

mediated signalling is largely distinct, crosstalk does occur through a few key components. Indeed, a

recent study demonstrated such a connection between HXK1 and PIF4 (Kelly et al 2021). The study

showed HXK1 located in guard cells controls hypocotyl elongation through PIF4, in white and blue

light. The authors suggest that in contrast to red light, blue light is not effective in driving

photosynthesis, and therefore in blue light conditions sucrose derived from seed storage is a major

carbon source. The study has strong parallels to our low light study where hypocotyl growth is largely

fuelled by seed reserves. In both instances, HXK1 has an important metabolic role in phosphorylating

the sucrose cleavage products, glucose and fructose. Like HXK1, PIF transcription factors are primed

to operate under low light conditions, and thus offer a route to couple carbon resource management to

growth prior to the establishment of photoautotrophic growth.

HXK1-dependent glucose signalling requires high amounts of glucose to operate.

Previous work has shown *VHA-B1* and *RPT5B* form a nuclear complex with HXK1, which directly

couples glucose sensing to nuclear signalling and hypocotyl growth (Cho et al, 2006). In agreement

with this notion, we observe a hypocotyl defect in *vha-B1, rpt5b* mutants, but only when exogenous

glucose is supplied. In contrast the *gin2-1/hxk1-3* short hypocotyl phenotype is observed both with

and without glucose (Fig. 4a). Collectively the data indicate hypocotyl elongation can be controlled

through HXK1 glycolytic and sensor-signalling pathways, however, prior to the establishment of

photoautotrophic growth the glycolytic pathway necessarily dominates. These observations are

backed by qPCR analysis of *CAB2* and *CAA1*, both HXK1/VHA-B1- /RPT5B-dependent glucose

repressed genes. Interestingly, we only observed HXK1-mediated repression at 55mM glucose (which

elevates internal glucose levels by 12.5-fold) and not the lower concentration of 28mM glucose (6.5-

fold increase) (Fig. 4b). This suggests HXK1-dependency is only observed when glucose levels

exceed normal physiological levels, which may be more indicative of a stress response. The ABA

signalling genes *ABI4* and *ABI5* have been shown to have key roles in abiotic stress and the glucose

signalling response (Dijkwel et al, 1997, Arenas-Huertero et al, 2000). As ABI4 is known to directly

regulate the expression of *CAB2* it will be interesting to establish whether the HXK1-VHA-BI-RPT5B

complex operates with ABI4 or other ABA signalling components.

Our study provides strong evidence for HXK1 glycolytic control of hypocotyl growth, yet

counterintuitively, the $hxk1^{S177A}$ catalytically inactive mutant does not have the expected short

hypocotyl phenotype in low light (Moore et al, 2003; Fig. 4d, S4). We therefore speculated that in a

similar manner to *gin2-1* and *hxk1-3*, loss of enzymatic activity could lead to enhanced internal

glucose levels. This would effectively activate glucose signalling and hypocotyl elongation, which

would effectively mask any growth impairment caused by loss of enzymatic function (Fig. 4e-f,). Our

analysis, which revealed elevated glucose levels in $hxk1^{S177A}$ and a shifted glucose dose-response,

provided support for this hypothesis, offering a rational explanation for the $hxk1^{S177A}$ phenotype.

Corroborative evidence for the importance of glycolysis, comes from genetic analysis demonstrating

the glycolytic enzyme mutant *pfk1-1* phenocopies *hxk1-3*, while both mutants are rescued by

exogenous pyruvate application (Fig. 4g).

In conclusion, this study has shown HXK1 performs an important role in post-germinative hypocotyl

growth, by enabling the consumption of seed reserves, prior to establishment of photosynthetic

competence. By controlling the expression of enzymes in the BCAA pathway, HXK1 provides a route

to generate alternative respiratory substrates when carbon resources are limited. HXK1 control of the

plastome, provides a means to link photosynthetic machinery establishment to carbon needs. While

HXK1 control of light signalling components provides a potential route to couple resource availability

to the regulation of growth.

Acknowledgements

M.L. and A.G. were supported by the University of Edinburgh Darwin Trust Scholarship. A.R was

supported by Biotechnology and Biological Sciences Research Council (BBSRC) grant

BB/N005147/1, awarded to K.J.H.)

Author Contributions

M.J.L., A.G., A.R. and K.J.H. planned and designed the research. A.G designed and performed, and

analysed the mRNA-seq experiment, A.R. conducted the bioinformatics analysis of the mRNAseq

data. All other experiments were performed and analysed by M.J.L and A.G. All authors contributed

to manuscript preparation.

References

Arenas-Huertero F, Arroyo A, Zhou L, Sheen J, León P. 2000**.** Analysis of Arabidopsis glucose
insensitive mutants, gin5 and gin6, reveals a central role of the plant hormone ABA in the
regulation of plant vegetative development by sugar. *Genes Dev*. **14**(16): 2085-96.
Baud S, Dubreucq B, Miquel M, Rochat C, Lepiniec L. 2008**.** Storage reserve accumulation in
Arabidopsis: metabolic and developmental control of seed filling. *Arabidopsis Book* **6**: e0113.
Bian Y, Chu L, Lin H, Qi Y, Fang Z, Xu D. 2022**.** PIFs- and COP1-HY5-mediated temperature
signaling in higher plants. *Stress Biology* **2**(1): 35.
Binder S. 2010**.** Branched-Chain Amino Acid Metabolism in Arabidopsis thaliana. *Arabidopsis*
Book **8**: e0137.
Boron AK, Vissenberg K. 2014 The Arabidopsis thaliana hypocotyl, a model to identify and
study control mechanisms of cellular expansion. *Plant Cell Rep*. **33**(5):697-706.
Chen GH, Liu MJ, Xiong Y, Sheen J, Wu SH. 2018**.** TOR and RPS6 transmit light signals to
enhance protein translation in deetiolating Arabidopsis seedlings. *Proc Natl Acad Sci U S A*
115(50): 12823-12828.
Cho YH, Yoo SD, Sheen J. 2006**.** Regulatory functions of nuclear hexokinase1 complex in
glucose signaling. *Cell* **127**(3): 579-589.
Claeyssen E, Rivoal J. 2007**.** Isozymes of plant hexokinase: occurrence, properties and functions.
Phytochemistry **68**(6): 709-731.
Contento AL, Kim SJ, Bassham DC. 2004**.** Transcriptome profiling of the response of
Arabidopsis suspension culture cells to Suc starvation. *Plant Physiol* **135**(4): 2330-2347.
de Wit, M., Martınez-Cero, N. C., Fankhauser, C. & Pierik, R. 2016 Integration of phytochrome
and cryptochrome signals determines plant growth during competition for light. Curr. Biol.
26(24)**:** 3320–3326.
Dijkwel PP, Huijser C, Weisbeek PJ, Chua NH, Smeekens SC. 1997**.** Sucrose control of
phytochrome A signaling in Arabidopsis. *Plant Cell*. **9**(4): 583-95.
Eastmond PJ, Germain V, Lange PR, Bryce JH, Smith SM, Graham IA. 2000**.** Postgerminative
growth and lipid catabolism in oilseeds lacking the glyoxylate cycle. *Proc Natl Acad Sci U S A*
97(10): 5669-5674.
Feng J, Zhao S, Chen X, Wang W, Dong W, Chen J, Shen JR, Liu L, Kuang T. 2015**.**
Biochemical and structural study of Arabidopsis hexokinase 1. *Acta Crystallogr D Biol*
Crystallogr **71**(Pt 2): 367-375.
Franklin KA, Quail PH. 2010 Phytochrome functions in Arabidopsis development. *J Exp Bot*.
61(1):11-24.
Gendreau E, Traas J, Desnos T, Grandjean O, Caboche M, Hofte H. 1997**.** Cellular basis of
hypocotyl growth in Arabidopsis thaliana. *Plant Physiol* **114**(1): 295-305.
Giegé P, Sweetlove LJ, Cognat V, Leaver CJ. 2005**.** Coordination of nuclear and mitochondrial
genome expression during mitochondrial biogenesis in Arabidopsis. *Plant Cell* **17**(5): 1497-1512.
Granot D. 2007**.** Role of tomato hexose kinases. *Functional Plant Biology* **34**: 564-570
Graham IA. 2008**.** Seed storage oil mobilization. *Annu Rev Plant Biol* **59**(1): 115-142.
Grübler B, Merendino L, Twardziok SO, Mininno M, Allorent G, Chevalier F, Liebers M,
Blanvillain R, Mayer KFX, Lerbs-Mache S, et al. 2017**.** Light and Plastid Signals Regulate
Different Sets of Genes in the Albino Mutant Pap7-1. *Plant Physiol* **175**(3): 1203-1219.
Harrington GN, Bush DR. 2003**.** The bifunctional role of hexokinase in metabolism and glucose
signaling. *Plant Cell* **15**(11): 2493-2496.
Heinemann B, Hildebrandt TM. 2021**.** The role of amino acid metabolism in signaling and
metabolic adaptation to stress-induced energy deficiency in plants. *J Exp Bot* **72**(13): 4634-4645.
Hildebrandt TM, Nunes Nesi A, Araujo WL, Braun HP. **2015.** Amino Acid Catabolism in Plants.
Mol Plant **8**(11): 1563-1579.

Hwang G, Kim S, Cho JY, Paik I, Kim JI, Oh E. 2019**.** Trehalose-6-phosphate signaling regulates thermoresponsive hypocotyl growth in Arabidopsis thaliana. *EMBO Rep* **20**(10): e47828.

Ivakov A, Flis A, Apelt F, Funfgeld M, Scherer U, Stitt M, Kragler F, Vissenberg K, Persson S, Suslov D. 2017**.** Cellulose Synthesis and Cell Expansion Are Regulated by Different Mechanisms in Growing Arabidopsis Hypocotyls. *Plant Cell* **29**(6): 1305-1315.

Jang JC, Sheen J. 1994**.** Sugar sensing in higher plants. *Plant Cell* **6**(11): 1665-1679.

Jang JC, León P, Zhou L, Sheen J. **1997.** Hexokinase as a sugar sensor in higher plants. *Plant Cell* **9**(1): 5-19.

Kanehisa M, Goto S, Sato Y, Furumichi M, Tanabe M. 2012**.** KEGG for integration and interpretation of large-scale molecular data sets. *Nucleic Acids Res* **40**(Database issue): D109-114.

Karve A, Rauh BL, Xia X, Kandasamy M, Meagher RB, Sheen J, Moore BD. **2008.** Expression and evolutionary features of the hexokinase gene family in Arabidopsis. *Planta* **228**(3): 411-425.

Kelly G, Brandsma D, Egbaria A, Stein O, Doron-Faigenboim A, Lugassi N, Belausov E, Zemach H, Shaya F, Carmi N, et al. 2021**.** Guard cells control hypocotyl elongation through HXK1, HY5, and PIF4. *Communications Biology* **4**(1): 765.

Kim D, Paggi JM, Park C, Bennett C, Salzberg SL. 2019**.** Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**(8): 907-915.

Kolesnikov N, Hastings E, Keays M, Melnichuk O, Tang YA, Williams E, Dylag M, Kurbatova N, Brandizi M, Burdett T, et al. 2015**.** ArrayExpress update--simplifying data submissions. *Nucleic Acids Res* **43**(Database issue): D1113-1116.

Krahmer J, Abbas A, Mengin V, Ishihara H, Romanowski A, Furniss JJ, Moraes TA, Krohn N, Annunziata MG, Feil R, et al. 2021 Phytochromes control metabolic flux, and their action at the seedling stage determines adult plant biomass. *J Exp Bot*. **72**(8):3263-3278.

Leivar P, Monte E, Al-Sady B, Carle C, Storer A, Alonso JM, Ecker JR, Quail PH. 2008**.** The Arabidopsis Phytochrome-Interacting Factor PIF7, Together with PIF3 and PIF4, Regulates Responses to Prolonged Red Light by Modulating phyB Levels. *The Plant Cell* **20**(2): 337-352.

Lilley JLS, Gee CW, Sairanen I, Ljung K, Nemhauser JL. 2012**.** An Endogenous Carbon-Sensing Pathway Triggers Increased Auxin Flux and Hypocotyl Elongation   *Plant Physiology* **160**(4): 2261-2270.

Liu Z, Zhang Y, Liu R, Hao H, Wang Z, Bi Y. 2011**.** Phytochrome interacting factors (PIFs) are essential regulators for sucrose-induced hypocotyl elongation in Arabidopsis. *Journal of Plant Physiology* **168**(15): 1771-1779.

Lun AT, Chen Y, Smyth GK. 2016**.** It's DE-licious: A Recipe for Differential Expression Analyses of RNA-seq Experiments Using Quasi-Likelihood Methods in edgeR. *Methods Mol Biol* **1418**: 391-416.

Mancini E, Rabinovich A, Iserte J, Yanovsky MC, Ariel   2021**.** ASpli: Integrative analysis of splicing landscapes through RNA-Seq assays. *Bioinformatics* **37**(17): 2609-2616.

Moore B, Zhou L, Rolland F, Hall Q, Cheng W-H, Liu Y-X, Hwang I, Iones T, Sheen J. 2003**.** Role of the Arabidopsis glucose sensor HXK1 in nutrient, light, and hormonal signalling. *Science* **300**(5617): 332-336.

Mustroph A, Sonnewald U, Biemelt S. 2007**.** Characterisation of the ATP-dependent phosphofructokinase gene family from Arabidopsis thaliana. *FEBS Letters* **581**(13): 2401-2410.

Nicolaï M, Roncato MA, Canoy AS, Rouquié D, Sarda X, Freyssinet G, Robaglia C. 2006**.** Large-Scale Analysis of mRNA Translation States during Sucrose Starvation in Arabidopsis Cells Identifies Cell Proliferation and Chromatin Structure as Targets of Translational Control. *Plant Physiology* **141**(2): 663-673.

Oh E, Zhu JY, Bai MY, Arenhart RA, Sun Y, Wang ZY. 2014. Cell elongation is regulated through a central circuit of interacting transcription factors in the Arabidopsis hypocotyl. *eLife* **3**: e03031.

Park E, Park J, Kim J, Nagatani A, Lagarias JC, Choi G. 2012**.** Phytochrome B inhibits binding of phytochrome-interacting factors to their target promoters. *Plant J* **72**(4): 537-546.

Park EK, Yeojae Choi, Giltsu. 2018**.** Phytochrome B Requires PIF Degradation and Sequestration to Induce Light Responses across a Wide Range of Light Conditions. *Plant Cell* **30**(6): 1277-1292.

Pedrotti L, Weiste C, Nägele T, Wolf E, Lorenzin F, Dietrich K, Mair A, Weckwerth W, Teige M, Baena-González E, et al. 2018**.** Snf1-RELATED KINASE1-controlled C/S1-bZIP signaling activates alternative mitochondrial metabolic pathways to ensure plant survival in extended darkness. *The Plant Cell* **30**(2): 495-509.

Penfield S, Rylott EL, Gilday AD, Graham S, Larson TR, Graham IA. 2004**.** Reserve Mobilization in the Arabidopsis Endosperm Fuels Hypocotyl Elongation in the Dark, Is Independent of Abscisic Acid, and Requires PHOSPHOENOLPYRUVATE CARBOXYKINASE1. *Plant Cell* **16**(10): 2705-2718.

Pfeiffer A, Janocha D, Dong Y, Medzihradszky A, Schöne S, Daum G, Suzaki T, Forner J, Langenecker T, Rempel E, et al. 2016**.** Integration of light and metabolic signals for stem cell activation at the shoot apical meristem. *eLife* **5**: e017023.

Pires MV, Pereira Júnior AA, Medeiros DB, Daloso DM, Pham PA, Barros KA, Engqvist MK, Florian A, Krahnert I, Maurino VG, et al. 2016**.** The influence of alternative pathways of respiration that utilize branched-chain amino acids following water shortage in Arabidopsis. *Plant Cell Environ* **39**(6): 1304-1319.

Robinson MD, McCarthy DJ, Smyth GK. 2010**.** edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**(1): 139-140.

Rolland F, Moore B, Sheen J. 2002**.** Sugar Sensing and Signaling in Plants. *Plant Cell* **14**(Suppl): s185-205.

Sairanen I, Novák O, Pěnčík A, Ikeda Y, Jones B, Sandberg G, Ljung K. 2013**.** Soluble Carbohydrates Regulate Auxin Biosynthesis via PIF Proteins in Arabidopsis. *The Plant Cell* **24**(12): 4907-4916.

Singh M, Gupta A, Singh D, Khurana JP, Laxmi A. 2017**.** Arabidopsis RSS1 Mediates Cross-Talk between Glucose and Light Signaling during Hypocotyl Elongation Growth. *Sci Rep* **7**: 16101.

Stewart JL, Maloof JN, Nemhauser JL. 2011**.** PIF genes mediate the effect of sucrose on seedling growth dynamics. *PLOS ONE* **6**(5): e19894.

Ueda H, Ito T, Inoue R, Masuda Y, Nagashima Y, Kozuka T, Kusaba M. 2020**.** Genetic Interaction Among Phytochrome, Ethylene and Abscisic Acid Signaling During Dark-Induced Senescence in Arabidopsis thaliana. *Frontiers in Plant Science* **11**.

van Dingenen J, Vermeersch M, De Milde L, Hulsmans S, De Winne N, Van Leene J, Gonzalez N, Dhondt S, De Jaeger G, Rolland F, et al. 2019**.** The role of HEXOKINASE1 in Arabidopsis leaf growth. *Plant Molecular Biology* **99**(1): 79-93.

Vanlerberghe GC. 2013**.** Alternative Oxidase: A Mitochondrial Respiratory Pathway to Maintain Metabolic and Signaling Homeostasis during Abiotic and Biotic Stress in Plants. *International Journal of Molecular Sciences* **14**(4): 6805-6847.

Yang DS, Daniel D., Krahmer J, Halliday KJ. 2016**.** Photoreceptor effects on plant biomass, resource allocation, and metabolic state. *PNAS* **113**(27): 7667-7672.

Yang Y, Benning C. 2018**.** Functions of triacylglycerols during plant development and stress. *Current Opinion in Biotechnology* **49**: 191-198.

Yoon J, Cho LH, Tun W, Jeon JS, An G. 2021**.** Sucrose signaling in higher plants. *Plant Science* **302**: 110703.

Yu G, Wang LG, Han Y, He QY. 2012. clusterProfiler: an R package for comparing biological themes among gene clusters. *Omics* **16**(5): 284-287.

Zhang R, Calixto CPG, Marquez Y, Venhuizen P, Tzioutziou NA, Guo W, Spensley M, Entizne JC, Lewandowska D, Have ST, et al. 2017**.** A high quality Arabidopsis transcriptome for accurate transcript-level analysis of alternative splicing. *Nucleic Acids Research* **45**(9): 5061-5073.

Zhang Y, Mayba O, Pfeiffer A, Shi H, Tepperman JM, Speed TP, Quail PH. 2013**.** A Quartet of PIF bHLH Factors Provides a Transcriptionally Centered Signaling Hub That Regulates Seedling Morphogenesis through Differential Expression-Patterning of Shared Target Genes in Arabidopsis. *PLOS Genetics* **9**(1): e1003244.

Zhang R, Calixto CPG, Marquez Y, Venhuizen P, Tzioutziou NA, Guo W, Spensley M, Entizne JC, Lewandowska D, Ten Have S, et al. 2017**.** A high quality Arabidopsis transcriptome for accurate transcript-level analysis of alternative splicing. *Nucleic Acids Res* **45**(9): 5061-5073.

Fig. 1 Analysis of the HXK1-dependent hypocotyl growth phenotype and dependence on G6P production (a) White light fluence response. Seedlings under continuous white light in decreasing fluences. (b) Images of seedlings. Seedlings were grown in low light (left), or high light (right). (c) Cotyledon Area. Continuous light seedlings. Numbers above represent fluence rate in $m^{-2}s^{-1}$. (d) Cell size of *gin2-1* etiolated seedlings. Etiolated seedlings. (e) Cell file of *gin2-1* etiolated seedlings. Etiolated seedlings. (f) Glucose dose-response curve. Etiolated seedlings with increasing amounts of glucose. (g) G6P dose-response curve. Etiolated seedlings with increasing amounts of G6P. (h) Competitive treatment of HXK1 inhibitor molecule mannoheptulose and glucose-6-phosphate. Short day low light seedlings without or with G6P, mannoheptulose (MHL), and both. (i) Cell size and count of *pifQ* seedlings treated with G6P. Short day low light without (left) or with 5mM G6P (right).

Fig. 2 *gin2-1* sequencing data reveals importance of HXK1 in translation and carbon stress metabolism (a) mRNAseq volcano plot. Yellow points represent logFC≥0.5 and blue points represent logFC≤-0.5, pval≤0.01. Red line marks pval≤0.01. (b) Bubble plot of selected GO terms in mRNAseq. Genes of interest were collected and displayed as described in the materials and methods. (c) Diagram of branched chain amino acid metabolism in Arabidopsis. Adapted from Neinast et al, 2018 with genes represented in (d) highlighted. (d) qPCR expression of core BCAA genes misregulated in *gin2-1*. Transcript abundance of BCAA pathway genes in etiolated WT and *gin2-1* seedlings grown with or without 0.5% w/v glucose or 0.125% w/v G6P. Numbers above represent fold-change between *gin2-1* and WT.

Fig. 3 mRNAseq of *gin2-1-1* implicate a significant role in regulating the plastid genome and demonstrate crosstalk with PIF-dependent signalling (a) Percentage of chromosome misregulation in *gin2-1*. Numbers above bar represent the percentage of genes significantly (P≤ 0.01) misregulated. (b) qPCR expression of RPO family genes, collected at ZT4. (c) Chlorophyll content of *gin2-1* and WT seedlings. Seedlings grown in darkness (left) or transferred to either 3 (middle) or 100 (right)100μmol m^{-2}s^{-1} white light for 1 hour. (d) Transcriptome comparison between *gin2-1* and *pifQ*. Genes significantly misregulated in *gin2-1* were compared to genes significantly misregulated in Zhang et al (2013, were categorized as being either up or downregulated in either *pifQ* or *gin2-1* and sorted into overlap and exclusive subcategories. Numbers above the bars represent count. (e) BCAA genes representation in *gin2-1* and *pifQ*. (f) Genes of interest comparison between *gin2-1* and *pifQ* with published correlations with PIF. For all figures except (c), seedlings were grown in darkness.

Fig. 4 HXK1-dependent glucose signaling requires high amounts of exogenous glucose to regulate expansion and genetic regulation (a) HXK1 signaling complex mutants grown without and with additional glucose. Short day seedlings in low light, without or with 28mM glucose. (b) Expression of CAB2 and CAA in *gin2-1* with added substrate. Transcript abundance of CAB2 (left) and CAA (right) relative to PP2A concentration in Short day seedlings in high light, without or with 28mM glucose, 55mM glucose, or 5mM G6P. Plants sampled at 4 hours after dawn (ZT4). (c) Internal glucose concentrations of Wt plants under variable light and nutrient conditions. Glucose measured at ZT 4 in Wt seedlings grown in 100, 300, and 600 µmol/m²s, or at high light with increasing glucose. Numbers above horizontal bars represent the fold change. (d) *hxk1S177A* seedlings grown with and without glucose. Etiolated seedlings without or with glucose. (e) Internal glucose concentration comparison between Wt and *gin2-1* seedlings. Glucose measured at ZT in etiolated seedlings. (f) Internal glucose concentration of *hxk1S177A* seedlings. Glucose measured at ZT4 in etiolated seedlings. (g) Glycolysis component mutant *pfk1* grown without or with G6P or sodium pyruvate (NaPyr) in 3 µmol/m²s WL.