HEXOKINASE 1 Glycolytic Action Fuels Post-Germinative Seedling Growth

Ashwin Ganpudi[1], Andrés Romanowski[1], Karen J. Halliday[1]

1. Institute for Molecular Plant Sciences, School of Biological Sciences, University of

Edinburgh, Edinburgh EH9 3BF, United Kingdom.

ORCID IDs: 0000-0002-9515-3899 (A.G.); 0000-0003-0737-2408 (A.R.); 0000-0003-0467-

104X (K.J.H.)

Corresponding author:

Karen J. Halliday

Institute for Molecular Plant Sciences, School of Biological Sciences, University of

Edinburgh, Edinburgh EH9 3BF, United Kingdom

+44-0131-651-9083

karen.halliday@ed.ac.uk

Abstract

HEXOKINASE 1 (HXK1) is an evolutionarily conserved glycolytic enzyme that has a

separate nuclear glucose-sensor signalling function. In Arabidopsis, HXK1 mutant seedlings

have a stunted growth defect, yet how this relates to HXK1 function remains inconclusive.

We show that the HXK1-glycolytic pathway performs a fundamental role in catabolising seed

reserves to fuel post-germinative cell division and expansion. This function is particularly

important in dim light which delays the switch to photoautotrophic growth. RNAseq analysis

reveals that HXK1 imposes strong repressive control on plastome gene expression, and

regulates nuclear-encoded genes that drive energy-consuming processes. Earlier studies

have implicated HXK1 sensor-signalling in the repression of *CAB2* and *CAA* gene

expression by exogenous glucose. We show that over a wide range of irradiances this

pathway is not operative in seedlings. Our study therefore defines a new operational model

for HXK1 action in catabolising carbon resources and tuning gene expression to optimize

seedling growth.

Introduction

The first days of life are critical for plant survival. Successful seedling establishment relies on

seed reserves to support emergence before the switch to photosynthetic growth. The

endosperm of oil seed plants, such as Arabidopsis, contains mainly triacylglycerols (TAG)

and storage proteins, which are mobilized during germination (**Baud et al., 2005; Penfield**

et al., 2006). Several studies have highlighted the critical role for TAGs in gluconeogenic

sugar production, which fuels post-germinative seedling growth (**Graham, 2008;**

Theodoulou and Eastmond, 2012). When germination occurs in darkness, or dim light

seedlings adopt a skotomorphogenic-type program where growth is largely confined to the

seedling stem (hypocotyl), at the expense of seedling leaf (cotyledon) and root development.

A large body of work has shown that members of the light regulated PHYTOCHROME

INTERACTING FACTOR (PIF) gene family operate cooperatively to regulate

skotomorphogenesis (**Gommers and Monte, 2018; Leivar and Monte, 2014; Chaiwanon**

et al., 2016). This growth strategy allows seedlings to forage for light when occluded by soil,

debris or vegetation cover. It is energy-conserving, as it restricts growth to one organ,

reducing the pull-on finite seed reserves.

Following gluconeogenesis, glucose is either used as a building block for protein, fatty acid

or cellulose production, or it is catabolized to generate energy for cellular metabolism and

growth. A fundamental enzyme in this latter process is the evolutionarily conserved

HEXOKINASE 1 (HXK1) that catalyses the first glycolytic step of glucose phosphorylation to

generate glucose-6-phosphate (G6P) (**Cárdenas et al., 1998; Claeyssen and Rivoal,**

2007). In addition to its enzymatic role, Arabidopsis HXK1 is reported to have a conserved

glucose activated signalling function where HXK1 operates in a nuclear-located complex

with VACUOLAR H(+)-ATPase B1 (VHA-B1) and the 19S regulatory particle of proteasome

subunit (RPT5B) (**Yanagisawa et al., 2003; Moore et al., 2003; Cho et al., 2007**). The

HXK1 complex was shown to be necessary for glucose-induced repression of

CHLOROPHYLL A/B BINDING PROTEIN 2 (*CAB2)* and *CARBONIC ANHYDRASE (CAA).*

ChIP-qPCR analysis indicated that at least for *CAB2* this transcriptional suppression was

direct. This sugar-dependent, nuclear regulatory role of HXK1 was postulated to function

during feedback inhibition of photosynthesis (*Moore et al., 2003; Cho et al., 2007*). Other

reports have also implicated HXK1 in the transcriptional repression of developmental genes

in response to exogenous sugar application (*Yang et al., 2013; Yu et al., 2013; de Jong et*

al., 2014; Hsu et al., 2014; Kunz et al., 2015).

A prominent feature of the *glucose insensitive 2 (gin2)* mutant that lacks HXK1, is its

diminutive seedling phenotype (*Moore et al., 2003; Cho et al., 2007*). Earlier work showing

mutated catalytic domain constructs, HXK1^{S177A} and HXK1^{G104D}, could complement the *gin2*

seedling phenotype (**Moore et al., 2003**) suggested *gin2* stunting arose from impaired

glucose-activated HXK1 signalling, and not HXK1 enzyme activity. However, several reports

present data that are incongruous with this interpretation, so the precise role of HXK1 in

seedlings remains unclear. Our study establishes that the HXK1-glycolytic pathway fulfils an

important function in supporting early seedling growth. This is particularly important during

light limited conditions that attenuate the switch to photoautotrophic growth. Under

physiologically relevant conditions, glucose repression of the photosynthetic genes *CAB2*

and CAA does not require HXK1. Rather, HXK1 boosts the expression of genes involved in

aerobic respiration and energy-consuming processes and strongly represses the Branched

Chain Amino Acid (BCAA) alternative respiratory pathway and the plastome. Thus, HXK1

plays a central role not just in glycolysis but in coordinating the transcriptional regulation of

cellular metabolism in developing seedlings.

Results

HXK1 supports post germination growth in darkness and low light. The *gin2* mutant

was previously reported to have impaired hypocotyl growth in low light and nutrient

conditions (*Moore et al., 2003; Cho et al., 2007*). We therefore wanted to establish the

fluence rate range in which HXK1 operates. This was assessed by growing HXK1 mutants

gin2 (Ler) and *hxk1-3* (Col) in darkness or increasing irradiances of continuous white light.

Our data show both *gin2* and *hxk1-3* exhibit shorter hypocotyls than their respective isogenic

wild types in darkness and low fluence rates (**Figure 1A**). Likewise, *gin2* and *hxk1-3*

cotyledon expansion is impaired at low, but not high fluence rates (**Figure 1B,C**). This

indicates that HXK1 has an important role in supporting seedling growth in darkness and

light limiting conditions.

In nature, plants do not commonly experience uninterrupted continuous light, so we wanted

to establish whether the *gin2* / *hxk1-3* seedling phenotypes were evident in photoperiodic

conditions. We found that this was indeed the case in Short Days (SD), but not in Long Days

(LD). However, by lowering the fluence rate from 100 to 5 µmol m^{-2} s^{-1} in LD we were able to

restore the *gin2* / *hxk1-3* short hypocotyl phenotype (**Figure1-figure supplement 1**). These

data illustrate that HXK1 is required for hypocotyl extension in short photoperiods or in LDs

when access to light is restricted.

We next quantified hypocotyl epidermal cell length and number to establish the basis of the

hypocotyl defect in *gin2*. On day 4 *gin2* had less elongated cells, and this was most

noticeable in the upper elongation region (34% decrease) (**Figure 1D**). We also established

that on day 4 and day 14, after growth cessation, *gin2* hypocotyls had fewer epidermal cells

than wild type (**Figure 1E**). Interestingly, we also observed that in darkness *gin2* mutants

had 55% higher levels of glucose than wild type (**Figure 1F**), which is indicative of impaired

HXK1 catalytic function. Supporting this notion, G6P but not glucose was able to rescue the

gin2 short etiolated hypocotyl phenotype (**Figure 1G**). Application of sodium pyruvate (the

salt of glycolytic end product pyruvate) was nearly as effective as G6P (**Figure1-figure**

supplement 2). Thus, our data imply HXK1 catalytic activity has an important function in

supporting cell proliferation and expansion in the seedling hypocotyl under light-restricted

conditions.

RNA-seq reveals role for HXK1 in nutrient resource management. To gain an

appreciation of how HXK1 influences gene expression we performed RNA-seq on 4-day-old

etiolated WT and *gin2* seedlings. This data revealed that 2353 genes were mis-regulated (±

1.5 FC) in *gin2* vs wild type, with 1276 genes down regulated, and 1077 upregulated (***Figure***

2-figure supplement 1, ***Figure 2-source data 1***). Amongst the downregulated subset we

noted a predominance of energy demanding pathways, for instance over-represented GO

terms included ribosome biogenesis, protein synthesis, translation, ATP driven cytoskeletal

motor proteins, microtubules (kinesins) and microfilaments (myosins) (***Figure 2A***).

Cytoskeletal motor proteins require microtubules and microfilaments to mobilize the

organization of various cytoskeletal arrays during cell division, cell expansion and cell growth

in plant tissues (***Hashimoto, 2015***). Indeed, several *CYCLIN (CYC)* genes that regulate cell

cycle progression and cell wall remodelling *XYLOGLUCAN ENDOTRANSGLUCOSYLASE /*

HYDROLASE (XTH) genes are also down-regulated in *gin2* (***Figure 2-figure supplement***

2). Collectively, these observations are consistent with the *gin2* phenotype that is

characterized by a reduction of cell division, expansion and growth (***Figure 1D,E***).

Amongst the upregulated transcript subset, enriched processes include abscisic acid and

auxin response pathways, water and photoperception responses. However, by far the most

prominent of these categories is of the branched chain amino acids (BCAA), leucine (Leu),

isoleucine (Ile) and valine (Val) (***Figure 2B***). BCAA degradation is an evolutionary conserved

mitochondrial pathway induced by intense carbon starvation (***Binder, 2010; Hildebrandt et***

al., 2015) (Fig. 2C). Oxidation of BCAAs directly feeds electrons into the electron transport

chain. The carbon skeletons are further converted to precursors of the tricarboxylic acid

(TCA) cycle through a series of biochemical steps for subsequent ATP production. Hence,

enhanced regulation of this pathway in *gin2* could help support cellular respiration and

subsequently energy production, as glycolysis is impaired.

Our qRT-PCR data show that 0.5% glucose application suppresses the expression of BCAA

catabolic pathway genes (*BRANCHED-CHAIN AMINO ACID TRANSAMINASE 2 (BCAT2),*

DARK INDUCIBLE 2 (DIN2), THIAMIN DIPHOSPHATE-BINDING FOLD PROTEIN (THDP),

ISOVALERYL-COA-DEHYDROGENASE (IVD) and *3-METHYLCROTONYL-COA*

CARBOXYLASE subunits (MCCA and MCCB) in both wild type and *gin2* seedlings (***Figure***

2D). Thus, glucose suppression of *BCAA* genes does not require HXK1. In contrast, G6P

application was ineffective in wild type, but fully restored the elevated expression of *BCAA*

genes in *gin2* to wild type levels (**Figure 2D**). As G6P can substitute for HXK1 deficiency this

suggests that reduced glycolysis in *gin2* leads to the activation of the BCAA pathway as an

alternative energy source for respiration.

HXK1 represses the plastome in the dark. An interesting observation from our RNA-seq

data is that HXK1 loss leads to a dramatic upregulation of 94% chloroplast-encoded and

24% mitochondrial genes in etiolated seedlings (**Figure 3A**). Chloroplastic genes represent

the most highly upregulated category in *gin2*, suggesting HXK1 has an important role in

global suppression of the chloroplast genome in darkness. This finding provided an

opportunity to test whether HXK1 is also able to regulate non-nuclear genes through the

G6P pathway. Here we chose to analyse the chloroplast-encoded *RNA POLYMERASE*

SUBUNIT genes *RPOA, RPOB, RPOC1* and *RPOC2* that are required for chloroplast gene

transcription. We found that G6P induced expression of *RPOA, RPOB, RPOC1* and *RPOC2*

in both wild type and *gin2*. However, as transcript levels were already elevated in *gin2*, its

G6P-induced response was attenuated (**Figure 3B**). These results suggest that while HXK1

most likely regulates these plastid genes through a G6P-dependent pathway, it is not the

sole regulator. It is possible that HXK1 homologue HXK2 or the chloroplast located HXK3

operate redundantly with HXK1 to regulate these plastid genes. Collectively, these data

provide genetic evidence that HXK1 regulates plastid-encoded as well as nuclear-encoded

genes most likely by enzymatic catalysis of the first step in glycolysis that yields G6P.

HXK1 regulates a small subset of PIF-targets. Similar to *gin2* and *hxk1-3*, *pif* mutations

can impair hypocotyl elongation in etiolated seedlings, and this effect is pronounced in the

higher order *pifQ* mutant that lacks *PIF1, PIF3, PIF4* and *PIF5* (**Leivar et al., 2008; 2012**).

We therefore wanted to establish the degree of overlap between the HXK1 and PIF

regulated transcriptomes by comparing RNA-seq data from our study (4-day old etiolated

gin2 seedlings) and a published dataset (2-day old etiolated *pifQ* mutants) (**Pfeiffer et al.,**

2014). In total, 4223 (± 1.5 FC) genes were misregulated in *pifQ*/wt compared to 2353 (± 1.5

FC) in *gin2*/Wt, with 748 common genes, which represents 11% of the total gene population,

but 32% of HXK1-regulated genes (**Figure 3C**). Interestingly, like *gin2*, the *pifQ*

transcriptome was enriched for BCAA catabolic pathway genes. However, while BCAA

genes *BCAT2, DIN2, THDP, IVD, MCCA, MCCB* are elevated in *gin2*, they are severely

suppressed in the *pifQ* mutant (**Figure 3-figure supplement 1**). As the promoters of these

genes are not identified as PIF-bound in these conditions (**Pfeiffer et al. 2014; Zhang et al.,**

2013), this suggests that the differences in *BCAA* gene expression in *pifQ* and *gin2* could

reflect differences in their respective metabolic states.

Next, we examined the likelihood of HXK1-PIF cross-talk by identifying the common gene

subset that are directly bound by PIFs based on ChIP-seq analysis (**Pfeiffer et al., 2014**).

Interestingly, this subset represents only 4.6% of common genes, suggesting that potential

cross-talk may be limited. By far the largest groups of genes were those upregulated by both

HXK1 and PIFs which are predominantly involved in metabolism (**Figure 3-figure**

supplement 2). In contrast, direct PIF targets that are HXK1-repressed are primarily

transcription factors and hormone regulated genes. Shared genes include *PIL2 (PIF6)* and

LONG HYPOCOTYL IN FAR-RED (*HFR1*), two well characterized PIF-targets (**Pfeiffer et**

al., 2014; Zhang et al., 2013; Hornitschek et al., 2012). Thus, we were keen to establish if

these genes are regulated by the HXK1-G6P pathway. We found that *PIL2* expression is

lower in *gin2*, and that G6P application can reduce expression to similarly low levels in wild

type and *gin2* (**Figure 3D**). This data indicates that HXK1 can promote *PIL2* expression, but

G6P suppresses *PIL2* levels independently of HXK1. On the other hand, *HFR1* expression

appears to be repressed by the HXK1-G6P pathway (**Figure 3D**). Our analysis implies that

PIFs and HXK1 both promote *PIL2* expression, but PIFs and HXK1-G6P have opposing

roles in the regulation of *HFR1*.

HXK1 is functionally important under low light conditions. Our fluence response and

short-day data (**Figure 1 and Figure 1-figure supplement 1**) illustrates that HXK1 loss

curtails seedling growth in light limiting conditions as well as in darkness. Therefore, we were

interested to establish if HXK1 was important for catabolizing photosynthesis-derived

glucose in low light. We found that G6P application enhanced *gin2* hypocotyl length in a

dose-dependent manner with full restoration of *gin2* hypocotyl length occurring at a

concentration of 5mM (0.125% w/v) G6P (**Figure 4A**). Likewise, application of sodium

pyruvate is able to restore *gin2* seedlings to wild type dimensions (**Figure 4-figure**

supplement 1). This confirms that as for dark conditions, HXK1 enzymatic function is

important for supporting seedling growth in low irradiance light. The data also confirm that

the 0.125% concentrations that we have used across experiments lie at the upper limit of the

physiological range.

HXK1 does not mediate glucose-induced hypocotyl elongation. Earlier work has

implicated HXK1 in glucose induced signalling, indeed the *gin2* mutant was identified in a

glucose insensitive germination assay (**Moore et al., 2003**). However, HXK1 does not

appear to control glucose-induced hypocotyl elongation, a response mediated by PIFs, and

BRASSINAZOLE-RESISTANT 1 (BZR1) (**Stewart et al., 2011; Zhang et al., 2016**). In

accord with these studies we have shown that (1%) glucose-mediated hypocotyl extension is

greatly attenuated in the *pifQ* mutant but not in *gin2* which has a near wild type response

(**Supplementary file 1**). In conditions that mimic the original *gin2* mutant screen, we can

show that, as expected, 6% glucose blocks wild type seedling development, while *gin2* is

unaffected. At these high glucose concentrations, like the wild type, *pifQ* fails to thrive

(**Supplementary file 1**). These observations indicate that HXK1 and PIFs operate in distinct

glucose-activated signalling pathways and that HXK1 is not the primary sensor for glucose

activated hypocotyl elongation induced by exogenous glucose.

HXK1 does not suppress photosynthetic gene repression in seedlings. While our data

show HXK1 enzymatic action supports post-germinative seedling growth, we wanted to

establish if we could also find evidence for a glucose-activated HXK1 signalling function.

Previous studies have implicated HXK1 signalling in glucose-mediated transcriptional

suppression of the photosynthetic genes *CAB2* and *CARBONIC ANHYDRASE (CAA)*

(**Moore et al., 2003; Cho et al., 2007**). Consistent with these earlier studies, we also

observe glucose mediated suppression of *CAB2* and *CAA* expression in wild type seedlings,

and the severity of suppression is dose dependent (**Figure 4B**). Likewise, our data illustrate

that HXK1 is required for glucose-suppression of *CAB2* and *CAA* expression, but only at the

higher 1% concentration (and not 0.5% glucose). Application of G6P does not substantially

change *CAB2* or *CAA* expression in either wild type or *gin2*, implying that in seedlings *CAB2*

and *CCA* are not regulated by the HXK1-G6P pathway (**Figure 4B**). Rather, these

photosynthetic genes are negatively regulated by an alternative HXK1-requiring pathway

that is activated by exogenously applied glucose above a concentration threshold.

Given this finding we wanted to establish whether commonly applied glucose concentrations

(0.5-2%) were compatible with or exceeded physiological levels (**Figure 4C**). This was

accomplished by quantifying internal glucose levels in seedlings grown in SD under

increasing light fluence rates: 100, 300 or 600 µmol m^{-2}s^{-1} (referred to henceforth as 100,

300 and 600) and those grown in 100, and supplied with 0.5%, 1% or 2% exogenous

glucose. We aimed to include seedlings grown at higher fluence rates (800µmol m^{-2}s^{-1}), but

they photobleached and perished. In irradiances of 100, internal glucose was recorded at 0.4

mg/g FW and rose by 1.9-fold to 0.75 mg/g FW in seedlings grown at 600 (**Figure 4C**). In

comparison, the exposure to the lowest dose (0.5%) increased internal glucose levels to

2.59 mg/g FW, which amounted to a 6.5-fold increase when compared to control (100) or

3.45-fold when compared to seedlings grown in 600. Application of 1% or 2% greatly

surpasses the normal range of endogenous glucose levels in young seedlings. This data

illustrates that even the lowest (0.5%) dose of glucose exceeds normal physiological levels

by some margin (**Figure 4C**).

We then tested whether *CAB2* and *CAA* expression could be repressed in plants grown in

increased light levels that generate physiological increases in glucose. We found that growth

in 600 compared to 100 led to a small upregulation in *CAB2* in wild type but not in *gin2*

(**Figure 4-figure supplement 2**)*.* The response of *CAA* to increased fluence rates was

relatively unchanged in both genotypes. This indicates that in our conditions, high light

stimulates *CAB2* expression in a HXK1-dependent manner, however this response is very

subtle.

As the above analysis was conducted in SD conditions that restrict the photosynthetic

period, we extended this line of investigation to seedlings grown in long day (LD)

photoperiods at a slightly warmer temperature to maximize photosynthetic potential. In LD

(100), end-of-day seedling glucose levels were nearly two-fold higher than SD (100) but fell

substantially below the 0.5% exogenous glucose dose (**Figure 4C,D**). We also recorded an

incremental fall in glucose levels in 100, 600 and 800. Although we expect flux to glucose to

be dynamic, the internal free-glucose pool is highly regulated (**Mengin et al., 2017**). Indeed,

we found that end-of-day glucose levels decreased slightly with increased fluence rate

(**Figure 4D**). Despite this we recorded a light dose dependent decrease in *CAB2* expression

in both wild type and *gin2* (**Figure 4E**). As for SD, in LD *CAA* levels are relatively unchanged

by light dose in wild type and *gin2*. Earlier studies have reported that *HXK1* transcript

abundance is upregulated by glucose application (**Price et al., 2004; Yang et al., 2013;**

Kelly et al., 2017). In both our SD and LD conditions we do not observe a fluence rate

dependent regulation of *HXK1* transcript levels. Collectively our data illustrate that internal

glucose levels do not appear to rise above the threshold that is required for HXK1-mediated

repression of *CAB2* and *CAA* expression (**Figure 4E and Figure 4-figure supplement 2**).

This indicates that in seedlings the glucose-activated HXK1 suppression of these

photosynthetic genes may not operate.

Discussion

The glucose sensor HXK1 has been shown to function as a glycolytic enzyme, and a

glucose-activated nuclear signalling component proposed to operate during seedling and

adult developmental stages (**Moore et al., 2003; Cho et al., 2007**). Our study delineates an

important glycolytic role for HXK1 in the emerging seedling. The *gin2* and *hxk1-3* mutations

cause stunting in seedlings grown in darkness or low light conditions (**Figure1 A,B and**

Figure 1-figure supplement 1). Quantification of cell number and dimensions reveal that

HXK1 depletion restricts hypocotyl cell division and expansion (**Figure 1D,E**). The *gin2*

mutant accumulated higher levels of glucose than the wild type, indicative of impaired HXK1

catalytic function (**Figure 1F**). Further, application of G6P (or sodium pyruvate), but not

glucose, restored *gin2* seedlings to wild type dimensions (**Figure 1G and Figure 1-figure**

supplement 2). These data point to loss of HXK1 enzymatic function as the cause of *gin2*

and *hxk1-3* seedling stunting. In etiolated seedlings, seed endosperm reserves are used to

fuel hypocotyl elongation. Indeed, like HXK1 mutants, the glyoxylate cycle mutant, *isocitrate*

lyase (icl) and the *phosphoenolpyruvate carboxykinase1 (pck1)* mutant, that blocks carbon

export from the endosperm, also exhibit short hypocotyl phenotypes in the dark (**Eastmond**

et al., 2000; Penfield et al., 2004). Our findings therefore indicate that HXK1 is an important

component of the carbon respiratory machinery that supports post-germinative growth and

this is particularly evident in light-limiting conditions. Lending further support to our findings,

the catalytic activity of an ancestral type B KnHXK1, was recently shown to rescue the

diminutive *gin2* adult phenotype (**Ulfstedt et al., 2018**).

HXK1 is known to regulate gene expression through both glucose-activated signalling and

glycolysis-dependent pathways. Through RNA-seq analysis we established that in darkness

HXK1 loss leads to altered expression of 2189 nuclear genes which represents 12% of the

transcriptome. Just over half (54%) exhibited reduced expression in *gin2 vs* WT, a subset

strongly enriched for energy demanding processes such as ribosome biogenesis,

translation, motor activity and cytoskeleton-dependent transport (**Figure 2A**). In accordance

with the reduction of cell division and expansion in *gin2* compared to wild type, we recorded

a concomitant reduction in the expression of several core *CYC* and *XTH* genes (**Figure**

1D,E and Figure 2-figure supplement 2). In contrast, upregulated genes in *gin2* are

strongly enriched for BCAA enzymes and to a lesser extent hormone response pathways

(**Figure 2B**). Catabolism of the BCAA's, Ile, Leu and Val, is known to support respiration

under carbohydrate-limiting conditions (**Binder, 2010; Hildebrandt et al., 2015**). Recent

work has shown that the expression of BCAA pathway genes are directly activated in

extended darkness by SnRK1 mediated S_1-bZIP signalling (**Pedrotti et al., 2018**; **Baena-**

González et al., 2007). Our data indicate that the enhanced expression of BCAA catabolism

enzymes in *gin2* results from depleted HXK1-glycolytic signalling, as G6P application

restores BCAA gene expression to wild type levels (**Figure 2D**). Interestingly, exogenous

glucose application represses BCAA gene expression in wild type, but also in *gin2*,

indicating this glucose-mediated response does not require HXK1 (**Figure 2D**). Our results

therefore show that BCAA gene expression is regulated through the HXK1-glycolytic

pathway and when HXK1 is fully operational the BCAA carbon starvation-activated pathway

is suppressed.

Remarkably, our RNA-seq analysis revealed that the majority (94%) of the chloroplast

genome and 27% of mitochondrial genes were up-regulated in *gin2* (**Figure 3A**). This

identifies an important novel function for HXK1 in the repression of mitochondrial and

chloroplastic gene expression in darkness. Like the BCAA catabolic pathway genes, we

showed that the chloroplast-encoded polymerase subunits *RPOA, RPOB, RPOC1* and

RPOC2 were also regulated by the HXK1 glycolytic pathway. However, in contrast to *BCAA*

genes, G6P application enhanced polymerase subunit gene expression in wild type and to a

lesser extent in *gin2*, suggesting HXK2 and/or the plastid localized HXK3, may operate

redundantly with HXK1 in the regulation of these genes. Indeed, HXK1, HXK2, and

particularly HXK3 have been implicated in the regulation of sugar-dependent retrograde

signalling, whereby chloroplast gene expression imposes control on nuclear-encoded

photosynthetic gene *LHCB1* (**Zhang et al., 2010**). Our data implies HXK genes operate

redundantly to modulate chloroplastic gene expression and this occurs at least partly

through the glycolytic pathway.

As the light regulated PIF transcription factors control skotomorphogenic growth, therefore

we sought to establish the extent to which HXK1 and PIF mediated gene expression

overlapped by comparing our RNA-seq data with published *pifQ* transcriptome data (**Pfeiffer**

et al. 2014; Zhang et al., 2013). 32% of HXK1 controlled genes were also regulated by

PIFs. BCAA pathway genes were amongst the highly enriched category of PIF-controlled

pathways, but it seems that here PIFs act antagonistically with HXK1 by promoting *BCAA*

gene expression (**Figure 3-figure supplement 1**). From the common gene-set we identified

a smaller group of genes that are directly regulated by PIFs (**Zhang et al., 2013; Pfeiffer et**

al., 2014). In this group, "PIF- and HXK1-induced genes" was the largest individual category,

which was strongly enriched for metabolic pathway genes (*Figure 3-figure supplement 2*).
Direct PIF targets that were repressed by HXK1 contained a higher proportion of
transcription factors and hormone genes. We established that *HFR1*, a gene that is
transcriptionally activated by PIFs is repressed by the HXK1-G6P pathway (*Figure 3D*). This
finding illustrates that carbon status can influence the expression of this key light signalling
component. HFR1 action may therefore be governed by both the external light and internal
metabolic environment.

Several studies have shown sugar application can control hypocotyl elongation, and that
PIFs and BZR1 are important transcriptional modulators of this response (*Stewart et al.,
2011; Zhang et al., 2016*). Our data and that of other labs (*Singh et al., 2017; Zhang et al.,
2016; Simon et al., 2018*) suggest that HXK1 is not the primary effector. Instead the nutrient
sensor Target Of Rapamycin (TOR) kinase is required for sugar induced stabilization of
BZR1 protein and seedling growth (*Zhang et al., 2016*). Further, a recent study has shown
the TOR-translation pathway, via ribosomal S6 kinase (S6K) and its substrate, ribosomal
protein S6 (RPS6), is photoreceptor-activated during the switch to photomorphogenic
seedling growth (*Chen et al., 2018*). Thus, it appears that TOR has a prominent role in
integrating sugar and light sensory signals in the developing seedling.

To investigate more thoroughly the potential for HXK1 glucose signalling in seedlings we
analysed photosynthetic genes *CAB2* and *CAA*, shown previously to be directly targeted by
HXK1 following glucose activation (*Cho et al., 2007*). As for earlier reports, we showed that
exogenously applied glucose inhibited *CAB2* and *CAA* expression in a HXK1-dependent
manner, but there is a threshold glucose concentration (1%) over which HXK1 operates
(*Figure 4B*). By measuring internal glucose levels, we established that the effective 1%
glucose, and even lower 0.5% dose, both greatly exceed physiological levels under a range
of light regimes (*Figure 4C,D*). Indeed, we found that even at high light irradiances HXK1
does not participate in the repression of *CAB2* and *CAA* (*Figure 4E*). It is possible that
glucose-activated HXK1 signalling is more prevalent in adult plants that have increased
capacity for photosynthesis and therefore potential for glucose-activated signalling. Indeed,

in addition to its proposed nuclear signalling role, a novel sucrose-dependent cytosolic

function has been proposed for HXK1 involving the direct interaction with KINγ, a SUCROSE

NON-FERMENTING1 (SNF1)-RELATED KINASE (SNRK1) subunit (***Van Dingenin et al.,***

2018). Our study re-casts HXK1 as a catabolic enzyme that plays a central role in supporting

post-germination growth. An important function of the HXK1-glycolytic pathway is to couple

the metabolism transcriptome to the energy status of the seedling.

Materials and Methods

Plant material, growth conditions, and treatments. The wild-type Arabidopsis thaliana

ecotypes used in this study are Landsberg *erecta* (Ler) and Columbia-0 (Col). The EMS

derived HXK1 nonsense mutant allele *gin2-1* (Ler) (***Moore et al., 2003***) and SALK T-DNA

insertion null allele *hxk1-3* (CS861759) (Col) (***Huang et al., 2015***) were obtained from The

Nottingham Arabidopsis Stock Centre (NASC), UK. For all experiments, seeds were surface-

sterilized as previously described in (***Fankhauser and Casal, 2004***), sown on 0.5X MS

plates ((Duchefa Biochemie, M0221), 0.8% agar, pH 5.7) and stratified in darkness for 2-3

days at 4°C. Specific details on growth conditions are mentioned in the respective figure

legends. Polylux XLR FT8/18W/835 fluorescent tubes (GE, Belgium) were used as a light

source at fluence rates indicated in the respective figure legends.

Seedling hypocotyl length and cotyledon area measurements. Images of seedlings laid

flat on growth media was used to quantify hypocotyl length and cotyledon area using ImageJ

(NIH, Maryland) and Adobe Photoshop CS6 (Adobe, California) respectively. For glucose-6-

phosphate (G6P) (Sigma G7879) and sodium pyruvate (Sigma P2256) treatments, desired

concentrations from filter sterilized stocks were added to cool sterilized media and seeds

directly sown. For determination of etiolated hypocotyl epidermal cell lengths/numbers,

seedlings were cleared overnight as described in (***Weigel and Glazebrook, 2002***), mounted

on slides and visualized using Eclipse E600 Nikon DIC microscope. Individual cell lengths

from each section of the hypocotyl (basal, middle, upper) was measured using ImageJ and

cell number per file was obtained by manual count.

Glucose quantification. Seedlings were harvested in liquid nitrogen, finely ground and ethanol extracted thrice. Glucose was then quantified using an enzymatic assay at 340nm wavelength and normalized to material fresh weight. The procedure has been described in detail previously (*Hendriks et al., 2003*).

Gene expression analysis. For qRT-PCR experiments, seedlings harvested in liquid nitrogen was ground into fine powder. Total RNA was extracted using the RNeasy Plant Mini Kit (Qiagen) with on-column DNase digestion. cDNA synthesis was performed using the qScript cDNA SuperMix (Quanta Biosciences) as described by the manufacturer. The qRT-PCR was set up as a 10µL reaction using SYBR Green (Roche) in a 384-well plate, performed with a Lightcycler 480 system (Roche). Results were analysed using the Light Cycler 480 software. The primers used in this study are listed in Supplementary file 2.

cDNA library preparation and high throughput sequencing. Total RNA was extracted from 4-day old etiolated Ler and *gin2* seedlings (biological duplicates) as described above. Samples were then sent to Edinburgh Genomics (University of Edinburgh, UK) for QC check and sequencing. Briefly, quality check of the samples was performed using Qubit with the broad range RNA kit (Thermo Fisher Scientific) and Tapestation 4200 with the RNA Screentape for eukaryotic RNA analysis (Agilent). Libraries were prepared using the TruSeq Stranded mRNA kit (Illumina), and then validated. Samples were pooled to create 4 multiplexed DNA libraries, which were paired-end sequenced on an Illumina HiSeq 4000 platform (Machine name K00166, Run number 346, flowcell AHT2HKBBXX, lanes 5 and 6). On average 26.6 million 150nt PE reads were obtained for each sample.

Processing of RNA sequencing reads. Sequence reads were aligned against the Arabidopsis thaliana genome (TAIR10) with TopHat v2.1.1 (*Kim et al., 2013*) with default parameters, except in the case of the maximum intron length parameter, which was set at 5000. Count tables for the different feature levels were obtained from bam files using the ASpli package version 1.6.0 (*Mancini et al. 2018*) with custom R scripts and considering the AtRTDv1 transcriptome (*Zhang et al., 2015*). Count tables at the gene level presented a good correlation overall between replicates and samples. Raw sequences (fastq files) and

used in this paper have been deposited in the ArrayExpress (**Kolesnikov et al., 2015**)

database at EMBL-EBI (www.ebi.ac.uk/arrayexpress) under accession number E-MTAB-

7654.

Differential gene expression analysis. Differential gene expression analysis was

conducted for 18,225 genes whose expression was above a minimum threshold level (read

density > 0.05) in at least one experimental condition. Read density (rd) was computed as

the number of reads in each gene divided by its effective width. The term effective width

corresponds to the sum of the length of all the exons of a given gene. Differential gene

expression was estimated using the edgeR package version 3.22.3 (**Robinson et al., 2009**)

and resulting p values were adjusted using a false discovery rate (FDR) criterion. Genes with

FDR values lower than 0.1 and a log2 fold change > 0.6 were considered to be differentially

expressed. Heatmaps were generated using R.

Acknowledgements. A.G was supported by a University of Edinburgh Darwin Trust

Scholarship. This work was funded by the Biotechnology and Biological Sciences Research

Council (BBSRC)-National Science Foundation Grant BB/M025551/1 (awarded to K.J.H)

and 14 ERA-CAPS PHYTOCAL Grant BB/N005147/1 (awarded to K.J.H).

References

Baena-González E, Rolland F, Thevelein JM, Sheen J. 2007. A central integrator of transcription

networks in plant stress and energy signalling. *Nature*. **448**:938-42.

Baud S, Dubreucq B, Miquel M, Rochat C, Lepiniec L. 2008. Storage reserve accumulation in

Arabidopsis: metabolic and developmental control of seed filling. *Arabidopsis Book* **6**:e0113. doi:

10.1199/tab.0113.

Binder S. 2010. Branched-Chain Amino Acid Metabolism in Arabidopsis thaliana. *Arabidopsis Book*.

8:e0137. doi: 10.1199/tab.0137.

Cárdenas ML, Cornish-Bowden A, Ureta T. 1998. Evolution and regulatory role of the hexokinases.

Biochim Biophys Acta. **1401**:242-64.

Chaiwanon J, Wang W, Zhu JY, Oh E, Wang ZY. 2016. Information Integration and Communication in

Plant Growth Regulation. *Cell* **164**:1257-1268. doi: 10.1016/j.cell.2016.01.044.

Chen GH, Liu MJ, Xiong Y, Sheen J, Wu SH. 2018. TOR and RPS6 transmit light signals to enhance

protein translation in deetiolating Arabidopsis seedlings. *Proc Natl Acad Sci U S A*. **115**:12823-12828.

doi: 10.1073/pnas.1809526115.

Cho YH, Yoo SD, Sheen J. 2006. Regulatory functions of nuclear hexokinase1 complex in glucose

signalling. *Cell* **127**:579-89.

Claeyssen E, Rivoal J. 2007. Isozymes of plant hexokinase: occurrence, properties and functions.

Phytochemistry. **68**:709-31.

de Jong F, Thodey K, Lejay LV, Bevan MW. 2014. Glucose elevates NITRATE TRANSPORTER2.1 protein

levels and nitrate transport activity independently of its HEXOKINASE1-mediated stimulation of

NITRATE TRANSPORTER2.1 expression. *Plant Physiol.* **164**:308-20. doi: 10.1104/pp.113.230599.

Eastmond PJ, Germain V, Lange PR, Bryce JH, Smith SM, Graham IA. 2000. Postgerminative growth

and lipid catabolism in oilseeds lacking the glyoxylate cycle. *Proc Natl Acad Sci U S A*. **97**:5669-74.

Fankhauser C, Casal JJ. 2004. Phenotypic characterization of a photomorphogenic mutant. *Plant J.*

39:747-60.

Gommers CMM, Monte E. 2018. Seedling Establishment: A Dimmer Switch-Regulated Process

between Dark and Light Signalling. *Plant Physiol.* **176**:1061-1074. doi: 10.1104/pp.17.01460.

Graham IA. 2008. Seed storage oil mobilization. *Annu Rev Plant Biol.* **59**:115-42. doi:

10.1146/annurev.arplant.59.032607.092938.

Hashimoto T. 2015. Microtubules in plants. *Arabidopsis Book.* **13**:e0179. doi: 10.1199/tab.0179.

Hendriks JH, Kolbe A, Gibon Y, Stitt M, Geigenberger P. 2003. ADP-glucose pyrophosphorylase is

activated by posttranslational redox-modification in response to light and to sugars in leaves of

Arabidopsis and other plant species. *Plant Physiol.* **133**:838-49.

Hildebrandt TM, Nunes Nesi A, Araújo WL, Braun HP. 2015. Amino Acid Catabolism in Plants. *Mol*

Plant. **8**:1563-79. doi: 10.1016/j.molp.2015.09.005.

Hornitschek P, Kohnen MV, Lorrain S, Rougemont J, Ljung K, López-Vidriero I, Franco-Zorrilla JM,

Solano R, Trevisan M, Pradervand S, Xenarios I, Fankhauser C. 2012. Phytochrome interacting factors

4 and 5 control seedling growth in changing light conditions by directly controlling auxin signalling.

Plant J. **71**:699-711. doi: 10.1111/j.1365-313X.2012.05033.x.

Hsu YF, Chen YC, Hsiao YC, Wang BJ, Lin SY, Cheng WH, Jauh GY, Harada JJ, Wang CS. 2014. AtRH57,

a DEAD-box RNA helicase, is involved in feedback inhibition of glucose-mediated abscisic acid

accumulation during seedling development and additively affects pre-ribosomal RNA processing with

high glucose. *Plant J.* **77**:119-35. doi: 10.1111/tpj.12371.

Huang JP, Tunc-Ozdemir M, Chang Y, Jones AM. 2015. Cooperative control between AtRGS1 and

AtHXK1 in a WD40-repeat protein pathway in Arabidopsis thaliana. *Front Plant Sci.* **6**:851. doi:

10.3389/fpls.2015.00851.

Kelly G, Sade N, Doron-Faigenboim A, Lerner S, Shatil-Cohen A, Yeselson Y, Egbaria A, Kottapalli J,

Schaffer AA, Moshelion M, Granot D. 2017. Sugar and hexokinase suppress expression of PIP

aquaporins and reduce leaf hydraulics that preserves leaf water potential. *Plant J.* **91**:325-339. doi:

10.1111/tpj.13568.

Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. 2013. TopHat2: accurate alignment of

transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol.* **14**:R36. doi:

10.1186/gb-2013-14-4-r36.

Kolesnikov N, Hastings E, Keays M, Melnichuk O, Tang YA, Williams E, Dylag M, Kurbatova N,

Brandizi M, Burdett T, Megy K, Pilicheva E, Rustici G, Tikhonov A, Parkinson H, Petryszak R, Sarkans

U, Brazma A. 2015. ArrayExpress update--simplifying data submissions. *Nucleic Acids Res.* **43**:D1113-

6. doi:10.1093/nar/gku1057

Kunz S, Gardeström P, Pesquet E, Kleczkowski LA. 2015. Hexokinase 1 is required for glucose-

induced repression of bZIP63, At5g22920, and BT2 in Arabidopsis. *Front Plant Sci.* **6**:525. doi:

10.3389/fpls.2015.00525

Leivar P, Monte E, Oka Y, Liu T, Carle C, Castillon A, Huq E, Quail PH. 2008. Multiple phytochrome-

interacting bHLH transcription factors repress premature seedling photomorphogenesis in darkness.

Curr Biol. **18**:1815-23. doi: 10.1016/j.cub.2008.

Leivar P, Monte E. 2014. PIFs: systems integrators in plant development. *Plant Cell.* **26**:56-78. doi:

10.1105/tpc.113.120857.

Leivar P, Tepperman JM, Cohn MM, Monte E, Al-Sady B, Erickson E, Quail PH. 2012. Dynamic

antagonism between phytochromes and PIF family basic helix-loop-helix factors induces selective

reciprocal responses to light and shade in a rapidly responsive transcriptional network in

Arabidopsis. *Plant Cell.* **24**:1398-419. doi: 10.1105/tpc.112.095711.

Mancini E, Iserte J, Yanovsky M, Chernomoretz A. 2018. ASpli: Analysis of alternative splicing using

RNA-Seq. R package version 1.6.0.

Mengin V, Pyl ET, Alexandre Moraes T, Sulpice R, Krohn N, Encke B, Stitt M. 2017. Photosynthate

partitioning to starch in Arabidopsis thaliana is insensitive to light intensity but sensitive to

photoperiod due to a restriction on growth in the light in short photoperiods. *Plant Cell Environ.*

40:2608-2627. doi:10.1111/pce.13000.

Moore B, Zhou L, Rolland F, Hall Q, Cheng WH, Liu YX, Hwang I, Jones T, Sheen J. 2003. Role of the

Arabidopsis glucose sensor HXK1 in nutrient, light, and hormonal signalling. *Science.* **300**:332-6.

Pedrotti L, Weiste C, Nägele T, Wolf E, Lorenzin F, Dietrich K, Mair A, Weckwerth W, Teige M, Baena-

González E, Dröge-Laser W. 2018. Snf1-RELATED KINASE1-Controlled C/S1-bZIP Signalling Activates

Alternative Mitochondrial Metabolic Pathways to Ensure Plant Survival in Extended Darkness. *Plant*

Cell **30**:495-509. doi: 10.1105/tpc.17.00414.

Penfield S, Pinfield-Wells HM, Graham IA. 2006. Storage reserve mobilisation and seedling

establishment in Arabidopsis. *Arabidopsis Book* **4**:e0100. doi:10.1199/tab.0100.

Penfield S, Rylott EL, Gilday AD, Graham S, Larson TR, Graham IA. 2004. Reserve mobilization in the

Arabidopsis endosperm fuels hypocotyl elongation in the dark, is independent of abscisic acid, and

requires PHOSPHOENOLPYRUVATE CARBOXYKINASE1. *Plant Cell*. **16**:2705-18.

Pfeiffer A, Shi H, Tepperman JM, Zhang Y, Quail PH. 2014. Combinatorial complexity in a

transcriptionally centered signalling hub in Arabidopsis. *Mol Plant*. **7**:1598-1618. doi:

10.1093/mp/ssu087.

Price J, Laxmi A, St Martin SK, Jang JC. 2004. Global transcription profiling reveals multiple sugar

signal transduction mechanisms in Arabidopsis. *Plant Cell*. **16**:2128-50.

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential

expression analysis of digital gene expression data. *Bioinformatics*. **26**:139-40. doi:

10.1093/bioinformatics/btp616.

Simon NML, Kusakina J, Fernández-López Á, Chembath A, Belbin FE, Dodd AN. 2018. The Energy-

Signalling Hub SnRK1 Is Important for Sucrose-Induced Hypocotyl Elongation. *Plant Physiol*.

176:1299-1310. doi: 10.1104/pp.17.01395.

Singh M, Gupta A, Singh D, Khurana JP, Laxmi A. 2017. Arabidopsis RSS1 Mediates Cross-Talk

Between Glucose and Light Signalling During Hypocotyl Elongation Growth. *Sci Rep*. **7**:16101. doi:

10.1038/s41598-017-16239-y.

Stewart JL, Maloof JN, Nemhauser JL. 2011. PIF genes mediate the effect of sucrose on seedling

growth dynamics. *PLoS One*. **6**:e19894. doi:10.1371/journal.pone.0019894.

Theodoulou FL, Eastmond PJ. 2012. Seed storage oil catabolism: a story of give and take. *Curr Opin*

Plant Biol. **15**:322-8. doi: 10.1016/j.pbi.2012.03.017.

Ulfstedt M, Hu GZ, Eklund DM, Ronne H. 2018. The Ability of a Charophyte Alga Hexokinase to

Restore Glucose Signalling and Glucose Repression of Gene Expression in a Glucose-Insensitive

Arabidopsis Hexokinase Mutant Depends on Its Catalytic Activity. *Front Plant Sci*. **9**:1887. doi:

10.3389/fpls.2018.01887.

Van Dingenen J, Vermeersch M, De Milde L, Hulsmans S, De Winne N, Van Leene J, Gonzalez N,

Dhondt S, De Jaeger G, Rolland F, Inzé D. 2019. The role of HEXOKINASE1 in Arabidopsis leaf growth.

Plant Mol Biol. **99**:79-93. doi: 10.1007/s11103-018-0803-0

Weigel D & Glazebrook J. 2002. Arabidopsis: A Laboratory Manual (Cold Spring Harbor Laboratory

Press, Cold Spring Harbor, NY).

Yanagisawa S, Yoo SD, Sheen J. 2003. Differential regulation of EIN3 stability by glucose and

ethylene signalling in plants. *Nature* **425**:521-5.

Yang L, Xu M, Koo Y, He J, Poethig RS. 2013. Sugar promotes vegetative phase change in Arabidopsis

thaliana by repressing the expression of MIR156A and MIR156C. *Elife.* **26**;2:e00260. doi:

10.7554/eLife.00260

Yu S, Cao L, Zhou CM, Zhang TQ, Lian H, Sun Y, Wu J, Huang J, Wang G, Wang JW. 2013. Sugar is an

endogenous cue for juvenile-to-adult phase transition in plants. *Elife.* **26**;2:e00269. doi:

10.7554/eLife.00269

Zhang R, Calixto CP, Tzioutziou NA, James AB, Simpson CG, Guo W, Marquez Y, Kalyna M, Patro R,

Eyras E, Barta A, Nimmo HG, Brown JW. 2015. AtRTD - a comprehensive reference transcript dataset

resource for accurate quantification of transcript-specific expression in Arabidopsis thaliana. *New*

Phytol. **208**:96-101. doi: 10.1111/nph.13545.

Zhang Y, Mayba O, Pfeiffer A, Shi H, Tepperman JM, Speed TP, Quail PH. 2013. A quartet of PIF bHLH

factors provides a transcriptionally centered signalling hub that regulates seedling morphogenesis

through differential expression-patterning of shared target genes in Arabidopsis. *PLoS Genet.*

9:e1003244. doi: 10.1371/journal.pgen.1003244.

Zhang Z, Zhu JY, Roh J, Marchive C, Kim SK, Meyer C, Sun Y, Wang W, Wang ZY. 2016. TOR Signalling

Promotes Accumulation of BZR1 to Balance Growth with Carbon Availability in Arabidopsis. *Curr Biol.*

26:1854-60. doi:10.1016/j.cub.2016.05.005.

Zhang ZW, Yuan S, Xu F, Yang H, Zhang NH, Cheng J, Lin HH. 2010. The plastid hexokinase pHXK: a

node of convergence for sugar and plastid signals in Arabidopsis. *FEBS Lett.* **584**:3573-9. doi:

10.1016/j.febslet.2010.07.024.

Figure 1. HXK1 operates in the dark and low light during seedling development. (**A)** Fluence

rate response curves for hypocotyl elongation in 7 day old WT, *gin2* (Ler) and *hxk1-3* (Col)

seedlings grown in continuous light at 18°C. Cotyledon area (**B**) and images of seedlings (**C**)

grown in low or high light (3 or 130 µmol m^{-2} s^{-1}). (**D**) Epidermal cell length at the basal,

middle and upper regions of hypocotyl in etiolated WT (Ler) and *gin2* seedlings at 4 days.

(**E**) Hypocotyl epidermal cell number in etiolated WT (Ler) and *gin2* seedlings at 4 and 14

days. Red circles in the dot plots indicate mean value. (**F**) Glucose content (mg/g FW) in 4

day old etiolated WT (Ler) and *gin2* seedlings. (**G**) Hypocotyl length of etiolated 7 day old

WT (Ler) and *gin2* seedlings grown at 18°C without or with 0.5% (w/v) glucose, 0.125% and

0.25% (w/v) G6P. Error bars indicate ±SEM; **P ≤ 0.01 (Student's t-test) of mutants relative

to respective WT for each condition.

Figure 2. HXK1 regulates key energy regulated pathways**.** Hierarchical clustering and Gene
Ontology analysis of (**A**) upregulated and (**B**) downregulated genes in etiolated WT (Ler) and
gin2 biological duplicates at 4 days. Scale bar denotes log counts per million, low (blue) to
high (yellow). Bar plots of statistically significant (p≤0.01, Fisher Exact Test) GO terms
derived from VirtualPlant database; orange line denotes an observed:expected frequency
ratio of 1. (**C**) Branched Chain Amino Acid (BCAA) degradation pathway schematic with
enzymes upregulated in gin2 highlighted in red. (**D**) Transcript abundance (qPCR) of BCAA
pathway genes in 4 day old etiolated WT (Ler) and *gin2* seedlings grown at 18°C with or
without 0.5% w/v glucose or 0.125% w/v G6P. Error bars indicate ±SE, *P ≤ 0.05, **P≤ 0.01
(Student's t-test) of *gin2* relative to WT for each treatment.

Figure 3. HXK1 represses plastome genes and operates mainly independently of PIFs during skotomorphogenesis. (**A**) Proportion of upregulated chloroplast and mitochondrial genes (from RNAseq) in etiolated *gin2* seedlings relative to WT (Ler) at 4 days. (**B**) Transcript abundance (qPCR) of plastid encoded polymerases (RPOs) in 4 day old etiolated WT (Ler) and *gin2* seedlings grown at 18ºC without or with 0.125% w/v G6P. Fold change induced by treatment is shown. (**C**) Comparison of *gin2*/WT and *pifQ*/WT transcriptomes from etiolated seedlings (**Pfeiffer et al., 2014**). (**D**) *PIL2* and *HFR1* transcript abundance (qPCR) in 4 day old etiolated WT (Ler) and *gin2* seedlings grown at 18ºC without or with 0.125% w/v G6P. Error bars indicate ±SE, **P≤ 0.01 (Student's t-test) of *gin2* relative to WT (Ler) for each treatment.

Figure 4. HXK1 does not control *CAB2* or *CAA* expression under physiological conditions.

(**A**) Hypocotyl length of 7 day old WT (Ler) and *gin2* seedlings grown in low light (3μmol m^{-2}

s^{-1}) at 18ºC without or with increasing concentrations of G6P. (**B**) Transcript abundance of

CAB2 and *CAA* in WT (Ler) and *gin2* seedlings grown in 100μmol m^{-2} s^{-1} light, without or

with 0.5%, 1% w/v glucose, or 0.125% w/v G6P. (**C**) Glucose (mg/g FW) levels in WT (Ler)

seedlings grown at increasing light intensities (100, 300 and 600μmol m^{-2} s^{-1}) or at 100μmol

m^{-2} s^{-1} plus increasing concentrations of exogenous glucose (0%, 0.5%, 1% and 2% (w/v)).

For both (**B**) and (**C**) seedlings were grown in SD photoperiods at 18°C and sampled on day

4 at ZT4. (**D**) WT (Ler) Glucose (mg/g FW) levels at increasing light intensities (100, 600 and

800μmol m^{-2} s^{-1}). (**E**) *HXK1*, *CAA* and *CAB2* transcript abundance (qPCR) in Ler and *gin2*

seedlings grown at increasing light intensities. For both (**D**) and (**E**) seedlings were grown in

LD photoperiods at 22°C and sampled after 7 days at EoD (ZT16). Error bars indicate ±SE,

**P≤ 0.01 (Student's t-test) of *gin2* relative to WT (Ler) for each treatment.

 Supplementary Information

Figure 1–figure supplement 1. HXK1 operates in the dark and suboptimal light during seedling

establishment. Hypocotyl length of 7-day old WT, *gin2* (Ler) and *hxk1-3* (Col) seedlings grown in

Short Day (SD) and long day (LD) photoperiods at 100µmol m^{-2}s^{-1} or in LD at 5µmol m^{-2}s^{-1} light at

18°C. Horizontal bars, boxes, and whiskers show medians, interquartile ranges (IQR), and data

ranges, respectively. Different letters denote statistical differences (P > 0.05) among samples as

assessed by one-factorial ANOVA and Tukey HSD.

Figure 1-figure supplement 2. Hypocotyl length of 7 day old etiolated WT (Ler) and *gin2* seedlings

grown in the absence and presence of 0.125% G6P, 0.125% and 0.25% (w/v) sodium pyruvate at

18°C. Error bars indicate ±SE, **P≤ 0.01 (Student's *t*-test) of *gin2* relative to WT (Ler) for each

treatment.

Figure 2-figure supplement 1. Percentage of genes down and up-regulated (± 1.5 FC) in 4 day old

etiolated *gin2* vs WT (Ler).

Figure 2-figure supplement 2. Fold decrease of (**A**) core cell cycle/cell division genes, (**B**) cell wall modifying XYLOGLUCOSYL TRANSFERASE (XTH) genes in *gin2* relative to WT (Ler) derived from the RNAseq experiment. Error bars represent the propagated error value.

Figure 3-figure supplement 1. HXK1 and PIFs largely operate independently during

skotomorphogenesis. **(A)** Statistically significant (p≤0.01, Fisher Exact Test) GO terms derived from

VirtualPlant database for 748 overlapping genes mis-regulated in the *gin2* and *pifQ* mutant are plotted.

Orange line indicate representation factor of 1 (observed/expected frequency ratio). **(B)** Normalised

counts for BCAA catabolic genes in WT and *pifQ* (Raw RNAseq data derived from ***Zhang et al.,***

2013).

Figure 3-figure supplement 2. HXK1 and PIFs largely operate independently during

skotomorphogenesis. Comparison between direct PIF targets (data derived from *Pfeiffer et al., 2014*)

with genes **(A)** upregulated and **(B)** downregulated in *gin2*. Genes regulated in a similar or opposite

manner are depicted below venn diagrams.

Figure 4-figure supplement 1. Restoring energy recues the *gin2* mutant seedling phenotype in low

light. **(A)** Hypocotyl length and **(B)** cotyledon area of 7 day old WT (Ler) and *gin2* seedlings grown in

the absence and presence of 0.125% w/v G6P and 0.125% and 0.25% (w/v) sodium pyruvate under

continuous low light (3μmol m^{-2} s^{-1}) conditions at 18°C. Error bars indicate ±SE, **P≤ 0.01 (Student's

t-test) of *gin2* relative to WT (Ler) for each treatmen

Figure 4-figure supplement 2. Photosynthetic feedback inhibition mediated by HXK1 signalling

does not occur in seedlings. qPCR measuring *CAB2*, *CAA and HXK1 transcript abundance* at ZT4 in

4 day old WT (Ler) and *gin2* seedlings grown under increasing light intensities (100, 300 and

600µmol m^{-2} s^{-1}) in SD photoperiods at18°C. Different letters denote statistical differences (P > 0.05)

among samples as assessed by one-factorial ANOVA and Tukey HSD.

Supplement figure 1. HXK1 and PIFs operate in independent sugar signalling pathways. (*A*)

Hypocotyl length of 7 day old WT, *pifQ* (Col) and *gin2* (Ler) seedlings grown in short days (100μmol

m^{-2} s^{-1}) in the absence (black bar) and presence of 1% (w/v) glucose at 18°C. (*B*) Images of 7 day old

seedlings grown under continuous light (130μmol m^{-2} s^{-1}) in the presence of 6% (w/v) glucose or

mannitol at 18°C. Error bars indicate ±SE, **P≤ 0.01 (Student's *t*-test) of glucose treated relative to

untreated

Supplement figure 2. Primers used in this work.

Primer name	Sequence (5'-3')
AG_2250_RPOA_F	GCGATGCGAAGAGCTTTACT
AG_2251_RPOA_R	CCAGGACCTTGGACACAAA
AG_2252_RPOB_F	GATGTGAGGTGGGTTCAGAA
AG_2253_RPOB_R	GGTCTCCCGTCTTGCAAATA
AG_2254_RPOC1_F	TTCTTCCTCCCGAGTTGAGA
AG_2255_RPOC1_R	CCACGGCTTCTTGTACCAAT
AG_2256_RPOC2_F	CGCGTCGACTTGTTGAAGTA
AG_2257_RPOC2_R	CGTCTGCTAAGACACGACCA
AG_2236_BCAT2_F	GGGATAATCTCGGGTTTGGT
AG_2237_BCAT2_R	CTTCATCCGGATAGCGTTGT
AG_2232_THDP_F	GACGAAGACGGACGAATCAT

AG_2233_THDP_R	TGCTGAAGCGATGTTAATGG
AG_2234_DIN2_F	CGGTCGTCGGAGAGAGTAAC
AG_2235_DIN2_R	GCCTTGCAAAACACCAAAAT
AG_2238_MCCA_F	CCCGTCTACAGGTCGAACAT
AG_2239_MCCA_R	ACCCGAACTGATGGTGAGAC
AG_2240_IVD_F_	ACTCTGTTGCGAGGGACTGT
AG_2241_IVD_R	CTTAGAAGGCGTCCTGTTGC
AG_2242_MCCB_F	CTTTGCCTTCAGGTGGGATA
AG_2243_MCCB_R	ACCGAGCAGCAATCTCTTGT
AG_1267_CAB2_F	CCCTGGAGACTACGGATG
AG_1267_CAB2_R	TCCAAACTTGACTCCGTTCC
AG_1428_CAA_F	TGAATACGCTGTCTTGCACC
AG_1429_CAA_R	TGTGATGGTGGTGGTAGCGA
DY_1166_HXK1_F	GGTTTCACTTTCTCGTTTCCTG
DY_1167_HXK1_R	CTTGTCCAACTGCTTCTTCG

690   *Figure 2—source data 1*

691   List of genes altered in 4-day-old etiolated *gin2* mutants

692   HXK1 regulated genes were defined by 1.5-fold difference between *gin2* and WT (Ler) with

693   p<0.01.