1   **Urinary glucocorticoids in harbour seal (*Phoca vitulina*) pups during rehabilitation**

2   Susan C. Wilson[1], Stella Villanueva[2], Lilia Dmitrieva[3], William Hayes[4] and Wesley Smyth[5]

3   [1]Tara Seal Research, Killyleagh, Co. Down, N. Ireland, UK

     [2] Vetagro-Sup (School of Veterinary Science), Lyon, France

     [3]School of Biology, University of Leeds, UK

[4] University of Lincoln, Riseholme Park, Lincolnshire, England, UK

[5] Chemical and Immunodiagnostic Sciences Branch, Veterinary Sciences Division, Agri-Food &

Biosciences Institute, Belfast BT4 3SD, N. Ireland

Short title: harbour seal pup urinary glucocorticoids in rehabilitation

Corresponding author: Susan C. Wilson, Tara Seal Research, 14 Bridge Street, Killyleagh, Co. Down

BT30 9QN, N. Ireland, UK. Tel. +44-28-44-821107, +44-7742-451852.

email: suewilson@sealresearch.org

12
13
14

Abstract

The aim of this empirical study was to investigate whether aspects of the care and conditions of

harbour seal (*Phoca vitulina*) pups in rehabilitation centres may be reflected in urinary concentrations

of glucocorticoid hormones (GCs). Urine samples were collected non-invasively from pups at five

different rehabilitation centres, three on the Irish Sea and two in the southern North Sea.

Concentrations of urinary cortisol (CL) were analysed by ELISA from one centre and subsequently

concentrations of CL, cortisone (CN), prednisolone (PL) and prednisone (PN) were analysed by mass

spectrometry. Urinary concentrations of PN and PL were found to be of similar magnitude to CL. CL,

CN and PL concentrations were reciprocally correlated, but PN was not correlated with CL and only

weakly correlated with the other three GCs. The strongest associations between GC concentrations and

pup conditions and procedures were negative correlations of PN with growth rate of individual pups

and body weight. Differences in GC concentrations, particularly CL, between pups in different social

groupings (pups alone, paired or in groups of more than two) were also detected. On the basis of our

results, we suggest that harbour seal pups should be fed a diet – from first entry to a rescue centre –

which maximises weight gain; this should not only help to prevent chronically elevated GC levels

during the rehabilitation period, but would also minimise time spent in captivity.

Key words

Seal pups, glucocorticoids, rehabilitation, nutrition, welfare, captive environment

INTRODUCTION

New-born harbour seal pups found stranded on the shoreline without the mother (termed 'orphan' pups) are typically taken to 'seal sanctuary' centres for mother-substitute care (usually termed 'rehabilitation') until they are considered well-enough grown to be released back into the sea, usually after a period varying between about six weeks to 4 months.

Harbour seal pups are highly precocial, and free-living pups spend around 60% of their time in the water from birth [1, 2]. Average birth weight is about 11 kg and pups nurse from their mother usually for 3–4 weeks, or up to 6 weeks in some populations [2–4]. The fat content of the milk increases from about 41% at birth to 50% by day 7 until weaning [5], with pups typically achieving an average net weight gain of 0.4 to 0.6 kg/day [2–4]. During the nursing period, free-living pups are constantly in close proximity to their mother [6–8] until about 10 days of age, when mothers sometimes leave their pups onshore in the haul-out group for a few hours while they make foraging trips offshore [9, 10]. Reproducing – for orphan pups in rehabilitation ('rehab') – this concentrated nutrition and key features of the amphibious and social environment of free-living pups is obviously challenging.

Harbour seal pups in rehab will usually not suckle from a bottle. Some centres feed pups by force-feeding with whole fish or tube-feeding fish 'soup', with little or no growth or weight gain by pups until they are past natural weaning age and feeding freely on whole fish [11, 12]. Other centres enhance pup weight gain by tube-feeding pups enriched formulae using dedicated high fat, multi-milk based formulations; however, average weight gain nevertheless rarely achieves 0.3 kg/day [12–14].

Rehab facilities in the British Isles and N. America are usually designed to maintain orphan harbour seal pups in isolation and in dry pens for some weeks after admission [12, 13, 15], a principal reason for this being to prevent potential transmission of infectious agents [16]. Less often, centres house these pups with access to a pool, either in pairs [14] or in small groups [17], where they are able to experience species-typical sensory stimulation and muscle movement during swimming and social interaction.

The glucocorticoid (GC) hormone cortisol is widely considered to be a physiological marker of an animal's 'stress' status [18, 19], with high GC levels usually considered to be an indicator of poor

animal welfare in captivity [20], although with the caveat that studies have shown inconsistency

between high cortisol levels and other welfare measures [21, 22]. A study of harbour seal pups in rehab

found higher plasma cortisol levels in pups being tube-fed than in pups (at a later rehab stage) feeding

on fish, suggesting that the tube-feeding procedures might be stressful [23]. The same study found that

plasma cortisol levels in the rehab pups were higher than in free-ranging pups and suggested that pups

from rehab are not physiologically equivalent to free-ranging weaned pups [23].

Because restraining animals to obtain blood samples for cortisol (CL) assays is stressful *per se*, urine

sampling has been suggested as being a less invasive procedure [19, 20] and has been trialled

successfully in free-living Weddell seals [24]. GC measurement in urine has the advantage that protein-

bound GCs are excluded from the kidney filtrate and so urinary GCs are a reflection of the free

hormone in blood over the time that the urine collects between voiding [20, 25].

The present study is a preliminary report on urinary GC concentrations in harbour seal pups in five

rehab centres with different feeding and care systems. In addition to cortisol, we analysed urine

concentrations of the three other closely related GCs, cortisone (CN), prednisolone (PL) and

prednisone (PN). To achieve this, we used mass spectrometry for multiple steroids, as recommended

by Koren and colleagues [26].

ANIMALS AND METHODS

Pup background

This study involved harbour seal pups which had been admitted to five participating rehabilitation

centres, three on the Irish Sea coast of Ireland and two in the southern North Sea (Lincolnshire, UK and

northern Germany). All the pups had been found stranded on the shoreline during the pup birthing

season. On admittance to the centres, the pups were found to be mostly at or below average birth

weight of approximately 11kg for harbour seals [2–4], with only two pups (in centre IV) being slightly

above this weight, at 12–13kg (Table 1). The pups in this study were all therefore believed to have

been permanently separated from their mothers in the neonatal or early post-natal period, which we

consider to be an operational definition of 'orphans'. None of the pups were of sufficient size to have

been healthy pups aged 10 days or more, mistakenly taken into captivity when their mothers had left

them onshore temporarily while foraging offshore [10]. The pups in this study were all therefore

believed to be genuine 'orphans'.

 Table 1. Summary of individual pup data and results of urinary glucocorticoid analysis (GC results in ng/mg creatinine)

Centre	Pup name	Sex	Date of entry to rehab	Entry weight (kg)	Feeding method	Growth rate (kg/d)	social group	n (samples)	CORTISOL range	CORTISOL median or single	CORTISONE range	CORTISONE median or single	PREDNISOLONE range	PREDNISOLONE median or single	PREDNISONE range	PREDNISONE median or single
Rehab day 0–1																
I	Orlando	M	21/07/12	10.1	Fish FF-HF	NA	1	1	221.23		NA	NA	NA	NA	NA	NA
I	"	"	"	"	"	"	"	"	211†							
I	Phocina	F	25/07/12	8.2	Fish FF-HF	NA	1	1	287†		NA	NA	NA	NA	NA	NA
II	Ula	F	05/07/14	7.1	Tube	NA	1	1		8062		4853.00		46.00		249.00
III	Rowan	M	24/06/14	7.8	Tube	NA	1	2	5-16	10.50	405-418	411.80	38-57	47.49	14-26	19.56
III	Beech	M	06/07/14	10.3	Tube	NA	1	1		14.28		407.00		30.00		13.62
III	Polly*	F	18/07/15	6.5	Tube	NA	1	1		486.6		455.03		114.93		236.59
IV	Urmel	F	05/07/15	10.6	Tube	NA	1	2	68-124	96.43	61-240	150.64	88-143	115.56	55-132	98.95
IV	Julio	M	08/07/15	11.5	Tube	NA	1	3	110-447	145.14	279-788	483.99	30-79	57.42	49-124	52.65
IV	Schultze	F	08/07/15	6.9	Tube	NA	1	1		4028.9		459.75		206.4		25.42
Rehab day 2 onwards																
I	Barry	M	20/06/12	8.94	Fish FF	0.08	1	7	33.6-106.8	62.83	NA	NA	NA	NA	NA	NA
I	"	"	"	"	"	"	"	9	61–134†	98†	NA	NA	NA	NA	NA	NA
I	Pascal	M	23/06/12	9.07	Fish FF	0.03	1	3	21.8-123.3	110.35	NA	NA	NA	NA	NA	NA
I	"	"	"	"	"	"	"	5	64-1164†	84†	NA	NA	NA	NA	NA	NA
II	Maxi	M	19/07/13	11.1	Tube	0.28	2	7	9.4-36.4	15.78	NA	NA	NA	NA	NA	NA
II	Mini	F	19/07/13	9	Tube	0.29	2	1		14.8	NA	NA	NA	NA	NA	NA
II	Ula	F	05/07/14	7.1	Tube	0.28	2	9	0.0-49.0	20.78	123.7-871.7	293.00	6.3-47.8	21.74	0.0-40.7	15.45
II	Earendil	M	07/07/14	7.4	Tube	0.27	2	3	6.6-21.3	20.16	231.2-693.4	444.82	9.2-14.8	11.76	8.8-11.2	9.09
II	Coral	F	03/07/16	8.5	Tube	0.29	2	6	9.2-206.7	39.14	69.5-571.2	153.34	5.4-27.0	20.35	10.4-45.2	32.46
II	Pearl	F	05/07/16	8.3	Tube	0.3	2	6	2.7-365.3	14.83	20.9-403.0	59.05	6.7-34.4	10.43	8.2-53.3	11.76
III	Elm	M	20/06/14	8	Tube	0.22	2	2	20.7-51.6	23.86	320.4-850.1	585.20	20.7-51.6	36.18	0.0-16.7	8.35
III	Birch	M	13/06/14	7.5	Tube	0.26	1	1		0.00		291.03		63.84		0.00
III	Fern	F	02/07/14	7.75	Tube	0.18	1	1		20.51		445.16		23.09		19.31
III	Finn	M	03/07/15	8.5	Tube	0.25	1	2	1.0-1.2	1.09	101.9-187.1	144.50	5.3-9.7	7.51	24.0-51.5	37.73
III	Isla	F	16/07/15	8	Tube	0.1	1	1		13.85		67.54		10.61		90.21
III	Klamath	M	30/06/15	7	Tube	0.23	1	1		11.73		327.63		39.22		91.30
III	Rowan	M	24/06/14	7.75	Tube	0.25	1	1		1.83		384.96		35.92		7.32
IV	Bärchen	M	29/06/15	12.2	Fish HF	0.16	3	1	10.67	10.67		259.85		22.97		27.31
IV	Beate	F	03/07/15	10.6	Tube	0.26	2	2	0.0-175.1	87.54	782.2-984.2	883.22	21.9-23.8	22.85	0.0-21.5	10.75
IV	Chigoma	M	03/07/15	10.3	Tube	0.01	2/3	2	66.7-178.6	122.67	371.8-827.1	599.47	40.3-48.0	44.18	36.7-57.3	47.00
IV	Daggi	F	25/06/15	10.7	Fish HF	0.17	3	1		68.43		357.34		45.78		26.83

	Name	Sex	Date		Method				Range 1	Value 1	Range 2	Value 2	Range 3	Value 3	Range 4	Value 4
IV	Eddi	M	01/07/15	13.1	Tube	0.17	2/3	4	44.4-79.9	62.26	281.6-741.0	592.26	25.2-60.9	28.21	31.6-66.7	40.23
IV	Henry	M	03/07/15	10.1	Tube	0.18	3	3	89.2-243.6	103.84	588.9-896.6	668.15	34.8-61.7	42.89	37.6-51.2	39.38
IV	Jens	M	26/06/15	8.8	Fish HF	0.23	3	3	41.5-60.3	44.10	383.3-515.5	464.13	30.8-36.3	34.35	36.8-42.2	39.40
IV	Julio	M	08/07/15	11.5	Tube	0.03	1	1		82.6		442.35		66.01		76.38
IV	Kawamba	F	06/07/15	10.4	Tube	0.13	1	1		84.19		417.66		37.67		55.18
IV	Lisi	F	01/07/15	11	Tube	0.14	2/3	4	31.0-704.8	61.11	320.1-2233.2	633.94	37.94-57.97	42.92	35.6-58.2	48.52
IV	Schultze	F	08/07/15	6.9	Tube	0.1	1/3	2	298.4-2030.9	1164.60	2109.4-2787.4	2448.40	72.6-101.3	86.96	23.1-74.0	48.57
IV	Urmel	F	05/07/15	10.6	Tube	0.11	1/2	4	90.3-236.9	178.24	440.5-1295.4	862.47	38.4-95.8	62.50	46.7-70.6	56.62
IV	Yukari	F	01/07/15	10	Tube	0.22	3	2	79.6-104.7	92.13	304.0-395.8	349.91	23.0-31.4	27.19	47.5-53.4	50.43
V	Aquaman	M	07/07/15	10.2	Tube	0.1	1	3	5.56-90.48	36.87	291.1-904.8	407.08	34.9-43.3	38.09	20.3-40.8	20.65
V	Blackbird	M	04/07/15	7.4	Tube	0.05	2	3	0.0-37.2	8.38	223.7-958.6	363.30	23.1-80.5	74.87	0.0-45.5	12.52
V	Hawkeye	M	09/07/15	8.35	Tube	0.11	1	4	0.0-138.3	47.00	162.1-941.4	586.78	0.0-92.7	70.00	31.9-139.7	59.64
V	Medusa	F	27/06/15	7	Tube	-0.1	2	2	0.0-11.5	5.76	236.7-521.1	378.87	42.0-55.8	48.90	0.0-25.3	12.65
V	Mystique	F	21/06/15	9.6	Fish HF	0.06	2	1		61.25		599.28		33.871		32.70
V	Wolverine	M	06/06/15	7.8	Fish HF	0.19	2	2	27.7-28.1	27.89	337.6-374.3	355.91	32.3-49.7	41.02	28.8-45.5	37.12

93   † results from ELISA, FF – force-fed, HF – hand-fed, Tube – stomach gavage

Pup rehabilitation centres

Five seal pup rehabilitation centres participated in this study. Two were on the southern North Sea coast: centre I in Lincolnshire, UK and centre IV in northern Germany. The other three centres (II, III and V) were located on the Irish Sea coast of Ireland. There were therefore differences between centres in the population of the pups' birth in addition to differences in pup care procedures, including the rehab captive environment (S1 Photos), food constituents and feeding methods during the sampling period (Table 1). In centres I and III, the pups were kept alone in dry pens, in centres II and IV the pups were usually in pairs or groups with free access to water, and in centre V the pups were either alone or paired and in dry pens (S1 Photos), see also Table 1 for individual pup details).

In this study we considered *pup growth rate* (kg/day) recorded by the centre personnel during our sampling period. We also recorded three categories of each pup's *social group* environment at the time of sampling, i.e. (1) alone, (2) paired or (3) more than two pups, and whether the pups had free access to *water* (omitted from Table 1; only pups at centres II and IV had free access to water during the sampling period). Pups were weighed by centre personnel on alternate days at centre II and approximately once a week at the other centres. Estimated pup weight on sampling day was calculated from the nearest pup weighing and average growth rate during the sampling period.

Urine sampling

This study used urine rather than blood sampling in order for the sampling to be completely non-invasive. The urine samples were collected opportunistically if the pups urinated while being handled during routine feeding or weighing. The authors (SW and SV) collected samples at centres II, IV and V while centre personnel collected samples at centres I and III. Collection was usually with a syringe from the ground beside the pup or directly from the stream of urine, although one sample from centre II (TSR-14-18; S2-dataset) was collected on a 'puppy pad' (plastic-backed tissue layers). Some samples at centre I were collected from a tray under the slatted floor of the pen during morning cleaning (representing urine voided at any time since the previous evening). Samples were transferred to standard plastic sample pots with a record for each sample of the source pup, date and collection time.

Samples were immediately transferred to a freezer (at least -18°C) and stored until analysis. Only

samples visibly uncontaminated with faeces and with at least 10 ml were used in the analyses.

Since a study of plasma cortisol levels in captive harbour seals found a diurnal pattern of plasma

cortisol levels, peaking at 01:00 and waning around 13:00 [27], the time of day of each sample

collection was recorded and samples used in the analyses (n=112) were assigned to time categories:

0700–12:59; 13:00–18:59; 19:00–00:59 (Fig 1); no samples were collected during the overnight period.

Fig 1. Distribution of urine sample collection between three 6-h periods at each centre (from all
samples day 2 onwards).

Glucocorticoid (GC) analysis

The urinary glucocorticoid (GC) analyses were carried out at three laboratories. Lab 1was at Riseholme

Park, University of Lincoln; Lab 2 was at the Institute of Global Food Research, Queens University,

Belfast, and Lab 3 was the Chemical and Immunodiagnostic Sciences Branch, Veterinary Sciences

Division, Agri-Food & Biosciences Institute (AFBI), Belfast. Seventeen samples from Centre I (2012)

were analysed by ELISA for cortisol (CL) at Lab 1. Eleven duplicate samples from Centre 1 and a

further 8 from centre II (2013) were analysed at Lab 2 for CL by ultra-performance liquid

chromatography – tandem mass spectrometry (UPLC−MS/MS) [28, 29]. Ninety-six further samples

collected from centres II, III, IV and V in 2014−15 were analysed at Lab 3 for CL, cortisone (CN),

prednisolone (PL) and prednisone (PN) by UPLC−MS/MS. These 96 samples included seven in which

the sample donor pup (one of a pair) was not identified; these seven samples were included in analyses

of the overall range of GC concentrations recorded and of the possible effect of the time of day of

sampling, but excluded from subsequent analyses. The full list of samples for each of the three

analytical laboratories and methods are given in S2 Dataset.

Urinary creatinine concentrations.

In order to control for the water content of the sample, urinary GC concentrations are expressed as a

ratio to the creatinine concentration (ng GC mg/Cr) in each sample [24]. Creatinine (Cr) concentrations

at all three laboratories were analysed using a colorimetric method modified from the Jaffe reaction, in

which creatinine in an alkaline solution reacts with picric acid to form a coloured complex [30]. For the

basic method, a standard Cr solution is prepared for use as a reference. A picric acid solution is added

to the urine sample and the Cr standard. The colour reaction is started by adding sodium hydroxide.

The resulting absorbance of each sample is then measured using a spectrophotometer in order to finally

obtain the Cr concentration of the urine sample. The detailed methods of Cr determination followed by

each laboratory are given in S3 Protocol and S4 Protocol.

Analysis of urinary cortisol concentrations by ELISA

The principle of the ELISA assay is the binding of a specific antibody to an enzyme-labelled target

antigen (CL in this case). The enzyme activity is measured using a solid substrate that changes colour

when modified by the enzyme. The intensity of the colour depends on the CL concentration and is read

on a microplate.

Urinary CL concentrations in 17 samples from centre I in 2012 were determined at Lab 1 using a

cortisol ELISA kit (Cat No ADI-900-071) for colorimetric detection from Enzo Life Sciences (UK)

Ltd. The details of the procedures undertaken are given in S3 Protocol. This method is stated to be for

plasma, saliva, serum or urine from any species. The cross-reactivity with other steroid hormones is

stated by the manufacturer to be Cortisol (100%), Prednisolone (122.35%), Corticosterone (27.68%),

11-deoxycortisol (4%), Progesterone (3.64%), Prednisone (0.85%), Testosterone (0.12%) and $\leq 0.10\%$:

Androstenedione, Cortisone and Estradiol.

The method (S3 Protocol) followed the kit instructions. Because the concentration range of CL in

harbour seal urine was not known, a test ELISA run was initially carried out to estimate the urine

sample dilutions needed for the middle calibrator of the ELISA kit. After converting the ELISA colour

changes on a microplate, the CL concentrations in the original sample were obtained by multiplying by

the dilution factors, and converted to mM/L using the molecular mass of cortisol. These values were

then divided by mM creatinine to obtain the cortisol/creatinine (CL/Cr) molar ratio.

Analysis of urinary glucocorticoid concentrations by Ultra Performance Liquid Chromatography–

tandem mass spectrometry (UPLC–MS/MS)

Immunoassays – as the ELISA described above – may be used to determine the concentration of a single

steroid hormone (assuming cross-reactivity is not considered to be high), but if the concentrations of

several steroids are required, either separate kits for each steroid must be used (requiring splitting of the

sample), or the steroids need to be separated by chromatography. Liquid chromatography coupled with

tandem mass spectrometry (LC–MS/MS) can quantify either one or multiple steroids at wide-ranging

concentrations, while avoiding the issue of antibody cross-reactivity [26].

For this study of seal pup urine glucocorticoid steroids, ultra-performance liquid chromatography

coupled with tandem mass spectrometry (UPLC–MS/MS) was employed by Lab 2 to determine urinary

cortisol (CL) concentrations of 19 samples collected in 2012–13 and by Lab 3 to determine

concentrations of the four GCs, CL, cortisone (CN), prednisolone (PL) and prednisone (PN) in 96

samples collected in 2014–2016. The CL analyses of the 2012-2013 samples by Lab 2 were carried out

alongside a separate study of CL and CN concentrations in cattle [31], while the analyses of CL, CN, PL

and PN in the 2014–2016 samples by Lab 3 were carried out according to the procedures for analysis of

bovine liver as part of the routine statutory Veterinary Drug Residue testing programme in N. Ireland.

The UPLC–MS/MS methods employed by both Labs 2 and 3 are described in full in S4 Protocol.

Method validation. The seal pup urine samples were analysed by UPLC-MS/MS at laboratories where

the analytical method validation was previously undertaken for analysis accuracy/recovery, intra- and

inter-day repeatability and linearity for bovine urine in Lab 2 [31] and bovine, porcine and ovine liver in

Lab 3, in accordance with EU guidelines (2002/657/EC). The method validation procedures are described

in S4 Protocol.

Statistical analyses of GC results

Statistical analyses were carried out in the XLSTAT-pro statistical package, using ANOVA, correlation

and regression, and non-parametric Kruskal-Wallis or Mann-Whitney tests. Statistics were carried out

both (*1*) on median GC concentrations for individual pups and (2) on all GC concentrations from all

samples (*2*). To take into account the rehab environment variables, analysis of covariance (ANCOVA)

was carried out using the R statistical package [32].

RESULTS

Range and median concentrations and correlation matrix of four glucocorticoid hormones (GCs)

by UPLC-MS/MS

The range of creatinine (Cr) concentrations in the 115 samples analysed by MS was 115–15576 µM/L,

(mean 3219, median 2273 µM/L). The range of glucocorticoid (GC) concentrations (ng/mg Cr) was

greatest for cortisol (CL) (0–8062 ng/mg; next for cortisone (CN) (0–4853 ng/mg), with a lower range

for prednisolone (PL) (0–236.6 ng/mg) and prednisone (PN) (0–142.5ng/mg). The complete set of raw

data is given in S2 Dataset, a summary of the dataset is given in Table 1 and the range of MS results on

a $\log_{10}$ scale is presented in Fig 2.

Fig 2. Scattergram of urinary concentrations of four GCs from all samples analysed by UPLC MS/MS.
(n=CL–115; CN, PL, PN – 96). Concentrations ng/mg creatinine on $\log_{10}$ scale. Horizontal lines indicate
medians and + markers indicate means.
211

The median values for CL, PN and PL were similar to each other (36.9, 31.1 and 31.57 ng/mg Cr

respectively), while median CN concentrations were an order of magnitude higher (390.4 ng/mg Cr).

There were seven samples with CL and PL concentrations below the detection limit (Fig 2). Six of

these samples had low Cr concentrations (range 115–683 µM/L). One sample which gave a very high

CL concentration (2030 ng/mg Cr) also had a low Cr concentration (444 µM/L). Because some very

high or very low results for concentrations from dilute samples, the median concentrations rather than

the means are used in continuing analysis of the data.

The correlation matrix (Table 2) for all four GCs for the samples measured by UPLC MS/MS indicates

that CL, CN and PL were significantly correlated with each other. PN was not correlated with CL and

correlated relatively weakly with PL and CN.

Table 2. Correlation (Pearson coefficient) between concentrations of four GCs (ng/mg creatinine) by
UPLC MS/MS (n=96 samples)

	Cortisol	Cortisone	Prednisolone	Prednisone
Cortisol	1	0.751	0.726	0.070
Prednisone	0.070	0.262	0.362	1
Cortisone	0.751	1	0.504	0.262
Prednisolone	0.726	0.504	1	0.362

225   The ratios of mean and median CL concentrations to the other three GCs from all samples from which

226   all four GCs were analysed are given in Table 3.

228 Table 3. Ratios of mean and median concentrations (ng/mg Cr) of Cortisol to each of the other GCs in
229 all samples for which all four GCs were analysed (n=96 samples).
230

	Cortisol	Cortisone	Prednisolone	Prednisone
Means	214.37	526.19	39.91	36.68
SD	922.39	617.28	43.62	26.73
Median	35.96	390.38	30.42	31.72
Ratio of CL: other GC (mean)		0.41	5.37	5.84
Ratio of CL: other GC (median)		0.09	1.18	1.13

Comparison of cortisol/creatinine concentrations analysed by ELISA and UPLC MS/MS

233 11 of the 17 samples from centre I were analysed for cortisol by both UPLC-MS/MS and by ELISA,

234 The correlation (Pearson coefficient) between the MS and ELISA results was 0.795 (2-tailed P <0.01),

235 although the mean concentrations for the ELISA results were about 13% higher on average than for

236 MS (Table 4).

237 Table 4. Cortisol concentrations (ng/mg creatinine) comparing UPLC-MS/MS with ELISA

	n	Minimum	Maximum	Median	Mean	SD	Correlation (Pearson coefft)
MS/MS	11	21.8	221.2	73.5	86.9	54.5	0.795
ELISA		46.0	211.0	98.0	98.1	47.0	

Comparison of GC concentrations of newly-admitted pups with those from rehab day 2 onwards

Since pups' GC concentrations when they first arrive at the centre are likely to reflect their stranding

conditions, handling and transport to the centre, concentrations in samples taken on rehab days 0–1

were tabulated separately from samples from rehab days 2 onwards (Table 1). The single or median

concentrations for each pup in days 0–1 (n= 8 pups sampled for CL, 6 pups for PN, CN and PL) were

compared with single or median concentrations in day 2 onwards (n= 34 pups sampled for CL, 34 for

PN, CN and PL). CL, PN and PL concentrations were generally higher on days 0-1 (Mann-Whitney U

test; 1-tailed P <0.05 for each GC), although CN concentrations did not differ (P = 0.328). Since the

aim of the study was to investigate the influences of the rehabilitation environment on pup GC levels,

and not the pups' pre-admission history, the 12 samples taken from 8 pups on days 0–1 (Table 1 and

Supplementary file S2) were separated from the continuing analyses, which considered pups from

rehab days 2 onwards.

Effect of time of day of sampling on GC concentrations

Distribution of sampling from rehab day 2– indicated that the majority of samples were taken in the

morning 6-hour period (0:700–12:59) and afternoon (13:00–18:59), with fewer in the evening (19:00–

00:59) (Fig 1). Comparison of concentrations of CL and PN in each of the three 6-hour periods

between 07:00 and 00:59 indicated no significant differences in concentrations of either of these GCs

according to time of day of sampling (Table 5).

Table 5. Median concentrations of urinary cortisol and prednisone (ng/mg creatinine) and significance
of differences between each of three 6-hour time periods between 07:00 and 00:59. (samples from pups
day 2 onwards).
261

Time period	07:00–12:59	13:00–18:59	19:00–00:59	
CENTRE	Median concentration ng/mg creatinine			P value
Cortisol				
I	84.0†	97.0†	no samples	0.846
II	22.27	11.50	15.78	0.258
IV	82.6	298.38	87.17	0.994
V	27.67	45.01	no samples	0.859
Prednisone				
II	26.88	11.55	8.72	0.091
IV	41.52	36.29	34.85	0.426
V	41.97	49.55	no samples	0.680

262   †– concentrations from ELISA; P values from Kruskal-Wallis 3-sample test or Mann-Whitney 2-
263   sample test; number of samples as in Fig 1.

Analysis of variance (ANOVA) and covariance (ANCOVA) of GC concentrations (ng/mg

creatinine) according to variables considered in study

The results for goodness of fit, R^2, by the ANOVA were similar for median GC concentrations for

individual pups and for all samples from all pups (Table 6): 47% and 49% of the variability of PN

concentrations was explained by the four and seven variables respectively, while less (25–36%) of the

variability of CL, CN and PL was explained by the same variables. The P value of the F test indicates

significant influence of the four variables only for PN for (*1*) the median individual pup concentrations.

The F test for GC concentrations for all samples indicates significant influence of the seven variables

for (*2*) all GC concentrations, but the highest level of significance for PN (Table 6).

Table 6. ANOVA for GC concentrations (ng/mg creatinine) results
275

	df	**CORTISOL (CL)** N = 34(*1*); 96(*2*)			df	**CORTISONE (CN)** N=30(*1*); 78(*2*)			**PREDNISOLONE (PL)** N=30(*1*); 78(*2*)			**PREDNISONE (PN)** N=30 (*1*); 78(*2*)		
Goodness of fit R²														
1	25	**0.289**			22	**0.340**			**0.286**			**0.466**		
2	84	**0.246**			67	**0.356**			**0.296**			**0.487**		
		Mean sq	**F value**	**Pr(>F)**		**Mean sq**	**F value**	**Pr(>F)**	**Mean sq**	**F value**	**Pr(>F)**	**Mean sq**	**F value**	**Pr(>F)**
ANOVA-Fisher's F test														
1	11	45857	1.27	0.302	10	247642	1.618	0.183	51440	1.26	0.315	781	2.74	**0.033**
2	11	104240	2.49	**0.009**	10	600346	3.71	**0.001**	1369	2.81	**0.006**	1996	6.35	**<0.0001**

276   *1*: Results from median concentrations for individual pups; 4 variables considered (*growth rate*, *centre*, *social group*, *sex*)
277   *2*: Results from concentrations in all samples from all pups; 7 variables considered (*growth rate*, *weight on sample day*, *rehab day*, *Centre*, *social group*, *free water access*,
278   *pup sex*)
279

GC concentrations according to social group and rehab centre

Because five pups from centre IV changed *social group* during the sampling period (Table 1), the GC concentrations from all samples (rather than medians for each pup) from pups in each of the social groups are visualised in Fig 3. The CL, PN and PL concentrations differed according to social group (P=0.006, 0.008 and 0.004 respectively, Kruskal-Wallis test), with social group 2 (paired pups) having relatively lower concentrations (Fig 3). The CN concentrations did not differ between the social groups (P=0.402).

Fig 3. Scattergrams of distribution of urinary concentrations of four GCs according to social group

Social group categories: 1 (alone), 2(paired) and 3 (≥3 pups). Samples from rehab d. 2 onwards; n=96 samples for CL, n=78 samples for PN, CN and PL. Horizontal lines indicate medians and + markers indicate means

The distribution of each GC (considering all samples from all pups) differed between *centres* (Fig 4; $P < 0.0001$ for each GC; Kruskal-Wallis test). Centre IV samples had overall the highest CL, CN and PL concentrations, whereas Centre II had the lowest CL, PN and PL concentrations. Scatterplots for median concentrations for individual pups from each centre (not shown) showed the same pattern, still with significance (CL – P = 0.003; PN – P = 0.011; CN – P =0.021, PL – P = 0.029; Kruskal-Wallis test).

Fig 4. Scattergrams of distribution of urinary concentrations of four GCs.
(all samples from rehab day 2 onwards) in each rehabilitation centre I–V. Horizontal lines indicate medians and + markers indicate means; n= 96 samples CL, 78 for PN, CN and PL.

The ANCOVA indicated which variables significantly influenced the variability in the GC concentrations. Considering (*1*) – median GC concentrations for individual pups – PN was significantly influenced by average *growth rate* during the sampling period, but not significantly by the other three variables; the other GCs were not significantly influenced by any of the four variables (Table 7). Considering (*2*) – GC concentrations of all samples – PN was most strongly influenced by *growth rate*, but also by *pup weight on sample day* (Table 7). However, the concentrations of the other GCs were

308   also apparently influenced by *growth rate* and *Centre* (PL), *growth rate*, *weight on sample day* and

309   *Centre* (CN) and *weight on sample day*, *Centre* and *social group* (CL) (Table 7).

Table 7. ANCOVA for GC concentration (ng/mg creatinine) results

Variables	Df	CORTISOL (CL) N=34(*1*); 96(*2*) Mean sq	F value	Pr(>F)	df	CORTISONE (CN) N=30(*1*); 78(*2*) Mean sq	F value	Pr(>F)	PREDNISOLONE (PL) N=30(*1*); 78(*2*) Mean sq	F value	Pr(>F)	PREDNISONE (PN) N=30(*1*); 78(*2*) Mean sq	F value	Pr(>F)
(*1*) MEDIAN VALUES FOR INDIVIDUAL PUPS (rehab day 2 onwards)														
Growth rate	1	45292	1.30	0.265	1	260203	1.74	0.200	766	1.96	0.175	3242	11.76	**0.002**
Centre	4	148981	1.07	0.392	3	252221	1.69	0.198	739	1.89	0.160	291	1.05	0.388
Social group	1	141227	4.05	0.055	1	583406	3.90	0.060	1558	3.98	0.058	1045	3.79	0.064
Sex	1	26977	0.77	0.387	1	62396	0.42	0.525	248	0.63	0.434	237	0.86	0.364
(*2*) VALUES FOR ALL SAMPLES (rehab day 2 onwards)														
Growth rate	1	97249	2.32	0.132	1	1586864	9.94	**0.002**	3927	7.51	**0.008**	13842	44.30	**5.9E-09**
Weight on sample day	1	265843	6.34	**0.014**	1	1287950	8.07	**0.006**	0	0	0.983	2274	7.28	**0.009**
Rehab day	1	19576	0.47	0.496	1	444196	2.78	0.100	1358	2.60	0.112	973	3.11	0.082
Centre	4	106516	2.54	**0.046**	3	705575	4.42	**0.007**	1747	3.34	**0.024**	742	2.38	0.078
Social group	1	182494	4.35	**0.040**	1	345182	2.16	0.146	11	0.02	0.884	143	0.46	0.501
Water access	1	7155	0.17	0.681	1	3246	0.02	0.887	86	0.16	0.687	319	1.02	0.316
Sex	1	105517	2.52	0.117	1	217899	1.37	0.247	120	0.23	0.634	6	0.02	0.893

1: Results from median concentrations for individual pups; 4 variables considered (*growth rate*, *centre*, *social group*, *sex*)

2: Results from concentrations in all samples from all pups; 7 variables considered (*growth rate*, *weight on sample day*, *rehab day*, *Centre*, *social group*, *free water access*, *pup sex*)

Pup growth rates

We considered the possibility that pup weight at entry to the rehab centre may have affected its subsequent growth rate at the centre. The entry weight of the pups ranged from 6.9 to 13.1 kg (n = 34 pups). Regression of individual pup growth rate during the sampling period by entry weight (Fig 5) indicated there was no relationship between entry weight and subsequent growth rate in our sample of pups ($R^2 = 0.010$, P = 0.576).

Fig 5. Regression of pup growth rate (kg/day) by entry weight (kg)

The entry weights of the six pups from centre II ranged from 7.1 to 11.1 kg (Table 1). Frequent weight records for these pups over a four-week period indicated similar linear growth rates for all the pups (S5 Supporting Information), irrespective of their entry weight (range 0.27–0.30; Table 1), $R^2 = 0.97$–0.99, P < 0.0001).

Regressions of GC concentrations by pup growth rates, pup weight on sample day and days since entry into rehab

Linear regressions of median GC concentrations for each pup (n = 34 for CL, n = 30 for PN, PL and CN) by *growth rate* (kg/day) showed a significant negative correlation only for PN, indicating generally lower concentrations of PN in pups with higher growth rates (Table 8a; Fig 6).

Fig 6. Regression of median GC concentrations (ng/mg creatinine) for each pup by growth rate (kg/day)

Linear regressions of GC concentrations in each sample (n=78) by pup estimated *weight* showed a significant negative correlation for all four GCs, with the strongest effect of pup weight (explaining 24% of the variability) on PN concentrations (Table 8a). However, because of the opportunistic nature of the urine sampling, pups at centres II–V were sampled at different weight ranges (Fig 7), with the

centre explaining a further 19% of the PN variability (combined R^2=0.436). There was no association found between the weight of pups in centre I and CL concentrations (Table 8c), but the range of pup weights in centre I over the sampling period was very small, owing to low growth rates (Table 1).

Table 8. Values of R^2 for regressions of concentrations (ng/mg creatinine) of each GC (Cortisol – CL, Cortisone – CN, Prednisolone – PL, Prednisone – PN) by pup growth rate, estimated pup weight on sample day and rehab day (day 2 onwards)

	(a) All 34 pups in the study n=34 (CL), 30 (CN, PL, PN) (pup medians) n= 96 (CL), 78 (CN, PL, PN) (all samples)				(b) Six pups in centre II n= 32 samples (CL), 24 (CN, PL, PN)				(c) Four pups in centre I n= 17 samples (ELISA)	
	CL	CN	PL	PN	CL	CN	PL	PN	CL	CN/PL/PN
Growth rate	0.036	0.051	0.056	**0.276**	NA	NA	NA	NA	NA	NA
Pr > F	*0.285*	*0.230*	*0.210*	*0.003*						
Weight	**0.051**	**0.145**	**0.073**	**0.242**	0.079	**0.193**	**0.190**	**0.348**	0.001	NA
Pr > F	*0.027*	*0.001*	*0.017*	*<0.0001*	*0.119*	*0.032*	*0.033*	*0.002*	*0.928*	
Rehab day	**0.05**	**0.145**	**0.126**	**0.16**	0.089	0.139	**0.278**	**0.369**	0.025	NA
Pr > F	*0.034*	*0.001*	*0.001*	*0.0002*	*0.098*	*0.072*	*0.008*	*0.002*	*0.544*	

Fig 7. Regression of Prednisone concentrations (ng/mg creatinine) by estimated pup weight (kg) on day of sampling for pups from each centre II–V

Pups generally gain weight during rehab, as exemplified by the six pups in Centre II, which were weighed regularly over a six-week period (S5 Supporting Information) Therefore, over all the pups in the five centres of this study, pup weight is conflated to a considerable extent with rehab day. The effect of rehab day (from day 2 onwards) was considered separately for the Centres I and II pups, for which there were samples taken over a relatively long period. For the centre II pups, the regressions showed a significant negative correlation with rehab day for PN and PL – most strongly for PN – but the correlation was non-significant for CL or CN (Table 8b). For the centre I pups, there was no association between rehab day and CL (Table 8c).

DISCUSSION

Choice of urine sampling and UPLC-MS/MS for sample analysis

Our choice or urine sampling was advantageous in that it caused no additional stress to the pups beyond their routine care. Its disadvantage was that samples could only be collected opportunistically when the pup happened to urinate during routine feeding or other care. The wide variability of GC concentrations found, even between samples from the same pup, is consistent with the conclusion of Cook [20] that the use of singly voided samples is not useful in assessing an animal's stress response. Although the variability might be in part due to the different times of day of sample collection [27], we found no significant difference between the samples collected at different times. The reason for this might be partly that the concentrations in a urine sample likely represent average circulating blood plasma concentrations over the preceding few hours [33].

Our GC analysis of the first set of samples intended for cortisol (CL) was carried out using an ELISA kit with known high cross-reactivity to prednisolone (PL), which undoubtedly explains at least in part why the ELISA CL concentration results for the centre I samples were on average 13% higher than the duplicate samples analysed by UPLC-MS/MS. This cross-reactivity with similarly-structured steroids is a recognised problem with any form of immune-assay (IA) for CL levels in humans, where other synthetic GCs may be present [34, 35], but at the time of our ELISA analysis in 2012, the MS analysis of the 2014–16 samples had not been carried out and therefore the existence of significant amounts of naturally-occurring PL in our seal pups was not suspected. As Koren and colleagues [26] have pointed out, distinguishing between similar steroids in a single sample requires a mass-spectrometric technique which can unequivocally discriminate between these similar GCs.

Comparison of seal pup urinary GC concentrations with other species

Cortisol. This is the first investigation into urinary GCs concentrations in harbour seals, and baseline levels of urinary CL for healthy pups are not known. Urinary CL has only been studied in one other

seal species, the Weddell seal (*Leptonychotes weddelli*) in the wild. Concentrations in juveniles <1

year were much higher than in the harbour seal pups of our study, with means of ~930–2370 ng/mg

Cr by radio-immuno-assay (RIA). However, it is thought that Weddell seal CL concentrations are

unusually high amongst phocid species [24].

The overall median and mean of our seal pup CL concentrations (overall median 37 ng/mg Cr) were

of the same order of magnitude as urinary concentrations found in cattle (~36–84 ng/mg Cr by UPLC-

MS/MS [31]) and in free-range lambs (~20–46 ng/mg Cr by RIA [36]).

Cortisone. CN is a downstream oxidised metabolite of CL [37], with less than half of CL's biological

activity [38, 39]. Except for a few samples with very high CL concentrations, CN concentrations in

our pup urine samples were an order of magnitude higher than for CL, with the CL:CN ratio of

median concentrations of 0.09: 1 (Table 3). This is much less than the average ratio for human

athletes of 0.64:1 [40], and also much less than that reported from median concentrations in free-

ranging Chianina cattle aged 1–2 years (1.43:1 in males, 1.02:1 in females [41]) and in young Friesian

bulls in pre-slaughter enclosures (1.35–1.68:1) [31].

Prednisolone. The ratio of CL:PL median concentrations in our seal pups was 1.2:1 (Table 3), i.e.

close to parity. In humans, urinary PL has been found to occur only in concentrations 100-times

lower than CL [42]. Naturally-occurring PL has been found in low concentrations in Chianina cows

(median CL:PL ratio of 25:1 (none present in bulls), and even lower concentrations in Friesian veal

calves (mean CL:PL ratio range 29–64:1 [43]). It has been suggested [44] that neo-formation of PL

due to microbial action may occur in bovine urine after collection if the samples are kept at a high

temperature (e.g. 37°) or if there is contamination (e.g. from faeces). However, we believe the PL in

our seal pup samples was endogenous since (i) the samples were not subjected to high temperatures,

were free from faecal contamination, and were transferred to the freezer within minutes, (ii)

concentrations were consistently at similar concentrations to cortisol and (iii) PL concentrations were

significantly correlated with CL, which would not be the case with post-collection microbial action.

Prednisone. PN is an oxidised metabolite (or precursor) of PL. Synthetic PN is widely used as an

anti-inflammatory agent in the treatment of human conditions and, often illegally, in domestic cattle.

Endogenous urinary PN has so far been reported to be absent in Piedmontese and Friesian cattle [45]

and Chianini cattle [41], present in trace amounts in a small proportion of Friesian veal calves (CL:

PN ratio 43:1 [43]), but has not been reported in other mammal species. Its occurrence in our seal pup

urine, with the median CL:PN ratio of only 1.13:1 (Table 3), i.e. close to parity, was therefore

unexpected. The lack of inter-correlation between PN and the other three GCs might be due in part to

a feedback effect by circulating PN suppressing adrenal function, possibly affecting secretion of the

other GCs [46].

Psychological stress and PL and PN concentrations in bovine urine. Concentrations of PL and PN in

bovine urine above the detection limit have been found to be more frequent in conditions of extreme

psychological stress, such as slaughterhouse conditions [43, 45]. The median concentrations of PL

and PN of a similar order of magnitude to CL routinely found in our seal pup urine in all five centres

are unlikely to be related to extreme psychological stress, since relaxed social interaction and play

behaviours between pups – which would be unlikely in a highly stressful environment [47, 48] –

were typical of pups held with companion pups, as in centres II, IV and V (S.Wilson, pers. obs.). In

harbour seal pups, therefore, it seems that the trigger for PN and PL secretion is different from the

bovines studied hitherto.

Possible association between levels of each GC and rehab procedures and conditions

Interpretation of our data is challenging because of the large number of variables considered in the

analysis as well as potential variables not considered, in relation to the relatively small dataset. These

limitations are a result of our studying pups opportunistically in existing rehab centres, each with their

independent procedures, rather than according to any controlled or experimental design.

Pup growth rate and weight on sample day. The ANCOVA and regression analyses indicated that the

variables chiefly influencing PN concentrations were pup growth rate and pup weight on the sampling

day, but that actual weight also influenced CN, PL and CL concentrations. Elevated CL has

previously been associated with poor nutrition and growth in human children [49] and with the post-

weaning fasting period in harp seal pups [50]. High endogenous CL levels in fasting post-weaning

grey seal pups have been shown experimentally to increase protein catabolism, which may

compromise pup survival during the period between weaning and successful post-weaning foraging

[51].

Since free-living harbour seal pups suckled by their mothers on milk typically have an average net

mass gain of ~0.4 to ~0.6 kg/day from a birth weight of ~11 kg [2–4], even our rehab pups with the

highest weight gain (0.27–0.30 kg/day in centre II) have barely half the net mass gain of free-living

pups. Because the urinary concentrations of PN appeared to be the most closely linked negatively

correlated) of the four GCs to pup growth rate and weight, we suggest that its presence in our rehab

pups of nursing age may be an age-specific response to the failure of pups in rehab to achieve the high

normal growth rate of >0.5 kg/day in naturally suckling pups of this species and therefore it is likely

that the PN concentrations seen in our rehab pups may be much higher than in healthy free-living

suckling pups.

Pup social group and Centre. *Social group* was to a considerable extent conflated with *Centre*, since

all pups in centres I and III were social group 1 (isolated pups), all pups in centre II were paired, and

all pups in groups of more than two pups were in Centre IV. CL was the only GC that the ANCOVA

indicated had a significant association with *social group*, with social group 2 (paired pups) overall

apparently having lower CL than either isolated or group pups.

Rehab conditions in which pups were kept alone in dry pens (*social group* 1) evidently did not mimic

key aspects of the pup's natural social or physical environment, and might therefore be interpreted as

relatively poor welfare with possible chronic stress or distress due to lack of stimulation [52].

However, although the isolated *social group* 1 pups in centre I did have relatively elevated CL levels,

the similarly isolated pups in centre III had relatively low (and sometimes undetectable) CL. This

significant difference between these two pup groups might be due to negative feedback mechanisms

(18, 53) operating in the centre III pups, but other factors – possibly including the very low growth

rate and body mass – being over-riding for the centre I pups.

Pup GC response to maternal separation and isolation. On losing contact with their mother, harbour

seal pups vocalise [54], swimming around and attempting to follow other mothers or pups (S. Wilson,

pers. obs.), while pups captured in the field while separated from their mothers have higher CL levels

than pups captured with their mothers [55]. The elevated CL levels typical of this first 'acute' stage of

maternal separation trigger the energy release necessary to fuel the typical agitated activity and

vocalisation shown by a pup trying to reconnect with its mother (56). By the time orphan seal pups

enter rehab centres, they will already have suffered this initial first stage of maternal separation.

CL levels are expected to reduce during the next "depressive" stage of separation, when the pup is

likely to show reduced locomotion and other more passive behaviours [56, 57, 58]. Pups kept alone in

pens without water access (centres I and III) do usually appear inactive and passive (S. Wilson, pers.

obs.) and may possibly be in a continuing depressive phase, possibly involving a feedback mechanism

which suppresses the cortisol response [18, 58]. By contrast, pups maintained socially with water

access (centres II and IV) show alternating rest and activity periods, with much behaviour typical of

pups with their mothers, and do not show signs of a 'depressive' stage of separation (S. Wilson, pers.

obs.).

Factors likely to affect urinary GC concentrations in orphan pups unrelated to rehab centres

conditions

Possible differences in pup GC concentrations due to regional population of origin. The pups with

generally relatively lower CN and PL levels were mostly from Irish Sea centres (II and V) and the

lower CL-level pups were also mostly from Irish Sea centres (II, III and V). One possible suggestion

for future investigation is whether the relatively elevated levels of GCs in pups from the southern

North Sea centres (I and IV) compared with centres in the Irish sea (centres II, III and V) might be, at

least in part, due to elevated levels of persistent organic pollutants (POPs) in the southern North Sea

and Wadden Sea [59]. POP contaminants in maternal tissues may be transferred to the foetus, posing

health risks during development [60]. Studies *in vitro* [61] and in adult polar bears [62] have

suggested that POPs can disrupt GC activity, potentially altering circulating CL levels. Future studies

might investigate the POP concentrations in tissues of pups currently entering seal centres in the

southern North Sea and Wadden Sea (Centres I and IV) compared with the Irish Sea and elsewhere.

Effects of interval between maternal separation and 'orphan' pup rescue. Several factors including

maternal separation time, body condition and infection status may all influence pups' GC cortisol

level on arrival at rehab centres. Pups separated for a few days may have opportunistic infections,

often causing pneumonia with respiratory distress (as 'Ula' in centre II), which is known to cause an

additional stress response in new-born babies [63]). We feel that our finding – that the GC

concentrations in newly arrived pups was generally higher than from rehab day 2 onwards – justifies

our omission of the results in days 0–1 from the main analyses. Our additional finding that body mass

at the time of stranding and entry into the centre was unrelated to growth rate during the sampling

period also suggests that poor body condition at stranding did not affect orphan pups' subsequent

physical development during rehab.

Conclusions and recommendations

This study has been preliminary and empirical, recording associations between urinary GCs and

variables in rehab conditions and procedures without a physiological explanation.

We found that the most significant association between GC concentrations and the variables

considered was a negative relationship between prednisone (PN) and growth rate or body weight. If

our preliminary findings should be confirmed by an extended, independent study with more controlled

variables, this would provide a strong argument for providing pups with an appropriate high fat

formula to maximise growth rate from an orphan pup's earliest days in rehab. Our results suggest that

slow or negative growth rates [12] for some weeks would likely result in chronically elevated

circulating PN levels, which could have a negative effect on a pup's future responses to stress as well

as possible neural, cognitive and immune system impairment [18, 64]. Increasing average growth rate

to an approximation of 0.3 kg/day, as in centre II, would have a positive and practical advantage of

enabling orphan pup release after 6–7 weeks [14]. Such a relatively early release potentially

maximises the opportunity for pups to develop their survival skills alongside free-living pups from

their original cohort at the time when the pups are initially dispersing from their natal sites and

learning to forage during their first post-weaning weeks [65].

Our study did not find a clear relationship between urinary GC levels and the pup's social or physical

environment in rehab, although – since a significant effect of social group on CL levels was found –

social and environmental variables as contributory factors could not be ruled out. We therefore

consider that the psychological welfare of seal pups in rehab may be best assessed using multimodal

indices of welfare, including behavioural as well as physiological components [25]. The latter could

in future investigate the benefits of keeping pups socially using markers of positive emotion, such as

oxytocin [66, 67].

ACKNOWLEDGEMENTS

We are grateful to Richard Roets at the University of Lincoln for carrying out the ELISA analyses,

Joanna Keenan for carrying out the initial UPLC-MS/MS analyses Queens University, N. Ireland, and

to Brett Greer for his help in facilitating the analyses. We would especially like to thank Professor

Daniel Mills at the University of Lincoln, Professor Chris Elliott at the Institute for Global Food

Research, Queens University, and Dr Robert McCracken at the Agri-Food & Biosciences Institute, N.

Ireland for permitting the analyses in their departmental laboratories.

We are indebted to all the personnel at the seal sanctuaries participating in this study for their

generosity in permitting and facilitating the urine sampling. In particular we would like to

acknowledge the invaluable support of Duncan, Richard and Nicki Yeadon and Curtis Jones at

Natureland, Skegness, England; Will Matthews during the study period at Skegness, Fleur Brochut at

Tara Seal Research, N. Ireland; Tania Singleton at Exploris, N. Ireland; Tanja Rosenberger and Janne

Sundermeyer at Seal Centre Friedrichskoog, N. Germany, and Ally McMillan, Lisa Becker and Sonja

Ciccaglione at Seal Rescue Ireland, Wexford, Ireland. Thanks to Janne Sundermeyer for helpful

comments on the draft manuscript.

CONFLICTS OF INTEREST

540 The authors have no conflicts of interest

FUNDING

542 This study has not received any grant from any funding agency.

ETHICS STATEMENT

The urine sampling in this study was carried out with the permission and collaboration of each of the

five seal centres. Four of the five centres (I, III, IV and V) were in the public domain and acting

within relevant government guidelines. Centre II was a voluntary private practice operating with local

government (N. Ireland) permission. The sampling did not cause additional disturbance to the pups

beyond their routine care and handling and did not affect their welfare. No specific license for this

study was therefore required.

REFERENCES

1. Bowen WD, Boness DJ, Iverson SJ. Diving behaviour of lactating harbour seals and their pups during maternal foraging trips. Can J Zool. 1999; 77: 978–988.

2. Skinner JP. Physical and behavioural development of nursing harbour seal (*Phoca vitulina*) pups in Maine. MSc thesis, University of Maine; 2006. Available from: https://digitalcommons.library.umaine.edu/cgi/viewcontent.cgi?article=1655&context=etd

3. Muelbert MMC, Bowen WD. Duration of lactation and postweaning changes in mass and body composition of harbour seal, *Phoca vitulina*, pups. Can J Zool. 1993; 71: 1405–1414.

4. Cottrell PE, Jeffries S, Beck B, Ross PS. Growth and development in free-ranging harbour seal (*Phoca vitulina*) pups from southern British Columbia, Canada. Mar Mamm Sci. 2002; 18: 721–733.

5. Lang SLC, Iverson SJ, BowenWD. Individual variation in milk composition over lactation in harbour seals (*Phoca vitulina*) and the potential consequences of intermittent attendance. Can J Zool. 2005; 83: 1525–1531. doi.org/10.1139/z05-149.

6. Venables UM, Venables LSV. Observations on a breeding colony of the seal *Phoca vitulina* in Shetland. Proc Zool Soc Lond. 1955; 125: 521–532. doi:10.1111/j.1096-642.1955.tb00614.x

7. Wilson S. Mother-young interactions in the common seal *Phoca vitulina vitulina*. Behaviour 1974; 48: 23–26.

8. Wilson SC, Jones KA. Behaviour of harbour seal (*Phoca vitulina vitulina*) mother-pup pairs in Irish Sea intertidal habitats. Biol Environ. 2018; 118: 1–15. doi:10.3318/BIOE.2018.02.

9. Wilson SC. Social organization and behavior of harbor seals, *Phoca vitulina concolor*, in Maine. Report to the Marine Mammal Commission PB-280 188. Contract no. MM6AC013, April 1978. xi + 103pp. Available from: http://www.sealresearch.org/attachments/article/580/Wilson%201978%20Maine_pup_sociali sation_1976.pdf

10. Boness DJ, Bowen WD, Oftedal OT. Evidence of a maternal foraging cycle resembling that

of otariid seals in a small phocid, the harbour seal. Behav Ecol Sociobiol. 1994; 34: 95–104.

11. Wilson SC, Johnston T, Corpe HM. Radiotelemetry study of four rehabilitated harbour seal

pups following their release in County Down, Northern Ireland. J Wildlife Rehabil. 1999; 22:

17–23.

12. MacRae AM, Haulena M, Fraser D. The effect of diet and feeding level on survival and

weight gain of hand-raised harbour seal pups (*Phoca vitulina*). Zoo Biol. 2011; 30: 532–541.

13. Robinson I. The rehabilitation of seals at the RSPCA Norfolk Wildlife Hospital. J British Vet

Zool Soc. 1995; 1:13-17.

14. Wilson SC (1999) Radiotelemetry study of two rehabilitated harbour seal pups released

close to the time of weaning in the wild. J Wildlife Rehabil. 1999; 22: 5–11.

15. Larmour LJ (1989) Hand-rearing and rehabilitation of common seal pups *Phoca vitulina* at

the Oban Sea Life Centre. Int Zoo Yearb. 1989; 28: 272–279. doi.org/10.1111/j.1748-

1090.1989.tb03299.x

16. Osinga N, Hart P 't. Harbour seals (*Phoca vitulina*) and rehabilitation. NAMMCO Sci Publ.

2010; 8: 355–372.

17. Müller G, Gröters S, Siebert U, Rosenberger T, Driver J, König M et al. Parapox infection in

harbour seals (*Phoca vitulina*) from the German North Sea. Vet Pathol. 2003; 40: 445–454.

18. Sapolsky RM, Romero LM, Munck AU. How do glucocorticoids influence stress responses?

Integrating permissive, suppressive, stimulatory and preparative actions. Endocr Rev. 2000;

21: 55–89.

19. Atkinson S, Crocker D, Houser D, Mashburn K. Stress physiology in marine mammals: how

well do they fit the terrestrial model? J Comp Physiol B. 2015. doi 10.1007/s00360-015-

0901-0.

20. Cook NJ. Review: Minimally invasive sampling media and the measurement of

corticosteroids as biomarkers of stress in animals. Can J Anim Sci. 2012; 92: 227–259.

21. Rushen J. Problems associated with the interpretation of physiological data in the assessment

of animal welfare. Appl Anim Behav Sci. 1991; 28: 381–386.

22. Menargues A, Urios V, Limiñand R. Seasonal pattern of salivary cortisol secretion in the

greater one-horned rhino (*Rhinoceros unicornis*). Anim Welf. 2013; 22: 467–472. doi:

10.7120/09627286.22.4.467

23. Trumble SJ, O'Neill DO, Cornick LA, Gulland FMD, Castellini MA, Atkinson S. Endocrine

changes in harbour seal (*Phoca vitulina*) pups undergoing rehabilitation. Zoo Biol. 2013; 32:

134–141.

24. Constable S, Parslow A, Dutton G, Rogers T, Hogg C. Urinary cortisol sampling: a non-

invasive technique for examining cortisol concentrations in the Weddell seal, *Leptonychotes*

weddelli. Zoo Biol. 2006; 25: 137–144.

25. Otovic P, Hutchinson E. Limits to using HPA axis activity as an indication of animal welfare.

ALTEX 32: 41–50. DOI: 10.14573/altex.1406161

26. Koren L, Ng ESM, Soma KK, Wynne-Edwards KE. Sample preparation and liquid

chromatography –tandem mass spectrometry for multiple steroids in mammalian and avian

circulation. PLoS ONE 2012; 7(2): e32496. doi:10.1371/journal.pone.0032496.

27. Gardiner KJ, Hall AJ. Diel and annual variation in plasma cortisol concentrations among

wild and captive harbour seals (*Phoca vitulina*). Can J Zool. 1997; 75: 1773–1779.

28. McWhinney BC, Briscoe SE, Ungerer JPJ, Pretorius CJ. Measurement of cortisol, cortisone,

prednisolone, dexamethasone and 11-deoxycortisol with ultra high performance liquid

chromatography–tandem mass spectrometry: application for plasma, plasma ultrafiltrate,

urine and saliva in a routine laboratory. J Chromatogr B. 2010; 878: 2863–2869.

29. De Clercq N, Van den Bussche J, Croubels S, De la Haut P, Van Haecke L. A validated

analytical method to study the long-term stability of natural and synthetic glucocorticoids in

livestock urine using ultra-high performance liquid chromatography coupled to Orbitrap-high

resolution mass spectrometry. J Chromatogr A. 2013; 1301: 111–121.

30. Taussky HH, Kurzmann R. A microcolorimetric determination of creatinine in urine by the

Jaffe reaction. J Biol Chem. 1954; 208: 853–862.

31. Keenan JB. Metabolomic approaches to assessing pre-slaughter stress in relation to beef

quality. PhD Thesis, University of Reading, UK. 2015. Available from: uk.bl.ethos.679250.

32. R Core Team. 2016. R: a language and environment for statistical computing. R Foundation

for Statistical Computing, Vienna, Austria.

33. Miller MW, Thompson Hobbs N, Sousa MC. Detecting stress in Rocky Mountain bighorn

sheep (*Ovis canadensis Canadensis*): reliability of cortisol concentrations in urine and feces.

Can J Zool. 1991; 69: 15–24. doi: 10.1139/z91-003.

34. Krasowski MD, Drees D, Morris CS, Maakestad J, Blau JL, Ekins S. Cross-reactivity of

steroid hormone immunoassays: clinical significance and two-dimensional molecular

similarity prediction.   Clin Pathol. 2014; 14: 33. doi:10.1186/1472-6890-14-33.

35. Taylor AE, Keevil B, Huhtaniemi IT. Mass spectrometry and immunoassay: how to measure

steroid hormones today and tomorrow. Eur J Endocrinol. 2015; 173: D1–D12. www.eje-

online.org.

36. Berman DG, Johnson DE, Phillips RW, Barry BP. Physiological and urinary metabolite

responses to cold shock and confinement of sheep. J Anim Sci. 1980; 50: 713–722.

37. Lupo S, Carroll F, Liang SH, Alagandula R, Kahler T. Analysis of urinary free cortisol and

cortisone by LC-MS/MS. 2018. Available from:

http://www.restek.com/pdfs/LC_28_Cortisol_Cortisone_Poster_2018.pdf

38. Bush IE, Mahesh VB. Metabolism of 11-oxygenated steroids. Biochem J. 1959; 71: 718–

742.

39. Jenkins JS, Sampson PA. Conversion of cortisone to cortisol and prednisone to prednisolone.

BMJ. 1967; 2: 205–207.

40. Park S, Kim Y, Pyo H, Park J. Analysis of corticosteroids in urine by HPLC and thermospray

LC/MS. J Anal Toxicol. 1990; 14: 102–108.

41. Chiesa L, Pavone S, Pasquale E, Pavlovic R, Panseri S., Valiani A et al. Study on cortisol,

cortisone and prednisolone presence in urine of Chianina cattle breed. J Anim Physiol Anim

Nutr (Berl). 2017; 101: 893–903.

42. Fidan M, Gamberini, MC, Pompa G, Mungiguerra F, Casati A, Arioli, F. Presence of

endogenous prednisolone in human urine. Steroids 2013; 78: 121–126.

43. Famele M, Ferranti C, Palleschi L, Abenavoli C, Fidente RM, Pezzolato M et al.

Quantification of natural and synthetic glucocorticoids in calf urine following different

growth-promoting prednisolone treatments. Steroids 2015; 104: 196–202.

44. Arioli F, Casati A, Fidani M, Silvestri M, Pompa G. Prednisolone and prednisone neo-

formation in bovine urine after sampling. Animal 2012; 6: 1023–1029.

doi:10.1017/S1751731111002497.

45. Vincenti M, Leporati M, Capra P, Gatto S, Attucci A, Barbarino G et al. A field survey on

the presence of prednisolone and prednisone in urine samples from untreated cows. Food

Addit Contam: Part A. 2013; 29: 1893–1900.

46. Friel PN, Alexander T, Wright JV. Suppression of adrenal function by low-dose prednisone:

assessment with 24-hour urinary steroid hormone profiles – a review of five cases. Altern

Med Rev. 2006; 11: 40–46.

47. Biben M, Chanpoux M. Play and stress: cortisol as a negative correlate of play in *Saimiri*.

Play Culture Studies 1999; 2: 191–208.

48. Fagen R, Fagen J. Juvenile survival and benefits of play behaviour in brown bears, *Ursus*

arctos. Evol Ecol Res. 2004; 6: 89–102.

49. Fernald LC, Grantham-McGregor SM. Stress response in school-age children who have been

growth retarded since early childhood. Am J Clin Nutr. 1998; 68: 691–8.

50. Engelhardt FR, Ferguson JM. Adaptive hormone changes in harp seals, *Phoca groenlandica*,

and gray seals, *Halichoerus grypus*, during the postnatal period. Gen Comp Endocrinol.

1980; 40: 434–445.

51. Bennett KA, Fedak MA, Moss SEW, Pomeroy PP, Speakman JR, Hall AJ (2013) The role

of glucocorticoids in naturally fasting grey seal (*Halichoerus grypus*) pups: dexamethasone

stimulates mass loss and protein utilisation, but not departure from the colony. J Exp Biol.

2013; 216: 984–991.

52. Wielebnowski N. Stress and distress: evaluating their impact for the well-being of zoo

animals. J Am Vet Med Assoc. 2003; 223: 973–977.

53. Linklater WL, Macdonald EA, Flamand JRB, Czekala NM. Declining and low fecal

corticoids are associated with distress, not acclimation to stress, during the translocation of

African rhinoceros. Anim Conserv. 2010; 13: 104–111. Doi:10.1111/j.1469-

1795.2009.00308.x

54. Perry EA, Renouf, D. Further studies of the role of harbour seal (*Phoca vitulina*) pup

vocalizations in preventing separation of mother-pup pairs. Can J Zool. 1988; 66: 934–938.

55. Di Poi C, Atkinson S, Hoover-Miller A, Blundell G. Maternal buffering of stress response in

free-ranging Pacific harbour seal pups in Alaska. Mar Mamm Sci. 2015; 31: 1098–1117.

56. Yusko B, Hawk K, Schiml PA, Deak T, Hennessy MB. Sensitization of depressive-like

behaviour during repeated maternal separation is associated with more-rapid increase in core

body temperature and reduced plasma cortisol levels. Physiol Behav. 2012; 105: 861–867.

57. Spencer-Booth Y, Hinde RA. Effects of brief separations from mothers during infancy on

behaviour of rhesus monkeys 6–24 months later. J Child Psychol Psychiatry 1971; 12: 157–

172.

58. Perkeybile AM, Schiml-Webb PA, O'Brien E, Deak T, Hennessy MB. Anti-inflammatory

influences on behavioural, but not cortisol, responses during maternal separation.

Psychoneuroendocrinology 2009; 34: 1101–1108.

59. Laane RWPM, Vethaak AD, Gandrass J, Vorkamp K, Koehler A, Larsen MM et al.

Chemical contaminants in the Wadden Sea: Sources, transport, fate and effects. J Sea Res.

2013; 82: 10–53.

60. Wang D, Atkinson S, Hoover-Miller A, Shelver WL, Li QL. Organic halogenated

contaminants in mother-fetus pairs of harbour seals (*Phoca vitulina richardii*) from Alaska,

2000–2002. J Hazard Mater. 2012; 223–224C: 72–78. doi:10.1016/j.jhazmat.2012.04.052.

61. Wilson J, Berntsen HF, Zimmer KE, Verhaegen S, Frizzell C, Ropstad E et al. Do persistent

organic pollutants interact with the stress response? Individual compounds, and their

mixtures, interaction with the glucocorticoid receptor. Toxicol Lett. 2016; 241: 121–132.

62. Beschøft TO, Sonne C, Dietz R, Born EW, Muir DCG, Letcher RJ et al. Associations

between complex OHC mixtures and thyroid and cortisol hormones in East Greenland polar

bears. Environ Res. 2012; 116: 26–35. doi:10.1016/j.envres.2012.04.010.

63. Economu G, Andronikou S, Choleras V, Lapatsanis P. Cortisol secretion in stressed babies

during the neonatal period. Horm Res. 1993; 40: 217–221.

64. Graham YP, Heim C, Goodman SH, Miller AH, Nemeroff CB. The effects of neonatal stress

on brain development: implications for psychopathology. Dev Psychopathol. 1999; 11: 545–

565.

65. Muelbert MMC, Bowen WD, Iverson SJ. Weaning mass affects changes in body

composition and food intake in harbour seal pups during the first month of independence.

Physiol Biochem Zool 2003; 76: 418–427.

66. Romero T, Nagasawa M, Mogi K, Hasegawa T, Kikusu T. Oxytocin promotes social bonding

in dogs. Proc Natl Acad Sci USA 2014; 111: 9085–9090. Doi/10.1073/pnas.1322868111.

67. Crockford C, Wittig RM, Langergraber K, Ziegler TE, Zuberbühler K, Dreschner T. Urinary

oxytocin and social bonding in related and unrelated wild chimpanzees. Proc R Soc B 2013;

280: 20122765. dx.doi.org/10.1098/rspb.2012.2765.

Supporting information

S1_pup_rehab_environment_photos
S2_dataset
S3_ELISA_protocol
S4_MS_protocol
S5_pup_growth_rates_centre_II

Figure

Figure

Figure

Figure

Figure

Figure

Figure