Abundance estimation from genetic mark-recapture data when not all sites are sampled: an example with the bowhead whale

Timothy R. Frasier[a], Stephen D. Petersen[b], Lianne Postma[c], Lucy Johnson[b], Mads Peter Heide-Jørgensen[d], Steven H. Ferguson[c]

[a] Department of Biology, Saint Mary's University, 923 Robie St., Halifax, NS, B3H 3C3 Canada

[b] Conservation and Research Department, Assiniboine Park Zoo, 2595 Roblin Blvd., Winnipeg Manitoba, R3R 0B8, Canada

[c] Fisheries and Oceans Canada, 501 University Crescent, Winnipeg, Manitoba, R3T 2N6, Canada

[d] Greenland Institute of Natural Resources, Box 570, 3900 Nuuk, Greenland

Corresponding Author:

Timothy R. Frasier

Saint Mary's University

Department of Biology

923 Robie Street

Halifax, NS B3H 3C3

Canada

Email: timothy.frasier@smu.ca

Tel: (902) 491-6382

Abstract

Estimating abundance is one of the most fundamental and important aspects of population

biology, with major implications on how the status of a population is perceived and thus on

conservation and management efforts. Although typically based on one of two methods

(distance sampling or mark-recapture), there are many individual identification methods

that can be used for mark-recapture purposes. In recent years, the use of genetic data for

individual identification and abundance estimation through mark-recapture analyses have

increased, and in some situations such genetic identifications are more efficient than their

field-based counterparts for population monitoring. One issue with mark-recapture

analyses, regardless of which method of individual identification is used, is that the study

area must provide adequate opportunities for "capturing" all individuals within a

population. However, many populations are unevenly and widely distributed, making it

unfeasible to adequately sample all necessary areas. Here we develop an analytical

technique that accounts for unsampled locations, and provides a means to infer "missing"

individuals from unsampled locations, and therefore obtain more accurate abundance

estimates when it is not possible to sample all sites. This method is validated using

simulations and is used to estimate abundance of the Eastern Canada-West Greenland

(EC-WG) bowhead whale population. Based on these analyses, the estimated size of this

population is 11,747 individuals, with a 95% highest density interval of 8,169-20,043.

Keywords: abundance estimation, mark-recapture, genetic identification, bowhead whale, whale,

genetic mark-recapture

1. Introduction

Mark-recapture analyses have become an invaluable tool not only as a means for abundance estimation, but also for obtaining a wealth of other information on population dynamics, such as population growth rates, life history, and survival probabilities (Kraus et al. 2001; Williams et al. 2002; Amstrup et al. 2005). Traditionally, mark-recapture methods involved the actual capture, marking, and recapture of individuals (Rice and Harder 1977; Hestbeck et al. 1991; Pradel et al. 1997), but their use quickly expanded to species where individuals could be identified based on natural markings, and thus "capturing" in these cases meant taking a picture that could be used for identification (Wilson et al. 1999; Matthews et al. 2001; Langtimm et al. 2004; Bradshaw et al. 2007). More recently, the increasing ease with which individual-specific genetic profiles can be obtained from small sources of DNA has made it possible (and sometimes more efficient) to use genetic data as the primary means of individual identification, and thus the basis for mark-recapture analyses (Mowat and Strobeck 2000; Miller et al. 2005; Petit and Valiere 2006).

One important requirement in mark-recapture studies, regardless of which method of identification is used, is ensuring that the study area provides adequate opportunities for "capturing" all individuals within the study population. This requirement may be easily addressed for populations with small ranges that can be surveyed in their entirety, or where all individuals utilize—or pass through—a single area. However, many populations are not so amenable, with individuals exhibiting substantial variation in site fidelity and habitat use patterns. Failure to account for this variation can bias abundance estimates either upwards or downwards. The former can occur if marked individuals move out of the study area and thus become unavailable for recapture. The latter can occur if the study area is only used by a portion

of the population, and thus estimates are only representative of some individuals rather than the

entire population.

For species with wide distributions and substantial heterogeneity in habitat use patterns it

may not be possible to obtain adequate coverage of all areas. In such cases abundance estimation

requires appropriate corrections during the analytical stages of the study. In 2005, Durban et al.

(2005) made substantial progress in this area by recognizing that analytically accounting for

missed sites is essentially a regression problem, where data from sampled areas can be used to

make predictions on the number of individuals not seen in any surveyed area. As an example,

suppose a study population of organisms where some, but not all, individuals can be identified

based on natural markings. Suppose the number of identifiable individuals has been counted at

each of three sites, but other unstudied sites exist. To estimate the total size of the identifiable

(marked) population, Durban et al. (2005) first organized the sighting histories of identifiable

individuals based on whether or not they were observed in each of the three surveyed areas (see

Table 1). The last row of such a table represents the identifiable individuals not seen in *any*

location, and can be inferred from the available data. In this case a regression model is fit with

the first seven rows of data, and the coefficient estimates from this process are then used to

predict the value for the last row. In their application of this approach to a study of bottlenose

dolphins (*Tursiops truncatus*) around Scotland, Durban et al. (2005) found that their approach

performed quite well and resulted in a similar estimate to that obtained through another study

based on a more rigorous and standardized survey design.

The eastern Canada-western Greenland (EC-WG) bowhead whale (*Balaena mysticetus*,

Fig. 1) population epitomizes the confounding factors associated with abundance estimation,

using either distance sampling or mark-recapture methods. First, it is an arctic-adapted

population with a wide distribution often associated with the ice edge (**Fig. 2**) (Reeves et al.

1983; Moore and Reeves 1993; Finley 2001; Ferguson et al. 2010). The necessary logistics and

monetary requirements make extensive surveys in the high arctic challenging. Second, the

population is highly segregated based on sex, age class, and perhaps other factors (Reeves et al.

1983; Finley 1990; Moore and Reeves 1993; Heide-Jørgensen et al. 2010). Indeed, the

segregation is so great that, until recently, multiple populations (and subsequent management

stocks) were thought to exist (Rugh et al. 2003). It was only when satellite tagging data recently

showed movement of whales across previously assumed population boundaries, and such

movements were also identified based on genetic mark-recapture analyses, that the presence of

only one population was recognized (Heide-Jørgensen et al. 2003, 2006; Wiig et al. 2011). These

confounding factors have led to very imprecise abundance estimates, as well as large differences

in estimates based on data from different locations and/or different analytical approaches, as

summarized in Koski et al. (2013).

Here, we extend the approach of Durban et al. (2005) in an important way. Specifically,

The model of Durban et al. (2005) was designed to estimate the number of missing *marked*

individuals in a population. However, our goal was to estimate the *total number* of missing

individuals (marked and unmarked), and combine this information with data from sampled sites

to result in one estimate of total abundance for the population of interest. If successful, this

approach could prove useful to a broad range of population studies. We then applied this method

to genetic mark-recapture data for the eastern Arctic-western Greenland bowhead whale

population. Genetic mark-recapture methods may be more efficient than photograph-based

identifications for this population because samples can be collected in collaboration with Arctic

communities, eliminating the need for extensive and expensive aerial surveys while also

fostering closer collaborations and relationships with Arctic communities.

2. Methods

2.1 Sampling

Intensive sampling effort has taken place out of the communities of Igloolik and Pangnirtung in

Canada, and in Disko Bay, Greenland (hereafter "Greenland") (**Fig. 2**, **Table S1**). Additionally,

more opportunistic sample collection has occurred in Repulse Bay and Arctic Bay, as well as

from 11 subsistence hunts by Canadian Inuit (Taloyoak, Kugaaruk, Cape Dorset, Pangnirtung,

Repulse Bay, Igloolik) (**Fig. 2**, **Table S1**). In total, 1,177 samples have been collected across 19

years.

In each location, small tissue (skin and blubber) samples were obtained from free-

swimming whales using a 150-lb draw-weight crossbow (Excalibur Vixen) with bolts and tips

modified specifically for this purpose (from Ceta-DART, Denmark). This is the most common

method for collecting small skin samples from free-swimming whales, and has no negative short

or long-term effects on individuals other than an initial "startle" response in some cases (Best et

al. 2005; Noren and Mocklin 2011; Kowarski et al. 2014). Biopsy samples were collected in

association with photo-identification data, as part of ongoing photo-identification studies in all

areas. Samples were preserved in either a salt-saturated 20% DMSO solution (Seutin et al. 1991),

flash frozen in liquid nitrogen, in RNAlater (Qiagen), or in Allprotect (Qiagen) and frozen upon

arrival in the laboratory. All samples were archived in -80ºC freezers at the Freshwater Institute

(DFO, Canada, Winnipeg) laboratories until further analysis. Bowhead whale biopsy samples

were collected under permit Department of Fisheries and Oceans License to Fish for Scientific

Purposes S-16/17 1005-NU and Animal Use Protocol FWI-ACC-2016-09.

2.2 Genetic Analyses

DNA was extracted using a modified version of the standard Qiagen extraction protocol (Wang

et al. 2008). DNA was quantified using a μQuant spectrophotometer (Bio-Tek) or a NanoDrop

spectrophotemeter (ThermoFisher Scientific) and normalized to 10–100 ng/μl. Sex was

determined through the amplification of a zinc finger gene intron using primers LGL331 and

LGL335 (Shaw et al. 2003). PCR fragments were size-separated in 1.5% agarose gels and

visualized with GelRed (Biotium). The resulting banding pattern of X (~975 bp) and Y (~1040

bp) fragments were used to infer sex.

Multiplex reactions were developed and used to amplify 23 microsatellite loci (**Table

S2**). The reaction mixtures contained 1.0 μl of template DNA, 1X AmpliTaq Gold Buffer, 2.0

mM MgCl$_2$, 0.25 mM each dNTP, 0.5 U AmpliTaq Gold DNA polymerase (Life Technologies),

and the primer concentrations indicated in **Table S2**. Thermal cycler profiles included a five

minute initial denaturing step at 95ºC followed by 35 cycles of: 95ºC for 30 seconds, annealing

temperature (**Table S2**) for 30 seconds, and 72ºC for 30 seconds; followed by a final extension

step of 72ºC for 30 minutes. PCR products were size-separated and visualized on an Applied

Biosystems 3130xl Genetic Analyzer (Life Technologies), with GeneScan 600 LIZ (Life

Technologies) as a size-standard. Size-standards were checked, and allele calls made using the

program GeneMarker (SoftGenetics).

2.3 Identifying Recaptures

The crux in identifying genetic recaptures is developing appropriate criteria for defining when

two genotypes can be considered the same, and thus representing the same individual. This

problem has a long history, and has been most thoroughly dealt with in human forensic DNA

typing (National Research Council 1996). The issues are two-fold. First, two genotypes may be

the same at the typed loci but represent different individuals if not enough resolution is available

with the chosen markers, if individuals are related, or just by chance (a false inclusion). Second,

two genotypes may be from the same individual but be scored differently due to genotyping

errors or null alleles (a false exclusion). Thus, the key is identifying what criteria minimize the

chances of each of these errors.

To identify recaptures, or samples likely originating from the same individual, we used

the R package ALLELEMATCH (Galpern et al. 2012). This package deals with these issues by

calculating the genetic similarity of all genotypes, and then creating clusters based on genetic

similarity. It then re-creates these clusters by sequentially allowing an incremental number of

mismatches, while keeping track of how many "unique" genotypes are present under each

number of allowed mismatches. The appropriate number of allowed mismatches is then

identified as the point when each genotype associates with only one cluster.

2.4 Estimating Abundance

We used Bayesian methods to estimate abundance from the genetic mark-recapture data, based

on the general approaches described in Williams et al. (2002) and Kéry and Schaub (2012). We

tried two general strategies, each with their own strengths and weaknesses. First, the main goal

of this study was to develop a method to use data from sampled locations to explicitly infer the

number of individuals that also exist in unsampled locations, and combine these to obtain an

estimate for overall abundance. We call this the "location-specific" approach, which was

modelled after that used in Durban et al. (2005). The strength of this approach is that it provides

an explicit and quantitative means to infer the number of individuals using unsampled locations.

The main weakness is that it requires dividing the data by locations, with the consequence that

many locations may not have enough samples to be included in the analyses, and thus some

information may be lost. The second and most simple approach involved ignoring the location

information and treating the data as one large capture-mark-recapture study (hereafter called the

"location-independent" approach). The main benefit of this approach is that it allows all samples

to be used, regardless of whether recaptures were found within each location. The main

weakness of this approach is that it does not explicitly account for the fact that not all locations

have been sampled, and therefore makes the assumption that the sampled locations represent

adequate sampling opportunities for all individuals. In this case, the location-specific analyses

serve as the focus of our work, whereas the location-independent analyses serve as references

against which the location-specific estimates can be compared. Both of these methods are

described in more detail below.

Again, the main goal of this study was to develop and test an approach that allows for the

inference of "missing" individuals when not all sites are sampled. For this approach, we first

needed abundance estimates for each sampled location. At this stage, we chose to do this using

closed population models for a number of reasons, with the realization that incorporating this

approach into open population models is desirable in the future. First, the simplest and clearest

way to incorporate our approach, and test its performance, was with closed population models.

We therefore felt that this simplest approach was the best way to initialize progress into inferring

missing individuals, with the recognition that incorporating it into more complex and realistic

models in the future will be required. Second, sampling effort varied widely across locations, in

terms of the number of years samples were collected in each location (**Table S1**), as well as the

effort put into collecting samples, violating assumptions of open multistate models (Williams et

al. 2002). Third, our method accounts for movement among locations, including unsampled

ones; given that estimating movement rates (but not explicitly accounting for unsampled

locations) is a key aspect of open population models, it is not immediately clear how these could

be combined. Therefore, we view this work as a starting point rather than an end point, where

future effort can be directed towards incorporating our approach into more complex (and

realistic) open population models. Additionally, the long life-span and generally slow life history

of bowhead whales mean that it is unlikely that there were dramatic changes to population size

(via births, immigration, deaths, or emigration) throughout the study period (Nerini et al. 1984;

George et al. 1999; Zeh et al. 2002; George and Bockstoce 2008). However, to try to account for

any potential changes over time, for each strategy (location-independent and location-specific)

we estimated population sizes based on two data sets: the full data set, and a subset containing

only five years of data. The rationale is that the population may have changed throughout the 19

years through which the samples were collected. Therefore, closed-population abundance

estimates based on the entire study period may be biased. The 5-year data set contains fewer data

but is also expected to be less biased by changes in population size within the sampling period.

Combined, this resulted in four estimates of abundance based on: (1) location-specific full data

set (LS-FD); (2) location-specific 5-year data set (LS-5Y); (3) location-independent full data set

(LI-FD); and (4) location-idependent 5-year data set (LI-5Y).

For all analyses, the recaptures identified by ALLELEMATCH were filtered to exclude

recaptures from the same location in the same year. Bowhead whales are known to vary their

migration and habitat use patterns depending on age, sex, and life-history stage (Moore and

Reeves 1993; Heide-Jørgensen et al. 2010). Additionally, once in an area, whales are likely to

remain there for some period of residency (e.g., Finley 1990). Thus, we thought that within-year

recaptures within locations would be artificially high and would not be representative of general

capture probabilities for the entire population throughout the study period.

2.4.1 Location-Specific Full Data Set (LS-FD)

Our location-specific approach for estimating abundance was modeled after that used by Durban

et al. (2005). The general idea is that location-specific abundance estimates can be used in

association with estimated movement rates between locations to identify (or infer) the number of

individuals with different sighting histories such as those shown in **Table 1**. When some areas

are unsampled, the goal is to estimate the number of individuals in the last row (who were not

captured in any area). This abundance estimate can then be used in association with estimates of

movement rates to obtain an estimate of total abundance even when some areas remain

unsampled. The entire process is summarized in **Fig. 3**, with details described below. There may

be more than one unsampled location (i.e., it need not represent a single actual location, but

rather could be a number of unsampled locations); however, these will be lumped into a single

estimate of "missing" individuals (across all unsampled locations) using this approach. This is

because the criterion for being counted as "missing" is that individuals are not present in any of

the sampled locations, and therefore the number of unsampled locations need not be defined.

The first step in this process was to estimate the movement rates between all pairs of sampled

locations. This is based on the number of recaptures within each considered location, and the

number of those recaptures that represent individuals originally marked in the alternative

location. We made the assumption that pairwise movement rates were symmetrical, largely

because the number of cross-location recaptures was very low (see Results), and therefore there

were few data available from which to estimate movement rates. However, it is possible to relax

this assumption in the future, which would just require the additional estimation of direction-

specific movement rates for each pair of locations. The number of recaptures within and between

each pair of locations was converted into a string of zeros and ones, where ones represented

recaptures across locations. The posterior probabilities for the movement rates were then

estimated using a hierarchical Bayesian approach with a Bernoulli likelihood function and a beta

prior with a uniform (flat) distribution (the code itself, as well as a step-by-step walk-through of

all analyses are available in the Supplementary Material). The result is a complete posterior

probability distribution for movement rates between all pairs of sampled locations, as well as an

overall posterior probability distribution for movement rates across all sites.

For the second step of the process, the probability distribution for the symmetrical

movement rate to and from the unsampled location(s) was estimated as the overall posterior

distribution from the hierarchical model described above across all pairs of locations. This

approach makes the implicit assumption that the rate at which individuals move among the

sampled locations is at least somewhat representative of the rate at which individuals move

between sampled and unsampled locations.

For the third step, abundance was estimated for each sampled location. The genetic

recapture data from ALLELEMATCH were first converted to typical sighting histories consisting of

ones and zeros for each sampling period where individuals were, or were not, captured,

respectively. However, these sighting histories are biased due to individuals moving to and from

the unsampled location(s). Individuals marked in the sampled location could move to an

unsampled location, and therefore not be available for recapture, artificially reducing the number

of recaptures. In the opposite direction, unmarked individuals from the unsampled location(s)

could move into the sampled location, which would also artificially reduce the proportion of

recaptures. Both of these processes reduce the proportion of individuals available for recapture,

and would therefore bias population size estimates upwards. To correct for this, for each sampled

location we estimated the number of individuals moving to and from the unsampled location(s)

and randomly add these to the sighting history data. For example, suppose at one location there

were 300 marked individuals and 68 recaptures. Suppose that the estimated movement rate to

and from the unsampled location(s) is 0.12. This means that we have "lost" 12% of marked

individuals to unsampled locations, and our number of unmarked individuals is also artificially

inflated by 12% due to movement of unmarked individuals from unsampled areas into this area.

This means that our proportion of recaptures is underestimated by two times the movement rate

(24%). Thus, our original 68 recaptures only represent (1 - 0.24 = 0.76) 76% of the recaptures we

"should" have had, if individuals did not move to and from unsampled locations (i.e., we should

have had 68 / 0.76 = 89 recaptures). To account for this, these (89 - 68 = 21) additional

recaptures were randomly added to the sighting histories for this location.

Once the sighting histories were corrected, abundance was estimated for each sampled

location using the same closed population approach. Note that although the model assumed a

closed population, movement to and from other locations has already been taken into account,

and therefore "closed" in this sense just refers to births and deaths. Specifically, to estimate

abundance for each location the sighting histories were augmented by 10,000-15,000 (based on

trials) to ensure that considered probabilities covered a wide enough range to ensure adequate

sampling of the posterior distribution (Tanner and Wong 1987; Royle et al. 2007). This

augmenting strategy was introduced by Royle et al. (2007) to deal with the fact that Bayesian

estimation of N has an unbounded upper end, which can make the efficient exploration of

parameter space difficult for many Markov Chain Monte Carlo samplers. Instead, they proposed

augmenting data sets by adding a large number of "uncaptured individuals" (individuals with all

zeroes in their sighting history), to the observed sighting histories in a manner so that the

augmented data set $\gg N$. This converts the problem from estimating an unbounded N directly

into one of estimating what proportion of the augmented data set should be included in the "true"

population size (N). The parameters of interest are the sighting probability (p), and the inclusion

probability (Ω), which is the probability that a member of the augmented data set is part of the

"true" population size (N). Abundance estimates were then obtained for a closed population

model (model M_0 from Otis et al. (1978)) using the approach described in Royle et al. (2007) and

Kéry and Schaub (2012). The priors for both the sighting probability (p) and the inclusion

probability (Ω) were uniform distributions ranging from 0 to 1. All Bayesian analyses were

conducted using a combination of R v.3.5.0 (R Core Team 2015), rstan v.2.18.2 (Stan

Development Team 2018), and Stan v.2.17.0 (Carpenter et al. 2017). The performance of models

for all analyses were assessed based on examination of effective sample sizes, trace plots, and

Rhat calculations. Specifically, for all analyses of all scenarios, all Rhat values were 1, all

effective sample sizes were greater than 4,000, and all traceplots showed that the chains quickly

converged onto one value (and the same value across all chains), and then mildly fluctuated

around that value for the vast majority of steps (except the very first few).

For step four, the posterior probabilities for abundance for each sampled location, and for

each pairwise movement rate, were used to estimate the number of individuals with each

possible habitat use pattern across the sampled locations (there are 3 such possible patterns for 2

locations – **Fig. 3**, 7 possible patterns for 3 locations, and so on). These data were then used in a

Bayesian regression analysis to estimate the relevant regression coefficients (see equation

below). In this case, y is a vector of integers containing the number of individuals with each

sighting history, and x_1 through x_4 are vectors of 1s and 0s indicating if each location (locations 1

through 4) is (1) or is not (0) part of each sighting history, respectively. Once estimates for the

coefficients (β_0 through β_4) have been obtained, these are used to predict the number of

individuals that are not seen in any of the sampled locations (i.e., where values for all x_1 though

x_4 are zero: the last row of the table shown in **Fig. 3**, step 4). The resulting posterior probability

distribution represents the estimate of the total number of individuals not seen in any sampled

location.

$$y \sim \beta_0 + \beta_1 x_1 + \beta_2 x_2 + \beta_3 x_3 + \beta_4 x_4$$

The estimated number of "missing" individuals from the previous step cannot just be

added to the estimates for each independent location to obtain an overall abundance estimate.

This is because the same individuals may use multiple locations, including a mixture of

unsampled and sampled locations (the estimate above is just an estimate of the individuals that

never use sampled locations, and will therefore often be an underestimate). Therefore, for the

fifth and final step, total abundance was estimated in a similar manner as that for the "missing"

individuals. Specifically, the complete posterior probabilities for abundance for each location,

and for each pairwise movement rate, were used to estimate the number of individuals with each

possible habitat use pattern across locations. The difference is that now this table includes data

for the unsampled location(s): effectively adding another location to the table and associated

movement rates (see **Fig. 3**). The sum of these values then represents the abundance estimate for

the total population. Note that one benefit of this approach is that the uncertainty in each estimate

is carried over into the uncertainty of the total abundance estimate. Put another way, rather than

estimating the total abundance based on the mean, mode, or some other summary metric of the

other parameters, this estimate is based on the entire probability distribution for each parameter.

In this way the uncertainty in each estimate of abundance and movement are incorporated into

the posterior probability distribution for the total abundance estimate. This is one of the attractive

properties of setting up the analyses in this way.

2.4.2 Location-Specific 5-Year Data Set (LS-5Y)

We conducted the same analyses as above (for the location-specific full data set, LS-FD), except

this time with a data set containing only those sampling events in the 5-year period from 2008-

2012. Again, the idea for this process was to reduce the potential biases in the longer-term data

set due to births and deaths occurring within the longer time frame, by analyzing data from a

much shorter time period (note that immigration and emigration are dealt with explicitly in the

model).

2.4.3 Location-Independent Full Data Set (LI-FD)

To estimate abundance for the entire data set, the recapture data from ALLELEMATCH, combining

data across all locations, were first converted to typical sighting histories consisting of ones and

zeros for each sampling period (year) where individuals were, or were not, sampled,

respectively. These sighting histories were augmented by 30,000 to ensure that considered

probabilities covered a wide enough range to ensure adequate sampling of the posterior

distribution. An estimate of abundance was then obtained using the same approach as described

for the closed population model aspect of the location-specific data set, where estimates of the

sighting probability (p), inclusion probability (Ω), and abundance (N) were obtained.

2.4.4 Location-Independent 5-Year Data Set (LI-5Y)

As with the location-specific 5-year data set, we conducted the same analyses as described above

for the location-independent full data set, but this time limiting the data to the years 2008-2013.

Again, the rationale was to reduce the potential biases in the longer-term data set changes in

population size during the longer sampling time.

2.5 Model Validation

A number of simulations were conducted to assess model performance, and to provide

information that would be useful for guiding sampling strategies in the field. All simulations

were based on the location-specific model, which, instead of estimating abundance for a single

population/location, is designed to estimate abundance for each location independently

(including unsampled locations), and then come up with one total abundance estimate for all

sampled and unsampled locations.

The first set of simulations focused on assessing the effects of varying sampling effort.

These simulations were based on four sampled locations and one unsampled location to resemble

the available bowhead whale data. For these, all population sizes (including that of the

unsampled location) were set to 1,000 individuals (resulting in a total population size of 5,000

individuals). All movement rates between locations were also set to 10%. These initial values

were selected to be fairly similar to the expected values for the bowhead whales, and therefore to

provide information regarding how this approach would perform with our actual data set.

Additionally, all simulations were run with one capture period and one recapture period, to

resemble the bowhead whale data, where no individual was recaptured more than once.

Abundance was then estimated based on sampling 5%, 10%, 15%, 20%, and 25% of individuals

from each sampled location, to assess how changing sampling effort influences the precision and

accuracy of abundance estimates.

A second set of five simulations were also run and analyzed to ensure that the model

performs well even when there are large differences in population sizes and sampling effort

between locations. As with the first set, these consisted of four sampled locations and one

unsampled location. However, this time the abundance at each location (including the unsampled

one) was randomly drawn from a list of 20 values ranging from 500 to 10,00, in increments of

500. The movement rates were all set to 10%, and 250 individuals were sampled at each

location. This means that the proportion of individuals sampled could range from 2.5% (for a

population size of 10,000 individuals) up to 50% (for a population size of 500 individuals).

A third set of two simulations were conducted, but in this case both the population sizes

and movement rates were randomly generated. These were conducted to assess how the model

performs when there are substantial differences in abundance at different locations, as well as in

the movement rates between these locations. Specifically, abundance for each of the four

sampled and one unsampled locations were randomly drawn from a list ranging from 250 to

10,000 in increments of 250. Movement rates were similarly randomly drawn from a list ranging

from 0.1 to 0.30 in increments of 0.01. The proportion of sampled individuals at each of the

sampled locations was set at 0.20.

A fourth set of six simulations was conducted to assess model performance if the

movement rate and/or abundance at a single unsampled location differed substantially from those

of the sampled locations. For the first scenario the abundance at the four sampled and one

unsampled location were all set to 5,000 individuals, but the movement rates among the sampled

locations was set to 0.20, and those between the sampled locations and unsampled location were

set to 0.05. We then conducted a second simulation reversing this pattern, where the movement

rate among the sampled locations was low (0.05), and those between the sampled and unsampled

locations was high (0.20). The third simulation consisted of the same movement rates among

sampled and unsampled locations (0.15), but all sampled locations had 5,000 individuals, and the

unsampled location had just 500. For the fourth simulation this pattern was reversed, with all

sampled locations having 500 individuals and the unsampled location having just 500. The

movement rate was uniform across all sites at 0.15. For the last pair of simulations under this

scenario, both the abundance and movement rates differed between sampled and unsampled

locations. Specifically, in the fifth simulation the sampled locations all had 5,000 individuals and

the usampled location had 500. The movement rate among the sampled locations was high

(0.20), whereas the movement rate between the sampled and unsampled locations was much

lower (0.05). For the sixth simulation the population sizes were the same as above, but the

movement rates were reversed: the movement rate among sampled locations was 0.05, and that

between the sampled locations and the unsampled location were 0.20. Under all six of these

scenarios 15% of individuals were sampled from each sampled location.

To gain insight into how the model performs when there is more than one unsampled

location, a fifth set of two simulations was conducted and analyzed to assess model performance

when there is more than one unsampled location. Specifically, these simulations extended the

strategy of randomly generating abundance at each of four sampled locations, as well as at two

unsampled locations. The range of values for abundance at each location were randomly drawn

from a list ranging from 250 to 10,000 in increments of 250. Movement rates were randomly

drawn from a list ranging from 0.05 to 0.30 in increments of 0.01. The proportion of sampled

individuals at each of the sampled locations was set at 0.15.

3. Results

3.1 Samples, Genotypes, and Recaptures

A total of 1,177 samples were genotyped. **Table 2** shows how these were divided among

locations. The final microsatellite data set contained 21 loci since two loci were dropped due to a

significant portion of the samples missing data that were profiled before 2008. Furthermore, a

sample was only included in the analysis if it had at least 10 of the 21 loci scored. This provided

an overall probability of identity of 1.16×10^{-9}, which corresponds to a 1 in 8.62×10^{-8} chance that

the match identified was due to two random animals matching at all loci. **Table S1** shows how

these samples were divided by year, location, and sex.

The ALLELEMATCH analysis suggested that the best fit for the data was to allow 3

mismatches. Based on this criterion, 992 unique genotypes were identified, along with 185

recaptures. However, 136 of these recaptures represented whales recaptured within the same area

and year in which they were marked, and were therefore removed from the analyses. This left 49

recaptures for subsequent analyses. **Table 3** summarizes the recaptures by location. Based on the

number of recaptures, only Greenland, Igloolik, Pangnirtung, and Repulse Bay were used for

estimating abundance based on the full data set.

3.2 Model Validation

The results for the simulations testing the effects of sampling effort are shown in **Fig. 4**. The

precision of the estimates (or the range between to upper and lower 95% HDI values) declines

rapidly until sampling reaches ~15%. Above this sampling effort, precision increases at a much

slower rate. Thus, ~15% seems to be a turning point above which diminishing returns are

achieved as sampling effort increases. The details of all of the estimates from these simulations

are shown in **Fig. 5**. These plots show that the model performs quite well in all aspects

(abundance estimation of sampled locations, abundance estimation of unsampled location(s), and

total abundance).

The results from the simulations testing model performance when population size and

sampling effort varies across locations are shown in **Fig. 6**. The results show that the model

performs quite well under a range of variation in population sizes and sampling efforts.

Similarly, the simulations testing model performance when both the abundance and the

movement rates varied across all locations showed that the model performs well under these

scenarios and recovers accurate estimates of total abundance when there is one (**Fig. S1**), or

multiple (**Fig. S2**) unsampled locations.

However, performance declined when the abundance within the unsampled location(s)

and/or the movement rates to and from the unsampled location(s) fell well outside the range of

values among the sampled locations (**Fig. 7**, **Fig. S3**). This makes intuitive sense because in

these cases the information gleaned from the sampled locations (and therefore used to infer

information about the unsampled location(s)), is not representative, and therefore leads to

inaccurate estimates. For example, when the movement rates to and from the unsampled location

were very low (0.05), but those among the sampled locations were high (0.20), the abundance in

all locations were biased downwards (**Fig. 7a**). Abundance estimates for the sampled locations

were biased downwards because the recapture rates were over-corrected. Specifically, the model

assumed a higher movement rate to/from the unsampled location, resulting in an overcorrection

of recapture rates for individuals presumed to move to/from the unsampled locations, and

therefore resulting in lower abundance estimates. Abundance was underestimated for the

unsampled location for the same reason: a higher rate of movement was assumed, resulting in

higher estimates of individuals that would be "captured" in the sampled locations, and a lower

estimate of those going "missing". Similarly, when this scenario was reversed and the movement

rates to/from the unsampled location were high (0.20), and those among the sampled locations

were low (0.05), abundance was overestimated in most locations, and particularly for the total

population size (**Fig. 7b**). The rationale is the same as the first scenario, but now recaptures

within the sampled locations were under corrected, resulting in overestimates, as well as an

underestimation of the probability that individuals would move into sampled areas and be

"captured", resulting in higher estimates of total population size. Similarly, abundance estimates

were biased if the actual abundance in the unsampled location(s) were vastly outside of the range

of those within the sampled locations (**Fig. 7c-d**, **Fig. S3**). Again, this is expected because the

sampled locations are not providing appropriate information on which inferences about the

unsampled locations can be made.

3.3 Bowhead Whale Abundance Estimates

The location-specific abundance estimates for each location, inferred for the unsampled

location(s), and for the total population, are shown in **Table 4** (rows 1–6). The mode was used as

the point estimate instead of the mean in all cases because the posterior probability distributions

for all abundance estimates were skewed, with heavy right-hand tails, and therefore the mode

represented the estimate with a higher probability than the mean. Note that the estimates for each

location are inclusive, meaning that they represent all whales that use each area (and may also be

counted for other areas), rather than representing whales that only use each area. The location–

independent abundance estimates based are also shown in **Table 4** (row 7).

4. Discussion

Analyses of simulated data sets suggest that our new approach performs quite well, and could be

a useful tool for estimating abundance of other populations where only a subset of possible

locations have been sampled. Simulations testing analysis performance with respect to sampling

effort show that sampling roughly 15% of the individuals at each location is desirable, whereas

sampling more intensively will result in diminishing returns. The approach still performs well

when fewer than 15% of individuals are sampled at each location, but there is decreasing

precision around the estimates as sample size decreases (**Fig. 5**).

The approach also performs well even when there is wide variation in abundance

between sampling locations. Although the simulations used in our tests were not exhaustive, they

do show that accurate abundance estimates can be obtained for the total population when only a

subset of locations are sampled, and when there is wide variation in abundance across locations

(**Fig. 6**). Combined, the analysis of simulated scenarios suggests that this new approach may

indeed be a viable option for estimating abundance from genetic mark-recapture data when not

all locations are sampled.

Performance of our approach declined, however, when the data from the sampled

locations were not representative of those for the unsampled locations. Specifically, if the

abundance in the unsampled locations and/or the movement rates to and from the unsampled

locations are well outside the range of those among the sampled locations, then our model was

unable to obtain accurate estimates. This is due to the fact that the information obtained from the

sampled locations is not representative of that in the unsampled locations, which leads to

incorrect inferences. Therefore, using this approach carries the implicit assumption that the

abundance of, and movement rates to and from, unsampled locations are at least somewhat

within the range of those among the sampled locations. Although this seems like a major

drawback, it suggests that our method may be a useful approach for studies when abundance

estimates are desired, but it has not yet been possible to sample from all areas. Using our

approach will produce more accurate abundance estimates than ignoring unsampled sites

altogether, but they will obviously not be as accurate as those possible if effort could be

distributed across all sites.

For analyses of the bowhead whale data specifically, the number of recaptures was very

low, despite the substantial effort in sample collection across a wide range of areas. This fact

alone suggests that the population size is quite large, and demonstrates the inherent difficulties in

estimating abundance for EC-WG bowhead whales. Given their movement dynamics and

heterogeneity, it seems that any estimate of population size is going to have its shortcomings,

and have to be taken with a fairly large degree of caution. However, here we have tried to be

comprehensive in our analyses to try to make the most of the data that are available.

For the location-specific estimates for sampled locations, our results are comparable to

previously published estimates that are available. First, Cosens and Innes (2000) report

abundance estimates from aerial line-transect surveys for a larger area centred around Repulse

Bay. Their estimate was 75 individuals, with a 95% confidence interval of 17–133. This agrees

well with our estimate of 35 (95% HDI 20–90), particularly when considering that they covered

a larger area than was available here. Second, Rekdal et al. (2014) report aerial survey line-

transect and genetic mark-recapture abundance estimates for West Greenland. The estimates

from these methods were 744 whales (95% CI: 357–1,461) and 1,538 whales (827–2,249),

respectively. However, the aerial surveys were conducted in a single year, and therefore the

resulting estimates do not include whales that may use the area, but did not do so in the year of

the survey, which may represent a substantial number. Our genetic mark-recapture abundance

estimate for West Greenland is a bit larger (2,615, 95% HDI: 1,353–6,114), and with a heavier

right-handed tail. This makes sense given that our approach attempts to infer whales that may be

missing, and therefore results in higher probabilities associated with larger estimates. Given the

heterogeneity in bowhead movement patterns, we think that basing estimates on a longer-term

data set increases the chances of including all whales that use specific areas, and will therefore

result in more accurate estimates.

For total abundance estimates, our values were fairly similar across approaches (location-

specific and location-independent), as well as across time periods used (full data set versus 5

years of data) (**Table 4**). In most cases the estimates for the 5-year data set were slightly lower

than those for the whole data set. This could result from two primary reasons. First, the higher

estimates may be correct, but only obtained with the larger data set due to the much larger

sample size available. Second, the higher estimates may be biased by population growth during

the study period, which would result in abundance estimates for the longer time period being

biased upwards. Precision was higher for the longer time period, which was expected due to the

larger number of recaptures available for the longer-term data set. Given these differences and

potential issues, we think that the location-specific estimate based on the full data set currently

represents the best overall estimate from these analyses (11,747 whales, 95% HDI: 8,169–

20,043). The precision of this estimate is quite low, indicating that effort should be improved in

the future to obtain more samples from these locations. Interestingly, Fisheries and Oceans

Canada (DFO) recently (in 2013) carried out large-scale aerial surveys of the arctic to estimate

abundance of narwhal and bowhead whale populations. The estimate they obtained for the EC-

WG bowhead population was 6,743 individuals (CV = 22%) (Doniol-Valcroze et al. 2014).

However, this is likely an underestimate given that several key areas were not covered by those

surveys (Foxe Basin, and northern Hudson Bay). Thus, we would expect our estimate to be

larger than that reported based on the aerial surveys.

These analyses show that, at least in the tested conditions, it is possible to use regression

techniques to infer "missing" individuals in mark-recapture abundance analyses, and therefore

improve abundance estimates for situations where it is not possible to sample all locations. This

advancement should be useful to a wide range of studies. We recognize that several

improvements could be made to this approach, such as incorporating it into open population

models, as opposed to just the closed population models considered here. However, we think that

it represents an important step forward - and a useful tool - in the quest to obtain accurate

estimates of abundance for populations that are difficult to study, but it is not meant to represent

a final solution. For studies and management of the EC-WG bowhead whale population, the

benefits of this approach may take many other forms as well. For example, the systematic

collection and analysis of biopsy samples from a few key locations is vastly less expensive than

conducting exhaustive aerial surveys of the high arctic. Thus, this may represent a more efficient

means to assess and monitor this population over time than the methods that are currently

employed.

Acknowledgements

Over the years numerous people have been involved in sample collection. This work would not have been possible without their participation. We sincerely thank the Hunters and Trappers Organizations and the communities of Igloolik, Pangnirtung, Arctic Bay, Repulse Bay, Kuugaaruk, Taloyoak and Cape Dorset who provided logistic support, field assistance and/or participation for the biopsy sampling and collection programs. We appreciate the work done by Denise Tenkula (DFO) for laboratory analyses and scoring of alleles. We are grateful to DFO staff for their efforts in receiving, handling and curating of samples. Funding for the sample collection and genetic analyses has been obtained from DFO, Polar Continental Shelf Project, Nunavut Implementation Fund, the Nunavut Wildlife Research Trust, ArcticNet Centre of Excellence, International Polar Year (Global Warming and Arctic Marine Mammals), Assiniboine Park Zoo, the University of Manitoba and Natural Sciences and Engineering Research Council. This work was funded by Fisheries and Oceans Canada (DFO). To comments of two anonymous reviewers greatly improved this manuscript and the degree to which this approach was tested.

Data Accessibility

The data, code, and instructions for combining the two to obtain the results are available at https://github.com/timothyfrasier/bowhead_abundance. Specifically, there is a pdf file containing "walk-through" instructions, explaining how to conduct the analyses and obtain the results

617 presented in the paper. There are also two folders: one containing the required data, and another

618 containing the required code. Users should be able to replicate our results by following the

619 instructions.

630   **Figure 1.** Bowhead whale (*Balaena mysticetus*). Photograph by Jeff Higdon.

644 **Figure 2.** Distribution and sampling locations of the eastern Arctic bowhead population.

645 Sampling locations are numbered as follows: 1 = Arctic Bay, 2 = Cape Dorset, 3 = Igloolik, 4 =

646 Kugaaruk, 5 = Pangnirtung, 6 = Repulse Bay, 7 = Taloyoak, and 8 = Disko Bay.

663 **Figure 3.** Illustration and brief description of the main steps of the model.

675 **Figure 4.** Range of abundance estimates (interval between the upper and lower 95% HDI values)

676 obtained for total abundance when 5, 10, 15, 20, and 25% of the individuals from each location

677 are sampled.

694 **Figure 5.** Abundance estimates and associated 95% HDIs for each simulated scenario testing the

695 effects of sampling effort. Actual values are shown in black circles. Here "Missing" refers to the

696 number of individuals in the unsampled location(s).

Figure 6. Abundance estimates and associated 95% HDIs for each simulated scenario testing the

effects of varying population sizes and sampling efforts across locations. Actual values are

shown in black circles. Here "Missing" refers to the number of individuals in the unsampled

location(s). Note that the "total" abundance estimate for all 5 tested scenarios are located in the

bottom-right panel. These were separated from the other results for each simulated condition

because the scale is so different for the total estimates versus those for each location

independently.

Figure 7. Actual (black circles), and estimated (open squares) abundance for four sampled

locations, and total abundance, when there is one unsampled location. Bars represent the 95%

HDIs. A. Abundance for each location (4 sampled and 1 unsampled) are 5,000 individuals, for a

total of 25,000. The movement rates among sampled locations was high (0.2), and those between

the sampled and unsampled locations was low (0.05). B. The reverse situation from A:

abundance at each location as 5,000 (for a total to 25,000), but movement rates among sampled

locations was low (0.05) and those between sampled and unsampled locations was high (0.20).

C. All movement rates were equal at 0.15, but abundance was 5,000 individuals for all sampled

locations and 500 for the unsampled location, for a total abundance of 20,500 individuals. D. The

reverse of C: all movement rates were equal at 0.15, but abundance was 500 individuals for all

sampled locations and 5,000 for the unsampled location, for a total abundance of 7,000

individuals.

Marked Individuals	Area 1	Area 2	Area 3
1	1	1	1
16	1	1	0
3	1	0	1
35	1	0	0
2	0	1	1
2	0	1	0
16	0	0	1
?	0	0	0

Table 1. A table showing the data layout as needed to infer the number of marked individuals in

unsampled sites, as shown in Durban et al. (2005). Each row represents individuals with a

particular capture history. For example, row one represents individuals seen in all three areas.

Row two represents individuals seen in areas #1 and #2, but not area #3. Row three represents

individuals seen in area #1 and area #3, but not area #2, and so on. The last row represents

identifiable individuals not seen in any of the surveyed areas, and the number of individuals with

this sighting history is to be inferred.

Location	Count
Arctic Bay	9
Cape Dorset	1
Greenland	295
Igloolik	546
Iqaluit	1
Kugaaruk	8
Pangnirtung	293
Repulse Bay	23
Taloyoak	1
Total	**1,177**

749   **Table 2**. Number of genotyped samples by location.

	Greenland	Igloolik	Pangnirtung	Repulse Bay
Greenland	5	0	2	0
Igloolik	0	23	6	0
Pangnirtung	0	0	3	3
Repulse Bay	0	0	0	1

753   **Table 3**. Identified recaptures by location. Only locations with recaptures are included. Location

754   of first capture is indicated by row, whereas location of recapture is indicated by column (i.e,.

755   there where two whale first "captured" in Greenland that were later recaptured in Pangnirtung).

Row #	Location	Full Data Set		5 Years of Data	
		Estimate	95% HDI	Estimate	95% HDI
1	Greenland	2,615	1,353–6,114	2,089	928–5,748
2	Igloolik	4,201	2,952–6,252	2,571	1,698–4,024
3	Pangnirtung	2,378	1,491–4,203	2,852	1,210–7,868
4	Repulse Bay	35	20–90	NA	NA
5	"Missing"	449	0–8,297	4,111	179–19,721
6	Total-LS	11,747	8,169–20,043	13,899	7,782–30,602
7	Total-LI	11,682	8,620–16,014	6,877	4,828–11,477

Table 4. Abundance estimates (modes of the posterior probabilities) for each location, as well as

for the full population. Included are these estimates and 95% highest density intervals (HDI).

Total-LS refers to the abundance estimate for the total population based on inference from the

location-specific analyses and Total-LI refers to estimates based on the location-independent

analyses.

References

Amstrup, S.C., McDonald, T.L., and Manly, B.F.J. 2005. Handbook of Capture-Recapture

Analysis. Princeton University Press, Princeton, NJ.

Best, P., Reeb, D., Rew, M., Palsbøll, P., Schaeff, C., and Brandão, A. 2005. Biopsying southern

right whales: Their reactions and effects on reproduction. The Journal of Wildlife Management:

1171–1180.

Bradshaw, C.J.A., Mollet, H.F., and Meekan, M.G. 2007. Inferring population trends for the

world's largest fish from mark-recapture estimates of survival. Journal of Animal Ecology **76**:

480–489.

Carpenter, B., Gelman, A., Hoffman, M.D., Lee, D., Goodrich, B., Betancourt, M., Brubaker,

M., Guo, J., Li, P., and Riddell, A. 2017. Stan: A Probabilistic Programming Language. Journal

of Statistical Software **76**: 1–32.

Cosens, S.E., and Innes, S. 2000. Distribution and numbers of bowhead whales (*Balaena*

mysticetus) in northwestern Hudson Bay in August 1995. Arctic **53**: 36–41.

Doniol-Valcroze, T., Gosselin, J.F., Pike, D., Lawson, J., Asselin, N., Hedges, K., and Ferguson,

S. 2014. Abundance estimate of the Eastern Canada - West Greenland bowhead whale

population based on the 2013 High Arctic Cetacean Survey. DFO Can. Sci. Advis. Sec. Res.

Doc. 2014/WP 2b, Ottawa, ON.

Durban, J.W., Elston, D.A., Ellifrit, D.K., Dickson, E., Hammond, P.S., and Thompson, P.M.

2005. Multisite mark-recapture for cetaceans: population estimates with Bayesian model

averaging. Marine Mammal Science **21**: 80–92.

Ferguson, S.H., Dueck, L., Loseto, L.L., and Luque, S.P. 2010. Bowhead whale *Balaena*

mysticetus seasonal selection of sea ice. Marine Ecology Progress Series **411**: 285–297.

Finley, K. 1990. Isabella Bay, Baffin Island: an important historical and present-day

concentration area for the endangered bowhead whale (*Balaena mysticetus*) of the eastern

Canadian Arctic. Arctic **43**: 137–152.

Finley, K.J. 2001. Natural history and conservation of the Greenland whale, or bowhead, in the

northwest Atlantic. Arctic **54**: 55–76.

Galpern, P., Manseau, M., Hettinga, P., Smith, K., and Wilson, P. 2012. Allelematch: an R

package for identifying unique multilocus genotypes where genotyping error and missing data

may be present. Molecular Ecology Resources **12**: 771–778.

George, J.C., and Bockstoce, J.R. 2008. Two historical weapon fragments as an aid to estimating

the longevity and movements of bowhead whales. Polar Biology **31**: 751–754.

George, J.C., Bada, J., Zeh, J., Scott, L., Brown, S.E., OHara, T., and Suydam, R. 1999. Age and

growth estimates of bowhead whales (*Balaena mysticetus*) via aspartic acid racemization.

Canadian Journal of Zoology **77**: 571–580.

Heide-Jørgensen, M.P., Laidre, K.L., Jensen, M.V., Dueck, L., and Postma, L.D. 2006.

Dissolving stock discreteness with satellite tracking: bowhead whales in Baffin Bay. Marine

Mammal Science **22**: 34–45.

Heide-Jørgensen, M.P., Laidre, K.L., Wiig, Ø., Jensen, M.V., Dueck, L., Maiers, L.D., Schmidt,

H.C., and Hobbs, R.C. 2003. From Greenland to Canada in ten days: tracks of bowhead whales,

Balaena mysticetus, across Baffin Bay. Arctic **56**: 21–31.

Heide-Jørgensen, M.P., Laidre, K.L., Wiig, Ø., Postma, L., Dueck, L., and Bachmann, L. 2010.

Large-scale sexual segregation of bowhead whales. Endangered Species Research **13**: 73–78.

Hestbeck, J.B., Nichols, J.D., and Malecki, R.A. 1991. Estimates of movement and site fidelity

using mark-resight data of wintering Canada geese. Ecology and Evolution **72**: 523–533.

Kéry, M., and Schaub, M. 2012. Bayesian Population Analysis Using WinBUGS. Academic

Press, Waltham, MA.

Koski, W.R., Da-Silva, C.Q., Zeh, J., and Reeves, R.R. 2013. Evaluation of the potential to use

capture-recapture analyses of photographs to estimate the size of the eastern Canada west

Greenland bowhead whale (*Balaena mysticetus*) population. Canadian Wildlife Biology

Management **2**: 23–35.

Kowarski, K.A., Augusto, J.F., Frasier, T.R., and Whitehead, H. 2014. Effects of remote biposy

sampling on long-finned pilot whales (*Globicephala melas*) in Nova Scotia. Aquatic Mammals

40: 117–125.

Kraus, S.D., Hamilton, P.K., Kenney, R.D., Knowlton, A.R., and Slay, C.K. 2001. Reproductive

parameters of the North Atlantic right whale. Journal of Cetacean Research and Management

Special Issue 2: 231–236.

Langtimm, C.A., Beck, C.A., Edwards, H.H., Fick-Child, K.J., Ackerman, B.B., Barton, S.L.,

and Hartley, W.C. 2004. Survival estimates for Florida manatees from the photo-identification of

individuals. Marine Mammal Science **20**: 438–463.

Matthews, J.N., Steiner, L., and Gordon, J. 2001. Mark-recapture analysis of sperm whale

(*Physeter macrocephalus*) photo-id data from the Azores (1987-1995). Journal of Heredity **3**:

219–226.

Miller, C.R., Joyce, P., and Waits, L.P. 2005. A new method for estimating the size of small

populations from genetic mark-recapture data. Molecular Ecology **14**: 1991–2005.

Moore, S.E., and Reeves, R.R. 1993. Distribution and movement. *In* The bowhead whale. *Edited*

by J.J. Burns, J.J. Montague, and C.J. Cowles. The Sociey of Marine Mammalogy, Lawrence,

KS. pp. 313–386.

Mowat, G., and Strobeck, C. 2000. Estimating population size of grizzly bears using hair

capture, DNA profiling, and mark-recapture analysis. The Journal of Wildlife Management **64**:

183–193.

National Research Council. 1996. The Evaluation of Forensic DNA Evidence. National

Academy Press, Washington, D.C.

Nerini, M.K., Braham, H.W., Marquette, W.M., and Rugh, D.J. 1984. Life history of the

bowhead whale, *Balaena mysticetus* (Mammalia: Cetacea). Journal of Zoology **204**: 443–468.

Noren, D.P., and Mocklin, J.A. 2011. Review of cetacean biopsy techniques: Factors

contributing to successful sample collection and physiological and behavioral impacts. Marine

Mammal Science **28**: 154–199.

Otis, D.L., Burnham, K.P., White, G.C., and Anderson, D.R. 1978. Statistical inference from

capture data on closed animal populations. Wildlife monographs **62**: 3–135.

Petit, E., and Valiere, N. 2006. Estimating population size with noninvasive capture-mark-

recapture data. Conservation Biology **20**: 1062–1073.

Pradel, R., Johnson, A.R., Viallefont, A., Nager, R.G., and Cézilly, F. 1997. Local recruitment in

the greater flamingo: a new approach using capture-mark-recapture data. Ecology and Evolution

78: 1431–1445.

R Core Team. 2015. R: A language and environment for statistical computing. R Foundation for

Statistical Computing, Vienna, Austria. Available from http://www.R-project.org/.

Reeves, R., Mitchell, E., Mansfield, A., and McLaughlin, M. 1983. Distribution and migration of

the bowhead whale, *Balaena mysticetus*, in the eastern North American Arctic. Arctic **36**: 5–64.

Rekdal, S.L., Hansen, R.G., Borchers, D., Bachmann, L., Laidre, K.L., Wiig, Ø., Nielsen, N.H.,

Fossette, S., Tervo, O., and Heide-Jørgensen, M.P. 2014. Trends in bowhead whales in West

Greenland: Aerial surveys vs. genetic capture-recapture analyses. Marine Mammal Science **31**:

133–154.

Rice, W.R., and Harder, J.D. 1977. Application of multiple aerial sampling to a mark-recapture

census of white-tailed deer. The Journal of Wildlife Management **41**: 197–206.

Royle, J.A., Dorazio, R.M., and Link, W.A. 2007. Analysis of multinomial models with

unknown index using data augmentation. Journal of Computational and Graphical Statistics **16**:

67–85.

Rugh, D., Demaster, D., Rooney, A., Breiwick, J., Shelden, K., and Moore, S. 2003. A review of

bowhead whale (*Balaena mysticetus*) stock identity. Journal of Cetacean Research and

Management **5**: 267–279.

Seutin, G., White, B., and Boag, P. 1991. Preservation of avian blood and tissue samples for

DNA analyses. Canadian Journal of Zoology **69**: 82–90.

Shaw, C., Wilson, P., and White, B. 2003. A reliable molecular method of gender determination

for mammals. Journal of Mammalogy **84**: 123–128.

Stan Development Team. 2018. RStan: The r interface to stan. R package version 2.17.3.

Tanner, M.A., and Wong, W.H. 1987. The calculation of posterior distributions by data

augmentation. Journal of the American Statistical Association **82**: 528–540.

Wang, J.Y., Frasier, T.R., Yang, S.C., and White, B.N. 2008. Detecting recent speciation events:

the case of the finless porpoise (genus *Neophocaena*). Heredity **101**: 145–155.

Wiig, Ø., Heide-Jørgensen, M.P., Lindqvist, C., Laidre, K.L., Postma, L.D., Dueck, L., Palsbøll,

P.J., and Bachmann, L. 2011. Recaptures of genotyped bowhead whales *Balaena mysticetus* in

eastern Canada and West Greenland. Endangered Species Research **14**: 235–242.

Williams, B.K., Nichols, J.D., and Conroy, M.J. 2002. Analysis and Management of Animal

Populations. Academic Press, Sand Diego, CA.

Wilson, B., Hammond, P.S., and Thompson, P.M. 1999. Estimating size and assessing trends in

a coastal bottlenose dolphin population. Ecological Applications **9**: 288–300.

Zeh, J., Poole, D., Miller, G., Koski, W., Baraff, L., and Hugh, D. 2002. Survival of bowhead

whales, *Balaena mysticetus*, estimated from 19811998 photoidentification data. Biometrics **58**:

832–840.

887