Bayesian abundance estimation from genetic mark-recapture

data when not all sites are sampled: an example with the

bowhead whale

Timothy R. Frasier[1], Stephen D. Petersen[2], Lianne Postma[3], Lucy Johnson[2], Mads Peter

Heide-Jørgensen[4] & Steven H. Ferguson[3]

[1]Department of Biology, Saint Mary's University, 923 Robie St., Halifax, NS, B3H 3C3 Canada,

[2]Conservation and Research Department, Assiniboine Park Zoo, 2595 Roblin Blvd., Winnipeg Manitoba, R3R 0B8,

Canada,

[3]Fisheries and Oceans Canada, 501 University Crescent, Winnipeg, Manitoba, R3T 2N6, Canada,

[4]Greenland Institute of Natural Resources, Box 570, 3900 Nuuk, Greenland.

<u>Corresponding Author:</u>

Timothy R. Frasier

Saint Mary's University

Department of Biology

923 Robie Street

Halifax, NS B3H 3C3

Canada

E-mail: timothy.frasier@smu.ca

Tel: (902) 491-6382

Fax: (902) 420-5046

Abstract

Estimating abundance is one of the most fundamental and important aspects of population biology, with major implications on how the status of a population is perceived and thus on conservation and management efforts. Although typically based on one of two methods (distance sampling or mark-recapture), there are many individual identification methods that can be used for mark-recapture purposes. In recent years, the use of genetic data for individual identification and abundance estimation through mark-recapture analyses have increased, and in some situations such genetic identifications are more efficient than their field-based counterparts for population monitoring. One issue with mark-recapture analyses, regardless of which method of individual identification is used, is that the study area must provide adequate opportunities for "capturing" all individuals within a population. However, many populations are unevenly and widely distributed, making it unfeasible to adequately sample all necessary areas. Here we develop an analytical technique that accounts for unsampled locations, and provides a means to infer "missing" individuals from unsampled locations, and therefore obtain more accurate abundance estimates when it is not possible to sample all sites. This method is validated using simulations, and is used to estimate abundance of the Eastern Canada-West Greenland (EC-WG) bowhead whale population. Based on these analyses, the estimated size of this population is 9,089 individuals, with a 95% highest density interval of 5,107–17,079.

Keywords: abundance estimation, mark-recapture, genetic identification, whale

Introduction

Mark-recapture analyses have become an invaluable tool not only as a means for

abundance estimation, but also for obtaining a wealth of other information on population

dynamics, such as population growth rates, life history, and survival probabilities [1-3].

Traditionally, mark-recapture methods involved the actual capture, marking, and

recapture of individuals [4-6], but their use quickly expanded to species where

individuals could be identified based on natural markings, and thus "capturing" in these

cases meant taking a picture that could be used for identification [7-10]. More recently,

the increasing ease with which individual-specific genetic profiles can be obtained from

small sources of DNA has made it possible (and sometimes more efficient) to use genetic

data as the primary means of individual identification, and thus the basis for mark-

recapture analyses [11-13].

One important requirement in mark-recapture studies, regardless of which method

of identification is used, is ensuring that the study area provides adequate opportunities

for "capturing" all individuals within the study population. This requirement may be

easily addressed for populations with small ranges that can be surveyed in their entirety,

or where all individuals utilize—or pass through—a single area. However, many

populations are not so amenable, with individuals exhibiting substantial variation in site

fidelity and habitat use patterns. Failure to account for this variation can bias abundance

estimates either upwards or downwards. The former can occur if marked individuals

move out of the study area and thus become unavailable for recapture. The latter can

occur if the study area is only used by a portion of the population, and thus estimates are

only representative of some individuals rather than the entire population.

For species with wide distributions and substantial heterogeneity in habitat use

patterns it may not be possible to obtain adequate coverage of all sites. In such cases

abundance estimation requires appropriate corrections during the analytical stages of the

study. In 2005, Durban et al. made substantial progress in this area by recognizing that

analytically accounting for missed sites is essentially a regression problem, where data

from sampled areas can be used to make predictions on the number of individuals not

seen in any surveyed area [14]. As an example, suppose a study population of organisms

where some, but not all, individuals can be identified based on natural markings. Suppose

the number of identifiable individuals has been counted at each of three sites, but other

unstudied sites exist. To estimate the total size of the identifiable (marked) population,

Durban et al. first organized the sighting histories of identifiable individuals based on

whether or not they were observed in each of the three surveyed areas (see **Table 1**). The

last row of such a table represents the identifiable individuals not seen in *any* location,

and can be inferred from the available data. In this case a regression model is fit with the

first seven rows of data, and the coefficient estimates from this process are then used to

predict the value for the last row. In their application of this approach to a study of

bottlenose dolphins (*Tursiops truncatus*) around Scotland, Durban et al. found that their

approach performed quite well and resulted in a similar estimate to that obtained through

another study based on a more rigorous and standardized survey design.

The eastern Canada-western Greenland (EC-WG) bowhead whale (*Balaena*

mysticetus) (**Figure 1**) population epitomizes the confounding factors associated with

abundance estimation, using either distance sampling or mark-recapture methods. First, it

is an arctic-adapted population with a wide distribution often associated with the ice edge

(**Figure 2**) [15-18]. The necessary logistics and monetary requirements make extensive

surveys in the high arctic challenging. Second, the population is highly segregated based

on sex, age class, and perhaps other factors [15,16,19,20]. Indeed, the segregation is so

great that, until recently, multiple populations (and subsequent management stocks) were

thought to exist [21]. It was only when satellite tagging data recently showed movement

of whales across previously assumed population boundaries, and such movements were

also identified based on genetic mark-recapture analyses, that the presence of only one

population was recognized [22-24]. These confounding factors have led to very imprecise

abundance estimates, as well as large differences in estimates based on data from

different locations and/or different analytical approaches, as summarized in Koski et al.

[25].

Here, we outline and utilize two new approaches that may improve abundance

estimates for the eastern Arctic-western Greenland bowhead whale population. First, our

analyses are based on genetic identifications. This approach may prove more efficient

than photograph-based identifications because samples can be collected in collaboration

with Arctic communities, eliminating the need for extensive and expensive aerial surveys

while also fostering closer collaborations and relationships with Arctic communities.

Second, we extend the approach of Durban et al. [14] in an important way. The model of

Durban et al. was designed to estimate the number of "missing" *marked* individuals in a

population. However, our goal was to estimate the *total number* of missing individuals

(marked and unmarked), and combine this information with data from sampled sites to

result in one estimate of total abundance for the population of interest. If successful, this

latter approach could prove useful to a broad range of population studies.

Materials and Methods

Sampling

Intensive sampling effort has taken place out of the communities of Igloolik and
Pangnirtung in Canada, and in Disko Bay, Greenland (hereafter "Greenland") (**Figure 2**,
Table S1). Additionally, more opportunistic sample collection has occurred in Repulse
Bay and Arctic Bay, as well as from 11 subsistence hunts by Canadian Inuit (Taloyoak,
Kugaaruk, Cape Dorset, Pangnirtung, Repulse Bay, Igloolik) (**Figure 2**, **Table S1**). In
total, 1,177 samples have been collected across 19 years.

In each location, small tissue (skin and blubber) samples were obtained from free-
swimming whales using a 150 lb draw-weight crossbow (Excalibur Vixen) with bolts and
tips modified specifically for this purpose (from Ceta-DART, Denmark). This is the most
common method for collecting small skin samples from free-swimming whales, and has
no negative short or long-term effects on individuals other than an initial "startle"
response in some cases [26-28]. Biopsy samples were collected in association with photo-
identification data, as part of ongoing photo-identification studies in all areas. Samples
were preserved in either a salt-saturated 20% DMSO solution [29], flash frozen in liquid
nitrogen, in RNAlater (Qiagen), or in Allprotect (Qiagen) and frozen upon arrival in the
laboratory. All samples were archived in -80°C freezers at the Freshwater Institute (DFO,
Canada, Winnipeg) laboratories until further analysis. Bowhead whale biopsy samples
were collected under permit Department of Fisheries and Oceans License to Fish for
Scientific Purposes S-16/17 1005-NU and Animal Use Protocol FWI-ACC-2016-09.

Genetic Analyses

DNA was extracted using a modified version of the standard Qiagen extraction protocol [30]. DNA was quantified using a μQuant spectrophotometer (Bio-Tek) or a NanoDrop spectrophotemeter (ThermoFisher Scientific) and normalized to 10–100 ng/μl. Sex was determined through the amplification of a zinc finger gene intron using primers LGL331 and LGL335 [31]. PCR fragments were size-separated in 1.5% agarose gels and visualized with GelRed (Biotium). The resulting banding pattern of X (∼ 975 bp) and Y (∼ 1040 bp) fragments were used to infer sex.

Multiplex reactions were developed and used to amplify 23 microsatellite loci (**Table S2**). The reaction mixtures contained 1.0 μl of template DNA, 1X AmpliTaq Gold Buffer, 2.0 mM $MgCl_2$, 0.25 mM each dNTP, 0.5 U AmpliTaq Gold DNA polymerase (Life Technologies), and the primer concentrations indicated in **Table S2**. Thermal cycler profiles included a five minute initial denaturing step at 95°C followed by 35 cycles of: 95°C for 30 seconds, annealing temperature (**Table S2**) for 30 seconds, and 72°C for 30 seconds; followed by a final extension step of 72°C for 30 minutes. PCR products were size-separated and visualized on an Applied Biosystems 3130xl Genetic Analyzer (Life Technologies), with GeneScan 600 LIZ (Life Technologies) as a size-standard. Size-standards were checked, and allele calls made using the program GeneMarker (SoftGenetics).

Identifying Recaptures

The crux in identifying genetic recaptures is developing appropriate criteria for defining

when two genotypes can be considered "the same", and thus representing the same

individual. This problem has a long history, and has been most thoroughly dealt with in

human forensic DNA typing [32]. The issues are two-fold. First, two genotypes may be

the same at the typed loci, but represent different individuals if not enough resolution is

available with the chosen markers, if individuals are related, or just by chance (a false

inclusion). Second, two genotypes may be from the same individual but be scored

differently due to genotyping errors or null alleles (a false exclusion). Thus, the key is

identifying what criteria minimize the chances of each of these errors.

To identify recaptures, or samples likely originating from the same individual, we

used the R package ALLELEMATCH [33]. This package deals with these issues by

calculating the genetic similarity of all genotypes, and then creating clusters based on

genetic similarity. It then re-creates these clusters by sequentially allowing an

incremental number of mismatches, while keeping track of how many "unique"

genotypes are present under each number of allowed mismatches. The appropriate

number of allowed mismatches is then identified as the point when each genotype

associates with only one cluster.

Estimating Abundance

We used Bayesian methods to estimate abundance from the genetic mark-recapture data,

based on the general approaches described in Kéry & Schaub [34] and Kruschke [35].

We tried two general strategies, each with their own strengths and weaknesses. The first

and most simple approach involved ignoring the location information and treating the

data as one large capture-mark-recapture study (hereafter called the "location-

independent" approach). The main benefit of this approach is that it allows all samples to

be used, regardless of whether recaptures were found within each location. The main

weakness of this approach is that it does not explicitly account for the fact that not all

locations have been sampled, and therefore makes the assumption that the sampled

locations represent adequate sampling opportunities for all individuals. Our second

strategy was modelled after that of Durban et al. [14]. Here, location-specific estimates of

abundance and movement rates were obtained and used to explicitly infer the number of

individuals in unsampled locations (hereafter called the "location-specific" approach).

The strength of this approach is that it provides an explicit and quantitative means to infer

the number of individuals using unsampled locations. The main weakness is that it

requires dividing the data by locations, with the consequence that many locations may not

have enough samples to be included in the analyses, and thus some information may be

lost. Both of these methods are described in more detail below.

The number of recaptures for this data set was small, and no individual was

sampled more than twice (see Results). We therefore estimated abundance using closed

population models because we did not feel that enough information was available to

obtain useful estimates of all the parameters included in an open population model.

Additionally, the long life-span and generally slow life history of bowhead whales mean

that it is unlikely that there were dramatic changes to population size throughout the

study period [36-39]. However, to try to account for any potential changes over time, for

each strategy (location-independent and location-specific) we estimated population sizes

based on two data sets: the full data set, and a subset containing only five years of data.

The rationale is that the population may have changed throughout the 19 years through

which the samples were collected. Therefore, closed-population abundance estimates

based on the entire study period may be biased. The 5-year data set contains fewer data,

but is also expected to be less biased by changes in population size within the sampling

period. Combined, this resulted in four estimates of abundance based on: (1) location-

independent full data set (LI-FD); (2) location-independent 5-year data set (LI-5Y); (3)

location-specific full data set (LS-FD); and (4) location-specific 5-year data set (LS-5Y).

For all analyses, the recaptures identified by ALLELEMATCH were filtered to

exclude recaptures from the same location in the same year. Bowhead whales are known

to vary their migration and habitat use patterns depending on age, sex, and life-history

stage [16,20]. Additionally, once in an area, whales are likely to remain there for some

period of residency [19]. Thus, we thought that within-year recaptures within locations

would be artificially high and would not be representative of general capture probabilities

for the entire population throughout the study period.

Location-Independent Full Data Set (LI-FD)

To estimate abundance for the entire data set, the recapture data from ALLELEMATCH

were first converted to typical sighting histories consisting of ones and zeros for each

sampling period (year) where individuals were, or were not, sampled, respectively. These

sighting histories were augmented by 30,000 to ensure that considered probabilities

covered a wide enough range to ensure adequate sampling of the posterior distribution

[40, 41]. The parameters of interest are the sighting probability (p), and the inclusion

probability (Ω), which is the probability that a member of the augmented data set is part

of the "true" population size (N). Abundance estimates were then obtained for a closed

population model (model M_0 from Otis et al. [42]) using the approach described in Royle

et al. [41] and Kéry & Schaub [34]. The priors for both the sighting probability (p) and

the inclusion probability (Ω) were uniform distributions ranging from 0 to 1. All

Bayesian analyses were conducted using a combination of R [43], runjags [44], and

JAGS [45]. The performance of models for all analyses were assessed based on

examination of trace plots, autocorrelation plots, the Gelman and Rubin diagnostic [46],

and the effective sample size, using the coda package [47].

Location-Independent 5-Year Data Set (LI-5Y)

Although it would make intuitive sense to select the most recent 5-year period for these

analyses, we did not do this because most locations did not have data for the most recent

year (2013). Instead, we queried the data to identify which 5-year subset would contain

the most sampling events. This turned out to be years 2008-2012, and therefore

represents a recent sampling period. Once this subset of data had been identified, an

estimate of abundance was obtained using the same approach as described for the full

data set.

Location-Specific Full Data Set (LS-FD)

Our location-specific approach for estimating abundance was modeled after that used by

Durban et al. [14]. The general idea is that location-specific abundance estimates can be

used in association with estimated movement rates between locations, to identify (or

infer) the number of individuals with different sighting histories such as those shown in

Table 1. When some areas are unsampled, the goal is to estimate the number of

individuals in the last row (who were not captured in any area). This abundance estimate

can then be used in association with estimates of movement rates to obtain an estimate of

total population size even when some areas remain unsampled. The entire process is

summarized in **Figure 3**, with details described below.

The first step in this process was to estimate the movement rates between all pairs

of sampled locations. This is based on the number of recaptures within each considered

location, and the number of those recaptures that represent individuals originally marked

in the alternative location. We made the assumption that pairwise movement rates were

symmetrical, largely because the number of cross-location recaptures was very low (see

Results), and therefore there were few data available from which to estimate movement

rates. However, it is possible to relax this assumption in the future. The number of

recaptures within and between each pair of locations was converted into a string of zeros

and ones, where ones represented recaptures across locations. The mode of the movement

rates, and associated 95% highest density intervals (HDIs), were then estimated using a

Bernoulli likelihood function and a beta prior with a uniform (flat) distribution. The result

is a complete posterior probability distribution for movement rates between all pairs of

sampled locations.

For the second step of the process, the probability distribution for the symmetrical

movement rate to and from the unsampled location(s) is then estimated by averaging the

posterior probabilities of all sampled locations. This approach makes the explicit

assumption that the rate at which individuals move between sampled and unsampled

locations is within the same range as the movement rates among sampled locations. We

feel that this assumption is valid in most cases, but even if not, the model still performs

well overall due to a balancing effect of how the model works. Briefly, estimating the

movement rates themselves are not the goal of the model, but rather they are used to

estimate the abundance of individuals using different locations. Therefore, if the

estimated movement rate to and from the unsampled location(s) is *lower* than the true

value, then this will result in the abundance estimates for the sampled locations being

lower than their true values (because it will be assumed that fewer individuals are missed

by moving to unsampled locations), whereas the abundance estimate for the unsampled

location(s) will be *higher* than its true value (because it will be assumed that fewer

individuals are moving into and out of that area, and thus also captured in other areas).

However, because the estimated movement rate is an average, these over- and under-

estimates will balance each other out, and the overall estimate for the total population

size should still be accurate. The same will be true for the opposite scenario where the

estimated movement rate to and from the unsampled location(s) is *higher* than the true

value.

For the third step, abundance is estimated for each sampled location. The genetic

recapture data were first converted to typical sighting histories consisting of ones and

zeros for each sampling period where individuals were, or were not, captured,

respectively. However, these sighting histories are biased due to individuals moving to

and from the unsampled location(s). Individuals marked in the sampled location could

move to an unsampled location, and therefore not be available for recapture, artificially

reducing the number of recaptures. In the opposite direction, unmarked individuals from

the unsampled location(s) could move into the sampled location, which would also

artificially reduce the proportion of recaptures. Both of these processes reduce the

proportion of individuals available for recapture, and would therefore bias population size

estimates upwards. To correct for this, for each sampled location we estimate the number

of individuals moving to and from the unsampled location(s) and randomly add these to

the sighting history data. For example, suppose at one location there were 300 marked

individuals and 68 recaptures. Suppose that the estimated movement rate to and from the

unsampled location(s) is 0.12. This means that we have "lost" 12% of marked individuals

to unsampled locations, and our number of unmarked individuals is also artificially

inflated by 12% due to movement of unmarked individuals from unsampled areas into

this area. This means that our proportion of recaptures is underestimated by two times the

movement rate (24%). Thus, our original 68 recaptures only represent (1 - 0.24 = 0.76)

76% of the recaptures we "should" have had, if individuals did not move to and from

unsampled locations (i.e., we should have had 68 / 0.76 = 89 recaptures). To account for

this, these (89 - 68 = 21) additional recaptures were randomly added to the sighting

histories for this location. Once the sighting histories are corrected, abundance was

estimated for each sampled location using the same closed population approach described

for the location-independent data sets. Note that although the model assumed a closed

population, movement to and from other locations has already been taken into account,

and therefore "closed" in this sense just refers to births and deaths.

For step four, the complete posterior probabilities for abundance for each sampled

location, and for each pairwise movement rate, are used to estimate the number of

individuals with each possible habitat use pattern across the sampled locations (there are

3 such possible patterns for 2 locations - **Figure 3**, 7 possible patterns for 3 locations, and

so on). These data are then used in a Bayesian regression analysis to estimate the relevant

regression coefficients. These coefficients are then used to predict the number of

individuals that are not seen in any of the sampled locations (i.e., the last row of the table

shown in **Figure 3**, step 4). The resulting posterior probability distribution represents the

estimate of the total number of individuals not seen in any sampled location.

For the fifth and final step, total abundance is estimated in a similar manner as

that for the "missing" individuals. Specifically, the complete posterior probabilities for

abundance for each location, and for each pairwise movement rate, are used to estimate

the number of individuals with each possible habitat use pattern across locations. The

difference is that now this table includes data for the unsampled location(s): effectively

adding another location to the table and associated movement rates (see **Figure 3**). The

sum of these values then represents the abundance estimate for the total population. Note

that one benefit of this approach is that the uncertainty in each estimate is carried over

into the uncertainty of the total abundance estimate. Put another way, rather than

estimating the total abundance based on the mean, mode, or some other summary metric

of the other parameters, this estimate is based on the entire probability distribution for

each parameter. In this way the uncertainty in each estimate of abundance and movement

are incorporated into the posterior probability distribution for the total abundance

estimate. This is one of the attractive properties of setting up the analyses in this way.

Location-Specific 5-Year Data Set (LS-5Y)

As with the location-independent 5-year data set, we conducted the same analyses as

above (for the location-specific full data set, LS-FD), except this time with a data set

containing only those sampling events in the 5-year period from 2008-2012. Again, the

idea for this process was to reduce the potential biases in the longer-term data set due to

births and deaths occurring within the longer time frame, by analyzing data from a much

shorter time period.

Model Validation

A number of simulations were conducted to assess model performance, and to provide

information that would be useful for guiding sampling strategies in the field. All

simulations were based on the location-specific model, which, instead of estimating

abundance for a single population/location, is designed to estimate abundance for each

location independently (including unsampled locations), and then come up with one total

abundance estimate for all sampled and unsampled locations. A simulation function was

written that generates data for four sampled locations and one unsampled location, where

the abundance and movement rates between locations/populations can be set by the user.

The first set of simulations focused on assessing the effects of varying sampling

effort. For these simulations all population sizes (including that of the unsampled

location) were set to 1,000 individuals (resulting in a total population size of 5,000

individuals). All movement rates between locations were also set to 10%. Abundance was

then estimated based on sampling 5%, 10%, 15%, 20%, and 25% of individuals from

each sampled location, to assess how changing sampling effort influences the precision

and accuracy of abundance estimates.

A second set of 5 simulations were also run and analyzed, but this time the

abundance at each location (including the unsampled one) was randomly drawn from a

list of 20 values ranging from 500 to 10,000, in increments of 500. The movement rates

were all set to 10%, and 250 individuals were sampled at each location. This means that

the proportion of individuals sampled could range from 2.5% (for a population size of

10,000 individuals) up to 50% (for a population size of 500 individuals). These

simulations were conducted to ensure that the model performs well even when there are

large differences in population sizes and sampling effort between locations.

Results

Samples, Genotypes, and Recaptures

A total of 1,177 samples were genotyped. **Table 2** shows how these were divided among

locations. The final microsatellite data set contained 21 loci since two loci were dropped

due to a significant portion of the samples missing data that were profiled before 2008.

Furthermore, a sample was only included in the analysis if it had at least 10 of the 21 loci

scored. This provided an overall probability of identity of 1.16×10^{-9}, which corresponds

to a 1 in 8.62×10^{8} chance that the match identified was due to two random animals

matching at all loci. **Supplementary File 1** shows how these samples were divided by

year, location, and sex.

The ALLELEMATCH analysis suggested that the best fit for the data was to allow 3

mismatches. Based on this criterion, 992 unique genotypes were identified, along with

185 recaptures. However, 136 of these recaptures represented whales recaptured within

the same area and year in which they were marked, and were therefore removed from the

analyses. This left 49 recaptures for subsequent analyses. **Table 3** summarizes the

recaptures by location. Based on the number of recaptures, only Greenland, Igloolik,

Pangnirtung, and Repulse Bay were used for estimating abundance based on the full data

set.

Model Validation

The results for the simulations testing the effects of sampling effort are shown in **Figure**

4. The precision of the estimates (or the range between to upper and lower 95% HDI

values) declines rapidly until sampling reaches ~15%. Above this sampling effort,

precision increases at a much slower rate. Thus, ~15% seems to be a turning point above

which diminishing returns are achieved as sampling effort increases. The details of all of

the estimates from these simulations are shown in **Figure 5**. These plots show that the

model performs quite well in all aspects (abundance estimation of sampled locations,

abundance estimation of unsampled location(s), and total abundance).

The results from the simulations testing the performance when population size

and sampling effort varies across locations are shown in **Figure 6**. The results show that

the model performs quite well under a range of variation in population sizes and sampling

efforts. This is particularly true for the estimates of "total" abundance, which tend to be

more accurate than the estimate for each specific location. This is due to differences

between the true and estimated movement rates as described in the Methods section.

It should be noted that all of the simulation results are based on only one iteration

of the simulation under each condition. Ideally, simulations would be run 10, 100, or

1000 times under each set of conditions to obtain information about the variance around

expectations under each conditions. However, given the way the model works, it takes

approximately one week to analyze the data from one simulation. Therefore, conducting

and analyzing a more desirable number simulations would take a prohibitive amount of

time. As a result, these results should be taken as a representative, but not a

comprehensive, evaluation of model performance under the simulated conditions.

Bowhead Whale Abundance Estimates

The location-specific abundance estimates for each location, inferred for the unsampled

location(s), and for the total population, are shown in **Table 4** (rows 1–6). The mode was

used as the point estimate instead of the mean in all cases because the posterior

probability distributions for all abundance estimates were skewed, with heavy right-hand

tails, and therefore the mode represented the estimate with a higher probability than the

mean. Note that the estimates for each location are inclusive, meaning that they represent

all whales that use each area (and may also be counted for other areas), rather than

representing whales that *only* use each area. The location–independent abundance

estimates based are also shown in **Table 4** (row 7).

Discussion

Analyses of simulated data sets suggest that our new approach performs quite well, and

could be a useful tool for estimating abundance of other populations where only a subset

of possible locations have been sampled. Simulations testing analysis performance with

respect to sampling effort show that sampling roughly 15% of the individuals at each

location is desirable, whereas sampling more intensively will result in diminishing

returns. The approach still performs well when fewer than 15% of individuals are

sampled at each location, but there is decreasing precision around the estimates as sample

size decreases (**Figure 5**).

The approach also performs well even when there is wide variation in abundance

between sampling locations. Although the simulations used in our tests were not

exhaustive, they do show that accurate abundance estimates can be obtained for the total

population when only a subset of locations are sampled, and when there is wide variation

in abundance across locations (**Figure 6**). Combined, the analysis of simulated scenarios

validates this new approach, and shows that it is indeed a viable option for estimating

abundance from genetic mark-recapture data when not all locations are sampled.

For analyses of the bowhead whale data specifically, the number of recaptures

was very low, despite the substantial effort in sample collection across a wide range of

areas. This fact alone suggests that the population size is quite large, and demonstrates

the inherent difficulties in estimating abundance for EC-WG bowhead whales. Given

their movement dynamics and heterogeneity, it seems that any estimate of population size

is going to have its shortcomings, and have to be taken with a fairly large grain of salt.

However, here we have tried to be comprehensive in our analyses to try to make the most

of the data that are available.

For the location-specific estimates for sampled locations, our results are

comparable to previously published estimates that are available. First, Cosens & Innes

[48] report abundance estimates from aerial line-transect surveys for a larger area centred

around Repulse Bay. Their estimate was 75 individuals, with a 95% confidence interval

of 17–133. This agrees well with our estimate of 30 (95% HDI 19–71), particularly when

considering that they covered a larger area than was available here. Second, Rekdal et al.

[49] report aerial survey line-transect and genetic mark-recapture abundance estimates for

West Greenland. The estimates from these methods were 744 whales (95% CI: 357–

1,461) and 1,538 whales (827–2,249), respectively. However, the aerial surveys were

conducted in a single year, and therefore the resulting estimates do not include whales

that may use the area, but did not do so in the year of the survey, which may represent a

substantial number. Our genetic mark-recapture abundance estimate for West Greenland

is (1,777, 95% HDI: 855–5,536), which is very similar to that of Rekdal et al. [49], but

with a heavier right-handed tail. This makes sense given that our approach attempts to

infer whales that may be missing, and therefore results in higher probabilities associated

with larger estimates. Given the heterogeneity in bowhead movement patterns, we think

that basing estimates on a longer-term data set increases the chances of including all

whales that use specific areas, and will therefore result in more accurate estimates.

For total abundance estimates, our values were fairly similar across approaches

(location-independent and location-dependent), as well as across time periods used (full

data set versus 5 years of data) (**Table 4**). In most cases the estimates for the 5-year data

set were slightly lower than those for the whole data set. This could result for two

primary reasons. First, the higher estimates may be correct, but only obtained with the

larger data set due to the much larger sample size available. Second, the higher estimates

may be biased by population growth during the study period, which would result in

abundance estimates for the longer time period being biased upwards. Precision was

higher for the longer time period, which was expected due to the larger number of

recaptures available for the longer-term data set. Given these differences and potential

issues, we think that the location-dependent estimate based on the 5-year data set

currently represents the best overall estimate from these analyses (9,089 whales, 95%

HDI: 5,107–17,079). This estimate is higher than that based on the location-independent

approach, but this is expected because the location-independent approach cannot account

for "missing" whales that do not utilize unsampled areas. The precision of this estimate is

quite low, indicating that effort should be improved in the future to obtain samples from

these locations. Interestingly, Fisheries and Oceans Canada (DFO) recently (in 2013)

carried out large-scale aerial surveys of the arctic to estimate abundance of narwhal and

bowhead whale populations. The estimate they obtained for the EC-WG bowhead

population was 6,743 individuals (CV = 22%) [50]. However, this is likely an

underestimate given that several key areas were not covered by those surveys (Foxe

Basin, and northern Hudson Bay). Thus, we would expect our estimate to be larger than

that reported based on the aerial surveys.

These analyses show that, at least in the tested conditions, it is possible to use

regression techniques to infer "missing" individuals in mark-recapture abundance

analyses, and therefore improve abundance estimates for situations where it is not

possible to sample all locations. This advancement should be useful to a wide range of

studies, and although it has its limitations and issues, and is by no means a final solution,

it does represent a step forward in the quest to obtain accurate estimates of abundance for

populations that are difficult to study. For studies and management of the EC-WG

bowhead whale population, the benefits of this approach may take many other forms as

well. For example, the systematic collection and analysis of biopsy samples from a few

key locations is vastly less expensive than conducting exhaustive aerial surveys of the

high arctic. Thus, this may represent a more efficient means to assess and monitor this

population over time than the methods that are currently employed.

Data Availability

The data, code, and instructions for combining the two to obtain the results are available

at https://github.com/timothyfrasier/bowhead_abundance. Specifically, there is a pdf file

containing "walk-through" instructions, explaining how to conduct the analyses and

obtain the results presented in the paper. There are also two folders: one containing the

required data, and another containing the required code. Users should be able to replicate

our results by following the instructions.

Acknowledgements

Over the years numerous people have been involved in sample collection. This work

would not have been possible without their participation. We sincerely thank the Hunters

and Trappers Organizations and the communities of Igloolik, Pangnirtung, Arctic Bay,

Repulse Bay, Kuugaaruk, Taloyoak and Cape Dorset who provided logistic support, field

assistance and/or participation for the biopsy sampling and collection programs. We

appreciate the work done by Denise Tenkula (DFO) for laboratory analyses and scoring

of alleles. We are grateful to DFO staff for their efforts in receiving, handling and

curating of samples. Funding for the sample collection and genetic analyses has been

obtained from DFO, Polar Continental Shelf Project, Nunavut Implementation Fund, the

Nunavut Wildlife Research Trust, ArcticNet Centre of Excellence, International Polar

Year (Global Warming and Arctic Marine Mammals), Assiniboine Park Zoo, the

University of Manitoba and Natural Sciences and Engineering Research Council. This

work was funded by Fisheries and Oceans Canada (DFO).

References

1.   Kraus SD, Hamilton PK, Kenney RD, Knowlton AR, Slay CK. Reproductive

parameters of the North Atlantic right whale. J. Cetacean Res. Manag. 2001;

Special Issue 2: 231-236.

2.   Williams BK, Nichols JD, Conroy MJ. Analysis and Management of Animal

Populations. Sand Diego: Academic Press; 2002.

3.   Amstrup SC, McDonald TL, Manly BFJ. Handbook of Capture-Recapture

Analysis. Princeton: Princeton University Press; 2005.

4.   Rice WR, Harder JD. Application of multiple aerial sampling to a mark-recapture

census of white-tailed deer. J. Wildl. Manage. 1977;41: 197-206.

5.   Hestbeck JB, Nichols JD, Malecki RA. Estimates of movement and site fidelity

using mark-resight data of wintering Canada geese. Ecol. Evol. 1991;72: 523-533.

6.   Pradel R, Johnson AR, Viallefont A, Nager RG, Cézilly F. Local recruitment in the

greater flamingo: a new approach using capture-mark-recapture data. Ecol. Evol.

1997;78: 1431-1445.

7.   Wilson B, Hammond PS, Thompson PM. Estimating size and assessing trends in a

coastal bottlenose dolphin population. Ecol. Appl. 1999;9: 288-300.

8.   Matthews JN, Steiner L, Gordon J. Mark-recapture analysis of sperm whale

(*Physeter macrocephalus*) photo-id data from the Azores (1987-1995). J. Hered.

2001;3: 219-226.

9.   Langtimm CA, Beck CA, Edwards HH, Fick-Child KJ, Ackerman BB, Barton SL,

et al. Survival estimates for Florida manatees from the photo-identification of

individuals. Mar. Mamm. Sci. 2004;20: 438-463.

10.   Bradshaw CJA, Mollet HF, Meekan MG. Inferring population trends for the

world's largest fish from mark-recapture estimates of survival. J. Anim. Ecol.

2007;76: 480-489.

11.   Mowat G, Strobeck C. Estimating population size of grizzly bears using hair

capture, DNA profiling, and mark-recapture analysis. J. Wildl. Manage. 2000;64:

183-193.

12.   Miller CR, Joyce P, Waits LP. A new method for estimating the size of small

populations from genetic mark-recapture data. Mol. Ecol. 2005;14: 1991-2005.

13.   Petit E, Valiere N. Estimating population size with noninvasive capture-mark-

recapture data. Conserv. Biol. 2006;20: 1062-1073.

14.   Durban JW, Elston DA, Ellifrit DK, Dickson, E. Hammond PS, Thompson PM.

Multisite mark-recapture for cetaceans: population estimates with Bayesian model

averaging. Mar. Mamm. Sci. 2005;21: 80-92.

15.   Reeves R, Mitchell E, Mansfield A, McLaughlin M. Distribution and migration of

the bowhead whale, *Balaena mysticetus*, in the eastern North American Arctic.

Arctic 1983;36: 5-64.

567 16. Moore SE, Reeves RR. Distribution and movement. In: Burns JJ, Montague JJ,

568   Cowles CJ, editors. The bowhead whale. Lawrence: The Society of Marine

569   Mammalogy; 1993. pp. 313-386.

570 17. Finley KJ. Natural history and conservation of the Greenland whale, or bowhead, in

571   the northwest Atlantic. Arctic 2001;54: 55-76.

572 18. Ferguson SH, Dueck L, Loseto LL, Luque SP. Bowhead whale *Balaena mysticetus*

573   seasonal selection of sea ice. Mar. Ecol. Prog. Ser. 2010;411: 285-297.

574 19. Finley K. Isabella Bay, Baffin Island: an important historical and present-day

575   concentration area for the endangered bowhead whale (*Balaena mysticetus*) of the

576   eastern Canadian Arctic. Arctic 1990;43: 137-152.

577 20. Heide-Jørgensen MP, Laidre KL, Wiig Ø, Postma L, Dueck L, Bachmann L.

578   Large-scale sexual segregation of bowhead whales. Endanger. Species Res.

579   2010;13: 73-78.

580 21. Rugh D, Demaster D, Rooney A, Breiwick J, Shelden K, Moore S. A review of

581   bowhead whale (*Balaena mysticetus*) stock identity. J. Cetacean Res. Manag.

582   2003;5: 267-279.

583 22. Heide-Jørgensen MP, Laidre KL, Wiig Ø, Jensen MV, Dueck L, Maiers LD, et al.

584   From Greenland to Canada in ten days: tracks of bowhead whales, *Balaena*

585   *mysticetus*, across Baffin Bay. Arctic 2003;56: 21-31.

23.   Heide-Jørgensen MP, Laidre KL, Jensen MV, Dueck L, Postma LD. Dissolving

stock discreteness with satellite tracking: bowhead whales in Baffin Bay. Mar.

Mamm. Sci. 2006;22: 34-45.

24.   Wiig Ø, Heide-Jørgensen MP, Lindqvist C, Laidre KL, Postma LD, Dueck L, et al.

Recaptures of genotyped bowhead whales *Balaena mysticetus* in eastern Canada

and West Greenland. Endanger. Species Res. 2011;14: 235-242.

25.   Koski WR, Da-Silva CQ, Zeh J, Reeves RR. Evaluation of the potential to use

capture-recapture analyses of photographs to estimate the size of the eastern

Canada west Greenland bowhead whale (*Balaena mysticetus*) population. Can.

Wild. Biol. Manage. 2013;2: 23-35.

26.   Best P, Reeb D, Rew M, Palsbøll PJ, Schaeff C, Brandão A. Biopsying southern

right whales: Their reactions and effects on reproduction. J. Wildl. Manage.

2005;69: 1171-1180.

27.   Noren DP, Mocklin JA. Review of cetacean biopsy techniques: Factors

contributing to successful sample collection and physiological and behavioral

impacts. Mar. Mamm. Sci. 2011;28: 154-199.

28.   Kowarski KA, Augusto JF, Frasier TR, Whitehead H. Effects of remote biposy

sampling on long-finned pilot whales (*Globicephala melas*) in Nova Scotia. Aquat.

Mamm. 2014;40: 117-125.

29.   Seutin G, White B, Boag P. Preservation of avian blood and tissue samples for

DNA analyses. Can. J. Zool. 1991;69: 82-90.

30.   Wang JY, Frasier TR, Yang SC, White BN. Detecting recent speciation events: the
case of the finless porpoise (genus *Neophocaena*). Heredity. 2008;101: 145-155.

31.   Shaw C, Wilson P, White B. A reliable molecular method of gender determination
for mammals. J. Mammal. 2003;84: 123-128.

32.   National Research Council. The evaluation of forensic DNA evidence. Washington,
D.C.: National Academy Press; 1996.

33.   Galpern P, Manseau M, Hettinga P, Smith K, Wilson P. Allelematch: an R package
for identifying unique multilocus genotypes where genotyping error and missing
data may be present. Mol. Ecol. Resour. 2012;12: 771-778.

34.   Kéry M, Schaub M. Bayesian Population Analysis Using WinBUGS. Waltham:
Academic Press; 2012.

35.   Kruschke JK. Doing Bayesian data analysis. Burlington: Academic Press; 2011.

36.   George JC, Bada J, Zeh J, Scott L, Brown SE, O'Hara T, et al. Age and growth
estimates of bowhead whales (*Balaena mysticetus*) via aspartic acid racemization.
Can. J. Zool. 1999;77L 571-580.

37.   George JC, Bockstoce JR. Two historical weapon fragments as an aid to estimating
the longevity and movements of bowhead whales. Polar Biol. 2008;31: 751-754.

38.   Nerini MK, Braham HW, Marquette WM, Rugh DJ. Life history of the bowhead
whale, *Balaena mysticetus* (Mammalia: Cetacea). J. Zool. 1984;204: 443-468.

39.   Zeh J, Poole D, Miller G, Koski W, Baraff L, Rugh D. Survival of bowhead

627   whales, *Balaena mysticetus*, estimated from 1981-1998 photoidentification data.

628   Biometrics 2002;58: 832-840.

629   40.   Tanner MA, Wong WH. The calculation of posterior distributions by data

630   augmentation. J. Am. Stat. Assoc. 1987;82: 528-540.

631   41.   Royle JA, Dorazio RM, Link WA. Analysis of multinomial models with unknown

632   index using data augmentation. J. Comput. Graph. Stat. 2007;16: 67-85.

633   42.   Otis DL, Burnham KP, White GC, Anderson DR. Statistical inference from capture

634   data on closed animal populations. Wildl. Monogr. 1978;62: 3-135.

635   43.   R Core Team. *R:* A language and environment for statistical computing. Vienna: R

636   Foundation for Statistical Computing; 2015.

637   44.   Denwood MJ. runjags: An R package providing interface utilities, model templates,

638   parallel computing methods and additional distributions for MCMC models in

639   JAGS. J. Stat. Softw. 2015;21: 80-92.

640   45.   Plummer M. JAGS: A program for analysis of Bayesian graphical models using

641   Gibbs sampling. Proceedings of the 3rd International Workshop on Distributed

642   Statistical Computing (DSC 2003), Vienna, Austria; 2003.

643   46.   Gelman A, Rubin DB. Inference from iterative simulation using multiple

644   sequences. Stat. Sci. 1992;7: 457-511.

645   47.   Plummer M, Best N, Cowles K, Vines K. CODA: convergence diagnosis and

646   output analysis for MCMC. R News 2006;6: 7-11.

48.   Cosens SE, Innes S. Distribution and numbers of bowhead whales (*Balaena*

mysticetus) in northwestern Hudson Bay in August 1995. Arctic 2000;53: 36-41.

49.   Rekdal SL, Hansen RG, Borchers D, Bachmann L, Laidre KL, Wiig Ø, et al.

Trends in bowhead whales in West Greenland: Aerial surveys vs. genetic capture-

recapture analyses. Mar. Mamm. Sci. 2014;31: 133–154.

50.   Doniol-Valcroze T, Gosselin JF, Pike D, Lawson J, Asselin N, Hedges K, et al.

Abundance estimate of the Eastern Canada - West Greenland bowhead whale

population based on the 2013 High Arctic Cetacean Survey. DFO Can. Sci. Advis.

Sec. Res. Doc. 2014/WP 2b, Ottawa, ON.

Table 1. A table showing the data layout as needed to infer the number of marked

individuals in unsampled sites, as shown in Durban et al. (2005). Each row represents

individuals with a particular capture history. For example, row one represents individuals

seen in all three areas. Row two represents individuals seen in areas #1 and #2, but not

area #3. Row three represents individuals seen in area #1 and area #3, but not area #2,

and so on. The last row represents identifiable individuals not seen in any of the surveyed

areas, and the number of individuals with this sighting history is to be inferred.

Marked Individuals	Area 1	Area 2	Area 3
1	1	1	1
16	1	1	0
3	1	0	1
35	1	0	0
2	0	1	1
2	0	1	0
16	0	0	1
??	0	0	0

666 **Table 2**. Number of genotyped samples by location.

Location	Count
Arctic Bay	9
Cape Dorset	1
Greenland	295
Igloolik	546
Iqaluit	1
Kugaaruk	8
Pangnirtung	293
Repulse Bay	23
Taloyoak	1
Total	**1,177**

670   **Table 3**. Identified recaptures by location. Only locations with recaptures are included.

	Greenland	Igloolik	Kugaaruk	Pangnirtung	Repulse Bay
Greenland	5	0	0	2	0
Igloolik	-	23	1	6	0
Kugaaruk	-	-	0	0	0
Pangnirtung	-	-	-	3	3
Repulse Bay	-	-	-	-	1

673   **Table 4**. Abundance estimates for each location, as well as for the full population.

674   Included are these estimates and 95% highest density intervals (HDI). Total-LS refers to

675   the abundance estimate for the total population based on inference from the location-

676   specific analyses and Total-LI refers to estimates based on the location-independent

677   analyses.

		Full Data Set		5 Years of Data	
Row #	Location	Estimate	95% HDI	Estimate	95% HDI
1	Greenland	2,254	1,250–5,097	1,777	855–5,536
2	Igloolik	3,629	2,638–5,098	2,055	1,424–3,013
3	Pangnirtung	2,179	1,257–3,659	2,510	1,150–6,102
4	Repulse Bay	30	19–71	NA	NA
5	"Missing"	1,694	0–3,940	2,122	0–10,095
6	Total-LS	10,331	7,639–14,558	9,089	5,107–17,079
7	Total-LI	12,220	8,680–16,200	7,660	4,500–11,100

Figure 1. Bowhead whale (*Balaena mysticetus*). Photograph by Jeff Higdon.

Figure 2. Distribution and sampling locations of the eastern Arctic bowhead population.

Sampling locations are numbered as follows: 1 = Arctic Bay, 2 = Cape Dorset, 3 =

Igloolik, 4 = Kugaaruk, 5 = Pangnirtung, 6 = Repulse Bay, 7 = Taloyoak, and 8 = Disko

Bay.

Figure 3. Illustration and brief description of the main steps of the model.

Figure 4. Range of abundance estimates (interval between the upper and lower 95% HDI

values) obtained for total abundance when 5, 10, 15, 20, and 25% of the individuals from

each location are sampled.

Figure 5. Abundance estimates and associated 95% HDIs for each simulated scenario

testing the effects of sampling effort. Actual values are shown in black circles. Here

"Missing" refers to the number of individuals in the unsampled location(s).

Figure 6. Abundance estimates and associated 95% HDIs for each simulated scenario

testing the effects of varying population sizes and sampling efforts across locations.

Actual values are shown in black circles. Here "Missing" refers to the number of

individuals in the unsampled location(s). Note that the "total" abundance estimate for all

5 tested scenarios are located in the bottom-right panel. These were separated from the

other results for each simulated condition because the scale is so different for the total

estimates versus those for each location independently.

Figure 1

Figure 2

Step 1:

Estimate movement rates between all pairwise combinations of sampled locations. Assume movements are symmetrical. Bayesian estimation provides complete probability distributions for possible values.

Step 2:

Estimate movement rate to unsampled location(s) as the average across all pairwise combinations. Again, this results in a complete probability distribution for this estimated movement rate.

Step 3:

Estimate abundance for each sampled location, correcting for the movement of individuals to and from unsampled location(s). Bayesian estimation provides complete probability distributions for possible values.

Step 4:

Use Pattern	Loc. 1	Loc. 2
1	YES	YES
2	YES	NO
3	NO	YES
4	NO	NO

Use abundance estimates for each sampled location and estimated movement rates to estimate the number of individuals with all possible habitat use patterns across sampled areas (example table with just 2 locations shown). Build regression model from data, and predict the number of individuals that don't use any sampled location (last row of table). This is the estimate of "missing" individuals.

Step 5:

Use Pattern	Loc. 1	Loc. 2	Unsampled
1	YES	YES	YES
2	YES	YES	NO
3	YES	NO	YES
4	NO	YES	YES
5	YES	NO	NO
6	NO	YES	NO
7	NO	NO	YES

Use abundance estimates for all locations (including unsampled ones) and estimated movement rates to estimate the number of individuals with all possible habitat use patterns. The sum of these is the abundance estimate for the total population.

Figure 3

Figure 4

Figure 5

Figure 6