Emergence of multidrug-resistant *Mycobacterium simiae*: An in vitro study from a regional tuberculosis reference laboratory in Iran

Mohammad Javad Nasiri[1*], Sirus Amini[2], Zahra Nikpor[1], Samaneh Arefzadeh[1], Mohammad Mosavi[1], Hossein Dabiri[1], Mehdi Goudarzi[1], Hossein Goudarzi[1], Payam Tabarsi[3], Davood Darban Sarokhalil[4], Abbas Ali Imani Fooladi[5] and Mohammad Mehdi Feizabadi[5]

[1]Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

[2]Regional Tuberculosis Reference Laboratory, Tehran University of Medical Science, Tehran, Iran

[3]Clinical TB and Epidemiology Research Center, National Research Institute of Tuberculosis and Lung Diseases, Shahid Beheshti University of Medical Sciences, Tehran, Iran

[4]Department of Microbiology, School of Medicine, Iran University of Medical Sciences, Tehran, Iran

[5]Microbiology Research Center, Baqiyatallah University of Medical Sciences, Tehran, Iran

[6]Department of Medical Microbiology, School of Medicine, Tehran University of Medical Sciences, Tehran, Iran

*Corresponding author:

Department of Microbiology, School of Medicine, Shahid Beheshti University of Medical Sciences, Tehran, Iran

E-mail: mj.nasiri@hotmail.com

Abstract

Introduction: *Mycobacterium simiae* is an emerging pathogen in Iran and little is known about drug susceptibility patterns of this pathogen.

Materials and methods: Twenty five clinical isolates of *M. simiae* from 80 patients with confirmed NTM pulmonary disease were included in this study. For drug susceptibility testing (DST), proportional and broth microdilution methods were used according to the clinical and laboratory standards institute (CLSI) guideline.

Results: All clinical isolates of *M. simiae* were resistant to isoniazid, rifampicin, ethambutol, streptomycin, amikacin, kanamycin, ciprofloxacin and clarithromycin. They also were highly resistant to ofloxacin (80%). Susceptibility to ofloxacin was only noted in the 5 isolates.

Conclusions: Clinical isolates of *M. simiae* were multidrug resistant, and had different drug susceptibility patterns than previously published studies. DST results can assist in selecting more appropriate treatment regimens. Newer drugs with proven clinical efficacy correlating with in vitro susceptibility should be substituted with first- and second line anti-TB drug testing.

Keywords: *Mycobacterium simiae,* drug resistance, Iran

Introduction

Infections caused by nontuberculous mycobacteria (NTM) have been recently reported as an important public health problem in many parts of the world, especially in developing countries [1]. In Iran, a country with a moderate rate of tuberculosis (TB), *Mycobacterium fortuitum*, *M. kansasii*, and *M. simiae* are the most common causes of NTM diseases, respectively [2, 3]. *M. simiae* is becoming an emerging pathogen in Iran and has been recognized as a causative agent of pulmonary and disseminated infections [4, 5]. Treatment of *M. simiae* disease is time consuming and often complicated [6, 7]. Based on previous studies, *M. simiae* exhibited the highest level of natural drug resistance to several anti-TB agents in vitro [7, 8]. Although the association between in vitro susceptibility and the in vivo treatment outcome has not been established for most drugs, treatment of individual patients should preferably be based on drug susceptibility testing (DST) results [7]. Given the absence of drug susceptibility patterns of clinical *M. simiae* isolates in Iran, we aimed to report the DST results of *M. simiae* isolated from clinical samples and discuss the implications of these findings for treatment regimens.

Materials and Methods

Samples and setting

Twenty five (31.2%) clinical isolates of *M. simiae* were included in this study. These samples were recovered from 80 patients with NTM pulmonary diseases who referred to Tehran regional TB reference laboratory (TRTB-RL) from March 2014 to January 2018. The mean age of patients was 51.6 ± 19.3 years. *M. simiae* was isolated from respiratory specimens (sputum, bronchoalveolar lavage). All clinical isolates form patients with NTM met ATS/IDSA diagnostic criteria for

inclusion and were considered as the causative pathogen for the disease [9]. TRTB-RL is well-equipped with biosafety level III facilities and is able to perform DST. The Ethics Committee of Shahid Beheshti University of Medical Sciences approved the study (approval number: IR.SBMU.RETECH.REC.1397.245).

Phenotypic and molecular Identification of M. simiae

Growth rate, macroscopic and microscopic morphological features, growth at different temperatures and also a set of biochemical tests including tween 80 hydrolysis, nitrate reduction, niacin production, arylsulfatase, urease production, tellurite reduction, salt tolerance ,and catalase production were used for identification of *M. simiae* [10].

For molecular identification, PCR restriction analysis (PRA) and sequencing of 16S rRNA were used as described previously [11, 12].

DST

For DST two methods were used, proportional assay and broth microdilution method. The following antibiotics were selected: isoniazid, rifampicin, ethambutol, streptomycin, amikacin, kanamycin, ciprofloxacin, ofloxacin and clarithromycin were selected. The stock solutions of each agent were prepared by dissolving pure powder of individual drugs in sterile distilled water or another suitable solvent.

Proportional DST

In proportional DST, resistance was expressed as the percentage of colonies that grew on critical concentrations of the drugs according to the guidelines established by the clinical and laboratory standards institute (CLSI) [13]. Interpretation was made according to the usual criteria for resistance, i.e., 1% for all drugs. *M. tuberculosis* H37Rv strain (ATCC 27294) was used for quality control testing in DST [14].

Broth microdilution

For each isolate, a suspension in broth was prepared from an agar medium with adequate growth of the isolate and the inoculum was standardized for susceptibility testing to a turbidity equivalent to a 0.5 McFarland standard. Suspensions were then diluted and inoculated into 96-well microtiter plates to achieve a final organism concentration from 1×10^5 to 5×10^5 CFU/ml. The antimicrobial agent minimum inhibitory concentrations (MICs) were interpreted according to the guidelines established by the CLSI [13]. Serial double dilutions of antimicrobial agents were prepared ranging from 0.06 to 512 mg/L, added to cation-supplemented Mueller–Hinton broth plus 5% OADC, and then 0.1 ml was dispensed into the wells of microdilution plates. Following inoculation, all cultures were incubated aerobically at 37 °C. Growth was examined weekly up to 4 weeks. The MIC was defined as the lowest concentration of the drug that inhibited visible growth. Quality control of MICs was performed by testing recommended reference strains, including *Enterococcus faecalis* ATCC 29212, *Mycobacterium peregrinum* ATCC 700686, *Pseudomonas aeruginosa* ATCC 27853, and *Staphylococcus aureus* ATCC 29213.

Data storage and analysis

Information and laboratory data were entered in Microsoft Excel 2017 software and were imported into SPSS (version 22; IBM) software. Frequency distributions were analyzed by use of the SPSS (version 22) software.

Results

Based on the results of proportional DST, all 25 (100%) isolates of *M. simiae* were resistant to isoniazid, rifampicin, ethambutol, streptomycin, amikacin, kanamycin, ciprofloxacin and clarithromycin (Table 1 and 2). They also were highly resistant to ofloxacin (80%). Susceptibility to ofloxacin was only noted in the 5 isolates. In vitro susceptibility to first-line anti-TB drugs was not found among clinical isolates of *M. simiae*. Acquired resistance, defined as a resistant follow-up isolate after a susceptible primary isolate, was noted in 10 patients; acquired rifampicin resistance was most frequent, followed by acquired and isoniazid resistance.

Discussion

M. simiae as an emerging pathogen in Iran accounts for more than 30.0% of all pulmonary diseases caused by NTM in our study. Based on the previous studies, 10 to 47% of the patients with NTM had true *M. simiae* infection [5, 15, 16]. We have recently indicated the clinical isolation of *M. simiae* by some studies in Iran, but the ecology behind this is not well understood [17]. We hypothesize that the high rate of *M. simiae* in Iran, may be due to the contamination of water supplies with this pathogen, a finding that was previously reported in studies conducted in Iran [18, 19]. Temperature and humidity have also been reported to be associated with NTM infections [16, 20]. Another explanatory factor may be the presence of underlying diseases among patients with *M. simiae* as described by Coolen-Allou [20].

Another issue raised by *M. simiae* pulmonary disease is its difficult treatment. Since little is known about drug susceptibility patterns of clinical isolates of *M. simiae* in Iran, there is little basis to predict potentially successful treatment regimens. Based on our study, all *M. simiae* species showed resistance to isoniazid, rifampicin, ethambutol, streptomycin, amikacin, kanamycin, ciprofloxacin,

clarithromycin and, to a lesser extent ofloxacin. Van Ingen et al in the Netherlands, showed resistant rates of 86% to amikacin, 77% to ciprofloxacin, 36% to moxifloxacin and 9% to clarithromycin in clinical isolates of *M. simiae* [7]. *M. simiae* isolates from Hamieh et al, study in the Lebanon were mostly susceptible to amikacin and clarithromycin and less susceptible to moxifloxacin and ciprofloxacin [16]. Likewise, in study conducted by Coolen-Allou et al, in the France, *M. simiae* isolates were found to have low susceptibility to antibiotics, except for amikacin, fluoroquinolones, and clarithromycin [20]. These findings indicate the variable drug susceptibility patterns of *M. simiae* in different geographic locations and emphasizes on the need to perform DST before initiating therapy.

This study has some limitations. First, the small numbers of evaluated patients and events might have reduced the power of the study. Second, the potential effect of predisposing factors could not be analyzed because of the limited information obtained from our data sources. The results of this single-center study cannot be extrapolated for the whole country.

In conclusions, clinical isolates of *M. simiae* were multidrug resistant, and had different drug susceptibility patterns than previously published studies. DST results can assist in selecting more appropriate treatment regimens. Newer drugs with proven clinical efficacy correlating with in vitro susceptibility should be substituted with first- and second line anti-TB drug testing. Further molecular studies on environmental and clinical isolates are needed to better assess the epidemiology and ecology of *M. simiae*.

Conflict of interest:

There is no conflict of interest.

Acknowledgements:

This study was supported by Shahid Beheshti University of Medical Sciences, Tehran, Iran.

References:

1. Prevots, D.R. and T.K. Marras, *Epidemiology of human pulmonary infection with non-tuberculous mycobacteria: a review*, in *Clinics in chest medicine*2015. p. 13.
2. Nasiri, M.J., et al., *Prevalence of non-tuberculosis mycobacterial infections among tuberculosis suspects in Iran: systematic review and meta-analysis*, in *PloS one*2015. p. e0129073.
3. Nasiri, M.J., et al., *High rates of nontuberculous mycobacteria isolation from patients with presumptive tuberculosis in Iran*, in *New microbes and new infections*2018. p. 12-17.
4. Hamieh, A., R. Tayyar, and S. Kanj, *Growing Knowledge about Mycobacterium simiae from Two Recent Studies.* Antimicrobial Agents and Chemotherapy, 2018. **62**(11): p. e01454-18.
5. Hashemi-Shahraki, A., et al., *Mycobacterium simiae: a possible emerging pathogen in Iran.* Japanese journal of infectious diseases, 2013. **66**(6): p. 475-479.
6. Philley, J.V. and D.E. Griffith, *Treatment of slowly growing mycobacteria.* Clinics in chest medicine, 2015. **36**(1): p. 79-90.
7. van Ingen, J., et al., *Drug susceptibility testing and pharmacokinetics question current treatment regimens in Mycobacterium simiae complex disease.* International journal of antimicrobial agents, 2012. **39**(2): p. 173-176.
8. Van Ingen, J., et al., *In vitro drug susceptibility of 2275 clinical non-tuberculous Mycobacterium isolates of 49 species in The Netherlands.* International journal of antimicrobial agents, 2010. **35**(2): p. 169-173.
9. Griffith, D.E., et al., *An official ATS/IDSA statement: diagnosis, treatment, and prevention of nontuberculous mycobacterial diseases*, in *American journal of respiratory and critical care medicine*2007. p. 367-416.
10. Kent, P.T., G.P. Kubica, and C.f.D. Control, *Public health mycobacteriology: a guide for the level III laboratory*. 1985: US Department of Health and Human Services, Public Health Service, Centers for Disease Control.
11. Telenti, A., et al., *Rapid identification of mycobacteria to the species level by polymerase chain reaction and restriction enzyme analysis.* Journal of Clinical Microbiology, 1993. **31**(2): p. 175-178.
12. Rogall, T., T. Flohr, and E.C. Böttger, *Differentiation of Mycobacterium species by direct sequencing of amplified DNA.* Microbiology, 1990. **136**(9): p. 1915-1920.
13. Woods, G.L., S.-Y.G. Lin, and E.P. Desmond, *Susceptibility test methods: mycobacteria, nocardia, and other actinomycetes*, in *Manual of Clinical Microbiology, 10th Edition*. 2011, American Society of Microbiology. p. 1215-1238.
14. Nasiri, M.J., et al., *Drug resistance pattern of Mycobacterium tuberculosis isolates from patients of five provinces of Iran.* Asian Pacific journal of tropical medicine, 2014. **7**(3): p. 193-196.
15. Van Ingen, J., et al., *Clinical relevance of Mycobacterium simiae in pulmonary samples.* European Respiratory Journal, 2008. **31**(1): p. 106-109.
16. Hamieh, A., et al., *Emergence of Mycobacterium simiae: A retrospective study from a tertiary care center in Lebanon.* PloS one, 2018. **13**(4): p. e0195390.
17. Nasiri, M.J., et al., *Mycobacterium simiae pulmonary disease in Iran: A systematic review and meta-analysis.* New microbes and new infections, 2018.

18.   Azadi, D., et al., *Species diversity and molecular characterization of nontuberculous mycobacteria in hospital water system of a developing country, Iran.* Microbial pathogenesis, 2016. **100**: p. 62-69.

19.   Khosravi, A.D., et al., *Prevalence of non-tuberculous mycobacteria in hospital waters of major cities of Khuzestan Province, Iran.* Frontiers in cellular and infection microbiology, 2016. **6**: p. 42.

20.   Coolen-Allou, N., et al., *Clinical, radiological, and microbiological characteristics of Mycobacterium simiae infection in 97 patients.* Antimicrobial Agents and Chemotherapy, 2018. **62**(7): p. e00395-18.

Table 1. In vitro resistance of the 25 *M. simiae* isolates to the tested antibiotics by proportional method.

Antibiotic	No. (%) of isolates		
	Susceptible	Intermediate	Resistant
Isoniazid	0 (0)	-	25 (100)
Rifampicin	0 (0)	-	25 (100)
Ethambutol	0 (0)	-	25 (100)
Streptomycin	0 (0)	-	25 (100)
Amikacin	0 (0)	-	25 (100)
Kanamycin	0 (0)	-	25 (100)
Ciprofloxacin	0 (0)	-	25 (100)
Ofloxacin	5 (20)	-	20 (80)
Clarithromycin	0 (0)	-	25 (100)

Table 2. In vitro resistance of the 25 *M. simiae* isolates to the tested antibiotics by broth microdilution method.

Antibiotic	Range	MIC (lg/ml)		No. (%) of isolates		
		MIC50	MIC90	Susceptible	Intermediate	Resistant
Isoniazid	16–128	32	64	0 (0)	-	25 (100)
Rifampicin	1-128	4	64	0 (0)	-	25 (100)
Ethambutol	4-64	16	32	0 (0)	-	25 (100)
Streptomycin	2-64	16	32	0 (0)	-	25 (100)
Amikacin	0.25-64	16	32	0 (0)	-	25 (100)
Kanamycin	0.25-64	16	32	0 (0)	-	25 (100)
Ciprofloxacin	0.25-64	8	32	0 (0)	-	25 (100)
Ofloxacin	1-64	8	32	5 (20)	-	20 (80)
Clarithromycin	32–128	32	64	0 (0)	-	25 (100)