Detection of classic and cryptic *Strongyloides* genotypes by deep amplicon

sequencing: A preliminary survey of dog and human specimens collected

from remote Australian communities

Meruyert Beknazarova[1]*, Joel L. N. Barratt[2,3]*, Richard S. Bradbury[2], Meredith Lane[2,4], Harriet

Whiley[1], Kirstin Ross[1]

* corresponding authors

[1] Environmental Health, Flinders University, South Australia

[2] Division of Parasitic Diseases and Malaria, Centers for Disease Control and Prevention

[3] Oak Ridge Associated Universities, Oak Ridge, Tennessee

[4] Synergy America Inc.

Abstract

Strongyloidiasis is a neglected disease caused by the human infective nematodes *Strongyloides stercoralis, Strongyloides fuelleborni* subsp. *fuelleborni* and *Strongyloides fuelleborni* subsp. *kellyi*. The zoonotic potential of *S. stercoralis* and the potential role of dogs in the maintenance of strongyloidiasis transmission has been a topic of interest and discussion for many years. In Australia, strongyloidiasis is prevalent in remote socioeconomically disadvantaged communities in the north of the continent. Being an isolated continent that has been separated from other regions for a long geological period, knowledge on the diversity of Australian *Strongyloides* genotypes is an important addition to our understanding of the genetic diversity within the genus. Using PCR enrichment combined with Illumina sequencing technology, we sequenced the *Strongyloides SSU* 18S rDNA hyper-variable I and hyper-variable IV regions using *Strongyloides*-specific primers, and a fragment of the mtDNA *cox*1 gene using primers that are broadly specific for *Strongyloides* sp. and hookworms. These loci were amplified from DNA extracted from human and dog faeces, and one human sputum sample. Using this approach, we confirm for the first time that potentially zoonotic *S. stercoralis* genotypes are present in Australia, suggesting that dogs represent a potential reservoir of human strongyloidiasis in remote Australian communities.

Author summary

Strongyloides stercoralis is a soil-transmitted nematode that causes the disease strongyloidiasis. Due to the autoinfective nature of this parasite, it can re-infect a host causing chronic infection. If not diagnosed and treated it can be highly detrimental to human health and has a high mortality rate. Strongyloidiasis is common in remote communities in the north of Australia and has been an issue for decades. Despite various successful intervention programs to treat human strongyloidiasis, the disease remains endemic in those communities. Here for the

40   first time we looked at the Australian dogs' potential to infect humans and found that they carry

41   two genetically distinct strains of *Strongyloides* spp., one of which also infects humans. This

42   supports the hypothesis that dogs are a potential source for human strongyloidiasis. We also

43   found that dogs in Australia might be carrying unique haplotypes. Whether these new

44   haplotypes are also human infective is to be confirmed by further research.

49   **Keywords:** *Strongyloides*, genotypes, dogs, Australia, zoonoses, hookworms, *Ancylostoma*

Introduction

Strongyloidiasis is a neglected disease caused by the human infective nematodes *Strongyloides stercoralis, Strongyloides fuelleborni* subsp. *fuelleborni* and *Strongyloides fuelleborni* subsp. *kellyi* (Grove, 1996). Worldwide, it is estimated to infect up to 370 million people and is predominately found in socioeconomically disadvantaged communities (Olsen et al., 2009, Bisoffi et al., 2013, Beknazarova et al., 2016). While *S. stercoralis* is a globally distributed nematode, *S. f. fuelleborni* has thus far only reported in Africa and Southeast Asia and *S. f. kellyi* from Papua New Guinea (PNG) (Thanchomnang et al., 2017, Pampiglione and Ricciardi, 1971, Ashford et al., 1992).

In Australia, strongyloidiasis is prevalent in remote communities located across the Northern Territory, Queensland, Western Australia, northern South Australia and northern New South Wales (Page et al., 2016). Current estimates indicate that infection rates in some communities are up to 41% or 60% based on microscopy or serology respectively (Heydon and Green, 1931, Sampson et al., 25-26 June 2003, Miller et al., 2018, Holt et al., 2017). Despite initially successful intervention programs targeting treatment to eliminate human strongyloidiasis in remote Australian communities, the disease remains endemic (Kearns et al., 2017, Page et al., 2006). Reappearance of the infection could possibly be as a result of zoonotic transmission from dog reservoirs given that dogs and humans share a close and intimate cultural bond in rural and remote Indigenous communities of Australia (Constable et al., 2010).

The zoonotic potential of *S. stercoralis* infected dogs and their potential role in the maintenance of strongyloidiasis transmission has been a topic of interest and discussion for many years (Beknazarova et al., 2017b, Goncalves et al., 2007, Takano et al., 2009). Molecular data from human and dog derived *S. stercoralis* are useful in understanding the nature of cross infection. There are two regions of the *S. stercoralis* nuclear and mitochondrial genome that are

considered to be conserved within the *Strongyloides* genus and can be used as markers for

molecular typing of *S. stercoralis*. Hyper-variable regions (HVR) I and IV of the small Subunit

(*SSU*) 18s ribosomal DNA and the cytochrome c-oxidase subunit 1 (*cox*1) gene of the

mitochondrial DNA (mtDNA) have been widely used to study relationships between *S.*

stercoralis from different hosts and different geographic locations (Hasegawa et al., 2009,

Hasegawa et al., 2010, Jaleta et al., 2017, Nagayasu et al., 2017, Basso et al., 2018). Based on

genetic analysis of these genetic loci, it has been recently found that there are two genetically

different *S. stercoralis* strains, one is dog and human infective, and the other is dog specific.

These data were collected from dogs and humans in Cambodia and Japan (Jaleta et al., 2017,

Nagayasu et al., 2017).

Being an isolated continent that has been separated from other regions for a long

geological period, Australia could represent an interesting addition to our understanding of the

genetic diversity within *S. stercoralis* and the *Strongyloides* genus more generally. Indigenous

Australians have inhabited the continent for at least 40,000 years and dogs (in the form of

dingoes) were likely introduced up to 12,000 years ago (Clutton-Brock, 1995). Given this long

period of relative isolation, it might be expected that Australia could harbor unique endemic

genotypes or unique sub-species of *S. stercoralis* that have evolved within dog and human

populations over this period.

Using PCR enrichment combined with Illumina sequencing technology, we sequenced

the *Strongyloides SSU* 18S rDNA hyper-variable I and hyper-variable IV regions using

Strongyloides-specific primers, and a fragment of the mtDNA *cox*1 gene using primers that are

broadly specific for *Strongyloides* sp. and hookworms. This approach was applied to DNA

extracted from human and dog faeces, and one human sputum sample. The main focus of this

study was to genotype Australian human and dog *S. stercoralis* strains to see whether dogs

carry human *S. stercoralis* strains and/or vice versa. To our knowledge this is the first time

human and dog *S. stercoralis* have been studied in Australia on a molecular level.

Methods

Study area and faeces collection

Dog faecal samples were collected from communities in the Northern Territory,

Australia. Dog faeces were collected from the environment (i.e., the ground) in the selected

communities and stored in the DESS (dimethyl sulfoxide, disodium EDTA, and saturated

NaCL) solution to preserve the DNA (Beknazarova et al., 2017a). For those dog faeces that

were collected from the private land, the consent forms were received to collect the samples.

The preserved faecal samples were express posted to the Environmental Health laboratory,

Flinders University, South Australia, for further analysis.

Human faecal and sputum samples were provided by our colleagues at the Royal

Darwin Hospital, NT, AusDiagnostics PTY LTD, NSW, and Townsville Hospital, Queensland.

While the personal information of the patients remains confidential and unknown, the infection

is known to have been locally acquired. Ethics approval from the Social and Behavioural

Research Ethics Committee (SBREC) No 6852 dated 1st June 2015 was obtained for collecting

dog faeces from the remote communities. Human ethical approval from the Southern Adelaide

Clinical Human Research Ethics Committee (SAC HREC) No 309.17 dated 24th January 2018

was obtained for comparing *S. stercoralis* DNA extracted from human and dog tissues.

DNA extraction

Prior to DNA extraction, faecal samples containing DESS were centrifuged for three

minutes at 3000 x g rpm using an Orbital 400 Clements (Phoenix, Lidcombe, Australia). The

supernatant consisting of the preservative solution was removed. The remaining faecal sample

was washed with sterile saline solution. DNA was extracted using the Power Soil DNA

isolation kit (QIAGEN, Hilden, Germany) following the manufacturer's instructions with slight

modifications that included incubating samples at 56ºC overnight after the cell lysis step,

followed by vortexing of samples for three minutes. Approximately 250 milligrams of the pellet

was placed into a PowerBead tube containing lysis buffer (included in the Power Soil DNA

extraction kit). The remainder of the extraction process was performed according to the

manufacturer's instructions. Extracted DNA was stored at -20°C prior to real-time PCR (qPCR)

analysis.

Real-time PCR

The samples (273 dog and 4 human DNA samples) were first screened for *Strongyloides*

spp. using qPCR in the Environmental Health laboratory at Flinders University, SA, Australia.

The real-time PCR assay was adopted from Verweij *et al.* (2009) using *S. stercoralis* - specific

primers targeting a 101 base pair region of 18S rRNA. The 20 µL reaction contained 10 µL

Supermix (SSoAdvanced, Universal Probes Supermix, Foster City, CA, USA), 1 µL primers

and probe mixture (Stro18S-1530F, Stro18S-1630R and Stro18S-1586T FAM) (Bio-Rad

Laboratories, CA, USA), 4 µL deionised H_2O, and 5 µL DNA template. All qPCR reactions

were performed in triplicate on the Corbett Rotor-Gene 6000 machine (QIAGEN, Hilden,

Germany). *S. stercoralis* primers and probes and qPCR conditions used are shown in Table 1.

A sample was considered positive when the Ct value was lower than the mean negative Ct

minus 2.6 standard deviations of a mean negative control Ct. Positive samples were amplified in

every qPCR reaction.

Conventional PCR for amplification of *SSU* HVR-I and HVR-IV, and *cox*1 sequences

Extracted DNA from samples that were qPCR positive for *Strongyloides* spp. (47 dog

and four human DNA samples) were shipped on dry ice to the Centers of Disease Control and

Prevention (CDC), Georgia, USA for conventional PCR, sequencing and bioinformatics

analysis. Hyper-variable regions (HVR) I and IV in the small Subunit (*SSU*) 18S ribosomal

DNA and a fragment of the mitochondrial cytochrome c-oxidase subunit 1 (*cox*1) gene were

amplified using conventional PCR and then sequenced using Illumina technology. All PCR

reactions were performed on a GeneAmp PCR System 9700 Thermo Cycler, version 3.12

(Applied Biosystems, USA). *S. stercoralis* primers and PCR conditions used for qPCR and

conventional PCR are shown in Table 1. For the *cox*1 gene, PCR reactions were performed in a

total volume of 50 µL containing 10 µL NEB 5X Q5® Buffer (New England BioLabs, USA),

10 µL NEB 5X Q5® High GC Enhancer (New England BioLabs, USA), 4 µL NEB

Deoxynucleotide Solution Mix (10 mM each nt) (New England BioLabs, USA), 1 µL Q5®

High-Fidelity DNA Polymerase (New England BioLabs, USA), 2.5 µL forward primer

(SSP_COX1_F), 2.5 µL reverse primer (SSP_COX1_R), 18 µL deionised H_2O, and 2 µL DNA

template. For the HVR-I and HVR-IV regions, PCR reactions were performed in a 25 µL

reaction containing 12.5 µL of NEBNext® Q5® Hot Start HiFI PCR Mastermix, MO543L

(New England BioLabs, USA), 1.5 µL forward primer (NEW_HVR_I_F or NEW_HVR_IV_F),

1.5 µL reverse primer (NEW_HVR_I_R or NEW_HVR_IV_R), 7.5 µL deionised H_2O, and 2

µL DNA template.

164   *Insert Table 1 here*

The amplified PCR products were separated by 1.5% agarose gel electrophoresis and stained

with ethidium bromide. The stained DNA bands were visualised by UV illumination using a

Ugenious 3 (SYNGENE, Japan). For quality control, each PCR run was accompanied by a

positive control containing *Strongyloides* genomic DNA as template, a non-template control

containing autoclaved sterile water instead of template, and a negative control containing DNA

extracted from a parasite-free specimen. Amplicons for each of the three markers were also

generated for *Strongyloides ratti* as an additional control for the sequencing and *in silico*

analysis steps.

Next Generation Sequencing (NGS)

Ten microliters of PCR amplicon was purified and normalized for concentration prior to

library preparation using SequalPrep Normalization Plate Kit (Thermo Fisher Scientific, USA).

DNA libraries were prepared using the NEBNext Ultra DNA Library Prep Kit for Illumina

(New England BioLabs, USA), and NEBNext Multiplex Oligos for Illumina Index kit (New

England BioLabs, USA). The sequencing reactions were prepared using the MiSeq reagent

Nano Kit v2 (PE250bp), and performed on the Illumina MiSeq platform (Illumina).

In silico **analysis**

The Illumina reads were analyzed using Geneious (www.geneious.com) by means of a

workflow that performed read quality control, assembly of contigs and genotype assignment. As

part of this workflow, quality trimming to a minimum phred score of 20 and removal of adapter

sequence was performed using BBDuk (v 37.64). Reads less than 50 bases in length were

discarded. Paired reads were then merged using BBMerge (v 37.64) and all reads (merged and

unmerged) were mapped to a reference sequence. For *cox*1 amplicons, reads were mapped to a

S. stercoralis sequence with the GenBank (GB) accession number LC050212.1. For *SSU* HVR-

I and *SSU* HVR-IV, reads were mapped to a *S. stercoralis* with the accession AF279916.2.

Prior to mapping, each reference sequence was trimmed to the length of the amplicon,

excluding the primer sequences. Mapping was performed using the Geneious mapper under the

Medium sensitivity / Fast default settings. Reads that successfully mapped (and their mates)

were retained for *de novo* assembly. These reads were assembled using the Geneious *de novo*

assembler with the following customized parameters; minimum overlap: 50 bases, minimum

overlap identity: 100%, maximum number of mismatches per read: 0%, and the maximum

number of ambiguities: 1. Contigs were split if coverage fell below 50 and sub-variants with

coverage less than 50 were not considered. Contigs were trimmed to the primer sequence using

the 'Trim Ends' function in Geneious. Contigs were validated by taking the trimmed and

merged reads and mapping them back to the contigs generated using the Geneious mapper, and

the following custom parameters: minimum mapping quality: 30, minimum overlap: 150 bases,

minimum overlap identity: 100%, maximum number of mismatches: 0%, and the maximum

number of ambiguities: 1. Contigs were split or discarded if the coverage fell below 300 to 500

depending on the depth obtained for a particular specimen (500 for specimens that obtained

>10k reads, 300 for specimens with <10k reads). Finally, genotypes were assigned by

performing a local BLASTN search (within Geneious) against a database constructed from all

unique *Strongyloides* sp. 18S and *cox*1 sequences available in GenBank and the DNA Databank

of Japan. A set of homologous sequences from several other roundworms (parasitic and free-

living) were also included in this database. Sequences were only added to the BLASTN

database if they overlapped our 217 base pair *cox*1 amplicon by more than 95%. This included

several previously published *cox*1 haplotypes that had matching *SSU* HVR-I and HVR-IV

genotypes assigned (Jaleta et al., 2017).

Any specimens for which *Strongyloides ratti* sequences were detected as part of this

workflow were considered to be at risk of cross contamination from our positive control and

potentially from other specimens included in the study. Any contaminated specimens were

excluded from further analysis.

Construction of a *cox*1 cluster dendrogram

Sequences of *cox*1 were exported from Geneious as a fasta file and were aligned using

the 'msa' package in R. The 'dist.alignment' function from the 'seqinr' package was used to

compute a pairwise identity matrix, considering gaps in the identity measure. The resulting

matrix was clustered using the agglomerative nested clustering approach performed with the

'agnes' R package, using "euclidean" distances and the "average" clustering method. A cluster

dendrogram was generated using the 'ggtree' R package. Vector images (i.e. graphics) used for

annotation of the dendrogram were either generated in house at CDC or obtained from PhyloPic

(http://phylopic.org).

Results

Real-time PCR, conventional PCR and sequencing

We screened 273 dog and four human samples using real-time and conventional PCR.

47 dog and four human samples that were positive by qPCR were selected for sequencing of

their *cox*1, *SSU* HVR-I and HVR-IV regions. In some cases, useable sequences were not

generated due to poor amplification of the PCR product. Sequence data was obtained for 24

specimens including four human specimens and twenty dog specimens. The complete genotype

of all specimens analyzed in this study is shown in Figure 1 and summarized in Table 2.

Insert Figure 1 here

Insert Table 2 here

238   *SSU* HVR-I haplotypes detected among *S. stercoralis* from Australian dogs and humans

Jaleta *et al.* (2017) previously sequenced the *SSU* HVR-I region of *S. stercoralis* worms

from Cambodian dog and human specimens and identified five different haplotypes (HVR-I

haplotypes I-V) (Jaleta et al., 2017). A recent study of European dogs identified a new

haplotype from the HVR-I region (haplotype VI) (Basso et al., 2018). In our Australian samples

we found haplotype II in both human and dog samples and haplotype IV in dog samples only,

which is consistent with the findings from Jaleta *et al.* (2017). We also identified haplotype VI

in a single Australian dog. Following the genotype nomenclature developed by Jaleta *et al.*

(2017) and Basso *et al.* (2018), in addition to these six haplotypes previously described (Jaleta

et al., 2017, Basso et al., 2018), we discovered two new HVR-I haplotypes; haplotypes VIII and

X (Figure 2, Table 1). Due to the existence of noteworthy similarities (> 99% in all cases)

between sequences of *S. stercoralis*, *Strongyloides procyonis*, a sequence assigned to

Strongyloides sp. Okayama (GB: LC038066.1), and our novel dog sequences, we expanded the

Jaletta *et al*. typing scheme to include these sequences. This involved inclusion of haplotypes

that could not be confidently assigned to *S. stercoralis* given the information on hand, yet are

highly similar to known *S. stercoralis* 18S haplotypes. This adjustment was also required

because a sequence attributed to *S. procyonis* (GB: AB272234.1) possesses HVR-I haplotype

IV, which is identical to an *S. stercoralis* genotype assigned to a Cambodian dog (GB:

KU724124.1). For details, refer to Figure 2 and Table 3.

Insert Figure 2 here

Insert Table 3 here

***SSU* HVR-IV haplotypes detected among *S. stercoralis* from Australian dogs and humans**

In the Australian samples we identified HVR-IV haplotype A in both humans and dogs,

and haplotype B only in dogs, as previously observed (Jaleta et al., 2017). Supporting the

findings of Jaleta *et al*. (2017), our results also showed that haplotype II of HVR-I is found in

combination with haplotype A of HVR-IV, and haplotype IV of the HVR-I region is only found

in combination with haplotype B of the HVR-IV region (Jaleta et al., 2017). We observed that a

unique sequence attributed to *S. stercoralis* had been submitted to GenBank in 1993, and this

was assigned to haplotype C due to its historic value (GB: M84229.1). Given the strong

similarity between HVR-IV sequences of *S. procyonis* and *S. stercoralis* and the fact that HRV-

I haplotye IV (Jaleta *et al*. 2017) is also found in *S. procyonis* SSU sequences, we assigned the

HVR-IV sequence from *S. procyonis SSU* DNA to haplotype D (GB: AB272234.1 and

AB205054.1). A HVR-IV genotype 99% similar to the *Strongyloides* sp. Okayama (GB:

LC038066.1) was detected in Dog 45. Therefore, the HVR-IV sequence of *Strongyloides* sp.

Okayama was assigned to haplotype H, and the sequence from Dog 45 was assigned to

Haplotype I. Consequently, four new haplotypes detected in four Australian dog samples were

assigned to HVR-IV haplotypes E, F, G and I (Figure 1, Table 4).

278   *Insert Table 4 here*

Clustering of *Strongyloides stercoralis* based on *cox*1 sequences

A 217 base pair fragment of *cox*1 was sequenced from 20 Australian specimens

including those from 18 dogs and two humans, plus the *S. ratti* control (21 *cox*1 sequences in

total). Multiple *cox*1 types were obtained from a single specimen in many cases, revealing

infections caused by multiple helminth species and multiple *S. stercoralis* genotypes in a single

host (Figure 1). Dendrogram construction by agglomerative nested clustering revealed three

distinct *S. stercoralis* clades, including one occupied predominantly by worms possessing the

II/A *SSU* genotype, which constituted sequences obtained from dogs and humans. Four *cox*1

sequences obtained in this study (one from each of human 1, human 3, dog 6, and dog 18), were

assigned to the dog and human-infecting *S. stercoralis* clade. A *S. stercoralis* clade occupied

mostly by specimens possessing the I/B and V/B *SSU* genotypes was also apparent (one

specimen possessed the IV/B genotype), representing dog infections only (dog clade 1). None

of the Australian specimens were assigned to this clade (Figure 1). A final *S. stercoralis* clade

containing *cox*1 sequences obtained from only dogs (dog clade 2) was also dominated by

specimens possessing the I/B and V/B genotypes, though two specimens were also assigned the

IV/B. A single *cox*1 sequence from each of dogs 6 and 7 was assigned to this clade (Figure 1).

Cryptic *cox*1 sequences potentially from *Strongyloides* sp. helminths that could not be

assigned to a genus or species

Sequences were obtained from two dog fecal specimens (dogs 22 and 43) that potentially

belong to a *Strongyloides* sp. helminth but could not be confidently assigned to a species given

the information available. Two *cox*1 sequences were obtained for dog 22. One of these clustered

with a *cox*1 sequences from *Strongyloides mirzai* (GB: AB526307.1), a helminth that infects a

Japanese pit viper. The second sequence from dog 22 clustered between the *S. mirzai* clade and

the *S. fuelleborni* clade yet also clustered in a position immediately basal to all hookworms

(Figure 1). This sequence also obtained a nearest BLASTN hit to a *Necator* sp. sequence (GB:

AB793563.1), though its next best hit based on an online BLASTN search was to a sequence

from *S. stercoralis* (GB: LC179452.1). Three unique *cox*1 sequences were obtained from dog

43 (GB: MK434220, MK434221; MK434222), one clustering with *S. mirzai* and a second

clustering with two sequences of the *Strongyloides* sp. 'loris' clade (GB: LC197958.1,

LC197946.1). When submitted to an online BLASTN search, the sequence clustering alongside

S. mirzai also obtained top hits to free living nematodes (*Ektaphelenchus* sp. and

Bursaphelenchus populi, GB: JX979197.1 and HQ699854.1 respectively), yet several of its top

hits were also to *S. fuelleborni cox*1 sequences. The third sequence from dog 43 clustered

between the *Strongyloides* sp. 'loris' clade and a clade containing all *S. stercoralis* sequences

(Figure 1).

Mixed genotype infections with *Strongyloides stercoralis*

Two samples had mixed *S. stercoralis* infections where a complete genotype was

obtained (a sequence for *cox*1, HVR-I and HVR-IV); dog 6 and dog 18. When examining the

number of reads that mapped to each haplotype for these specimens (not shown), for dog 6

approximately 20% of reads were assigned to haplotype II and 80% to haplotype IV for the

HVR-I region. For the HVR-IV region, approximately 20% of reads were assigned to haplotype

A and 80% to haplotype B. With this information it was deduced that this dog was infected with

two strains of *S. stercoralis*, one from the human / dog clade (genotype II/A) and another from a

dog-specific clade (genotype IV/B). Interestingly, two *cox*1 sequences were obtained from this

dog, one assigned to the human / dog clade and another assigned to dog clade 2, supporting our

deduction. Dog 18 was also infected with two types of *S. stercoralis*, with a genotype of II + VI

/ A + E assigned to this specimen. Given that the number of reads assigned to each of these

types fell between 40% and 50%, it is difficult to link the HVR-I types identified here to their

corresponding HVR IV type. It should also be noted that while the specimen from dog 18

possessed two 18S genotypes, evidence was only found for a single *cox*1 sequence. There were

two *S. stercoralis* strains found in the HVR-IV region of the dog 13. Approximately 50% of

reads were assigned to the haplotype A and 50% to a new haplotype G.

Detection of non-*Strongyloides* sp. *cox*1 sequences and mixed helminth infections

Twenty-three *cox*1 sequences were attributed to *Ancylostoma* spp. and one dog (dog 6)

was infected with an *Ancylostoma* sp. clustering closely with *Anyclostoma ceylanicum* (Figure

1, cyan clade), and two distinct genotypes of *S. stercoralis*. Sequences were obtained from dogs

14, 15 and 16 that belong to a *Metastrongylus*-like helminth, possibly *Metastrongylus salmi*.

Two sequences attributed to *Ancylostoma caninum* were also obtained from dog 15, and a

fourth sequence belonging to an *Ancylostoma* sp. was also detected in this dog. A sequence was

obtained from dog 37 that obtained BLASTN hits to *S. fuelleborni* sequences (e.g., GB:

AB526303.1, 86.2% identity). Agglomerative nested clustering placed this sequence in a

position basal to all *Strongyloides* and hookworm sequences included in this analysis. This *cox*1

sequence also obtained close BLASTN hits (87% identity) to *Aphelenchoides* sp. (GB:

KX356839.1) and *Bursaphelenchus luxuriosae* (AB097863.1) which are free living

mycophagous and/or potentially plant parasitic nematodes. The *cox*1 from dog 28 does not

appear to be helminth in origin and most closely resembles a *cox*1 sequence from a rotifer;

Macrotrachela quadricornifera (GB: JX184003.1), which served as a convenient outgroup for

the clustering analysis (Figure 1).

The presence of *Strongyloides* and other helminths in remote Australian communities

Strongyloides stercoralis was detected in 10 specimens from seven remote communities in

Australia by genotypic analysis. While all specimens included in this study tested positive for *S.*

stercoralis using a published real-time PCR assay (Verweij *et al.* 2009), deep sequencing found

no evidence of *S. stercoralis* infection in several cases where instead, an infection with another

helminth (usually *Ancylostoma* spp.) was confirmed. Some of these sequences clustered closely

to *A. ceylanicum* yet appeared to be distinct. However, it should be noted that these sequences

were identical to a published sequence assigned to *A. caninum* (GB: AJ407962.1) which did not

overlap completely with our amplicon (50 bases short), so we cannot be sure they possess the

same sequence type. This necessitates the conservative assignment of these sequences to

Ancylostoma sp. Alternatively, most non-*Strongyloides* spp. sequences obtained from dogs were

more confidently assigned to *Ancylostoma caninum* (Figure 2). Two dogs from community 2

were also infected with *S. stercoralis* while in one dog from community 5, a cryptic

Strongyloides sp. possessing a unique *SSU* genotype for both HVR-I and HVR-IV was detected

(dog 22, genotype VIII/F). Interestingly, a *Metastrongylus*-like *cox*1 sequence was detected in

all dogs from community 3 that were tested, and a single dog (dog 15) from this same

community was infected with *A. caninum* and at least one other *Ancylostoma* sp. (Figure 2, cyan

clade). We propose that sequences obtained from community 6 are from environmental

organisms, possibly representing extraneous contaminants given that one represents a rotifer-

like sequence (dog 28) (GB: MK434225) and the other obtained BLASTN hits to free-living

nematodes (dog 37) (GB: MK434224). Community 4 and community 7 are represented by a

single typed specimen each (dog 18 and dog 45 respectively), that include unique sequences

from a helminth that we can confidently assign to the genus *Strongyloides* (Figure 1).

Discussion and conclusions

In a recent study on *S. stercoralis* from humans and dogs in Cambodia, two genetically

distinguishable *S. stercoralis* populations were identified based on the HVR-IV region. The

HVR-IV haplotype A strain was found to be dog and human infective, while HVR-IV

haplotype B strain was shown to be dog specific (Jaleta et al., 2017). In our study we observed

the same phenomenon, that HVR-IV haplotype A is associated with strains infective for both

humans and dogs, while HVR-IV haplotype B is restricted to strains that are only infectious to

dogs. Supporting earlier findings, our results also showed that haplotype II of HVR-I is found in

combination with haplotype A of the HVR-IV region, and haplotype IV of the HVR I region is

only found in combination with haplotype B, which is specific to dogs. One dog possessing the

HVR-I haplotype IV (dog 6) had a mixed *S. stercoralis* infection, presumably with worms of

the genotype II/A (a type infectious to dogs and humans) and others with the genotype IV/B (a

dog specific genotype). The detection of two *cox*1 sequences that cluster in the dog / human and

dog specific clades respectively supports this assessment. In an Australian dog, we also

identified HVR-I haplotype VI which has only been previously reported in European dogs.

Interestingly, this dog (dog 18) also had a mixed genotype infection that included a novel *S.*

stercoralis HVR-IV genotype (haplotype E), that was linked to a *cox*1 sequence clustering in

the dog / human *S. stercoralis* clade.

In agreement with previous reports, the current study demonstrated that the *SSU* HVR

IV region in the *SSU* rDNA can be used to detect within species differences that correspond

with the genetic clades that appear when the same specimens are analyzed at the *cox*1 locus

(Hasegawa et al., 2016, Jaleta et al., 2017) (Figure 1). To support analysis of the *cox*1 locus by

deep sequencing, the *cox*1 PCR assay described here was designed so that merged paired-end

Illumina reads span the entire length of the amplicon. This greatly reduces the complexity of *in*

silico analysis when mixed *cox*1 genotypes are encountered. A trade-off of using a short

amplicon for this analysis is that it may capture less diversity. Additionally, short sequences are

of limited use in phylogenetic analysis. However, it should be noted that phylogenies are only

truly relevant for constructing the evolutionary history of taxa and because evolutionary

analysis was not the objective of this study, agglomerative nested clustering was used to group

*cox*1 sequences based on their pairwise sequence identity. Despite its limitations, our *cox*1

assay clearly resolved the dog and human infective *S. stercoralis* genotypes into a clade that is

distinct from the dog-specific types, which was our primary objective, allowing us to compare

our results to those obtained in previous studies. Furthermore, we show that the *cox*1 fragment

amplified here clearly distinguishes *S. stercoralis*-derived *cox*1 amplicons from other helminth

species including *S. fuelleborni* and multiple species of hookworm.

As part of this study, we modified the typing scheme initially reported by Jaleta *et al*.

(2017) to include certain *SSU* sequences previously published in GenBank in order to

accommodate our novel genotypes. The rationale for these additions was several fold (Table 2

and Table 3). Firstly, we observe that a sequence published in 1993 by Putland *et al*. (1993)

(GB: M84229.1) was distinct from other *S. stercoralis* sequences that have been mentioned in

the literature since, differing from HVR-IV haplotype A by one SNP at position 125 (Figure 2)

(Putland et al., 1993). Consequently, the HVR-IV region of this sequence was added to the

typing scheme as haplotype C due to its historic value (Figure 2). The HVR-I sequence of

M84229.1 is so drastically different to that of other *S. stercoralis* haplotypes (and to that of any

other *Strongyloides* spp. in general), and given that it has not been reported in the literature

since its initial publication, it was not added to the typing scheme. Next, we observed that *S.*

stercoralis SSU HVR-I haplotype IV (reportedly a dog-specific type) is also found in sequences

assigned to *S. procyonis* from a Japanese badger (GB: AB272234.1). To reconcile this

observation, we incorporated the HVR-IV region of sequences assigned to *S. procyonis* into the

typing scheme, referring to them as haplotype D (Table 3, GB: AB272234.1 and AB205054.1).

This also meant that the HVR-I sequence of the *S. procyonis SSU* (GB: AB205054.1) became

HVR-I haplotype VII. Hence, the new *SSU* HVR-I genotype from dog 22 (GB: MK468661)

became type haplotype VIII. A novel SSU HVR-I genotype from dog 45 was also discovered as

part of this study (GB: MK468662), and its sequence was most similar to the *SSU* HVR-I region

from *Strongyloides* sp. Okayama, isolated from a Japanese striped snake (GB: LC038066.1). As

this sequence was already in GenBank prior to the commencement of this study, the HVR-I

region of LC038066.1 was assigned to haplotype IX, while the novel sequence obtained from

dog 45 was assigned to haplotype X (GB: MK468662). As haplotypes A to D for HVR-IV had

been assigned to other sequences, the novel genotypes discovered in dogs 18, 22 and 13 were

assigned to HVR-IV haplotype E, F and G respectively (GB: MK468674, MK468675, and

MK468676). Finally, the HVR-IV sequence from *Strongyloides* sp. Okayama

(GB:LC038066.1) was assigned to haplotype H because it obtained a nearest match to the

HVR-IV regions sequenced from dog 45, which was consequently assigned to haplotype I (GB:

MK468677). Also note that all HVR-I and HVR-IV types discussed above (both novel and

previously published) are more similar to each other than they are to the corresponding *SSU*

regions from *S. ratti*. Consequently, they do not provide enough information on their own to

make confident species assignments (Figure 2).

Ultimately, while the typing scheme developed by Jaleta *et al.* (2017) was originally

designed to consider *S. stercoralis* genotypes alone, the detection of several novel cryptic

genotypes that: (1) cannot be confidently assigned to a species, (2) are nonetheless greater than

99% similar to each other and to known *S. stercoralis* genotypes and (3), are genotypes detected

in the same host (dogs), means that the adjustments made here represent the most

straightforward solution to the issue at hand. Moreover, being an isolated continent, it is

possible that Australian dogs might be infected with genetically distinct *S. stercoralis* strains

(Cawood and Korsch, 2008). Consequently, it is not unreasonable to suggest that some of the

cryptic dog *Strongyloides* genotypes described herein (i.e., from dogs 18, 22, 13 and 45) might

be attributable to truly novel *S. stercoralis* genotypes restricted to Australia. However, a larger

sample number is needed along with additional sequences and morphological analysis of

multiple specimens before these sequences can be assigned to a species of helminth. Similarly,

it is plausible that the *cox*1 sequences obtained from dog 43 are attributable to a helminth

belonging to the genus *Strongyloides*, yet the lack of any *SSU* sequences associated with this

specimen makes it difficult to draw any solid conclusions in that regard.

This study employed an alteration of the Jaleta *et al.* (2017) genotyping assay developed

at the Centers for Disease Control and Prevention (J. Barratt) for adaptation to NGS

technologies. The assay was designed to genotype *Strongyloides* sp. and potentially detect

mixed helminth infections (e.g., hookworm and *Strongyloides*) when applied directly to DNA

extracted from faeces and other biological specimens. This method has great advantages over

previous genotyping techniques in that it can be undertaken directly from faecal DNA extracts

and does not require culture of larvae for extraction of DNA from individual worms.

Furthermore, the depth of sequencing provided by NGS allows the detection of all genotypes in

a single sample (Zahedi et al., 2017).

This study had a number of limitations including the collection of dog stool from the

environment where it could have potentially become contaminated with extraneous

environmental organisms or their DNA. As noted in Table 1, the *cox*1 PCR employed in this

study also detects multiple hookworm species, and was even found to amplify the *cox*1

sequence of a *Metastrongylus*-like helminth. This may represent an advantage of the method if

simultaneous detection and genotyping of multiple pathogenic intestinal nematodes from dogs

and humans is required, although results such as this should be viewed with caution. Given the

sensitivity of deep sequencing, we suspect that detection of *cox*1 sequences resembling those of

Metastrongylus salmi could be attributable to the consumption of pig offal by dogs in

community 3. While *Metastrongylus* sp. are known to occasionally infect other species

including humans (Calvopina et al., 2016), the genus is generally thought to be specific to pigs.

We also note that the genotype from dog 45 closely resembles that of a reptile-infecting

Strongyloides sp. and its presence in a dog could be due to consumption of reptiles or reptile

feces by the dog. Consequently, these cases may represent an incidental finding rather than true

infections with a *Metastrongylus*-like helminth or a *Strongyloides* sp. resembling those found in

reptiles. Similarly, the *cox*1 assay described here detected DNA from potentially free – living

nematodes. A sequence obtained from dog 37 received a BLASTN hit to *S. fuelleborni* (GB:

AB526306.1), though with only 86.2% identity. However, agglomerative nested clustering

placed this sequence in a position basal to all *Strongyloides* and hookworm sequences included

in this analysis. This sequence also obtained close BLASTN hits (87% identity) to

Aphelenchoides sp. (GB: KX356839.1) and *Bursaphelenchus luxuriosae* (AB097863.1) which

are mycophagous and/or potentially plant parasitic nematodes. This sequence could therefore

represent a free-living nematode that came into contact with the fecal specimens in the

environment between when the stool was passed and collected. Surprisingly, a sequence similar

to one obtained from a rotifer; a free-living extremely distant relative of nematodes, was

detected in the specimen from dog 28 using this assay. This also likely represents contamination

of the stool specimen from the local environment prior to its collection.

Several unique sequences assigned to *Strongyloides* spp., *Ancylostoma* spp.,

Metastrongylus sp. and some unknown helminth species were detected for the first time using

the technology described here. We propose that some of the cryptic *Strongyloides* sp.

genotypes we encountered are potentially unique to the Australian continent and may have

diverged from southeast Asian *Strongyloides* populations as a result of vicariance. Discovery of

a *cox*1 sequence that clusters most closely to a *Strongyloides* sp. identified from a slow loris

might support this (dog 43, GB: MK434221), given that lorises are endemic to southeast Asia.

Aside from these cryptic types, several examples of the classic *S. stercoralis* genotypes

described by Jaleta *et al.* (2017) were detected. Consequently, our study was able to

independently support previous reports of at least two genetically distinct groups of *S.*

stercoralis; one infecting both dogs and humans and another group that is specific to dogs.

While this does not prove the potential of dogs as a reservoir of *S. stercoralis*, it supports the

hypothesis of zoonotic transmission in remote Australian communities. As discussed previously

and with respect to the One Health approach, we suggest that humans and dogs should be

treated concomitantly in these communities to control strongyloidiasis (Beknazarova et al.,

2017b, Jaleta et al., 2017). Ultimately, we confirm for the first time that potentially zoonotic *S.*

stercoralis genotypes are present in Australia and suggest that dogs represent a potential

reservoir of human strongyloidiasis in remote Australian communities.

Author Contributions

MB: Designed study, obtained ethics approvals, collected specimens, extracted DNA, ran real-

time PCR, optimized and ran PCR assays for *SSU* HVR-I, HVR-IV and *cox*1, performed

Illumina library preparation, performed *in-silico* analysis, data analysis and interpretation, wrote

and reviewed draft manuscripts; JB: Designed PCR assay for *SSU* HVR-I, HVR-IV and *cox*1,

performed *in silico* analysis and data interpretation, constructed cluster dendrogram, wrote and

reviewed manuscript drafts; RB: Designed study, reviewed manuscript drafts, obtained funding,

reviewed manuscript drafts; ML: Carried out PCR product normalization, prepared Illumina

libraries for deep sequencing, and sequenced amplicons; HW: Provided PhD supervision of

MB, designed study, obtained ethics approvals, obtained funding, reviewed manuscript drafts;

KR: Provided PhD supervision of MB, designed study, obtained ethics approvals, obtained

funding, reviewed manuscript drafts.

Acknowledgments

The authors would like to thank Professor Robert W. Baird and Dr Richard Sullivan at the

Royal Darwin Hospital, Darwin Pathology, NT and Dr Gemma Robertson at the Health Support

Queensland, QLD for providing us human *S. stercoralis* DNA samples We would also like to

acknowledge the contribution of Animal Management in Rural and Remote Indigenous

Communities, Environmental Health Branch at the Department of Health, NT and all the other

lovely veterinarians across Australia for helping us collecting dog faecal samples in the remote

communities. We acknowledge the help of Dr Rogan Lee, Dr Matthew Watts, John Clancy and

Vishal Ahuja at the Westmead Hospital, NSW with sending us *S. ratti* infected rat faeces. We

wish to thank staff at the Centers for Disease Control and Prevention in Atlanta, USA.

The work has been supported by the Australian Government Research Training Program

Scholarship and Flinders University Overseas Travelling Fellowship.

References

531 ASHFORD, R., BARNISH, G. & VINEY, M. 1992. *Strongyloides fuelleborni kellyi*: Infection and disease in
532   Papua New Guinea. *Parasitology Today,* 8**,** 314-318.
533 BASSO, W., GRANDT, L.-M., MAGNENAT, A.-L., GOTTSTEIN, B. & CAMPOS, M. 2018. *Strongyloides*
534   *stercoralis* infection in imported and local dogs in Switzerland: from clinics to molecular genetics.
535   *Parasitology research,* 118**,** 255-266.
536 BEKNAZAROVA, M., MILLSTEED, S., ROBERTSON, G., WHILEY, H. & ROSS, K. 2017a. Validation of
537   DESS as a DNA Preservation Method for the Detection of *Strongyloides* spp. in Canine Feces.
538   *International Journal of Environmental Research and Public Health,* 14**,** 624.
539 BEKNAZAROVA, M., WHILEY, H. & ROSS, K. 2016. Strongyloidiasis: A Disease of Socioeconomic
540   Disadvantage. *International Journal of Environmental Research and Public Health,* 13**,** 517.
541 BEKNAZAROVA, M., WHILEY, H. & ROSS, K. 2017b. Mass drug administration for the prevention human
542   strongyloidiasis should consider concomitant treatment of dogs. *PLoS Neglected Tropical Diseases,* 11**,**
543   e0005735.
544 BISOFFI, Z., BUONFRATE, D., MONTRESOR, A., REQUENA-MÉNDEZ, A., MUÑOZ, J., KROLEWIECKI,
545   A. J., GOTUZZO, E., MENA, M. A., CHIODINI, P. L. & ANSELMI, M. 2013. *Strongyloides stercoralis*:
546   a plea for action. *PLoS Neglected Tropical Diseases,* 7**,** e2214.
547 CALVOPINA, M., CABALLERO, H., MORITA, T. & KORENAGA, M. 2016. Human pulmonary infection by
548   the zoonotic *Metastrongylus salmi* Nematode. The first reported case in the Americas. *The American*
549   *Journal of Tropical Medicine and Hygiene,* 95**,** 871-873.
550 CAWOOD, P. A. & KORSCH, R. 2008. Assembling Australia: Proterozoic building of a continent. *Precambrian*
551   *Research,* 166**,** 1-35.
552 CLUTTON-BROCK, J. 1995. Origins of the dog: domestication and early history. *The domestic dog: Its evolution,*
553   *behaviour and interactions with people***,** 7-20.
554 CONSTABLE, S., DIXON, R. & DIXON, R. 2010. For the love of dog: the human–dog bond in rural and remote
555   Australian indigenous communities. *Anthrozoös,* 23**,** 337-349.
556 GONCALVES, A., MACHADO, G., GONCALVES-PIRES, M., FERREIRA-JUNIOR, A., SILVA, D. &
557   COSTA-CRUZ, J. 2007. Evaluation of strongyloidiasis in kennel dogs and keepers by parasitological and
558   serological assays. *Veterinary Parasitology,* 147**,** 132-139.
559 GROVE, D. I. 1996. Human strongyloidiasis. *Advances in Parasitology.* Elsevier.
560 HASEGAWA, H., HAYASHIDA, S., IKEDA, Y. & SATO, H. 2009. Hyper-variable regions in 18S rDNA of
561   *Strongyloides* spp. as markers for species-specific diagnosis. *Parasitology Research,* 104**,** 869-874.

HASEGAWA, H., KALOUSOVA, B., MCLENNAN, M. R., MODRY, D., PROFOUSOVA-PSENKOVA, I., SHUTT-PHILLIPS, K. A., TODD, A., HUFFMAN, M. A. & PETRZELKOVA, K. J. 2016. *Strongyloides infections* of humans and great apes in Dzanga-Sangha Protected Areas, Central African Republic and in degraded forest fragments in Bulindi, Uganda. *Parasitology International,* 65**,** 367-370.

HASEGAWA, H., SATO, H., FUJITA, S., NGUEMA, P. P. M., NOBUSUE, K., MIYAGI, K., KOORIYAMA, T., TAKENOSHITA, Y., NODA, S. & SATO, A. 2010. Molecular identification of the causative agent of human strongyloidiasis acquired in Tanzania: dispersal and diversity of *Strongyloides* spp. and their hosts. *Parasitology International,* 59**,** 407-413.

HEYDON, G. & GREEN, A. 1931. Some Worm Infestations of Man. in Australia. *Medical Journal of Australia,* 1.

HOLT, D. C., SHIELD, J., HARRIS, T. M., MOUNSEY, K. E., ALAND, K., MCCARTHY, J. S., CURRIE, B. J. & KEARNS, T. M. 2017. Soil-Transmitted Helminths in Children in a Remote Aboriginal Community in the Northern Territory: Hookworm is Rare but *Strongyloides stercoralis* and *Trichuris trichiura* Persist. *Tropical Medicine and Infectious Disease,* 2**,** 51.

JALETA, T. G., ZHOU, S., BEMM, F. M., SCHÄR, F., KHIEU, V., MUTH, S., ODERMATT, P., LOK, J. B. & STREIT, A. 2017. Different but overlapping populations of *Strongyloides stercoralis* in dogs and humans—Dogs as a possible source for zoonotic strongyloidiasis. *PLoS Neglected Tropical Diseases,* 11**,** e0005752.

KEARNS, T. M., CURRIE, B. J., CHENG, A. C., MCCARTHY, J., CARAPETIS, J. R., HOLT, D. C., PAGE, W., SHIELD, J., GUNDJIRRYIRR, R. & MULHOLLAND, E. 2017. *Strongyloides* seroprevalence before and after an ivermectin mass drug administration in a remote Australian Aboriginal community. *PLoS Neglected Tropical Diseases,* 11**,** e0005607.

MILLER, A., YOUNG, E., TYE, V., CODY, R., MUSCAT, M., SAUNDERS, V., SMITH, M., JUDD, J. & SPEARE, R. 2018. A community-directed integrated *Strongyloides* control program in Queensland, Australia. *Tropical Medicine and Infectious Disease,* 3**,** 48.

NAGAYASU, E., HTWE, M. P. P. T. H., HORTIWAKUL, T., HINO, A., TANAKA, T., HIGASHIARAKAWA, M., OLIA, A., TANIGUCHI, T., WIN, S. M. T. & OHASHI, I. 2017. A possible origin population of pathogenic intestinal nematodes, *Strongyloides stercoralis*, unveiled by molecular phylogeny. *Scientific Reports,* 7**,** 4844.

OLSEN, A., VAN LIESHOUT, L., MARTI, H., POLDERMAN, T., POLMAN, K., STEINMANN, P., STOTHARD, R., THYBO, S., VERWEIJ, J. J. & MAGNUSSEN, P. 2009. Strongyloidiasis–the most neglected of the neglected tropical diseases? *Transactions of the Royal Society of Tropical Medicine and Hygiene,* 103**,** 967-972.

PAGE, W., SHIELD, J., O'DONAHOO, F., MILLER, A., JUDD, J. & SPEARE, R. 2016. Strongyloidiasis in Oceania. *Neglected Tropical Diseases-Oceania.* Springer.

PAGE, W. A., DEMPSEY, K. & MCCARTHY, J. S. 2006. Utility of serological follow-up of chronic strongyloidiasis after anthelminthic chemotherapy. *Transactions of the Royal Society of Tropical Medicine and Hygiene,* 100**,** 1056-1062.

PAMPIGLIONE, S. & RICCIARDI, M. 1971. The presence of *Strongyloides fiilleborni* von Linstow, 1905, in man in Central and East Africa. *Parassitologia,* 13.

PUTLAND, R., THOMAS, S., GROVE, D. & JOHNSON, A. 1993. Analysis of the 18S ribosomal RNA gene of *Strongyloides stercoralis*. *International Journal for Parasitology,* 23**,** 149-151.

SAMPSON, I., SMITH, D. & MACKENZIE, B. Serological diagnosis of *Strongyloides stercoralis* infection. Second National Workshop on Strongyloidiasis, Royal Brisbane Hospital, Hernston, Australia, 25-26 June 2003. 25-26.

TAKANO, Y., MINAKAMI, K., KODAMA, S., MATSUO, T. & SATOZONO, I. 2009. Cross infection of *Strongyloides* between humans and dogs in the Amami Islands, Japan. *Tropical Medicine and Health,* 37**,** 149-152.

THANCHOMNANG, T., INTAPAN, P. M., SANPOOL, O., RODPAI, R., TOURTIP, S., YAHOM, S., KULLAWAT, J., RADOMYOS, P., THAMMASIRI, C. & MALEEWONG, W. 2017. First molecular identification and genetic diversity of *Strongyloides stercoralis* and Strongyloides fuelleborni in human communities having contact with long-tailed macaques in Thailand. *Parasitology Research,* 116**,** 1917-1923.

ZAHEDI, A., GOFTON, A. W., JIAN, F., PAPARINI, A., OSKAM, C., BALL, A., ROBERTSON, I. & RYAN, U. 2017. Next Generation Sequencing uncovers within-host differences in the genetic diversity of *Cryptosporidium* gp60 subtypes. *International Journal for Parasitology,* 47**,** 601-607.

S1 Table. GenBank accession numbers of the *SSU* HVR I and IV and *cox*1 haplotypes.

The GenBank accession numbers are provided for the *SSU* HVR I and IV haplotypes generated in this study (MK468654-MK468677). The GenBank accession numbers are provided for the *cox*1 haplotypes generated in this study (MK434217- MK434258).

Branch colors

- *Strongyloides stercoralis* dog / human genotypes
- *Strongyloides stercoralis* dog genotypes, clade 1
- *Strongyloides stercoralis* dog genotypes, clade 2
- *Strongyloides* sp. 'loris' and similar types
- *Strongyloides fuelleborni*
- *Strongyloides mirzai* and similar types
- *Strongyloides ratti*
- *Matastrongylus salmi* and similar types
- *Macrotrachela quadricornifera* (a rotifer) and similar types
- *Ancylostoma caninum*
- *Ancylostoma duodenale*
- *Ancylostoma* sp.
- *Ancylostoma ceylanicum*
- *Necator* sp. clade 1
- *Necator americanus*
- *Necator* sp. clade 2
- Other taxa

Specimen shading

Community 1 - Northern Territory, Australia
Community 2 - Northern Territory, Australia
Community 3 - Northern Territory, Australia
Community 4 - Northern Territory, Australia
Community 5 - Northern Territory, Australia
Community 6 - Northern Territory, Australia
Community 7 - Northern Territory, Australia
Human - New South Wales, Australia
Human - Northern Territory, Australia
Human - Queensland, Australia

***Strongyloides* 18S rDNA types with absent *cox*1 sequence**

	HVR I	HVR IV
Dog 13	-	A, G
Dog 32	-	A
Dog 45	X	I
Human 2	-	A
Sputum	II	A

Figure Legends

Figure 1. Dendrogram of clustered *cox*1 amplicons from Australian dog and human specimens

This dendrogram includes *cox*1 sequences generated in this study and a selection of previously published *cox*1 sequences that overlap with our 217 base pair *cox*1 amplicon by 100%. Specimens analyzed as part of this study are shaded according to their site of collection. Branches are color coded according to their identity; either a species assignment, a proposed genus assignment, or their *S. stercoralis* genotype. When available, *Strongyloides* sp. *cox*1 sequences are annotated with their associated *SSU* genotypes, with their HVR-I type shown in blue and their HVR-IV type shown in black. Specimens for which a *cox*1 sequence was not obtained are shown in a table embedded in the figure (bottom right), which includes two specimens possessing unique 18S genotypes; dog 13 (HVR-IV, type G) and dog 45 (HVR-I, type X). A dash (-) shown in this table indicates failed amplification and/or sequencing of that marker. 'Sputum' refers to the sole sputum sample from a human patient (human 4) included in this study. Sequences published in previous studies that are not from *S. stercoralis* are labelled with their GenBank and/or DNA Data Bank of Japan accession numbers followed by their species name. *Strongyloides stercoralis* sequences from previous studies are labelled with their accession number, host species, and country of origin. Note that 'CAR' means Central African Republic. Names of specimens collected as part of this study begin with a host name and a unique number assigned in this study, followed by a percentage similarity to (~) a near BLASTN hit identifiable by its accession number.

Figure 2. Schematic detailing the proposed modifications to the previously described *S. stercoralis* genotyping scheme

A graphical representation of the novel *Strongyloides* genotypes discovered in this study compared to haplotypes identified in previous reports. The location of sites that were genotypically informative based on the original genotyping method described by Jaletta *et al.* (2017) and Basso *et al.* (2018) are indicated, as well as new SNP/indel sites that have been incorporated into the typing scheme based on the results of this study (Jaleta et al., 2017, Basso et al., 2018). For hypervariable region I, we introduce two novel types (VIII and X), and assign new haplotype names to published sequences that had not been previously considered in this typing scheme (VII and IX). For hypervariable region IV, we introduce three novel types (E, F, G and I), and assign new haplotype names to published sequences that had not been previously considered in this typing scheme (C, D and H) (see Tables 3 and 4 for details).

Tables

Table 1: Primers and probes and PCR conditions

Primer	Amplicon	Sequence	Reaction conditions
Stro18S-1530F Stro18S-1630R Stro18S-1586T FAM	rDNA 101 bp	5'-GAATTCCAAGTAAACGTAAGTCATTAGC-3' 5'-TGCCTCTGGATATTGCTCAGTTC-3' 5'-ACACACCGGCCGTCGCTGC-3'-BHQ1	**Step 1:** 95°C for 15 min, **Step 2:** 95°C for 15 s, **Step 3:** 60°C for 30 s. Repeat steps two and three 40 times.
SSP_COX1_F * SSP_COX1_R *	mtDNA 270 bp	5'-TTTGATCCTAGTTCTGGTGGTAATCC-3' 5'-GTAGCAGCAGTAAAATAAGCACGAGA-3'	**Step 1:** 98°C for 30 s, **Step 2:** 98°C for 10 s, **Step 3:** 60°C for 10 s, **Step 4:** 72°C for 10 s, **Step 5:** 72 °C for 2 min. Repeat steps two to four 45 times.
NEW_HVR_I_F NEW_HVR_I_R	rDNA 500 bp	5'-GCTCATTATAACAGCTATAGACTACACGGTA-3' 5'-CCACAACAATCATTTTATGCACTTGG-3'	**Step 1:** 98°C for 30 s, **Step 2:** 98°C for 10 s, **Step 3:** 60°C for 10 s, **Step 4:** 72°C for 10 s, **Step 5:** 72 °C for 2 min. Repeat steps two to four 45 times.
NEW_HVR_IV_F NEW_HVR_IV_R	rDNA 320 bp	5'-CGGGCCGGACACTATAAGG-3' 5'-ATCTCTAAACAGGAACATAATGATCACTAC-3'	**Step 1:** 98°C for 30 s, **Step 2:** 98°C for 10 s, **Step 3:** 63°C for 10 s, **Step 4:** 72°C for 10 s, **Step 5:** 72 °C for 2 min. Repeat steps two to four 45 times.

*Broadly specific primers for amplification of *Strongyloides* sp. *cox*1 sequences as well as those of several hookworms

Table 2: Human and dog samples analyzed in this study and their complete genotype

Sample	HVR-I haplotype	HVR-IV haplotype	*Cox*1 Accessions
Human 1	Haplotype II	Haplotype A	MK434219†
Human 2	NA (excluded due to *S. ratti* contamination)	Haplotype A	NA
Human 3	Haplotype II	Haplotype A	MK434218
Sputum (Human 4)	Haplotype II	Haplotype A	NA
Dog 4	NA	NA	MK434258†
Dog 6	Haplotype II and Haplotype IV	Haplotype A and Haplotype B	MK434255 MK434256† MK434257†
Dog 7	Haplotype IV	Haplotype B	MK434254
Dog 10	NA	NA	MK434251 MK434252† MK434253†
Dog 12	NA	NA	MK434249 MK434250
Dog 13	NA	Haplotype A and Haplotype G*	MK434246 MK434247 MK434248†
Dog 14	NA (excluded due to *S. ratti* contamination)	NA	MK434244† MK434245†
Dog 15	NA	NA	MK434240† MK434241† MK434242† MK434243†
Dog 16	NA	NA	MK434238† MK434239†
Dog 18	Haplotype II and Haplotype VI	Haplotype A and Haplotype E*	MK434237†
Dog 19	NA	NA	MK434233 MK434234 MK434235† MK434236†
Dog 20	NA	NA	MK434230

			MK434231†
			MK434232†
Dog 21	NA	NA	MK434228
			MK434229†
Dog 22	Haplotype VIII*	Haplotype F*	MK434226†
			MK434227†
Dog 28	NA	NA	MK434225†
Dog 32	NA	Haplotype A	NA
Dog 37	NA	NA	MK434224†
Dog 41	NA	NA	MK434223†
			MK434220†
Dog 43	NA	NA	MK434221†
			MK434222†
Dog 45	Haplotype X*	Haplotype I*	NA

* Novel *SSU* haplotypes, †Novel *cox*1 sequences

Note: All GenBank Accession numbers associated with the sequences generated in this study are provided in Supplementary Table S1

Table 3. HVR I genotypes assigned to *Strongyloides* sp. based on current data

Haplotypes	GenBank Accession/s	Notes
I	AB923888.1, AF279916.2, AJ417023.1, MK468654	Found in dogs and humans.
II	KF926659.1, MK468655, MK468656, MK468657, MK468658	Found in dogs and humans.
III	AB453315.1	Fund in dogs and humans.
IV	AB272234.1, KU724124.1, MK468663	This haplotype has been detected in dogs though not in humans. It has also been described in a badger where it was assigned to *S. procyonis*.
V	Jaleta *et al*. (2017)	Described by Jaleta *et al*. (2017) but the sequence of this type was not available in GenBank prior to this study. This is potentially a dog-specific haplotype.
VI	AB453316.1, AB453314.1, MH932098.1, MH932099.1, MH932100.1, MK468660	Assigned only to *S. stercoralis*. Identified in dog 18 from this study. Described predominantly in dogs but also in a chimpanzee (AB453314.1).
VII	AB205054.1	A *S. procyonis* sequence greater than 99% similar to *S. stercoralis* Haplotype I and Haplotype IV.
VIII	MK468661	Novel *Strongyloides* sp. sequence from Dog 22 most similar to Haplotype VII.
IX	LC038066.1	*Strongyloides* sp. Okayama isolated from a Japanese striped snake. This sequence is greater than 99% identical to sequences of *S. stercoralis* and *S. procyonis*. Similar to Haplotype X (see below) identified in an Australian dog.
X	MK468662	Sequence identified in dog 45 in present study. Most similar to *Strongyloides* sp. Okayama (Haplotype IX).

Table 4. HVR IV genotypes assigned to *Strongyloides* sp. based on current data

Haplotypes	GenBank Accession/s	Notes
A	KY081221.1, KU724128.1, KU724125.1, LC085483.1, LC085482.1, LC085481.1, KU962182.1, KU962181.1, KU962180.1, KU962179.1, AB923888.1, KF926662.1, KF926661.1, AF279916.2, MH932097.1, MH932097.1, MH932095.1, KY081223.1, AB526826.1, AB453316.1, AB453315.1, AB453314.1, MK468664 - MK468671	Identified in dogs, humans and chimpanzees. A *S. stercoralis*- specific haplotype
B	KU724129.1	Identified only in dogs. Consistently assigned to *S. stercoralis*
C	M84229.1	This sequence was published in GenBank in 1993 and assigned to *S. stercoralis*. It has not appeared in the literature since based on our knowledge, though we assigned to genotype C for its historic value. It shares one SNP difference to type A (Figure 2).
D	AB272234.1, AB205054.1	Includes two sequences assigned to *S. procyonis*. This type is 99% similar to Haplotype B.
E	MK468674	Novel genotype identified in dog 18. Most similar to Haplotype A.
F	MK468675	Novel genotype identified in dog 22. Most similar to Haplotype D.
G	MK468676	Novel genotype identified in dog 13. Most similar to Haplotype A.
H	LC038066.1	*Strongyloides* sp. Okayama isolated from a Japanese striped snake.
I	MK468677	Novel genotype identified in dog 45. Most similar to Haplotype H.

Figure 1

Figure 2