Title: Pesticide exposure and adverse health effects associated with farm work in Northern Thailand

Authors: Chanese A. Forté[1], Justin Colacino[1,2,3], Katelyn Polemi[1], Andrea Guytingco[1], Nicholas J. Peraino[4], Siripond Jindaphong[5], Tharinya Kaviya[5], Judy Westrick[4], Rick Neitzel[1], Kowit Nambunmee[5]

Affiliations: [1]Department of Environmental Health Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA

[2]Department of Nutritional Sciences, University of Michigan School of Public Health, Ann Arbor, MI, USA

[3]Center for Computational Medicine and Bioinformatics, University of Michigan, Ann Arbor, MI, USA

[4]Lumigen Instrument Center, Wayne State University, Detroit, MI, USA

[5]Occupational Health and Safety, Mae Fah Luang University, Chiang Rai, TH

Contributions: Kowit Nambunmee, Rick Neitzel, Justin Colacino, and Chanese Forte developed the study design. Katelyn Polemi and Chanese Forte completed the literature review. Andrea Guytinco, Siripond Jindhapong, and Tharinya Kaviya collected samples and administering the survey to farmworkers. Judy Westrick and Nick Peraino completed the air sample analyses and Nick Peraino drafted the methods section for the air samples. Chanese Forte analyzed the data under the supervision of Justin Colacino, Kowit Nambunmee and Rick Neitzel. Chanese Forte drafted the paper with all authors contributing with edits.

Acknowledgements: Participant recruitment, Data Collection and Translation between English and Thai: Kanlaya Rattannasuwan, Anong Mungmueang, Aurai Channgoen, Rungthong Bunta, Rungtiwa N Lampang, Phornphan Saenchangmai, Kith Pakdeepong, Tharinya Kavuya, Nattaya Singkaew, Pimprapai Junsang, Phadungsuk Muanguk, Sratchaya Poolperm, Or-raya Onsongchan, and Wanwisa Kansa. We thank Dr. Kirsten Herold of the SPH Writing Lab, University of Michigan School of Public Health for her final review and editing.

Funding: This publication was supported by the Grant or Cooperative Agreement Number, T42 OH008455, funded by the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention or the Department of Health and Human Services. This study was also supported by the Environmental Toxicology and Epidemiology Program (ETEP) training grant by the National Institutes of Health/National Institute for Environmental Health Sciences (NIH/NIEHS) (Grant# T32 ES007062), and the University of Michigan Center for the Southeast Asian Studies and the Thai Studies Committee: Thai Studies Grant Program at the Center for Southeast Asian Studies.

Keywords: Occupational Health; Organophosphates; Carbamates; Pesticides; Occupational Exposure; Agriculture

Corresponding Author: Dr. Kowit Nambunmee, Occupational Health and Safety

School of Health Science, Mae Fah Luang University, 333 Moo 1, Tasud, Muang, Chiang Rai, 57100, Thailand, Phone: +6653-917225, Fax: +6653-916821, Email: kowit.nam@mfu.ac.th

Competing Interests: Authors have none to disclose.

Abstract: Objectives—To assess pesticide exposure and understand the resultant health effects relative to non-agriculture workers in Northern Thailand.

Methods—Collaborating with hospital staff and health care volunteers, we surveyed agricultural (n=70) and comparison workers (n=27) residing in Northern Thailand. We quantified exposure to pesticides including chlorpyrifos, methomyl, and metalaxyl, by air sampling and liquid chromatography/mass spectrometry. We estimated differences in self-reported health outcomes, complete blood counts, cholinesterase activity, and serum and urine calcium and creatinine concentrations at baseline between farmworkers and comparison workers, and after pesticide spraying in farmworkers only.

Results—None of the farmworkers wore standardized PPE for the concentrated chemicals they were working with. Methomyl, ethyl chlorpyrifos, and metalaxyl were detected air samples in varying amounts. When it came to reporting confidence in ability to handle personal problems, only 43% of farmworkers reported feeling confident, reflecting higher stress levels, in comparison to 78% of comparison workers (p = 0.028). Conventional farmworkers also had significantly lower monocyte counts (p=0.01), serum calcium (p=0.01), red blood count (p=0.01), white blood cell count (p=0.04), and butyrylcholinesterase activity (p<0.0001), relative to comparison workers. After adjusting for BMI, age, and smoking, methomyl air concentrations were associated with a decrease in farmworker acetylcholinesterase activity (beta= -0.327, p =0.016).

Conclusions—Farmworkers' blood analytes, acetylcholinesterase, and self-reported symptoms differed from comparison workers. Improving PPE use presents a likely route for preventive intervention in this population.

Introduction

Agriculture is a major contributor to Thailand's economy[1,2]. Among surveyed farmers residing in Northern Thailand, most (97%) reported using pesticides on their crops [3–6]. Over 93% of agriculture workers in Thailand work through the informal sector, and are not defined as employees under the Labor Protection Act; thus, these workers are exempt from numerous safety laws surrounding labor and social security rights[7,8]. Thailand ranked third out of 15 Asian countries for pesticide use per unit area and fourth in annual pesticide use[2]. Although there have been increases in organic food consumption in Thai markets, there is no evident reduction in pesticide use[2,7]. In fact, pesticide use in Thailand has increased from 110,000 tons in 2007 to roughly 172,000 tons in 2013[2]. For the pesticides being imported into Thailand, a third are considered extremely, highly, or moderately hazardous based on the World Health Organization's (WHO) hazard categories[2,9]. Insecticides are most often associated with negative human health effects[10]. Overall, research is lacking on low and middle income countries (LMIC) pesticide exposure and related health outcomes[11].

Pesticide pollution in the environment is associated with poisonings, oxidative stress, neurological dysfunction, birth defects, reproductive disorders, and cancers including colorectal cancer, prostate, and non-Hodgkin lymphoma[8,12–14]. Although Thailand does not have a poisonings registry, a poison center located in Bangkok recorded more than 15,000 patients over a three-year period, 42% of whom had poisonings related to pesticide exposure[8]. The pesticides most associated with these poisonings were insecticides: Carbamates, organophosphates, and pyrethroids[8].

This study was motivated by a group of contract farmers residing in Wiang Pa Pao, Chiang Rai Province. These farmers reported concerns about their health related to spraying

pesticides to Chiang Rai Prachanukroh hospital employees. Our project was launched in response to this concern. The purpose of this project was to assess the pesticide exposure of these farmworkers in Northern Thailand, and to understand the resultant health effects when compared to workers residing in a non-agricultural area (Chiang Rai, Thailand). This project adopted a mixed-methods approach during assessment of pesticide exposure and the resulting health effects, including: 1) personal air sampling, 2) biomarker sampling, and 3) perceived exposure and health effects assessed by questionnaire.

Material and Method

Study Population

The study protocol was approved by the Institutional Review Boards of Mae Fah Luang University and the University of Michigan (UM Submission ID: HUM00128091/CR00068767). All study participants were 21 years and older, male, and resided in Thailand. Participants were recruited in Chiang Rai, Thailand, using a recruitment script in Thai administered by health volunteers—laypersons trained through the Chiang Rai Prachanukroh hospital on patient care and interactions. Comparison workers were recruited through word of mouth at Mae Fah Luang University (MFU), with a focus on older males in work fields unrelated to agriculture with no commercial pesticide spraying experience.

Sample Collection

Participants completed the consent form by oral confirmation due to literacy differences. Conventional farmworkers consented to allow work observation, active air sampling, and pre- and post-spray urine and blood samples to be taken. Each study participant also completed a

researcher-administered survey. Participants received 600 Thai baht for participation and completion of the study.

We quantified exposure and potential adverse health effects among workers by self-report, biological, and environmental sampling. A 69-item occupational health questionnaire was translated into Thai from English by MFU researchers. The questionnaire was based on a previous study of worker health and occupational noise[16]. The survey was administered to study participants at the local community center and took roughly 35 to 45 minutes to complete. Questionnaire data were collected from July 2017 to the end of August 2017 (conventional farmworkers and comparison workers) and again in January 2018 (sample of comparison workers from Chiang Rai).

Nurses collected 10 mL of urine and 10 mL of blood from workers at the local hospital (Chiang Rai or Wiang Pa Pao). For farmworkers, blood and urine samples were collected within one week prior to pesticide spraying and again within 3 days after pesticide spraying. Whole blood samples were collected in red-top tubes with no anticoagulant, lavender-top tubes with EDTA, and green-top tubes with heparin.

Blood and Urine Analysis

Urine creatinine, urinary calcium, serum creatinine, serum calcium, and complete blood counts were quantified by Meng Rai Laboratory in Chiang Rai. Acetylcholinesterase (AChE) and butyrylcholinesterase (BuChE) were analyzed using the Ellman method, from whole blood and serum, respectively[18]. AChE was analyzed since it is directly related to AChE inhibition by the pesticides, and BuChE since it can be a sensitive biomarker of exposure to AChE inhibitors[17]. Concentrations of the health biomarkers were compared between conventional

farmworkers and comparison study participants, as well as within farmworkers before and after spraying.

Air Sample Collection and Analysis

Personal pesticide air exposure levels were measured in conventional farmworkers only. GilAir Plus Personal Air Sampling Pumps by Sensidyne, Inc. were calibrated each day with the Gilbrator Calibrator before heading to the field. Air samples were collected during the time farmers were spraying pesticides (14 to 63 minutes) using XAD-2 sorbent tubes (SKC Ltd) based on the standard NIOSH method 5600 at a flowrate of one liter per minute[19]. One field blank was collected by opening tubes away from the farms for a similar period of time; no air was drawn through the blanks, but they were otherwise handled identically to actual air samples.

The analysis of pesticide residues was completed by the Lumigen Instrument Center at Wayne State University. Laboratory controls from a spiked and blank filter were extracted each batch. A calibration check at 10 ng/mL and a solvent blank were run every 10 samples. If a check did not pass within 20% error, the entire 10 sample section was re-run. Calibration curves were prepared the same day as every batch as concentration vs area ratio of analytes to deuterated surrogates. A linear regression was taken, and percent error calculated at each calibration point. If a point did not fall within 20% error of the predicted value, it was dropped from the end of the curve.

Air sampling tubes were extracted as either top portions or bottom portions. Top portions of the tubes contained the retainer ring, filter paper, foam pad, and XAD fill. Bottom portions contained the middle foam pad, and the bottom portion of XAD fill; the bottom foam was discarded. Portions were transferred to a 2 dram vial followed by 1960 µL of methanol and 40

μL of a solution containing 1 μg/mL each of deuterated surrogates. The vials were tightly

capped and sonicated for 30 minutes, allowing the bath to heat naturally. Heating was found

necessary to achieve equilibration of the surrogates and standards to the XAD fill, and improve

recoveries. Samples were centrifuged to settle any particulates and 900 μL of sample was diluted

with 100 μL of water containing 100 mM ammonium formate to give a final sample containing

10% water and 10 mM ammonium formate. Liquid chromatography mass spectroscopy (LCMS)

analysis was completed using a Nexera-X2 ultra performance liquid chromatography (UPLC)

with Shimadzu 8040 triple quadrupole mass analyzer. Chromatography was achieved using a

Waters acquity UPLC Ethylene-Bridged Hybrid (BEH) C18 (1.2 μm, 2.1 x 50 mm) column

eluting with 10 mM ammonium formate in water (mobile phase A) and 10 mM ammonium

formate in methanol (mobile phase B).

Worker Observations

MFU and UM student researchers observed the farmworkers during spraying activities

and recorded worker practices for mixing, spraying, and storage of the pesticides on a worker

observation form. This included information on the pump used, pump calibration, personal

protective equipment (PPE) used, and notes on common practices. Detailed notes on PPE used

such as item and material were noted by observers (e.g. gloves made of rubber, latex, or cloth).

Work observations were not completed for comparison workers since they do not perform

commercial pesticide spraying.

Statistical Analyses

For questionnaire data, we calculated summary statistics of demographic data across the

entire population. We first tested for crude differences across the measures between the

conventional farmers and the comparison workers by t-test for continuous variables or by

Fisher's Exact test for categorical variables. Analysis of variance (ANOVA) and multivariate

regression analyses were used to compare serum and urinary biomarker concentrations between

farmworkers and comparison workers, adjusting for the potential confounders age, smoking

status, alcohol consumption, and BMI. We also conducted multivariate linear regression analyses

of the association between pesticide levels quantified in the personal air samples and serum and

urinary biomarkers in the farmworkers collected after spraying, adjusting for the same covariates

as described above. Any variables that were not normally distributed were log transformed prior

to regression. SAS 9.4 was used to complete regressions and demographic tables, and R 3.5.1

was used to create figures and graphics.

Results

Overall, farmworkers and comparison workers had similar demographics (Table 1),

except for BMI, which was significantly lower for farmworkers. Farmworkers had a median age

of 50 years with a range of 22 to 76 years of age, and comparison workers had a median age of

49 with a range of 39 to 68 years. Most of the workers were married, attended primary and

secondary school, drank alcohol 1-3 days per week, and were current smokers (Table 1). With

respect to self-reported health concerns, farmworker and comparison worker self-reported

symptom responses only statistically significantly differed for 'dizziness' and 'shaking or

trembling of hands' (Supplemental Table 1).

None of the farmworkers wore chemical proof aprons, chemical proof gloves, or a

respirator. The farmworkers did wear an item that we named 'face hats' which included both

home-made and commercial versions. The home-made version was usually two towels or scarves (e.g. durags) stitched together to cover the head and face, and the commercial version was a canvas brimmed hat with a canvas face covering that could be removed. The use of gloves, long sleeve clothing, and any sort of clothes covering (e.g. rain poncho or plastic sheet) were usually used items with some damage and use was not consistent across workers.

Supplemental Table 2 presents worker self-reported stress levels by worker category based on the Cohen's Perceived Stress Scale. Overall, comparison workers more frequently reported stress in comparison to farmworkers, although not statistically significantly so. However, when it came to reporting confidence in ability to handle personal problems, only 43% of farmworkers reported feeling confident often in comparison to 78% of comparison workers (p-value = 0.028).

A comparison of cholinesterase activity between the two groups indicated that although there were some farmworkers with higher AChE activity, the medians of farmworkers and comparison workers were not significantly different. AChE and BuChE concentrations were not found to be correlated (0.05, 95% CI: -0.199, 0.294). Comparison workers had higher BuChE activity levels in comparison to farmworkers (Figure 1). Based on the ANOVA with cholinesterase measurement at baseline as the outcome and worker category as the predictor, worker AChE activitiy did not differ (p=0.20), while BuChE activity did differed significantly (p<0.0001). Within farmworkers only, pre and post-spray activity of AChE (0.31, p-value=0.0127) and BuChE (0.31, p-value=0.0129) were correlated.

Measurements of pesticide residues on air samples are displayed in Figure 2. Overall, ethyl chlorpyrifos and metalaxyl were detected at the highest frequency, while methomyl was less frequently detected. Most chlorantraniliprole (N=60) and methyl-chlorpyrifos (N=61)

measurements were below the limit of detection. Ethyl chlorpyrifos followed by metalaxyl and methomyl, was found in the highest concentrations in the air filter samples.

Table 2 presents linear regression parameter estimates and confidence intervals comparing biomarker concentrations between farmworkers and comparison workers. Eosinophil (p=0.02), urine creatinine (p=0.03), and MCV (p=0.04) concentrations were statistically significantly higher in farmworkers than the comparison group. Monocytes (p=0.01), red blood cell counts (p=0.01), white blood cell count (p=0.04), and serum calcium (p=0.02) were statistically significantly lower in farmworkers than comparison workers. Table 3 presents the linear regression beta estimates between biomarker concentrations and air sample pesticide content for farmworkers only. The ratio of pre- and post-spray AChE was significantly lower for increased concentrations of methomyl in air samples (p=0.02). Urinary creatinine and serum calcium were inversely associated with air sample concentrations of metalaxyl.

Discussion

This pilot study was designed to better characterize pesticide exposure and resultant health effects among farmworkers residing in Northern Thailand. We used a mixed-methods approach which included collection of self-reported data, biological samples, and environmental air sampling. Farmworker and comparison worker demographics were not significantly different, with the exception of BMI, where farmworkers had significantly lower values most likely due to full work days doing physical labor. Based on worker observations, PPE was never suffificient for protection based on US Occupatonal Saftety and Health Agency regulations for protective gear. Farmworkers differed from comparison workers in self-reported stress levels such as the

ability to handle personal problems, which was reported less in farmworkers. Farmworkers and comparison workers responded statistically significantly different on dizziness and in shaking or trembling of the hands. Biomarker concentrations also differed significantly between farmworkers and comparison workers. Farmworker concentations of urinary creatinine, MCV, Eosinophil were higher, whereas monocytes,WBC, serum calcium, and BuChE ratio were lower. Based on the regression analyses of only exposed farmworkers, serum calcium and urine creatinine were statistically significant among farmworkers exposed to metalaxyl. AChE ratio was also statistically lower for methomyl in the regression model controlling for age, BMI, current smoking status and former smoking status among farmworkers.

The most commonly used insecticides in Thailand are AChE inhibitors such as organophosphates and carbamates like chlorpyrifos and methomyl [2]. In this study, 59 farmworkers self-reported using chlorpyrifos during pesticide spraying, 17 reported using metalaxyl, and 1 reported using methomyl. The analytic method utilized in this study quantified ethyl-chlorpyrifos, methyl-chlorpyrifos, metalaxyl, methomyl, and chlorantraniliprole. However, only ethyl chlorpyrifos, methomyl, and metalaxyl were detected on workplace air samples. Methomyl is a carbamate and insecticide and metalaxyl is a benzene compound used as a fungicide[20,21].

We observed workers during day 8 of edamame growing because they reported using chlorpyrifos during that time. This sampling timepoint likely resulted in our air samples not containing chemicals that are used at other times in the edamame growth cycle, based on farmworker self-reported chemical use and the spray schedule. Also, exposure to carbamates or OPPs can alter AChE levels from baseline[22–24]. Considering the cross-sectional nature of this study and the frequency of pesticide spraying among our cohort, we cannot determine whether

the cholinesterase biomarker results were due directly to the insecticides monitored or from another pesticide sprayed during another agriculture job or spraying another plant on the same farm.

On average, the farmworkers in our cohort worked 50.95 hour work weeks. The Labor Protection Act limit jobs that are deemed hazardous to human health a maximum of 42-hour work weeks. However, since 93% of all farmworkers in Thailand are informal sector workers they are not working under the Labor Protection Act of 1998[7,8]. Our cohort pilot study clearly demonstrates there are informal sector farmworkers working above the legal limit for hazardous jobs. For migrant farmworkers, this means they are likely working unhealthy hours and also lack access to the Thai healthcare system[25].

In a study by Sapbamrer and Nata (2017), rice farmers and non-farmer controls residing in Northern Thailand were interviewed and had blood samples taken to ascertain their overall pesticide exposure[27]. When looking at self-reported health outcomes in our study, farmworkers did not have differences in respiratory symptoms relative to comparison workers. Farmworkers reported trembling in their hands less often in comparison to other workers, whereas with exposure to AChE inhibitors, we would expect farmworkers would report trembling more often.

Other studies have assessed the relationship between pesticide exposure and cholinesterase activity outside of Thailand. Strelitz et al. measured baseline to post-spraying whole blood and serum cholinesterase levels of 215 farmworkers from the Washington State Cholinesterase Monitoring Program. Ellman colorimetric enzymatic assays were used by two different laboratories to measure BuChE and AChE using the same methodology (Washington State Public Health Laboratories in 2006 and Pathology Associates Medical Laboratories in 2007 to 2011)[17]. In this same study, cholinesterase depression was defined as 20% or greater decrease

from baseline to post-exposure cholinesterase exposure[17]. The authors found AChE and BuChE

activity to be negatively correlated (-0.14, 95% CI: -0.27, -0.01)[17]. Our study also found the

correlation between AChE and BuChE to be weak, but the correlation was non-significantly

positive (0.05, 95% CI: -0.199, 0.294). The ratio of pre-and post- spray AChE activity was also

significant lower with increasing concentrations of methomyl in air samples. Others have

reported BuChE activity as a more sensitive biomarker of exposure to cholinesterase inhibitors

than AChE[17,28,29]. In our study, BuChE activity was significantly lower in conventional

farmworkers relative to comparison workers.

Garcia-Garcia et al. 2015 completed a similar study comparing 169 green house workers

who were exposed to pesticides and controls who were unexposed to pesticides in southeastern

Spain[30]. Among this cohort, researchers controlled for sex, age, and BMI , but not smoking

status. They found BuChE inhibition, RBC, MCV, and neutrophil levels significantly increased,

whereas eosinophils significantly decreased and monocytes were not different[30]. Our study

assessed hematological parameters by measuring differences in complete blood counts between

farmworkers and comparison workers, while controlling for BMI, age, former smoker status,

current smoker status, worker category, and pesticide exposure levels. Serum calcium was

statistically significant by both worker category and was also significantly different in

farmworker. MCV, monocytes, RBC, eosinophils, urine creatinine and WBC were significantly

different between our Northern Thailand farmworkers and comparison worker. In our study,

farmworkers had a reduction, in RBC and neutrophils, whereas MCV increased (opposite of

Garcia-Garcia et al.). Monocytes and eosinophils reduced and increased, respectively in our

study and the Garcia-Garcia et al. study. In our study, an association existed between pesticide

exposure and all of the aforementioned blood counts when looking at farmworkers and comparison workers.

Finally, a cohort study on farmworkers in China before and after pesticide spraying points to the effects of pesticides on complete blood count (CBC) having an effect that can be categorized as long term (3 years) or short term (10 days or less). Long term results include decreased white blood cell count, and short term results such as alterations in complete blood count, hepatic and renal functions, nerve conduction velocities, and on monocytes, hemoglobin and platelet counts[31]. When comparing air sample measurements of methomyl, chlorpyrifos and metalaxyl, we did not identify differences in blood count analytes by chemical exposure, although we identified a range of differences when comparing farmworkers and comparison workers. Prior research has reported significant changes to hematological biochemistry as a result of pesticide exposures causing oxidative stress[30]. Oxidative stress due to pesticide spraying has also been related to changes in CBC such as monocyotsis, leukocytosis, neutrophilia, and lymphocytopenia which were thought to also be closely related to patients in oxidative stress[30,32].

Our study has some limitations. We focused on farmworkers who contracted to numerous farms, and therefore many workers were likely exposed to other chemicals that were not quantified in this study. These farmworkers may have been exposed to chemicals that were not captured by this study because they sprayed pesticides on other farms and also sprayed other plants on the same farms. Due to this discrepancy, our farmworkers' baseline cholinesterase measurements may not actually represent a true baseline measurement due to their overlapping work schedules. The US Occupational Safety and Health Administration (OSHA) generally recommends testing for baseline cholinesterase levels after not working with organophosphates for at least 30 days[33]. OSHA also recommends taking a baseline measurement before and after

organophosphate use (with at least a 30 day washout period for both)[33]. This pilot study did not take multiple baseline measurements, and the one baseline that was taken was likely taken before the OSHA recommended guideline of 30 days since pesticide use. However, this population does not appear to ever experience thirty days of no pesticide use. Our study also focused on workplace sampling at a time when the specific farm of interest was expected to be spraying chlorpyrifos, and therefore the study results show an over-representation of chlorpyrifos.

Overall, this is the first study of its type that took a mixed-methods approach using survey, biomarker, and workplace observation data to analyze farmworker pesticide health effects in comparison to other workers in Northern Thailand. Our study is also one of the larger studies to use a combination of air sampling and biomarker data. These data can inform the methods for future global occupational health research on farmworkers. Work observations also included a more detailed outline of PPE used by the farmworkers and could inform future studies on PPE in relation to pesticide exposure and preventive interventions. We are hopeful this study will contribute to the building literature on farmworker pesticide exposure and resultant health outcomes.

References

1.   Panuwet P, Siriwong W, Prapamontol T, et al. Agricultural Pesticide Management in

        Thailand: Situation and Population Health Risk. *Env sci policy*. 2012;17:72-81.

        doi:10.1016/j.envsci.2011.12.005.Agricultural.

2.   Tawatsin A., Thavara, U & Siriyasatien P. Pesticides used in Thailand and toxic effects to

        human health. *Med Res Arch*. 2015;(3):1-10. doi:10.18103/mra.v0i3.176.

3.   Kearns JP, Knappe DRU, Summers RS. Synthetic organic water contaminants in

        developing communities: an overlooked challenge addressed by adsorption with locally

        generated char. *J Water Sanit Hyg Dev*. 2014;4(3):422-436.

        doi:10.2166/washdev.2014.073.

4.   Thapinta A, Hudak PF. Pesticide use and residual occurrence in Thailand. *Environ Monit

        Assess*. 2000;60(1):103-114. doi:10.1023/A:1006156313253.

5.   Kunstadter P, Prapamontal T, Sirirojn BO, Sontirat A, Tansuhaj A, Khamboonruang C.

        Pesticide exposures among Hmong farmers in Thailand. *Int J Occup Environ Health*.

        2001;7(4):313-325. doi:10.1179/oeh.2001.7.4.313.

6.   Plianbangchang P, Jetiyanon K, Wittaya-areekul S. Pesticide use patterns among small-

        scale farmers: A case study from phitsanulok, Thailand. *Southeast Asian J Trop Med

        Public Health*. 2009.

7.   Vasquez C. Illegal Pesticide Trade in Mekong Countries: Pesticide Action Network Asia

        and the Pacific (PANAP) Sustainable Agriculture and Environment Development

        Association (SAEDA). 2011.

8.   Kaewboonchoo O, Kongtip P, Woskie S. Occupational health and safety for agricultural workers in Thailand: gaps and recommendations, with a focus on pesticide use. *New Solut*. 2015;25(1):102-120. doi:10.1177/1048291115569028.

9.   Panuwet P, Prapamontol T, Chantara S, et al. Concentrations of urinary pesticide metabolites in small-scale farmers in Chiang Mai Province, Thailand. *Sci Total Environ*. 2008;407(1):655-668. doi:10.1016/j.scitotenv.2008.08.044.

10.   Wang L, Liu Z, Zhang J, Wu Y, Sun H. Chlorpyrifos exposure in farmers and urban adults: Metabolic characteristic, exposure estimation, and potential effect of oxidative damage. *Environ Res*. 2016;149:164-170. doi:10.1016/j.envres.2016.05.011.

11.   Weiss F, Leuzinger M, Zurbrügg C, Eggen R. Chemical Pollution in Low- and Middle-Income Countries. *Swiss Fed Inst Aquat Sci Technol*. 2016:164.

12.   Lee WJ, Sandler DP, Blair A, Samanic C, Cross AJ, Alavanja MCR. Pesticide use and colorectal cancer risk in the Agricultural Health Study. *Int J cancer*. 2007;121(2):339-346. doi:10.1002/ijc.22635.

13.   Oddone E, Modonesi C, Gatta G. Occupational exposures and colorectal cancers: A quantitative overview of epidemiological evidence. *World J Gastroenterol*. 2014;20(35):12431-12444. doi:10.3748/wjg.v20.i35.12431.

14.   Lee WJ, Hoppin JA, Blair A, et al. Cancer Incidence among Pesticide Applicators Exposed to Alachlor in the Agricultural Health Study. *Am J Epidemiol*. 2004;159(4):373-380. doi:10.1093/aje/kwh040.

15.   Mostafalou S, Abdollahi M. Pesticides and human chronic diseases: Evidences,

mechanisms, and perspectives. *Toxicol Appl Pharmacol*. 2013;268(2):157-177. doi:10.1016/j.taap.2013.01.025.

16. Burns KN, Sun K, Fobil JN, Neitzel RL. Heart rate, stress, and occupational noise exposure among electronic waste recycling workers. *Int J Environ Res Public Health*. 2016;13(1). doi:10.3390/ijerph13010140.

17. Strelitz J, Engel LS, Keifer MC, Clinic M. Blood acetylcholinesterase and butyrylcholinesterase as biomarkers of cholinesterase depression among pesticide handlers. *Occup Env Med*. 2014;71(12):842-847. doi:10.1136/oemed-2014-102315.Blood.

18. Ellman GL, Courtney KD, Andres V, Featherstone RM. A new and rapid colorimetric determination of acetylcholinesterase activity. *Biochem Pharmacol*. 1961;7:88-95.

19. National Institute of Occupational Safety and Health (NIOSH). Organophosphorus Pesticides Method 5600. *NIOSH Man Anal Methods*. 1993;4(1):419-473. doi:10.1002/0471435139.tox095.pub2.

20. Torres CM, Picó Y, Mañes J. Determination of pesticide residues in fruit and vegetables. *J Chromatogr A*. 1996;754(1-2):301-331. doi:10.1016/S0021-9673(96)00407-4.

21. Gupta RC, Crissman JW. *Agricultural Chemicals*. Third Edit. Elsevier; 2013. doi:10.1016/B978-0-12-415759-0.00042-X.

22. Wilson BW, Arrieta DE, Henderson JD. Monitoring cholinesterases to detect pesticide exposure. *Chem Biol Interact*. 2005;157-158(November):253-256. doi:10.1016/j.cbi.2005.10.043.

23. Stefanidou M, Athanaselis S, Spiliopoulou H. Butyrylcholinesterase: Biomarker for

exposure to organophosphorus insecticides. *Intern Med J*. 2009;39(1):57-60.

doi:10.1111/j.1445-5994.2008.01779.x.

24.   Hofmann JN, Keifer MC, De Roos AJ, et al. Occupational determinants of serum

cholinesterase inhibition among organophosphate-exposed agricultural pesticide handlers

in Washington State. *Occup Environ Med*. 2010;67(6):375-386.

doi:10.1136/oem.2009.046391.

25.   Thetkathuek A, Daniell W. Migrant workers in agriculture: a view from Thailand. *J*

Agromedicine. 2016;21(1):106-112. doi:10.1080/1059924X.2015.1043169.

26.   Thetkathuek A, Yenjai P, Jaidee W, Jaidee P, Sriprapat P. Pesticide Exposure and

Cholinesterase Levels in Migrant Farm Workers in Thailand. *J Agromedicine*.

2017;22(2):118-130. doi:10.1080/1059924X.2017.1283276.

27.   Sapbamrer R, Nata S. Health symptoms related to pesticide exposure and agricultural

tasks among rice farmers from northern Thailand. *Environ Health Prev Med*.

2014;19(1):12-20. doi:10.1007/s12199-013-0349-3.

28.   Andreescu S, Marty JL. Twenty years research in cholinesterase biosensors: From basic

research to practical applications. *Biomol Eng*. 2006;23(1):1-15.

doi:10.1016/j.bioeng.2006.01.001.

29.   Amine A, Mohammadi H, Bourais I, Palleschi G. Enzyme inhibition-based biosensors for

food safety and environmental monitoring. *Biosens Bioelectron*. 2006;21(8):1405-1423.

doi:10.1016/j.bios.2005.07.012.

30.   García-García CR, Parrón T, Requena M, Alarcón R, Tsatsakis AM, Hernández AF.

Occupational pesticide exposure and adverse health effects at the clinical, hematological and biochemical level. *Life Sci*. 2016;145:274-283. doi:10.1016/j.lfs.2015.10.013.

31.   Hu R, Huang X, Huang J, et al. Long- and short-term health effects of pesticide exposure: A cohort study from China. *PLoS One*. 2015;10(6):1-13. doi:10.1371/journal.pone.0128766.

32.   Dundar ZD, Ergin M, Koylu R, Ozer R, Cander B, Gunaydin YK. Neutrophil-lymphocyte ratio in patients with pesticide poisoning. *J Emerg Med*. 2014;47(3):286-293. doi:10.1016/j.jemermed.2014.01.034.

33.   MCN's Environmental and Occupational Health Advisory Committee and AgriSafe's Quality Assurance Committee. Cholinesterase Testing Protocols for Healthcare Providers. (2010): 22284.

What this paper adds

1. What is already known about this subject?
   - Agriculture is classified under the informal sector in Thailand which means many of the workers are not protected under worker safety laws.
   - No known study has used a mixed-methods approach (self-report, biological markers, and environmental exposures) to assess farmworker pesticide exposure in this population.

2. What are the new findings?
   - Methomyl, ethyl chlorpyrifos, and metalaxyl were detected in varying amounts in air monitor sampling.
   - Conventional farmworkers also had significantly lower monocyte counts (p=0.01), serum calcium (p=0.01), red blood count (p=0.01), white blood cell count (p=0.04), and butyrylcholinesterase activity (p<0.0001), relative to comparison workers.
   - After adjusting for BMI, age, and smoking, methomyl air monitor concentrations were associated with a decrease in farmworker acetylcholinesterase activity (beta= -0.327, p =0.016).

3. How might this impact on policy or clinical practice in the foreseeable future?
   - This study provides a new mixed methods approach to assessing worker exposure to chemicals, and therefore can highlight more adverse effects such as mental health, biomarkers, and PPE which were previously not possible in one study design.
   - Farmworkers on average worked 50.96 hours per week which is above the Royal Thai Government Labor Protection Act of 1998 that limits work weeks to 42 hours.
   - Farmworkers are exposed to high levels of pesticides without access to the appropriate PPE which could cause new company policies for PPE access or laws by the Royal Thai Government.

Table 1
Study participant demographics, males only

Variable	Variable Outcome	Comparison Worker		Farmworker		Fisher's Exact Significance
		N=27	%	N=70	%	p-value
Marital Status	single	6	22.2	7	10.0	0.11
	married	16	59.3	55	78.6	
	divorced	3	11.1	2	2.9	
	living with partner	2	7.4	6	8.6	
Education	none	0	--	3	4.3	0.48
	primary	18	66.7	49	70.0	
	secondary	6	22.2	15	21.4	
	some college	1	3.7	2	2.9	
	4 year degree	1	3.7	1	1.4	
	graduate Level	1	3.7	0	--	
Alcohol	never	7	26.9	12	17.4	0.64
	not much	4	15.4	14	20.3	
	1-3 days per week	9	34.6	18	26.1	
	4-6 days per week	4	15.4	14	20.3	
	daily	2	7.7	11	15.9	
Tobacco Use	never smoked	8	30.8	25	35.7	0.67
	former smoker	8	30.8	15	21.4	
	currently smokes	10	38.5	30	42.9	
		Median	STD	Min.	Max.	
Comparison Worker Age		49.0	8.1	39.0	68.0	0.61
Farmworker Age		50.0	11.0	22.0	76.0	
Comparison Worker BMI		24.7	3.7	20.0	34.1	0.02[a]
Farmworker BMI		22.2	3.5	16.1	33.1	

[a]Signifies a p-value less than 0.05

Table 2

Linear regression analyses comparing biomarker concentrations in farmworkers and comparison workers

Analytes	Units	Farmworkers=1 vs. Comparison=0			
		β Estimate[b]	p-value	95% CI	
log(Basophil)	%	0.12	0.13	-0.04	0.29
log(Eosinophil)	%	0.24[a]	0.02	0.04	0.44
log(Hemoglobin)	g/dl	-0.01	0.16	-0.03	0.01
log(Hematocrit)	%	-0.01	0.28	-0.03	0.01
log(Lymphocyte)	%	-0.03	0.41	-0.10	0.04
log(Mean corpuscular hemoglobin concentration (MCHC))	pg	<0.01	0.53	-0.01	0.01
log(Mean Corpuscular Volume)	fl	0.03 [a]	0.04	<0.01	0.06
Monocyte	%	-1.33 [a]	0.01	-2.30	-0.37
Neutrophil	%	-1.34	0.69	-8.08	5.39
log(Plate Count)	cells/uL	-0.03	0.39	-0.10	0.04
log(Red Blood Cell (RBC) Count)	cells/uL	-0.04 [a]	0.01	-0.07	-0.01
log(RBC Distribution Width (RDW))	%	0.01	0.19	-0.01	0.03
log(White Blood Cell Count)	cells/uL	-0.06 [a]	0.04	-0.12	<0.01
Serum Calcium	mg/dl	-0.79 [a]	<0.01	-1.31	-0.27
log(Serum Creatinine)	mg/dl	-0.03	0.22	-0.07	0.02
log(Urine Calcium)	mg/dl	0.14	0.14	-0.05	0.32
Urine Creatinine	mg/dl	36.29 [a]	0.03	4.17	68.41
Acetylcholinesterase	U/L	0.004	0.47	-0.01	0.02
Butyrylcholinesterase	U/L	-0.29 [a]	<0.01	-0.40	-0.19

[a]Signifies a p-value less than 0.05

[b]These values were adjusted for age, former smoker status, current smoking status, alcohol use, and BMI.

Table 3

Linear regression models of association between blood analytes and pesticide air sample concentrations

Analytes	Units	Log(Methomyl)[c]				Log(Metalaxyl)[c]				Log(Ethyl Chlorpyrifos)[c]			
		β	p-value	95% CI		β	p-value	95% CI		β	p-value	95% CI	
log(Basophil)	%	0.28	0.52	-0.61	1.16	-0.19	0.05	-0.39	0.00	-0.03	0.83	-0.30	0.25
log(Eosinophil)	%	-0.25	0.64	-1.33	0.84	-0.07	0.56	-0.33	0.19	-0.22	0.17	-0.54	0.10
log(Hemoglobin)	g/dl	-0.01	0.79	-0.09	0.07	-0.01	0.41	-0.03	0.01	-0.01	0.40	-0.03	0.01
log(Hematocrit)	%	-0.02	0.65	-0.11	0.07	0.00	0.80	-0.02	0.01	-0.01	0.38	-0.04	0.02
log(Lymphocyte)	%	0.18	0.33	-0.19	0.54	0.00	0.94	-0.09	0.09	-0.08	0.15	-0.19	0.03
log(MCHC)	pg	0.01	0.68	-0.04	0.06	-0.01	0.37	-0.02	0.01	0.00	0.83	-0.01	0.02
log(MCV)	fl	-0.07	0.28	-0.22	0.07	0.02	0.27	-0.02	0.05	0.01	0.50	-0.03	0.06
Monocyte	%	-1.38	0.58	-6.49	3.73	0.32	0.59	-0.91	1.55	0.18	0.82	-1.40	1.76
Neutrophil	%	-1.61	0.93	-38.75	35.53	0.93	0.83	-7.98	9.83	6.35	0.24	-4.69	17.40
log(Plate Count)	cells/uL	0.13	0.37	-0.16	0.41	0.00	0.98	-0.07	0.07	-0.07	0.12	-0.15	0.02
log(RBC)	cells/uL	0.05	0.49	-0.11	0.22	-0.02	0.26	-0.06	0.02	-0.03	0.27	-0.07	0.02
log(RDW)	%	0.05	0.26	-0.04	0.14	0.00	0.86	-0.02	0.03	0.00	0.93	-0.03	0.03
log(WBC)	cells/uL	-0.12	0.31	-0.37	0.12	0.01	0.83	-0.05	0.07	-0.02	0.62	-0.10	0.06
Serum Calcium	mg/dl	0.13	0.62	-0.38	0.63	-0.55[a]	0.02	-1.02	-0.08	0.18	0.59	-0.48	0.83
log(Serum Creatinine)	mg/dl	-0.02	0.20	-0.06	0.01	-0.01	0.55	-0.05	0.03	0.03	0.22	-0.02	0.08
log(Urine Calcium)	mg/dl	0.02	0.81	-0.13	0.17	-0.05	0.52	-0.19	0.10	0.07	0.45	-0.12	0.26
Urine Creatinine	mg/dl	6.39	0.64	-20.60	33.38	-30.17[a]	0.02	-54.94	-5.41	5.70	0.74	-29.21	40.61
AChE Ratio[b]	U/L	-0.37[a]	0.02	-0.66	-0.07	0.05	0.72	-0.25	0.36	0.35	0.08	-0.04	0.74
BuChE Ratio[b]	U/L	-0.13	0.35	-0.40	0.15	-0.05	0.70	-0.33	0.22	-0.12	0.50	-0.48	0.24

[a]Signifies a p-value less than 0.05

[b]Cholinesterase ratios were calculated by dividing post measures by pre measures.

[c]These values were adjusted for age, former smoker status, current smoking status, alcohol use, and BMI.

Figure 1

Distribution of acetyl- and butyrylcholinesterase activities in comparison workers (one time point only) and farmworkers (pre- and post-spray)[a]

[a]In previous research completed in Northern Thailand, the national reference range is 6,400 U/L to 8,200 U/L[27].

Figure 2

Pesticide measurements captured by personal air monitors on Northern Thailand farmworkers (N=51) with histograms showing distributions of detected values of (A) metalaxyl and (B) ethyl chlorpyrifos

Pesticides[a]	LOD (ppb)	<LOD (N)	Minimum	1st Quantile	Median	Mean	3rd Quantile	Maximum
Methyl Chlorpyrifos	0.31	51	--	--	--	--	--	--
Chlorantraniliprole	0.25	51	--	--	--	--	--	--
Methomyl	0.28	44	8.4	9.7	12.9	469.2	21.5	13,483.9
Metalaxyl	0.44	11	13.9	359.2	1,063.6	2,797.6	2,613.2	41,191.3
Ethyl Chlorpyrifos	0.33	3	11.6	4,403.5	8,379.1	12,670.1	14,500.2	67,759.8

[a]The units of these measurements are ng/m^3.

Measurements below the limit of detection (LOD) have been removed from the histograms since they have falsified values due to calculations.

Measurements below the LOD have been removed from the histograms and the values for the calculated ng of pesticide per cube meter of air (calculation: [(filter*2) / ((calibration of liters per minute)*minutes of spraying) (1/1000)])

Supplemental Table 1
Northern Thailand Participant Self-reported Symptoms by worker category

Variable Outcome		Comparison Worker		Farmworker		Fisher's Exact Test
		N=27	%	N=70	%	p-value
In the last two weeks, how often have you had the following conditions…						
stuffy, itchy, runny nose?	rarely or never	18	66.7	51	72.9	0.31
	occasionally	8	29.6	15	21.4	
	frequently	--	--	2	2.9	
watery, itchy eyes?	rarely or never	20	74.1	47	67.1	0.12
	occasionally	4	14.8	20	28.6	
	frequently	2	7.4	2	2.9	
sinusitis or sinus problems?	rarely or never	25	92.6	68	97.1	0.19
	occasionally	1	3.7	---	--	
	frequently	--	--	1	1.4	
pneumonia?	rarely or never	26	96.3	62	88.6	0.14
	occasionally	--	--	5	7.1	
	frequently	--	--	3	4.3	
dizziness?	rarely or never	23	85.2	45	64.3	0.03
	occasionally	3	11.1	23	32.9	
	frequently	--	--	1	1.4	
nausea/vomiting?	rarely or never	25	92.6	61	87.1	0.2
	occasionally	1	3.7	8	11.4	
	frequently	--	--	--	--	
Being absentminded, forgetful, or confused?	rarely or never	12	44.4	31	44.3	0.55
	occasionally	11	40.7	29	41.4	
	frequently	3	11.1	10	14.3	
headache?	rarely or never	17	63.0	48	68.6	0.4
	occasionally	9	33.3	20	28.6	
	frequently	--	--	1	1.4	
loss of appetite?	rarely or never	23	85.2	62	88.6	0.38
	occasionally	3	11.1	7	10.0	
	frequently	--	--	--	--	
fast heart rate?	rarely or never	25	92.6	57	81.4	0.14
	occasionally	1	3.7	10	14.3	
	frequently	--	--	1	1.4	
difficulty with balance?	rarely or never	21	77.8	59	84.3	0.24

	occasionally	5	18.5	10	14.3	
	frequently	--	--	--	--	
blurred vision or double vision?	rarely or never	16	59.3	41	58.6	0.15
	occasionally	9	33.3	25	35.7	
	frequently	--	--	3	4.3	
numbness or pins-and-needles in your hands or feet?	rarely or never	15	55.6	41	58.6	0.32
	occasionally	11	40.7	25	35.7	
	frequently	--	--	4	5.7	
shaking or trembling of your hands?	rarely or never	18	66.7	61	87.1	0.01
	occasionally	8	29.6	6	8.6	
	frequently	--	--	1	1.4	
twitches, jerks, or involuntary movements of your arms or legs?	rarely or never	20	74.1	54	77.1	0.39
	occasionally	6	22.2	12	17.1	
	frequently	--	--	1	1.4	

Supplemental Table 2
Northern Thailand Participant Stressor Responses 2017, by worker category

Variable	Variable Outcome	Comparison Worker		Farmworker		Fisher's Exact Test
		N=27	%	N=70	%	p-value
In the last month, how often have you felt that you were unable to control the important things in your life?	never	17.0	63.0	48.0	68.6	0.57
	almost never	4.0	14.8	13.0	18.6	
	sometimes	5.0	18.5	8.0	11.4	
	fairly often	1.0	3.7	1.0	1.4	
	Often	--	--	--	--	
In the last month, how often have you felt confident about your ability to handle your personal problems?	never	--	--	4.0	5.7	0.03
	almost never	--	--	2.0	2.9	
	sometimes	2.0	7.4	6.0	8.6	
	fairly often	4.0	14.8	28.0	40.0	
	Often	21.0	77.8	30.0	42.9	
In the last month, how often have you felt that things were going your way?	never	--	--	--	--	0.27
	almost never	--	--	1.0	1.4	
	sometimes	8.0	29.6	28.0	40.0	
	fairly often	6.0	22.2	22.0	31.4	
	Often	13.0	48.2	19.0	27.1	
In the last month, how often have you felt difficulties were piling up so high that you could not overcome them?	never	16.0	59.3	49.0	70.0	0.34
	almost never	8.0	29.6	11.0	15.7	
	sometimes	3.0	11.1	10.0	14.3	
	fairly often	--	--	--	--	
	Often	--	--	--	--	
	range	median		standard deviation		
How many hours do you work in a typical week in total at all of your jobs?	6,140	50.96		25.73		