# External location of touch is constructed post-hoc based on limb choice

Femke Maij[#3], Christian Seegelke[#1,2], W. Pieter Medendorp[3], Tobias Heed[1,2]

[1]Faculty of Psychology and Sports Science, Bielefeld University, Bielefeld, Germany

[2]Center of Excellence Cognitive Interaction Technology (CITEC), Bielefeld University, Bielefeld, Germany

[3]Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, Nijmegen, The Netherlands

#These authors contributed equally to this work

*Running head*: Post-hoc construction of tactile external location

*Correspondence*: Tobias Heed, tobias.heed@uni-bielefeld.de

## 1    **Abstract**

When humans indicate on which hand a tactile stimulus occurred, they

often err when their hands are crossed. This finding seemingly supports the

view that the automatically determined touch location in external space affects

limb assignment: the crossed right hand is localized in left space, and this

conflict presumably provokes hand assignment errors. Here, participants

judged on which hand the first of two stimuli, presented during a bimanual

movement, had occurred, and then indicated its external location by a reach-

to-point movement. When participants incorrectly chose the hand stimulated

second, they pointed to where that hand had been at the correct, first time point,

though no stimulus had occurred at that location. This behavior suggests that

stimulus localization depended on hand assignment, not vice versa. It is, thus,

incompatible with the notion of automatic computation of external stimulus

location upon occurrence. Instead, humans construct external touch location

post-hoc and on demand.

## 17 Introduction

Spatial perception and actions rely on multiple spatial codes, often
associated with different reference frames. For instance, the accuracy of
pointing or reaching with an arm or finger to a visual target depends not only on
the position of target relative to gaze (Fiehler et al., 2011; Thompson et al.,
2014), but also on salient world-centered landmarks (Schütz et al., 2013).
Similarly, judgment of visual location during whole-body movement is
influenced by a target's position relative to gaze, as well as by the location of
the target relative to the body (Tramper & Medendorp, 2015).

In touch, too, space is coded in several reference frames. Touch
activates specialized sensory receptors embedded in the skin, and the
arrangement of the peripheral sensors is reflected in the homuncular
organization of primary somatosensory cortex (Penfield & Boldrey, 1937; Roux
et al., 2018), referred to as a skin-based or somatotopic layout. However,
because our body can take various postures, the stimulus location in space –
often referred to as its external location – must be derived by combining skin
location and body posture, a process termed tactile remapping (Heed, Buchholz,
et al., 2015). Indeed, there is evidence that external tactile locations can be
coded in a gaze-centered reference frame (Harrar & Harris, 2010; Mueller &
Fiehler, 2014a, 2014b), but also relative to anchors such as the head, torso,
and hand (Alsmith et al., 2017; Heed et al., 2016).

It is less clear, however, according to which principles these different
spatial codes are employed. Both bottom-up features such as the availability of
sensory information (Bernier & Grafton, 2010) and the spatial reliability of a
sensory channel (Ernst & Banks, 2002; van Beers et al., 2002), as well as top-
down information such as task-constraints (Badde et al., 2015; Schubert et al.,
2017), action context (Mueller & Fiehler, 2014b), and cognitive load (Badde et
al., 2014) can affect the relative contributions of different reference frames,
presumably in a weighted manner (Angelaki et al., 2009; Atsma et al., 2016;
Badde & Heed, 2016; Ernst & Di Luca, 2011; Kayser & Shams, 2015; Lohmann
& Butz, 2017; Tramper & Medendorp, 2015). Yet, whereas there is widespread

consensus that each spatial code can have more or less influence depending

on the specific situation, it is currently not known whether all putative codes are

always constructed, or whether they are only computed based on demand.

For touch, it has been suggested that the construction of spatial location

is an automatic process, implying that any tactile input is remapped into an

external code, irrespective of its relevance (Heed & Azañón, 2014; Röder et al.,

2004). The most common experimental manipulation underlying this claim is

limb crossing. Crossing, say, a right arm over to the left side of space leads to

different skin-based (here: right body side) and external (here: left side of

space) spatial codes of a tactile stimulus delivered to the right hand. A task-

irrelevant tactile stimulus delivered to a crossed right hand accelerates visual

discrimination in the right visual field if it precedes the visual target stimulus by

60 ms, but on the left side if it leads by 180 ms or more (Azañón & Soto-Faraco,

2008). Thus, responses to visual targets were faster after anatomically

congruent tactile cues (e.g., tactile stimulus on crossed right hand, visual target

in right hemifield) at short cue-stimulus intervals, but after externally congruent

tactile cues (e.g., tactile stimulus on the left hand crossed over to the right side,

visual target in right field) at long cue-stimulus intervals. Such effects are

usually interpreted as evidence that tactile remapping – the precise

computation of the external tactile stimulus location – is automatic and forms

the basis for the performance enhancement at this external location.

The same conclusion has also been drawn from results obtained with

the tactile temporal order judgment (TOJ) task; in this task, participants report

which of two successive tactile stimuli, each presented to a different body part

– typically the two hands – occurred first (Heed & Azañón, 2014; Shore et al.,

2002; Yamamoto & Kitazawa, 2001). When the time interval between the two

stimuli is short, participants sometimes choose the wrong stimulus. Notably,

stimulus confusion is much more prominent when the arms are in a crossed

than uncrossed posture. This is surprising because the TOJ task asks about

the identity of the touched limb, and, in theory, it would be irrelevant to this

question where the hand was in space. That limb crossing, nevertheless,

affects TOJ implies that posture cannot be strategically ignored, but is
automatically incorporated into the hand assignment.

Several explanations have been put forward to account for crossing
effects in tactile localization. First, it has been suggested that touch location,
once it is remapped, is retained only in an external spatial code, and the original
skin location is discarded in the process. To report which body part has been
touched, the brain must then reversely determine which limb was located at the
computed external location at the time of the touch (Kitazawa, 2002; Kitazawa
et al., 2008; Yamamoto & Kitazawa, 2001). We refer to this suggestion as the
*space-to-limb reconstruction hypothesis*. When applied to errors in the TOJ task,
this hypothesis implies that participants correctly remap the two tactile stimuli
into external space, but then reconstruct erroneously which hand was at the
first spatial location.

A second explanation assumes that TOJ errors reflect the conflict
between different codes used for stimulus location. When the limbs are
crossed, skin-based and external spatial codes point to different sides of space,
and this conflict must be resolved, a process that takes time and is error-prone
(Röder, Kusmierek, Spence, & Schicke, 2007; Simon, Hinrichs, & Craft, 1970).
In this view, the TOJ crossing effect is a marker for the presence of conflict and,
thus, for the fact that remapping into an external spatial code has taken place.
Notably, the interpretation that the TOJ crossing effect derives from a remapped
stimulus location is indirect because participants only report a binary decision
about which hand was stimulated, not the spatial location of the perceived
stimulus. Increasing the distance between the uncrossed hands can slightly
reduce errors in TOJ (Gallace & Spence, 2005; Roberts et al., 2003; Shore et
al., 2005), and the TOJ crossing effect is smaller when the hands' positions
additionally differ in height or depth (Azañón et al., 2016). These graded
modulations of the TOJ have led to the claim that the TOJ paradigm is an
implicit index of tactile remapping (Azañón et al., 2015; Badde & Heed, 2016;
Heed & Azañón, 2014). We refer to this suggestion as the *stimulus switch*
*hypothesis*. It implies that participants have correctly remapped the two stimuli

into space, but have incorrectly resolved the conflict between the different

spatial codes of the first stimulus, consequently assigning the incorrect stimulus

to the first time point; as a consequence, participants incorrectly report the hand

that received the second stimulus.

Importantly, both hypotheses outlined above assume that touch is

automatically remapped to its veridical external location. However, recent

experiments have cast doubt on whether this is actually the case. For instance,

if a tactile stimulus is presented during an arm movement, and participants

indicate the stimulus's location by pointing to its external location after the

movement, they make systematic localization errors (Dassonville, 1995; Maij,

Grave, et al., 2011; Maij et al., 2013, 2017; Watanabe et al., 2009). Importantly,

because these errors differ for fast and slow movements, it has been suggested

that participants do not compute the precise spatial location of a stimulus when

it occurs, but instead infer spatial location post-hoc by estimating hand location

at the perceived time of the tactile stimulus (Maij et al., 2017). We refer to this

suggestion as the *time reconstruction hypothesis*. Accordingly, errors in the

TOJ task would occur because participants first choose the incorrect hand, and

then derive stimulus location based on that hand's position at the time of the

first stimulus. Note, that here participants merge the correct, first stimulus's time

with the incorrect, second stimulus's hand. For the present study, the key claim

of the *time reconstruction hypothesis* is that stimulus location is only computed

after the hand has been chosen. This feature is at odds with the idea that tactile

judgments are based on spatial remapping (Heed, Buchholz, et al., 2015; Shore

et al., 2002; Yamamoto & Kitazawa, 2001), according to which the stimulus

location is determined first and then used to make the hand assignment – in

fact, the time reconstruction hypothesis reverses the dependency between

localization and limb assignment proposed by other the theoretical accounts.

**Figure 1.** *Experimental conditions of Experiment 1 and predictions of the tested tactile localization hypotheses. A-B. Experimental procedure. A. The arms moved from an uncrossed or crossed start posture to an uncrossed or crossed arm end posture. B. Representative example TOJ trial showing the bimanual movement (grey, left hand; yellow, right hand) for the four combinations of uncrossed and crossed start and end postures, as well as the reach-to-point movement of the hand at which the first tactile stimulus was reported. C. Illustation of a correct TOJ trial: the stimulus is assigned to the correct hand, which points to the correct location. Grey (yellow) traces illustrate the left (right) hand's movement toward the body, here during a trial from an uncrossed start to an uncrossed end posture. The blue arrow indicates the movement of the correctly assigned hand towards the location of the first stimulus (cross). D-F. Illustration of the three hypotheses that may account for TOJ errors. The red arrows indicate the movement of the incorrectly chosen hand. D. Space-to-limb reconstruction hypothesis: participants point with the incorrect hand at the external location of the first stimulus. E. Stimulus switch hypothesis: participants point with the incorrect hand at the external location*

*of the second stimulus (star). F. Time reconstruction hypothesis: participants point with the*

*incorrect hand at the location at which that hand was at the time of the first stimulus.*

Here, we assessed hand assignment and spatial localization of tactile

stimuli presented during movement. Our objective was to test whether TOJ

responses mark the use of the stimulus's external-spatial location constructed

in response to the stimulus, or whether instead participants estimate stimulus

location post-hoc by integrating the hand movement trajectory with stimulus

time. In other words, we aimed to directly contrast the three discussed

hypotheses for tactile localization: the space-to-limb reconstruction hypothesis,

the stimulus switch hypothesis, and the time reconstruction hypothesis.

We presented human participants with two tactile stimuli during a

bimanual movement and assessed which hand participants perceived to have

been stimulated first (TOJ hand assignment), as well as exactly where in space

participants had perceived the first stimulus (tactile stimulus localization). The

experimental logic, and its relation to the three tested tactile localization

hypotheses, are illustrated in Fig. 1. Because tactile stimuli were presented

shortly before, after, and during the time of movement, their spatial location

depended on their timing relative to the movement. This allowed us to

determine which tactile location participants had perceived when they had

made a hand assignment error in the TOJ task. Contrary to common opinion,

TOJ errors were not associated with the location of the second, incorrect

stimulus. Instead, when participants chose the incorrect hand, they reported its

location at the time point at which the first, correct stimulus had occurred. Thus,

participants constructed stimulus location by combining the position of the

incorrectly chosen hand with the stimulus timing that belonged to the other, non-

chosen hand's stimulus, resulting in reported locations at which no stimulus had

ever occurred. This finding invalidates current explanations of crossing effects

as being based on the remapped external spatial location of the tactile stimulus.

## Results

### Experiment 1

Participants adopted a start posture with their hands resting on a table and their arms stretched out in an uncrossed or crossed posture (see Fig. 1A,B for an illustration of experimental conditions and trial timing). A tone then instructed a movement of the two hands about 30 cm towards their body, bringing the arms into either an uncrossed or crossed arm end posture (see Fig. 1A). Shortly before, during, or shortly after the movement, participants received two tactile stimuli, one on each hand, with a stimulus onset asynchrony (SOA) of 110 ms. At this SOA, participants often misreport which of the two stimuli occurred first, both when the arms are still (Heed & Azañón, 2014; Shore et al., 2002; Yamamoto & Kitazawa, 2001) and during movement (Heed, Möller, et al., 2015; Hermosillo et al., 2011). After the bimanual movement, participants reported on which of the two hands the first stimulus had occurred by reaching with this hand to the perceived external location of the stimulus (see Fig. 1B). The response, thus, contained two components: the hand to which the first stimulus was assigned, and explicit spatial localization of this stimulus.

### Hand assignment

In a first step, we verified that hand assignment in the TOJ task was modulated by hand crossing and timing of stimuli relative to the movement (Heed, Möller, et al., 2015; Hermosillo et al., 2011). We measured TOJ performance as the percentage of correct reports of which hand had been stimulated first in the TOJ task, as indicated by the hand that participants used for their localization response (see Fig. 2). Stimuli could occur during all times (see Methods for details), so we binned the binary (correct/ incorrect) TOJ response data into four movement phases – stimulation before movement onset, during first and second half of movement, and after movement offset – to assess the modulation of TOJ performance by stimulus time relative to the bimanual movement.

**Figure 2.** *Proportion of correct hand assignment across movement conditions (uncrossed-uncrossed, uncrossed-crossed, crossed-uncrossed, crossed-crossed) in the four phases of the bimanual movement (before movement, during start posture, during end posture, after movement) in Experiment 1. For conditions without a postural change (i.e., uncrossed-uncrossed, crossed-crossed), trials were assigned as "during start posture" if the first stimulus occurred during the first temporal half of the movement and they were assigned as "during end posture" if the first stimulus occurred during the second temporal half of the movement. For conditions with a postural change (i.e., uncrossed-crossed, crossed-uncrossed), trials were assigned as "during start posture" if the first stimulus occurred before the postural change and as "during end posture" if the first stimulus occurred after the postural change. Error bars denote 2 s.e. from the mean; asymmetry is due to nonlinear conversion from the GLMM's logit scale to percentage correct. Large symbols are group means, small symbols are individual participants' performance.*

In accordance with previous findings, TOJ performance declined in the crossed compared to the uncrossed posture (Heed & Azañón, 2014), and depended on the posture at the time of stimulation (Heed, Möller, et al., 2015; Hermosillo et al., 2011). For instance, for the uncrossed-uncrossed movement condition (see Fig. 2, dark-green data points), the probability of a correct response was high compared to the crossed-crossed movement condition (see Fig. 2, light magenta data points) throughout all movement phases. For the conditions with a postural change (uncrossed-crossed, crossed-uncrossed, see Fig. 2, dark-magenta, light-green data points) the probability of correct responses was modulated by the posture at the time of stimulation. A generalized mixed model (GLMM) with factors Start Posture, End Posture, and Movement Phase revealed significance for all main effects and interactions (see Supplementary

Table 1). With movement phase, the effect of Start Posture (see Fig. 2, dark vs.

light colors) declined, whereas the effect of End Posture (see Fig. 2, green vs.

magenta colors) increased. For instance, for movements from an uncrossed to

a crossed posture, TOJ performance was better during the first two movement

phases, that is, when the hands were still uncrossed, than during the last two

movement phases, that is, when the hands were crossed.

In sum, TOJ performance in our first experiment reflected known

modulations of hand posture and movement timing. Participants made, on

average, more than 15% TOJ errors even with uncrossed hands. This high error

rate is due to the use of the short SOA of 110 ms (Heed, Möller, et al., 2015;

Heed & Azañón, 2014; Shore et al., 2002; Yamamoto & Kitazawa, 2001), and

is an intended outcome of the experimental paradigm, allowing, as a next step,

comparison of spatial localization responses for incorrect TOJ trials across all

postures.

**Explicit tactile localization in space is unaffected by hand posture**

Having verified that TOJ hand assignment showed the well-known

effects of posture, we next turned to tactile stimulus localization. Localization

errors are computed as the spatial difference (calculated as signed difference

in the direction along the path of the reporting hand, see Methods for details) of

the perceived stimulus location and the hand's true position at stimulus

presentation. From previous studies involving single stimuli and unimanual

movements, it is known that participants make systematic localization errors

when they retrospectively point to the spatial location of a tactile stimulus that

was presented while the target limb was moving. More specifically, localization

is systematically biased in the direction of the movement during the initial part

of a movement, and in the opposite direction during the final part of the

movement (Dassonville, 1995; Maij et al., 2013, 2017; Watanabe et al., 2009),

resulting in systematic localization error curves with positive values indicating

errors in movement direction and negative values indicating errors in the

opposite direction. This pattern of movement time-related directional biases

was evident also in the present data (see Fig. 3A for an example of a single

participant). Critically, bias was comparable across all four posture conditions

(see Fig. 3B).

To validate that localization behavior in our task was not biased by the

specifics of the TOJ task, participants performed a simpler 1-stimulus control

task in separate blocks of the experiment. While making bimanual movements

with uncrossed and crossed start and end postures, they received a single

tactile stimulus and pointed to it, as in the 2-stimulus task (see Methods for

details). Participants virtually always indicated correctly which hand had

received the stimulus (average percentage correct, 99.5%). Critically,

localization error curves were indistinguishable from the task with two stimuli

(see Fig. 3C), indicating that tactile localization was affected neither by task

difficulty nor by the nature of the TOJ task.

**Figure 3.** *Localization errors systematically vary with the progression of the movement. A.*

*Mean localization error (teal line) of all correct TOJ trials of a single participant in Experiment*

*1. Each dot represents the localization error of a single trial, that is the difference between*

*actual hand position at the time of tactile stimulation and indicated hand position at the end of*

*the trial. Blue shading represents the average movement time, with 0 ms = movement start.*

*Note, that the localization error is positive at the beginning of the movement, indicating error in*

*the direction of the movement. Localization error is negative towards the end of the movement,*

*indicating error against the direction of movement. B-C. The localization error pattern of the*

*correct trials in A was evident across all subjects for both the 2 Stimulus experimental task (B)*

*and for the 1 Stimulus control task (C) and for all posture combinations. Traces reflect the mean,*

*shaded areas around the traces reflect s.e.m. The shaded regions in the background indicate*

*the average movement duration, which differed shlightly between conditions (see*

*Supplementary Table 2).*

**Explicit tactile localization is directed towards the assigned hand**

We have so far assessed performance in trials in which participants had
made a correct TOJ hand assignment (referred to as correct TOJ trials from
hereon). We now turn to localization errors in incorrect TOJ trials. These errors
allow differentiating between the three hypotheses about how participants
determine stimulus localization in tactile decision paradigms (see Fig. 1D-F).

We first turn to the *space-to limb reconstruction hypothesis*. It posits that
tactile perception takes place in space rather than on the body; thus, a limb
assignment entails computing which limb was at the first spatial location. Thus,
in our task, responses with the incorrect hand would result from assigning the
incorrect hand to the correct spatial location of the first tactile stimulus (see Fig.
1D). Accordingly, the assigned, incorrect hand should be directed to the
location at which the stimulus of the other, correct hand had occurred, and the
reported stimulus location in incorrect TOJ trials should scatter around the
movement trajectory of the correct hand. Contrary to this prediction, participants
consistently pointed to locations scattered around the movement trajectory of
the assigned, incorrect hand, indicating that the chosen stimulus had been
perceived on the incorrect hand (see Fig. 4 for the localization responses of the
participant with the largest variability in localization errors). Thus, localization
behavior did not support the implication of the *space-to-limb reconstruction*
*hypothesis* that the correct external spatial location is simply assigned to a
wrong limb.

**Figure 4.** *Localization responses (i.e., finger positions in the horizontal plane at the end of the reach-to-point movement indicating the location where the participant perceived the first stimulus) for the different movement conditions. Data are from a single participant with the largest variability. Ellipses represent 95% of the variability and show large overlap for correct and incorrect TOJ trials. The space-to-limb reconstruction hypothesis would predict that, during error trials, participants point with the incorrectly assigned hand to the location of the correct stimulus; thus, if this hypothesis were correct, orange ellipses should overlay with light grey ellipses, and dark grey ellipses should overlay with yellow ellipses.*

## Localization aims at the assigned hand's position at the time of the first tactile stimulus

Given that participants appear to have perceived the stimulus at the assigned hand, two possibilities remain as to which stimulus location was associated with erroneous responses (see Fig. 1). The *stimulus switch hypothesis* posits that the two stimuli were localized correctly, and one is chosen for the response. In incorrect TOJ trials, participants would confuse the two stimuli and report the second stimulus by pointing at its location with the respective, incorrect hand. In this case, participants should point to where the hand was positioned at the time point of the *second*, erroneously chosen

stimulus (referred to as time 2 from hereon). In contrast, the *time reconstruction*

*hypothesis* assumes that participants always use the correct, first time point

(time 1 from hereon), and determine the position of the assigned response limb

at this time point. This hypothesis predicts that, in incorrect TOJ trials,

participants point to where the incorrectly assigned hand was positioned at the

correct time, that is, time 1. Note, that no tactile stimulus occurred at this

external spatial location, because it combines the time of the first, correct

stimulus with the movement trajectory of the second, incorrect stimulus's hand.

We test between the predictions of these hypotheses by comparing the

localization error curves in correct and incorrect TOJ trials. In the case of correct

TOJ trials, we assume that participants aimed, as instructed, at the position of

the correct hand at time 1. Therefore, we derive the localization error curve as

the spatial difference of perceived location and hand position at time 1 (see Fig.

5, dark blue lines). However, we can also derive a localization error curve for

correct trials under the assumption that participants pointed towards the hand's

position at time 2. To derive this hypothetical curve, we calculated the spatial

difference of participants' localization responses and the hand's position at time

2, rather than time 1 (see Fig. 5, light blue lines). The time 2 error curve is

shifted to the left, or "backwards" in time, relative to the time 1 error curve. This

is because, for the time 2 curve, the assumed "true" target location is the hand's

position 110 ms further into the movement, due to the SOA between the two

tactile stimuli. Accordingly, each assumed target location is closer to the

movement's end by the trajectory the hand has moved during the 110 ms

interval between the two stimuli.

The first, time 1 error curve can now serve as a template of a localization

error curve if the participant truly aimed at the hand's position at time 1. The

second, time 2 error curve, in contrast, serves as a template of a localization

error curve if the participants had truly aimed at the hand's position at time 2.

For incorrect TOJ trials, we do not know whether participants aimed at where

the incorrectly chosen hand was positioned at time 1 or at time 2. The rationale

of our analysis, thus, is to compare the localization error curves of incorrect TOJ

trials with the template localization error curves derived from correct TOJ trials

(see Fig. 5 and Methods).

**Figure 5**. *Localization curves, averaged across participants, for each of the four posture*

*conditions in Experiment 1. Curves of incorrect TOJ trials (red) show a similar pattern as the*

*localization curves of the correct TOJ trials at time 1 (dark blue), but not as the localization*

*curves of the correct TOJ trials at time 2 (light blue). Traces reflect the mean localization error,*

*shaded areas around the traces reflect s.e.m. across participants. The shaded regions in the*

*background represent the average movement time.*

Localization errors of incorrect TOJ trials overlapped with localization

errors of correct TOJ trials at time 1 for each of the four start and end posture

combinations. To quantify this further, we computed the temporal shift required

to align localization errors of incorrect TOJ trials with those of correct TOJ trials

(see Maij et al., 2009). If, in incorrect TOJ trials, participants aimed for the

incorrect hand's position at time 1, then the temporal shift should be zero

relative to the localization error curve in correct TOJ trials relative to time 1;

furthermore, it should be about -110 ms (negative denoting a shift towards left,

see above) compared to the localization error curve of correct TOJ trials relative

to time 2. If, however, participants aimed for the incorrect hand's position at

time 2, the shift pattern should be exactly reverse, that is, zero compared to the

second template curve, and around +110 ms compared to the first template

curve.

*Figure 6.* *Time shift of stimulus localization error in TOJ error trials relative to time 1 (left panel)*

*and time 2 (middle panel) for the four posture conditions in Experiment 1. The temporal shift of*

*the localization error curve was significantly different from zero when calculated relative to time*

*2, but not relative to time 1; this result is consistent with the time reconstruction hypothesis, but*

*not with the stimulus switch hypothesis. This pattern was similar across all participants as*

*demonstrated by the differences in time shift between time 2 and time 1 (right panel). Data are*

*visualized with raincloud plots (Allen et al., 2019) that display probability density estimates,*

*condition averages (large symbols), and individual participants (small symbols). Error bars*

*denote 95% confidence intervals.*

Fig. 6 displays the temporal shift between the localization error curve of

incorrect TOJ trials and the error curves at time 1 and time 2 derived from

correct TOJ trials.

We fitted a linear mixed model with two factors: factor Posture Condition

coded the four combinations resulting from uncrossed and crossed start and

end postures. Factor Reference Time Point coded whether the localization error

curve for correct TOJ trials was computed relative to time 1 or time 2. The

dependent variable was the time shift that best aligns the localization error

curve between correct and incorrect TOJ. This analysis revealed a significant

effect of Reference Time Point ($\chi^2(9,10) = 15.55$, p < 0.001), indicating that the

time shift required to align the localization error curves of correct and incorrect

TOJ trials differed depending on whether localization error of correct TOJ trials

was computed based on time 1 or time 2. In contrast, there was no effect of

Posture Condition ($\chi^2(7,10) = 1.57$, p = 0.67) or interaction ($\chi^2(7,10) = 4.53$, p

= 0.21) between the two factors. This latter result indicates that the relationship

of localization in correct and incorrect TOJ trials held across all postures; this

result is also illustrated by the very similar relationship of the different

localization error curves in the four panels of Fig. 5. Thus, whereas limb posture

affected hand assignment, it did not affect tactile localization.

To assess whether localization in incorrect TOJ trials aimed at a location

related to time 1 or time 2, we tested the respective times shifts required to align

the localization errors of the two types of trials against zero. The time shift

between localization errors for correct TOJ trials at time 1 and incorrect TOJ

trials, averaged across the four posture conditions, was 6 (s.e., 3) ms; if this

value is not significantly different from 0, then an LMM of only this condition

should not improve by inclusion of an intercept, as the latter would model the

deviation of the average shift away from 0. The comparison of a model with and

without intercept did not provide statistical evidence to reject a zero time shift

($\chi^2(1) = 0.84$, p = 0.36). Null findings are difficult to interpret in the context of

frequentist statistics. Therefore, we complemented our analysis by a Bayesian

analysis comparing a model with only a random participant factor with a model

that, in addition, included a population intercept, equivalent to the linear mixed

model reported above. The population-level intercept estimate was 6 ms and

the 95% confidence range [-8.28; 19.27 ms] included 0. Model comparison via

leave-one-out cross-validation found themodel without intercept to be more

credible than the model with the population intercept, that would have been

indicative of a non-zero localization error shift between correct and incorrect

TOJ tials (difference of expected log predictive density, ELPD, for second as

compared to first model: -0.6, s.e. 0.9).

We ran the same analyses for the time shifts required to align localization

error curves between correct and incorrect TOJ trials when correct trials' error

curve had been calculated relative to time 2. In contrast to the results for time

1, the average time shift between localization errors for correct TOJ trials at

time 2 and incorrect TOJ trials was -105 (s.e., 7) ms, and a model without

intercept fit this condition siginificantly worse than a model with the intercept

($\chi^2(1)$ = 19.38, p < 0.001). The significant difference to time 2 suggests that

participants did not aim at the position of the second stimulus in incorrect TOJ

trials.

While neither the non-significant difference to time 1 in the LMM analysis,

nor the Bayesian parameter estimate including 0 statistically imply equality of

the error curves in correct and incorrect TOJ trials, these statistical results are

consistent with the two conditions being equal, and they suggest that, if a

difference exists, it is small. Furthermore, the time shift of -105 ms for time 2

closely matches the stimulus SOA of 110 ms, further suggesting that, in error

trials, participants did not aim for hand location at the second, but rather at the

first time point. Corroborating this conclusion, the Bayesian 95% interval [-138;

-76 ms] of the intercept estimate includes -110 ms, and comparison of Bayesian

models for time 2 with and without intercept strongly favored the model

including the intercept (difference in ELPD from first to second model, -1.4, s.e.

1.9).

**Experiment 2**

The results of Experiment 1 suggest that when asked to localize the external

location of the first of to two tactile stimuli applied in succession to different

hands, participants chose which hand received the stimulus and then inferred

the position of the chosen limb at the time point of the first stimulus.

Consequently, when participants chose the incorrect limb, stimulus location

was determined as the location at which the incorrect hand was at the correct

(first) time point. While these results support the *time reconstruction hypothesis*,

it should be realized that Experiment 1 tested only a single SOA of 110 ms

between the two tactile stimuli. If our conclusions drawn from Experiment 1 are

correct, then localization of stimuli assigned to the incorrect hand should always

depend on the first stimulus's time, independent of SOA.

Experiment 2 tested this conjecture. Again, participants judged which hand had

received the first of two tactile stimuli during a bimanual movement and then

located the stimulus perceived to have occurred first. We presented tactile

stimuli with four different SOAs: 60, 85, 110, and 135 ms. As explained in

Experiment 1, the shift between the time 1 and time 2 curves of correct TOJ

trials reflects the SOA of the two tactile stimuli. Accordingly, the two template

curves are further apart the larger the SOA (compare light vs. dark blue lines in

panels A-D of Fig. 7). As the estimated localization error curves in Experiment

1 were similar for all combinations of the hands' start and end posture,

Experiment 2 involved only reaches from an uncrossed to an uncrossed posture

and from a crossed to a crossed posture. This strategy minimized marker

obstruction and homogenized movement time across conditions (see

Supplementary Table 3). Experiment 2 was conducted in a different lab than

Experiment 1, using different equipment, re-written experimental code, different

experimenters, and new analysis scripts (see Material and methods for details).

To further scrutinize the reliability of our results, we increased our sample size

and acquired a higher number of trials.

**Hand assigment**

In accordance with Experiment 1 and previous findings (Heed, Möller, et al.,

2015; Hermosillo et al., 2011), TOJ performance in Experiment 2 was

modulated by hand posture and SOA. At all SOA, participants made large

amounts of errors, ensuring that a sufficient number of trials were available to

analyze incorrect TOJ trials. Detailed results are reported in the Supplementary

Information (Supplementary Figure 1 and Supplementary Table 4).

**Explicit stimulus localization in space**

Complementing the findings from Experiment 1 and further corroborating the

*time reconstruction hypothesis*, localization errors of the incorrect TOJ trials

largely overlapped with the localization errors of correct TOJ trials at time 1 for

each of the four SOAs (see Fig. 7) and for each subject and posture condition

(see Supplementary Fig. 2-4).

***Figure 7.*** *Localization curves, averaged across participants and posture, for each of the four*

*SOAs in Experiment 2. Curves of incorrect TOJ trials (red) show a similar pattern as the*

*localization curves of the correct TOJ trials at time 1 (dark blue), but not as the localization*

*curves of the correct TOJ trials at time 2 (light blue). This pattern was highly similar across all*

*subjects and also when calculated separately for each posture condition (see Supplementary*

*Information). Traces reflect the mean, shaded areas around the traces reflect s.e.m. The*

*shaded regions in the background represent the average movement time.*

Fig. 8 shows the temporal shift between the localization error curves of incorrect

TOJ trials and the error curves of correct TOJ trials relative to time 1 and time

2 four all four SOAs.

*Figure 8.* Time shift of stimulus localization error in incorrect TOJ trials relative to time 1 (left panel) and time 2 (middle panel) for the four SOAs in Experiment 2. For all SOAs, the temporal shift relative to time 1 was not significantly different from 0. In contrast, the time shift was significantly different from zero for all SOAs when calculated relative to time 2, and it was numerically similar to the respective SOA. These results are consistent with the time reconstruction hypothesis, but not with the stimulus switch hypothesis. This pattern was similar across all participants as demonstrated by the differences in time shift between time 2 and time 1 (right panel). Data are visualized with raincloud plots (Allen et al., 2019) displaying probability density estimates, condition averages (large symbols), and individual participants (small symbols). Error bars denote 95% confidence intervals.

A linear mixed model with factors SOA and Reference Time Point (localization errors for correct TOJ trials computed relative to time 1 vs. time 2) and time shift of the error curve between correct and incorrect TOJ trials as dependent variable revealed a significant main effect of Reference Time Point ($\chi^2(9,10) = 21.33$, p < 0.001) and a significant Reference Time Point × SOA interaction ($\chi^2(7,10) = 9.90$, p = 0.02). The time shift between localization errors for correct TOJ trials at time 1 and incorrect TOJ trials, averaged across all four SOA conditions, was 8 ms. As in Experiment 1, a LMM with a common intercept for all SOAs did not fit the data better than a model without an intercept ($\chi^2(1)$ = 1.24, p = 0.27). Similarly, allowing for individual intercepts per SOA did not improve the goodness of fit, ($\chi^2(3) = 2.74$, p = 0.43). Thus, none of the tested models provided statistical evidence to reject a zero time shift with respect to time 1. These results were again supported by parameter estimates of Bayesian models equivalent to the afore-mentioned LMMs, which estimated both an intercept across all SOAs and individual intercepts per SOA to lie in

intervals that included 0 (see Supplementary Information). Model comparison

via leave-one-out cross-validation found a model without population intercept

to be more credible than a model with a common intercept (difference of

expected log predictive density, ELPD: -0.2, s.e. 0.6) and a model with

individual intercepts per SOA (ELPD: -2.2, s.e. 2.2).

In contrast, the average time shifts between localization errors for correct

TOJ trials at time 2 and incorrect TOJ trials were -52 (s.e., 12) ms, -75 (s.e.,

12) ms, -87 (s.e., 14) ms, and -104 (s.e., 13) ms for the SOA 60 ms, 85 ms, 110

ms, and 135 ms, respectively. A model with a common intercept for all SOAs

explained the data significantly better than a model without an intercept ($\chi^2(1)$

= 29.27, p < 0.001). A model allowing for different intercepts for each SOA

further improved the goodness of fit ($\chi^2(3)$ = 14.3, p = 0.003), indicating that

localization curve's time shift relative to time 2 depended on the respective SOA

(see Supplementary Table 5 for Bayesian model estimates). Model comparison

via leave-one-out cross-validation found a model with individual SOA intercepts

to be more credible than a model with a common intercept (difference of

expected log predictive density, ELPD: -5.4, s.e. 3.8) and a model without

intercept (ELPD: -7.2, s.e. 4.7).

In sum, localization curves reflected the increase of, and shift values

were numerically close to, the stimulus SOAs. Yet, the time shift value for the

110 ms SOA in Experiment 2 was smaller than that of Experiment 1 (-87 vs. -

105 ms). In fact, average time shift values of Experiment 2 seemed to

underestimate the true SOA in Experiment 2, although the Bayesian 95%

intervals of the intercept estimates included the true SOA for all but the largest

SOA (i.e., 135 ms, see Supplementary Table 5). Furthermore, at time 1, the

estimated intercepts were all slightly (albeit non-significantly) positive (Fig. 8 left

panel, Supplementary Table 5), and when one considers the difference in time

shift (i.e., Time 2 – Time 1), the estimated values match the true SOA more

closely (-63 ms, -81 ms, -88 ms, and -120 ms for SOAs 60 ms, 85 ms, 110 ms ,

and 135 ms, respectively; see Fig. 8, right panel). We note that time shift

calcluations are based on a sliding Gaussian average across noisy, non-linear

patterns of localization errors, and so absolute shift values may not exactly

reflect the stimulus SOAs.

## Discussion

The aim of our study was to test whether participants represent the

remapped spatial location of tactile stimuli when they make spatial decisions

about tactile stimuli. Participants indicated both the target limb and the

perceived location in space of the first of two tactile stimuli in a tactile TOJ task.

Presentation of stimuli during movement implied that stimulus location

depended on stimulation time, allowing us to determine the relationship of

stimulus timing and perceived stimulus location in space. If participants had first

computed the spatial location, and then derived which limb had occupied this

location at time of stimulation (space-to-limb hypothesis, see Kitazawa, 2002),

incorrect hand assignment should have been associated with external

localization along the trajectory of the correct hand; we did not find any

evidence for such behavior. If participants had represented stimulus location

and stimulated limb together, and simply confused the two due to conflict

between different spatial codes (*stimulus switch hypothesis*), incorrect hand

assignment should have been associated with spatial localization at the location

of the incorrect limb at the second stimulus's time. Localization error curves

were incompatible with this view, as their systematic bias differed significantly

from a hypothetical localization curve, derived from correct trials, relating to

stimulus time 2. Instead, when participants chose the incorrect hand, their

localization errors implied that they had aimed at that hand's location at the time

of the first, correct stimulus, evident in a close match of localization curves of

correct and erroneous TOJ trials when computed relative to stimulus time 1. In

other words, participants derived the reported stimulus location by combining

the time of the first, correct stimulus with the trajectory of the second, incorrectly

chosen hand, effectively indicating a location at which no stimulus had occurred

– consistent with the *time reconstruction hypothesis*. This behavior was evident

for all combinations of uncrossed and crossed start and end postures, as well

as for all tested SOAs between the two tactile stimuli. TOJ errors, thus, did not

simply reflect temporal confusion of two stimuli; instead, localization in TOJ

error trials marks the computation of tactile stimulus location based on correct

stimulus timing and movement information of a (correctly or incorrectly) implied

body part. Accordingly, limb crossing affected hand assignment, but not

stimulus localization.

The pattern of hand assignment errors was in line with previous studies:

Participants made more TOJ hand assignment errors in conditions that involved

hand crossing than in conditions in which the hands were uncrossed (see Fig.

3; Heed, Möller, et al., 2015; Hermosillo et al., 2011; Shore et al., 2002;

Yamamoto & Kitazawa, 2001). These reliable findings support the interpretation

that categorical decisions in touch, such as choosing which limb was

stimulated, are affected by weighted integration of different spatial aspects of

the tactile stimulus and configuration of the body (Badde et al., 2019; Badde &

Heed, 2016; Heed & Azañón, 2014). In contrast, localization error patterns were

similar across uncrossed and crossed start and end hand posture conditions,

suggesting that arm posture during stimulation did not affect localization

responses (see Fig. 4, 5). In particular, localization errors exhibited comparable

spatial biases over time in uncrossed and crossed conditions. Furthermore,

localization error scattered around the chosen hand was not biased towards the

other hand (see Fig. 4), an effect one might have expected if, like hand

assignment, spatial localization was subject to weighted influence of the tactile

stimulus's anatomical origin as coded by a body-based reference frame.

The dissociation between TOJ hand assignment and localization

responses indicates that the two phenomena do not reflect the same process.

It is widely assumed that the weighted integration of spatial factors reflected by

tactile limb crossing effects trades off the anatomical and the external location

of a tactile stimulus (Badde & Heed, 2016; Cadieux & Shore, 2013; Kitazawa,

2002; Shore et al., 2002; Yamamoto & Kitazawa, 2001). This assumption

requires that an external location is constructed as a prerequisite for assigning

a stimulus to a hand. Our present finding that participants incorrectly localize

tactile stimuli associated with incorrect limb choice, in contrast, implies that not

stimulus location determines hand assignment, but vice versa, hand

assignment determines perceived stimulus location.

This conclusion is incompatible with the common view that crossing

effects, obtained in experiments that require categorical decisions such as TOJ,

are an implicit indicator of precise tactile localization and tactile remapping. This

is a strong claim that may invalidate the experimental logic of numerous papers

that have applied this logic. However, the present results are corroborated by

another recent study that has challenged the view that errors in tactile

categorical response paradigms reflect a conflict between anatomical and

external spatial coordinates. In that study, participants performed TOJ of tactile

stimuli presented to uncrossed and crossed hands and feet (Badde et al.,

2019). In each trial, two stimuli were randomly presented to two of the four

limbs. In some trials, participants reported the first touch on a limb that had not

been stimulated in this trial. For instance, after stimulation of the left hand and

the right foot, a participant may have indicated that the first stimulus had

occurred on the right hand. These TOJ errors systematically depended on

different anatomical features such as the type (hand or foot) of the correct limb

and its body side. Critically, neither the side of space of the limb that had

received the correct stimulus, nor the spatial distance between stimulus and

response limb affected TOJ errors in this task. Like the present results, these

findings are incompatible with the prevailing view that crossing effects reflect

conflict during the integration of anatomical and spatial stimulus location. In fact,

the two studies complement each other in that we show here that stimulus

location is not used for hand assignment, and Badde et al. (2019) suggest

which information is instead used to choose between the two hands in a

categorical tactile-spatial task.

Notably, other manipulations that have been used to argue for the

relevance of precise spatial representations in tactile decisions can be framed

in such a feature-based account as well. For instance, the TOJ crossing effect

was reduced when the two hands' positions differed in height or in depth

(Azañón et al., 2016). However, this manipulation may have simply introduced

an additional, non-metric feature that helped representing the two choice

options as different, thus improving TOJ choice based on a categorical feature

rather than on metric distance. Similarly, whereas effects of distance between

the two hands during a TOJ have been interpreted as implying a metric

representation of stimulus location (Roberts et al., 2003; Shore et al., 2005),

others have reported such effects only for very small (3cm), but not other (10

cm and larger), distances between the hands (Kim & Cruse, 2001), again

suggesting that they may reflect categorically coded spatial features, not metric

stimulus location.

The reason for the apparent contradiction between previous research

and these new findings stems from the fact that typical experimental designs,

in which tactile stimuli are presented to the two stationary hands, do not allow

disentangling the prevailing view of crossing effects indicating conflict between

tactile anatomical and (precise) spatial stimulus location from the view that the

conflict apparent in limb crossing must indicate other spatial aspects, such as

those identified in our previous study (Badde et al., 2019). Thus, limb crossing

paradigms that require limb choices about the origin of touch likely reflect the

integration of categorical, tactile-spatial stimulus features. We propose that

automatic effects, such as crossmodal, tactile-visual cueing (Azañón & Soto-

Faraco, 2008), too, are based on such feature-based processing. In contrast,

precise stimulus location is, contrary to what has regularly been implied, not

among the pieces of information that are integrated for automatic, tactile-spatial

coding. Instead, precise location of the tactile stimulus in space is inferred post-

hoc only when required.

Furthermore, our proposal contrasts with the suggestion that TOJ errors

are due to temporal confusion of the two stimuli, hypothesized to occur due to

slowing of a neural clock mechanism because crossed postures induce higher

cognitive load as compared to uncrossed postures (Kitazawa et al., 2008). This

hypothesis is based on the assumption that tactile locations are represented

correctly but ordered incorrectly in time. It is incompatible with our finding that

participants localized touch in incorrect TOJ trials at the location of the incorrect

hand at the correct, first stimulus time point. Moreover, an account based on

time confusion is specific to the TOJ paradigm, in which participants compare

two stimuli. In contrast, a feature-based account generalizes to other

experimental paradigms, including ones that present only a single tactile

stimulus (Azañón et al., 2010; Azañón & Soto-Faraco, 2008; Badde et al., 2015,

2019).

701           Our study exploited systematic localization errors when a stimulus is

presented during a movement of the arm. A modulation of spatial localization

by movement of the respective sensors is not unique to touch. When a brief

flash is shown during a smooth pursuit or saccadic eye movement, its

localization is perceived with a bias in the direction of the eye movement shortly

before and during the first half of the eye movement, and in the opposite

direction at the end of the eye movement (Matin & Pearce, 1965; reviewed by

Schlag & Schlag-Rey, 2002). As suggested here for touch, visual

mislocalization during saccades, too, depends on temporal processing. For

instance, irrelevant auditory temporal information can influence the perceived

location of a flash near the time of saccades and result in a temporal shift of the

visual localization error curve (Binda et al., 2010; Maij et al., 2009). Moreover,

when a red flash was presented around the time of a saccade on a split

green/red background, participants sometimes reported that the red flash had

occurred on the (same-color) red background; these reports of an objectively

impossible perception (flash on same-colored background) were best explained

by integration of temporal uncertainty of the flash's timing and eye position

(Maij, Brenner, et al., 2011). A computational model for these temporal-spatial

phenomena faithfully replicates the observed spatial biases for both vision and

touch. Illustrated for the case of tactile localization on the arm, the model

assumes temporal uncertainty about tactile stimulus occurrence relative to the

arm movement and combines a probability distribution of the possible stimulus

time with the perceived arm movement trajectory (Maij et al., 2013, 2017; see

Maij, Grave, et al., 2011, for the visual analogue). The present results, too, are

compatible with the temporal uncertainty model. Independent of which hand the

stimulus was assigned to, the temporal estimate of the tactile stimulus was

identical, resulting in identical localization error profiles, based on the trajectory

of the chosen hand, in correct and incorrect TOJ trials. Thus, the time-based

mechanism that leads to the seemingly surprising perception of spatial

locations at which no stimulus really occurred may be task- and domain-

general.

To summarize, we observed the typical dependence of tactile TOJ

responses on limb posture, with higher error rates when the hands are crossed

rather than uncrossed. Explicit localization responses of the stimulus chosen

as having occurred first were incompatible with theoretical accounts that posit

confusion of yoked stimulus representations that encompass the independently

determined external-spatial location of tactile stimuli, or projection of a body

part onto the determined spatial location of a stimulus. Instead, participants

chose one hand presumably based on categorical, spatial stimulus

characteristics such as the stimulated body side (Badde et al., 2019), and then

combined the time point associated with the first stimulus with the chosen arm's

trajectory. After hand assignment errors, participants, thus, effectively

referenced a post-hoc constructed spatial location at which no stimulus had

ever occurred.

## Materials & methods

746   Data for the presented analyses as well as code to run analyses and create

747   figures are provided at the Open Science Framework website,

748   https://osf.io/ybxn5/.

### Participants

Experiment 1 was performed at the Faculty of Psychology and Human

Movement Science of the University of Hamburg. Twelve right-handed

participants (aged 19-31 years, 7 female) gave informed consent to take part in

the experiment. The study was part of a research program approved by the

ethics committee of the German Psychological Society (DGPs). Experiment 2

was        preregistered      at       the       Open       Science      Framework      website

(https://osf.io/qyzgb). A sample size of 20 participants was defined a priori. We

collected data from 20 individuals from Bielefeld University. We excluded data

of 1 participant from analyses as s/he did not follow the instructions and most

of the time localized the tactile stimulus at the start or end position, but not along

the movement trajectory. Furthermore, we excluded data of another participant

as s/he only completed 384 trials in total. As any form of data acquisition was

stopped in our lab beginning of March 2020 due to the spread of the corona

virus, we did not collect data from replacement participants. Our sample thus

constisted of 18 participants (aged 18-25 years, 15 female). The experiment

was approved by the ethics committee at Bielefeld University (Ethical

Application Ref: 2017-114).

Participants provided written informed consent and were compensated with

€7/hr or received course credit. All participants had normal or corrected-to-

normal vision and did not have any known perceptual, motor, or neurological

disorders. Participants took part only if, in a screening experiment, they

exhibited a TOJ crossing effect at the SOA used in the main experiment. We

used this screening procedure because individual response patterns in tactile

experiments involving hand crossing are quite variable (Badde et al., 2015;

Cadieux et al., 2010; Yamamoto & Kitazawa, 2001); however, crossing effects

are highly reliable across the entire population, so that our screening procedure

does not preclude generalization.

**General setup**

Participants were blindfolded. They sat on a chair at a table. A tactile stimulator

(Oticon BC 461-0/12, Oticon Ltd., London, UK) was attached to the phalanx

media of each index finger. Stimulation consisted of 200 Hz vibration for 10 ms.

To mask any noise of the vibrators, participants wore earplugs and heard white

noise through speakers (Experiment 1) or wore sound-attenuating headphones

(Superlux HD669, Superlux Enterprise Development, Shanghai, China;

Experiment 2).

**Experiment 1**

***Apparatus, task and procedure***

The position of each index finger in space was recorded with an Optotrak active,

infrared marker motion tracking system (Northern Digital Inc., Waterloo,

Ontario, Canada) at a sampling rate of 1000 Hz. One marker was positioned

on the nail of each index finger, directly next to the tactile stimulator. A data

acquisition unit (Odau; Northern Digital Inc., Waterloo, Ontario, Canada;

sampling rate 1000 Hz) synchronized marker position and timing of the tactile

stimuli. The experiment was controlled with Matlab (Mathworks, Natick, MA,

USA), using the Psychophysics Toolbox (Brainard, 1997) and the Optotrak

Toolbox (http://webapp6.rrz.uni-hamburg.de/allpsy/vf/OptotrakToolbox).

*2 Stimulus Task:* Participants moved both hands from a position of about 40 cm

away from their body towards their body to a position about 10 cm away from

their body. Hand start and end posture were uncrossed and crossed, varied in

blocks of 50 trials in pseudo-randomized order (see Fig. 1A). In each trial, a

tone instructed the movement start. At a random time (presented between 50-

800 ms after the tone, drawn from a square distribution) before, during, or after

the movement, two tactile stimuli were applied, one to each hand, at an SOA of

110 ms; the left-right order of stimuli was pseudo-random. Upon movement

completion, participants moved the index finger that they had perceived to have

been stimulated first to the location of the first stimulus on the table. The hand

remained in this location until a tone, presented 2.5 s after the initial movement

cue instructed them to lift the index finger; this finger lift was used to identify the

response hand during trajectory analysis. Subsequently, participants

repositioned the hands to their start locations. We acquired 300 trials of each

posture combination. To compensate for obstruction of motion tracking

markers, we acquired more trials for 2 participants in the uncrossed to crossed

posture and vice versa movement condition. The experiment took

approximately 4 hours, split in two-hour sessions held on different days.

Practice trials were included on each day before the experiment started until

the participant had understood, and felt confident with, the task. In total we
acquired 14.776 trials.

*1 Stimulus Task:* The procedure was identical to the 2 Stimulus task except that
participants only received one stimulus at either hand and than indicated the
perceived location with the respective index finger. Participants performed 300
trials in each posture combination split in blocks of 50 trials. In 99,5% of the
trials participants were using the correct arm when localizing the stimulus.

**Analysis**

*Data preprocessing.* Start and end of the movement were determined based on
a velocity threshold of 5 cm/s. We interpolated missing motion tracking data,
for instance due to obstruction when the hands passed each other or due to
rotation of the hands, using splines, with the restriction that movement onset
and offset could be determined. Trials were discarded when (1) missing marker
data could not be adequadetely interpolated (1 Stimulus Task: 13%; 2 Stimulus
Task: 11.9%); (2) no stimulus localization response, indicated by finger lifting,
could be detected (1 Stimulus Task: 2.8%; 2 Stimulus Task: 5.4%); (3) and
when participants did not perform smooth, continuous, and synchronous
movements with movement duration less than 200 ms or more 1000 ms (1
Stimulus Task: 3.7%; 2 Stimulus Task: 0.2%). In total, 19% (1 Stimulus Task)
and 17.5% (2 Stimulus Task) of trials were removed.

*Analysis of Temporal Order Judgements (TOJ).* We considered the TOJ to be
correct when the hand used for the localization response had indeed been
stimulated first.

*Localization error.* We calculated the localization error, that is, the difference
between the true location of the index finger at the time of stimulation and the
reported location, that is, index finger pointing location just before finger lifting.
Errors are reported relative to the direction of the movement as a straight line
between start and end position of the hand, with positive values indicating
errors in movement direction towards the end position of the hand. The
localization error varies systematically with the stimulus time relative to

movement onset (Dassonville, 1995; Maij et al., 2013, 2017; Maij, Grave, et al.,

2011; Watanabe et al., 2009). The localization errors were converted to an

estimated localization curve by averaging errors using a moving Gaussian

window of 75 ms across a time window of -200 to 600 ms (step size 1ms) with

respect to movement onset. For each participant and posture condition, we

calculated localization curves for correct TOJ trials relative to onset of the first

stimulus, for correct TOJ trials relative to onset of the second stimulus, and for

incorrect TOJ trials relative to onset of the first stimulus.

*Comparison of localization errors in correct and incorrect TOJ trials.* To

determine whether participants localized the stimulus relative to hand position

at the first or the second stimulus timepoint, we calculated – separately for each

participant and posture condition –  the temporal shift that would produce the

smallest deviations around a single, common localization curve of the

compared conditions 1) between the incorrect TOJ localization curve and the

correct TOJ localization curve relative to the first stimulus time point and 2)

between the incorrect TOJ localization curve and the correct TOJ localization

curve relative to the second stimulus time point. Specifically, we shifted the data

points of the incorrect TOJ localization curves in time from 300 – 300 ms in

steps of 1 ms and calculated the squared localization error differences with an

localization curve calculated from data points of the two compared conditions

using a common moving Gaussian average for each time shift. We refer to the

time value that minimized the summed squared error differences with this

overall construction curve as time shift (Maij et al., 2009, 2017).

We assessed the time shift for both stimulus times and, accordingly, obtained

two shift values per participant and posture condition. In some instances, we

were unable to construct a time shift due to a low number of incorrect TOJ trials

for that specific condition (18 cases out of 4 postures x 2 stimulus times x 12

participants = 96); these data points were treated as missing data in the linear

mixed model analysis.

*Statistical analysis.* We assessed statistical significance of the reported

results using (Generalized) Linear Mixed Models ([G]LMM) (Bolker et al., 2009,

2009) as implemented in R version 3.6.1 (R Core Team, 2014) using packages

lme4, version 1.1-21 (Bates et al., 2015), and afex version 0.26-0 (Singmann,

2015). We estimated intercept parameters of Bayesian mixed factorial models

equivalent to LMM using packages brms, version 2.12.0 (Bürkner, 2017, 2018),

and loo, version 2.2.0 (Vehtari, Gabry, et al., 2017; Vehtari, Gelman, et al.,

2017).

GLMM are adequate for analysis of binary variables such as correct vs.

incorrect responses in our TOJ task (Jaeger, 2008). Furthermore, (G)LMM and

are robust against missing data and account for differences in trial numbers

across conditions, as present in our data. All reported statistics were computed

using type 3 sums of squares, as implemented in afex. For the random structure

of LMM and GLMM for TOJ analysis, we included only random intercepts,

because models did not reliably converge when random slopes were included.

Models that tested time shifts against zero used only the data corresponding to

one particular reference time point of correct trials (localization error relative to

time 1 or time 2); given that posture did not significantly modulate time shift, the

respective models excluded this factor. Accordingly, we compared a model

without intercept [shift ~ 0 + (1 l participant] against a model with intercept [shift

~ 1 + (1 l participant], effectively testing whether a non-zero intercept

significantly improved the time shift fit.

The brms R package uses STAN as backend. We ran LMM to estimate

the 95% interval of the intercepts in the different time shift models. We

compared Bayesian models using the loo_compare() function of the loo R

package. This function uses leave-one-out cross-validation to compare models

by assessing the models' predictive density when each data point is omitted

from fitting (Vehtari, Gelman, et al., 2017).

**Experiment 2**

*Apparatus, task and procedure*

Kinematic data of the fingers was recorded using an optical motion capture

system (Visualeyez II VZ4000v, Phoenix Technologies Inc, Vancouver, BC,

Canada) at 250Hz sampling frequency with markers placed on the nail of the

two index fingers. The experiment was controlled with Matlab (The MathWorks

Version R2015a; Natick, MA, USA) using the Psychophysics Toolbox

(Brainard, 1997). Stimulus presentation was controlled via custom-made

hardware and triggered through a digital acquisition card (PCI-6509, National

Instruments, Austin, USA).

The procedure was largely similar to Experiment 1, except that 1) participants

made only reaches from an uncrossed to an uncrossed posture or from a

crossed to a crossed posture and 2) tactile stimuli were separated by SOAs of

60, 85, 110, or 135 ms. Posture was varied in blocks of 64 trials in a pseudo-

randomized order. Within each block, SOA and which hand was stimulated first

were pseudo-randomized. Participants performed 28 blocks (14 of each

posture combination) for a total of 1792 trials. To compensate for marker

obstruction and failure to reliably detect finger lifting, 2 participants performed

10 additional blocks (5 of each posture combination). As we stopped any form

of data acquisition in our lab for an indefinite time period beginning of March

2020, 2 participants performed only 19 and 23 blocks, respectively. The

experiment took about 5-6 hours to complete, split in two-hour sessions held

on different days. Practice trials were included prior to each experimental

session.

***Analysis***

*Data preprocessing.* We used custom-written Matlab scripts for processing of

kinematic data. We first interpolated missing data points and resampled the

data to 1000 Hz using splines, and low-pass filtered the data using a second-

order butterworth filter with a cut-off frequency of 6 Hz. We determined

movement onset/ offset of each hand as the time of the sample in which the

resultant velocity of the respective finger marker exceeded/ dropped below 5

cm/s. We excluded trials when missing marker data could not be adequadetely

interpolated (8%), when no stimulus localization response, indicated by finger

lifting, could be detected (4.2%), and when participants did not perform smooth,

continuous, and synchronous movements (6.8%). In total we removed 18.2%
of the trials.

*Analysis of Temporal Order Judgements (TOJ) and localization error.* TOJ and
localization errors were determined as in Experiment 1. Because localization
error curves were similar regardless of start and end posture in Experiment 1,
we collapsed across postures in Experiment 2 to calculate the localization error
curves; we calculated individual curves for each participant and SOA (60, 85,
110, and 135 ms). Localization error curves calculated separately for
Experiment 2's two posture conditions yielded qualitatively similar results (see
Supplementary Information).

*Comparison of localization errors in correct and incorrect TOJ trials.* Time shift
947  values were calculated as in Experiment 1 (separately for each participant and
948  SOA)  using a shifting window of -300 to 300 ms (step size 1ms).

*Statistical analysis.* We assessed TOJ performance using a generalized mixed
model (GLMM) with factors Posture (uncrossed-uncrossed, crossed-crossed),
and SOA (60, 85, 110, 135 ms). The analysis approach for the dependence of
time shifts on SOAs followed a similar logic as Experiment 1. We first assessed
the significance of main effects and interaction of the experimental design with
afex. We then assessed whether time shifts were 0 relative to time 1 and time
2. To this end, we compared models with a random participant factor but no
fixed factors and intercept [shift ~ 0 + (1 | participant], with a common intercept
for all SOAs [shift ~1 + (1 | participant], and with individual intercepts per SOA
[shift ~ SOA + (1 | participant] separately for shift values relative to time 1 and
time 2, respectively.

## Acknowledgements

We thank Maie Stein, Franziska Rudzik, and Nina Held for help with data
acquisition, Volker Franz, Rainer Schäfer, and Renate Kiesewalter for their help
with the Optotrak and ODAU setup, Conrad Alting for help with programming of
Experiment 2, and Mathieu G.M. Koppen for help with LMM analysis. FM was

supported by the Netherlands Organization for Scientific Research (NWO Veni

Grant 451-12-009). TH was supported by an Emmy Noether grant of the

German Research Foundation (DFG) (He 6368/1-1). WPM received support

from the European Research Council (EU-ERC-283567) and the Netherlands

Organization for Scientific Research (NWO-VICI: 453-11-001). Data and code

for the present paper are available at the Open Science Framework website,

https://osf.io/ybxn5/.

## References

Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R., & Kievit, R. A. (2019). Raincloud plots: A multi-platform tool for robust data visualization. *Wellcome Open Research, 4*, 63. https://doi.org/10.12688/wellcomeopenres.15191.1

Alsmith, A. J. T., Ferrè, E. R., & Longo, M. R. (2017). Dissociating contributions of head and torso to spatial reference frames: The misalignment paradigm. *Consciousness and Cognition, 53*, 105–114. https://doi.org/10.1016/j.concog.2017.06.005

Angelaki, D. E., Gu, Y., & DeAngelis, G. C. (2009). Multisensory integration: Psychophysics, neurophysiology, and computation. *Current Opinion in Neurobiology, 19*(4), 452–458. https://doi.org/10.1016/j.conb.2009.06.008

Atsma, J., Maij, F., Koppen, M., Irwin, D. E., & Medendorp, W. P. (2016). Causal Inference for Spatial Constancy across Saccades. *PLOS Computational Biology, 12*(3), e1004766. https://doi.org/10.1371/journal.pcbi.1004766

Azañón, E., Camacho, K., & Soto-Faraco, S. (2010). Tactile remapping beyond space. *European Journal of Neuroscience, 31*(10), 1858–1867. https://doi.org/10.1111/j.1460-9568.2010.07233.x

Azañón, E., Mihaljevic, K., & Longo, M. R. (2016). A three-dimensional spatial characterization of the crossed-hands deficit. *Cognition, 157*, 289–295. https://doi.org/10.1016/j.cognition.2016.09.007

Azañón, E., & Soto-Faraco, S. (2008). Changing reference frames during the encoding of tactile events. *Current Biology, 18*(14), 1044–1049. https://doi.org/10.1016/j.cub.2008.06.045

Azañón, E., Stenner, M.-P., Cardini, F., & Haggard, P. (2015). Dynamic Tuning of Tactile Localization to Body Posture. *Current Biology, 25*(4), 512–517. https://doi.org/10.1016/j.cub.2014.12.038

Badde, S., & Heed, T. (2016). Towards explaining spatial touch perception: Weighted integration of multiple location codes. *Cognitive Neuropsychology, 33*(1–2), 1–22. https://doi.org/10.1080/02643294.2016.1168791

Badde, S., Heed, T., & Röder, B. (2014). Processing load impairs coordinate integration for the localization of touch. *Attention, Perception, & Psychophysics, 76*(4), 1136–1150. https://doi.org/10.3758/s13414-013-0590-2

Badde, S., Heed, T., & Röder, B. (2015). Integration of anatomical and external response mappings explains crossing effects in tactile localization: A probabilistic modeling approach. *Psychonomic Bulletin & Review, 23*(2), 387–404. https://doi.org/10.3758/s13423-015-0918-0

Badde, S., Röder, B., & Heed, T. (2019). Feeling a Touch to the Hand on the Foot. *Current Biology, 29*(9), 1491-1497.e4. https://doi.org/10.1016/j.cub.2019.02.060

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects Models Using lme4. *Journal of Statistical Software, 67*(1). https://doi.org/10.18637/jss.v067.i01

Bernier, P.-M., & Grafton, S. T. (2010). Human posterior parietal cortex flexibly determines feference frames for reaching based on sensory context. *Neuron, 68*(4), 776–788. https://doi.org/10.1016/j.neuron.2010.11.002

Binda, P., Morrone, M. C., & Burr, D. C. (2010). Temporal auditory capture does not affect the time course of saccadic mislocalization of visual stimuli. *Journal of Vision, 10*(2), 7–7. https://doi.org/10.1167/10.2.7

Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R., Stevens, M. H. H., & White, J.-S. S. (2009). Generalized linear mixed models: A practical guide for ecology and evolution. *Trends in Ecology & Evolution, 24*(3), 127–135. https://doi.org/10.1016/j.tree.2008.10.008

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision, 10*(4), 433–436. https://doi.org/10.1163/156856897X00357

Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using Stan. *Journal of Statistical Software, 80*(1), 1–28. https://doi.org/10.18637/jss.v080.i01

Bürkner, P.-C. (2018). Advanced Bayesian Multilevel Modeling with the R Package brms. *The R Journal, 10*(1), 395–411.

Cadieux, M. L., Barnett-Cowan, M., & Shore, D. I. (2010). Crossing the hands is more confusing for females than males. *Experimental Brain Research, 204*(3), 431–446. https://doi.org/10.1007/s00221-010-2268-5

Cadieux, M. L., & Shore, D. I. (2013). Response Demands and Blindfolding in the Crossed-Hands Deficit: An Exploration of Reference Frame Conflict. *Multisensory Research, 26*(5), 465–482. https://doi.org/10.1163/22134808-00002423

Dassonville, P. (1995). Haptic localization and the internal representation of the hand in space. *Experimental Brain Research, 106*(3), 434–448.

Ernst, M. O., & Banks, M. S. (2002). Humans integrate visual and haptic information in a statistically optimal fashion. *Nature, 415*(6870), 429–433. https://doi.org/10.1038/415429a

Ernst, M. O., & Di Luca, M. (2011). Multisensory perception: From integration to remapping. In *Sensory cue integration* (pp. 224–250). Oxford University Press, Oxford.

Fiehler, K., Schütz, I., & Henriques, D. Y. P. (2011). Gaze-centered spatial updating of reach targets across different memory delays. *Vision Research, 51*(8), 890–897. https://doi.org/10.1016/j.visres.2010.12.015

1057    Gallace, A., & Spence, C. (2005). Visual capture of apparent limb position
1058            influences tactile temporal order judgments. *Neuroscience Letters*,
1059            *379*(1), 63–68. https://doi.org/10.1016/j.neulet.2004.12.052

1060    Harrar, V., & Harris, L. R. (2010). Touch used to guide action is partially
1061            coded in a visual reference frame. *Experimental Brain Research*,
1062            *203*(3), 615–620. https://doi.org/10.1007/s00221-010-2252-0

1063    Heed, T., & Azañón, E. (2014). Using time to investigate space: A review of
1064            tactile temporal order judgments as a window onto spatial processing
1065            in touch. *Frontiers in Psychology*, *5*, 76.
1066            https://doi.org/10.3389/fpsyg.2014.00076

1067    Heed, T., Backhaus, J., Röder, B., & Badde, S. (2016). Disentangling the
1068            External Reference Frames Relevant to Tactile Localization. *PLOS*
1069            *ONE*, *11*(7), e0158829. https://doi.org/10.1371/journal.pone.0158829

1070    Heed, T., Buchholz, V. N., Engel, A. K., & Röder, B. (2015). Tactile
1071            remapping: From coordinate transformation to integration in
1072            sensorimotor processing. *Trends in Cognitive Sciences*, *19*(5), 251–
1073            258. https://doi.org/10.1016/j.tics.2015.03.001

1074    Heed, T., Möller, J., & Röder, B. (2015). Movement Induces the Use of
1075            External Spatial Coordinates for Tactile Localization in Congenitally
1076            Blind Humans. *Multisensory Research*, *28*(1–2), 173–194.
1077            https://doi.org/10.1163/22134808-00002485

1078    Hermosillo, R., Ritterband-Rosenbaum, A., & van Donkelaar, P. (2011).
1079            Predicting Future Sensorimotor States Influences Current Temporal
1080            Decision Making. *The Journal of Neuroscience*, *31*(27), 10019–10022.
1081            https://doi.org/10.1523/JNEUROSCI.0037-11.2011

1082    Jaeger, T. F. (2008). Categorical data analysis: Away from ANOVAs
1083            (transformation or not) and towards logit mixed models. *Journal of*
1084            *Memory and Language*, *59*(4), 434–446.
1085            https://doi.org/10.1016/j.jml.2007.11.007

1086    Kayser, C., & Shams, L. (2015). Multisensory Causal Inference in the Brain.
1087            *PLOS Biology*, *13*(2), e1002075.
1088            https://doi.org/10.1371/journal.pbio.1002075

1089    Kim, D.-H., & Cruse, H. (2001). Two kinds of body representation are used to
1090            control hand movements following tactile stimulation. *Experimental*
1091            *Brain Research*, *139*(1), 76–91.
1092            https://doi.org/10.1007/s002210100741

1093    Kitazawa, S. (2002). Where conscious sensation takes place. *Consciousness*
1094            *and Cognition*, *11*(3), 475–477. https://doi.org/10.1016/S1053-
1095            8100(02)00031-4

1096    Kitazawa, S., Moizumi, S., Okuzumi, A., Saito, F., Shibuya, S., Takahashi, T.,
1097            Wada, M., & Yamamoto, S. (2008). Reversal of subjective temporal
1098            order due to sensory and motor integrations. In P. Haggard, Y.

1099   Rossetti, & M. Kawato (Eds.), *Sensorimotor foundations of higher*
1100   *cognition* (pp. 73–97). Oxford University Press.

1101 Lohmann, J., & Butz, M. V. (2017). Lost in space: Multisensory conflict yields
1102   adaptation in spatial representations across frames of reference.
1103   *Cognitive Processing*, 1–18. https://doi.org/10.1007/s10339-017-0798-
1104   5

1105 Maij, F., Brenner, E., & Smeets, J. B. J. (2009). Temporal Information Can
1106   Influence Spatial Localization. *Journal of Neurophysiology*, *102*(1),
1107   490–495. https://doi.org/10.1152/jn.91253.2008

1108 Maij, F., Brenner, E., & Smeets, J. B. J. (2011). Temporal Uncertainty
1109   Separates Flashes from Their Background during Saccades. *Journal of*
1110   *Neuroscience*, *31*(10), 3708–3711.
1111   https://doi.org/10.1523/JNEUROSCI.5097-10.2011

1112 Maij, F., Grave, D. D. J., Brenner, E., & Smeets, J. B. J. (2011). Misjudging
1113   where you felt a light switch in a dark room. *Experimental Brain*
1114   *Research*, *213*(2–3), 223–227. https://doi.org/10.1007/s00221-011-
1115   2680-5

1116 Maij, F., Wing, A. M., & Medendorp, W. P. (2013). Spatiotemporal integration
1117   for tactile localization during arm movements: A probabilistic approach.
1118   *Journal of Neurophysiology*, *110*(11), 2661–2669.
1119   https://doi.org/10.1152/jn.00971.2012

1120 Maij, F., Wing, A. M., & Medendorp, W. P. (2017). Afferent motor feedback
1121   determines the perceived location of tactile stimuli in the external space
1122   presented to the moving arm. *Journal of Neurophysiology*, *118*(1),
1123   187–193. https://doi.org/10.1152/jn.00286.2016

1124 Matin, L., & Pearce, D. G. (1965). Visual Perception of Direction for Stimuli
1125   Flashed During Voluntary Saccadic Eye Movements. *Science*,
1126   *148*(3676), 1485–1488. https://doi.org/10.1126/science.148.3676.1485

1127 Mueller, S., & Fiehler, K. (2014a). Gaze-dependent spatial updating of tactile
1128   targets in a localization task. *Frontiers in Psychology*, *5*, 66.
1129   https://doi.org/10.3389/fpsyg.2014.00066

1130 Mueller, S., & Fiehler, K. (2014b). Effector movement triggers gaze-
1131   dependent spatial coding of tactile and proprioceptive-tactile reach
1132   targets. *Neuropsychologia*, *62*, 184–193.
1133   https://doi.org/10.1016/j.neuropsychologia.2014.07.025

1134 Penfield, W., & Boldrey, E. (1937). Somatic motor and sensory representation
1135   in the cerebral cortex of man as studied by electrical stimulation. *Brain*,
1136   *60*(4), 389–443. https://doi.org/10.1093/brain/60.4.389

1137 R Core Team. (2014). *R: A Language and Environment for Statistical*
1138   *Computing*. R Foundation for Statistical Computing. http://www.R-
1139   project.org/

Roberts, R. D., Wing, A. M., Durkin, J., & Humphreys, G. W. (2003). Effects of
posture on tactile temporal order judgements. *Eurohaptics Conference,*
*Dublin, Ireland*, 300–304.

Röder, B., Kusmierek, A., Spence, C., & Schicke, T. (2007). Developmental
vision determines the reference frame for the multisensory control of
action. *Proceedings of the National Academy of Sciences*, *104*(11),
4753–4758. https://doi.org/10.1073/pnas.0607158104

Röder, B., Rösler, F., & Spence, C. (2004). Early vision impairs tactile
perception in the blind. *Current Biology*, *14*(2), 121–124.
https://doi.org/10.1016/j.cub.2003.12.054

Roux, F., Djidjeli, I., & Durand, J. (2018). Functional architecture of the
somatosensory homunculus detected by electrostimulation. *The*
*Journal of Physiology*, *596*(5), 941–956.
https://doi.org/10.1113/JP275243

Schlag, J., & Schlag-Rey, M. (2002). Through the eye, slowly; Delays and
localization errors in the visual system. *Nature Reviews Neuroscience*,
*3*(3), 191–191. https://doi.org/10.1038/nrn750

Schubert, J. T. W., Badde, S., Röder, B., & Heed, T. (2017). Task demands
affect spatial reference frame weighting during tactile localization in
sighted and congenitally blind adults. *PLOS ONE*, *12*(12), e0189067.
https://doi.org/10.1371/journal.pone.0189067

Schütz, I., Henriques, D. Y. P., & Fiehler, K. (2013). Gaze-centered spatial
updating in delayed reaching even in the presence of landmarks.
*Vision Research*, *87*, 46–52.
https://doi.org/10.1016/j.visres.2013.06.001

Shore, D. I., Gray, K., Spry, E., Spence, C., & others. (2005). Spatial
modulation of tactile temporal-order judgments. *Perception*, *34*(10),
1251–1262.

Shore, D. I., Spry, E., & Spence, C. (2002). Confusing the mind by crossing
the hands. *Cognitive Brain Research*, *14*(1), 153–163.
https://doi.org/10.1016/S0926-6410(02)00070-8

Simon, J. R., Hinrichs, J. V., & Craft, J. L. (1970). Auditory SR compatibility:
Reaction time as a function of ear-hand correspondence and ear-
response-location correspondence. *Journal of Experimental*
*Psychology*, *86*(1), 97–102.

Singmann, H. (2015). *afex: Analysis of Factorial Experiments*. http://R-
Forge.R-project.org/projects/afe/

Thompson, A. A., Byrne, P. A., & Henriques, D. Y. P. (2014). Visual Targets
Aren't Irreversibly Converted to Motor Coordinates: Eye-Centered
Updating of Visuospatial Memory in Online Reach Control. *PLoS ONE*,
*9*(3), e92455. https://doi.org/10.1371/journal.pone.0092455

Tramper, J. J., & Medendorp, W. P. (2015). Parallel updating and weighting of
multiple spatial maps for visual stability during whole body motion.

*Journal of Neurophysiology, 114*(6), 3211–3219.
1184                  https://doi.org/10.1152/jn.00576.2015

1185   van Beers, R. J., Wolpert, D. M., & Haggard, P. (2002). When Feeling Is More
1186                  Important Than Seeing in Sensorimotor Adaptation. *Current Biology,*
1187                  *12*(10), 834–837. https://doi.org/10.1016/S0960-9822(02)00836-9

1188   Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., & Gelman, A. (2017). *loo:*
1189                  *Efficient Leave-One-Out Cross-Validation and WAIC for Bayesian*
1190                  *Models.* https://mc-stan.org/loo/

1191   Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model
1192                  evaluation using leave-one-out cross-validation and WAIC. *Statistics*
1193                  *and Computing, 27*(5), 1413–1432. https://doi.org/10.1007/s11222-
1194                  016-9696-4

1195   Watanabe, J., Nakatani, M., Ando, H., & Tachi, S. (2009). Haptic localizations
1196                  for onset and offset of vibro-tactile stimuli are dissociated. *Experimental*
1197                  *Brain Research, 193*(3), 483–489. https://doi.org/10.1007/s00221-009-
1198                  1711-y

1199   Yamamoto, S., & Kitazawa, S. (2001). Reversal of subjective temporal order
1200                  due to arm crossing. *Nat. Neurosci., 4*(7), 759–765.
1201                  https://doi.org/10.1038/89559

1202