External location of touch is constructed post-hoc based on limb choice

Femke Maij[#3], Christian Seegelke[#1,2], W. Pieter Medendorp[3], Tobias Heed[1,2]

[1]Faculty of Psychology and Sports Science, Bielefeld University, Bielefeld, Germany

[2]Center of Excellence Cognitive Interaction Technology (CITEC), Bielefeld University, Bielefeld, Germany

[3]Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, Nijmegen, The Netherlands

[#]These authors contributed equally to this work

Running head: Post-hoc construction of tactile external location

Correspondence: Tobias Heed, tobias.heed@uni-bielefeld.de

Abstract (150 words)

When humans indicate on which hand a tactile stimulus occurred, they often
err when their hands are crossed. This finding seemingly supports the view that
the automatically determined touch location in external space affects limb
assignment: the crossed right hand is localized in left space, and this conflict
presumably provokes hand assignment errors. Here, participants judged on which
hand the first of two stimuli, presented during a bimanual movement, had occurred,
and then indicated its external location by a reach-to-point movement. When
participants incorrectly chose the hand stimulated second, they pointed to where
that hand had been at the correct, first time point, though no stimulus had occurred
at that location. This behavior suggests that stimulus localization depended on
hand assignment, not vice versa. It is, thus, incompatible with the notion of
automatic computation of external stimulus location upon occurrence. Instead,
humans construct external touch location post-hoc and on demand.

Introduction

Spatial perception and actions rely on multiple spatial codes, often

associated with different reference frames. For instance, the accuracy of pointing

or reaching with an arm or finger to a visual target depends not only on the position

of target relative to gaze (Fiehler et al., 2011; Thompson et al., 2014), but also on

salient world-centered landmarks (Schütz et al., 2013). Similarly, judgment of

visual location during whole-body movement is influenced by a target's position

relative to gaze, as well as by the location of the target relative to the body

(Tramper & Medendorp, 2015).

In touch, too, space is coded in several reference frames. Touch activates

specialized sensory receptors embedded in the skin, and the arrangement of the

peripheral sensors is reflected in the homuncular organization of primary

somatosensory cortex (Penfield & Boldrey, 1937; Roux et al., 2018), referred to as

a skin-based or somatotopic layout. However, because our body can take various

postures, the stimulus location in space – often referred to as its external location

– must be derived by combining skin location and body posture, a process termed

tactile remapping (Heed, Buchholz, et al., 2015). Indeed, there is evidence that

external tactile locations can be coded in a gaze-centered reference frame (Harrar

& Harris, 2010; Mueller & Fiehler, 2014a, 2014b), but also relative to anchors such

as the head, torso, and hand (Alsmith et al., 2017; Heed et al., 2016).

It is less clear, however, according to which principles these different spatial

codes are employed. Both bottom-up features such as the availability of sensory

information (Bernier & Grafton, 2010) and the spatial reliability of a sensory

channel (Ernst & Banks, 2002; van Beers et al., 2002), as well as top-down

information such as task-constraints (Badde et al., 2015; Schubert et al., 2017),

action context (Mueller & Fiehler, 2014b), and cognitive load (Badde et al., 2014)

can affect the relative contributions of different reference frames, presumably in a

weighted manner (Angelaki et al., 2009; Atsma et al., 2016; Badde & Heed, 2016;

Ernst & Di Luca, 2011; Kayser & Shams, 2015; Lohmann & Butz, 2017; Tramper

& Medendorp, 2015). Yet, whereas there is widespread consensus that each

spatial code can have more or less influence depending on the specific situation,

it is currently not known whether all putative codes are always constructed, or

whether they are only computed based on demand.

For touch, it has been suggested that the construction of spatial location is

an automatic process, implying that any tactile input is remapped into an external

code, irrespective of its relevance (Heed & Azañón, 2014; Röder et al., 2004). The

most common experimental manipulation underlying this claim is limb crossing.

Crossing, say, a right arm over to the left side of space leads to different skin-

based (here: right body side) and external (here: left side of space) spatial codes

of a tactile stimulus delivered to the right hand. A task-irrelevant tactile stimulus

delivered to a crossed right hand accelerates visual discrimination in the right

visual field if it precedes the visual target stimulus by 60 ms, but on the left side if

it leads by 180 ms or more (Azañón et al., 2010; Azañón & Soto-Faraco, 2008).

Thus, responses to visual targets were faster after anatomically congruent tactile

cues (e.g., tactile stimulus on [crossed] right hand, visual target in right hemifield)

at short cue-stimulus intervals, but after externally congruent tactile cues (e.g.,

tactile stimulus on the left hand crossed over to the right side, visual target in right

field) at long cue-stimulus intervals. Such effects are usually interpreted as

evidence that tactile remapping – the precise computation of the external tactile

stimulus location – is automatic and forms the basis for the performance

enhancement at this external location.

The same conclusion has also been drawn from results obtained with the

tactile temporal order judgment (TOJ) task; in this task, participants report which

of two successive tactile stimuli, each presented to a different body part – typically

the two hands – occurred first (Heed & Azañón, 2014; Shore et al., 2002;

Yamamoto & Kitazawa, 2001). When the time interval between the two stimuli is

short, participants sometimes choose the wrong stimulus. Notably, stimulus

confusion is much more prominent when the arms are in a crossed than uncrossed

posture. This is surprising because the TOJ task asks about the identity of the

touched limb, and, in theory, it would be irrelevant to this question where the hand

was in space. That limb crossing, nevertheless, affects TOJ implies that posture

cannot be strategically ignored, but is automatically incorporated into the hand

assignment.

Several explanations have been put forward to account for crossing effects

in tactile localization. First, it has been suggested that touch location, once it is

remapped, is retained only in an external spatial code, and the original skin location

is discarded in the process. To report which body part has been touched, the brain

must then reversely determine which limb was located at the computed external

location at the time of the touch (Kitazawa, 2002; Kitazawa et al., 2008; Yamamoto

& Kitazawa, 2001). We refer to this suggestion as the *space-to-limb reconstruction*

hypothesis. When applied to errors in the TOJ task, this hypothesis implies that

participants correctly remap the two tactile stimuli into external space, but then

reconstruct erroneously which hand was at the first spatial location.

A second explanation assumes that TOJ errors reflect the conflict between

different codes used for stimulus location. When the limbs are crossed, skin-based

and external spatial codes point to different sides of space, and this conflict must

be resolved, a process that takes time and is error-prone (Röder, Kusmierek,

Spence, & Schicke, 2007; Simon, Hinrichs, & Craft, 1970). In this view, the TOJ

crossing effect is a marker for the presence of conflict and, thus, for the fact that

remapping into an external spatial code has taken place. Notably, the interpretation

that the TOJ crossing effect derives from a remapped stimulus location is indirect

because participants only report a binary decision about which hand was

stimulated, not the spatial location of the perceived stimulus. Increasing the

distance between the uncrossed hands can slightly reduce errors in TOJ (Gallace

& Spence, 2005; Roberts et al., 2003; Shore et al., 2005), and the TOJ crossing

effect is smaller when the hands' positions additionally differ in height or depth

(Azañón et al., 2016). These graded modulations of the TOJ have led to the claim

that the TOJ paradigm is an implicit index of tactile remapping (Azañón et al., 2015;

Badde & Heed, 2016; Heed & Azañón, 2014). We refer to this suggestion as the

stimulus switch hypothesis. It implies that participants have correctly remapped the

two stimuli into space, but have incorrectly resolved the conflict between the

different spatial codes of the first stimulus, consequently assigning the incorrect

stimulus to the first time point; as a consequence, participants incorrectly report

the hand that received the second stimulus.

110   Importantly, both hypotheses outlined above assume that touch is

automatically and accurately remapped to its external location. However, recent

experiments have cast doubt on whether this is actually the case. For instance, if

a tactile stimulus is presented during an arm movement, and participants indicate

the stimulus's location by pointing to its external location after the movement, they

make systematic localization errors (Dassonville, 1995; Maij et al., 2011, 2013,

2017; Watanabe et al., 2009). Importantly, because these errors differ for fast and

slow movements, it has been suggested that participants do not compute the

precise spatial location of a stimulus when it occurs, but instead infer spatial

location post-hoc by estimating hand location at the perceived time of the tactile

stimulus (Maij et al., 2017). We refer to this suggestion as the *time reconstruction*

hypothesis. Accordingly, errors in the TOJ task would occur because participants

first choose the incorrect hand, and then derive stimulus location based on that

hand's location at the time of the first stimulus. Note, that here participants merge

the correct, first stimulus's time with the incorrect, second stimulus's hand.

Importantly, the claim of the *time reconstruction hypothesis* that tactile location is

not determined by integrating posture and skin location at the time of stimulus

occurrence – in the case of TOJ errors, this is the time point of the second stimulus

Figure 1. *Hand assignment and tactile localization in a TOJ task with bimanual movements. Grey*
trace, left arm trajectory; yellow trace: right arm trajectory; red trace: reach-to-point movement to
external location of the tactile stimulus. A. Correct TOJ trial: the stimulus is assigned to the correct
hand, which points to the correct location. B-D Three hypotheses for tactile location in incorrect
TOJ trials. B. Space-to-limb reconstruction hypothesis: participants point with the incorrect hand at
the external location of the first stimulus. C. Stimulus switch hypothesis: participants point with the
incorrect hand at the external location of the second stimulus. D. Time reconstruction hypothesis:
participants point with the incorrect hand at the location at which that hand was at the time of the
first stimulus.

– is at odds with the idea that tactile judgments are based on precise spatial
remapping.

Here, we assessed hand assignment and spatial localization of tactile
stimuli presented during movement. Our objective was to test whether TOJ
responses mark the use of the stimulus's precise, external-spatial location
constructed in response to the stimulus, or whether instead participants estimate
the precise stimulus location post-hoc by integrating the hand movement trajectory
with stimulus time. In other words, we aimed to directly contrast the three

discussed hypotheses for tactile localization: the space-to-limb reconstruction
hypothesis, the stimulus switch hypothesis, and the time reconstruction
hypothesis.

We presented human participants with two tactile stimuli during a bimanual
movement and assessed which hand participants perceived to have been
stimulated first (TOJ hand assignment), as well as exactly where in space
participants had perceived it (tactile stimulus localization). The experimental logic,
and its relation to the three tested tactile localization hypotheses, are illustrated in
Fig. 1. Because tactile stimuli were presented shortly before, after, and during the
time of movement, their precise spatial location depended on their timing relative
to the movement. This allowed us to determine the precise tactile spatial location
participants had perceived when they had made a hand assignment error in the
TOJ task. Contrary to common opinion, TOJ errors were not associated with the
location of the second, incorrect stimulus. Instead, when participants chose the
incorrect hand, they reported its location at the time point at which the other, correct
stimulus had occurred. Thus, participants constructed stimulus location by
combining the position of the incorrectly chosen hand with the stimulus timing that
belonged to the other, non-chosen hand's stimulus, resulting in reported locations
at which no stimulus had ever occurred. This finding invalidates current
explanations of crossing effects as being based on the precise, remapped external
spatial location of the tactile stimulus.

Results

Experiment 1

Participants assumed a start posture with their hands resting on a table and
their arms stretched out in an uncrossed or crossed posture (see Fig. 2 for an
illustration of experimental conditions and trial timing). A tone then instructed a
movement of the two hands about 30 cm towards their body, bringing the arms

into either an uncrossed or crossed arm end posture (see Fig. 2A). Shortly

Figure 2. *Experimental conditions. A. The arms moved from an uncrossed (u) or crossed (c) start*
175
posture to either an uncrossed or crossed arm end posture. B. Example TOJ trial of a representative
subject, showing the bimanual movement (grey: left arm, yellow, right arm) and the reach-to-point
movement of the assigned hand for the tactile localization response.

before, during, or shortly after the movement, participants received two tactile

stimuli, one on each hand, with a stimulus onset asynchrony (SOA) of 110 ms. At

this SOA, participants often misreport which of the two stimuli occurred first, both

when the arms are still (Heed & Azañón, 2014; Shore et al., 2002; Yamamoto &

Kitazawa, 2001) and during movement (Heed, Möller, et al., 2015; Hermosillo et

al., 2011). After the bimanual movement, participants reported on which of the two

hands the first stimulus had occurred by reaching with this hand to the perceived

external location of the stimulus (see Fig. 2B). The response, thus, contained two

components: the hand to which the first stimulus was assigned, and explicit spatial

localization of this stimulus.

Hand assignment

190   We measured TOJ performance as the percentage of correct reports of

which hand had been stimulated first in the TOJ task, as indicated by the hand that

participants used for their localization response (see Fig. 3). Stimuli could occur

during all times (see Methods for details), so we binned the binary

Figure 3. *Proportion of correct hand assignment across movement conditions (uu, uc, cu, uu, with*

c=crossed, u=uncrossed posture) in the four phases of the bimanual movement (before movement,

during start posture, during end posture, after movement). For uc and cu conditions, stimuli were

classified according to whether the first stimulus occurred while the hands were still in the start

posture, or already in the end posture. For uu and cc conditions, trials were assigned as "during

start posture" if stimuli had been presented during the first temporal half of the movement, and as

"during end posture" otherwise. Error bars denote 2 s.e. from the mean; asymmetry is due to

nonlinear conversion from the GLMM's logit scale to percentage correct. Large symbols are group

means, small symbols are individual participants' performance.

(correct/incorrect) TOJ response data into four movement phases – stimulation

before movement onset, during first and second half of movement, and after

movement offset – to assess the modulation of TOJ performance by stimulus time

relative to the bimanual movement.

In accordance with previous findings, TOJ performance declined in the

crossed compared to the uncrossed posture (Heed & Azañón, 2014), and

depended on the posture at the time of stimulation (Heed, Möller, et al., 2015;

Hermosillo et al., 2011). For instance, for the uncrossed-uncrossed movement

condition (see Fig. 3, uu), the probability of correct response was high compared

to the crossed-crossed movement condition (see Fig. 3, cc) throughout all

movement phases. For the conditions with a postural change (see Fig. 3, uc & cu)

the probability of correct responses was modulated by the posture at the time of

stimulation. A generalized mixed model (GLMM) with factors Start Posture, End

Posture, and Movement Phase revealed significance for all main effects and

Figure 4. *Reach end points, as measures of localization responses, in the horizontal plane for the*

different movement conditions. Data are from a single participant with substantial end point

variability. Ellipses cover 95% of the variability. Ellipses cover virtually identical spaces when

participants give correct and incorrect responses. The space-to-limb reconstruction hypothesis

would predict that, during error trials, participants point with the incorrectly chosen hand to the

location of the correct stimulus; thus, if this hypothesis were correct, orange ellipses should overlay

with grey ellipses, and red ellipses should overlay with yellow ellipses.

interactions (see Supplementary Table 1). With movement phase, the effect of

Start Posture (see Fig. 3, dark vs. light colors) declined, whereas the effect of End

Posture (see Fig. 3, green vs. magenta colors) increased. For instance, for

movements from an uncrossed to a crossed posture, TOJ performance was better

during the first two movement phases, that is, when the hands were still uncrossed,

than during the last two movement phases, that is, when the hands were crossed.

In sum, TOJ performance in our first experiment reflected known

modulations of hand posture and movement timing. Participants made, on

average, more than 15% TOJ errors even with uncrossed hands. This high error

rate is due to the use of the short SOA of 110 ms (Heed, Möller, et al., 2015; Heed

& Azañón, 2014; Shore et al., 2002; Yamamoto & Kitazawa, 2001), and is an

intended outcome of the experimental paradigm, allowing, as a next step,

comparison of spatial localization responses for erroneous TOJ trials across all

postures.

Explicit stimulus localization in space

Localization responses in TOJ error trials allow differentiating between the

three hypotheses about how participants determine stimulus localization in tactile

decision paradigms (see Fig. 1). The *space-to limb reconstruction hypothesis*

posits that the spatial location of the first tactile stimulus is represented correctly,

and that a limb assignment entails computing which limb was at that spatial

location when the stimulus occurred. Thus, in our task, responses with the

incorrect hand would result from assigning the incorrect hand to the correct spatial

location of the first tactile stimulus (see Fig. 1B). Accordingly, the chosen, incorrect

hand should be directed to the location at which the stimulus of the other, correct

hand had occurred, and the reported stimulus location in TOJ error trials should

scatter around the movement trajectory of the correct hand. Contrary to this

prediction, participants pointed to locations scattered around the movement

trajectory of the chosen, incorrect hand (see also Fig. 2B), indicating that the

chosen stimulus had been perceived on the incorrect hand (see Fig. 4 for the

localization responses of the participant with the largest variability in localization

errors). Thus, localization behavior did not support the implication of the *space-to-*

limb reconstruction hypothesis that the correct external spatial location is simply

assigned to a wrong limb.

259   Having established that participants appear to have perceived the stimulus

at the reporting hand, two possibilities remain as to which stimulus location was

associated with erroneous responses (see Fig. 1). The *stimulus switch hypothesis*

posits that the two stimuli were localized correctly, and one is chosen for the

response. In TOJ error trials, participants would confuse the two stimuli and report

the second stimulus by pointing at its location with the respective, incorrect hand.

In this case, spatial localization errors in the reach-to-point task should scatter

around the position of the hand at the time point of the *second*, erroneously chosen

stimulus (referred to as time 2 from hereon). In contrast, the *time reconstruction*

hypothesis assumes that participants always use the correct, first time point (time

1 from hereon), and determine the chosen response limb's location at this time

point. This hypothesis predicts that, in TOJ error trials, participants point to the

location of the erroneously chosen, incorrect hand at the correct time, that is, time

1. Note, that no tactile stimulus occurred at this external spatial location, because

it combines the time of the first, correct stimulus with the movement trajectory of

the second, incorrect stimulus's hand.

Figure 5. *Localization errors systematically vary in dependence of the progression of the*
movement. A. Mean localization error (teal line) of all correct localization trials of a single participant.
Each dot represents the error of a single trial, that is the difference between actual hand position
at the time of tactile stimulation and indicated hand position at the end of the trial. Blue shading
represents the average movement time, with 0 = movement start. Note, that localization bias is
positive at the beginning of the movement, but negative towards the end of the movement. These
biases reflect mislocalization in the direction of movement at the beginning, and in the opposite
direction at the end of the movement. B. The error pattern in A was evident across all subjects for
all posture combinations. Traces reflect the mean, shaded areas around the traces reflect s.e.m.
The shaded regions in the background indicate movement times, which differed slightly between
conditions.

Assessment of whether participants pointed to the hand's position at the
first or second time point during TOJ error trials is based on comparison of the
localization error curve in correct and erroneous TOJ trials. When participants
retrospectively point to the spatial location of a tactile stimulus presented while the
target limb was moving, their location judgment errors are systematically biased
in the direction of the movement during the initial part of a movement, and in the
opposite direction during the final part of the movement (Dassonville, 1995; Maij
et al., 2013, 2017; Watanabe et al., 2009), resulting in systematic localization error
curves. Such movement time-related biases were evident also in the present data
(see Fig. 5). In the case of correct TOJ trials, we assume that participants aimed,
as instructed, at the location of the correct hand at time 1. Therefore, we derive the
localization error curve as the spatial difference of reaching endpoint and hand
location at time 1 (see Fig. 6, dark blue lines). However, we can also derive a

second localization error curve by computing localization error relative to time 2

Figure 6. *Localization curves, averaged across participants, for each of the four posture conditions.*
Curves of incorrect TOJ trials (red) show a similar pattern as the localization curves of the correct
TOJ trials at time 1 (dark blue), but not as the localization curves of the correct TOJ trials at time 2
(light blue). Traces reflect the mean, shaded areas around the traces reflect s.e.m. The shaded
regions in the background represent the average movement time.

(see Fig. 6, light blue lines). This hypothetical error curve is shifted to the left of
the first curve, because time 2 is later in time relative to movement start (and,
accordingly, closer in time to movement end), so that the movement time-related
localization errors should be shifted backwards in time by the 110 ms time
difference of the two stimuli. In other words, this second error curve reflects the
pattern we would expect if participants had aimed for hand location at time 2.

The two localization error curves computed on the basis of correct TOJ trials
serve as templates against which we compare the localization error curve of TOJ
error trials (see Fig. 6 and Methods). Localization errors of incorrect TOJ trials
overlapped with localization errors of correct TOJ trials at time 1, for each of the

four start and end posture combinations. To quantify this further, we computed by

how much reaching endpoints of TOJ error trials had to be shifted to best match

either template curve derived from correct TOJ trials. If, in incorrect TOJ trials,

participants aimed for the incorrect hand's position at time 1, then the temporal

shift should be zero relative to the localization error curve in correct TOJ trials

relative to time 1; furthermore, it should be about -110 ms (negative denoting a

shift towards left, see above) compared to the localization error curve of correct

TOJ trials relative to time 2. If, however, participants aimed for the incorrect hand's

position at time 2, the shift pattern should be exactly reverse, that is, zero

compared to the second template curve, and near +110 ms compared to the first

template curve.

Figure 7. *Time shift of stimulus localization error in TOJ error trials relative to time 1 (left panel)*

and time 2 (middle panel) for the four posture conditions. The temporal shift of the localization error

curve was significantly different from zero when calculated relative to time 2, but not relative to time

1; this result is consistent with the time reconstruction hypothesis, but not with the stimulus switch

hypothesis. This pattern was similar across all participants as demonstrated by the differences in

time shift between time 2 and time 1 (right panel). Data are visualized with raincloud plots (Allen et

al., 2019) displaying probability density estimates, condition averages (large symbols), and

individual participants (small symbols). Error bars denote 95% confidence intervals.

Fig. 7 displays the temporal shift between the localization error curve of

incorrect TOJ trials and the error curves at time 1 and time 2, as derived from

correct TOJ trials. A linear mixed model with factors Posture Condition and

Reference Time Point (localization errors for correct TOJ trials computed relative

to time 1 or time 2) and time shift of the error curve between correct and erroneous

TOJ trials as dependent variable revealed a significant effect of Reference Time

Point ($\chi^2(9,10) = 15.55$, $p < 0.001$), but no effect of Posture Condition ($\chi^2(7,10) =$

1.57, p = 0.67) or interaction ($\chi^2(7,10) = 4.53$, p = 0.21) between the two factors.

The time shift between localization errors for correct TOJ trials at time 1 and TOJ

error trials, averaged across all four posture conditions, was 6 (s.e., 3) ms; if this

value is not significantly different from 0, then an LMM of only this condition should

not improve by inclusion of an intercept, as the latter would model the deviation of

the average shift away from 0. Indeed, comparison of a model with and without

intercept did not provide statistical evidence to reject a zero time shift ($\chi^2(1) = 0.84$,

p = 0.36). In line with this result, a Bayesian analysis of a model with a random

participant factor and a population-level intercept estimated the population

intercept at 6 ms, and the 95% confidence range [-8.28; 19.27 ms] included 0.

Model comparison via leave-one-out cross-validation found a model without

population intercept to be more credible (difference of expected log predictive

density, ELPD, for second as compared to first model: -0.6, s.e. 0.9). In contrast,

the average time shift between localization errors for correct TOJ trials at time 2

and TOJ error trials was -105 (s.e., 7) ms, and a model without intercept fit this

condition siginificantly worse than a model that included the intercept ($\chi^2(1) =$

19.38, p < 0.001). The significant difference to time 2 suggests that participants

did not aim at the position of the second stimulus in TOJ error trials. While neither

the non-significant difference to time 1 in the LMM analysis, nor the Bayesian

parameter estimate including 0 statistically imply equality of the error curves in

correct and incorrect TOJ trials, these statistical results are consistent with the two

conditions being equal, and they suggest that, if a difference exists, it is small.

Furthermore, the time shift of -105 ms for time 2 closely matches the stimulus SOA

of 110 ms, further suggesting that, in error trials, participants did not aim for hand

location at the second, but rather at the first time point. Corroborating this

conclusion, the Bayesian 95% interval [-138; -76 ms] of the intercept estimate

includes -110 ms, and comparison of Bayesian models for time 2 with and without

intercept strongly favored the model including the intercept (difference in ELPD

from first to second model, -1.4, s.e. 1.9).

Experiment 2

The results of Experiment 1 suggest that when asked to localize the external

location of the first of to two tactile stimuli applied in succession to different hands,

participants chose which hand received the stimulus and then inferred the location

of the chosen limb at the time point of the first stimulus. Consequently, when

participants chose the incorrect limb, stimulus location was determined as the

location at which the incorrect hand was at the correct (first) time point. While these

results support the *time reconstruction hypothesis*, it should be realized that

Experiment 1 tested only a single SOA of 110 ms between the two tactile stimuli.

If our conclusions drawn from Experiment 1 are correct, then localization of stimuli

assigned to the incorrect hand should always depend on the first stimulus's time,

independent of SOA.

Experiment 2 tested this conjecture. Again, participants judged which hand had

received the first of two tactile stimuli during a bimanual movement and then

located the stimulus perceived to have occurred first. We presented tactile stimuli

with four different SOAs: 60, 85, 110, and 135 ms. As the estimated localization

error curves in Experiment 1 were similar for all combinations of the hands' start

and end posture, Experiment 2 involved only reaches from an uncrossed to an

uncrossed posture and from a crossed to a crossed posture. This strategy

minimized marker obstruction and homogenized movement time across conditions.

Experiment 2 was conducted in a different lab than Experiment 1, using different

equipment, re-written experimental code, different experimenters, and new

analysis scripts (see Material and methods for details). To further scrutinize the

reliability of our results, we increased our sample size and acquired a higher

number of trials.

Hand assigment

In accordance with Experiment 1 and previous findings (Heed, Möller, et al., 2015;

Hermosillo et al., 2011), TOJ performance in Experiment 2 was modulated by hand

posture and SOA. At all SOA, participants made large amounts of errors, ensuring

that a sufficient number of trials were available to analyze incorrect hand choice

trials. Detailed results are reported in the Supplementary Information.

Explicit stimulus localization in space

Complementing the findings from Experiment 1 and further corroborating the *time*

reconstruction hypothesis, localization errors of the incorrect TOJ trials largely

overlapped with the localization errors of correct TOJ trials at time 1 for each of the

four SOAs (see Fig. 8) and for each subject and posture condition (see

Supplementary Fig. 2-7).

Figure 8. *Localization curves, averaged across participants and posture, for each of the four SOAs.*

Curves of incorrect TOJ trials (red) show a similar pattern as the localization curves of the correct

TOJ trials at time 1 (dark blue), but not as the localization curves of the correct TOJ trials at time 2

(light blue). This pattern was highly similar across all subjects (see Supplementary Information) and

also when calculated separately for each posture condition (see Supplementary Information).

Traces reflect the mean, shaded areas around the traces reflect s.e.m. The shaded regions in the

background represent the average movement time.

Fig. 9 shows the temporal shift between the localization error curves of incorrect

TOJ trials and the error curves of correct TOJ trials relative to time 1 and time 2

four all four SOAs.

Figure 9. *Time shift of stimulus localization error in TOJ error trials relative to time 1 (left panel)*
and time 2 (middle panel) for the four SOAs. For all SOAs, the temporal shift relative to time 1 was
not significantly different from 0. In contrast, the time shift was significantly different from zero for
all SOAs when calculated relative to time 2, and it was numerically similar to the respective SOA.
These results are consistent with the time reconstruction hypothesis, but not with the stimulus
switch hypothesis. This pattern was similar across all participants as demonstrated by the
differences in time shift between time 2 and time 1 (right panel). Data are visualized with raincloud
plots (Allen et al., 2019) displaying probability density estimates, condition averages (large
symbols), and individual participants (small symbols). Error bars denote 95% confidence intervals.

A linear mixed model with factors SOA and Reference Time Point
(localization errors for correct TOJ trials computed relative to time 1 vs. time 2) and
time shift of the error curve between correct and erroneous TOJ trials as
dependent variable revealed a significant main effect of Reference Time Point
($\chi^2(9,10) = 21.33$, p < 0.001) and a significant Reference Time Point × SOA
interaction ($\chi^2(7,10) = 9.90$, p = 0.02). The time shift between localization errors
for correct TOJ trials at time 1 and TOJ error trials, averaged across all four SOA
conditions, was 8 ms. As in Experiment 1, a LMM with a common intercept for all
SOAs did not fit the data better than a model without an intercept ($\chi^2(1) = 1.24$, p
= 0.27). Similarly, allowing for individual intercepts per SOA did not improve the
goodness of fit, ($\chi^2(3) = 2.74$, p = 0.43). Thus, none of the tested models provided
statistical evidence to reject a zero time shift with respect to time 1. These results
were again supported by parameter estimates of Bayesian models equivalent to

the afore-mentioned LMMs, which estimated both an intercept across all SOAs
and individual intercepts per SOA to lie in intervals that included 0 (see
Supplementary Information). Model comparison via leave-one-out cross-validation
found a model without population intercept to be more credible than a model with
a common intercept (difference of expected log predictive density, ELPD: -0.2, s.e.
0.6) and a model with individual intercepts per SOA (ELPD: -2.2, s.e. 2.2).

In contrast, the average time shifts between localization errors for correct
TOJ trials at time 2 and TOJ error trials were -52 (s.e., 12) ms, -75 (s.e., 12) ms, -
87 (s.e., 14) ms, and -104 (s.e., 13) ms for the SOA 60 ms, 85 ms, 110 ms, and
135 ms, respectively. A model with a common intercept for all SOAs explained the
data significantly better than a model without an intercept ($\chi^2(1) = 29.27$, $p < 0.001$).
A model allowing for different intercepts for each SOA further improved the
goodness of fit ($\chi^2(3) = 14.3$, $p = 0.003$), indicating that localization curve's time
shift relative to time 2 depended on the respective SOA (see Supplementary Table
3 for Bayesian model estimates). Model comparison via leave-one-out cross-
validation found a model with individual SOA intercepts to be more credible than a
model with a common intercept (difference of expected log predictive density,
ELPD: -5.4, s.e. 3.8) and a model without intercept (ELPD: -7.2, s.e. 4.7).

Somewhat unexpectedly, the time shift value for the 110 ms SOA in
Experiment 2 (-87 ms) diverged more from the SOA compared to Experiment 1 (-
105 ms, averaged across all four posture conditions). However, in Experiment 2,
average time shift values generally underestimated the true SOA. Nontheless, the
Bayesian 95% intervals of the intercept estimates included the true SOA for all but
the largest SOA (i.e., 135 ms, see Supplementary Table 3). Furthermore, at time
1, the estimated intercepts were all slightly (albeit non-significantly) positive (Fig.
9 left panel, Supplementary Table 3), and when one considers the difference in
time shift (i.e., Time 2 – Time 1), the estimated values match the true SOA more
closely (-63 ms, -81 ms, -88 ms, and -120 ms for SOAs 60 ms, 85 ms, 110 ms ,
and 135 ms, respectively; Fig. 9 right panel).

Discussion

The aim of our study was to test whether participants represent the

remapped spatial location of tactile stimuli when they make spatial decisions about

tactile stimuli. Participants indicated both the target limb and the perceived location

in space of the first of two tactile stimuli in a tactile TOJ task. Presentation of stimuli

during movement implied that stimulus location depended on stimulation time,

allowing us to determine the relationship of stimulus timing and perceived stimulus

location in space. If participants had first computed the spatial location, and then

derived which limb had occupied this location at time of stimulation (*space-to-limb*

reconstruction hypothesis, see Kitazawa, 2002), incorrect hand assignment should

have been associated with external localization along the trajectory of the correct

hand; we did not find any evidence for such behavior. If participants had

represented stimulus location and stimulated limb together, and simply confused

the two due to conflict between different spatial codes (*stimulus switch hypothesis*),

incorrect hand choice should have been associated with spatial localization at the

location of the incorrect limb at the second stimulus's time. Localization error

curves were incompatible with this view, as their systematic bias differed

significantly from a hypothetical localization curve, derived from correct trials,

relating to stimulus time 2. Instead, when participants chose the incorrect hand,

their localization errors implied that they had aimed at that hand's location at the

time of the first, correct stimulus, evident in a close match of localization curves of

correct and erroneous TOJ trials when computed relative to stimulus time 1. In

other words, participants derived the reported stimulus location by combining the

time of the first, correct stimulus with the trajectory of the second, incorrectly

chosen hand, effectively indicating a location at which no stimulus had occurred –

consistent with the *time reconstruction hypothesis*. This behavior was evident for

all combinations of uncrossed and crossed start and end postures, as well as for

all tested SOAs between the two tactile stimuli. TOJ errors, thus, did not simply

reflect temporal confusion of two stimuli; instead, localization in TOJ error trials

marks the computation of tactile stimulus location based on correct stimulus timing

and movement information of a (correctly or incorrectly) implied body part.

Accordingly, limb crossing affected hand assignment, but not stimulus localization.

The pattern of hand assignment errors was in line with previous studies:

Participants made more TOJ hand assignment errors in conditions that involved

hand crossing than in conditions in which the hands were uncrossed (see Fig. 3;

Heed, Möller, et al., 2015; Hermosillo et al., 2011; Shore et al., 2002; Yamamoto

& Kitazawa, 2001). These reliable findings support the interpretation that

categorical decisions in touch, such as choosing which limb was stimulated, are

affected by weighted integration of different spatial aspects of the tactile stimulus

and configuration of the body (Badde et al., 2019; Badde & Heed, 2016; Heed &

Azañón, 2014). In contrast, localization error patterns were similar across

uncrossed and crossed start and end hand posture conditions, suggesting that arm

posture during stimulation did not affect reach-to-point localization (see Fig. 4, 5).

In particular, localization errors exhibited comparable spatial biases over time in

uncrossed and crossed conditions. Furthermore, localization error scattered

around the chosen hand was not biased towards the other hand (see Fig. 4), an

effect one might have expected if, like hand assignment, spatial localization was

subject to weighted influence of the tactile stimulus's anatomical origin as coded

by a body-based reference frame.

The dissociation between TOJ hand choice and localization responses

indicates that the two phenomena do not reflect the same process. It is widely

assumed that the weighted integration of spatial factors reflected by tactile limb

crossing effects trades off the anatomical and the external location of a tactile

stimulus (Badde & Heed, 2016; Cadieux & Shore, 2013; Kitazawa, 2002; Shore et

al., 2002; Yamamoto & Kitazawa, 2001). This assumption requires that an external

location is constructed as a prerequisite for assigning a stimulus to a hand. Our

present finding that participants incorrectly localize tactile stimuli associated with

incorrect limb choice, in contrast, implies that not stimulus location determines
hand choice, but vice versa, hand choice determines perceived stimulus location.

In turn, this conclusion is incompatible with the common view that crossing
effects, obtained in experiments that require categorical decisions such as TOJ,
are an implicit indicator of precise tactile localization and, thus, tactile remapping.
This is a seemingly surprising conclusion given that limb crossing quite obviously
creates conflict between the anatomical body side and the side of space, and it is
a strong claim that seems to invalidate the experimental logic of numerous papers
that have applied this logic. Our conclusion is, however, in fact compatible with
that of another recent study in which participants performed TOJ of tactile stimuli
presented to the hands and feet (Badde et al., 2019). In that study, stimuli were
presented to two of the four limbs (hands and feet), but it was unknown to the
participant which two limbs would be stimulated in a given trial. Accordingly, four
rather than just two response options were always available. In a percentage of
trials, participants systematically assigned the first touch to a limb that had not
received a tactile stimulus at all during the respective trial. For instance, after
stimulation of the left hand and the right foot, a participant may have indicated that
the first stimulus had occurred on the right hand. These errors depended on
whether the arms and/or legs were crossed. Importantly, neither the side of space
of the limb that had received the correct stimulus, nor the spatial distance between
stimulus and response limb, were indicative of the erroneously chosen,
unstimulated limb – again incompatible with the prevailing view that crossing
effects reflect conflict during the integration of anatomical and spatial stimulus
location. Instead, participants responded with unstimulated limbs that occupied the
space anywhere around the body side of the correct, first tactile stimulus,
independent of whether that limb was crossed or not. Thus, what mattered for limb
assignment was not the precise, spatial location of the stimulus, but much broader,
global spatial stimulus features such as an entire side of the body and its
surrounding space. Critically, these global features explained behavior without

having to recur to the use of external-spatial coding of touch location, consistent

with the present finding that the latter is only computed when necessary.

Notably, other manipulations that have been used to argue for the relevance

of precise spatial representations in tactile decisions can be framed in such a

feature-based account as well. For instance, the TOJ crossing effect was reduced

when the two hands' positions differed in height or in depth (Azañón et al., 2016).

However, this manipulation may have simply introduced an additional, non-metric

feature that helped representing the two choice options as different, thus improving

TOJ choice based on a categorical feature rather than on metric distance.

Similarly, whereas effects of distance between the two hands during a TOJ have

been interpreted as implying a metric representation of stimulus location (Roberts

et al., 2003; Shore et al., 2005), others have reported such effects only for very

small (3cm), but not other (10 cm and larger), distances between the hands (Kim

& Cruse, 2001), again suggesting that they may reflect categorically coded spatial

features, not metric stimulus location.

The reason for the apparent contradiction between previous research and

these new findings stems from the fact that typical experimental designs, in which

tactile stimuli are presented to the two stationary hands, do not allow disentangling

the prevailing view of crossing effects indicating conflict between tactile anatomical

and (precise) spatial stimulus location from the view that the conflict apparent in

limb crossing must indicate other spatial aspects, such as those identified in our

previous study (Badde et al., 2019). Thus, limb crossing paradigms that require

limb choices about the origin of touch likely reflect the integration of categorical,

tactile-spatial stimulus features. We propose that automatic effects, such as

crossmodal, tactile-visual cueing (Azañón & Soto-Faraco, 2008), too, are based

on such feature-based processing. In contrast, precise stimulus location is,

contrary to what has regularly been implied, not among the pieces of information

that are integrated for automatic, tactile-spatial coding. Instead, precise location of

the tactile stimulus in space is inferred post-hoc only when required.

Furthermore, our proposal contrasts with the suggestion that TOJ errors are

due to temporal confusion of the two stimuli, hypothesized to occur due to slowing

of a neural clock mechanism because crossed postures induce higher cognitive

load as compared to uncrossed postures (Kitazawa et al., 2008). This hypothesis

is based on the assumption that tactile locations are represented correctly but

ordered incorrectly in time. It is incompatible with our finding that participants

localized touch in incorrect TOJ trials at the location of the incorrect hand at the

correct, first stimulus time point. Moreover, an account based on time confusion is

specific to the TOJ paradigm, in which participants compare two stimuli. In

contrast, a feature-based account generalizes to other experimental paradigms,

including ones that present only a single tactile stimulus (Azañón et al., 2010;

Azañón & Soto-Faraco, 2008; Badde et al., 2015, 2019).

Our study exploited known variations in localization bias when touch is

presented during movement. We have previously proposed a computational model

for tactile localization errors that faithfully replicates these biases assuming that

participants are uncertain about the time at which the stimulus occurred relative to

the movement, and combine a probability distribution of the possible stimulus time

with the perceived arm movement trajectory (Maij et al., 2013, 2017, see also Maij

et al., 2011 for a visual analogue). The present results are compatible with the

temporal uncertainty model. Independent of which hand the stimulus was assigned

to, the temporal estimate of the tactile stimulus was identical, resulting in identical

localization error profiles, based on the trajectory of the chosen hand, in correct

and erroneous TOJ trials.

To summarize, we observed the typical dependence of tactile TOJ

responses on limb posture, with higher error rates when the hands are crossed

rather than uncrossed. Explicit localization responses of the stimulus chosen as

having occurred first were incompatible with theoretical accounts that posit

confusion of yoked stimulus representations that encompass the independently

determined external-spatial location of tactile stimuli, or projection of a body part

onto the determined spatial location of a stimulus. Instead, participants chose one

hand presumably based on categorical, spatial stimulus characteristics such as

the stimulated body side (Badde et al., 2019), and then combined the time point

associated with the first stimulus with the chosen arm's trajectory. After hand

assignment errors, participants, thus, effectively referenced a post-hoc

constructed spatial location at which no stimulus had ever occurred.

Materials & methods

627   Data for the presented analyses as well as code to run analyses and create figures

628   are provided at the Open Science Framework website, https://osf.io/ybxn5/.

Participants

Experiment 1 was performed at the Faculty of Psychology and Human Movement

Science of the University of Hamburg. Twelve right-handed participants (aged 19-

31 years, 7 female) gave informed consent to take part in the experiment. The

study was part of a research program approved by the ethics committee of the

German Psychological Society (DGPs). Experiment 2 was preregistered at the

Open Science Framework website (https://osf.io/qyzgb). A sample size of 20

participants was defined a priori. We collected data from 20 individuals from

Bielefeld University. We excluded data of 1 participant from analyses as s/he did

not follow the instructions and most of the time localized the tactile stimulus at the

start or end position, but not along the movement trajectory. Furthermore, we

excluded data of another participant as s/he only completed 384 trials in total. As

any form of data acquisition was stopped in our lab beginning of March 2020 due

to the spread of the corona virus, we did not collect data from replacement

participants. Our sample thus constisted of 18 participants (aged 18-25 years, 15

female). The experiment was approved by the ethics committee at Bielefeld

University (Ethical Application Ref: 2017-114).

Participants provided written informed consent and were compensated with €7/hr

or received course credit. All participants had normal or corrected-to-normal vision

and did not have any known perceptual, motor, or neurological disorders.

Participants took part only if, in a screening experiment, they exhibited a TOJ

crossing effect at the SOA used in the main experiment. We used this screening

procedure because individual response patterns in tactile experiments involving

hand crossing are quite variable (Badde et al., 2015; Cadieux et al., 2010;

Yamamoto & Kitazawa, 2001); however, crossing effects are highly reliable across

the entire population, so that our screening procedure does not preclude

generalization.

General setup

Participants were blindfolded. They sat on a chair at a table. A tactile stimulator

(Oticon BC 461-0/12, Oticon Ltd., London, UK) was attached to the phalanx media

of each index finger. Stimulation consisted of 200 Hz vibration for 10 ms. To mask

any noise of the vibrators, participants wore earplugs and heard white noise

through speakers (Experiment 1) or wore sound-attenuating headphones

(Superlux HD669, Superlux Enterprise Development, Shanghai, China;

Experiment 2).

Experiment 1

Apparatus, task and procedure

The position of each index finger in space was recorded with an Optotrak active,

infrared marker motion tracking system (Northern Digital Inc., Waterloo, Ontario,

Canada) at a sampling rate of 1000 Hz. One marker was positioned on the nail of

each index finger, directly next to the tactile stimulator. A data acquisition unit

(Odau; Northern Digital Inc., Waterloo, Ontario, Canada; sampling rate 1000 Hz)

synchronized marker position and timing of the tactile stimuli. The experiment was

controlled with Matlab (Mathworks, Natick, MA, USA), using the Psychophysics

Toolbox (Brainard, 1997) and the Optotrak Toolbox (http://webapp6.rrz.uni-

hamburg.de/allpsy/vf/OptotrakToolbox).

Participants moved both hands from a position of about 40 cm away from their

body towards their body to a position about 10 cm away from their body. Hand

start and end posture were uncrossed and crossed, varied in blocks of 50 trials in

pseudo-randomized order (see Fig. 2A). In each trial, a tone instructed the

movement start. At a random time (presented between 50-800 ms after the tone,

drawn from a square distribution) before, during, or after the movement, two tactile

stimuli were applied, one to each hand, at an SOA of 110 ms; the left-right order

of stimuli was pseudo-random. Upon movement completion, participants moved

the index finger that they had perceived to have been stimulated first to the location

of the first stimulus on the table. The hand remained in this location until a tone,

presented 2.5 s after the initial movement cue instructed them to lift the index

finger; this finger lift was used to identify the response hand during trajectory

analysis. Subsequently, participants repositioned the hands to their start locations.

We acquired 300 trials of each posture combination. To compensate for

obstruction of motion tracking markers, we acquired more trials for 2 participants

in the uncrossed to crossed posture and vice versa movement condition. The

experiment took approximately 4 hours, split in two-hour sessions held on different

days. Practice trials were included on each day before the experiment started until

the participant had understood, and felt confident with, the task. In total we

acquired 14.776 trials.

Analysis

Data preprocessing. Start and end of the movement were determined based on a

velocity threshold of 5 cm/s. We interpolated missing motion tracking data, for

instance due to obstruction when the hands passed each other or due to rotation

of the hands, using splines, with the restriction that movement onset and offset

could be determined. Trials were discarded when (1) marker data were missing at

movement onset and/or movement offset (11.9±2.6%; mean±sem); (2) no stimulus

localization response, indicated by finger lifting, could be detected (5.4±1.3%;

mean±sem); (3) movement duration is less than 200 ms (0.2±0.05%; mean±sem).

In total, 17.5±2.9% (mean±sem) of trials were removed.

Analysis of Temporal Order Judgements (TOJ). We considered the TOJ to be

correct when the hand used for the localization response had indeed been

stimulated first.

Localization error. We calculated the localization error, that is, the difference

between the true location of the index finger at the time of stimulation and the

reported location, that is, index finger pointing location just before finger lifting.

Errors are reported relative to the direction of the movement as a straight line

between start and end position of the hand, with positive values indicating errors

in movement direction towards the end position of the hand. The localization error

varies systematically with the stimulus time relative to movement onset

(Dassonville, 1995; Maij et al., 2011, 2013, 2017; Watanabe et al., 2009). The

localization errors were converted to an estimated localization curve by averaging

errors using a moving Gaussian window of 75 ms across a time window of -200 to

600 ms (step size 1ms) with respect to movement onset. For each participant and

posture condition, we calculated localization curves for correct TOJ trials relative

to onset of the first stimulus, for correct TOJ trials relative to onset of the second

stimulus, and for incorrect TOJ trials relative to onset of the first stimulus.

Comparison of localization errors in correct and incorrect TOJ trials. To determine

whether participants localized the stimulus relative to hand position at the first or

the second stimulus timepoint, we calculated – separately for each participant and

posture condition – the temporal shift that would produce the smallest deviations

around a single, common localization curve of the compared conditions 1) between

the incorrect TOJ localization curve and the correct TOJ localization curve relative

to the first stimulus time point and 2) between the incorrect TOJ localization curve

and the correct TOJ localization curve relative to the second stimulus time point.

Specifically, we shifted the data points of the incorrect TOJ localization curves in

time from 300 – 300 ms in steps of 1 ms and calculated the squared localization

error differences with an localization curve calculated from data points of the two

compared conditions using a common moving Gaussian average for each time

shift. We refer to the time value that minimized the summed squared error

differences with this overall construction curve as time shift (Maij et al., 2009,

2017).

We assessed the time shift for both stimulus times and, accordingly, obtained two

shift values per participant and posture condition. In some instances, we were

unable to construct a time shift due to a low number of incorrect TOJ trials for that

specific condition (18 cases out of 4 postures x 2 stimulus times x 12 participants

= 96); these data points were treated as missing data in the linear mixed model

analysis.

Statistical analysis. We assessed statistical significance of the reported

results using (Generalized) Linear Mixed Models ([G]LMM) (Bolker et al., 2009,

2009) as implemented in R version 3.6.1 (R Core Team, 2014) using packages

lme4, version 1.1-21 (Bates et al., 2015), and afex version 0.26-0 (Singmann,

2015). We estimated intercept parameters of Bayesian mixed factorial models

equivalent to LMM using packages brms, version 2.12.0 (Bürkner, 2017, 2018),

and loo, version 2.2.0 (Vehtari et al., 2017).

GLMM are adequate for analysis of binary variables such as correct vs.

incorrect responses in our TOJ task (Jaeger, 2008). Furthermore, (G)LMM and are

robust against missing data and account for differences in trial numbers across

conditions, as present in our data. All reported statistics were computed using type

3 sums of squares, as implemented in afex. For the random structure of LMM and

GLMM for TOJ analysis, we included only random intercepts, because models did

not reliably converge when random slopes were included. Models that tested time

shifts against zero used only the data corresponding to one particular reference

time point of correct trials (localization error relative to time 1 or time 2); given that

posture did not significantly modulate time shift, the respective models excluded

this factor. Accordingly, we compared a model without intercept [shift ~ 0 + (1 |

participant] against a model with intercept [shift ~ 1 + (1 | participant], effectively

testing whether a non-zero intercept significantly improved the time shift fit.

The brms R package uses STAN as backend. We ran LMM to estimate the

95% interval of the intercepts in the different time shift models. We compared

Bayesian models using the loo_compare() function of the loo R package. This

function uses leave-one-out cross-validation to compare models by assessing the

models' predictive density when each data point is omitted from fitting (Vehtari et

al., 2017).

Experiment 2

Apparatus, task and procedure

Kinematic data of the fingers was recorded using an optical motion capture system

(Visualeyez II VZ4000v, Phoenix Technologies Inc, Vancouver, BC, Canada) at

250Hz sampling frequency with markers placed on the nail of the two index fingers.

The experiment was controlled with Matlab (The MathWorks Version R2015a;

Natick, MA, USA) using the Psychophysics Toolbox (Brainard, 1997). Stimulus

presentation was controlled via custom-made hardware and triggered through a

digital acquisition card (PCI-6509, National Instruments, Austin, USA).

The procedure was largely similar to Experiment 1, except that 1) participants

made only reaches from an uncrossed to an uncrossed posture or from a crossed

to a crossed posture and 2) tactile stimuli were separated by SOAs of 60, 85, 110,

or 135 ms. Posture was varied in blocks of 64 trials in a pseudo-randomized order.

Within each block, SOA and which hand was stimulated first were pseudo-

randomized. Participants performed 28 blocks (14 of each posture combination)

for a total of 1792 trials. To compensate for marker obstruction and failure to

reliably detect finger lifting, 2 participants performed 10 additional blocks (5 of each

posture combination). As we stopped any form of data acquisition in our lab for an

indefinite time period beginning of March 2020, 2 participants performed only 19

and 23 blocks, respectively. The experiment took about 5-6 hours to complete,

split in two-hour sessions held on different days. Practice trials were included prior

to each experimental session.

Analysis

Data preprocessing. We used custom-written Matlab scripts for processing of

kinematic data. We first interpolated missing data points and resampled the data

to 1000 Hz using splines, and low-pass filtered the data using a second-order

butterworth filter with a cut-off frequency of 6 Hz. We determined movement onset/

offset of each hand as the time of the sample in which the resultant velocity of the

respective finger marker exceeded/ dropped below 5 cm/s. We excluded trials

when missing marker data could not be adequadetely interpolated (8 %), when no

stimulus localization response, indicated by finger lifting, could be detected (4.2

%), and when participants did not perform smooth, continuous, and synchronous

movements (6.8 %). In total we removed 18.2 % of the trials.

Analysis of Temporal Order Judgements (TOJ) and localization error. TOJ and

localization errors were determined as in Experiment 1. Because localization error

curves were similar regardless of start and end posture in Experiment 1, we

collapsed across postures in Experiment 2 to calculate the localization error

curves; we calculated individual curves for each participant and SOA (60, 85, 110,

and 135 ms). Localization error curves calculated separately for Experiment 2's

two posture conditions yielded qualitatively similar results (see Supplementary

Information).

Comparison of localization errors in correct and incorrect TOJ trials. Time shift

values were calculated as in Experiment 1 (separately for each participant and

SOA) using a shifting window of -300 to 300 ms (step size 1ms).

Statistical analysis. We assessed TOJ performance using a generalized mixed

model (GLMM) with factors Posture (uncrossed-uncrossed, crossed-crossed), and

SOA (60, 85, 110, 135 ms). The analysis approach for the dependence of time

shifts on SOAs followed a similar logic as Experiment 1. We first assessed the

significance of main effects and interaction of the experimental design with afex.

We then assessed whether time shifts were 0 relative to time 1 and time 2. To this

end, we compared models with a random participant factor but no fixed factors and

intercept [shift ~ 0 + (1 | participant], with a common intercept for all SOAs [shift ~1

+ (1 | participant], and with individual intercepts per SOA [shift ~ SOA + 0 + (1 |

participant] separately for shift values relative to time 1 and time 2, respectively.

823   Acknowledgements

We thank Maie Stein, Franziska Rudzik, and Nina Held for help with

dataacquisition, Volker Franz, Rainer Schäfer, and Renate Kiesewalter for their

help with the Optotrak and ODAU setup, Conrad Alting for help with programming

of Experiment 2, and Mathieu G.M. Koppen for help with LMM analysis. FM was

supported by the Netherlands Organization for Scientific Research (NWO Veni

Grant 451-12-009). TH was supported by an Emmy Noether grant of the German

Research Foundation (DFG) (He 6368/1-1). WPM received support from the

European Research Council (EU-ERC-283567) and the Netherlands Organization

for Scientific Research (NWO-VICI: 453-11-001). Data and code for the present

paper are available at the Open Science Framework website, https://osf.io/ybxn5/.

834   **References**

Allen, M., Poggiali, D., Whitaker, K., Marshall, T. R., & Kievit, R. A. (2019).
Raincloud plots: A multi-platform tool for robust data visualization.
Wellcome Open Research, 4, 63.
https://doi.org/10.12688/wellcomeopenres.15191.1

Alsmith, A. J. T., Ferrè, E. R., & Longo, M. R. (2017). Dissociating contributions
of head and torso to spatial reference frames: The misalignment
paradigm. *Consciousness and Cognition, 53*, 105–114.
https://doi.org/10.1016/j.concog.2017.06.005

Angelaki, D. E., Gu, Y., & DeAngelis, G. C. (2009). Multisensory integration:
Psychophysics, neurophysiology, and computation. *Current Opinion in*
Neurobiology, 19(4), 452–458. https://doi.org/10.1016/j.conb.2009.06.008

Atsma, J., Maij, F., Koppen, M., Irwin, D. E., & Medendorp, W. P. (2016). Causal
Inference for Spatial Constancy across Saccades. *PLOS Computational*
Biology, 12(3), e1004766. https://doi.org/10.1371/journal.pcbi.1004766

Azañón, E., Camacho, K., & Soto-Faraco, S. (2010). Tactile remapping beyond
space. *European Journal of Neuroscience, 31*(10), 1858–1867.
https://doi.org/10.1111/j.1460-9568.2010.07233.x

Azañón, E., Mihaljevic, K., & Longo, M. R. (2016). A three-dimensional spatial
characterization of the crossed-hands deficit. *Cognition, 157*, 289–295.
https://doi.org/10.1016/j.cognition.2016.09.007

Azañón, E., & Soto-Faraco, S. (2008). Changing reference frames during the
encoding of tactile events. *Current Biology, 18*(14), 1044–1049.
https://doi.org/10.1016/j.cub.2008.06.045

Azañón, E., Stenner, M.-P., Cardini, F., & Haggard, P. (2015). Dynamic Tuning of
Tactile Localization to Body Posture. *Current Biology, 25*(4), 512–517.
https://doi.org/10.1016/j.cub.2014.12.038

Badde, S., & Heed, T. (2016). Towards explaining spatial touch perception:
Weighted integration of multiple location codes. *Cognitive*
Neuropsychology, 33(1–2), 1–22.
https://doi.org/10.1080/02643294.2016.1168791

Badde, S., Heed, T., & Röder, B. (2014). Processing load impairs coordinate
integration for the localization of touch. *Attention, Perception, &*
Psychophysics, 76(4), 1136–1150. https://doi.org/10.3758/s13414-013-
0590-2

Badde, S., Heed, T., & Röder, B. (2015). Integration of anatomical and external
response mappings explains crossing effects in tactile localization: A
probabilistic modeling approach. *Psychonomic Bulletin & Review, 23*(2),
387–404. https://doi.org/10.3758/s13423-015-0918-0

Badde, S., Röder, B., & Heed, T. (2019). Feeling a Touch to the Hand on the
874   Foot. *Current Biology*, *29*(9), 1491-1497.e4.
875   https://doi.org/10.1016/j.cub.2019.02.060

876   Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-
877   Effects Models Using lme4. *Journal of Statistical Software*, *67*(1).
878   https://doi.org/10.18637/jss.v067.i01

879   Bernier, P.-M., & Grafton, S. T. (2010). Human posterior parietal cortex flexibly
880   determines feference frames for reaching based on sensory context.
881   *Neuron*, *68*(4), 776–788. https://doi.org/10.1016/j.neuron.2010.11.002

882   Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R.,
883   Stevens, M. H. H., & White, J.-S. S. (2009). Generalized linear mixed
884   models: A practical guide for ecology and evolution. *Trends in Ecology &*
885   *Evolution*, *24*(3), 127–135. https://doi.org/10.1016/j.tree.2008.10.008

886   Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–
887   436. https://doi.org/10.1163/156856897X00357

888   Bürkner, P.-C. (2017). brms: An R Package for Bayesian Multilevel Models Using
889   Stan. *Journal of Statistical Software*, *80*(1), 1–28.
890   https://doi.org/10.18637/jss.v080.i01

891   Bürkner, P.-C. (2018). Advanced Bayesian Multilevel Modeling with the R
892   Package brms. *The R Journal*, *10*(1), 395–411.

893   Cadieux, M. L., Barnett-Cowan, M., & Shore, D. I. (2010). Crossing the hands is
894   more confusing for females than males. *Experimental Brain Research*,
895   *204*(3), 431–446. https://doi.org/10.1007/s00221-010-2268-5

896   Cadieux, M. L., & Shore, D. I. (2013). Response Demands and Blindfolding in the
897   Crossed-Hands Deficit: An Exploration of Reference Frame Conflict.
898   *Multisensory Research*, *26*(5), 465–482.
899   https://doi.org/10.1163/22134808-00002423

900   Dassonville, P. (1995). Haptic localization and the internal representation of the
901   hand in space. *Experimental Brain Research*, *106*(3), 434–448.

902   Ernst, M. O., & Banks, M. S. (2002). Humans integrate visual and haptic
903   information in a statistically optimal fashion. *Nature*, *415*(6870), 429–433.
904   https://doi.org/10.1038/415429a

905   Ernst, M. O., & Di Luca, M. (2011). Multisensory perception: From integration to
906   remapping. In *Sensory cue integration* (pp. 224–250). Oxford University
907   Press, Oxford.

908   Fiehler, K., Schütz, I., & Henriques, D. Y. P. (2011). Gaze-centered spatial
909   updating of reach targets across different memory delays. *Vision*
910   *Research*, *51*(8), 890–897. https://doi.org/10.1016/j.visres.2010.12.015

Gallace, A., & Spence, C. (2005). Visual capture of apparent limb position
influences tactile temporal order judgments. *Neuroscience Letters*, *379*(1),
63–68. https://doi.org/10.1016/j.neulet.2004.12.052

Harrar, V., & Harris, L. R. (2010). Touch used to guide action is partially coded in
a visual reference frame. *Experimental Brain Research*, *203*(3), 615–620.
https://doi.org/10.1007/s00221-010-2252-0

Heed, T., & Azañón, E. (2014). Using time to investigate space: A review of
tactile temporal order judgments as a window onto spatial processing in
touch. *Frontiers in Psychology*, *5*, 76.
https://doi.org/10.3389/fpsyg.2014.00076

Heed, T., Backhaus, J., Röder, B., & Badde, S. (2016). Disentangling the
External Reference Frames Relevant to Tactile Localization. *PLOS ONE*,
11(7), e0158829. https://doi.org/10.1371/journal.pone.0158829

Heed, T., Buchholz, V. N., Engel, A. K., & Röder, B. (2015). Tactile remapping:
From coordinate transformation to integration in sensorimotor processing.
Trends in Cognitive Sciences, *19*(5), 251–258.
https://doi.org/10.1016/j.tics.2015.03.001

Heed, T., Möller, J., & Röder, B. (2015). Movement Induces the Use of External
Spatial Coordinates for Tactile Localization in Congenitally Blind Humans.
Multisensory Research, *28*(1–2), 173–194.
https://doi.org/10.1163/22134808-00002485

Hermosillo, R., Ritterband-Rosenbaum, A., & van Donkelaar, P. (2011).
Predicting Future Sensorimotor States Influences Current Temporal
Decision Making. *The Journal of Neuroscience*, *31*(27), 10019–10022.
https://doi.org/10.1523/JNEUROSCI.0037-11.2011

Jaeger, T. F. (2008). Categorical data analysis: Away from ANOVAs
(transformation or not) and towards logit mixed models. *Journal of
Memory and Language*, *59*(4), 434–446.
https://doi.org/10.1016/j.jml.2007.11.007

Kayser, C., & Shams, L. (2015). Multisensory Causal Inference in the Brain.
PLOS Biology, *13*(2), e1002075.
https://doi.org/10.1371/journal.pbio.1002075

Kim, D.-H., & Cruse, H. (2001). Two kinds of body representation are used to
control hand movements following tactile stimulation. *Experimental Brain
Research*, *139*(1), 76–91. https://doi.org/10.1007/s002210100741

Kitazawa, S. (2002). Where conscious sensation takes place. *Consciousness
and Cognition*, *11*(3), 475–477. https://doi.org/10.1016/S1053-
8100(02)00031-4

Kitazawa, S., Moizumi, S., Okuzumi, A., Saito, F., Shibuya, S., Takahashi, T.,
Wada, M., & Yamamoto, S. (2008). Reversal of subjective temporal order

due to sensory and motor integrations. In P. Haggard, Y. Rossetti, & M.
Kawato (Eds.), *Sensorimotor foundations of higher cognition* (pp. 73–97).
Oxford University Press.

Lohmann, J., & Butz, M. V. (2017). Lost in space: Multisensory conflict yields
adaptation in spatial representations across frames of reference. *Cognitive*
Processing, 1–18. https://doi.org/10.1007/s10339-017-0798-5

Maij, F., Brenner, E., & Smeets, J. B. J. (2009). Temporal Information Can
Influence Spatial Localization. *Journal of Neurophysiology*, *102*(1), 490–
495. https://doi.org/10.1152/jn.91253.2008

Maij, F., Grave, D. D. J., Brenner, E., & Smeets, J. B. J. (2011). Misjudging
where you felt a light switch in a dark room. *Experimental Brain Research*,
213(2–3), 223–227. https://doi.org/10.1007/s00221-011-2680-5

Maij, F., Wing, A. M., & Medendorp, W. P. (2013). Spatiotemporal integration for
tactile localization during arm movements: A probabilistic approach.
Journal of Neurophysiology, *110*(11), 2661–2669.
https://doi.org/10.1152/jn.00971.2012

Maij, F., Wing, A. M., & Medendorp, W. P. (2017). Afferent motor feedback
determines the perceived location of tactile stimuli in the external space
presented to the moving arm. *Journal of Neurophysiology*, *118*(1), 187–
193. https://doi.org/10.1152/jn.00286.2016

Mueller, S., & Fiehler, K. (2014a). Gaze-dependent spatial updating of tactile
targets in a localization task. *Frontiers in Psychology*, *5*, 66.
https://doi.org/10.3389/fpsyg.2014.00066

Mueller, S., & Fiehler, K. (2014b). Effector movement triggers gaze-dependent
spatial coding of tactile and proprioceptive-tactile reach targets.
Neuropsychologia, *62*, 184–193.
https://doi.org/10.1016/j.neuropsychologia.2014.07.025

Penfield, W., & Boldrey, E. (1937). Somatic motor and sensory representation in
the cerebral cortex of man as studied by electrical stimulation. *Brain*,
60(4), 389–443. https://doi.org/10.1093/brain/60.4.389

R Core Team. (2014). *R: A Language and Environment for Statistical Computing*.
R Foundation for Statistical Computing. http://www.R-project.org/

Roberts, R. D., Wing, A. M., Durkin, J., & Humphreys, G. W. (2003). Effects of
posture on tactile temporal order judgements. *Eurohaptics Conference,*
Dublin, Ireland, 300–304.

Röder, B., Kusmierek, A., Spence, C., & Schicke, T. (2007). Developmental
vision determines the reference frame for the multisensory control of
action. *Proceedings of the National Academy of Sciences*, *104*(11), 4753–
4758. https://doi.org/10.1073/pnas.0607158104

Röder, B., Rösler, F., & Spence, C. (2004). Early vision impairs tactile perception
991   in the blind. *Current Biology*, *14*(2), 121–124.
992   https://doi.org/10.1016/j.cub.2003.12.054

993   Roux, F., Djidjeli, I., & Durand, J. (2018). Functional architecture of the
994   somatosensory homunculus detected by electrostimulation. *The Journal of*
995   *Physiology*, *596*(5), 941–956. https://doi.org/10.1113/JP275243

996   Schubert, J. T. W., Badde, S., Röder, B., & Heed, T. (2017). Task demands
997   affect spatial reference frame weighting during tactile localization in
998   sighted and congenitally blind adults. *PLOS ONE*, *12*(12), e0189067.
999   https://doi.org/10.1371/journal.pone.0189067

1000   Schütz, I., Henriques, D. Y. P., & Fiehler, K. (2013). Gaze-centered spatial
1001   updating in delayed reaching even in the presence of landmarks. *Vision*
1002   *Research*, *87*, 46–52. https://doi.org/10.1016/j.visres.2013.06.001

1003   Shore, D. I., Gray, K., Spry, E., Spence, C., & others. (2005). Spatial modulation
1004   of tactile temporal-order judgments. *Perception*, *34*(10), 1251–1262.

1005   Shore, D. I., Spry, E., & Spence, C. (2002). Confusing the mind by crossing the
1006   hands. *Cognitive Brain Research*, *14*(1), 153–163.
1007   https://doi.org/10.1016/S0926-6410(02)00070-8

1008   Simon, J. R., Hinrichs, J. V., & Craft, J. L. (1970). Auditory SR compatibility:
1009   Reaction time as a function of ear-hand correspondence and ear-
1010   response-location correspondence. *Journal of Experimental Psychology*,
1011   *86*(1), 97–102.

1012   Singmann, H. (2015). *afex: Analysis of Factorial Experiments*. http://R-Forge.R-
1013   project.org/projects/afe/

1014   Thompson, A. A., Byrne, P. A., & Henriques, D. Y. P. (2014). Visual Targets
1015   Aren't Irreversibly Converted to Motor Coordinates: Eye-Centered
1016   Updating of Visuospatial Memory in Online Reach Control. *PLoS ONE*,
1017   *9*(3), e92455. https://doi.org/10.1371/journal.pone.0092455

1018   Tramper, J. J., & Medendorp, W. P. (2015). Parallel updating and weighting of
1019   multiple spatial maps for visual stability during whole body motion. *Journal*
1020   *of Neurophysiology*, *114*(6), 3211–3219.
1021   https://doi.org/10.1152/jn.00576.2015

1022   van Beers, R. J., Wolpert, D. M., & Haggard, P. (2002). When Feeling Is More
1023   Important Than Seeing in Sensorimotor Adaptation. *Current Biology*,
1024   *12*(10), 834–837. https://doi.org/10.1016/S0960-9822(02)00836-9

1025   Vehtari, A., Gabry, J., Magnusson, M., Yao, Y., & Gelman, A. (2017). *loo:*
1026   *Efficient Leave-One-Out Cross-Validation and WAIC for Bayesian Models*.
1027   https://mc-stan.org/loo/

Vehtari, A., Gelman, A., & Gabry, J. (2017). Practical Bayesian model evaluation
using leave-one-out cross-validation and WAIC. *Statistics and Computing*,
27(5), 1413–1432. https://doi.org/10.1007/s11222-016-9696-4

Watanabe, J., Nakatani, M., Ando, H., & Tachi, S. (2009). Haptic localizations for
onset and offset of vibro-tactile stimuli are dissociated. *Experimental Brain*
Research, *193*(3), 483–489. https://doi.org/10.1007/s00221-009-1711-y

Yamamoto, S., & Kitazawa, S. (2001). Reversal of subjective temporal order due
to arm crossing. *Nat. Neurosci.*, *4*(7), 759–765.
https://doi.org/10.1038/89559

1037