External location of touch is constructed post-hoc based on limb choice

Femke Maij[3], W. Pieter Medendorp[3], Tobias Heed[1,2]

[1]Faculty of Psychology and Movement Science, Bielefeld University, Bielefeld, Germany

[2]Center of Excellence Cognitive Interaction Technology (CITEC), Bielefeld University, Bielefeld, Germany

[3]Donders Institute for Brain, Cognition and Behaviour, Radboud University Nijmegen, Nijmegen, The Netherlands

Running head: Post-hoc construction of tactile external location

Correspondence: Tobias Heed, tobias.heed@uni-bielefeld.de

Abstract (150 words)

When humans indicate on which hand a tactile stimulus occurred, they often

err when their hands are crossed. This finding seemingly supports the view that

the automatically determined touch location in external space affects limb

assignment: the crossed right hand is localized in left space, and this conflict

presumably provokes hand assignment errors. Here, participants judged on which

hand the first of two stimuli, presented during a bimanual movement, had occurred,

and then indicated its external location by a reach-to-point movement. When

participants incorrectly chose the hand stimulated second, they pointed to where

that hand had been at the correct, first time point, though no stimulus had occurred

at that location. This behavior suggests that stimulus localization depended on

hand assignment, not vice versa. It is, thus, incompatible with the notion of

automatic computation of external stimulus location upon occurrence. Instead,

humans construct external touch location post-hoc and on demand.

Introduction

Spatial perception and actions rely on different spatial codes, often

associated with different reference frames. For instance, the accuracy of pointing

or reaching with an arm or finger to a visual target depends not only on the position

of target relative to gaze (Fiehler, Schütz, & Henriques, 2011; Thompson, Byrne,

& Henriques, 2014), but also on salient world-centered landmarks (Schütz,

Henriques, & Fiehler, 2013). Similarly, judgment of visual location during whole-

body movement is influenced by a target's position relative to gaze, as well as by

the location of the target relative to the body (Tramper & Medendorp, 2015).

In touch, too, space is coded in several reference frames. Touch activates

specialized sensory receptors embedded in the skin, and the arrangement of the

peripheral sensors is reflected in the homuncular organization of primary

somatosensory cortex (Penfield & Boldrey, 1937; Roux, Djidjeli, & Durand, 2018),

referred to as a skin-based or somatotopic layout. However, because our body can

take various postures, the stimulus location in space – often referred to as its

external location – must be derived by combining skin location and body posture,

a process termed tactile remapping (Heed, Buchholz, Engel, & Röder, 2015).

Indeed, there is evidence that external tactile locations can be coded in a gaze-

centered reference frame (Harrar & Harris, 2010; Mueller & Fiehler, 2014b, 2014a),

but also relative to anchors such as the head, torso, and hand (Alsmith, Ferrè, &

Longo, 2017; Heed, Backhaus, Röder, & Badde, 2016).

It is less clear, however, according to which principles these different spatial

codes are employed. Both bottom-up features such as the availability of sensory

information (Bernier & Grafton, 2010) and the spatial reliability of a sensory

channel (Ernst & Banks, 2002; van Beers, Wolpert, & Haggard, 2002), as well as

top-down information such as task-constraints (Badde, Heed, & Röder, 2015;

Schubert, Badde, Röder, & Heed, 2017), action context (Mueller & Fiehler, 2014a),

and cognitive load (Badde, Heed, & Röder, 2014) can affect the relative

contributions of different reference frames, presumably in a weighted manner

(Angelaki, Gu, & DeAngelis, 2009; Atsma, Maij, Koppen, Irwin, & Medendorp,

2016; Badde & Heed, 2016; Ernst & Di Luca, 2011; Kayser & Shams, 2015;

Lohmann & Butz, 2017; Tramper & Medendorp, 2015). Yet, whereas there is

widespread consensus that each spatial code can have more or less influence in

dependence of the specific situation, it is currently not known whether all putative

codes are always constructed, or whether they are only computed based on

demand.

For touch, it has been suggested that the construction of spatial location is

an automatic process, implying that any tactile input is remapped into an external

code, irrespective of its relevance (Heed & Azañón, 2014; Röder, Rösler, &

Spence, 2004). The most common experimental manipulation underlying this

claim is limb crossing. For a right-hand tactile stimulus, part of the right body side

lies on the left side of space when the right arm is crossed over. A task-irrelevant

tactile stimulus delivered to a crossed right hand accelerates visual discrimination

in the right visual field if it precedes the visual target stimulus by 60 ms, but on the

left side if it leads by 180 ms or more (Azañón, Camacho, & Soto-Faraco, 2010;

Azañón & Soto-Faraco, 2008). Such effects are usually interpreted as evidence

that tactile remapping – the precise computation of the external tactile stimulus

location – is automatic and forms the basis for the performance enhancement at

this external location.

The same conclusion has also been drawn from results obtained with the

tactile temporal order judgment (TOJ) task; in this task, participants report which

of two successive tactile stimuli, each presented to a different body part – typically

the two hands – occurred first (Heed & Azañón, 2014; Shore, Spry, & Spence,

2002; Yamamoto & Kitazawa, 2001). When the time interval between the two

stimuli is short, participants sometimes choose the wrong stimulus. Notably,

stimulus confusion is much more prominent when the arms are in a crossed than

in an uncrossed posture. This is surprising because the TOJ task asks about the

identity of the touched limb, and, in theory, it would be irrelevant to this question

where the hand was in space. That limb crossing, nevertheless, affects TOJ

implies that posture cannot be strategically ignored, but is automatically

incorporated into the hand assignment.

Several explanations have been put forward to account for crossing effects

in tactile localization. First, it has been suggested that touch location, once it is

remapped, is retained only in an external spatial code, and the original skin location

is discarded in the process. To report which body part has been touched, the brain

must then reversely determine which limb was located at the computed external

location at the time of the touch (Kitazawa, 2002; Kitazawa et al., 2008; Yamamoto

& Kitazawa, 2001). We refer to this suggestion as the *space-to-limb reconstruction*

hypothesis. When applied to errors in the TOJ task, this hypothesis implies that

participants correctly remap the two tactile stimuli into external space, but then

reconstruct erroneously which hand was at the first spatial location.

A second explanation assumes that TOJ errors reflect the conflict between

different codes used for stimulus location. Skin-based and external spatial codes

point to different sides of space, and this conflict must be resolved, a process that

takes time and is error-prone (Röder, Kusmierek, Spence, & Schicke, 2007; Simon,

Hinrichs, & Craft, 1970). In this view, the TOJ crossing effect is a marker for the

presence of conflict and, thus, for the fact that remapping into an external spatial

code has taken place. Notably, the interpretation that the TOJ crossing effect

derives from a remapped stimulus location is indirect because participants only

report a binary decision about which hand was stimulated, not the spatial location

of the perceived stimulus. Increasing the distance between the uncrossed hands

slightly reduces errors in TOJ (Gallace & Spence, 2005; Roberts, Wing, Durkin, &

Humphreys, 2003; Shore, Gray, Spry, Spence, & others, 2005), and the TOJ

crossing effect is smaller when the hands' positions additionally differ in height or

depth (Azañón, Mihaljevic, & Longo, 2016). These graded modulations of the TOJ

have led to the claim that the TOJ paradigm is an implicit index of tactile remapping

(Azañón, Stenner, Cardini, & Haggard, 2015; Badde & Heed, 2016; Heed &

Azañón, 2014). We refer to this suggestion as the *stimulus switch hypothesis*. It

implies that participants have correctly remapped the two stimuli into space, but

have incorrectly resolved the conflict between the different spatial codes of the first

stimulus, consequently assigning the incorrect stimulus to the first time point; as a

consequence, participants incorrectly report the hand that received the second

stimulus.

Importantly, both hypotheses outlined above assume that touch is

automatically and accurately remapped to its external location. However, recent

experiments have cast doubt on whether this is actually the case. For instance, if

a tactile stimulus is presented during an arm movement, and participants indicate

the stimulus's location by pointing to its external location after the movement, they

make systematic localization errors (Dassonville, 1995; Maij, Grave, Brenner, &

Smeets, 2011; Maij, Wing, & Medendorp, 2013, 2017; Watanabe, Nakatani, Ando,

& Tachi, 2009). Importantly, because these errors differ for fast and slow

movements, it has been suggested that participants do not compute the precise

spatial location of a stimulus when it occurs, but instead infer spatial location post-

hoc by estimating hand location at the perceived time of the tactile stimulus (Maij

et al., 2017). We refer to this suggestion as the *time reconstruction hypothesis*.

Accordingly, errors in the TOJ task would occur because participants first choose

the incorrect hand, and then derive stimulus location based on that hand's location

at the time of the first stimulus. Note, that here participants merge the correct, first

stimulus's time, together with the incorrect, second stimulus's hand. Importantly,

the claim of the *time reconstruction hypothesis* that tactile location is not

determined by integrating posture and skin location at the time of stimulus

occurrence – in the case of TOJ errors, this is the time point of the second stimulus

Figure 1. Hand assignment and tactile localization in a TOJ task with bimanual movements. Grey trace, right arm trajectory; yellow trace: left arm trajectory; red trace: reach-to-point movement to external location of the tactile stimulus. A. Correct TOJ trial: the stimulus is assigned to the correct hand, which points to the correct location. B-D Three hypotheses for tactile location in incorrect TOJ trials. B. Space-to-limb reconstruction hypothesis: participants point with the incorrect hand at the external location of the first stimulus. C. Stimulus switch hypothesis: participants point with the incorrect hand at the external location of the second stimulus. D. Time reconstruction hypothesis: participants point with the incorrect hand at thelocation at which that hand was at the time of the first stimulus.

– is at odds with the idea that tactile judgments are based on precise spatial

remapping.

Here, we assessed, in a single experiment, hand assignment and spatial

localization of tactile stimuli presented during movement. Our objective was to test

whether TOJ responses mark the use of the stimulus's precise, external-spatial

location constructed in response to the stimulus, or whether instead participants

estimate the precise stimulus location post-hoc by integrating the hand movement

trajectory with stimulus time. In other words, we aimed to directly contrast, in a

single experiment, the three discussed hypotheses for tactile localization, the

space-to-limb reconstruction hypothesis, the stimulus switch hypothesis, and the

time reconstruction hypothesis.

We presented human participants with two tactile stimuli during a bimanual

movement and assessed which hand participants perceived to have been

stimulated first (TOJ hand assignment), as well as exactly where in space

participants had perceived it (tactile stimulus localization). The experimental logic,

and its relation to the three tested tactile localization hypotheses, are illustrated in

Figure 1. Because tactile stimuli were presented near the time of a movement,

their precise spatial location depended on their timing relative to the movement.

This allowed us to determine the precise tactile spatial location participants had

perceived when they had made a hand assignment error in the TOJ task. Contrary

to common opinion, TOJ errors were not associated with the location of the

second, incorrect stimulus. Instead, when participants chose the wrong hand, they

reported stimulus location at where that hand been at the time point at which the

other, correct stimulus had occurred. Thus, participants constructed stimulus

location by combining the position of the incorrectly chosen hand with the stimulus

timing that belonged to the other, non-chosen hand's stimulus, resulting in reported

locations at which no stimulus had ever occurred. This finding invalidates current

explanations of crossing effects as being based on the precise, remapped external

spatial location of the tactile stimulus.

Results

Participants assumed a start posture with their hands resting on a table and

their arms stretched out in an uncrossed or crossed posture (see Fig. 2 for an

illustration of experimental conditions and trial timing). A tone then instructed a

movement of the two hands about 30 cm towards their body, bringing the arms

into either an uncrossed or crossed arm end posture (see Figure 2A). Shortly

Figure 2. *Experimental conditions. A. The arms moved from an uncrossed (u) or crossed (c) start posture to either an uncrossed or crossed arm end posture. B. Example TOJ trial of a representative subject, showing the bimanual movement (grey: left arm, yellow, right arm) and the reach-to-point movement of the assigned hand for the tactile localization response.*

before, during, or shortly after the movement, participants received two tactile stimuli, one on each hand, with a stimulus onset asynchrony (SOA) of 110 ms. At this SOA, participants often misreport which of the two stimuli occurred first, both when the arms are still (Heed & Azañón, 2014; Shore et al., 2002; Yamamoto & Kitazawa, 2001) and during movement (Heed, Möller, & Röder, 2015; Hermosillo, Ritterband-Rosenbaum, & van Donkelaar, 2011). After the bimanual movement, participants reported on which of the two hands the first stimulus had occurred by reaching with this hand to the perceived external location of the stimulus (see Figure 2B). The response, thus, contained two components: the hand to which the first stimulus was assigned, and explicit spatial localization of this stimulus.

Hand assignment

We measured TOJ performance as the percentage of correct reports of which hand had been stimulated first in the TOJ task, as indicated by the hand that participants used for their localization response (see Figure 3). Stimuli could occur during all times (see Methods for details), so we binned the binary (correct/incorrect) TOJ response data into four movement phases – stimulation before movement onset, during first and second half of movement, and after

Figure 3. *Probability of correct hand assignment across movement conditions (uu, uc, cu, uu, with c=crossed, u=uncrossed posture) in the four phases of the bimanual movement (before movement, during start posture, during end posture, after movement). For uc and cu conditions, stimuli were classified according to whether the first stimulus occurred while the hands were still in the start posture, or already in the end posture. For uu and cc conditions, trials were assigned as "during start posture" if stimuli had been presented during the first temporal half of the movement, and as "during end posture" otherwise. Error bars denote 2 s.d. from the mean; asymmetry is due to nonlinear conversion from the GLMM's logit scale to percentage correct. Large symbols are group means, small symbols are individual participants' performance.*

movement offset – to assess the modulation of TOJ performance by stimulus time

relative to the bimanual movement.

In accordance with previous findings, TOJ performance declined in the

crossed compared to the uncrossed posture (Heed & Azañón, 2014), and

depended on the posture at the time of stimulation (Heed, Möller, et al., 2015;

Hermosillo et al., 2011). For instance, for the uncrossed-uncrossed movement

condition (see Fig. 3, uu), the probability of correct response was high compared

to the crossed-crossed movement condition (see Fig. 3, cc) throughout all

movement phases. For the conditions with a postural change (see Fig. 3, uc & cu)

the probability of correct responses was modulated by the posture at the time of

stimulation. A generalized mixed model (GLMM) with factors Start Posture, End

Posture, and Movement Phase revealed significance for all main effects and

interactions (see Table 1 at the end of this manuscript). With movement phase, the

effect of Start Posture (see Fig. 3, dark vs. light colors) declined, whereas the effect

Figure 4. *Reach end points, as measures of localization responses, in the horizontal plane for the different movement conditions. Data are from a single participant with substantial end point variability. Ellipses cover 95% of the variability. Ellipses cover virtually identical spaces when participants give correct and incorrect responses. The space-to-limb reconstruction hypothesis would predict that, during error trials, participants point with the incorrectly chosen hand to the location of the correct stimulus; thus, if this hypothesis were correct, orange ellipses should overlay with grey ellipses, and red ellipses should overlay with yellow ellipses.*

of End Posture (see Fig. 3, blue vs. red colors) increased, over time. Note that,

due to our randomized stimulus timing strategy, this fourth movement phase

contained fewer trials than the other phases (see Materials and methods).

In sum, TOJ performance in the present experiment reflected known

modulations of hand posture and movement timing. Participants made, on

average, more than 15% TOJ errors even with uncrossed hands. This high error

rate is due to the use of the short SOA of 110 ms (Heed & Azañón, 2014; Heed,

Möller, et al., 2015; Shore et al., 2002; Yamamoto & Kitazawa, 2001), and is an

intended outcome of the experimental paradigm, allowing, as a next step,

comparison of spatial localization responses for erroneous TOJ trials across all

postures.

Explicit stimulus localization in space

Localization responses in TOJ error trials allow differentiating between the

three hypotheses about how participants determine stimulus localization in tactile

decision paradigms (see Figure 1). The *space-to limb reconstruction hypothesis*

posits that the spatial location of the first tactile stimulus is represented correctly,

and that a limb assignment entails computing which limb was at that spatial

location when the stimulus occurred. Thus, in our task, responses with the

incorrect hand would result from assigning the incorrect hand to the correct spatial

location of the first tactile stimulus (see Figure 1A). Accordingly, the chosen,

incorrect hand should be directed to the location at which the stimulus of the other,

correct hand had occurred, and the reported stimulus location in TOJ error trials

should scatter around the movement trajectory of the correct hand. Contrary to this

prediction, participants pointed to locations scattered around the movement

trajectory of the chosen, incorrect hand (see also Figure 2B), indicating that the

chosen stimulus had been perceived on the incorrect hand (see Figure 4 for the

localization responses of the participant with the largest variability in localization

errors). Thus, localization behavior did not support the implication of the *space-to-*

limb reconstruction hypothesis that the correct external spatial location is simply

assigned to a wrong limb.

Having established that participants appear to have perceived the stimulus

at the reporting hand, two possibilities remain as to which stimulus location was

associated with erroneous responses (see Fig. 1). The *stimulus switch hypothesis*

assumes that the two stimuli were localized correctly, and one is chosen for the

response. In TOJ error trials, participants would confuse the two stimuli and report

the second stimulus by pointing at its location with the respective, incorrect hand.

In this case, spatial localization errors in the reach-to-point task should scatter

around the position of the hand at the time point of the *second,* erroneously chosen

Figure 5. *Localization errors systematically vary in dependence of the progression of the movement. A. Mean localization error (blue line) of all correct localization trials of a single participant. Each dot represents the error of a single trial, that is the difference between true hand position at the time of tactile stimulation and indicated hand position at the end of the trial. Blue shading represents the average movement time, with 0 = movement start. Note, that localization bias is positive at the beginning of the movement, but negative towards the end of the movement. These biases reflect mislocalization toward the movement start point at the beginning, and toward the movement endpoint at the end of the movement. B. The error pattern in A was evident across all subjects for all posture combinations. Traces reflect the mean, shaded areas around the traces reflect s.e.m. The shaded regions in the background indicate movement times, which differed slightly between conditions.*

stimulus (referred to as time 2 from hereon). In contrast, the *time reconstruction*

hypothesis assumes that participants always use the correct, first time point (time

1 from hereon), and determine the chosen response limb's location at this time

point. This hypothesis predicts that, in TOJ error trials, participants point to the

location of the erroneously chosen, incorrect hand at the correct time, that is, time

1. Note, that no tactile stimulus occurred at this external spatial location, because

it combines the time of the first, correct stimulus with the movement trajectory of

the second, incorrect stimulus's hand.

Assessment of whether participants pointed to the hand's position at the

first or second time point during TOJ error trials is based on comparison of the

localization error curve in correct and erroneous TOJ trials. When participants

Figure 6. *Localization curves of incorrect TOJ trials (in red), averaged across subjects, for each of the four movement conditions. Curves show a similar pattern as the localization curves of the correct TOJ performance (light blue), but shifted by the stimulus onset asynchrony (dark blue).*

retrospectively point to the spatial location of a tactile stimulus presented while the

target limb was moving, their location judgment errors are systematically biased

towards the movement start point during the initial part of a movement, and

towards the movement end point during the final part of the movement (Dassonville,

1995; Maij et al., 2013, 2017; Watanabe et al., 2009), resulting in systematic

localization error curves. Such movement time-related biases were evident also in

the present data (see Figure 5). In the case of correct TOJ trials, we assume that

participants aimed, as instructed, at the location of the correct hand at time 1.

Therefore, we derive the localization error curve as the spatial difference of

reaching endpoint and hand location at time 1 (see Fig. 6, dark blue lines).

However, we can also derive a second localization error curve by computing

localization error relative to time 2 (see Fig. 6, light blue lines). This hypothetical

error curve is shifted to the left of the first curve, because time 2 is later in time

Figure 7. *Time shift of individual participants (small symbols) and average time shift (large symbols) in the four movement conditions when stimulus localization error in TOJ error trials is calculated relative to time 1 (left panel) and time 2 (right panel). Localization error was siginificantly different from zero when calculated relative to time 2, but not relative to time 1; this result is consistent with the time reconstruction hypothesis, but not with the stimulus switch hypothesis.*

relative to movement start (and, accordingly, closer in time to movement end), so

that the movement time-related localization errors should be shifted backwards in

time by the 110 ms time difference of the two stimuli. In other words, this second

error curve reflects the pattern we would expect if participants had aimed for hand

location at time 2.

The two localization error curves computed on the basis of correct TOJ trials

serve as templates against which we compare the localization error curve of TOJ

error trials (see Fig. 6 and Methods). Localization errors of incorrect TOJ trials

overlapped with localization errors of correct TOJ trials at time 1, for each of the

four start and end posture combinations. To quantify this further, we computed by

how much reaching endpoints of TOJ error trials had to be shifted to best match

either template curve derived from correct TOJ trials. If, in incorrect TOJ trials,

participants aimed for the incorrect hand's position at time 1, then the temporal

shift should be zero relative to the localization error curve in correct TOJ trials

relative to time 1; furthermore, it should around -110 ms (negative denoting a shift

towards left, see above) – compared the localization error curve of correct TOJ

trials relative to time 2. If, however, participants aimed for the incorrect hand's

position at time 2, the shift pattern should be exactly reverse, that is, zero

compared to second template curve, and near +110 ms compared to first template

curve.

Figure 7 displays the temporal shift between the localization error curve of

incorrect TOJ trials and the error curves at time 1 and time 2, as derived from

correct TOJ trials. A linear mixed model with factors Posture Condition and

Reference Time Point (localization errors for correct TOJ trials computed relative

to time 1 or time 2) and time shift of the error curve between correct and erroneous

TOJ trials as dependent variable revealed a significant effect of Reference Time

Point ($\chi^2(1) = 22.67$, $p < 0.001$), but no effect of Posture Condition ($\chi^2(3) = 6.82$,

$p = 0.078$) or interaction ($\chi^2(3) = 2.40$, $p = 0.49$) between the two factors. The time

shift between localization errors for correct TOJ trials at time 1 and TOJ error trials,

averaged across all four posture conditions, was -5 ms; if this value is not

significantly different from 0, then an LMM of only this condition should not improve

by inclusion of an intercept, as the latter would model the deviation of the average

shift away from 0. Indeed, comparison of a model with and without intercept did

not provide statistical evidence to reject a zero time shift ($\chi^2(1) = 0.36$, $p = 0.55$).

In contrast, the average time shift between localization errors for correct TOJ trials

at time 2 and TOJ error trials was -109 ms, and a model without intercept fit this

condition siginifcantly worse than a model that included the intercept ($\chi^2(1) =$

26.68, $p < 0.001$). The significant difference to time 2 suggests that participants

did not aim at the position of the second stimulus in TOJ error trials. While the non-

significant difference to to time 1 does not statistically imply equality of the error

curves in correct and incorrect TOJ trials, it suggests that, if a difference exists, it

is small.

Discussion

The aim of our study was to test whether participants represent the

remapped spatial location of tactile stimuli when they make spatial decisions about

tactile stimuli. Participants indicated both the target limb and the perceived location

in space of the first of two tactile stimuli in a tactile TOJ task. Presentation of stimuli

during movement implied that stimulus location depended on stimulation time,

allowing us to determine the relationship of stimulus timing and perceived stimulus

location in space. Had participants first computed the spatial location, and then

derived which limb had occupied this location at time of stimulation (*space-to-limb*

reconstruction hypothesis, see Kitazawa, 2002), incorrect hand assignment should

have been associated with external localization along the trajectory of the correct

hand; we did not find any evidence for such behavior. Had participants represented

stimulus location and stimulated limb together, and simply confused the two due

to conflict between different spatial codes (*stimulus switch hypothesis*), incorrect

hand choice should have been associated with spatial localization at the location

of the incorrect limb at the second stimulus's time. Localization error curves were

incompatible with this view, as their systematic bias differed significantly from a

hypothetical localization curve, derived from correct trials, relating to stimulus time

2. Instead, when participants chose the incorrect hand, their localization errors

implied that they had aimed at that hand's location at the time of the first, correct

stimulus, evident in a close match of localization curves of correct and erroneous

TOJ trials when computed relative to stimulus time 1. In other words, participants

derived the reported stimulus location by combining the time of the first, correct

stimulus with the trajectory of the second, incorrectly chosen hand, effectively

indicating a location at which no stimulus had occurred – consistent with the *time*

reconstruction hypothesis. This behavior was evident for all combinations of

uncrossed and crossed start and end postures. TOJ errors, thus, did not simply

reflect temporal confusion of two stimuli; instead, localization in TOJ error trials

marks the computation of tactile stimulus location based on correct stimulus timing

and movement information of a (correctly or incorrectly) implied body part.

Accordingly, limb crossing affected hand assignment, but not stimulus localization.

The pattern of hand assignment errors was in line with previous studies:
Participants made more TOJ hand assignment errors in conditions that involved
hand crossing than in conditions in which the hands were uncrossed (see Fig. 3;
Heed, Möller, et al., 2015; Hermosillo et al., 2011; Shore et al., 2002; Yamamoto
& Kitazawa, 2001). These reliable findings support the interpretation that
categorical decisions in touch, such as choosing which limb was stimulated, are
affected by weighted integration of different spatial aspects of the tactile stimulus
and configuration of the body (Badde & Heed, 2016; Heed & Azañón, 2014). In
contrast, localization error patterns were similar across uncrossed and crossed
start and end hand posture conditions, suggesting that arm posture during
stimulation did not affect reach-to-point localization (see Figures 4, 5). In particular,
localization errors exhibited comparable spatial biases over time in uncrossed and
crossed conditions. Furthermore, localization error scattered around the chosen
hand was not biased towards the other hand (see Figure 4), an effect one might
have expected if, like hand assignment, spatial localization was subject to
weighted influence of the tactile stimulus's anatomical origin as coded by a body-
based reference frame.

The dissociation between TOJ hand choice and localization responses
indicates that the two phenomena do not reflect the same process. It is widely
assumed that the weighted integration of spatial factors reflected by tactile limb
crossing effects trades off the anatomical and the external location of a tactile
stimulus (Badde & Heed, 2016; Badde et al., 2015; Cadieux & Shore, 2013; Heed
& Azañón, 2014; Kitazawa, 2002; Shore et al., 2002; Yamamoto & Kitazawa, 2001).
This assumption requires that an external location is constructed as a prerequisite
for assigning a stimulus to a hand. Our present finding that participants incorrectly
localize tactile stimuli associated with incorrect limb choice, in contrast, implies that
not stimulus location determines hand choice, but vice versa, hand choice
determines perceived stimulus location.

In turn, this conclusion is incompatible with the common view that crossing

effects, obtained in experiments that require categorical decisions such as TOJ,

are an implicit indicator of precise tactile localization and, thus, tactile remapping.

This is a seemingly surprising conclusion given that limb crossing quite obviously

creates conflict between the anatomical body side and the side of space, and it is

a strong claim that seems to invalidate the experimental logic of numerous papers

that have applied this logic. Our conclusion is, however, in fact compatible with

that of another recent study in which participants performed TOJ of tactile stimuli

presented to the hands and feet (Badde, Röder, & Heed, 2018). In that study, the

limbs that would receive the two stimuli for the TOJ were unknown prior to a given

trial, so that four, rather than just two, response options were always available. In

a percentage of trials, participants systematically assigned the first touch to a limb

that had not received a tactile stimulus at all during the respective trial. For

instance, after stimulation of the left hand and the right foot, a participant may have

indicated that the first stimulus had occurred on the right hand. These errors

depended on whether the arms and/or legs were crossed. Importantly, neither the

side of space of the limb that had received the correct stimulus, nor the spatial

distance between stimulus and response limb, were indicative of the erroneously

chosen, unstimulated limb – again incompatible with the prevailing view that

crossing effects reflect conflict during the integration of anatomical and spatial

stimulus location. Instead, participants responded with unstimulated limbs that

occupied the space anywhere around the body side of the correct, first tactile

stimulus, independent of whether that limb was crossed or not. Thus, what

mattered for limb assignment was not the precise, spatial location of the stimulus,

but much broader, global spatial stimulus features such as an entire side of the

body and its surrounding space. Critically, these global features explained

behavior without having to recur to the use of external-spatial coding of touch

location, consistent with the present finding that the latter is only computed when

necessary.

Notably, other manipulations that have been used to argue for the relevance

of precise spatial representations in tactile decisions can be framed in such a

feature-based account as well. For instance, the TOJ crossing effect was reduced

when the two hands' positions differed in height or in depth (Azañón et al., 2016).

However, this manipulation may have simply introduced an additional, non-metric

feature that helped representing the two choice options as different, thus improving

TOJ choice based on a categorical feature rather than on metric distance.

Similarly, whereas effects of distance between the two hands during a TOJ have

been interpreted as implying a metric representation of stimulus location (Roberts

et al., 2003; Shore et al., 2005), others have reported such effects only for very

small (3cm), but not other (10 cm and larger), distances between the hands (Kim

& Cruse, 2001), again suggesting that such effects may reflect categorically coded

spatial features, not metric stimulus location.

The reason for the apparent contradiction between previous research and

these new findings stems from the fact that typical experimental TOJ designs, in

which stimuli are presented to the two stationary hands, do not allow disentangling

the prevailing view of crossing effects indicating conflict between tactile anatomical

and (precise) spatial stimulus location from the view that the conflict apparent in

limb crossing must indicate other spatial aspects, such as those identified in our

previous study (Badde et al., 2018). Thus, limb crossing paradigms that require

limb choices about the origin of touch likely reflect the integration of categorical,

tactile-spatial stimulus features. We propose that automatic effects, such as

crossmodal, tactile-visual cueing (Azañón & Soto-Faraco, 2008), too, are based

on such feature-based processing. In contrast, precise stimulus location is,

contrary to what has regularly been implied, not among the pieces of information

that are integrated for automatic, tactile-spatial coding. Instead, precise location of

the tactile stimulus in space is inferred post-hoc only when required.

Furthermore, our proposal contrasts with the suggestion that TOJ errors are

due to temporal confusion of the two stimuli, hypothesized to occur due to slowing

of a neural clock mechanism because crossed postures induce higher cognitive

load as compared to uncrossed postures (Kitazawa et al., 2008). This hypothesis

is based on the assumption that tactile locations are represented correctly but

ordered incorrectly in time. It is incompatible with our finding that participants

localized touch in incorrect TOJ trials at the location of the incorrect hand at the

correct, first stimulus time point. Moreover, an account based on time confusion is

specific to the TOJ paradigm, in which participants compare two stimuli. In

contrast, a feature-based account generalizes to other experimental paradigms,

including ones that present only a single tactile stimulus (Azañón et al., 2010;

Azañón & Soto-Faraco, 2008; Badde et al., 2015, 2018).

Our study exploited known variations in localization biases when touch is

presented during movement. We have previously proposed a computational model

for tactile localization errors that faithfully replicates these biases assuming that

participants are uncertain about the time at which the stimulus occurred relative to

the movement, and combine a probability distribution of the possible stimulus time

with the perceived arm movement trajectory (Maij et al., 2013, 2017, see also Maij

et al., 2011 for a visual analague). The present results are compatible with the

temporal uncertainty model. Independent of which hand the stimulus was assigned

to, the temporal estimate of the tactile stimulus was identical, resulting in identical

localization error profiles, based on the trajectory of the chosen hand, in correct

and erroneous TOJ trials.

To summarize, we observed the typical dependence of tactile TOJ

responses on limb posture, with higher error rates when the hands are crossed

rather than uncrossed. Explicit localization responses of the stimulus chosen as

having occurred first were incompatible with theoretical accounts that posit

confusion of yoked stimulus representations that encompass the independently

determined external-spatial location of tactile stimuli, or projection of a body part

onto the determined spatial location of a stimulus. Instead, participants chose one

hand presumably based on categorical, spatial stimulus characteristics such as

the stimulated body side (Badde et al., 2018), and then combined the time point

associated with the first stimulus with the chosen arm's trajectory. After hand

assignment errors, participants, thus, effectively referenced a post-hoc

constructed spatial location at which no stimulus had ever occurred.

Materials & methods

Data for the presented analyses as well as code to run analyses and create figures

are provided at the Open Science Framework website, https://osf.io/ybxn5/.

The study was performed at the Faculty of Psychology and Human Movement

Science of the University of Hamburg. Twelve right-handed participants (aged 19-

31 years, 7 female) gave informed consent to take part in the experiment. All

participants were free of any known sensory, perceptual, or motor disorders. The

study was part of a research program approved by the ethics committee of the

German Psychological Society (DGPs). Participants were compensated with €7/hr

or received course credit. Participants took part only if, in a screening experiment,

they exhibited a TOJ crossing effect at the SOA used in the main experiment (see

Supplementary Information).

Experimental setup. Participants were blindfolded. They sat on a chair at a table.

A tactile stimulator (Oticon BC 461-0/12, Oticon Ltd., London, UK) was attached

to each index finger, close to the fingernail. Stimulation consisted of 200 Hz

vibration for 10 ms. To mask any noise of the vibrators, participants wore earplugs

and heard white noise through speakers.

The position of each index finger in space was recorded with an Optotrak active,

infrared marker motion tracking system (Northern Digital Inc., Waterloo, Ontario,

Canada) at a sampling rate of 1000 Hz. One marker was positioned on the nail of

each index finger, directly next to the tactile stimulator. A data acquisition unit

(Odau; Northern Digital Inc., Waterloo, Ontario, Canada; sampling rate 1000 Hz)

synchronized marker position and timing of the tactile stimuli. The experiment was

controlled with Matlab (Mathworks, Natick, MA, USA), using the Psychophysics

Toolbox [17] and the Optotrak Toolbox (http://webapp6.rrz.uni-

hamburg.de/allpsy/vf/OptotrakToolbox).

Experimental procedure. Participants moved both hands from a position of about

40 cm away from their body towards their body to a position about 10 cm away

from their body. Hand start and end posture were uncrossed and crossed, varied

in blocks of 50 trials in pseudo-randomized order (see Fig. 2A). In each trial, a tone

instructed the movement start. At a random time (drawn from a square distribution)

before, during, or after the movement, two tactile stimuli were applied, one to each

hand, at an SOA of 110 ms; the left-right order of stimuli was pseudo-random.

Upon movement completion, participants moved the index finger that they had

perceived to have been stimulated first to the location of the first stimulus on the

table. The hand remained in this location until a tone, presented 2.5 s after the

initial movement cue instructed them to lift the index finger; this finger lift was used

to identify the response hand during trajectory analysis. Subsequently, participants

repositioned the hands to their start locations. We acquired 300 trials of each

posture combination. To compensate for obstruction of motion tracking markers,

we acquired more trials for 2 participants in the uncrossed to crossed posture and

vice versa movement condition. The experiment took approximately 4 hours, split

in two-hour sessions held on different days. Practice trials were included on each

day before the experiment started until the participant had understood, and felt

confident with, the task. In total we acquired 14.776 trials.

Data analysis. Start and end of the movement were determined based on a velocity

threshold of 5 cm/s. We interpolated missing motion tracking data, for instance due

to obstruction when the hands passed each other or due to rotation of the hands,

using splines, with the restriction that movement onset and offset could be

determined. Trials were discarded when (1) marker data were missing at

movement onset and/or movement offset ($11.9\pm2.6\%$; mean$\pm$sem); (2) no stimulus

localization response, indicated by finger lifting, could be detected ($5.4\pm1.3\%$;

mean±sem); (3) movement duration is less than 200 ms (0.2±0.05%; mean±sem).

In total, 17.5±2.9% (mean±sem) of trials were removed.

Analysis of Temporal Order Judgements (TOJ). We considered the TOJ to be

correct when the hand used for the localization response had indeed been

stimulated first.

Localization error. We calculated the localization error, that is, the difference

between the true location of the index finger at the time of stimulation and the

reported location, that is, index finger pointing location just before finger lifting.

Errors are reported relative to the direction of the movement as a straight line

between start and end position of the hand, with positive values indicating errors

in movement direction towards the end position of the hand. The localization error

varies systematically with the stimulus time relative to movement onset

(Dassonville, 1995; Maij et al., 2011, 2013, 2017; Watanabe et al., 2009); for each

participant and posture condition, the localization errors were converted to an

estimated localization curve by averaging errors using a moving Gaussian window

of 75 ms.

Comparison of localization errors in correct and incorrect TOJ trials. To determine

whether participants localized the stimulus relative to the first or the second

stimulus, we calculated the temporal shift that minimized the squared difference

between the localization error curve constructed for the correct TOJ trials with the

moving window relative to the first and second stimulus time points, and the

reaching endpoints of the incorrect TOJ trials. In other words, we shifted the

endpoints of incorrect TOJ trials in time until the correct TOJ trial localization error

curve provided the best achievable fit for localization in incorrect TOJ trials. We

restricted this search to a time window of [-200 to 600 ms] around the respective

correct TOJ localization error curve. We refer to the time value that minimized the

squared difference as time shift. As the time shift was assessed for both stimulus

times, we obtained two shift values per participant. In some instances, we were

unable to construct a time shift due to a low number of incorrect TOJ trials for that

specific condition (11 cases out of 4 postures x 2 stimulus times x12 participants

= 96); these data points were treated as missing data in the linear mixed model

analysis.

Statistical analysis. We assessed statistical significance of the reported

results using (Generalized) Linear Mixed Models ([G]LMM) (Bolker et al., 2009,

2009) as implemented in R version 3.4.3 (R Core Team, 2014) through the

package lme4, version 1.1-7 (Bates, Mächler, Bolker, & Walker, 2015), and afex

version 0.21-2 (Singmann, 2015). GLMM are adequate for analysis of binary

variables such as correct vs. incorrect responses in our TOJ task (Jaeger, 2008).

Furthermore, (G)LMM and are robust against missing data and account for

differences in trial numbers across conditions, as present in our data. All reported

statistics were computed using type 3 sums of squares, as implemented in afex.

GLMM for TOJ analysis used only random intercepts, because models did not

reliably converge across all trial phases when random slopes were included. LMM

for time shift analysis used main effects, because a there was not enough data to

include the interaction. Models that tested time shifts against zero used only the

data corresponding to one particular reference time point of correct trials

(localization error relative to time 1 or time 2); given that posture did not

significantly modulate time shift, the respective models excluded this factor.

Accordingly, we compared a model with against a model without an intercept,

effectively testing whether a non-zero intercept significantly improved the time shift

fit.

Acknowledgements

We thank Maie Stein and Franziska Rudzik for help with data

acquisition,Volker Franz, Rainer Schäfer, and Renate Kiesewalter for their help

with the Optotrak and ODAU setup, and Mathieu G.M. Koppen for help with LMM

analysis. FM was supported by the Netherlands Organization for Scientific

Research (NWO Veni Grant 451-12-009). TH was supported by an Emmy Noether

grant of the German Research Foundation (DFG) (He 6368/1-1). WPM received

support from the European Research Council (EU-ERC-283567) and the

Netherlands Organization for Scientific Research (NWO-VICI: 453-11-001). Data

and code for the present paper are available at the Open Science Framework

website, https://osf.io/ybxn5/.

References

Alsmith, A. J. T., Ferrè, E. R., & Longo, M. R. (2017). Dissociating contributions of head and torso to spatial reference frames: The misalignment paradigm. *Consciousness and Cognition*, *53*, 105–114. https://doi.org/10.1016/j.concog.2017.06.005

Angelaki, D. E., Gu, Y., & DeAngelis, G. C. (2009). Multisensory integration: psychophysics, neurophysiology, and computation. *Current Opinion in Neurobiology*, *19*(4), 452–458. https://doi.org/10.1016/j.conb.2009.06.008

Atsma, J., Maij, F., Koppen, M., Irwin, D. E., & Medendorp, W. P. (2016). Causal Inference for Spatial Constancy across Saccades. *PLOS Computational Biology*, *12*(3), e1004766. https://doi.org/10.1371/journal.pcbi.1004766

Azañón, E., Camacho, K., & Soto-Faraco, S. (2010). Tactile remapping beyond space. *European Journal of Neuroscience*, *31*(10), 1858–1867. https://doi.org/10.1111/j.1460-9568.2010.07233.x

Azañón, E., Mihaljevic, K., & Longo, M. R. (2016). A three-dimensional spatial characterization of the crossed-hands deficit. *Cognition*, *157*, 289–295. https://doi.org/10.1016/j.cognition.2016.09.007

Azañón, E., & Soto-Faraco, S. (2008). Changing reference frames during the encoding of tactile events. *Current Biology*, *18*(14), 1044–1049. https://doi.org/10.1016/j.cub.2008.06.045

Azañón, E., Stenner, M.-P., Cardini, F., & Haggard, P. (2015). Dynamic Tuning of Tactile Localization to Body Posture. *Current Biology*, *25*(4), 512–517. https://doi.org/10.1016/j.cub.2014.12.038

Badde, S., & Heed, T. (2016). Towards explaining spatial touch perception: Weighted integration of multiple location codes. *Cognitive Neuropsychology*, *33*(1–2), 1–22. https://doi.org/10.1080/02643294.2016.1168791

Badde, S., Heed, T., & Röder, B. (2014). Processing load impairs coordinate integration for the localization of touch. *Attention, Perception, & Psychophysics*, *76*(4), 1136–1150. https://doi.org/10.3758/s13414-013-0590-2

Badde, S., Heed, T., & Röder, B. (2015). Integration of anatomical and external response mappings explains crossing effects in tactile localization: A probabilistic modeling approach. *Psychonomic Bulletin & Review*, *23*(2), 387–404. https://doi.org/10.3758/s13423-015-0918-0

Badde, S., Röder, B., & Heed, T. (2018). Feeling a touch to the hand on the foot: phantom percepts in healthy adults. *Submitted*.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-
Effects Models Using lme4. *Journal of Statistical Software*, *67*(1).
https://doi.org/10.18637/jss.v067.i01

Bernier, P.-M., & Grafton, S. T. (2010). Human posterior parietal cortex flexibly
determines feference frames for reaching based on sensory context.
Neuron, *68*(4), 776–788. https://doi.org/10.1016/j.neuron.2010.11.002

Bolker, B. M., Brooks, M. E., Clark, C. J., Geange, S. W., Poulsen, J. R.,
Stevens, M. H. H., & White, J.-S. S. (2009). Generalized linear mixed
models: a practical guide for ecology and evolution. *Trends in Ecology &*
Evolution, *24*(3), 127–135. https://doi.org/10.1016/j.tree.2008.10.008

Cadieux, M. L., & Shore, D. I. (2013). Response Demands and Blindfolding in the
Crossed-Hands Deficit: An Exploration of Reference Frame Conflict.
Multisensory Research, *26*(5), 465–482.
https://doi.org/10.1163/22134808-00002423

Dassonville, P. (1995). Haptic localization and the internal representation of the
hand in space. *Experimental Brain Research*, *106*(3), 434–448.

Ernst, M. O., & Banks, M. S. (2002). Humans integrate visual and haptic
information in a statistically optimal fashion. *Nature*, *415*(6870), 429–433.
https://doi.org/10.1038/415429a

Ernst, M. O., & Di Luca, M. (2011). Multisensory perception: from integration to
remapping. In *Sensory cue integration* (pp. 224–250). Oxford University
Press, Oxford.

Fiehler, K., Schütz, I., & Henriques, D. Y. P. (2011). Gaze-centered spatial
updating of reach targets across different memory delays. *Vision*
Research, *51*(8), 890–897. https://doi.org/10.1016/j.visres.2010.12.015

Gallace, A., & Spence, C. (2005). Visual capture of apparent limb position
influences tactile temporal order judgments. *Neuroscience Letters*, *379*(1),
63–68. https://doi.org/10.1016/j.neulet.2004.12.052

Harrar, V., & Harris, L. R. (2010). Touch used to guide action is partially coded in
a visual reference frame. *Experimental Brain Research*, *203*(3), 615–620.
https://doi.org/10.1007/s00221-010-2252-0

Heed, T., & Azañón, E. (2014). Using time to investigate space: a review of
tactile temporal order judgments as a window onto spatial processing in
touch. *Frontiers in Psychology*, *5*, 76.
https://doi.org/10.3389/fpsyg.2014.00076

Heed, T., Backhaus, J., Röder, B., & Badde, S. (2016). Disentangling the
External Reference Frames Relevant to Tactile Localization. *PLOS ONE*,
11(7), e0158829. https://doi.org/10.1371/journal.pone.0158829

Heed, T., Buchholz, V. N., Engel, A. K., & Röder, B. (2015). Tactile remapping:
639   from coordinate transformation to integration in sensorimotor processing.
640   *Trends in Cognitive Sciences*, *19*(5), 251–258.
641   https://doi.org/10.1016/j.tics.2015.03.001

642   Heed, T., Möller, J., & Röder, B. (2015). Movement Induces the Use of External
643   Spatial Coordinates for Tactile Localization in Congenitally Blind Humans.
644   *Multisensory Research*, *28*(1–2), 173–194.
645   https://doi.org/10.1163/22134808-00002485

646   Hermosillo, R., Ritterband-Rosenbaum, A., & van Donkelaar, P. (2011).
647   Predicting Future Sensorimotor States Influences Current Temporal
648   Decision Making. *The Journal of Neuroscience*, *31*(27), 10019–10022.
649   https://doi.org/10.1523/JNEUROSCI.0037-11.2011

650   Jaeger, T. F. (2008). Categorical data analysis: Away from ANOVAs
651   (transformation or not) and towards logit mixed models. *Journal of
652   Memory and Language*, *59*(4), 434–446.
653   https://doi.org/10.1016/j.jml.2007.11.007

654   Kayser, C., & Shams, L. (2015). Multisensory Causal Inference in the Brain.
655   *PLOS Biology*, *13*(2), e1002075.
656   https://doi.org/10.1371/journal.pbio.1002075

657   Kim, D.-H., & Cruse, H. (2001). Two kinds of body representation are used to
658   control hand movements following tactile stimulation. *Experimental Brain
659   Research*, *139*(1), 76–91. https://doi.org/10.1007/s002210100741

660   Kitazawa, S. (2002). Where conscious sensation takes place. *Consciousness
661   and Cognition*, *11*(3), 475–477. https://doi.org/10.1016/S1053-
662   8100(02)00031-4

663   Kitazawa, S., Moizumi, S., Okuzumi, A., Saito, F., Shibuya, S., Takahashi, T., …
664   Yamamoto, S. (2008). Reversal of subjective temporal order due to
665   sensory and motor integrations. In P. Haggard, Y. Rossetti, & M. Kawato
666   (Eds.), *Sensorimotor foundations of higher cognition* (pp. 73–97). Oxford:
667   Oxford University Press.

668   Lohmann, J., & Butz, M. V. (2017). Lost in space: multisensory conflict yields
669   adaptation in spatial representations across frames of reference. *Cognitive
670   Processing*, 1–18. https://doi.org/10.1007/s10339-017-0798-5

671   Maij, F., Grave, D. D. J., Brenner, E., & Smeets, J. B. J. (2011). Misjudging
672   where you felt a light switch in a dark room. *Experimental Brain Research*,
673   *213*(2–3), 223–227. https://doi.org/10.1007/s00221-011-2680-5

674   Maij, F., Wing, A. M., & Medendorp, W. P. (2013). Spatiotemporal integration for
675   tactile localization during arm movements: a probabilistic approach.
676   *Journal of Neurophysiology, 110*(11), 2661–2669.
677   https://doi.org/10.1152/jn.00971.2012

Maij, F., Wing, A. M., & Medendorp, W. P. (2017). Afferent motor feedback
determines the perceived location of tactile stimuli in the external space
presented to the moving arm. *Journal of Neurophysiology*, *118*(1), 187–
193. https://doi.org/10.1152/jn.00286.2016

Mueller, S., & Fiehler, K. (2014a). Effector movement triggers gaze-dependent
spatial coding of tactile and proprioceptive-tactile reach targets.
Neuropsychologia, *62*, 184–193.
https://doi.org/10.1016/j.neuropsychologia.2014.07.025

Mueller, S., & Fiehler, K. (2014b). Gaze-dependent spatial updating of tactile
targets in a localization task. *Frontiers in Psychology*, *5*, 66.
https://doi.org/10.3389/fpsyg.2014.00066

Penfield, W., & Boldrey, E. (1937). Somatic motor and sensory representation in
the cerebral cortex of man as studied by electrical stimulation. *Brain*,
60(4), 389–443. https://doi.org/10.1093/brain/60.4.389

R Core Team. (2014). *R: A Language and Environment for Statistical Computing*.
Vienna, Austria: R Foundation for Statistical Computing. Retrieved from
http://www.R-project.org/

Roberts, R. D., Wing, A. M., Durkin, J., & Humphreys, G. W. (2003). Effects of
posture on tactile temporal order judgements. In *Eurohaptics conference,*
Dublin, Ireland (pp. 300–304).

Röder, B., Kusmierek, A., Spence, C., & Schicke, T. (2007). Developmental
vision determines the reference frame for the multisensory control of
action. *Proceedings of the National Academy of Sciences*, *104*(11), 4753–
4758. https://doi.org/10.1073/pnas.0607158104

Röder, B., Rösler, F., & Spence, C. (2004). Early vision impairs tactile perception
in the blind. *Current Biology*, *14*(2), 121–124.
https://doi.org/10.1016/j.cub.2003.12.054

Roux, F., Djidjeli, I., & Durand, J. (2018). Functional architecture of the
somatosensory homunculus detected by electrostimulation. *The Journal of*
Physiology, *596*(5), 941–956. https://doi.org/10.1113/JP275243

Schubert, J. T. W., Badde, S., Röder, B., & Heed, T. (2017). Task demands
affect spatial reference frame weighting during tactile localization in
sighted and congenitally blind adults. *PLOS ONE*, *12*(12), e0189067.
https://doi.org/10.1371/journal.pone.0189067

Schütz, I., Henriques, D. Y. P., & Fiehler, K. (2013). Gaze-centered spatial
updating in delayed reaching even in the presence of landmarks. *Vision*
Research, *87*, 46–52. https://doi.org/10.1016/j.visres.2013.06.001

Shore, D. I., Gray, K., Spry, E., Spence, C., & others. (2005). Spatial modulation
of tactile temporal-order judgments. *Perception*, *34*(10), 1251–1262.

Shore, D. I., Spry, E., & Spence, C. (2002). Confusing the mind by crossing the
718   hands. *Cognitive Brain Research*, *14*(1), 153–163.
719   https://doi.org/10.1016/S0926-6410(02)00070-8

720   Simon, J. R., Hinrichs, J. V., & Craft, J. L. (1970). Auditory SR compatibility:
721   Reaction time as a function of ear-hand correspondence and ear-
722   response-location correspondence. *Journal of Experimental Psychology*,
723   *86*(1), 97–102.

724   Singmann, H. (2015). *afex: Analysis of Factorial Experiments*. Retrieved from
725   http://R-Forge.R-project.org/projects/afe/

726   Thompson, A. A., Byrne, P. A., & Henriques, D. Y. P. (2014). Visual Targets
727   Aren't Irreversibly Converted to Motor Coordinates: Eye-Centered
728   Updating of Visuospatial Memory in Online Reach Control. *PLoS ONE*,
729   *9*(3), e92455. https://doi.org/10.1371/journal.pone.0092455

730   Tramper, J. J., & Medendorp, W. P. (2015). Parallel updating and weighting of
731   multiple spatial maps for visual stability during whole body motion. *Journal
732   of Neurophysiology*, *114*(6), 3211–3219.
733   https://doi.org/10.1152/jn.00576.2015

734   van Beers, R. J., Wolpert, D. M., & Haggard, P. (2002). When Feeling Is More
735   Important Than Seeing in Sensorimotor Adaptation. *Current Biology*,
736   *12*(10), 834–837. https://doi.org/10.1016/S0960-9822(02)00836-9

737   Watanabe, J., Nakatani, M., Ando, H., & Tachi, S. (2009). Haptic localizations for
738   onset and offset of vibro-tactile stimuli are dissociated. *Experimental Brain
739   Research*, *193*(3), 483–489. https://doi.org/10.1007/s00221-009-1711-y

740   Yamamoto, S., & Kitazawa, S. (2001). Reversal of subjective temporal order due
741   to arm crossing. *Nat. Neurosci.*, *4*(7), 759–765.
742   https://doi.org/10.1038/89559

745 **Tables**

746 *Table 1. Generalized Linear Mixed Models of temporal order judgment accuracy.*

	df	χ^2	p
all trial phases (full model)			
start posture	16, 17	21.92	<0.001
end posture	16, 17	82.94	<0.001
movement phase	14, 17	69.36	<0.001
start posture : end posture	16, 17	10.48	0.001
start posture : movement phase	14, 17	8.26	.04
end posture : movement phase	14, 17	15.41	0.001
start posture : end posture : movement phase	14, 17	12.34	0.006
phase 1: before movement onset			
start posture	1	23.09	<0.001
end posture	1	8.48	0.004
start posture : end posture:	1	< 0.01	0.96
phase 2: first half of movement			
start posture	1	8.85	0.003
end posture	1	23.20	<0.001
start posture : end posture:	1	2.55	0.11
phase 3: second half of movement			
start posture	1	1.54	0.21
end posture	1	68.24	<0.001
start posture : end posture:	1	24.96	<0.001
phase 4: after movement offset			
start posture	1	4.15	0.04
end posture	1	22.67	<0.001
start posture : end posture:	1	2.33	0.13

747