# Unique-region phosphorylation targets LynA for rapid degradation, tuning its expression and signaling in myeloid cells

Ben F. Brian IV[1], Myra G. Nunez[1], Kathryn L. Schwertfeger[1,2,3], Tanya S. Freedman[1,2,4,*]

[1]Department of Pharmacology, University of Minnesota, Minneapolis, MN 55455, USA.

[2]Center for Immunology and Masonic Cancer Center, University of Minnesota, Minneapolis, MN 55455, USA.

[3]Department of Laboratory Medicine and Pathology, University of Minnesota, Minneapolis, MN 55455, USA.

[4]Center for Autoimmune Diseases Research, University of Minnesota, Minneapolis, MN 55455, USA.

[*]To whom correspondence should be addressed: tfreedma@umn.edu

## Abstract

The activity of Src-family kinases (SFKs), which phosphorylate immunoreceptor tyrosine-based activation motifs (ITAMs), is critical factor regulating myeloid-cell activation. In a previous paper (Freedman *et al.*, 2015) we showed in macrophages that the SFK LynA is uniquely susceptible to rapid ubiquitin-mediated degradation, functioning as a rheostat regulating ITAM signaling. We now report the mechanism by which LynA is preferentially targeted for degradation and how cell specificity is built into the LynA rheostat. Using genetic and biochemical analysis, we found that the E3 ubiquitin ligase c-Cbl preferentially targets LynA via tyrosine 32 in its unique insert region. This orthogonal mode of c-Cbl recognition depresses the steady-state level of macrophage LynA. Mast cells, however, express little c-Cbl and have correspondingly high steady-state levels of LynA. Upon activation, mast-cell LynA is not rapidly degraded, and SFK-mediated signaling is amplified relative to macrophages. Cell-specific c-Cbl expression therefore builds cell specificity into the LynA checkpoint.

## Introduction

Phosphorylation of immunoreceptor tyrosine-based activation motifs (ITAMs) by Src-family kinases (SFKs) is the first step in the activation of an innate immune response during a pathogen encounter. Initiation of cell-activating signaling typically occurs within clusters of ITAM-coupled receptors such as

FcγR (1) or the hemi-ITAM Dectin-1 (2, 3), nucleated by highly multivalent binding to immunoglobulin-

G-decorated pathogens or fungal cell-wall β-glucans respectively. Together, SFKs and phosphorylated

ITAMs trigger activation of the tyrosine kinase Syk (4). The SFKs and Syk then drive activation of

membrane-proximal signaling through adaptor proteins, cytoskeleton-modulating proteins such as the

FAK/Pyk2 kinases and Paxillin, and Tec kinases, which activate second-messenger pathways via

phosphoinositide 3-kinase and phosphoinositide phospholipases $C\gamma$ (PI3K and PLC$\gamma$1/2 respectively).

Ultimately, these early events lead to downstream signaling through Erk1/2 and other pathways (4). As

the byproducts of activated macrophages can be toxic (*e.g.* release of reactive oxygen species) and

drive inflammation (*e.g.* release of tumor necrosis factor α), the responsiveness of innate immune cells

must be tightly regulated (5-8).

Multiple mechanisms work together to tune the responsiveness of macrophages and other myeloid

cells, including negative regulation by the phosphatases CD45 and CD148 (5, 9, 10), cytoskeletal

barriers to diffusion (11), signaling via immunoreceptor tyrosine inhibitory motifs (ITIMs) (12) and

inhibitory ITAMs (13, 14), and degradation and mislocalization of signaling molecules targeted for

polyubiquitination by ubiquitin ligases (15, 16). The SFKs, which in myeloid cells typically include Fgr,

Fyn, and two splice forms each of Hck and Lyn, may also have positive and negative functions (12, 17-

19). Layered onto the traditional positive- and negative-regulatory roles of the SFKs, we have shown

that activated LynA (the longer of the two Lyn splice forms) is rapidly and specifically targeted for

polyubiquitination and degradation, forming the basis of a signaling checkpoint that blocks spurious

macrophage activation (20). The mechanism preferentially targeting LynA for polyubiquitination and the

molecular basis of this selectivity have not been elucidated previously.

Unlike Dectin-1 and FcγR signaling in macrophages, which is typically triggered in the context of

pathogen-induced µm-scale clusters of receptors (5, 9, 20), mast-cell FcεR signaling has a low

threshold for activation-- small or even monovalent antigen-IgE complexes are sufficient to induce a

signaling response (21, 22). Like macrophages, mast cells express LynA, and this disparity in receptor

sensitivity has not previously been explained.

This paper describes the mechanism by which active LynA is selectively recognized and rapidly degraded, tuning its activation kinetics and its steady-state protein levels in a cell-specific manner. To reveal the requirements for LynA degradation, we synchronized receptor-independent SFK activation using the designer inhibitor 3-IB-PP1 (23-25), which specifically inhibits a variant of the SFK-inhibitory kinase Csk (Csk[AS]) (20, 25, 26). Csk is responsible for phosphorylating a key inhibitory tyrosine in the C-terminus of all the SFKs. Inhibiting Csk[AS] leads to rapid and robust SFK activation due to the loss of the dynamic equilibrium between Csk and the phosphatases CD45 and CD148 (27, 28). We showed previously that SFK activation induced via 3-IB-PP1 treatment leads to the phosphorylation of the E3 ubiquitin ligase c-Cbl and the preferential polyubiquitination and degradation of the SFK LynA, but it was not clear whether these two events were linked. Using knockdown, overexpression, and genetic models, we now demonstrate that c-Cbl, but not its homolog Cbl-b, controls the steady-state expression of LynA and mediates its rapid degradation upon activation in macrophages. LynA is targeted by c-Cbl via a phosphorylation site, Tyr32 within its unique region, a mode of interaction distinct from the slower-phase action of c-Cbl and other E3 ligases on LynB and the other SFKs. Finally, we have discovered that the LynA checkpoint is cell specific. In mast cells, which express very little c-Cbl but abundant Cbl-b, LynA is not rapidly degraded upon activation, helping to explain why mast cells have a lower activation threshold than macrophages.

## Results

**c-Cbl mediates steady-state and activation-induced degradation of LynA in macrophages**

We reported previously that activated LynA is rapidly polyubiquitinated and degraded in Csk[AS] macrophages treated with the Csk[AS] inhibitor (SFK activator) 3-IB-PP1 (20). To identify the responsible E3 ubiquitin ligase, we tested the effects of knocking down the expression of candidate ligases using small interfering (si)RNA. Csk[AS] bone-marrow-derived macrophages (BMDMs) were transfected with non-targeting control RNA or with siRNA constructs targeting c-Cbl or Cbl-b, known modulators of ITAM signaling in both adaptive and innate immune cells (12, 15, 29). Resting Csk[AS] BMDMs were treated

with 3-IB-PP1 for up to 5 min to activate the SFKs and initiate bulk degradation of LynA. Immunoblots

of whole-cell lysates showed efficient and specific knockdown **(Figure 1A)** of c-Cbl protein (70%

reduced) and Cbl-b protein (60% reduced) **(Figure 1B)**.

Knockdown of c-Cbl led to a delay in the degradation of activated LynA, with 1 min 3-IB-PP1 treatment

reducing LynA by 30% in c-Cbl knockdown BMDMs compared to 61% and 64% in Cbl-b knockdown

and control BMDMs, respectively **(Figure 1C)**. c-Cbl knockdown also increased the level of steady-

state LynA by 60% **(Figure 1D)**. The impaired degradation and increased steady-state LynA expression

together resulted in pronounced elevations in LynA protein in the first minutes of 3-IB-PP1 treatment;

after 1 min of 3-IB-PP1 treatment, LynA was elevated 2.9-fold in cCbl-deficient BMDMs relative to

control. Although it has been shown previously that c-Cbl and Cbl-b can have redundant functions (30,

31), we were unable to detect any role for Cbl-b in the steady-state or activation-induced degradation of

LynA.

To test whether the residual LynA degradation was due to incomplete knockdown, we also generated

$Csk^{AS}Cbl^{-/-}$ mice to test LynA degradation in BMDMs lacking any c-Cbl expression. As with siRNA

knockdown, total c-Cbl knockout impaired LynA degradation in BMDMs **(Figure 1E)**; Cbl-b expression

in Csk$^{AS}$c-Cbl$^{-/-}$ BMDMs remained comparable to wild-type levels **(Figure 1F)**. Complete loss of c-Cbl

caused a more profound impairment of LynA degradation during the first few minutes of 3-IB-PP1

treatment **(Figure 1G)** a dramatic increase in steady-state LynA protein (up to 6-fold higher than in c-

Cbl$^{+/+}$ cells, **Figure 1H**). As in the knockdown experiments, LynA degradation in c-Cbl$^{-/-}$ BMDMs was not

fully eliminated but occurred on a much slower timescale, similar to the degradation of the other SFKs

(20). From this observation we conclude that c-Cbl is solely responsible for the rapid-phase degradation

unique to LynA, whereas other E3 ubiquitin ligases can complement c-Cbl to mediate slower-phase

SFK degradation. Although activating Syk phosphorylation was increased in Csk$^{AS}$c-Cbl$^{-/-}$ BMDMs

relative to Csk$^{AS}$ BMDMs treated with 3-IB-PP1 **(Figure 1E)**, Erk phosphorylation after SFK activation

was paradoxically impaired **(Figure 1–figure supplement 1)**, likely due to the loss of the adaptor

function of c-Cbl in PI3K signaling (32, 33).

**Figure 1**. **Loss of c-Cbl expression in macrophages leads to steady-state upregulation and delayed degradation of LynA protein. (A)** Immunoblots showing LynA and Cbl protein content of Csk$^{AS}$ BMDMs transfected with non-targeting control RNA (ctrl, black) or siRNA sequences targeting c-Cbl (red) or Cbl-b (blue). After overnight incubation, cells were treated with 3-IB-PP1 for the indicated times; β-actin is shown as a loading control. **(B-D)** Densitometry quantification of relative Cbl and LynA protein levels, corrected for total protein content. Error bars reflect standard error of the mean (SEM), n=3. **(B)** Quantification of c-Cbl and Cbl-b relative to ctrl expression of each (dotted line). Significant differences (Sig.) from two-way ANOVA with Sidak's multiple comparison test (asterisks): [c-Cbl expression: c-Cbl vs. Cbl-b siRNA samples], [Cbl-b expression: c-Cbl vs. Cbl-b siRNA samples]; *

P=0.0217-0.0473. **(C-D)** Quantification of LynA relative to t=0 within each siRNA condition **(C)** or the

steady-state level in the ctrl **(D)**. Sig. from two-way ANOVA with Tukey's multiple comparison test: [c-

Cbl vs. ctrl, black asterisks], [c-Cbl vs. Cbl-b, blue asterisks]; **** P<0.0001, *** P=0.0001-0.0005, **

P=0.0015-0.0058, * P=0.0105-0.0451. No significant differences (ns) between other pairs. **(E)**

Immunoblots showing LynA and Cbl protein content of Csk$^{AS}$ and Csk$^{AS}$c-Cbl$^{-/-}$ BMDMs treated with 3-

IB-PP1; Erk1/2 is shown as a loading control. **(F-H)** Quantification of relative Cbl and LynA protein

levels, corrected for total protein content. SEM, n=5. **(F)** Quantification of c-Cbl and Cbl-b in Csk$^{AS}$c-Cbl$^{-}$

$^{/-}$ BMDMs relative to expression levels in Csk$^{AS}$ BMDMs (dotted line). **(G-H)** Quantification of LynA

relative to t=0 for each genotype **(G)** or the steady-state level in Csk$^{AS}$ BMDMs **(H)**. Sig. from two-way

ANOVA with Tukey's multiple comparison test: [Csk$^{AS}$ vs. Csk$^{AS}$c-Cbl$^{-/-}$, asterisks]; **** P<0.0001, ***

P=0.0004, ** P=0.0014-0.0048. Paradoxical loss of Erk signaling in c-Cbl-deficient cells is shown in

Figure 1–figure supplement 1. Refer to Figure 1–Source Data 1.

**A tyrosine residue in the unique-region insert of LynA is required for its rapid degradation**

Susceptibility to rapid, activation-induced degradation differentiates LynA from the splice variant LynB

and the other macrophage SFKs. Hck, Fgr, LynB, and Fyn (17, 19), are degraded >10-fold more slowly

than LynA during 3-IB-PP1 treatment (20) **(Figure 2A)**. Although not expressed in macrophages, Lck is

also degraded more slowly than LynA **(Figure 2B)**.

131

**Figure 2. Fyn and Lck are not rapidly degraded upon activation.** Immunoblots and quantification of

protein levels of Fyn, Lck, and LynB in 3-IB-PP1-treated Csk$^{AS}$ BMDMs **(A)** and Jurkat cells **(B)**. Total

Erk1/2 protein (Erk) is a loading control.

As we have been unable to overexpress LynA in BMDMs by transfection, by Amaxa nucleofection, or

by lentiviral transduction, we expressed LynA ectopically in Jurkat T cells by transient transfection along

with a Cbl construct and a membrane-localized construct of Csk$^{AS}$ (memCsk$^{AS}$) (25), which sensitizes

Jurkat cells to 3-IB-PP1 for synchronized SFK activation. As a proof of principle for this model system,

we transfected Jurkat cells with wild-type LynA, memCsk$^{AS}$, and empty vector (ctrl), c-Cbl, or Cbl-b

plasmid DNA **(Figure 3A)**, which increased c-Cbl expression 3-fold and Cbl-b 25-fold relative to

endogenous levels **(Figure 3B)**. Overexpressing one Cbl family member did not affect the abundance

of the other. As predicted, c-Cbl overexpression in Jurkat cells increased the rate of LynA degradation

(30% depleted after 5 min 3-IB-PP1 treatment in c-Cbl-overexpressing Jurkat cells vs. 10% and 0% in

ctrl and Cbl-b-overexpressing cells, respectively) **(Figure 3C)**. This observation is consistent with our

siRNA experiments in BMDMs, suggesting that Jurkat cells, which do not normally express Lyn, are

capable of supporting rapid, c-Cbl-mediated degradation of LynA.

LynB, which lacks a 21 amino-acid insert found in the unique region of LynA **(Figure 3D)**, was less

efficiently degraded in c-Cbl-transfected Jurkat cells **(Figure 3E)**, depleted by 10% (LynB) vs. 40%

(LynA) from baseline after 5 min 3-IB-PP1 **(Figure 3F)**. These data mirror our findings in Csk$^{AS}$ BMDMs

(20) and confirm that the unique sensitivity of LynA to rapid degradation is preserved in the Jurkat

model. We hypothesized that the targeting site in LynA must lie within the unique-region insert

(residues 23-43) lacking in LynB (34, 35), which contains a tyrosine phosphorylation site (Tyr32) (36), a

predicted ubiquitination site (Lys40) (UbPred, 37), and a predicted threonine phosphorylation site

(Thr27) (NetPhos 3.1, 38) **(Figure 3D)**.

Substituting LynA Tyr32 with either alanine or phenylalanine completely blocked its rapid degradation

**(Figure 3E)**, reducing the depletion after 5 min 3-IB-PP1 from 40% (wt) to 10% (LynA$^{Y32A}$) or 8%

(LynA$^{Y32F}$) from baseline, levels indistinguishable from LynB **(Figure 3F)**. In contrast, mutation of the

predicted ubiquitination (LynA$^{K40R}$) or threonine phosphorylation (LynA$^{T27A}$) sites **(Figure 3E)** did not

impair LynA degradation **(Figure 3F)**.

**Figure 3. A unique tyrosine residue is required for activation-induced degradation of LynA. (A)** Immunoblots of Jurkat cells cotransfected with V5-tagged LynA, Myc-tagged memCsk[AS], and either empty vector (ctrl, black dotted), c-Cbl (orange), or Cbl-b (blue) and treated with 3-IB-PP1. Activating phosphorylation (Thr202/Tyr204) of ERK1/2 (pERK) shows activating signaling, and total ERK1/2 (ERK) shows lane loading. **(B-C)** Quantification of relative Cbl and LynA protein, corrected for total protein content. SEM, n=4. **(B)** Quantification of overexpressed c-Cbl and Cbl-b relative to endogenous (ctrl) levels (dotted line). **(C)** Quantification of LynA protein during 3-IB-PP1 treatment relative to the steady-state level. Sig. from two-way ANOVA with Tukey's multiple comparison test: [c-Cbl vs. ctrl, black asterisks], [c-Cbl vs. Cbl-b, blue asterisks]; ** P=0.0092, * P=0.0270-0.0102. Other pairs ns. **(D)** N-terminal amino-acid sequences of mouse LynA and LynB (34, 35), including the 21 amino-acid unique insert of LynA. Tyrosine 32 (olive), threonine 27 (light gray), and lysine 40 (dark gray) are labeled. **(E)** Immunoblots of 3-IB-PP1-treated Jurkat cells cotransfected with c-Cbl, Myc-tagged memCsk[AS], and V5-tagged Lyn constructs, including wild-type LynA (black, dashed), LynA[Y32A] (olive),

LynB (teal, dotted), LynA$^{K40R}$ (dark gray), LynA$^{T27A}$ (light gray), or LynA$^{Y32F}$ (orange). **(F)** Quantification of LynA protein levels, corrected for total protein, relative to the steady-state level. SEM, n=7 for LynA, n=4 for LynA$^{Y32A}$ and LynB; n=3 for LynA$^{K40R}$, LynA$^{T27A}$, and LynA$^{Y32F}$. Sig. from two-way ANOVA with Tukey's multiple comparison test: [LynA$^{Y32A}$ vs. LynA, olive asterisks], [LynB vs. LynA, teal asterisks], [LynA$^{Y32F}$ vs. LynA, orange asterisks], [LynA$^{K40R}$ vs. LynA$^{Y32A}$, dark gray asterisks], [LynA$^{T27S}$ vs. LynA$^{Y32A}$, light gray asterisks]; **** P<0.0001, *** P=0.0006, * P=0.0166. ns: [LynA$^{T27S}$ and LynA$^{K40R}$ vs. each other and LynA] and [LynA$^{Y32A}$ and LynA$^{Y32F}$ vs. each other and LynB]. Note: the error bar for LynA$^{Y32F}$ is smaller than the line width. Relative expression levels of LynA variants and LynB are shown in Figure 3–figure supplement 1. Refer to Figure 3–Source Data 1.

As in BMDMs, the efficiency of Lyn degradation also affects its steady-state expression in Jurkat cells. LynB is 2- and LynA$^{Y32A}$ 3-fold more highly expressed than wild-type LynA, LynA$^{K40R}$, or LynA$^{T27A}$ in resting cells, resulting in 3- to 4-fold more protein remaining after 20 min treatment with 3-IB-PP1 **(Figure 3–figure supplement 1)**. Overall, we conclude that the unique tyrosine residue in the LynA insert, a known site of tyrosine phosphorylation (36), is the recognition site for rapid, c-Cbl-mediated degradation, and that this mechanism determines the half-life of activated LynA protein within an ITAM signaling complex and tunes the steady-state expression of LynA protein. This suggests that c-Cbl-mediated regulation of steady-state LynA protein levels occurs as a direct effect of c-Cbl protein activity rather than merely a transcriptional feedback effect.

**LynA kinase activity is required for its rapid degradation**

SFK activation is a requirement for rapid degradation of LynA protein in BMDMs (20), but whether this is direct effect of the LynA kinase itself or initiated by feedback from downstream signaling proteins has been unclear. To investigate the role of LynA kinase activity in its degradation, we tested two functionally impaired LynA constructs. First, we introduced a lysine residue in place of an activation-loop threonine (T410K, c-Src residue 429 (39)) to disrupt substrate recognition (40). Second, we introduced a phenylalanine residue in place of the activation-loop tyrosine, a known site of interaction

between the phosphotyrosine binding (PTB) domain of c-Cbl and Src (41) and the key

autophosphorylation site that stabilizes the active state (27, 42) (Y397F, c-Src residue 416).

**Figure 4. SFK catalytic activity is required for rapid degradation of LynA. (A)** Immunoblots

showing ectopic expression of wild-type (wt, black dotted), substrate-binding-impaired (T410K, gray), or

activation-loop-mutated (Y397F, orange) V5-tagged LynA protein coexpressed in Jurkat cells with

memCsk$^{AS}$ and c-Cbl and treated with 3-IB-PP1. pERK is shown as a control for 3-IB-PP1 treatment. β-

actin reflects lane loading. **(B)** Overexpression of wt and variants of LynA in JCaM1.6 cells. ZAP70

(Y319) and ERK are phosphorylated as a consequence of SFK activation. **(C)** Quantification of LynA

protein in transfected Jurkat cells during 3-IB-PP1 treatment, corrected for total protein, relative to the

steady-state level for each variant. SEM, n=3. Sig. from two-way ANOVA with Tukey's multiple

comparison test: [T410K vs. wt, gray asterisks], [Y397F vs. wt, orange asterisks]; ** P=0.0015-0.0083.

Other pairs ns. **(D)** Immunoblots showing expression of endogenous LynA protein in Csk$^{AS}$ BMDMs

treated with 3-IB-PP1 (3IB only, black dotted) or with 3-IB-PP1 plus the Syk inhibitor Bay-61-3606 (Bay,

light olive), the SFK inhibitor PP2 (light purple), or the FAK/Pyk2 inhibitor PF-431396 (green).

In untransfected Jurkat cells ERK1/2 is phosphorylated downstream of the endogenous SFKs Lck (43, 44) and Fyn (45), which are activated following treatment with 3-IB-PP1 (25). Since the endogenous kinases are not disrupted by substitutions in transfected LynA, ERK1/2 continues to be efficiently phosphorylated in cells transfected with LynA T410K or Y397F **(Figure 4A)**. In JCaM1.6 cells, an Lck-deficient Jurkat strain (43, 46), ERK1/2 phosphorylation depended on transfected LynA and reflected the functional impairment of LynA T410K and Y397F **(Figure 4B)**. As reported previously, Fyn activity in the absence of Lck cannot mediate ERK1/2 phosphorylation (43).

Functional LynA kinase was required for c-Cbl-mediated degradation in Jurkat cells. While wild-type LynA protein was 50% depleted after 20 min of 3-IB-PP1 treatment, degradation of either LynA T410K or Y397F was undetectable **(Figure 4C)**. The dependence on LynA kinase function was evident in both Lck- and Fyn-expressing Jurkat cells **(Figure 4A)** and Lck-deficient JCaM1.6 cells **(Figure 4B)**, indicating that, in this system, the other activated SFKs could not target LynA in trans to induce c-Cbl-mediated degradation. We do not yet know whether the unique insert of LynA is a substrate for the other macrophage SFKs, but the peptide containing Tyr32 is a predicted SFK substrate (NetPhos 3.1, 38).

In macrophages and other hematopoietic cells the SFKs are the first kinases to be activated upon antigen receptor engagement, phosphorylating intracellular ITAMs, ITAM-associated Syk or Zap70, and integrin-associated FAK or Pyk2 (4, 47). We have reported that Syk and FAK are phosphorylated on activating tyrosine residues in Csk$^{AS}$ BMDMs treated with 3-IB-PP1 (20). To test whether these downstream kinases might participate in negative feedback leading to LynA degradation, we treated

Csk[AS] macrophages with competitive inhibitors of Syk (48), FAK/Pyk2 (49), or the SFKs themselves

(50) in combination with 3-IB-PP1. As controls for on-target effects, we demonstrated that Erk1/2 and

SFK activation-loop phosphorylation was blocked in samples cotreated with the pan-SFK inhibitor PP2,

that Erk1/2 but not SFK activation-loop phosphorylation was blocked in samples cotreated with the Syk

inhibitor BAY-61-3606, and that Paxillin phosphorylation was blocked in samples cotreated with the

FAK/Pyk2 inhibitor PF-431396 **(Figure 4D)**. As with genetically impairing LynA kinase function,

inhibiting SFK activity with PP2 abrogated LynA degradation **(Figure 4D-E)**. In contrast, LynA

degradation in BMDMs was unaffected by cotreatment with BAY-61-3606 or PF-431396. Together

these data suggest that Syk and FAK/Pyk2 (and any off-target effects of BAY-61-3606 and PF-431396)

do not participate in a larger negative-feedback process, suggesting that LynA might target itself

directly through Tyr32 in cis or trans. Interestingly, since the T-cell SFKs Lck and Fyn cannot effect

LynA degradation in trans, this feedback process could be limited to a kinase effect in cis or to another

macrophage-expressed Src family member. This tight feedback circuit could explain the strikingly fast

kinetics of LynA degradation.

**Differential Cbl expression tunes LynA protein levels and signaling responses in macrophages**

**and mast cells**

Analysis of data from the Immunological Genome Project shows that expression levels of Lyn and c-Cbl

RNA are differentially regulated across different cell types. Mast cells, for instance, express very low

levels of c-Cbl mRNA **(Figure 5)** (51, 52), which corresponds to a low level of c-Cbl protein expression

(53). Cbl-b mRNA, in contrast, is abundant in mast cells.

As in macrophages, Lyn is a key regulator of mast-cell ITAM signaling (54), but the effect of disparate

Cbl-family expression on LynA in the two cell types remains unknown. We generated bone-marrow-

derived mast cells from Csk[AS] mice (hereafter referred to as "mast cells") and performed surface marker

analysis as we had done already for BMDMs (20). Csk[AS] mast cells were homogenous in their

expression of traditional mast-cell markers FcεR1α and c-Kit (CD117) (55) **(Figure 6A)**.

**Figure 5. LynA, c-Cbl, and Cbl-b mRNAs are regulated differentially in myeloid cells.** Mouse RNAseq data were derived from the Immunological Genome Project (ImmGen, http://rstats.immgen.org/Skyline/skyline.html, 51, 52). RNA levels of Lyn, c-Cbl, and Cbl-b are shown for macrophages, mast cells, neutrophils, dendritic cells, and blood monocytes. Refer to Figure 5–Source Data 1.

**Figure 6. c-Cbl expression correlates with LynA protein levels and pathway activation in macrophages and mast cells. (A)** Flow cytometry plots showing surface expression of the mast-cell markers FcεR1α and c-Kit (CD117) in Csk[AS] bone-marrow-derived mast cells compared to a peritoneal B-cell control. **(B)** Immunoblots of Csk[AS] bone-marrow-derived mast cells (mast resting, red) and BMDMs after resting (light teal) or IFN-γ priming (dark teal). LynA, c-Cbl, Cbl-b, and pErk1/2 are shown; total Erk1/2 reflects lane loading. **(C)** Quantification of c-Cbl and Cbl-b protein in mast cells and macrophages, corrected for total protein, relative to the level in primed macrophages (mΦ). SEM, n=3. Sig. from one-way ANOVA with Sidak multiple comparison test: [mast vs. BMDM resting, asterisk]; ** P= 0.0035. Others pairs ns. **(D-E)** Quantification of LynA, corrected for total protein, relative to the steady-state level in each cell condition **(D)** or in resting BMDMs at t=0 **(E)**. SEM, n=3 for mast cells and primed BMDMs, 4 for resting BMDMs. Sig. from two-way ANOVA with Tukey's multiple comparison test: [mast vs. BMDM resting, light teal asterisks], [mast vs. BMDM primed, dark teal asterisks]; *** P=0.0004-0.0008, ** P=0.0025-0.0088, * P=0.0134-0.0466. Other pairs ns. **(F)** Quantification of the maximum increase in pErk1/2 over the first 5 min of 3-IB-PP1 treatment, corrected for total Erk1/2 protein, relative to the level in resting BMDMs at t=0. SEM, n=3 for mast cells, 5 for resting BMDMs, and 4 for primed BMDMs. Sig. from one-way ANOVA with Tukey's multiple comparison test: [mast vs. BMDM resting, light teal asterisks]; ** P= 0.0085. Others pairs ns. Refer to Figure 6–Source Data 1.

We then used immunoblotting to compare the expression of Cbl and LynA protein in Csk[AS] mast cells

and BMDMs **(Figure 6B)** and found that mast cells express very little c-Cbl (20% of the level in resting

macrophages) and relatively high levels of Cbl-b (80% of the level in resting macrophages) **(Figure**

**6C)**. Strikingly, low c-Cbl expression was accompanied by a complete resistance of LynA to

degradation in the first 5 min of 3-IB-PP1 treatment **(Figure 6D)** and an increase in steady-state

expression of LynA (30% higher than in resting BMDMs) **(Figure 6E)**, resulting in 3-4x more LynA

protein in mast cells than in resting BMDMs and 2x more than in primed BMDMs between 3 and 5 min

of 3-IB-PP1 treatment. Mast cells also had a stronger signaling response to 3-IB-PP1. Although the

kinetics of Erk phosphorylation in mast cells varied among experiments, the fold increase in pErk1/2 in

the first 5 min of 3-IB-PP1 treatment was consistently higher than in macrophages, even primed

BMDMs (40x pErk increase in mast cells compared with 14x in primed BMDMs and no significant

increase in resting BMDMs) **(Figure 6F)**. We therefore conclude that c-Cbl expression tunes the

steady-state expression of LynA protein and the cell's ability to target LynA for rapid degradation upon

activation. Since mast cells are less dependent on c-Cbl for positive signaling, long-lived LynA

potentiates a strong response to 3-IB-PP1. Thus mast cells bypass the LynA signaling checkpoint by

maintaining low levels of c-Cbl and therefore high levels of activated LynA.

## Discussion

Both c-Cbl and Cbl-b negatively regulate ITAM signaling, recognizing activated signaling proteins via

PTB binding to activating phosphotyrosine sites. Cbl-mediated polyubiquitination leads to protein

degradation via the lysosome and proteasome (56). Cbl-family ligases must in turn be tightly regulated

to prevent hyperresponsive or aberrant signaling while maintaining proper pathogen clearance (57).

Although highly homologous, c-Cbl and Cbl-b target different subsets of signaling proteins for

degradation (58, 59), with c-Cbl targeting the SFKs, including Lyn (60, 61), and Cbl-b interacting with

ITAM-coupled receptors and other kinases (31, 53, 57). c-Cbl is thought to interact with active SFKs

through the SFK activation-loop phosphotyrosine residue, which binds the c-Cbl PTB domain (62), and

through the SFK SH3 domain, which binds the c-Cbl PXXP motifs (63). In this paper we report a new

mode of recognition specific to LynA. Although LynA and LynB have identical activation kinetics and

activation-loop and SH3-domain sequences, LynB is degraded 10-fold more slowly in macrophages

treated with 3-IB-PP1 (20). This rapid degradation profile is unique among the SFKs we have tested

(LynB, HckA, HckB, Fgr (20), and Fyn in BMDMs and Lck and Fyn in Jurkat cells). By exploiting the

similarity between LynA and LynB, we conclude that, while activation-loop phosphorylation and SH3-

domain interactions mediate slower-phase targeting by c-Cbl and other E3 ligases, rapid degradation of

LynA is triggered by an interaction unique to its insert region.

A single tyrosine residue (Tyr32) peculiar to its unique region marks activated LynA for rapid

degradation. This phenomenon is easily observed upon bulk SFK activation by 3-IB-PP1 but also

affects the steady-state level of LynA protein, likely due to the degradation of basally active LynA over

time. Phosphorylation of LynA Tyr32 has been observed in neuroblastoma cell lines upon receptor-

tyrosine-kinase activation (64). Further, in epidermoid carcinoma (A431) cells and breast tumor

samples LynA Tyr32 is phosphorylated by the epidermal growth factor receptor (EGFR), allowing it to

phosphorylate MCM-7 and stimulate cell proliferation (36). In macrophages this potentially positive-

regulatory function of Tyr32 paradoxically causes LynA to trigger its own c-Cbl-mediated degradation,

underlying its function as a rheostat in the LynA checkpoint. We can speculate about two possible

mechanisms. First, by analogy to a similarly situated but non-homologous transautophosphorylation

site in Hck (Tyr[29]) (65), LynA phospho-Tyr32 may be directly activating. The unique regions of LynA

and LynB make distinct SH3-domain contacts (66), and phospho-Tyr32 could alter these interactions.

Hyperactivated LynA could then efficiently phosphorylate and activate nearby or pre-complexed c-Cbl,

triggering its own degradation more aggressively than do the other SFKs. Alternatively, phospho-Tyr32

could directly or indirectly serve as a docking site for c-Cbl, constituting a unique interaction site to

efficiently recruit its degradation machinery. Bolstering this hypothesis is a published study showing that

the interactome of LynA[Y32F] transfected into mammary adenocarcinoma (MDA-MB-231) cells more

closely resembles the interactome of wild-type LynA than LynB, but some LynA interactions are lost

(67). It is also possible that LynA Tyr32 is not phosphorylated, but is required for an allosteric

intramolecular interaction or docking of interacting proteins.

We have shown that SFK activity is required for LynA degradation, and LynA Tyr32 is a predicted

substrate for Hck (NetPhos 3.1, 38) in addition to EGFR. It is therefore tempting to speculate that LynA

Tyr32 is a site of autophosphorylation, since the SFKs Lck and Fyn cannot mediate the degradation of

kinase-impaired LynA in trans. On the other hand, we have shown that LynA and LynB are activation-

loop phosphorylated 20x more quickly than Hck and Fgr in response to 3-IB-PP1 (20), and it is possible

that this delay could function as a timer, allowing a rapid burst of LynA activation with a short half-life,

either globally (in the case of 3-IB-PP1 treatment) or at the single-molecule scale within a phagocytic

synapse.

LynA and LynB have been reported to interact with different subsets of proteins in mast cells and in

triple-negative breast cancer cells, with LynA signaling via ITAM (18), cytoskeletal, and proliferative (67)

pathways and LynB initiating negative feedback via ITIMs and phosphatases (18). In mammary

epithelial cells the ratio of LynA and LynB is actively regulated by Epithelial Splicing Regulatory Protein

1 (ESRP1), with LynA-upregulated tumors having the more invasive phenotype (67). The positive-

regulatory roles reported for LynA in mast and tumor cells complements our own observations that

LynA degradation can block macrophage signaling through the Erk, Akt, and NFAT pathways, which

cannot be complemented by active LynB (20).

Aberrant activation driven by ITAM signaling pathways in macrophages is a known driver of

autoimmune and inflammatory disease, with activated macrophages accumulating in chronically

inflamed tissues in the absence of an active infection (68). Regulation of LynA protein expression and

deactivation kinetics by c-Cbl, could, via the LynA checkpoint, help to prevent the initiation of

pathological signaling. The threshold for macrophage activation is modifiable by changes to their local

environment. For example, IFN-$\gamma$ and LPS polarize macrophages for a pro-inflammatory response,

whereas IL-4 and IL-13 polarize macrophages for tissue repair (i.e. collagen deposition) and

inflammatory resolution (69, 70). We have reported that IFN-$\gamma$ decreases the macrophage signaling

threshold in part by increasing the expression of LynA (20). Dynamic changes in SFK and c-Cbl levels

could modulate the macrophage activation threshold, ensuring that macrophages respond appropriately

during times of infection (low threshold) and limiting aberrant activation in response to cellular debris

and small-scale antibody complexes during inflammatory resolution (high threshold).

Like macrophages, mast cells reside in nearly every bodily tissue and perform environment-specific

functions in addition to sensing non-self (69, 71). While macrophages have distinct anti-inflammatory

roles as professional phagocytes in the silent clearance of apoptotic cells and agents of wound healing,

mast cells are constitutively primed for ITAM-induced triggering, releasing preformed granules that

contain inflammatory cytokines, chemokines, prostaglandins, and proteases. Pursuant to these differing

functions, macrophages continuously gauge ITAM ligand valency (*i.e.* particle size) and have a

relatively high basal threshold for inflammatory activation (5, 20), while mast cells can be triggered by

small-scale receptor clustering induced by low-valency or even monovalent FcεR-IgE-allergen

complexes (21, 22). One striking difference between the macrophage and mast-cell ITAM regulatory

machinery is that mast cells express almost no c-Cbl. In spite of low reported mRNA levels (ImmGen,

51, 52), mast cells constitutively express a high level of LynA protein, likely due to impaired steady-

state degradation of basally active LynA. Furthermore, LynA is not degraded upon activation and 3-IB-

PP1 triggers a stronger Erk phosphorylation response in the absence of ITAM-receptor ligation in

resting mast cells than in resting macrophages.

Overall, we describe a model in which regulation of LynA and c-Cbl levels helps to determine an

immune cell's potential for inflammatory ITAM signaling. Regulated degradation tunes down the LynA

rheostat, blocking cell signaling via the LynA checkpoint at low cellular doses of LynA and overriding

the LynA checkpoint at higher doses. In macrophages, this occurs on a continuum where LynA dose is

highest in M1-polarized cells but still susceptible to rapid degradation by c-Cbl. Mast cells express

almost no c-Cbl and maintain high levels of LynA at steady state and over time, consistent with

permissive signaling in the absence of large-scale ITAM receptor clustering nucleated by multivalent

receptor ligation. Appropriate regulation of immune receptor thresholds is critical for maintaining the

function of innate immune cells; threshold dysregulation can lead to chronic feedback loops that drive

inflammatory signaling in autoimmune disease and conversely tumor-supporting immunosuppressive

signaling (68). Elucidating the mechanisms by which the LynA checkpoint is regulated may allow us to

tune immune-cell sensitivity and reprogram pathological cells without causing immunodeficiency.

## Materials and Methods

**Mice**

C57BL/6-derived $Csk^{AS}$ mice are hemizygous for the $Csk^{AS}$ BAC transgene on a $Csk^{-/-}$ background, as

described previously (20, 26). Due to sterility of c-Cbl$^{-/-}$ male mice, we were not able to produce a

sustained lineage of $Csk^{AS}Cbl^{-/-}$ mice, but we obtained three individuals by crossing $c$-$Cbl^{-/-}$ female mice

from E. Peterson (University of Minnesota) (72) with $Csk^{+/-}$ (26) and $Csk^{AS}$ male mice from our colony

and then crossing $Cbl^{+/-}Csk^{+/-}$ with $Cbl^{+/-}Csk^{AS}$ mice. All mice were housed in specific pathogen-free

conditions and genotyped using real-time PCR (Transnetyx, Inc., Memphis, TN).

**Jurkat cell lines and transfection**

The Jurkat T-cell strains E6-1 ("wild-type") (73) and JCaM1.6 (Lck-deficient) (43, 46) were gifts from the

laboratories of A. Weiss (University of California, San Francisco) and Y. Shimizu (University of

Minnesota), respectively. Jurkat cell lines were cultured in RPMI-1640 medium supplemented with 5%

fetal calf serum (Omega Scientific, Inc., Tarzana, CA) and 2 mM glutamine, penicillin and streptomycin

(Sigma-Aldrich, St. Louis, MO) as described previously *(74)*. Jurkat and JCaM1.6 cells were transiently

transfected via electroporation, as described previously (74). Briefly, cells were grown overnight in

antibiotic-free RPMI-1640 medium supplemented with 10% fetal bovine serum (Omega Scientific) and 2

mM glutamine (RPMI10). Batches of 15 M cells were resuspended in RPMI10 with 15 µg plasmid DNA

per construct. Cells were rested 20 min, electroporated 285 V/10 ms in a BTX square-wave

electroporator (Harvard Apparatus, Holliston, MA), resuspended in 10 mL RPMI10, and allowed to

recover overnight. One million live cells were then resuspended in phosphate-buffered saline, rested for

30 min at 37°C, and stimulated.

**DNA constructs and mutagenesis**

Plasmids containing V5+His-tagged mouse LynA and LynB were gifts from J. Rivera/R. Suzuki

(National Institutes of Health) (18). Myc-tagged mouse memCsk$^{AS}$ (also referred to as Lck$_{11}$-Csk$^{AS}$) (25)

and Xpress-tagged human c-Cbl and Cbl-b (75) were gifts from A. Weiss (University of California, San

Francisco). Site-directed mutagenesis was performed using QuikChange Lightning (Agilent

Technologies, Santa Clara, CA). Sequences of all constructs were verified by Sanger sequencing

(GENEWIZ, South Plainfield, NJ).

**Preparation and flow cytometry of myeloid cells**

BMDMs were prepared using standard methods (76). Briefly, bone marrow was extracted from femura

and tibiae of mice aged 6-8-weeks. After hypotonic lysis of erythrocytes, BMDMs were derived on

untreated plastic plates (BD Falcon, Sigma-Aldrich) by culturing in Dulbecco's Modified Eagle Medium

(DMEM, Corning Cellgro, Corning, NY) containing approximately 10% heat-inactivated fetal calf serum

(Omega Scientific), 0.11 mg/ml sodium pyruvate (Corning), 2 mM penicillin/streptomycin/L-glutamine

(Sigma-Aldrich), and 10% CMG-14-12-cell-conditioned medium as a source of M-CSF (77). After 6 or 7

days cells were resuspended in enzyme-free EDTA buffer and replated in untreated 6-well plates (BD

Falcon, Sigma-Aldrich) at 1 M cells per well in unconditioned medium with or without priming in 25 U

IFN-γ (PeproTech, Rocky Hill, NJ). Bone-marrow-derived mast cells were prepared as described

previously (55), as above except for the substitution of 10 ng/ml IL-3 (PeproTech) for CMG-14-12-

conditoned medium. After 5 weeks, cells were subjected to flow-cytometry analysis (LSRFortessa,

Becton Dickinson, Franklin Lakes, NJ) and found to be uniformly positive for FcεRIα (Mar-1, FITC-

labeled, #134305) and c-Kit (2B8, APC-labeled, #105811), both from BioLegend (San Diego, CA). Cells

were rested or primed overnight before stimulation.

**siRNA knockdown**

The siRNA sequence for knockdown of mouse c-Cbl was adapted from human siRNA. Cbl-b siRNA

was designed using the Integrated DNA Technologies double-stranded siRNA design tools (IDT,

Skokie, IL). Control double-stranded RNA from IDT was not predicted to be complementary to any

sequence in human, mouse, or rat transcriptomes. 2 M BMDMs in 100 $\mu$L opti-MEM (ThermoFisher,

Waltham, MA) were transfected with 1 $\mu$M siRNA via electroporation at 400 V/10 ms using a BTX

square-wave electroporator. Cells were then plated on 150 mm untreated cell culture dishes and rested

in 10 mL DMEM-10 for 30 min before adding 10 mL DMEM-10 supplemented with 10% CMG-14-12-

cell-conditioned medium as a source of M-CSF. Transfections were pooled from several cuvettes to

obtain enough cells for several stimulation conditions. After 24 h cells were resuspended in enzyme-

free EDTA buffer and replated in untreated 6-well plates (BD Falcon, Sigma-Aldrich) at 1 M cells per

well in unconditioned medium. Cells were used 48 h after transfection.

**Cell stimulation**

BMDM stimulations have been described previously (20). After resting as described above, 1 M live

Jurkat cells, mast cells, or adherent BMDMs were treated at 37°C in DMEM (myeloid cells) or PBS

(Jurkat cells) with 10 µM 3-IB-PP1, a gift from K. Shokat (University of California, San Francisco).

Signaling reactions were quenched by placing on ice and lysing cells in sodium dodecyl sulfate (SDS)

buffer (128 mM Tris base, 10% glycerol, 4% SDS, 50 mM dithiothreitol, pH 6.8). Whole-cell lysates

were prepared for immunoblotting by sonicating with a Bioruptor (Diagenode, Inc., Denville, NJ) for 3

min and boiling for 15 min.

**Antibodies and immunoblotting**

After sample preparation, 0.25 M cell equivalents were run on each lane of a 7% NuPage Tris-Acetate

gel (Invitrogen, Carlsbad, CA) and then transferred to an Immobilon-FL PVDF membrane (EMD

Millipore, Burlington, MA). REVERT Total Protein Stain was used according to the standard protocol to

quantify lane loading. After destaining, membranes were then treated with Odyssey® Blocking Buffer

(TBS) for at least 1 h. Blotting was performed using standard procedures, and blots were imaged on an

Odyssey CLx near-infrared imager (LI-COR Biosciences, Lincoln, NE). Antibodies for immunoblotting

were purchased from Cell Signaling Technology (Danvers, MA), ProMab Biotechnologies (Richmond,

CA), Thermo Fisher Scientific (Waltham, MA), LifeSpan Biosciences (Seattle, WA), Abcam (Cambridge,

UK), and LI-COR Biosciences as follows:

| Antibody | Source and Catalog No. |
|---|---|
| β-Actin (8H10D10) | Cell Signaling 3700 |
| c-Cbl (D4E10) | Cell Signaling 8447 |
| Cbl-b (D3C12) | Cell Signaling 9498 |
| pErk1/2 (phosphoThr$^{202}$/phosphoTyr$^{204}$, D13.14.4E) | Cell Signaling 4370 |
| Erk1/2 (3A7) | Cell Signaling 9107 |
| Fyn | Cell Signaling 4023 |
| Lck (D88) XP(R) | Cell Signaling 2984 |
| LynA (C13F9) | Cell Signaling 2796 |
| LynA + LynB (Lyn-01) | Abcam ab1890 |
| Myc (9B11) | Cell Signaling 2276 |
| pPaxillin (phosphoTyr$^{118}$) | Cell Signaling 2541 |
| pSFK$^{act}$ (phosphoTyr$^{416}$) | Cell Signaling 2101 |
| pZAP70/pSyk (phosphoTyr$^{319(Zap)/352(Syk)}$, 65E4) | Cell Signaling 2717 |
| V5 (D3H8Q) | Cell Signaling 13202 |
| Donkey anti-rabbit IgG 800CW | LI-COR 925-32213 |
| Donkey anti-rabbit IgG 680LT | LI-COR 925-68023 |
| Donkey anti-mouse IgG 800CW | LI-COR 925-32212 |
| Donkey anti-mouse IgG 680LT | LI-COR 925-68022 |

**Quantification, statistics, and image processing**

Immunoblots were analyzed by densitometry from ImageStudio software (LI-COR Biosciences). Images

were background-subtracted, and bands of the appropriate molecular weight were demarcated and

analyzed for each gel lane. Each value was corrected for the total lane protein content (REVERT stain,

above). Data were further normalized to a control condition to show relative changes in protein content.

Where phosphoproteins were quantified, these were corrected for the total amount of the individual

protein, rather than whole-lane protein content. For all figures, n values are biological replicates,

reflecting independent experiments from different batches of cells, stimulus preparations, and

experimental workflows. Statistical analyses were performed using Prism Software (Graphpad, La Jolla,

CA). Significance was assessed using one- or two-way ANOVA analysis with Tukey's or Sidak

correction for multiple comparisons as indicated in the figure legends, with analysis of the mean at each

time point or condition. Error bars in each figure represent standard errors of the mean from at least

three independent experiments. Asterisks reflect specified P-values. Figures were further prepared

using Adobe Creative Cloud software (San Jose, CA). Where appropriate, images were optimized by

applying brightness/contrast changes to the whole image. No gamma correction was used. Images

were only rotated after densitometry analysis was completed.

## References

1.      Guilliams M, Bruhns P, Saeys Y, Hammad H, Lambrecht BN. The function of Fcgamma
receptors in dendritic cells and macrophages. Nat Rev Immunol. 2014;14(2):94-108.

2.      Brown GD. Dectin-1: a signalling non-TLR pattern-recognition receptor. Nat Rev Immunol.
2006;6(1):33-43.

3.      Deng Z, Ma S, Zhou H, Zang A, Fang Y, Li T, et al. Tyrosine phosphatase SHP-2 mediates C-
type lectin receptor-induced activation of the kinase Syk and anti-fungal TH17 responses. Nat Immunol.
2015;16(6):642-52.

4.      Lowell CA. Src-family and Syk kinases in activating and inhibitory pathways in innate immune
cells: signaling cross talk. Cold Spring Harb Perspect Biol. 2011;3(3).

5.      Goodridge HS, Reyes CN, Becker CA, Katsumoto TR, Ma J, Wolf AJ, et al. Activation of the
innate immune receptor Dectin-1 upon formation of a 'phagocytic synapse'. Nature.
2011;472(7344):471-5.

6.      Takai T. Roles of Fc receptors in autoimmunity. Nat Rev Immunol. 2002;2(8):580-92.

7.      Sondermann P. The FcgammaR/IgG Interaction as Target for the Treatment of Autoimmune
Diseases. J Clin Immunol. 2016;36 Suppl 1:95-9.

8.      Chiffoleau E. C-Type Lectin-Like Receptors As Emerging Orchestrators of Sterile Inflammation
Represent Potential Therapeutic Targets. Front Immunol. 2018;9:227.

9.      Freeman SA, Goyette J, Furuya W, Woods EC, Bertozzi CR, Bergmeier W, et al. Integrins Form
an Expanding Diffusional Barrier that Coordinates Phagocytosis. Cell. 2016;164(1-2):128-40.

10.     Bakalar MH, Joffe AM, Schmid EM, Son S, Podolski M, Fletcher DA. Size-Dependent
Segregation Controls Macrophage Phagocytosis of Antibody-Opsonized Targets. Cell.
2018;174(1):131-42 e13.

11.     Jaumouille V, Farkash Y, Jaqaman K, Das R, Lowell CA, Grinstein S. Actin cytoskeleton
reorganization by Syk regulates Fcgamma receptor responsiveness by increasing its lateral mobility
and clustering. Dev Cell. 2014;29(5):534-46.

12.     Abram CL, Lowell CA. The diverse functions of Src family kinases in macrophages. Front
Biosci. 2008;13:4426-50.

13.     Hamerman JA, Lanier LL. Inhibition of immune responses by ITAM-bearing receptors. Sci
STKE. 2006;2006(320):re1.

14.     Hamerman JA, Ni M, Killebrew JR, Chu CL, Lowell CA. The expanding roles of ITAM adapters
FcRgamma and DAP12 in myeloid cells. Immunol Rev. 2009;232(1):42-58.

15.     Lutz-Nicoladoni C, Wolf D, Sopper S. Modulation of Immune Cell Functions by the E3 Ligase
Cbl-b. Front Oncol. 2015;5:58.

16.     Liyasova MS, Ma K, Lipkowitz S. Molecular pathways: cbl proteins in tumorigenesis and
antitumor immunity-opportunities for cancer treatment. Clin Cancer Res. 2015;21(8):1789-94.

17.     Mkaddem SB, Murua A, Flament H, Titeca-Beauport D, Bounaix C, Danelli L, et al. Lyn and Fyn
function as molecular switches that control immunoreceptors to direct homeostasis or inflammation. Nat
Commun. 2017;8(1):246.

18.     Alvarez-Errico D, Yamashita Y, Suzuki R, Odom S, Furumoto Y, Yamashita T, et al. Functional
analysis of Lyn kinase A and B isoforms reveals redundant and distinct roles in Fc epsilon RI-
dependent mast cell activation. J Immunol. 2010;184(9):5000-8.

19.     Scapini P, Pereira S, Zhang H, Lowell CA. Multiple roles of Lyn kinase in myeloid cell signaling
and function. Immunol Rev. 2009;228(1):23-40.

20.     Freedman TS, Tan YX, Skrzypczynska KM, Manz BN, Sjaastad FV, Goodridge HS, et al. LynA
regulates an inflammation-sensitive signaling checkpoint in macrophages. Elife. 2015;4.

21.     Andrews NL, Pfeiffer JR, Martinez AM, Haaland DM, Davis RW, Kawakami T, et al. Small,
mobile FcepsilonRI receptor aggregates are signaling competent. Immunity. 2009;31(3):469-79.

22.     Felce JH, Sezgin E, Wane M, Brouwer H, Dustin ML, Eggeling C, et al. CD45 exclusion- and
cross-linking-based receptor signaling together broaden FcepsilonRI reactivity. Sci Signal.
2018;11(561).

23.     Okuzumi T, Ducker GS, Zhang C, Aizenstein B, Hoffman R, Shokat KM. Synthesis and
evaluation of indazole based analog sensitive Akt inhibitors. Mol Biosyst. 2010;6(8):1389-402.

24.     Okuzumi T, Fiedler D, Zhang C, Gray DC, Aizenstein B, Hoffman R, et al. Inhibitor hijacking of
Akt activation. Nat Chem Biol. 2009;5(7):484-93.

25.     Schoenborn JR, Tan YX, Zhang C, Shokat KM, Weiss A. Feedback circuits monitor and adjust
basal Lck-dependent events in T cell receptor signaling. Sci Signal. 2011;4(190):ra59.

26.     Tan YX, Manz BN, Freedman TS, Zhang C, Shokat KM, Weiss A. Inhibition of the kinase Csk in
thymocytes reveals a requirement for actin remodeling in the initiation of full TCR signaling. Nat
Immunol. 2014;15(2):186-94.

27.     Chow LM, Veillette A. The Src and Csk families of tyrosine protein kinases in hemopoietic cells.
Semin Immunol. 1995;7(4):207-26.

28.     Zhu JW, Doan K, Park J, Chau AH, Zhang H, Lowell CA, et al. Receptor-like tyrosine
phosphatases CD45 and CD148 have distinct functions in chemoattractant-mediated neutrophil
migration and response to S. aureus. Immunity. 2011;35(5):757-69.

29.     Tang R, Langdon WY, Zhang J. Regulation of immune responses by E3 ubiquitin ligase Cbl-b.
Cell Immunol. 2018.

30.    Naramura M, Jang IK, Kole H, Huang F, Haines D, Gu H. c-Cbl and Cbl-b regulate T cell
responsiveness by promoting ligand-induced TCR down-modulation. Nat Immunol. 2002;3(12):1192-9.

31.    Purev E, Neff L, Horne WC, Baron R. c-Cbl and Cbl-b act redundantly to protect osteoclasts
from apoptosis and to displace HDAC6 from beta-tubulin, stabilizing microtubules and podosomes. Mol
Biol Cell. 2009;20(18):4021-30.

32.    Ueno H, Sasaki K, Honda H, Nakamoto T, Yamagata T, Miyagawa K, et al. c-Cbl is tyrosine-
phosphorylated by interleukin-4 and enhances mitogenic and survival signals of interleukin-4 receptor
by linking with the phosphatidylinositol 3'-kinase pathway. Blood. 1998;91(1):46-53.

33.    Hunter S, Burton EA, Wu SC, Anderson SM. Fyn associates with Cbl and phosphorylates
tyrosine 731 in Cbl, a binding site for phosphatidylinositol 3-kinase. J Biol Chem. 1999;274(4):2097-
106.

34.    Yi TL, Bolen JB, Ihle JN. Hematopoietic cells express two forms of lyn kinase differing by 21
amino acids in the amino terminus. Mol Cell Biol. 1991;11(5):2391-8.

35.    Stanley E, Ralph S, McEwen S, Boulet I, Holtzman DA, Lock P, et al. Alternatively spliced
murine lyn mRNAs encode distinct proteins. Mol Cell Biol. 1991;11(7):3399-406.

36.    Huang TH, Huo L, Wang YN, Xia W, Wei Y, Chang SS, et al. Epidermal growth factor receptor
potentiates MCM7-mediated DNA replication through tyrosine phosphorylation of Lyn kinase in human
cancers. Cancer cell. 2013;23(6):796-810.

37.    Radivojac P, Vacic V, Haynes C, Cocklin RR, Mohan A, Heyen JW, et al. Identification,
analysis, and prediction of protein ubiquitination sites. Proteins. 2010;78(2):365-80.

38.    Blom N, Sicheritz-Ponten T, Gupta R, Gammeltoft S, Brunak S. Prediction of post-translational
glycosylation and phosphorylation of proteins from the amino acid sequence. Proteomics.
2004;4(6):1633-49.

39.    Verhagen AM, Wallace ME, Goradia A, Jones SA, Croom HA, Metcalf D, et al. A kinase-dead
allele of Lyn attenuates autoimmune disease normally associated with Lyn deficiency. J Immunol.
2009;182(4):2020-9.

40.    Yokoyama N, Miller WT. Identification of residues involved in v-Src substrate recognition by site-
directed mutagenesis. FEBS Lett. 1999;456(3):403-8.

41.    Xu W, Doshi A, Lei M, Eck MJ, Harrison SC. Crystal structures of c-Src reveal features of its
autoinhibitory mechanism. Mol Cell. 1999;3(5):629-38.

42.    Brown MT, Cooper JA. Regulation, substrates and functions of src. Biochim Biophys Acta.
1996;1287(2-3):121-49.

43.    Straus DB, Weiss A. Genetic evidence for the involvement of the lck tyrosine kinase in signal
transduction through the T cell antigen receptor. Cell. 1992;70(4):585-93.

44.    Burgess KE, Odysseos AD, Zalvan C, Druker BJ, Anderson P, Schlossman SF, et al.
Biochemical identification of a direct physical interaction between the CD4:p56lck and Ti(TcR)/CD3
complexes. Eur J Immunol. 1991;21(7):1663-8.

45.    Samelson LE, Phillips AF, Luong ET, Klausner RD. Association of the fyn protein-tyrosine
kinase with the T-cell antigen receptor. Proc Natl Acad Sci U S A. 1990;87(11):4358-62.

46.     Goldsmith MA, Weiss A. Isolation and characterization of a T-lymphocyte somatic mutant with
altered signal transduction by the antigen receptor. Proc Natl Acad Sci U S A. 1987;84(19):6879-83.

47.     Han J, Zhang G, Welch EJ, Liang Y, Fu J, Vogel SM, et al. A critical role for Lyn kinase in
strengthening endothelial integrity and barrier function. Blood. 2013;122(25):4140-9.

48.     Yamamoto N, Takeshita K, Shichijo M, Kokubo T, Sato M, Nakashima K, et al. The orally
available spleen tyrosine kinase inhibitor 2-[7-(3,4-dimethoxyphenyl)-imidazo[1,2-c]pyrimidin-5-
ylamino]nicotinamide dihydrochloride (BAY 61-3606) blocks antigen-induced airway inflammation in
rodents. The Journal of pharmacology and experimental therapeutics. 2003;306(3):1174-81.

49.     Buckbinder L, Crawford DT, Qi H, Ke HZ, Olson LM, Long KR, et al. Proline-rich tyrosine kinase
2 regulates osteoprogenitor cells and bone formation, and offers an anabolic treatment approach for
osteoporosis. Proc Natl Acad Sci U S A. 2007;104(25):10619-24.

50.     Hanke JH, Gardner JP, Dow RL, Changelian PS, Brissette WH, Weringer EJ, et al. Discovery of
a novel, potent, and Src family-selective tyrosine kinase inhibitor. Study of Lck- and FynT-dependent T
cell activation. J Biol Chem. 1996;271(2):695-701.

51.     Heng TS, Painter MW, Immunological Genome Project C. The Immunological Genome Project:
networks of gene expression in immune cells. Nat Immunol. 2008;9(10):1091-4.

52.     Dwyer DF, Barrett NA, Austen KF, Immunological Genome Project C. Expression profiling of
constitutive mast cells reveals a unique identity within the immune system. Nat Immunol.
2016;17(7):878-87.

53.     Gustin SE, Thien CB, Langdon WY. Cbl-b is a negative regulator of inflammatory cytokines
produced by IgE-activated mast cells. J Immunol. 2006;177(9):5980-9.

54.     Shelby SA, Veatch SL, Holowka DA, Baird BA. Functional nanoscale coupling of Lyn kinase
with IgE-FcepsilonRI is restricted by the actin cytoskeleton in early antigen-stimulated signaling. Mol
Biol Cell. 2016;27(22):3645-58.

55.     Kalesnikoff J, Galli SJ. Antiinflammatory and immunosuppressive functions of mast cells.
Methods Mol Biol. 2011;677:207-20.

56.     Duan L, Reddi AL, Ghosh A, Dimri M, Band H. The Cbl family and other ubiquitin ligases:
destructive forces in control of antigen receptor signaling. Immunity. 2004;21(1):7-17.

57.     Zhu LL, Luo TM, Xu X, Guo YH, Zhao XQ, Wang TT, et al. E3 ubiquitin ligase Cbl-b negatively
regulates C-type lectin receptor-mediated antifungal innate immunity. J Exp Med. 2016;213(8):1555-70.

58.     Thien CB, Langdon WY. c-Cbl and Cbl-b ubiquitin ligases: substrate diversity and the negative
regulation of signalling responses. Biochem J. 2005;391(Pt 2):153-66.

59.     Mohapatra B, Ahmad G, Nadeau S, Zutshi N, An W, Scheffe S, et al. Protein tyrosine kinase
regulation by ubiquitination: critical roles of Cbl-family ubiquitin ligases. Biochim Biophys Acta.
2013;1833(1):122-39.

60.     Kyo S, Sada K, Qu X, Maeno K, Miah SMS, Kawauchi-Kamata K, et al. Negative regulation of
Lyn protein-tyrosine kinase by c-Cbl ubiquitin-protein ligase in FceRI-mediated mast cell activation.
Genes to Cells. 2003;8(10):825-36.

61.     Shao Y, Yang C, Elly C, Liu YC. Differential regulation of the B cell receptor-mediated signaling
by the E3 ubiquitin ligase Cbl. J Biol Chem. 2004;279(42):43646-53.

62.     Sanjay A, Houghton A, Neff L, DiDomenico E, Bardelay C, Antoine E, et al. Cbl associates with
Pyk2 and Src to regulate Src kinase activity, alpha(v)beta(3) integrin-mediated signaling, cell adhesion,
and osteoclast motility. J Cell Biol. 2001;152(1):181-95.

63.     Sanjay A, Miyazaki T, Itzstein C, Purev E, Horne WC, Baron R. Identification and functional
characterization of an Src homology domain 3 domain-binding site on Cbl. FEBS J. 2006;273(23):5442-
56.

64.     Palacios-Moreno J, Foltz L, Guo A, Stokes MP, Kuehn ED, George L, et al. Neuroblastoma
tyrosine kinase signaling networks involve FYN and LYN in endosomes and lipid rafts. PLoS Comput
Biol. 2015;11(4):e1004130.

65.     Johnson TM, Williamson NA, Scholz G, Jaworowski A, Wettenhall RE, Dunn AR, et al.
Modulation of the catalytic activity of the Src family tyrosine kinase Hck by autophosphorylation at a
novel site in the unique domain. J Biol Chem. 2000;275(43):33353-64.

66.     Teixeira J, Fuentes H, Bielskutė S, Gairi M, Żerko S, Koźmiński W, et al. The Two Isoforms of
Lyn Display Different Intramolecular Fuzzy Complexes with the SH3 Domain. Molecules. 2018;23(11).

67.     Tornillo G, Knowlson C, Kendrick H, Cooke J, Mirza H, Aurrekoetxea-Rodriguez I, et al. Dual
Mechanisms of LYN Kinase Dysregulation Drive Aggressive Behavior in Breast Cancer Cells. Cell Rep.
2018;25(13):3674-92 e10.

68.     Wynn TA, Chawla A, Pollard JW. Macrophage biology in development, homeostasis and
disease. Nature. 2013;496(7446):445-55.

69.     Galli SJ, Borregaard N, Wynn TA. Phenotypic and functional plasticity of cells of innate
immunity: macrophages, mast cells and neutrophils. Nat Immunol. 2011;12(11):1035-44.

70.     Bosurgi L, Cao YG, Cabeza-Cabrerizo M, Tucci A, Hughes LD, Kong Y, et al. Macrophage
function in tissue repair and remodeling requires IL-4 or IL-13 with apoptotic cells. Science.
2017;356(6342):1072-6.

71.     Gentek R, Ghigo C, Hoeffel G, Bulle MJ, Msallam R, Gautier G, et al. Hemogenic Endothelial
Fate Mapping Reveals Dual Developmental Origin of Mast Cells. Immunity. 2018;48(6):1160-71 e5.

72.     Rafiq K, Kolpakov MA, Seqqat R, Guo J, Guo X, Qi Z, et al. c-Cbl inhibition improves cardiac
function and survival in response to myocardial ischemia. Circulation. 2014;129(20):2031-43.

73.     Weiss A, Stobo JD. Requirement for the coexpression of T3 and the T cell antigen receptor on a
malignant human T cell line. J Exp Med. 1984;160(5):1284-99.

74.     Phee H, Abraham RT, Weiss A. Dynamic recruitment of PAK1 to the immunological synapse is
mediated by PIX independently of SLP-76 and Vav1. Nat Immunol. 2005;6(6):608-17.

75.     Myers MD, Sosinowski T, Dragone LL, White C, Band H, Gu H, et al. Src-like adaptor protein
regulates TCR expression on thymocytes by linking the ubiquitin ligase c-Cbl to the TCR complex. Nat
Immunol. 2006;7(1):57-66.

76.     Zhu JW, Brdicka T, Katsumoto TR, Lin J, Weiss A. Structurally distinct phosphatases CD45 and
CD148 both regulate B cell and macrophage immunoreceptor signaling. Immunity. 2008;28(2):183-96.

77.     Takeshita S, Kaji K, Kudo A. Identification and characterization of the new osteoclast progenitor
with macrophage phenotypes being able to differentiate into mature osteoclasts. Journal of bone and
mineral research : the official journal of the American Society for Bone and Mineral Research.
2000;15(8):1477-88.

## Acknowledgements

We thank Arthur Weiss, Kevan Shokat, Yoji Shimizu, Brandon Burbach, Erik Peterson, Juan Rivera, and Ryo Suzuki for reagents and discussion and Terri Kadlecek, Marianne Mollenauer, and Whitney Swanson for technical support. Many thanks also to Hai-Bin Ruan, John Connett, Marc Jenkins, Bryce Binstadt, Erik Peterson, Nicholas Levinson, Aditi Bapat, Frances Sjaastad, and Emily Ewan for valuable discussions and feedback.

## Funding

This work was supported by NIH grants R01AR073966 (TSF), R03AI130978 (TSF) and R01CA215052 (KLS) and the following University of Minnesota awards (all TSF): Masonic Cancer Center/American Cancer Society IRG-58-001-55; Grant-in-Aid of Research, Artistry, and Scholarship #92286; University of Minnesota Foundation Research and Equipment Award NF-0315-02; and Center for Autoimmune Diseases Research Pilot Grant 2017. Training support was provided by NIH award T32DA007097 (BFB).

## Author Contributions

Conceptualization, Methodology, and Data Curation: BFB, TSF

Investigation, Validation, and Visualization: BFB, MGN, TSF

Formal Analysis, Writing- Review and Editing: BFB, MGN, KLS, TSF

Writing- Original Draft: BFB, TSF

Funding Acquisition: BFB, KLS, TSF

690    Project Administration, Resources, Validation, and Software: TSF

691    Supervision: KLS, TSF

## Competing Interests

693    The authors state that they have no competing interests.

## Data and Materials Availability

695    Reagents and protocols will be provided upon request.

## Figure Supplements

697    **Figure 1–figure supplement 1. Loss of positive signaling in c-Cbl-deficient cells prevents**

698    **assessment of LynA contributions to Erk phosphorylation.**

699    **Figure 3–figure supplement 1. LynA$^{Y32A}$ and LynB are expressed more highly in Jurkat cells than**

700    **wild-type LynA.**

## Source Data

Figure 1–Source Data 1

Figure 3–Source Data 1

Figure 3–figure supplement 1-Source Data 1

Figure 4–Source Data 1

Figure 5–Source Data 1

Figure 6–Source Data 1