1 **Thrombin cleaves and activates the protease-activated receptor 2**
2 **dependent on thrombomodulin co-receptor availability**
3

4 Dorothea M. Heuberger,*[+] Alessandro G. Franchini,* Jerzy Madon,* Reto A. Schuepbach*

6 *Institute of Intensive Care Medicine, University Hospital Zurich, University of Zurich, Zurich,
7 Switzerland
8 [+]Surgical Research Division, University Hospital Zurich, University of Zurich, Zurich, Switzerland
9
10

Corresponding author:
Reto A. Schuepbach, MD, MSc
University Hospital Zurich, University of Zurich
Institute of Intensive Care Medicine
OPS B 17/Lab E 26
Raemistrasse 100
CH-8091 Zurich
Tel.: +41 44 255 1092
Email: reto.schuepbach@usz.ch
20

21 Short title: TM-bound thrombin cleaves PAR2

Abstract

Introduction

Protease-activated receptors (PARs) evolved to react to extracellular proteolytic activity. In mammals, three of the four PARs (PAR1, PAR3, and PAR4) that are expressed respond to the prototypical procoagulant enzyme thrombin, whereas PAR2 was assumed to resist activation by thrombin. To date, involvement of cell surface thrombin-recruiting co-receptors such as thrombomodulin (TM), which potentially facilitates PAR2 cleavage, has not been addressed. Thus, we examined whether TM-bound thrombin cleaved PAR2 and tested biological responses such as nuclear factor kappa B (NF-κB) DNA binding activity and cytokine release.

Materials and Methods

We examined 293T cells overexpressing PAR2 and TM for thrombin recruitment by TM promoting PAR2 cleavage. To test for the TM–thrombin interactions required for PAR2 cleavage and to map cleavage sites on PAR2, mutant constructs of TM or PAR2 were engineered. Biological effects because of PAR2 activation were investigated using an NF-κB reporter system and cytokine release.

Results and Conclusions

We identified that, at low to moderate concentrations, thrombin cleaved PAR2 in a TM co-receptor-dependent manner with cleavage efficiency comparable to that of trypsin. In TM's presence, thrombin efficiently cleaved both, PAR1 and PAR2, albeit kinetics differed. Whereas the majority of surface expressed PAR1 was immediately cleaved off, prolonged exposure to thrombin resulted in few additional cleavage. In contrast, PAR2 cleavage was sustained upon prolonged exposure to thrombin. However, TM EGF-like domain 5 was required and TM chondroitin sulfate (CS) proteoglycan sites serine 490 and serine 492 assisted in PAR2 cleavage, while thrombin preferentially cleaved at arginine 36 on PAR2's N-terminus. Note that thrombin-induced activation of NF-κB via PAR2 resulted in release of interleukin-8. Thus, we provide a novel concept of how thrombin efficiently cleaves PAR2 in a TM-dependent manner, resulting in pro-inflammatory interleukin-8 release. This unexpected pro-inflammatory role of TM, promoting cleavage and activation of PAR2

55 by thrombin, may lead to novel therapeutic options for treating inflammatory and
56 malignant diseases.

58 Keywords: Receptor; PAR 2; thrombin; thrombomodulin; trypsin; NF-κB, IL-8

Abbreviations

61 PAR: protease-activated receptor
62 TM: thrombomodulin
63 CS: chondroitin sulfate
64 EGF-like domain: epidermal growth factor-like domain
65

Introduction

Clotting proteases elicit multiple physiological and pathophysiological responses
beyond clot formation. The family of PARs has evolved to allow cells to react to
presence of clotting (serine) proteases [1]. PARs belong to the seven-
transmembrane G-protein-coupled receptor family. In mammals, the family of PARs
consists of four highly related serine protease receptors, i.e., PAR1–4 [2-7].

Previously, PAR2 has been linked to inflammation-driven diseases of the brain,
cardiovascular system, lungs, joints, and gastrointestinal tract [8-16], as well as
cancer growth [17, 18], migration, and metastasis [9, 19]. Despite its importance in
pathophysiological processes, no treatments based on interference with PAR2 have
been established [20]. There are potential obstacles for defining therapeutic targets
because of the existence of multiple activation mechanisms and a large variety of
causal activating proteases [21-23], as well as the complex nature of PAR2
compartmentalization and co-localization [19]. Recent studies that emphasize the
importance of PAR hetero-dimerization [24], requirement of PAR2 co-localized co-
receptors [22], and introduction of the concept of biased PAR agonists [25] further
demonstrate the complexity of PAR2-related functions under physiological or
pathophysiological conditions.

To date, PAR2 has been assumed to resist activation by the prototypical
procoagulant protease thrombin, as it has not been observed to activate
overexpressed PAR2 in the absence of co-receptors [1, 5, 26]. Moreover, only

extremely high, potentially supra-physiological concentrations of thrombin (up to
500 nM) cleave peptides that are homologous to PAR2's N-terminus and activate
PAR2 [27].

Unlike previous observations of PAR2 resisting thrombin activation, *in vitro* and *in*
vivo models have demonstrated multiple effects involving both, PAR2 and thrombin,
such as cancer cell invasion [18], signal transduction [28, 29], and late sepsis. To
explain these observations, PAR1 and PAR2 transactivation is the leading model;
the thrombin-generated N-terminus of PAR1 signaling via a PAR2 hetero-dimer
partner [24, 30].

To date, the activation of PAR2 by cofactor (or co-receptor)-bound thrombin has
not been systematically evaluated, although co-receptor-dependent mechanisms,
such as tissue factor-bound clotting factor VIIa activating PAR2 [22], endothelial
protein C receptor-bound activated protein C (aPC) cleaving PAR1 and PAR3 [31],
endothelial protein C receptor-bound clotting factor Xa cleaving PAR1 [32], and
activating PAR1 or PAR3 [33-35], are well known. Moreover, in mice, PAR3 recruits
thrombin which facilitates cleavage of PAR4 [36].

For thrombin-mediated activation of protein C, the cell surface glycoprotein TM
(also known as fetomodulin or CD141) serves as a cofactor [37,38]. TM is primarily
expressed on all vascular endothelial cells [39], keratinocytes [40], astrocytes [41],
monocytes and macrophages [42], and lung alveolar epithelial cells [43].

Because TM serves as the key re-director of procoagulant thrombin [44], which
limits clot formation and injury to the vascular bed [45], TM might play a role in re-
directing PAR activation by thrombin in situations of ongoing thrombin generation
such as chronic inflammation. Note that TM-bound thrombin changes substrate
specificity of thrombin to anticoagulant and anti-inflammatory signaling via protein C
activation. TM expression is linked to regulation of homeostatic and inflammatory
responses during injuries [46]. For acute injuries of the endothelium, TM
downregulation or shedding occurs; accordingly, soluble TM (sTM) serves as a
marker for acute inflammation [47]. Furthermore, compared with the endothelium,
less is known about the TM co-receptor's function in other tissues.

Therefore, we tested whether TM serves as a co-receptor for the thrombin-
dependent cleavage of PAR2. Moreover, we verified whether TM-bound thrombin
activation of PAR2 resulted in pro- or anti-inflammatory effects.

Materials and Methods

Reagents

Thrombin [EC 3.4.21.5] was purchased from Haematologic Technologies (Essex Junction, VT, USA) and trypsin [EC 3.4.21.4] from Gibco (Thermo Fisher Scientific, Reinach, Switzerland). Recombinant aPC (Xigris) was purchased from Eli Lilly and Company (Indianapolis, IN, USA). Peptides corresponding to the N-terminus of R41-cleaved PAR1 (TFLLRNPNDK), R36-cleaved PAR2 (SLIGRL), and PAR4 (AYPGKF) were custom-made (Synpeptide Co., Ltd., Shanghai, China; and ProteoGenix, Schiltigheim, France).

Recombinant sTM (SRP3172), Nordihydroguaiaretic acid (NDGA), TNFα and PI3K inhibitor LY294002 were purchased from Sigma (Buchs SG, Switzerland). We used lepirudin (Refludan; Schering, Berlin, Germany) to block thrombin. In fact, when added without thrombin, lepirudin had no effect in any of the reported assays and readouts. Heparin was purchased from Bichsel (Interlaken, Switzerland). Thrombin exosite I blocking aptamer (5′-GGTTGGTGTGGTTGG-3′ [48]) and exosite II blocking aptamer (5′-AGTCCGTGGTAGGGCAGGTTGGGGTGACT-3′ [49]) were custom-made (Microsynth, Balgach, Switzerland). Monoclonal anti-PAR1 ATAP2 was used as described previously [32, 50] and anti-PAR2 (SAM11; Thermo Fisher Scientific), anti-PAR2_#344222 (MAB3949; R&D Systems, Inc., MN, USA), anti-TM_#1009 [141C01(1009), Thermo Fisher Scientific], anti-TM_#2375 (American Diagnostics, Pfungstadt, Germany), anti-TM_#501733 (MAB3947; R&D Systems), anti-His_#AD1.1.10 (MAB050; R&D Systems), and goat anti-mouse IgG Alexa-594 (R37121; Thermo Fisher Scientific) were all used according to manufacturer's instructions unless stated otherwise. All oligonucleotides were purchased from Microsynth. For quantifying luciferase activity, the Bright-Glo™ Luciferase Assay System (Promega, Dübendorf, Switzerland) was used, in accordance to the manufacturer's instructions.

Peptide cleavage

A high-binding microplate (#9018; Corning, Tewksbury, MA, USA) was coated (o/n) with anti-His#AD1.1.10 (2 µg/ml) and anti-TM_#501733 (1 µg/ml). Then, the plate was washed and incubated with buffer or recombinant sTM (60 nM) to bind to anti-

TM for 30 minutes. Synthetic PAR2 peptide (10 µg/ml) was bound to the plate by the

N-terminal His-tag (1 h). We added pre-warmed thrombin (30 nM) to the washed

plate and incubated the plate for 1 h at 37°C. The uncleaved (full length) PAR2

peptide was detected via the C-terminal biotin-tag using streptavidin-HRP and

quantified by a peroxidase-based enzyme immunoassay.

Cell culture, plasmid transfection, and gene silencing

Epithelial A549 cells and human embryonic kidney cell-derived 293T cells (ATCC;

LGC Standards GmbH, Wesel, Germany) were cultivated and propagated as

described previously [32, 50]. In brief, cells were propagated in DMEM (Gibco)

containing 10% fetal calf serum (GE Healthcare, Glattbrugg, Switzerland), detached

using dissociation buffer (Gibco), transfected, seeded, and grown to confluence in

multi-well dishes. Two days later, experiments were performed using washed cells

and serum-free conditioning medium (DMEM) containing 0.04% BSA (Sigma).

Tagged and untagged PAR1, PAR2, TM constructs, and pGL4.32[*luc2P*/NF-κB-

RE/Hygro] (Promega) were transiently expressed in 293T cells by transfection with

Lipofectamine2000 (Thermo Fisher Scientific), as described previously [34]. Cells

used for cleavage assays were transfected with 1.6 µg/ml plasmid DNA; however, for

NF-κB luciferase and IL-8 enzyme-linked immunosorbent assay (ELISA) assays,

1.0 µg/ml plasmid DNA was used for transfection of the 293T cells. To reach

comparable amounts of PAR2 expression, in those cells that overexpressed several

proteins (such as with and without TM), the empty pcDNA3.1/Zeo(+) plasmid

(Thermo Fisher Scientific) was used as a mock plasmid. We performed gene

silencing using Lipofectamine RNAiMAX (Thermo Fisher Scientific) as described

previously [32], using 20 nM silencing RNA (siRNA). GGAACCAAUAGAUCCUCUAtt

(si PAR2) to silence PAR2 and AGGUAGUGUAAUCGCCUUGtt was used as a

control silencing RNA (si control). Furthermore, all oligonucleotides were synthetized

by Microsynth.

Overexpression constructs

The overexpression constructs were embedded within the pcDNA3.1/Zeo(+)

(Thermo Fisher Scientific) expression plasmid, as described previously [34]. TM was

cloned from cDNA (obtained from RNA isolated from EA.hy926 endothelial cells)

using primers containing restriction sites (italics) HindIII and XhoI (forward: 5′-

GCTA*AAGCTT*GACAGGAGAGGCTGTCGCCATC-3′   and   reverse: 5′-

GCTA*CTCGAG*GGGAATAAGTGGGGCTTGCT-3′). The HindIII/XhoI-digested PCR

product was directly introduced into the HindIII/XhoI-cut pcDNA3.1/Zeo(+) backbone.

For removal of N-terminal domains in TM, two unique BamHI restriction sites were

introduced by mutagenesis (NEB's plasmid mutation kit; Ipswich, MA, USA): the first

3′ from the signal sequence and the second 3′ from the sequence that encoded the

domain to be removed. Digestion of the mutated TM expression vector with BamHI

followed by plasmid re-ligation allowed dropping domains while maintaining the TM

native signal sequence (details are provided in Supplementary Table S1). Note that

TM chondroitin sulfate (CS) sites serine 490 and 492 were mutated to alanine with

the primer set (forward: 5′-CACGGCTCGACCTCAATG-3′, reverse: 5′-

GGGGCTCGCC*AGC*GCC*GGC*GTCGCCACCGTCC-3′).

PAR1 and PAR2 alkaline phosphatase-tagged reporter ("AP-PAR1 and AP-

PAR2") was as described elsewhere [34]. Sequence of the PAR4 alkaline

phosphatase-tagged reporter ("AP-PAR4") is described in Supplementary Table S4.

In brief, an alkaline phosphatase (AP) tag (pSEAP; Clontech, CA, USA) was linked

to the N-terminus of PAR1 cDNA (F2R, NM_001992.4), PAR2 cDNA (F2RL1,

NM_005242.5), and PAR4 cDNA (F2RL3, NM_003950.3). Unlike the PAR2

construct mentioned elsewhere [34], an arginine (italics) at the C-terminal end of AP

(potential thrombin cleavage site) was removed using phosphorylated mutagenic

primers (forward: 5′-CCCGGGTTACTCT*GCG*GCCCAAGGAACCAATAG-3′; reverse:

5′-CTATTGGTTCCTTGGGC*CGC*AGAGTAACCCGGG-3′; Supplementary Table S2).

We inserted a green fluorescent C-terminal tag (EGFP; Clontech) analogous to a

procedure used in a previous study [34] (for details, see Supplementary Table S2

and Supplementary Scheme S1). To test for preferred cleavage sites on PAR2's N-

terminal, arginine 36 was replaced by alanine ("PAR2 R36A;" protein sequence in

Supplementary Table S3) using mutagenic primers (forward:

5′-CCAATAGATCCTCTAAAGGA*GCA*AGCCTTATTGGTAAGG-3′; reverse:

5′-TTCCTTGGGCCGCAGAGTAACCCGGG-3′). Next, all potential thrombin

cleavage sites (arginine and lysine) were replaced by alanine ("PAR2 all-to-A"

mutant; protein sequence provided in Supplementary Table S3). Note that all

constructs that were used were verified by sequencing.

PAR cleavage reporter assay

The cleavage reporter assay was performed on 293T cells that transiently
expressing AP-PAR constructs (Supplementary Table S2), as described elsewhere
[32]. In brief, washed cells were incubated with an agonist (for 20 minutes, or if
indicated, for repeated periods of 20 minutes), supernatants were removed and
filtered (cellulose ester filter; pore size, 0.45 µm), and AP activity was quantified (1-
Step PNPP; Thermo Fisher Scientific) by spectrophotometry (Labsystems Multiskan
MCC/340; Fisher Scientific). For all experiments, AP-PAR expression levels were
confirmed to be comparable among constructs by quantifying AP activity of cell
surface. Data are presented in optical density (OD) after subtraction of signals
obtained from cells not incubated with the agonist.
232

Immunoassays
IL-8 was quantified by a commercial (R&D Systems) sandwich ELISA, according to
the manufacturer's instructions. In brief, 293T cells were transfected with plasmid
DNA and A549 cells that were silenced with siRNA 48 h prior to incubation with
agonists for 24 and 6 h, respectively. Then, IL-8 release was measured in
supernatants. As described previously, cell surface PAR2 and TM were quantified by
cell surface ELISA [50].
240

Immunofluorescence microscopy
EGFP-tagged protein (Supplementary Scheme S1) was overexpressed using the
same protocols as that described for PAR cleavage reporter constructs and
visualized using appropriate filter sets of an Axiovert-10 fluorescent microscope (Carl
Zeiss AG, Feldbach, Switzerland). As described for the cell surface ELISA, untagged
proteins were incubated with antibodies; however, the secondary antibody carried an
Alexa-594 fluorescent tag (Thermo Fisher Scientific) rather than HRP.
248

NF-κB luciferase assay
293T cells were transfected with the NF-κB firefly luciferase construct
pGL4.32[*luc2P*/NF-κB-RE/Hygro] (Promega) along with AP-PAR constructs and
seeded into black, clear-bottomed 96-well plates (Greiner Bio-One, Kremsmünster,
Austria). After transfection for 48 h, cells were washed and incubated with an agonist
(for 6 h) at 37°C. Bright-Glo™ Luciferase Assay System reagent was added to cells

and luciferase activity was detected via luminescence (SpectraMax i3; Molecular

Devices, San Jose, CA, USA), according to the manufacturer's instructions.

Furthermore luciferase activity was normalized to the buffer-induced relative light

units (RLUs).

Statistics

We analyzed and presented the data using GraphPad Prism5 (GraphPad Software,

La Jolla, CA, USA). To calculate the indicated P-values, a two-sample, two-tailed

homoscedastic t-test and one-way or two-way ANOVA with Bonferroni correction

were used.

Results

Effect of PAR2 activation by thrombin on IL-8 release in native A549 cells

Various studies showed expression of all PARs in lung epithelial cells promoting IL-8

release upon PAR agonist peptide stimulation due to PAR cis- or trans-activation

[51] [52]. In this study, we first tested the concentrations of PAR agonist peptides

required to induce IL-8 secretion in A549 cells. Similar to a previous study [53], we

found that A549 cells were induced to release IL-8 on incubation with the PAR2

agonist peptide SLIGRL, whereas the PAR1 agonist peptide TFLLRNPNDK and the

specific PAR4 agonist peptide AYPGKF demonstrated no such effect, even at high

concentrations. Unlike the PAR1-specific peptide, thrombin significantly induced IL-8

release from 30 to 100 nM (Figure 1A). To test whether thrombin induced IL-8

release could be linked to PAR2 activation, we relied on the PAR2-specific inhibitor

NDGA. NDGA did neither affect IL-8 release at baseline (buffer control) nor in highly

TNFα-induced A549 cells (not shown). However, NDGA significantly reduced IL-8

release by thrombin as well as by the PAR2-specific agonist SLIGRL (Figure 1B).

Consistently, silencing of PAR2 by siRNA abolished responses of A549 cells to

thrombin or SLIGRL induction (Figure 1C). In summary, IL-8 release by thrombin

stimulation was significantly reduced in PAR2-silenced or inhibited lung epithelial

cells, suggesting that thrombin plays a role in PAR2 signaling.

Figure 1: Increased IL-8 release upon thrombin and SLIGRL stimulation in a PAR2-dependent manner. (**A**) A549 cells were stimulated with thrombin, PAR2 agonist peptide SLIGRL, PAR1 agonist peptide TFLLRNPNDK, and PAR4 agonist peptide AYPGKF for 6 h. Then, IL-8 release was quantified by a peroxidase-based enzyme immunoassay. (**B**) A549 were pretreated with DMSO or NDGA (20 µM) for 30 min before stimulation with buffer, thrombin or SLIGRL (concentrations indicated in the graph). IL-8 release in the supernatant was quantified by ELISA after 6 h of agonist incubation. (**C**) A549 cells were transfected with siRNA targeting no specific mRNA (si control) or siRNA targeting PAR2 (si PAR2). Then, cells were stimulated with buffer, thrombin (30 nM), or SLIGRL (50 µM) for 6 h before IL-8 release was measured in cell supernatants. Data was presented as mean ± SEM; three independent experiments were each performed in triplicate; ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ using one-way ANOVA (A) and student's t test (B,C).

Role of TM on PAR2 cleavage by thrombin

To test whether IL-8 induction by thrombin in epithelial cells of the lung could be caused by direct cleavage of PAR2 using a co-receptor, we established a purified

system comprising immobilized PAR2 peptide with immobilized sTM and thrombin.
In this purified system, thrombin efficiently cleaved the PAR2 peptide in immobilized
sTM's presence; however, consistent with the established conclusion, thrombin
failed to directly cleave PAR2 in the co-receptor's absence (Figure 2A). To test on
human cells whether physiologically achievable concentrations of thrombin cleaved
PAR2 and whether co-receptor(s) could be involved, we relied on the previously
established PAR cleavage reporter system [32, 34, 54]. To allow verification of
construct expression by assessing the tag's cell surface AP activity and monitor
cleavage by quantifying the activity of AP released into the supernatant, we
overexpressed PAR2 containing an AP enzyme domain at the N-terminus
(Supplementary Table S2; Supplementary Scheme S1, [34]). PAR2 reporter and TM
constructs were expressed at the cell surface, which was further assessed by
fluorescence microscopy and cell surface ELISA. Co-expression of PAR2 and TM
did not affect expression of both constructs (Supplementary Fig. S1a and S1b). In
such an overexpression system, free thrombin at concentrations up to 30 nM failed
to cleave PAR2 (Figure 2B), unlike the results of a recent study [27]. However, in
TM's presence, cleavage efficiency was significantly enhanced and, at low
nanomolar concentrations of proteases, thrombin and trypsin showed comparable
cleavage efficiency (Figure 2C). To test whether thrombin/TM could mediate release
or activation of another protease, which ultimately cleaved PAR2, we added
thrombin-induced supernatant from TM-expressing cells to 1) PAR2- and 2) PAR2-
and TM-expressing cells, for assessing PAR2 cleavage. We identified that, in the
absence of TM, PAR2 was not cleaved and, wherever thrombin in supernatant was
blocked by the specific inhibitor lepirudin, cleavage was blunted (Figure 2D). Addition
of activated protein C or inhibition of metalloproteases did not affect PAR2 cleavage
by thrombin (Supplementary Fig. S3a and S3b). Therefore, overall, these results
cannot support a model whereby thrombin induces a PAR2-cleaving protease. To
simulate more physiological conditions where PAR1, PAR2 and TM are co-
expressed, and to test whether the co-expression impacts cleavage, we
overexpressed 1) PAR1 containing an N-terminal AP enzyme tag together with
untagged-PAR2 or 2) PAR2 containing an N-terminal AP enzyme tag together with
an untagged-PAR1 (Supplementary Table S2; Supplementary Scheme S1, [34]) in
presence or absence of TM. As expected, PAR1 was rapidly cleaved by thrombin (5
and 30 nM) while PAR2 - in TM's presence - was more efficiently cleaved upon

prolonged thrombin exposure (5 and 30 nM) (Figure 2E and 2F). The cleavage of

PAR2 at 60 minutes of exposure to thrombin (30 nM) was comparably efficient to the

PAR1 cleavage at 20 minutes (Figure 2F). Cleavage of PAR2 was sustained upon

prolonged exposure to thrombin for up to 25 hours (Supplementary Fig. S4a-c). Note

that, upon a exposure to thrombin for 3 hours, a broad range of thrombin

concentrations from 1 to 30 nM resulted in significant cleavage of PAR2

(Supplementary Fig. S4a-c). Our observations suggest that TM-bound thrombin

directly cleaves PAR2 in our overexpression system and that TM dependent

cleavage of PAR2 by thrombin is enhanced over time. Thus, in summary, our data

suggest that thrombin at moderate concentrations (5 and 30 nM) efficiently cleaves

overexpressed PAR2 reporter construct in TM's presence.

Figure 2: Thrombin efficiently cleaves overexpressed PAR2 in the presence of
TM. (**A**) A PAR2-derived synthetic peptide mimicking the entire extracellular N-
terminal domain of PAR2 was bound to the plate via a C-terminal 6xHis-tag and
incubated with thrombin in the presence or absence of immobilized sTM. Uncleaved
PAR2 was detected via ELISA peroxidase-based quantification of unremoved N-

terminal biotin tags. Stars indicate comparisons of * PAR2 peptide cleavage of
sTM+thrombin vs. thrombin without sTM or (*) sTM+thrombin vs. sTM without
thrombin. (**B**) 293T cells transiently overexpressing AP-PAR2 reporter construct
either alone or with TM. Where indicated, cells were incubated with thrombin (30 nM)
for 20 minutes before the released AP's activity was measured in the cells'
supernatants, serving as a surrogate for PAR2 cleavage. (**C**) Cells expressing TM
and AP-PAR2 reporter construct were incubated with agonists as indicated and
assayed as described in (B). Stars indicate the comparison of thrombin to buffer in
cells overexpressing AP-PAR2+TM. (**D**) 293T cells expressing only TM were
incubated with thrombin (20 min, 30 nM). Thereafter, the supernatant was swapped
to AP-PAR2- or AP-PAR2 and TM-expressing cells and the cells were incubated for
another 20 minutes. Where indicated, the supernatant was pre-incubated with the
thrombin inhibitor lepirudin (30 nM) before it was added to the cells expressing AP-
PAR2 or AP-PAR2 and TM. (**E**) and (**F**) 293T cells expressing AP-PAR1 together with
non-tagged PAR2 or AP-PAR2 together with non-tagged PAR1 with or without TM
were incubated with (**E**) thrombin (5 nM) or (**F**) thrombin (30 nM) for 20 minutes. AP
activity was measured in the supernatant and the cells were agonist-incubated for
repeated periods of 20 minutes. Stars indicate the comparison of thrombin to buffer
in cells overexpressing AP-PAR2+PAR1+TM. Data are presented as mean ± SEM;
(A-D) three or (E,F) four independent experiments, each performed in triplicate; ns p
> 0.05, * p < 0.05, ** p < 0.01, *** p < 0.001, using student's t test.
380
381

Importance of interactions between TM EGF-like domains and CS with
thrombin exosite I and exosite II on PAR2 cleavage

Next, we addressed domains of TM required for increasing PAR2 cleavage by

thrombin. TM contains six epidermal growth factor (EGF)-like domains, from which

EGF-like domain 4 was linked to protein C substrate acceptance, whereas EGF-like

domains 5 and 6 were linked to thrombin recruitment [55] to the cell surface. To test

whether these EGF-like domains were involved in promoting PAR2 cleavage, we

produced N-terminally truncated mutants of TM (Supplementary Table S1). The

mutants were expressed and recognized as expected by cell surface ELISA using

anti-TM targeting EGF-like domains 5 and 6 (Supplementary Fig. S5). Note that

PAR2 was efficiently cleaved when wild-type TM or mutants containing EGF-like

domains 5 and 6 were present. The TM mutant composed of only EGF-like domain 6

failed to promote the cleavage of PAR2 by thrombin (Figure 3A).

Moreover, interactions between thrombin exosite II and CS on TM were involved

for thrombin recruitment [56] and to enhance the endothelial protein C pathway. To

test whether such interactions between exosite II and CS facilitated PAR2 cleavage

by thrombin, we synthesized a TM mutant that lacked the serine 490 and 492

glycosylation sites. Similar to the conclusions from the protein C pathway that

interactions between exosite II and CS assist with thrombin recruitment toward TM,

CS-deficient mutant of TM was less efficient for supporting the cleavage of PAR2 by

thrombin (Figure 3B). Similarly, blocking the interactions of exosite II with CS with an

aptamer, a short single-stranded DNA specifically blocking exosite II [49], interfered

with PAR2 cleavage (Figure 3C). Supporting this observation, PAR2 cleavage was

reduced by exosite II binding agent heparin (Supplementary Fig. S2). Unlike exosite

II and TM interaction, the interaction of exosite I with EGF-like domain 5 and 6 of TM

is considered essential for thrombin recruitment [57]. Moreover, blocking exosite I by

an aptamer [48] blunted PAR2 cleavage by thrombin (Figure 3C).

To test whether sTM supports PAR2 cleavage by thrombin, thrombin was pre-

incubated with recombinant sTM that lacks the membrane anchor. The mixture was

then tested for PAR2 cleavage. In the absence of cell surface-expressed TM, sTM–

thrombin mixture failed to significantly cleave PAR2 (Figure 3D). This observation

supports either the lack of sTM's role in assisting PAR2 cleavage by thrombin or the

failure of our cells to recruit sTM–thrombin complexes onto the cell surface.

Thus, similar to our observations of the protein C pathway, our data suggest that,

for efficient PAR2 cleavage by thrombin, interactions between TM's EGF-like domain

5 and the thrombin exosite I is particularly important, in addition to the efficiency-

boosting effect of the interaction between CS and thrombin exosite II, as well as the

essential role played by cell surface-anchored TM.

Figure 3: For PAR2 cleavage by thrombin, interaction between thrombin exosite I and TM's EGF-like domain 5 is necessary, along with interaction between exosite II and TM's CS motif. (**A**) 293T cells overexpressing AP-PAR2 reporter construct alone or together with either wild-type TM (TM) or mutants of TM lacking N-terminal domain(s). Scheme of the expressed TM domains is provided (bottom of A) and the details are explained in Supplementary Table S1. After 20 minutes of incubation with thrombin (30 nM), PAR2 cleavage was measured. (**B**) AP-PAR2 reporter construct was co-expressed in 293T cells with wild-type TM or a TM mutant lacking the CS sites serine 490 and 492 (TM CS-). Moreover, PAR2 cleavage by thrombin (30 nM) was measured as in (A). (**C**) 293T cells overexpressing AP-PAR2 and TM were incubated (for 20 minutes) with buffer, thrombin (30 nM), or a mixture of thrombin (30 nM) and exosite blocking aptamers (2 µM); then, PAR2 cleavage efficiency was measured. (**D**) 293T cells overexpressed AP-PAR2 with or without TM. Where indicated, cells were incubated for 20 minutes with thrombin (30 nM) alone or a 1:1 mixture of thrombin and sTM (20 minutes of pre-incubation at 37°C for complex formation). Data are presented as mean ± SEM; three independent experiments were each performed in triplicate; ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, using student's t test.

Role of PAR2 N-terminal arginine 36 in cleavage by thrombin

Previous evidence suggested that PARs harbor a unique cleavage site that

promotes initiation of biological effects [1]. Recently, however, studies have revealed

that PARs behave as "multi-switches" with several cleavage sites, resulting in

cleavage-site-specific biological effects [23, 34, 35, 58] and referred to as biased

signaling. Thus, we investigated whether TM-bound thrombin cleaved PAR2 at

specific sites. Similar to our studies in PAR1 and PAR3 [34], we synthesized mutants

of PAR2 with alanine (A) substitutions for the positively charged amino acids arginine

(R) and lysine (K) in the extracellular N-terminal domain (Supplementary Table S3).

A mutant of PAR2 that was devoid of all R and K on the entire N-terminal domain

("PAR2 all-to-A" mutant; Supplementary Table S3) resisted cleavage by thrombin

(Figure 4A). Removal of canonical R36 ("PAR2 R36A" mutant; Supplementary Table

S3) yielded an almost thrombin-resistant mutant (Figure 4B); however, additional

sites at which cleavage occurs inefficiently might be present. Thus, our data suggest

that R36 is the preferred cleavage site of TM-bound thrombin; however, the

existence of additional cleavage site(s) cannot be ruled out.

Figure 4: TM-bound thrombin preferentially cleaves PAR2 at arginine 36. (A)

and (**B**) 293T cells overexpressing TM and wild-type or mutants of AP-PAR2 reporter

constructs. The mutant "AP-PAR2 all-to-A" contained no positively charged amino

acids in the N-terminal PAR2 domain and in the "AP-PAR2 R36A" mutant, arginine

36, was replaced by alanine (details provided in Supplementary Table S3). Cells

were incubated with thrombin (30 nM) for 20 minutes before PAR2 cleavage was

measured. Data are presented as mean ± SEM; three independent experiments

were each performed in triplicate; (A-B) ns $p > 0.05$, * $p < 0.05$, ** $p < 0.01$, *** $p <$
0.001, using student's t test.
470

Role of the NF-κB pathway downstream of thrombin-cleaved PAR2

Previously, pro-inflammatory signaling was reported to occur because of PAR2
activation by other proteases that cleaved PAR2 at R36 [11-13]. Thus, we
investigated whether TM-bound thrombin cleaved PAR2 and induced pro-
inflammatory signaling via NF-κB activation. In our commercial NF-κB reporter
system, thrombin failed to induce NF-κB activation when PAR2 and TM were absent
or expressed alone (Supplementary Fig. S6a). NF-κB DNA binding activity, however,
increased in a manner dependent on thrombin concentration due to the availability of
PAR2 and TM (Figure 5A). PAR2 agonist peptide SLIGRL induced NF-κB DNA
binding activity in PAR2 and TM overexpressing cells similarly to thrombin; however,
>1000 times higher concentrations of agonist peptide was required to obtain
comparable induction (Figure 5A). Similar to thrombin, the PAR1 agonist peptide
TFLLRNPNDK induced NF-κB activation only if PAR1 was overexpressed. Similarly,
the PAR4 agonist peptide AYPGKF induced NF-κB activation only in cells
overexpressing PAR4. As expected, the PAR4 activation by thrombin was less
efficient compared to PAR1 activation. In the absence of overexpressed PAR1 or
PAR4, agonist peptides showed no such effect, similar to TFLLRNPNDK and
AYPGKF failing to activate endogenous PAR1 or PAR4 in these cells and consistent
with TFLLRNPNDK and AYPGKF failing to activate overexpressed PAR2 in this
system. However, similar to the PAR2 agonist peptide SLIGRL, thrombin-induced
NF-κB activation via PAR2 only in the presence of TM (Supplementary Fig. S6b).

Moreover, consistent with our conclusions from the PAR2 cleavage reporter
assay, EGF-like domain 5 of TM was required for thrombin- and PAR2-mediated
induction of NF-κB activation (Figure 5B).

We then tested whether thrombin's preferred cleavage site R36 was important
for NF-κB induction. In R36's absence, our assays showed diminished cleavage of
PAR2 and no induction of NF-κB activation (Figure 5C).

We then addressed the signal pathways downstream of PAR2 but upstream of
NF-κB activation. Phosphatidylinositol 3-kinase (PI3K) pathway is an important
regulator of NF-κB activation and potentially links PAR2 cleavage and activation to
the NF-κB pathway. To test whether the induction of NF-κB activation after PAR2

cleavage occurred via PI3K, we used the PI3K inhibitor LY294002. Consistent with
PI3K linking PAR2 and NF-κB, LY294002 interfered with NF-κB activation by
thrombin and activation by PAR2-specific peptide SLIGRL (Figure 5D).
Note that 293T cells secreted IL-8 with the overexpression of PAR2 and SLIGRL
agonist induction. However, as shown previously [34] and in Supplementary Fig. S6b,
cells remained quiescent upon incubation with the PAR1 agonist peptide, which
confirms our previous conclusion that, without transfection, these cells would not
respond to PAR agonists. Note that thrombin-stimulated 293T cells overexpressing
PAR2 alone remained quiescent. However, in cells expressing TM and PAR2,
thrombin (30 nM) and SLIGRL (50 μM) were comparably efficient for inducing the
release of IL-8 (Figure 5E). Thus, our data suggest that PAR2 cleaved at R36 by
TM-bound thrombin induced NF-κB DNA binding activity via PI3K and ultimately
released IL-8 release (Figure 5F).
515

Figure 5: NF-κB DNA binding activity and IL-8 release is induced by TM-bound

thrombin in a PAR2-dependent manner. (**A**) 293T cells carrying an NF-κB reporter

overexpressed either AP-PAR2 alone or AP-PAR2 together with TM, were incubated

with an agonist as indicated and quantified for NF-κB luciferase activity by measuring
luminescence. Arrows indicate thrombin (30 nM) and SLIGRL (50 μM),
concentrations used in further NF-κB reporter assays. (**B**) Cells overexpressing NF-
κB reporter, AP-PAR2 together with wild-type TM, a mutant of TM lacking all domains
at the N-terminal side of EGF-like domain 5 ("TM EGF 5–6"), or a mutant lacking all
domains at the N-terminal side of EGF-like domain 6 ("TM EGF 6"). Induction of NF-
κB luciferase activity in response to the buffer, thrombin (30 nM), and SLIGRL (50
μM) was also measured. (**C**) Cells overexpressing NF-κB reporter, TM, and either
AP-PAR2 wild-type or AP-PAR2 mutants that lacked all positively charged N-terminal
amino acids ("AP-PAR2 all-to-A") or only arginine 36 ("AP-PAR2 R36A"). Cells were
incubated with buffer, thrombin (30 nM), and SLIGRL (50 μM), after which induction
via luciferase was measured. (**D**) Cells overexpressing NF-κB reporter, TM, and AP-
PAR2 were pre-incubated with vehicle (DMSO) or the PI3K inhibitor LY294002
(Sigma) for 2 h, followed by incubation with buffer, thrombin (30 nM), or SLIGRL (50
μM) for 6 h before NF-κB luciferase induction was measured. (**E**) 293T cells
expressing AP-PAR2 or AP-PAR2 and TM were incubated with an agonist for 24 h,
followed by quantification of the release of IL-8 into the supernatant. (**F**) Scheme of
proposed PAR2 activation pathway: thrombin (IIa) binds to TM, cleaves PAR2, and
activates the NF-κB pathway via PI3K activation. NF-κB activation induces IL-8
synthesis and release. Data are presented as fold induction of buffer RLU, presented
as mean ± SEM; three independent experiments were each performed in triplicate; *
$p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, using two-way ANOVA (B, C, E), one-way
ANOVA (A, D).

Discussion

Our data show, for the first time, efficient TM-dependent direct cleavage and

activation of PAR2 by thrombin. This breaks the dogma of PAR2 as the only PAR

that resists activation by an important clotting protease thrombin. In a manner

dependent on TM co-receptor availability, efficient PAR2 cleavage occurred even at

lower concentrations of thrombin. Unexpectedly, at low concentrations of proteases,

efficiency levels of PAR2 reporter cleavage between thrombin and trypsin were

comparable. PAR2 cleavage by thrombin was - in TM's presence of TM -

comparable to cleavage of PAR1, although kinetics differed. While PAR1 was rapidly

cleaved off with minimal sustained cleavage, PAR2 cleavage increased over time.

Thus, we suggest that PAR1 is responsible for rapid thrombin responses while PAR2

sustains the thrombin signaling following PAR1 desensitization. In fact, PAR2

cleavage by thrombin was most efficient at the canonical cleavage site arginine 36.

Note that this study establishes a direct link of cleavage of PAR2 by thrombin to pro-

inflammatory signaling such as activation of NF-κB pathway and release of IL-8 in

natively PAR2- and TM-expressing A549 lung epithelial cells. To date, PAR2 has

been assumed to be distinct in the family of PARs in terms of resisting cleavage and

activation by thrombin [1, 5, 26]. However, biological effects of thrombin have now

been linked to PAR2, although the exact mechanism of activation needs to be

clarified. Transactivation of PAR2 by thrombin-cleaved PAR1 is one such standard

model [24, 59], although it is inconsistent with observations demonstrating that

thrombin maintained signaling via PAR2 under conditions where PAR1's tethered

ligand was blocked [9].

As an alternative model, two studies have confirmed extremely high

concentrations of free thrombin directly cleaving PAR2 [27, 60]. During a thrombin

burst, the required concentration of thrombin of 100–500 nM is potentially achievable

[61]. However, in-depth studies in mice that lack the thrombin platelet surface

recruiting receptor PAR3 (a nonfunctional receptor in mice) bleed, not in line with the

physiologic occurrence of thrombin concentrations at some hundred nanomolar [62].

Studies involving blocking antibodies [63] support an upper limit of physiologically

relevant thrombin concentrations of <30 nM, thus ruling out the physiological

relevance of PAR2 activation at high concentrations of free thrombin.

Similar to our co-receptor-dependent activation of PARs, as shown in the case

of PAR1 by aPC [31] and clotting factor Xa [32], PAR2 by clotting factor FVIIa [22],

or platelet-expressed PAR4 by thrombin [62], we postulated that, for efficient PAR2

activation by thrombin, a co-receptor was required. Indeed, we discovered a novel

co-receptor function of TM, supporting the cleavage and activation of PAR2 by

thrombin. Indeed, TM's presence increased thrombin's efficiency for cleaving PAR2.

Other potential PAR2 cleaving proteases induced or activated by thrombin or the

thrombin-TM complex were ruled out. Highlighting the plausibility of the presence of

TM playing a physiological role, 5 nM thrombin and 5 nM trypsin cleaved PAR2 with

comparable efficiency. Because TM is not the only cell surface receptor for thrombin

[64], other surface recruiting receptors might contribute to further PAR2 activation by

thrombin and explain the PAR2-dependent effects of thrombin in cells that did not

express TM.   We identified that the EGF-like domain 5 of TM and exosite I in

thrombin were important for TM-promoted PAR2 cleavage and NF-κB pathway

activation by thrombin. Furthermore, this conclusion was supported by removing the

EGF-like domains on TM and by blocking thrombin exosite I site using a small

specific aptamer. Our observations reinforce previous research on TM [55], showing

that in the context of the protein C pathway the described interaction is required for

cell surface recruitment of thrombin. Similarly, interactions between exosite II and CS

help in thrombin recruitment [57]. Furthermore, we found that exosite II blocking

aptamers or exosite II binding heparin interfere with PAR2 cleavage by thrombin.

These results are in accordance with a recently published study that links the TM

glycosaminoglycan domain to immune responses [65].

Thus, our data confirms that thrombin activates NF-κB and induces IL-8 release

for overexpressing 293T cells and triggers IL-8 release in A549 cells in a TM- and

PAR2-dependent manner. The specific PAR2 agonist SLIGRL induced IL-8 release,

supporting PAR2 stimulation to be sufficient for IL-8 release. Complementary,

interfering with PAR2 either via inhibition (NDGA) or knock down (siRNA)

significantly reduced responsiveness to thrombin and SLIGRL of natively PAR2

expressing A549 cells. Although thrombin-induced IL-8 release was significantly

reduced in PAR2 silenced or inhibited cells, we failed to render our system fully

thrombin irresponsive. Possible explanations include incomplete PAR2 knock down,

or another (yet to be identified) thrombin receptor induced IL-8 release.

With such a well-characterized system, we provided evidence that co-receptor-

bound thrombin directly activates pro-inflammatory pathways via PAR2. Our

evidence confirms that A549 cells did not release IL-8 on stimulation with PAR1

agonist peptide. This result is consistent with that of a recent study [53]. Our results

support that prolonged thrombin signaling via PAR2 / TM could explain pro-

inflammatory signaling in A549 alveolar cells. In view of thrombin inhibition

ameliorating chronic inflammation of the lung as reviewed in [66] one is tempted to

speculate, that thrombin / TM / PAR2 might serve as underlying mechanism. Thus,

stimulation of PAR2 by extravascular thrombin might contribute to inflammatory

diseases of the lung.

Our conclusion that thrombin activates PAR2 via the co-receptor TM is heavily

reliant on overexpression systems; thus, the physiological relevance of this has yet

to be established completely. The confirmation of our concept in a single cell line, i.e.,

A549 natively expressing PAR2 and TM, cannot eliminate this important limitation.

However, to summarize, we provide the first evidence that TM-bound thrombin

cleaves and activates PAR2 efficiently at low to moderate nanomolar concentrations

and that this pathway of NF-κB activation results in the release of IL-8 in PAR2- and

TM-overexpressing and natively expressing cells. Further studies will be required to

link this novel signaling pathway to important pathophysiological processes,

particularly for chronic inflammatory diseases such as asthma, arthritis, and cancer.

Note that such studies may reveal novel therapeutic strategies for diseases that are

linked to PAR2.

Acknowledgments

We would like to express our sincere gratitude for the support given to RAS by the

Vontobel Stiftung, the Swiss National Science Foundation grant #PZ00P3_136639,

the Hartmann Müller-Stiftung, the University Hospital Zurich, and the University of

Zurich.

Authors' Contributions

Study designed by DMH and RAS. Experiments conducted by DMH, AGF, and JM.

Data analysis and interpretation by DMH and RAS. Manuscript preparation by DMH

and RAS. All authors read and approved the final manuscript.

Disclosure of Conflict of Interest

The authors state that they have no conflicts of interest.

References

[1] S.R. Coughlin, How the protease thrombin talks to cells, Proc Natl Acad Sci U S A
96(20) (1999) 11023-7.
[2] S.R. Coughlin, Thrombin signalling and protease-activated receptors, Nature 407(6801)
(2000) 258-64.
[3] H. Ishihara, A.J. Connolly, D. Zeng, M.L. Kahn, Y.W. Zheng, C. Timmons, T. Tram, S.R.
Coughlin, Protease-activated receptor 3 is a second thrombin receptor in humans, Nature
386(6624) (1997) 502-6.
[4] M.L. Kahn, Y.W. Zheng, W. Huang, V. Bigornia, D. Zeng, S. Moff, R.V. Farese, Jr., C.
Tam, S.R. Coughlin, A dual thrombin receptor system for platelet activation, Nature
394(6694) (1998) 690-4.
[5] S. Nystedt, K. Emilsson, C. Wahlestedt, J. Sundelin, Molecular cloning of a potential
proteinase activated receptor, Proc Natl Acad Sci U S A 91(20) (1994) 9208-12.

[6] T.K. Vu, D.T. Hung, V.I. Wheaton, S.R. Coughlin, Molecular cloning of a functional
thrombin receptor reveals a novel proteolytic mechanism of receptor activation, Cell 64(6)
(1991) 1057-68.
[7] W.F. Xu, H. Andersen, T.E. Whitmore, S.R. Presnell, D.P. Yee, A. Ching, T. Gilbert,
E.W. Davie, D.C. Foster, Cloning and characterization of human protease-activated receptor
4, Proc Natl Acad Sci U S A 95(12) (1998) 6642-6.
[8] E.B. Kelso, J.C. Lockhart, T. Hembrough, L. Dunning, R. Plevin, M.D. Hollenberg, C.P.
Sommerhoff, J.S. McLean, W.R. Ferrell, Therapeutic Promise of Par2 Antagonism in Joint
Inflammation, J Pharmacol Exp Ther (2005).
[9] X. Shi, B. Gangadharan, L.F. Brass, W. Ruf, B.M. Mueller, Protease-activated receptors
(PAR1 and PAR2) contribute to tumor cell motility and metastasis, Mol Cancer Res 2(7)
(2004) 395-402.
[10] N. Vergnolle, N.W. Bunnett, K.A. Sharkey, V. Brussee, S.J. Compton, E.F. Grady, G.
Cirino, N. Gerard, A.I. Basbaum, P. Andrade-Gordon, M.D. Hollenberg, J.L. Wallace,
Proteinase-activated receptor-2 and hyperalgesia: A novel pain pathway, Nat Med 7(7)
(2001) 821-6.
[11] F. Schmidlin, S. Amadesi, K. Dabbagh, D.E. Lewis, P. Knott, N.W. Bunnett, P.R. Gater,
P. Geppetti, C. Bertrand, M.E. Stevens, Protease-activated receptor 2 mediates eosinophil
infiltration and hyperreactivity in allergic inflammation of the airway, J Immunol 169(9)
(2002) 5315-21.
[12] V. Knight, J. Tchongue, D. Lourensz, P. Tipping, W. Sievert, Protease-activated receptor
2 promotes experimental liver fibrosis in mice and activates human hepatic stellate cells,
Hepatology 55(3) (2012) 879-87.
[13] N.W. Bunnett, Protease-activated receptors: how proteases signal to cells to cause
inflammation and pain, Semin Thromb Hemost 32 Suppl 1 (2006) 39-48.
[14] I. Seitz, S. Hess, H. Schulz, R. Eckl, G. Busch, H.P. Montens, R. Brandl, S. Seidl, A.
Schomig, I. Ott, Membrane-type serine protease-1/matriptase induces interleukin-6 and -8 in
endothelial cells by activation of protease-activated receptor-2: potential implications in
atherosclerosis, Arterioscler Thromb Vasc Biol 27(4) (2007) 769-75.
[15] S. Antoniak, M. Rojas, D. Spring, T.A. Bullard, E.D. Verrier, B.C. Blaxall, N. Mackman,
R. Pawlinski, Protease-activated receptor 2 deficiency reduces cardiac ischemia/reperfusion
injury, Arterioscler Thromb Vasc Biol 30(11) (2010) 2136-42.
[16] S.M. Jones, A. Mann, K. Conrad, K. Saum, D.E. Hall, L.M. McKinney, N. Robbins, J.
Thompson, A.D. Peairs, E. Camerer, K.J. Rayner, M. Tranter, N. Mackman, P. Owens, PAR2
(Protease-Activated Receptor 2) Deficiency Attenuates Atherosclerosis in Mice, Arterioscler
Thromb Vasc Biol 38(4) (2018).
[17] H.H. Versteeg, F. Schaffner, M. Kerver, H.H. Petersen, J. Ahamed, B. Felding-
Habermann, Y. Takada, B.M. Mueller, W. Ruf, Inhibition of tissue factor signaling
suppresses tumor growth, Blood 111(1) (2008) 190-9.
[18] M. Jaber, M. Maoz, A. Kancharla, D. Agranovich, T. Peretz, S. Grisaru-Granovsky, B.
Uziely, R. Bar-Shavit, Protease-activated-receptor-2 affects protease-activated-receptor-1-
driven breast cancer, Cell Mol Life Sci (2013).
[19] F. Schaffner, H.H. Versteeg, A. Schillert, N. Yokota, L.C. Petersen, B.M. Mueller, W.
Ruf, Cooperation of tissue factor cytoplasmic domain and PAR2 signaling in breast cancer
development, Blood (2010).
[20] H. Lin, J. Trejo, Transactivation of the PAR1-PAR2 heterodimer by thrombin elicits
beta-arrestin-mediated endosomal signaling, J Biol Chem 288(16) (2013) 11203-15.
[21] S. Nystedt, K. Emilsson, A.K. Larsson, B. Strombeck, J. Sundelin, Molecular cloning
and functional expression of the gene encoding the human proteinase-activated receptor 2,
Eur J Biochem 232(1) (1995) 84-9.

[22] E. Camerer, W. Huang, S.R. Coughlin, Tissue factor- and factor X-dependent activation
of protease-activated receptor 2 by factor VIIa, Proc Natl Acad Sci U S A 97(10) (2000)
5255-60.
[23] R. Ramachandran, K. Mihara, H. Chung, B. Renaux, C.S. Lau, D.A. Muruve, K.A.
DeFea, M. Bouvier, M.D. Hollenberg, Neutrophil elastase acts as a biased agonist for
proteinase-activated receptor-2 (PAR2), J Biol Chem 286(28) (2011) 24638-48.
[24] N.C. Kaneider, A.J. Leger, A. Agarwal, N. Nguyen, G. Perides, C. Derian, L. Covic, A.
Kuliopulos, 'Role reversal' for the receptor PAR1 in sepsis-induced vascular damage, Nat
Immunol 8(12) (2007) 1303-12.
[25] M.D. Hollenberg, K. Mihara, D. Polley, J.Y. Suen, A. Han, D.P. Fairlie, R.
Ramachandran, Biased signalling and proteinase-activated receptors (PARs): targeting
inflammatory disease, Br J Pharmacol 171(5) (2014) 1180-94.
[26] P.J. O'Brien, M. Molino, M. Kahn, L.F. Brass, Protease activated receptors: theme and
variations, Oncogene 20(13) (2001) 1570-81.
[27] K. Mihara, R. Ramachandran, M. Saifeddine, K.K. Hansen, B. Renaux, D. Polley, S.
Gibson, C. Vanderboor, M.D. Hollenberg, Thrombin-mediated direct activation of
proteinase-activated receptor-2 (PAR2): another target for thrombin signaling, Mol
Pharmacol (2016).
[28] H. Suzuki, E.D. Motley, K. Eguchi, A. Hinoki, H. Shirai, V. Watts, L.N. Stemmle, T.A.
Fields, S. Eguchi, Distinct roles of protease-activated receptors in signal transduction
regulation of endothelial nitric oxide synthase, Hypertension 53(2) (2009) 182-8.
[29] E.A. Lidington, R. Steinberg, A.R. Kinderlerer, R.C. Landis, M. Ohba, A. Samarel, D.O.
Haskard, J.C. Mason, A role for proteinase-activated receptor 2 and PKC-epsilon in
thrombin-mediated induction of decay-accelerating factor on human endothelial cells, Am J
Physiol Cell Physiol 289(6) (2005) C1437-47.
[30] R. Pawlinski, M. Holinstat, We can do it together: PAR1/PAR2 heterodimer signaling in
VSMCs, Arterioscler Thromb Vasc Biol 31(12) (2011) 2775-6.
[31] M. Riewald, R.J. Petrovan, A. Donner, B.M. Mueller, W. Ruf, Activation of endothelial
cell protease activated receptor 1 by the protein C pathway, Science 296(5574) (2002) 1880-2.
[32] R.A. Schuepbach, M. Riewald, Coagulation factor Xa cleaves protease-activated
receptor-1 and mediates signaling dependent on binding to the endothelial protein C receptor,
J Thromb Haemost 8(2) (2010) 379-88.
[33] H. Guo, D. Liu, H. Gelbard, T. Cheng, R. Insalaco, J.A. Fernandez, J.H. Griffin, B.V.
Zlokovic, Activated protein C prevents neuronal apoptosis via protease activated receptors 1
and 3, Neuron 41(4) (2004) 563-72.
[34] R.A. Schuepbach, J. Madon, M. Ender, P. Galli, M. Riewald, Protease Activated
Receptor-1 Cleaved at R46 Mediates Cytoprotective Effects, J Thromb Haemost (2012).
[35] L. Burnier, L.O. Mosnier, Novel mechanisms for activated protein C cytoprotective
activities involving noncanonical activation of protease-activated receptor 3, Blood 122(5)
(2013) 807-16.
[36] M. Nakanishi-Matsui, Y.W. Zheng, D.J. Sulciner, E.J. Weiss, M.J. Ludeman, S.R.
Coughlin, PAR3 is a cofactor for PAR4 activation by thrombin, Nature 404(6778) (2000)
609-13.
[37] K. Suzuki, H. Kusumoto, Y. Deyashiki, J. Nishioka, I. Maruyama, M. Zushi, S.
Kawahara, G. Honda, S. Yamamoto, S. Horiguchi, Structure and Expression of Human
Thrombomodulin, a Thrombin Receptor on Endothelium Acting as a Cofactor for Protein-C
Activation, Embo J 6(7) (1987) 1891-1897.
[38] D.J. Stearns, S. Kurosawa, C.T. Esmon, Microthrombomodulin. Residues 310-486 from
the epidermal growth factor precursor homology domain of thrombomodulin will accelerate
protein C activation, J Biol Chem 264(6) (1989) 3352-6.

[39] C.T. Esmon, W.G. Owen, Identification of an endothelial cell cofactor for thrombin-
catalyzed activation of protein C, Proc Natl Acad Sci U S A 78(4) (1981) 2249-52.
[40] T.J. Raife, D.J. Lager, K.C. Madison, W.W. Piette, E.J. Howard, M.T. Sturm, Y. Chen,
S.R. Lentz, Thrombomodulin Expression by Human Keratinocytes Induction of Cofactor
Activity during Epidermal Differentiation, J Clin Invest 93(4) (1994) 1846-1851.
[41] A. Pindon, M. Berry, D. Hantai, Thrombomodulin as a new marker of lesion-induced
astrogliosis: Involvement of thrombin through the G-protein-coupled protease activated
receptor-1, J Neurosci 20(7) (2000) 2543-2550.
[42] S.S. Mccachren, J. Diggs, J.B. Weinberg, W.A. Dittman, Thrombomodulin Expression
by Human Blood Monocytes and by Human Synovial Tissue Lining Macrophages, Blood
78(12) (1991) 3128-3132.
[43] L. Wang, J.A. Bastarache, N. Wickersham, X.H. Fang, M.A. Matthay, L.B. Ware, Novel
role of the human alveolar epithelium in regulating intra-alveolar coagulation, Am J Resp
Cell Mol 36(4) (2007) 497-503.
[44] C.T. Esmon, Thrombomodulin as a model of molecular mechanisms that modulate
protease specificity and function at the vessel surface, Faseb J 9(10) (1995) 946-55.
[45] D.A. Lane, H. Philippou, J.A. Huntington, Directing thrombin, Blood 106(8) (2005)
2605-12.
[46] T. Okamoto, H. Tanigami, K. Suzuki, M. Shimaoka, Thrombomodulin: a bifunctional
modulator of inflammation and coagulation in sepsis, Crit Care Res Pract 2012 (2012)
614545.
[47] H. Loghmani, E.M. Conway, Exploring traditional and nontraditional roles for
thrombomodulin, Blood 132(2) (2018) 148-158.
[48] R.F. Macaya, P. Schultze, F.W. Smith, J.A. Roe, J. Feigon, Thrombin-binding DNA
aptamer forms a unimolecular quadruplex structure in solution, Proc Natl Acad Sci U S A
90(8) (1993) 3745-9.
[49] K. Segers, B. Dahlback, P.E. Bock, G. Tans, J. Rosing, G.A. Nicolaes, The role of
thrombin exosites I and II in the activation of human coagulation factor V, J Biol Chem
282(47) (2007) 33915-24.
[50] R.A. Schuepbach, C. Feistritzer, L.F. Brass, M. Riewald, Activated protein C-cleaved
protease activated receptor-1 is retained on the endothelial cell surface even in the presence
of thrombin, Blood 111(5) (2008) 2667-73.
[51] N. Asokananthan, P.T. Graham, J. Fink, D.A. Knight, A.J. Bakker, A.S. McWilliam, P.J.
Thompson, G.A. Stewart, Activation of protease-activated receptor (PAR)-1, PAR-2, and
PAR-4 stimulates IL-6, IL-8, and prostaglandin E2 release from human respiratory epithelial
cells, J Immunol 168(7) (2002) 3577-85.
[52] H. Wang, Y. Zheng, S. He, Induction of release and up-regulated gene expression of
interleukin (IL)-8 in A549 cells by serine proteinases, BMC Cell Biol 7 (2006) 22.
[53] E. Ostrowska, E. Sokolova, G. Reiser, PAR-2 activation and LPS synergistically
enhance inflammatory signaling in airway epithelial cells by raising PAR expression level
and interleukin-8 release, Am J Physiol Lung Cell Mol Physiol 293(5) (2007) L1208-18.
[54] M. Ender, F. Andreoni, A.S. Zinkernagel, R.A. Schuepbach, Streptococcal SpeB
Cleaved PAR-1 Suppresses ERK Phosphorylation and Blunts Thrombin-Induced Platelet
Aggregation, PLoS One 8(11) (2013) e81298.
[55] K. Suzuki, T. Hayashi, J. Nishioka, Y. Kosaka, M. Zushi, G. Honda, S. Yamamoto, A
domain composed of epidermal growth factor-like structures of human thrombomodulin is
essential for thrombin binding and for protein C activation, J Biol Chem 264(9) (1989) 4872-
6.
[56] M.C. Bourin, U. Lindahl, Glycosaminoglycans and the regulation of blood coagulation,
Biochem J 289 (Pt 2) (1993) 313-30.

[57] J.W. Weisel, C. Nagaswami, T.A. Young, D.R. Light, The shape of thrombomodulin and
interactions with thrombin as determined by electron microscopy, J Biol Chem 271(49)
(1996) 31485-90.
[58] V. Trivedi, A. Boire, B. Tchernychev, N.C. Kaneider, A.J. Leger, K. O'Callaghan, L.
Covic, A. Kuliopulos, Platelet matrix metalloprotease-1 mediates thrombogenesis by
activating PAR1 at a cryptic ligand site, Cell 137(2) (2009) 332-43.
[59] L.M. Sevigny, K.M. Austin, P. Zhang, S. Kasuda, G. Koukos, S. Sharifi, L. Covic, A.
Kuliopulos, Protease-activated receptor-2 modulates protease-activated receptor-1-driven
neointimal hyperplasia, Arterioscler Thromb Vasc Biol 31(12) (2011) e100-6.
[60] D. Loew, C. Perrault, M. Morales, S. Moog, C. Ravanat, S. Schuhler, R. Arcone, C.
Pietropaolo, J.P. Cazenave, A. van Dorsselaer, F. Lanza, Proteolysis of the exodomain of
recombinant protease-activated receptors: prediction of receptor activation or inactivation by
MALDI mass spectrometry, Biochemistry 39(35) (2000) 10812-22.
[61] S. Butenas, C. van't Veer, K.G. Mann, "Normal" thrombin generation, Blood 94(7)
(1999) 2169-78.
[62] G.R. Sambrano, E.J. Weiss, Y.W. Zheng, W. Huang, S.R. Coughlin, Role of thrombin
signalling in platelets in haemostasis and thrombosis, Nature 413(6851) (2001) 74-8.
[63] M.L. Kahn, M. Nakanishi-Matsui, M.J. Shapiro, H. Ishihara, S.R. Coughlin, Protease-
activated receptors 1 and 4 mediate activation of human platelets by thrombin, J Clin Invest
103(6) (1999) 879-87.
[64] D. Dormann, K.J. Clemetson, B.E. Kehrel, The GPIb thrombin-binding site is essential
for thrombin-induced platelet procoagulant activity, Blood 96(7) (2000) 2469-78.
[65] V.C. Pai, I.C. Lo, Y.W. Huang, I.C. Tsai, H.P. Cheng, G.Y. Shi, H.L. Wu, M.J. Jiang,
The chondroitin sulfate moiety mediates thrombomodulin-enhanced adhesion and migration
of vascular smooth muscle cells, J Biomed Sci 25(1) (2018) 14.
[66] J.K. Shute, E. Puxeddu, L. Calzetta, Therapeutic use of heparin and derivatives beyond
anticoagulation in patients with bronchial asthma or COPD, Curr Opin Pharmacol 40 (2018)
39-45.
840
841