Branching Networks Can Have Opposing Influences on Genetic Variation in

Riverine Metapopulations

Ming-Chih Chiu[1,2], Bin Li[1,4], Kei Nukazawa[3], Vincent H. Resh[5], Thaddeus Carvajal[1], and Kozo

Watanabe[1*]

[1]Department of Civil and Environmental Engineering, Ehime University, Matsuyama,

790-8577, Japan

[2]Institute of Hydrobiology, Chinese Academy of Sciences, Wuhan, 430070, China

[3]Department of Civil and Environmental Engineering, University of Miyazaki, Miyazaki,

889-2192, Japan

[4]Institute of Environment and Ecology, Shandong Normal University, Jinan, 250000, China

[5]Department of Environmental Science, Policy & Management, University of California,

Berkeley, CA 94720, USA

[*]Corresponding Author:

E-mail address: watanabe_kozo@cee.ehime-u.ac.jp; Phone & Fax: +81 (0) 89-927-9847

Ming-Chih Chiu: mingchih.chiu@gmail.com; Bin Li: binglee527@sdnu.edu.cn; Kei Nukazawa:

kei.nukazawa@gmail.com; Vincent H. Resh: resh@berkeley.edu; Thaddeus Carvajal:

tads.carvajal@gmail.com

Running Head: Genetic effects of river branching

Acknowledgements

We are grateful to Maribet Gamboa for her constructive comments on the manuscript. This

study was partly funded by the Japan Society for the Promotion of Science (JSPS). MCC was

supported by the JSPS (JSPS Invitation Fellowship Program; L18522) and the Chinese

Academy of Sciences (CAS Taiwan Young Talent Programme; 2017TW2SA0004).

Author Contributions: All authors conceived the research. MCC, BI, KN, and TC performed

analyses, and MCC performed modeling work. MCC, KW, and VHR wrote the draft of the

manuscript, and all authors contributed substantially to revisions.

Conflict of Interest: All authors have no conflict of interest to declare.

Abstract

Aim: Fractal networks, represented by branching complexity in rivers, are ubiquitous in nature.

In rivers, the number of either distal (e.g., in headwater streams) or confluent (e.g., in

mainstems) locations can be increased along with their branching complexity. Distal- or

confluent-spatial locations can result in fewer or greater corridor linkages that can alter genetic

divergence at the metapopulation scale. These mechanisms underlying the resulting genetic

structuring remain poorly understood at the metapopulation scale, particularly in terms of the

roles of species-specific dispersal traits. The objective of this study is to mechanistically

understand how branching complexity can simultaneously influence genetic divergence in

opposite directions.

Location: Northeastern Japan

Methods: To evaluate the integrated influences of network complexity and species dispersal

on genetic divergence among populations at the catchment scale, we conducted simulation

modelling on a mechanistic framework based on Bayesian inference by adapting empirical

genetic data from four macroinvertebrate species. Simulations were then done using empirical

and virtual species-characteristics on virtual river networks.

Results: Our novel simulation showed that both greater landscape connectivity (resulting from

shorter watercourse distance) and greater isolation of distal locations occurred in the

more-branched river networks. These two spatial features have negative and positive

influences on genetic divergence, with their relative importance varying among different

species and dispersal characteristics. Specifically, genetic divergence at the metapopulation

scale increased for species having higher downstream-biased dispersal but decreased for

species having higher upstream-biased dispersal. Distal populations (e.g., in headwaters) have

higher genetic independence when downstream-biased asymmetry is higher.

Main conclusions: We found a strong association between species dispersal and evolutionary

processes such as gene flow and genetic drift. This association mediates the pervasive

influences of branching complexity on genetic-divergence in the metapopulation. It also

highlights the importance of considering species dispersal-patterns when developing

management strategies in the face of rapid environmental-change scenarios.

Keywords: Bayesian inference, distribution, fractal geometry, habitat fragmentation, isolation

by distance, landscape complexity, macroinvertebrates

33
34

Introduction

There is increasing interest in understanding how landscape architecture determines spatial

patterns of intraspecific genetic-diversity (Paz-Vinas, Loot, Stevens, & Blanchet, 2015;

Phillipsen & Lytle, 2013), although a mechanistic understanding of this process remains highly

challenging in the real world. For example, ecological and evolutionary evidence and theories

that are derived from simplified landscapes are insufficient for understanding spatial genetic

patterns in complex systems, such as rivers (Fourcade, Chaput-Bardy, Secondi, Fleurant, &

Lemaire, 2013; Thomaz, Christie, & Knowles, 2016). More in-depth explorations of the

integrated genetic effects of species dispersal and landscape connectivity on metapopulations

(here defined as groups of subpopulations with dispersal interactions) occurring in complex

habitats are needed. These efforts are especially important when fragmented landscapes that

result from global-change influences affect spatial genetic variability (Martins et al., 2016;

Prunier, Dubut, Loot, Tudesque, & Blanchet, 2018).

In nature, fractal branching-networks (e.g., those with "treelike" patterns) have similar

structural features to those seen in fluvial landscapes (Green, Klomp, Rimmington, & Sadedin,

2006). In these environments, landscape connectivity shapes evolutionary processes, such as

gene flow and genetic drift, which ultimately produce the spatial patterns observed in

intraspecific genetic variation (McRae, 2006; Phillipsen et al., 2015; Thomaz et al., 2016). In

analyzing the evolutionary action of landscape architecture, branching networks can be

characterised by being either distal or confluent locations (e.g., in headwater streams or

mainstems, respectively), which allow fewer or greater corridor linkages, respectively.

Therefore, these distal and confluent locations may have opposing influences by increasing or

decreasing genetic divergence at the metapopulation scale, respectively (Alp, Keller, Westram,

& Robinson, 2012).

The tendency of species to disperse among habitats is not necessarily occurring in a single

direction (i.e., toward distal or confluent areas) or in different directions. These differences can

result in dispersal asymmetry that can dictate the isolation processes between pairs of

populations within a metapopulation (Junker et al., 2012; Ma, Shen, Bearup, Fagan, & Liao,

2020; Morrissey & de Kerckhove, 2009). For example, riverine populations occurring in a

network's distal branches (e.g., different headwaters of a riverine system) can still be

connected to a common source population in downstream confluences through dispersal.

However, downstream-biased dispersal (a tendency for higher dispersal downstream than

upstream) may lead to weak connections among headwaters and result in a large genetic

divergence among riverine species of fish and macroinvertebrates (Alp et al., 2012; Paz-Vinas

& Blanchet, 2015; Paz-Vinas, Quéméré, Chikhi, Loot, & Blanchet, 2013). Theoretically, more

isolated tributaries within a highly-branched network would result in higher genetic

differentiation between local populations (Thomaz et al., 2016).

In contrast to the effects of dispersal asymmetry, network branching (e.g., in river systems)

can enhance connectivity between populations by naturally increasing the number of

confluences and by shortening the distance of their pathways (Labonne, Ravigné, Parisi, &

Gaucherel, 2008; Paz-Vinas & Blanchet, 2015). Stream-dwelling species with strong

tendencies to migrate upstream, such as aquatic insects that disperse by flying during their

terrestrial adult stages (Petersen, Masters, Hildrew, & Ormerod, 2004; Winterbourn,

Chadderton, Entrekin, Tank, & Harding, 2007), may exhibit low downstream- or

upstream-biased gene flow. In this case, there is weaker isolation between distal populations in

the river network when these populations receive more gene flow because of their shared

source-population that occurs at downstream confluences (Alp et al., 2012). Therefore,

dispersal asymmetry provides a mechanism that can underlie the widely observed patterns of

spatial genetic diversity and genetic differentiation that is found in river systems (Kawecki &

Holt, 2002; Pilger, Gido, Propst, Whitney, & Turner, 2017).

Based on simulation modelling, the objective of this study is to mechanistically

understand how branching complexity can influence genetic divergence, and how it can do this

in opposing ways depending on dispersal traits. Although dendritic structure and species

dispersal have been reported to determine spatial genetic patterns across a river network in

Switzerland (e.g., Alp et al., 2012), our study extends this observation to the metapopulation

level through the comparison of multiple networks having varying levels of branching

complexity (e.g., as in Thomaz et al., 2016). Through empirical data of riverine

macroinvertebrates in northeastern Japan, we constructed the genetic simulations using

empirical and virtual species-characteristics on virtual river networks. Herein, we hypothesised

that increased network branching has positive effects on genetic divergence at the

metapopulation scale for species with downstream-biased dispersal but has the opposite,

negative effect for those with symmetric dispersal or upstream-biased asymmetric dispersal.

This difference implies that the consequences of network branching, namely landscape

connectivity (via watercourse distance) and the isolation of distal locations (e.g., headwaters),

may have opposite effects on genetic structure and attributes of the metapopulation.

Materials and methods

Framework overview

The opposing influences of different dispersal modes on the genetic divergence within a

metapopulation can be theoretically explained but these influences have not been demonstrated

to occur along a gradient of river branching in multiple networks. To demonstrate these

influences, we followed two successive steps. First, by using a mechanistic model based on

evolutionary processes and asymmetric dispersal, we explored the spatial genetic variation

among four macroinvertebrate species with flying adult-stages that occur in a river network of

northeastern Japan. Second, we evaluated how branching complexity of random virtual

networks can differentially affect the broader genetic differentiation throughout catchments in

the context of different asymmetric gene-flow modes.

To achieve the first step, we used the empirical data (see the subsequent section

'*Empirical catchment and genetic data*') and developed a metapopulation genetic-model based

on isolation by distance (i.e., lower gene flow with greater separation in terms of distance

along the watercourse) and asymmetric dispersal (i.e., upstream- and downstream-biased

movements) between local populations in the river network (see the subsequent section

'*Development of metapopulation genetic model*'). Rather than using their actual locations, the

watercourse distances between local populations were used directly as the input data in the

model. Consequently, if the spatial connectivity among some local populations is high, then

their genetic structure would tend to have consistent differences from that of their

metapopulation (their genetic pool). In turn, this model was fitted to these empirical data with

its parameters estimated by Bayesian inference (see the subsequent section '*Bayesian fitting*

process'). To validate this inference process using the approach of Zurell et al. (2010), we

performed this metapopulation genetic modelling with known values of parameters to confirm

whether the Bayesian estimates matched these known values (Appendix S1). The Bayesian

filtering process can provide the reliable estimation of model parameters when these known

values were within the 95% confidence interval of the posterior distribution of estimates.

However, the variance parameters (σ_D^2 and σ_U^2) had relatively large uncertainty in the scenario

of equal scale parameters ($c_D = c_U$; Figure S2).

Bayesian estimation can provide realistic ranges of dispersal asymmetry and other

parameters. In our model, these parameters were related to biological features of the four

species and were not independent of each other. Because a particular species can be described

as having a certain combination of parameters based on a set of trait characteristics and through

its systematic clade, comparisons among these species can reveal the opposing influences of

branching complexity on metapopulation genetic-structure. This can be done by examining the

parameter space (i.e., the ranges of possible values of multiple parameters) and how each

region of this parameter space reacts to different branching contexts. We used these ranges in

our next step to achieve the main objective of this study, which was to evaluate the influence of

network branching, and make our simulations be closer to the empirical system than to an

arbitrary determination. The different parameter combinations (derived from Bayesian fitting

process for the four macroinvertebrate species) allowed the creation of virtual species.

For the second step, the metapopulation genetic model constructed above was used to

simulate the river branching influence on genetic divergence in different dispersal modes (i.e.,

by setting up the virtual species) in 1,000 virtual networks (see the subsequent section

'*Simulation of river-branching influences*'). These virtual networks were randomly created to

represent a variety of branching complexity.

Empirical catchment and genetic data

In an empirical catchment in the Natori and Nanakita Rivers located in northeastern Japan

(integrated catchment area c. 1200 km2; Figure 1), the flow regime exhibits a seasonal pattern,

with spring flooding from snowmelt. In the integrated catchment, the rivers flow from the

western headwaters at an elevation of 1350 m on Mount Kamuro to the east and into the

Pacific Ocean. Approximately 60% of this area is forested and mountainous. Two major

reservoir dams (Kamafusa and Okura dams) are located there on the rivers. The regional

lowlands are farmlands (13%, primarily with rice paddy fields) and a mixture of residential and

commercial areas (11%).

Genetic data were generated from a previous study (Watanabe, Kazama, Omura, &

Monaghan, 2014). We used neutral AFLP markers from four macroinvertebrate species (the

mayfly *Ephemera japonica*, and the caddisflies *Hydropsyche orientalis*, *Stenopsyche*

marmorata, and *Hydropsyche albicephala*) in this catchment (Appendix S1 and Figure S1).

These species all have substantially diverse habitat specificity and distributions within the

network (Nukazawa, Kazama, & Watanabe, 2015; Nukazawa, Kazama, & Watanabe, 2017;

Watanabe et al., 2014).

Development of metapopulation genetic model

This metapopulation genetic model has the two modules that correspond to observation and to

evolutionary processes. The observation module describes the statistical inference of the true

value of population's genetic-frequency based on genetic measurements derived from our

sampling of parts of individuals of population. We use Bayesian inference (see the subsequent

section '*Bayesian fitting process*') to fit the output of this module to our genetic measurements

and then to calibrate the model parameters. By connecting statistical frameworks with two key

processes of gene flow and genetic drift, the evolutionary module allows the genetic frequency

of each local population to be equivalent to the frequency of the metapopulation with an

estimate of the deviation of the frequency. The deviation is based on the genetic similarity of

distance-decay among local populations, with the values of deviation tending to be similar if

these populations have high ecological connectivity among them along watercourses. The

paragraphs below describe the statistical inference used for the descriptions of these two

modules.

The observation module describes how genetic measurements and sampling errors are

produced. A single AFLP locus has two allelic types, labelled '1' and '0' for the dominant and

recessive types, respectively. We denote by $z_{k,l}$ the number of individuals for which the

dominant type '1' corresponds to genotypes ('1', '1') or ('1', '0') in our diploid species, which

was detected at locus l from the total number of individuals (s_k; with types '1' and '0' together)

in local population k. Here, $z_{k,l}$ was used as the modelling output calibrated by our observed

data. This random sampling process can be characterised by a binomial distribution as follows

(Guillot, Vitalis, Rouzic, & Gautier, 2014):

187
$$z_{k,l} \sim \text{Binomial}\left(s_k, f'_{k,l}\right) \text{ with } f'_{k,l} = 2f_{k,l}\left(1 - f_{k,l}\right) + f^2_{k,l} \qquad [1]$$

where, in this local population, the frequency of allele '1' is denoted by $f_{k,l}$. The allele

frequencies are independent between loci. The detection probability $f'_{k,l}$ linked to $f_{k,l}$ was set

up for the allele type '1' corresponding to genotypes.

The evolutionary module describes the genetic structuring of local populations within a

metapopulation. For each locus, the frequency of allele '1' in the local populations is

determined by their genetic variation (i.e., that related to genetic drift), the watercourse

distance between local populations within the network (related to gene flow), and the allele

frequencies of the metapopulation. Without genetic drift and natural selection, gene flow leads

to genetic homogeneity among local populations, which results in allele frequencies at loci in

local populations matching those of their metapopulation (Andrews, 2010). We denote the

deviation from the expected frequency of allele '1' in a logit scale as $\theta_{k,l}$, which was derived

from the assumption of local populations being connected at the metapopulation scale. The

allele frequency $f_{k,l}$ is obtained from the inverse logit transformation as follows:

201
$$f_{k,l} = \text{invLogit}\left(\theta_{k,l} + m_l\right) = \frac{1}{1 + \exp\left(-\left(\theta_{k,l} + m_l\right)\right)} \qquad [2]$$

202   where m_l denotes the metapopulation's logit-transformed allele frequency. The deviation $\theta_{k,l}$

203   can be modelled by a multivariate normal distribution as follows (Bradburd, Ralph, & Coop,

204   2013):

205
$$\theta_{k,l} \sim \text{MultiNormal}(\mu, \Omega) \quad [3]$$

where μ denotes the vector of zero, and the covariance matrix Ω is a function of the

watercourse distance between local populations and their spatial relationships. These spatial

connections between local populations are either streamflow-connected (populations with

downstream–upstream relationships) or streamflow-disconnected (e.g., populations in different

headwaters). This statistical process allows similar values of deviation for local populations

having high ecological connectivity along watercourses. We modelled the covariance across

local populations as a function of the shortest watercourse distance, h_{ij}, along the river

network between populations *i* and *j* as follows (Ver Hoef & Peterson, 2010):

214
$$\Omega_{ij} = \begin{cases} \sigma_D^2 + \sigma_U^2 & \text{if flow-connected and } i = j \\ \sigma_D^2 \exp\left(-\left(c_D h_{ij}\right)\right) + \sigma_U^2 \exp\left(-\left(c_U h_{ij}\right)\right) & \text{if flow-connected and } i \neq j \\ 0 + \sigma_U^2 \exp\left(-\left(c_U h_{ij}\right)\right) & \text{if flow-disconnected} \end{cases} \quad [4]$$

where the first or second addend (i.e., for any number that's added to another; e.g., σ_D^2 and σ_U^2

are different addends of their summation $\sigma_D^2 + \sigma_U^2$) describes the autocovariance, with the

variance σ_D^2 or σ_U^2 and the scale parameter c_D or c_U related to the downstream (D) or

upstream (U) movement, respectively. In the first addend, the autocovariance is set to zero for

any two streamflow-disconnected populations. In other words, streamflow-disconnected

populations are assumed to be independent and have no gene flow between them via

downstream movement. In the second addend, the autocovariance can apply to both situations.

The genetic connections between streamflow-disconnected populations are attributed to the

upstream gene-flow coming from a common source population in the downstream confluences.

High values of variance and small values of the scale parameter lead to weak isolation as a

function of the watercourse distance.

Bayesian fitting process

In the Bayesian framework 'Stan' (Stan Development Team, 2018), the R interface 'RStan'

(Stan Development Team, 2019) was used for genetic modelling of the metapopulation and to

fit the model using our empirical data (see the above section '*Framework overview*'). For each

species, four Markov Chain Monte Carlo (MCMC) chains (which were used for numerical

approximations of Bayesian inference) were run with 20,000 iterations each, and the first half

of the iterations for each chain were discarded as "burn-in". This iteration setting was

determined by the convergence of MCMC chains when the R-hat statistic of each parameter

approached a value of 1. To estimate the model parameters, 2,000 samples obtained by

collecting one sample out of every 20 iterations for each chain were used to build each

parameter's posterior distribution. The target average for the proposal acceptance-probability

and initial discretization interval was set to 0.95 and 0.1, respectively, during Stan's adaptation

period.

Prior distributions were set up for the model parameters based on their data type (e.g.,

continuous or discrete), range of values (e.g., unlimited or positive), adjustments using an

informative feature to keep reasonable values but not eliminating those that might be

appropriate, parameter validation (through the virtual ecologist approach; see Appendix S1),

and model-convergence efficacy. The transformed allele frequency m_l has a normal

distribution with mean of 0 and variance τ, and a half-Cauchy distributed hyper-prior on the

variance τ with a location of 0 and a scale of 5. The variances (σ_D^2 and σ_U^2) each have a

half-Cauchy distribution with a location of 0 and a scale of 5. The scale parameters (c_D and

c_U) each have a half-normal distribution with mean of 0 and variance of 2. The four

macroinvertebrate species examined (*E. japonica*, *H. orientalis*, *S. marmorata* and *H.*

albicephala) each had both types of relationship between locations (i.e., streamflow-connected

and -disconnected pairs).

Creation of virtual river networks

We created 1,000 virtual river networks with varying branching-complexities (Terui et al.,

2018; Yeakel, Moore, Guimarães Jr., & de Aguiar, 2014). These virtual river networks (with

scale length *e* equal to 1 km between local populations) have the branching probability *P* and

metapopulation size N (which is an integer and represents the number of local populations in a

river network) randomly drawn from 0 to 1 and from 100 to 500, respectively (Figure S3). This

scale length was determined by referring to the local scale of stream macroinvertebrate

occurrence, e.g., a 100 m reach (Stoll, Breyer, Tonkin, Früh, & Haase, 2016). Natural river

networks in Japan have a range of P from 0.3 to 0.6 (Terui et al., 2018). The full range of P can

reflect general conditions and provide insights into networks that, because of human

intervention, depart from natural value of P (e.g., a network with $P = 0.1$ can represent a

human-created canal; Figure S3).

The virtual river networks comprised nodes with scale length e, with each node

representing a local population. These nodes were assigned to be either branching (or

representing an upstream terminal) or non-branching with a probability of P or $1 – P$,

respectively. A segment was defined as a series of non-branching nodes terminating at a

branching (or terminal) node. The number of nodes in a segment is a random variable that is

assumed to be geometrically distributed with the probability of success P (equal to the

branching probability). To achieve the target number of total nodes in a network, individual

segments were formed by consecutively drawing their number of nodes from the geometric

distribution. As a statistical reformulation of Horton's laws, a geometric distribution has been

used to describe the segment lengths (e.g., Peckham & Gupta, 1999). Before merging the

segments to create a river network, the whole drawing process was repeated if a non-equal

number of total nodes or a non-odd number of segments was attained. To create the river

network, these segments were put together as a hierarchically merged pool in three steps

(Figure S3). For Step 1, one segment, consisting of one or more nodes, was randomly selected

as the root, and its upstream end was merged with the downstream end of two randomly

selected segments. From this procedure, the semi-complete network had two unmerged

upstream ends each for the next possible merger. For Step 2 Two more segments were then

randomly selected, and their downstream ends were merged together to the random draw of

one of two (or even more at subsequent steps) unmerged upstream ends of the semi-complete

network. For Step 3, Step 2 was repeated until there were no available segments left in the

pool.

Simulation of river-branching influences

We conducted stochastic simulations to illustrate the genetic differentiation and diversity at the

metapopulation scale under river branching. Our metapopulation genetic model for each

species with the Bayesian-median estimates of the parameters added was then used to simulate

allele frequencies at loci in local populations in each of the 1,000 random river networks (see

the above section '*Creation of virtual river networks*'). The spatial genetic structure was

summarized to the global G_{ST} to be the metapopulation's genetic divergence. In addition, the

metapopulation's genetic diversity was calculated as global H_t (Nei, 1973). We performed

this simulation using the R packages 'stats' and 'base' (R Core Team, 2018).

We then built a regression model based on gradient boosting (GB) (Appendix S1) for each

of the four macroinvertebrate species, which identified the importance of the following spatial

features: (1) the fraction of any two local populations being streamflow-disconnected in all

possible combinations; (2) the mean watercourse distance between local populations under

different levels of river branching; and (3) the metapopulation size (i.e., number of local

populations) for genetic divergence (global G_{ST}). These features were considered to be

summarised aspects of complex spatial configuration of habitat network.

Through further simulations on the same set of river networks, we illustrated how the

downstream and upstream dispersal-related parameters of the scales (c_D and c_U, respectively)

influence river-branching effects on the genetic divergence of the metapopulation (global G_{ST})

and genetic diversity (global H_t). We considered 3×3 (nine total) combinations of the scale

parameters, with each having three values, which were the maximum, median, and minimum

values of range of their Bayesian pooled estimates where the posterior median of both c_D and

c_U across the four macroinvertebrate species were pooled together. Because of the shared pool,

the maximum, median, or minimum values set for the two scale parameters in the

combinations are the same. We set the metapopulation's logit-transformed frequency of allele

"1" at each locus to zero during each simulation. The number of loci was set as the median

across the four macroinvertebrate species. In addition, we set the variance parameter (σ_D^2 or

σ_U^2) to the median value of their Bayesian pooled estimates, with the two variance parameters

having the same values. We replicated the nine combinations in each of the 1,000 random river

networks.

Results

Metapopulation genetic modelling

When our Bayesian model was fitted to the empirical genetic data in the Natori and Nanakita

catchments, the R^2 values derived from the residuals (i.e., the differences between the observed

and predicted numbers of type '1' alleles at a locus from the number of alleles observed in a

local population; see Equation 1) were 0.98, 0.97, 0.97, and 0.93 for *H. orientalis*, *S.*

marmorata, *H. albicephala* and *E. japonica*, respectively (Figure S4).

The metapopulation allele frequencies at loci are species-specific, and the variation of

allele frequency was greater in the widespread *H. orientalis* and *S. marmorata* than in the other

two species that had narrower habitat distributions (Figure S1). Pairwise genetic differences

between empirical local populations tended to increase with their watercourse distances

throughout the four macroinvertebrate species (Figure 1). Despite substantial variation in the

scale parameter that amplified the isolating effect of distance across study species (Figure 2),

there was a consistent decline in the genetic correlation between populations (the covariance

Ω_{ij} divided by the variance $\sigma_D^2 + \sigma_U^2$; see Equation 4) with the increasing distance between

local populations being obvious for the three caddisfly species but not the mayfly having lower

value of c_D (Figure S5).

Simulation of river-branching influences

The increasing branching probability of river networks caused changes in the observed

spatial features (Figure 3). For example, the fraction of any two local populations being

streamflow-disconnected (e.g., occurring in different tributaries) across the metapopulations

occurs at high rates in the heavily branched river networks. However, we found shorter

watercourse distances occurred between local populations in river networks with higher

branching probability. As the size of the virtual network increases, the metapopulation size (i.e.,

the total number of local populations of the network) increases as do the values of both spatial

configurations under the same level of river branching.

Two spatial features (watercourse distance and fraction of flow-disconnected sites; Figure

3) summarized the spatial configuration of habitat network and were linked to the genetic

differentiation of metapopulations (G_{ST}; Figure 4) and total genetic diversity (H_t; Figure S6).

However, species-specific responses to the influence of river branching were evident. For

example, increased branching probability decreased the genetic divergence and diversity of the

metapopulation for one caddisfly (*H. albicephala*), but the mayfly *E. japonica* showed the

opposite response (higher genetic divergence) before a branching threshold. According to the

GB modelling results, the relative importance of streamflow-disconnected habitats, compared

to the landscape connectivity via a shorter watercourse distance, was higher for the mayfly than

in all the caddisflies examined (Table 1).

Branching complexity has various impacts on genetic divergence and diversity, which is

determined by the relative values of the upstream and downstream scale-parameters in a

genetic covariation function (Equation 4). Stronger dispersal tendency can be associated with a

lower value of the scale parameter in either direction, which will result in weak isolation by

watercourse distance within a network. When species dispersal was upstream-biased (c_U <

c_D), branching complexity enhanced genetic homogenization among local populations within a

metapopulation. In addition, lower genetic divergence and diversity are related to lower

downstream-biased and higher upstream-biased dispersals (relative higher c_D and lower c_U,

respectively; Figures 5 and S7).

Discussion

Using simulation modelling in virtual networks, we explored the integrated role of landscape

architecture and species dispersal asymmetry in shaping genetic divergence at the

metapopulation scale. We compared the spatial patterns of genetic differentiation of four

macroinvertebrate species in river networks based on our Bayesian model that explicitly

accounted for the effects of evolutionary processes among components of the metapopulations

on their spatial genetic-structure. Our simulations indicated that greater landscape connectivity

(via shortened watercourse distance) and higher distal isolation (e.g., in headwater streams)

simultaneously occur in more-branched river networks and have countervailing influences on

genetic divergence.

The two simplified, correlated aspects of complex networks (Figure S8) can have

nonequal importance for different species and this difference can be linked to varying genetic

divergence in the metapopulation. These results, derived from the simulation of virtual

networks, may not directly apply to all empirical systems but they can provide hypothesized

expectations for the relationship between genetic divergence and network branching. We

suggest that species dispersal can cause varying levels of overall genetic differentiation in river

networks having increased river branching, and can have either different or even opposite

effects. We believe that this result can provide insights into other complex networks (e.g.,

highly fragmented landscapes or those with corridors via ocean and atmospheric circulation).

In addition, these results highlight the fundamental importance of considering dispersal traits of

individual species, which can result in different contributions to genetic connectivity, for the

successful management of ecological corridors. Our theoretical evidence suggests the value of

conducting further studies using empirical comparisons across multiple natural networks based

on hypotheses generated from these modelling efforts.

Branching complexity and genetic divergence

The role of network branching has been documented, to some extent, in genetic studies of

riverscape conducted in situations where higher genetic diversity is observed in downstream

populations than in upstream ones (Paz-Vinas et al., 2015). Also, greater river branching can

increase the differences between such populations (Thomaz et al., 2016). In dendritic networks,

our simulation results showed a species-dependent change in global genetic-differentiation

levels that occur with increases in network complexity. This finding was also previously

observed for two sympatric salmonid species that were found to have remarkably different

spawning locations, mating systems, and population sizes, and for which biological traits

mediated the influences of riverscape features that shape their dispersal and genetic divergence

(Whiteley, Spruell, & Allendorf, 2004).

Little or no gene flow can occur with high network branching. This pattern of gene flow is

caused by strong genetic isolation from an increase in the number of distal populations (e.g., in

headwaters) when the same number of nodes in a habitat network is present. This pattern can

generally be observed for some riverine species with high or intermediate levels of

downstream-biased vagility, such as fish and molluscs (Osborne, Perkin, Gido, & Turner, 2014;

Pilger et al., 2017; Terui et al., 2014). In one, early theoretical study that did not consider

dispersal asymmetry (which would be analogous to having equal downstream and upstream

dispersals in our study), the dendritic network structure was also documented to promote low

genetic distances under conditions of high connectivity (Labonne et al., 2008). However, our

results show that the different downstream and upstream gene flows considered in our model

can act together to generate similar or even different relationships.

Importance of species dispersal

The architecture of river networks can be an important extrinsic factor in explaining the

observed and simulated genetic patterns in our study, but there was high variation among the

species with different dispersal traits in our study. By adopting our mechanistic model that was

fitted by empirical data on macroinvertebrate species with flying adult-stages, our simulations

reveal that their better dispersal abilities can overcome catchment constraints, leading to low

downstream-biased asymmetry. The opposite (negative) influence occurs under increased river

branching. In addition, our comprehensive consideration of various branching patterns in

river-network topologies in our simulations demonstrate the existence of opposing influences

that can co-occur under branching complexity.

In the river systems we examined, asymmetric (either downstream- or upstream-biased)

dispersal could determine the effect of branching complexity on genetic divergence of a

metapopulation. In our model, the scale parameters for upstream- and downstream-dispersal

tendencies (in the genetic covariation function; Equation 4) can be linked to dispersal

asymmetry. These two factors together provide a mechanism behind the countervailing

influences of river branching on the genetic divergence of metapopulations. Stronger dispersal

can be associated with a lower value of the scale parameter either in downstream or upstream

direction, which will result in weak isolation by distance within a metapopulation.

Our results suggested that, in the mayfly species examined (*E. japonica*) having high

downstream-biased gene flow (based on a higher value of the scale parameter for upstream

than for downstream dispersal), river branching has a positive influence on genetic divergence.

This phenomenon can be attributed to a higher number of isolated distal branches in river

networks (e.g., more headwaters) occurring under this dispersal asymmetry, as typically shown

in fish (e.g., Pilger et al., 2017). Larvae of mayflies also are susceptible to high levels of drift

during times of increased river flow and therefore have high potential to be strong downstream

dispersers (Nukazawa et al., 2017).

In our system, an opposite (negative) influence of the branching network on the three

caddisfly species with dispersal symmetry or even upstream-biased dispersal (based on the

scale parameter for upstream being similar to or higher than that for downstream dispersal,

respectively; Figures 2 and 4) was evident. This result could be attributable to their flying

adults generally providing a wide dispersal range and strong terrestrial movement (e.g., at least

in the upstream direction, which would compensate for downstream drift dispersal). This

upstream movement dampened the isolation between distal branches in river networks for

caddisflies compared to mayflies (and although not examined, likely stoneflies as well), which

exhibit restricted distributions to areas very close to their sources of emergence in a stream

(Winterbourn et al., 2007).

Unexplained genetic variations

The Bayesian fitting process is more likely to have failed in retrieving the assumed value of

variance parameters in the scenario of equal values of scale parameters than in other scenarios.

These stochastic effects are derived from evolutionary processes modelled by autocovariances

related to upstream and downstream dispersal along the habitat networks. The deviations from

stochasticity are not significant when compared to the large difference between the actions of

the autocovariances on the genetic structures. Thus, the Bayesian estimations are more easily

confounded when both autocovariances have equal or similar influence on the stochastic

processes. In general, more sampling of sites in the river network and additional collections of

individuals from local populations would result in higher constraints for parameter estimations,

which can decrease the stochastic effects. In river systems, this issue can occur from aquatic

insects having different aerial-dispersal abilities and may not apply in other riverine species

(e.g., fish and mollusc species) having downstream-biased dispersal.

463   Some bias in our modelling results could result from not considering overland dispersal,

particularly when asymmetric dispersals along the watercourse in our model explain a large

part of the genetic variation at a metapopulation scale. Overland dispersal across tributaries or

catchments has been demonstrated to influence genetic differentiation (Alp et al., 2012;

Chaput-Bardy, Fleurant, Lemaire, & Secondi, 2009). However, this type of dispersal may not

have made an important contribution in our empirical system when compared with in-stream

dispersal. Overland movement can be particularly important to strong fliers like damselflies

and riverine birds, e.g., dippers (Hernández et al., 2012). However, aquatic insects with weaker

flying strength (e.g., mayflies or caddisflies) have larger longitudinal tendency to use stream

corridors than to use lateral areas away from the stream channel (Petersen et al., 2004).

However, empirical genetic-evidence indicates that in-stream dispersal was the main dispersal

pathway (Chaput-Bardy, Lemaire, Picard, & Secondi, 2008).

Overview and management implications

From a conservation and management perspective that attempts to include considerations of

the spatial genetic-structure of metapopulations (e.g., Luque, Saura, & Fortin, 2012), an

understanding of the branching structure's role in driving metapopulation genetic divergence is

critical. Dispersal can dictate differences in genetic diversification in landscapes (Medina,

Cooke, & Ord, 2018) and our predictive modelling based on asymmetric dispersals across

networks sheds light on better management practices. For example, in the context of managing

native or even invasive species (Chaput-Bardy, Alcala, Secondi, & Vuilleumier, 2017), our

simulation results provide insight into the evolutionary importance of dispersal abilities and

modes, suggesting that these intrinsic factors should be considered in decision-making

processes where one management strategy does not fit all species (Figure S9). For example, the

same management and conservation practices can produce different or even opposite results for

species with varying levels of asymmetric gene flow and genetic drift (e.g., in dendritic river

systems). Moreover, by adopting a Bayesian fitted mechanistic framework into local

landscapes, simulations in virtual networks could result in insights gained via experimental or

empirical studies that can assist managers build specific strategies for different regions. For

example, the potential genetic influences from past or planned human alterations of river

systems (e.g., dam construction, loss of headwater habitats from pollution, branches diverted

for agriculture, or creation of flood-bypass channels) may be incorporated into assessments of

branching complexity. In doing the assessments, ecological connections rather than purely

hydrological linkages are considered to be the elements forming the structure of habitat

networks. In addition, this approach can be considered for incorporation into management

strategies (e.g., in-stream barrier removal and habitat restoration placed in the context of the

river network) and anticipated responses to environmental changes (e.g., drying out of

segments due to reduced precipitation under climate change).

Data Accessibility Statement: No new genetic data were created. AFLP genetic data,

including neutral and non-neutral loci, are available from the Dryad Digital Repository at:

http://dx.doi.org/10.5061/dryad.82br3. Stan code is uploaded as a separate supplemental file.

Literature cited

Alp, M., Keller, I., Westram, A. M., & Robinson, C. T. (2012). How river structure and

biological traits influence gene flow: a population genetic study of two stream

invertebrates with differing dispersal abilities. *Freshwater Biology, 57*(5), 969-981.

doi:10.1111/j.1365-2427.2012.02758.x

Andrews, C. A. (2010). Natural selection, genetic drift, and gene flow do not act in isolation in

natural populations. *Nature Education Knowledge, 3*(10), 5.

Bradburd, G. S., Ralph, P. L., & Coop, G. M. (2013). Disentangling the effects of geographic

and ecological isolation on genetic differentiation. *Evolution, 67*(11), 3258-3273.

doi:10.1111/evo.12193

Chaput-Bardy, A., Alcala, N., Secondi, J., & Vuilleumier, S. (2017). Network analysis for

species management in rivers networks: Application to the Loire River. *Biological*

Conservation, 210, 26-36. doi:https://doi.org/10.1016/j.biocon.2017.04.003

Chaput-Bardy, A., Fleurant, C., Lemaire, C., & Secondi, J. (2009). Modelling the effect of

in-stream and overland dispersal on gene flow in river networks. *Ecological Modelling,*

220(24), 3589-3598. doi:https://doi.org/10.1016/j.ecolmodel.2009.06.027

Chaput-Bardy, A., Lemaire, C., Picard, D., & Secondi, J. (2008). In-stream and overland

dispersal across a river network influences gene flow in a freshwater insect, *Calopteryx*

splendens. *Molecular Ecology, 17*(15), 3496-3505.

doi:10.1111/j.1365-294X.2008.03856.x

Fourcade, Y., Chaput-Bardy, A., Secondi, J., Fleurant, C., & Lemaire, C. (2013). Is local

selection so widespread in river organisms? Fractal geometry of river networks leads to

high bias in outlier detection. *Molecular Ecology, 22*(8), 2065-2073.

doi:doi:10.1111/mec.12158

Green, D., Klomp, N., Rimmington, G., & Sadedin, S. (2006). *Complexity in Landscape*

Ecology. Dordrecht, The Netherlands: Springer.

Guillot, G., Vitalis, R., Rouzic, A. l., & Gautier, M. (2014). Detecting correlation between

allele frequencies and environmental variables as a signature of selection. A fast

computational approach for genome-wide studies. *Spatial Statistics, 8*, 145-155.

doi:10.1016/j.spasta.2013.08.001

Hernández, M. Á., Campos, F., Santamaría, T., Corrales, L., Rojo, M. A., & Dias, S. (2012).

Genetic differences among iberian white-throated dipper *Cinclus cinclus* populations

based on the cytochrome b sequence. *Ardeola, 59*(1), 111-122.

doi:10.13157/arla.59.1.2012.111

Junker, J., Peter, A., Wagner, C. E., Mwaiko, S., Germann, B., Seehausen, O., & Keller, I.

(2012). River fragmentation increases localized population genetic structure and

enhances asymmetry of dispersal in bullhead (Cottus gobio). *Conservation Genetics,*

13(2), 545-556. doi:10.1007/s10592-011-0306-x

Kawecki, T. J., & Holt, R. D. (2002). Evolutionary consequences of asymmetric dispersal rates.

American Naturalist, 160(3), 333-347. doi:10.1086/341519

Labonne, J., Ravigné, V., Parisi, B., & Gaucherel, C. (2008). Linking dendritic network

structures to population demogenetics: The downside of connectivity. *Oikos, 117*(10),

1479-1490. doi:10.1111/j.0030-1299.2008.16976.x

Luque, S., Saura, S., & Fortin, M.-J. (2012). Landscape connectivity analysis for conservation:

insights from combining new methods with ecological and genetic data. *Landscape*

Ecology, 27(2), 153-157. doi:10.1007/s10980-011-9700-5

Ma, C., Shen, Y., Bearup, D., Fagan, W. F., & Liao, J. (2020). Spatial variation in branch size

promotes metapopulation persistence in dendritic river networks. *Freshwater Biology,*

65(3), 426-434. doi:10.1111/fwb.13435

Martins, K., Kimura, R. K., Francisconi, A. F., Gezan, S., Kainer, K., & Christianini, A. V.

(2016). The role of very small fragments in conserving genetic diversity of a common

tree in a hyper fragmented Brazilian Atlantic forest landscape. *Conservation Genetics,*

17(3), 509-520. doi:10.1007/s10592-015-0800-7

McRae, B. H. (2006). Isolation by resistance. *Evolution, 60*(8), 1551-1561.

doi:10.1111/j.0014-3820.2006.tb00500.x

Medina, I., Cooke, G. M., & Ord, T. J. (2018). Walk, swim or fly? Locomotor mode predicts

genetic differentiation in vertebrates. *Ecology Letters, 21*(5), 638-645.

doi:10.1111/ele.12930

Morrissey, Michael B., & de Kerckhove, Derrick T. (2009). The maintenance of genetic

variation due to asymmetric gene flow in dendritic metapopulations. *American*

Naturalist, 174(6), 875-889. doi:10.1086/648311

Nei, M. (1973). Analysis of gene diversity in subdivided populations. *Proceedings of the*

National Academy of Sciences of the United States of America, 70(12), 3321-3323.

Nukazawa, K., Kazama, S., & Watanabe, K. (2015). A hydrothermal simulation approach to

modelling spatial patterns of adaptive genetic variation in four stream insects. *Journal*

of Biogeography, 42(1), 103-113. doi:10.1111/jbi.12392

Nukazawa, K., Kazama, S., & Watanabe, K. (2017). Catchment-scale modeling of riverine

species diversity using hydrological simulation: application to tests of species-genetic

diversity correlation. *Ecohydrology, 10*(1), e1778. doi:10.1002/eco.1778

Osborne, M. J., Perkin, J. S., Gido, K. B., & Turner, T. F. (2014). Comparative riverscape

genetics reveals reservoirs of genetic diversity for conservation and restoration of Great

Plains fishes. *Molecular Ecology, 23*(23), 5663-5679. doi:10.1111/mec.12970

Paz-Vinas, I., & Blanchet, S. (2015). Dendritic connectivity shapes spatial patterns of genetic

diversity: a simulation-based study. *Journal of Evolutionary Biology, 28*(4), 986-994.

doi:10.1111/jeb.12626

Paz-Vinas, I., Loot, G., Stevens, V. M., & Blanchet, S. (2015). Evolutionary processes driving

spatial patterns of intraspecific genetic diversity in river ecosystems. *Molecular*

Ecology, 24(18), 4586-4604. doi:10.1111/mec.13345

Paz-Vinas, I., Quéméré, E., Chikhi, L., Loot, G., & Blanchet, S. (2013). The demographic

history of populations experiencing asymmetric gene flow: combining simulated and

empirical data. *Molecular Ecology, 22*(12), 3279-3291. doi:10.1111/mec.12321

Peckham, S. D., & Gupta, V. K. (1999). A reformulation of Horton's Laws for large river

networks in terms of statistical self-similarity. *Water Resources Research, 35*(9),

2763-2777. doi:10.1029/1999wr900154

Petersen, I., Masters, Z., Hildrew, A. G., & Ormerod, S. J. (2004). Dispersal of adult aquatic

insects in catchments of differing land use. *Journal of Applied Ecology, 41*(5), 934-950.

doi:10.1111/j.0021-8901.2004.00942.x

Phillipsen, I. C., Kirk, E. H., Bogan, M. T., Mims, M. C., Olden, J. D., & Lytle, D. A. (2015).

Dispersal ability and habitat requirements determine landscape-level genetic patterns in

desert aquatic insects. *Molecular Ecology, 24*(1), 54-69. doi:10.1111/mec.13003

Phillipsen, I. C., & Lytle, D. A. (2013). Aquatic insects in a sea of desert: population genetic

structure is shaped by limited dispersal in a naturally fragmented landscape. *Ecography,*

36(6), 731-743. doi:10.1111/j.1600-0587.2012.00002.x

Pilger, T. J., Gido, K. B., Propst, D. L., Whitney, J. E., & Turner, T. F. (2017). River network

architecture, genetic effective size and distributional patterns predict differences in

genetic structure across species in a dryland stream fish community. *Molecular Ecology,*

26(10), 2687-2697. doi:10.1111/mec.14079

Prunier, J. G., Dubut, V., Loot, G., Tudesque, L., & Blanchet, S. (2018). The relative

contribution of river network structure and anthropogenic stressors to spatial patterns of

genetic diversity in two freshwater fishes: A multiple-stressors approach. *Freshwater*

Biology, 63(1), 6-21. doi:doi:10.1111/fwb.13034

R Core Team. (2018). *R: A Language and Environment for Statistical Computing*. Vienna,

Austria. https://www.R-project.org/: R Foundation for Statistical Computing.

Stan Development Team. (2018). *Stan Modeling Language Users Guide and Reference Manual,*

Version 2.18.0. http://mc-stan.org. In.

Stan Development Team. (2019). RStan: the R interface to Stan. R package version 2.19.1.

http://mc-stan.org/. Retrieved from http://mc-stan.org/rstan.html

Stoll, S., Breyer, P., Tonkin, J. D., Früh, D., & Haase, P. (2016). Scale-dependent effects of

river habitat quality on benthic invertebrate communities--Implications for stream

restoration practice. *Science of the Total Environment, 553*, 495-503.

doi:10.1016/j.scitotenv.2016.02.126

Terui, A., Ishiyama, N., Urabe, H., Ono, S., Finlay, J. C., & Nakamura, F. (2018).

Metapopulation stability in branching river networks. *Proceedings of the National*

Academy of Sciences of the United States of America, 115(26), E5963-E5969.

doi:10.1073/pnas.1800060115

Terui, A., Miyazaki, Y., Yoshioka, A., Kaifu, K., Matsuzaki, S.-I. S., & Washitani, I. (2014).

Asymmetric dispersal structures a riverine metapopulation of the freshwater pearl

mussel *Margaritifera laevis*. *Ecology and Evolution, 4*(15), 3004-3014.

doi:10.1002/ece3.1135

Thomaz, A. T., Christie, M. R., & Knowles, L. L. (2016). The architecture of river networks

can drive the evolutionary dynamics of aquatic populations. *Evolution, 70*(3), 731-739.

doi:10.1111/evo.12883

Ver Hoef, J. M., & Peterson, E. E. (2010). A moving average approach for spatial statistical

models of stream networks. *Journal of the American Statistical Association, 105*(489),

6-18. doi:10.1198/jasa.2009.ap08248

Watanabe, K., Kazama, S., Omura, T., & Monaghan, M. T. (2014). Adaptive genetic

divergence along narrow environmental gradients in four stream insects. *PLoS ONE,*

9(3), e93055. doi:10.1371/journal.pone.0093055

Whiteley, A. R., Spruell, P., & Allendorf, F. W. (2004). Ecological and life history

characteristics predict population genetic divergence of two salmonids in the same

landscape. *Molecular Ecology, 13*(12), 3675-3688.

doi:10.1111/j.1365-294X.2004.02365.x

Winterbourn, M. J., Chadderton, W. L., Entrekin, S. A., Tank, J. L., & Harding, J. S. (2007).

Distribution and dispersal of adult stream insects in a heterogeneous montane

environment. *Fundamental and Applied Limnology, 168*(2), 127-135.

doi:10.1127/1863-9135/2007/0168-0127

Yeakel, J. D., Moore, J. W., Guimarães Jr., P. R., & de Aguiar, M. A. M. (2014).

Synchronisation and stability in river metapopulation networks. *Ecology Letters, 17*(3),

273-283. doi:10.1111/ele.12228

Zurell, D., Berger, U., Sarmento Cabral, J., Jeltsch, F., Meynard, C., Münkemüller, T., . . .

Grimm, V. (2010). The virtual ecologist approach: simulating data and observers. *Oikos,*

221, 98-105.

Table 1. Performance measures and relative importance of predictors in gradient-boosting models for *Ephemera japonica* (EJ), *Stenopsyche marmorata*

(SM), *Hydropsyche orientalis* (HO), and *Hydropsyche albicephala* (HA). The simulated genetic divergence of the metapopulation (global G_{ST}) is the

response variable and fraction for any two streamflow-disconnected local populations, the mean watercourse distance between populations, and

metapopulation size, all of which are predictor variables.

Species	Number of trees	Model performance		Relative importance (%)		
		RMSE	R^2	Streamflow-disconnected fraction	Watercourse distance	Metapopulation size
EJ	3800	0.03	0.95	84.2	9.1	6.7
HA	950	0.05	0.89	1.4	95.9	2.7
HO	600	0.05	0.59	4.0	82.6	13.5
SM	600	0.04	0.50	6.6	26.0	67.4

   Figure legends

Figure 1. Study catchment area and distribution of (a) *Ephemera japonica* (EJ), (b)

Hydropsyche albicephala (HA), (c) *Hydropsyche orientalis* (HO) and (d) *Stenopsyche*

marmorata (SM), and the predicted genetic divergence of their local populations on the basis

of nMDS (nonmetric multidimensional scaling) of the modeled pairwise G_{ST} between river

grid squares (500×500 m^2) in northeastern Japan (see Appendix S1). Gray squares and color

points indicate sampling sites and predictions, respectively.

Figure 2. Estimated values of (a) variances (σ_D^2 and σ_U^2) and (b) scales (c_D and c_U) of

covariance function (see Equation 4) with the posterior distribution in our Bayesian modeling

for *Ephemera japonica* (EJ), *Hydropsyche albicephala* (HA), *Hydropsyche orientalis* (HO),

and *Stenopsyche marmorata* (SM).

Figure 3. Theoretical predictions for relationships of (a) mean watercourse distance between

populations or (b) fraction of any two streamflow-disconnected populations (e.g., in

headwaters) in all combinations having branching complexity under differential

metapopulation sizes (range: 100 to 500 local populations).

Figure 4. Theoretical predictions for relationships between metapopulation genetic divergence

(global $G_{ST} = GG_{ST}$) and branching complexity under different metapopulation sizes (range:

100 to 500 local populations) for (a) *Ephemera japonica* (EJ), (b) *Hydropsyche albicephala*

(HA), (c) *Hydropsyche orientalis* (HO), and (d) *Stenopsyche marmorata* (SM). Relative GG_{ST}

$= (GG_{ST} - \text{minimum } GG_{ST}) / (\text{maximum } GG_{ST} - \text{minimum } GG_{ST})$ across the range for each

species.

Figure 5. Theoretical predictions of relationships between metapopulation genetic divergence

(global $G_{ST} = GG_{ST}$) and branching complexity under differential metapopulation sizes (range:

100 to 500 local populations) for combinations of dispersal-related scale parameters in a

genetic covariation function (Equation 4), including (a, e and i) $c_D = c_U$, (b, c and f) $c_D < c_U$,

and (d, g and h) $c_D > c_U$. The three values of each parameter are the maximum, median, and

minimum values of range of Bayesian pooled estimates when the posterior median of both c_D

and c_U across the four macroinvertebrate species were pooled. Relative $GG_{ST} = (GG_{ST} - $

minimum GG_{ST}) / (maximum GG_{ST} - minimum GG_{ST}) across the range for each parameter

combination.

Figure 1

Figure 2

Figure 3

695   **Figure 4**

697 **Figure 5**