Pharmacodynamic inoculum effect from the perspective of bacterial population modeling

Desiree Y. Baeder[1] and Roland R. Regoes[1]

[1]Institute of Integrative Biology, ETH Zurich, Universitätsstrasse 16, 8092 Zurich, Switzerland

1 Abstract

Efficacy assessment of antimicrobials is essential, both to determine the best clinical use of the antimicrobial and as input for predictive mathematical modeling. The pharmacodynamic (PD) function is a tool to describe antimicrobial efficacy and can be seen as an extension of the commonly used MIC. While the PD function describes the efficacy of a given dose of antimicrobials, it is based on one bacterial inoculum size only. Therefore, the PD function does not inform us about the change in efficacy due to changes in the inoculum size, also known as the inoculum effect. Here, we used mathematical modeling to quantify the inoculum effect in terms of PD parameters. In particular, we used the multi-hit model that describes the mechanism of action of antimicrobials: When a certain number of antimicrobial molecules have hit a bacterial cell, it dies. This framework allowed us to examine the effect of reversible binding and enzymatic degradation of antimicrobials on the pharmacodynamics. A change in bacterial inoculum size resulted in a change of the PD parameter A_{50}, MIC_{PD}, and therefore κ due to reversible binding. We determined an extended PD function that captured the inoculum effect as change in the PD parameters. When we allowed for degradation of antimicrobials by bacterial enzymes, the inoculum effect was intensified. The extended PD function could mimic long term population dynamics based on the multi-hit model with reversible binding only,

but deviated from long-term predictions based on a multi-hit model including degradation. We then used the multi-hit framework to estimate outcomes of competition experiments with a sensitive and resistant strain and compared it to predictions of the PD function. When we do not take the inoculum effect into account, our simulations underestimated the ability of the sensitive population to survive for a given PK regime and the competitiveness of the sensitive strain against the resistant strain. Our work emphasizes that the PD function — and in general any efficacy measure — should, at least, include information about the inoculum size and, ideally, account for the inoculum effect.

Introduction

Efficacy of antimicrobials against bacteria is quantified using time-kill curves and is subsequently captured in so-called pharmacodynamic (PD) functions, also known as E_{max} and Zhi models (*1–3*). A time-kill curve captures how a bacterial population changes over time in presence of a given concentration of an antimicrobial (fig. 1a). During exponential growth, time-kill curves are approximately linear on a logarithmic scale. Thus, the slopes of these log-linear time-kill curves describe the change of the bacterial population over time and are a direct measure of the bacterial net growth rate. The PD function returns the net growth rate for any given antimicrobial concentration. The parameters of the PD function are estimated based on several time-kill curves with the same initial bacterial population size (bacterial inoculum) (*1, 4–8*).

A more widespread but less informative way to express efficacy of antimicrobials is the minimum inhibitory concentration (MIC). The MIC is the lowest drug concentration that prevents visible bacterial growth (fig. 1b). For concentration-independent antimicrobials (steep PD curve), the MIC is comparable to the antimicrobial concentration that results in a time-kill curve with the

slope of zero (*8*). In this case, one of the four parameters of the PD function, which we call MIC_{PD}, is equivalent to the MIC. Although it is recognized that the MIC might change due to change in bacterial inoculum size (see for example the MIC protocol by Wiegand *et al.* (*9*)), both the CLSI and the EUCAST recommendation to measure the MIC is based on a standard bacterial inoculum of 5×10^5 CFU/ml only (*10, 11*). Thus, as in the case of time-kill curves, standardization of the inoculum neglects any effect on efficacy of antimicrobials due to changes in the bacterial inoculum.

Both experimental and theoretical work has put forward hypotheses explaining why the inoculum size matters when determining antimicrobial efficacy. In our opinion, the most intuitive explanation is that the amount of antimicrobials is finite in the system and a decrease in the amount of free antimicrobials changes the efficacy of the antimicrobial (*12–14*). One reason why the amount of antimicrobials decreases is binding of antimicrobial molecules to targets. Assuming mass action kinetics, the number of free antimicrobial molecules is reduced proportional to bacterial cell density, which results in changes in the efficacy of the antimicrobial. In addition to this binding kinetic effect, bacteria can produce extracellular enzymes that further decrease the concentration of antimicrobials (*15–20*). Some bacteria, for example, produce β-lactamase, which is able to degrade β-lactam antibiotics. Also AMPs, for example LL-37, can be degraded by proteases produced by bacteria (*21–23*). The higher the bacterial population, the more enzymes are produced and thus the higher the change in antimicrobial efficacy.

Changes in efficacy of antibacterial drugs caused by changing the initial bacterial inoculum size are summarized under the term *inoculum effect* (fig. 1). The inoculum effect was first described in the early 1940s (*18*). Since then, the inoculum effect has been measured experimentally for a variety of combinations

of bacterial species and antibiotics (13, 15, 20, 24–26). Recently, the inoculum effect became of interest in the field of antimicrobial peptides (AMPs) (27, 28). It is common to determine the inoculum effect in terms of the change in the MIC with inoculum size (13, 15, 20, 24, 25, 27). To our knowledge, the inoculum effect has not yet been determined using the more comprehensive PD function approach. In addition to its experimental quantification in the wet lab, the inoculum effect has been explored with mathematical models. These models either comprise a mechanistic description of the interaction between antimicrobial molecules and bacterial cells (12, 14, 26, 29, 30) or are based directly on MIC measurements (13). In the latter case, the PD function was extended to incorporate the inoculum effect for one of the four parameters based on MIC measurements.

Here we use mathematical modeling as a tool to explore how bacterial inoculum size affects all four parameters of the PD function. The advantage of employing mathematical models is the possibility to determine the efficacy of an antimicrobial for various bacterial inoculum sizes and to explore mechanisms of the inoculum effect individually and in combination, which would be challenging experimentally. We adopt a mathematical framework called the multi-hit model, which is based on the idea that antimicrobial molecules hit, i.e. bind to, cellular targets (31). In this approach, complex chemical reactions and molecular pathways are summarized in a simple mechanistic structure. This framework enables us to simulate bacterial population dynamics, from which we can infer the pharmacodynamic curve. We already used the multi-hit model in previous work (32) to describe the action of AMPs and a similar, albeit more antibiotic-specific, approach was used to model classic reaction kinetics of antibiotics (12).

A mechanistic explanation for the inoculum effect has already been provided

by Abel zur Wiesch *et al.* (*12*). In their model, binding of molecules decreases the pool of free antimicrobials and thus the antimicrobial efficacy. Here, we use the multi-hit model to extend this analysis and apply the results to the PD function. With the multi-hit framework, we are able to ask how the changes in the inoculum size affects the pharmacokinetics of antimicrobials and how this in turn affects the PD curve. We show that reversible binding alone can explain the inoculum effect. Further, we quantify the inoculum effect in terms of change in PD parameter values with increasing inoculum size. In a next step, we analyze how the PD curve changes due to adding degradation of antimicrobial molecules by bacterial enzymes and illustrate how well the PD function predicts long-term population dynamics. Last, we apply our findings to the population dynamics of competing bacterial strains and predict how the inoculum size influences competition outcomes when a resistant strain is introduced.

1   Methods

Multi-hit model

We used the multi-hit model introduced by Hedges (*31*) to explore bacterial population dynamics *in silico* in presence of antimicrobial peptides (AMPs). We have already applied a simplified version of the multi-hit model in previous work (*32*) and a similar approach has been introduced by Abel zur Wiesch *et al.* (*12*) to describe effects of antibiotics.

The multi-hit model describes a population N_{tot}. In the presence of antimicrobials, bacterial cells are hit by antimicrobial molecules. A hit is the mathematical simplification of adsorption and subsequent downstream processes connected to binding of one antimicrobial molecule to a bacterial target site and is described with the rate α. The parameter μ is the detachment of an antimi-

Figure 1: Inoculum effect changes efficacy of antimicrobials. Two common methods to quantify the efficacy of antimicrobials are (a) time-kill curves and (b) MIC measurements. (a) Time-kill curves illustrate the change in population size over time in presence of antimicrobials. The bacterial inoculum is the bacterial population size N at $t_0 = 0h$ ($N_{inoc} = N(t = 0)$). The slope of the curve results from fitting a log-linear function to the population dynamics. The slope of time-kill curve representing dynamics of bacteria starting at a high inoculum size (orange) is higher than dynamics of bacteria starting at a low inoculum (blue). In absence of an inoculum effect, the slope is independent of the inoculum (dashed orange line). (b) Alternative to measuring time-kill curves, efficacy can also be determined with the MIC. An inoculum effect is observed if increasing the inoculum size from low (blue) to high (orange) increases the MIC.

126 crobial from the bacterial target. Bacteria that have i antimicrobial molecules

127 adsorbed are categorized in class N_i. The number of absorbed molecules varies

128 from cell to cell, with $0 \leq i \leq n$. Thus, we have a total of $n + 1$ classes and

129 $N_{tot} = \sum_{i=0}^{n} N_i$. The rate of change of each subpopulation is described with

130 the following ordinary differential equation:

$$
\begin{aligned}
\frac{dN_i}{dt} = {} & \alpha_{i-1}AN_{i-1} - \alpha_i AN_i + \\
& \mu_{i+1}(i+1)N_{i+1} - \mu_i i N_i + \\
& \Omega(N_i) - d_i N_i
\end{aligned}
\tag{1}
$$

Each class N_i gains bacterial cells from the class N_{i-1} with the term $\alpha_{i-1}AN_{i-1}$.

Here, we assume that finding the bacterial cell is the rate limiting step for an

antimicrobial molecule (33). This is appropriate if the number of targets are

in abundance, e.g. when antimicrobial peptides adsorb to the cell membrane of

bacterial cells. For antimicrobials that act within the cell, a hit rate dependent

on the number of free targets is more appropriate (see (12)).

The term $\Omega(N_i)$ in eq. 1 summarizes replication processes, which we modeled

according to Abel zur Wiesch *et al.* (12). In short, cells of class N_i replicate

with the replication rate b_i. During replication, x bound antimicrobial molecules

of the mother cells are redistributed to the two daughter cells according to a

hypergeometric distribution:

$$
\Omega(N_i) = 2\sum_{x=i}^{n}(f_{x,i}b_x N_x) - b_i N_i
\tag{2}
$$

142 The parameter $f_{x,i}$ denotes the fraction of cells from class N_x that are redis-

143 tributed to N_i due to replication.

144   Bacteria in class N_i die with a natural death rate d_i and due to interaction

145 with the antimicrobial. We assume that the antimicrobial is bactericidal only.

The bactericidal effect becomes apparent in the zombie stage. Once a threshold number of antimicrobial molecules has hit the cell ($i = z$), the cell enters the zombie class. In the basic model, we assume complete multi-hit (31), with $z = n$. In the zombie class, bacteria are doomed to die and do so with an increased death rate $d_z > d$.

The multi-hit model allows us to keep track of the amount of antimicrobials A. In the basic model, we assume that there is no loss of antimicrobials due to non-reversible binding, non-targeted binding, or degradation, i.e. we assume that all antimicrobial molecules are recycled back into the system after death:

$$\frac{dA}{dt} = \sum_{i=0}^{n}(\mu_i i N_i + d_i i N_i - \alpha_i A N_i)$$

The variables N and A are used in a term describing mass action kinetics. Therefore, N and A are typically concentrations of bacteria and antimicrobial molecules, respectively (34). In the framework of the multi-hit model, it is however necessary to use the absolute amount of bacteria and antimicrobial molecules because we keep track of how many antimicrobial molecules have hit the bacterial cells (12). The volume of the system V is assumed constant, with $V = 1ml$. Resulting, the number of bacteria cells and antimicrobial molecules are equivalent to their concentration values.

Variations in the Multi-hit framework

We determined the multi-hit model based on the ODE system presented above. In the basic model (eq. 1–3), we assume the case of complete multi-hit (31), with $z = n$. The complete multi-hit model is appropriate for antimicrobials that need to hit a large fraction of the existing targets. An example of such a case are AMPs that kill according to the carpet model (35). Moreover we assume that the total amount of antimicrobials are stable and that antimicrobial molecules

are recycled back into the system when a cell dies. We relax these assumptions with (i) superfluous binding, (ii) non-reversible binding, (iii) non-targeted and non-reversible binding, and (iv) enzymatic degradation.

In the case of (i) superfluous binding , we implement the fractional multi-hit model by Hedges (*31*). In this model, $z < n$. Binding to targets is possible although the cell already reached the zombie stage. The fractional multi-hit model describes AMPs that kill by forming pores into the membrane of bacteria (*36*). Here, a cell has more targets, i.e. places where a pore can be formed in the membrane, than pores needed to kill a cell.

In the case of (ii) non-reversible binding, we set $\mu = 0$ and assume that antimicrobial molecules are not recycled back into the system, which results in

$$\frac{dA}{dt} \quad = \quad \sum_{i=0}^{n} -\alpha_i A N_i \tag{3}$$

In the case of (iii) non-reversible and non-targeted binding, we assume that in addition to non-reversible binding, molecules attach irreversibly to leaked cell content of dead cells. In our model, dead cells leak l targets per dying cell for non-targeted binding. L is the total number of targets available for non-targeted binding with AMPs with the rate α_d.

$$\frac{dL}{dt} \quad = \quad l \sum_{i=0}^{n} d_i N_i \tag{4}$$

$$\frac{dA}{dt} \quad = \quad \sum_{i=0}^{n} (-\alpha_i A N_i - \alpha_d A L) \tag{5}$$

Case (iv), enzymatic degradation, changes the absolute amount of total antimicrobials in the system, which also happens in the cases (ii) and (iii). Bacteria can produce proteases, e.g. aureolysin (*23*), which catalyze hydrolysis of AMPs. We assume that all bacterial cells produce enzymes E with the production rate

p and that enzymes degrade antimicrobial molecules with the rate d_E:

$$\frac{dE}{dt} \;=\; p\sum_{i=0}^{n} N_i - d_E E \tag{6}$$

In the presence of antimicrobials, enzymes catalyze antimicrobial degradation in a two step process. First, an enzyme binds to an antimicrobial molecule to form the complex $(E.A)$, then the bound enzyme catalyzes the degradation of the antimicrobial, resulting in enzymes and products: $\mathrm{E} + \mathrm{A} \;\underset{r_2}{\overset{r_1}{\rightleftharpoons}}\; \mathrm{E} \cdot \mathrm{A} \;\xrightarrow{r_3}\;$ $\mathrm{E} + \mathrm{P}$

$$\frac{dE}{dt} \;=\; p\sum_{i=0}^{n} N_i - d_E E - r_1 E A + r_2 E.A + r_3 E.A \tag{7}$$

$$\frac{dE.A}{dt} \;=\; r_1 E A - r_2 E.A - r_3 E.A \tag{8}$$

$$\frac{dA}{dt} \;=\; \sum_{i=0}^{n}(\mu_i i N_i + d_i i N_i - \alpha_i A N_i) - r_1 A E + r_2 E.A \tag{9}$$

Pharmacodynamic function

The pharmacodynamic (PD) function $\psi(A_0)$ (1, 2) describes the net growth rate for given antimicrobial concentration that was put in the system at $t = 0$ (A_0), with

$$\psi(A_0) \;=\; \psi_{max} - e(A_0) \tag{10}$$

$$\psi(A_0) \;=\; \psi_{max} - \frac{(\psi_{max} - \psi_{min})(A_0/MIC_{PD})^{\kappa}}{(A_0/MIC_{PD})^{\kappa} - \psi_{min}/\psi_{max}} \tag{11}$$

Here, the effect e describes the reduction of the net growth rate due to antimicrobials A_0. The MIC_{PD} is the antimicrobial concentration A_0 which results in a net growth rate of zero: $\psi(A_0 = MIC_{PD}) = 0$. Note that the MIC_{PD} is not necessary equal to the MIC (37). Other parameters of the PD function are the maximum net growth rate in absence of a drug $\psi_{max} = \psi(A_0 = 0)$,

the minimum net growth rate $\psi_{min} = \psi(A_0 \to \infty)$, and the slope parameter κ.

Alternatively, the PD function can be described with the parameters ψ_{max}, the maximum effect E_{max}, with $E_{max} = \psi_{max} - \psi_{min}$, the dose that results in half of the maximum effect A_{50}, with $\psi(A_{50}) = E_{max}/2$, and κ.

Together, the two definitions of th PD function allow us to calculate the slope parameter κ based on estimations of ψ_{max}, ψ_{min}, MIC_{PD}, and A_{50}, with

$$\kappa \;=\; \frac{\ln\left(-\frac{\psi_{max}}{\psi_{min}}\right)}{\ln\left(\frac{MIC_{PD}}{A_{50}}\right)} \tag{12}$$

The estimations of the parameter are based on simulated time-kill curves, in which we vary the antimicrobial concentration (fig. S1). Multiple such parameterized PD functions for varying bacterial inoculum sizes are then used to estimate the parameters for the extended PD function. Here, all estimates of the PD parameters are based on population dynamics on the ln-scale (natural logarithm).

We extend the PD function with information about the inoculum effect, $\psi(A_0, N_{inoc})$. The inoculum effect is quantified as slope m of the function

$$A_{0,\psi=y}(N_{inoc}) \;=\; mN_{inoc} + c, \tag{13}$$

with $y = 0$ and $y = E_{max}/2$ in the case of $\psi(A_0 = MIC_{PD})$ and $\psi(A_0 = A_{50})$, respectively.

The PD function is implemented in a population model to estimate population dynamics, with

$$\frac{dN_{tot}}{dt} \;=\; \psi_{max}N\left(1 - \frac{N}{K}\right) - e(A_0, N_{inoc})N \tag{14}$$

Here, $\psi_{max} = b$. We used this framework before (38).

Pharmacokinetic function

To model competition (see below), we use the pharmacokinetic (PK) function as described in Regoes *et al.* (*1*). Antimicrobials are put in the system with the concentration A_0. Antimicrobial molecules decay with the decay rate k_A. Without binding, the pharmacokinetic (PK) function is

$$A(t) = A_0 e^{k_A(t-p)} \tag{15}$$

with $n \times 8h \leq t \leq (n+1) \times 8h$.

Competition

To describe competition between a sensitive strain S and a resistant strain R, we extend the basic multi-hit model, with

$$\begin{aligned}
\frac{dS_i}{dt} &= \alpha_{i-1}AS_{i-1} - \alpha_i AS_i + \\
&\quad \mu_{i+1}(i+1)S_{i+1} - \mu_i i N_i + \\
&\quad \Omega(S_i) - d_i S_i \\
\frac{dR_j}{dt} &= \alpha_{j-1}AR_{j-1} - \alpha_j AR_j + \\
&\quad \mu_{j+1}(j+1)R_{j+1} - \mu_j j R_j + \\
&\quad \Omega(R_j) - d_j R_j \\
\frac{dA}{dt} &= \sum_{i=0}^{n}(\mu_i i S_i + d_i i S_i - \alpha_i AS_i) + \\
&\quad \sum_{j=0}^{n}(\mu_j i R_j + d_j j R_j - \alpha_i AR_i)
\end{aligned}$$

(16)
(17)
(18)

Here, i and j describe the number of targets hit on a sensitive and resistant bacterial cell, respectively, with $0 \leq i, j \leq n$. Strain R has a 10 times lower attachment rate ($\alpha_j = \alpha_i/10$) which reflects modification in the surface of bac-

terial cells, which is a common resistance mechanism against AMPs (21). To model realistic long term population dynamics, we also include a carrying capacity, as it is done in (12):

$$\Omega(N_i) \quad = \quad \left(2\sum_{x=i}^{n}(f_{x,i}b_x N_x) - b_i N_i\right)\left(1 - \frac{N_{tot}}{K}\right) \tag{19}$$

We simulated population dynamics in a window of 7 days.

Implementation

The multi-hit model was implemented in R (39) (version 3.4.0) in RStudio (40) (version 1.0.143). We solved the multi-hit model numerically with the package deSolve (41). For plotting, we used the following packages: sfsmisc (42) and RColorBrewer (43). The code is available upon request.

Results

We developed a mathematical model to describe the interaction between a bacterial population and antimicrobials. The model is based on the multi-hit concept of antimicrobial action (31). In the model, bacteria are killed as soon as a certain number of antimicrobial molecules hit the cell. The modeling framework allowed us to keep track of the number of bacteria and antimicrobial molecules over time. We assumed that the interaction occurs in a constant volume. Hence, the amounts of bacteria and antimicrobial molecules are equivalent to their concentration.

Model predictions of pharmacokinetics and pharmacodynamics

First we used our multi-hit modeling framework to investigate if an inoculum effect emerges simply as a result of concentration-dependent interactions between bacteria and antimicrobials and their binding kinetics. To this end, our basic model assumed reversible binding of antimicrobial molecules to target sites on the bacterial cells. In the model, antimicrobials were recycled back into the system once a bacterial cell died. As a consequence, only the amount of free antimicrobials, but not their total amount, decreased during the time course.

The decrease in the amount of free antimicrobials was dependent on the bacterial inoculum N_{inoc} (fig. 2a). The higher the bacterial inoculum, the more antimicrobial molecules hit bacteria due to the mass-action type binding kinetics that our model assumes, and the fewer antimicrobial molecules were free to hit in the following time step. The decrease in the amount of free antimicrobials due to initial binding to bacteria in the inoculum, in turn, changed the bacterial population dynamics. A bacterial population starting at a high bacterial inoculum grew over time, while a population exposed to the same amount of antimicrobials but starting at a low bacterial inoculum declined (fig. 2b). Thus, binding of antimicrobial molecules — even though it was reversible — caused an inoculum effect, i.e. the antimicrobial decreased in efficacy with increasing bacterial inoculum.

This inoculum effect altered the parameter of the pharmacodynamic (PD) function. The PD function describes the net growth rate of bacterial population for a given antimicrobial concentration and is a tool to describe efficacy of an antimicrobial. Here, we compared parameterized PD functions that are based on different inoculum sizes N_{inoc} with the PD plane (fig. 2c). The PD plane shows that the shape of the PD function changed with increasing in bacterial inoculum.

Figure 2: Inoculum size affects pharmacokinetics and pharmacodynamics. PD functions are estimated for a range of inoculum sizes. Increasing the inoculum size changes the efficacy of antimicrobials (b) due to reversible reduction of the number of free antimicrobials (a). (c) We determined the efficacy for varying inoculum sizes (N_{inoc}) with $z = 10$, resulting in a PD plane (note that in contrast to (d)-(f), the inoculum effect here is illustrated on a log_{10}-log_{10} scale). In the PD plane, solid gray contour lines show combinations of antimicrobials and inoculum sizes with the same net growth rate, both in the PD plane and projected above the PD plot. MIC_{PD} and A_{50} are indicated as thick solid and thick dashed line, respectively. For a number of inoculum sizes, we plotted the PD curve as dashed contour lines along the antimicrobial molecule number axis and again projected the contour lines, this time in front of the plot. The slopes of (b) are marked as dots in the corresponding colors. In (d)-(f) we plotted the PD parameters MIC_{PD}, A_{50}, and κ against the bacterial inoculum for varying number of hits needed to kill a bacterial cell (z). The bold black line (d) and dashed line (e) correspond to the same lines in (c). Parameters not mentioned here are listed in tab. S1.

To illustrate the magnitude of change, we plotted the net growth rates of the population dynamics pictured in fig. 2b in the PD plane. The net growth rate of a population based on a small inoculum size (blue point) is negative, while it is positive for a populations based on a large inoculum (orange point). Note that both populations were exposed to the same amount of antimicrobials at the beginning of the simulation (A_0). The example illustrates that the PD function and therefore the PD parameters are dependent on the bacterial inoculum.

In terms of specific PD parameters, both the MIC_{PD}, the amount of antimicrobials that results in zero growth, and A_{50}, the antimicrobial concentration that results in half of the maximum effect, increased with increasing inoculum (fig. 2d, 2e). The increase was linear and increased for increasing number of targets needed to kill a cell (z) (fig. 2d, 2e). Superfluous binding according to the fractional single hit model, i.e. antimicrobials can hit more than z targets, increased the inoculum effect, as well as non-reversible binding, and non-targeted binding combined with non-reversible binding (fig. S3). The steepness of the curve, parameterized with the PD parameter κ, increased non-linearly with increasing inoculum size (fig. 2f). The maximum effect of the antimicrobial E_{max} is defined by the parameters ψ_{max}, the maximum growth rate in absence of the antimicrobial ($\psi_{max} = \psi(A = 0)$), and ψ_{min}, the minimum net growth rate for $A \to \infty$. In previous work (32) we showed that ψ_{max} is only dependent on the multi-hit parameter replication and death rate of bacteria ($\psi_{max} = b - d$) and ψ_{min} is solely determined by the death rate of the zombie stages (d_z). This means, both PD parameters ψ_{max} and ψ_{min} and therefore also the parameter E_{max} were independent of any binding processes and the bacterial inoculum (fig. S4b, S4c).

Quantifying the inoculum effect

We quantified the inoculum effect in terms of the PD function by incorporating how the PD parameters change due to changes in the bacterial inoculum. We already determined that both MIC_{PD} and A_{50} increased linearly with increasing bacterial inoculum in our basic multi-hit model. The linear increase is a direct measure of the inoculum effect and therefore, we used the linear relationship $A_{0,\psi}(N_{inoc}) = mN_{inoc} + c$ to quantify the inoculum effect.

This linear relationship describes how many antimicrobial molecules A_0 we need to achieve an effect, specified in terms of net growth rate ψ, for a given bacterial inoculum size N_{inoc}. The slope m in this relation measures the inoculum effect. The inoculum effect increased with increasing number of hits needed to kill a cell (z), and was higher when we assumed superfluous binding, non-reversible binding, and non-reversible and non-targeted binding compared to the basic model of reversible binding (tab. 1). Moreover, it was higher for A_{50} than for MIC_{PD} in all the above cases. While the intersection c was influenced by the number of hits needed to kill a cell (tab. 1) it is per definition independent of the inoculum effect. The slope m and the intersection c describe how much antimicrobials we need to achieve a given pharmacodynamic efficacy. With these two parameters, we can also describe how much more antimicrobials we need to achieve the same pharmacodynamic efficacy when we increase the bacterial inoculum x-fold:

$$\frac{A_0(xN_{inoc})}{A_0(N_{inoc})} = \frac{mxN_{inoc} + c}{mN_{inoc} + c} \quad < \quad x \text{ for } c > 0 \tag{20}$$

Thus, the increase in antimicrobials required to achieve the same effect on an x-fold increased inoculum is, in general, less than x-fold.

The PD parameter κ was calculated in terms of the PD parameters MIC_{PD} and A_{50}. The bacterial inoculum size N_{inoc} affected the shape and position

Table 1: Results of fitting linear function $x = m_x N_{inoc} + c_x$ with $x = MIC_{PD}$ and $x = A_{50}$ to the simulated data. The data is plotted in fig. 2.

z	n	binding	MIC_{PD}		A_{50}	
			$c_{MIC_{PD}}$	$m_{MIC_{PD}}$	$c_{MIC_{PD}}$	$m_{A_{50}}$
1	1	reversible	2514516	0.167	8284221	0.258
2	2	reversible	6234471	0.728	14665506	0.812
10	10	reversible	37675418	6.508	61533236	6.577
10	39	reversible	37368958	6.672	61618874	6.902
10	10	non-reversible	57313826	9.081	62435986	9.944
10	10	non-reversible, non-targeted	36111621	9.416	111901448	11.080

of the pharmacodynamic curve, with the PD parameter MIC_{PD}, A_{50} and κ.

Temporary binding of antimicrobial molecules alone was enough to result in a

perceivable inoculum effect and the inoculum effect was intensified if we assumed

more complex binding processes. In agreement with this result, the inoculum

effect was abrogated if antimicrobial binding was not allowed in the simulation,

meaning the free antimicrobial concentration was assumed to be constant inde-

pendent of any binding processes (fig. S2). Using our results, we were able to

describe the net growth of bacteria for given N_{inoc} and A:

$$\psi(A_0, N_{inoc}) = \psi_{max} - \frac{E_{max}\left(\frac{A_0}{A50(N_{inoc})}\right)^{\kappa(N_{inoc})}}{1 + \left(\frac{A_0}{A50(N_{inoc})}\right)^{\kappa(N_{inoc})}} \tag{21}$$

$$\tag{22}$$

$$= \psi_{max} - \frac{(\psi_{max} - \psi_{min})\frac{A_0}{MIC_{PD}(N_{inoc})}^{\kappa(N_{inoc})}}{\frac{A_0}{MIC_{PD}(N_{inoc})}^{\kappa(N_{inoc})} - \frac{\psi_{min}}{\psi_{max}}}$$

339 with

$$A_{50}(N_{inoc}) = m_{A_{50}} N_{inoc} + c_{A_{50}} \tag{23}$$

$$MIC_{PD}(N_{inoc}) = m_{MIC_{PD}} N_{inoc} + c_{MIC_{PD}} \tag{24}$$

$$\kappa(N_{inoc}) = \frac{\ln\left(-\frac{\psi max}{\psi min}\right)}{\ln\left(\frac{MIC_{PD}(N_{inoc})}{A_{50}(N_{inoc})}\right)} \tag{25}$$

Degradation enhances the inoculum effect

Next, we extended our basic model to integrate proteolytic degradation of antimicrobials by extracellular enzymes. Bactria that are able to express extracellular enzymes have often been used to explore the inoculum effect (*15–20*). Assuming that each cell produces enzymes, the amount of enzymes is dependent on the bacterial density. In contrast to reversible binding, which we assumed in our basic model, degradation of antimicrobials decreased the number of antimicrobials irreversibly, i.e. the total number of antimicrobials in the system decreased.

Degradation of antimicrobial molecules in addition to reversible binding enhanced the effect the bacterial inoculum has on the dynamics of the antimicrobials (fig. 3a). While in the basic model the amount of free antimicrobials was only reduced due to binding, degradation decreased the amount of free antimicrobials further. The reduction of antimicrobials in turn enabled higher bacterial population sizes (fig. 3b) compared to the basic model. Not surprisingly, quantification of the inoculum effect based on PD parameters showed that the inoculum effect was stronger assuming degradation on top of reversible binding compared to reversible-binding only: The slopes of MIC_{PD} and A_{50}, $m_{MIC_{PD}}$ and $m_{A_{50}}$, were higher compared to reversible binding (fig. 3c, 3d). Again, due to the assumptions in the model, the size of the PD curve, i.e. parameter ψ_{max}, ψ_{min}, and therefore E_{max} were not affected by the inoculum (fig. S5a, S5b).

Evaluation of the extended PD function

The PD function $\psi(A_0)$ describes efficacy based on initial dynamics in the system and has been used to extrapolate long term bacterial population dynamics, e.g. (*1, 13, 38, 44–47*). Here, the underlying assumption is that the long-term bacterial population dynamics can be predicted on the basis of the initial short-

Figure 3: Degradation enhances the inoculum effect. (a) Antimicrobials are affected by degradation depending on the (b) bacterial population size. Here, we use 2 examples of bacterial inoculum sizes: low (blue) and high (orange). Dynamics with degradation: solid lines; dynamics without degradation (basic model): dashed lines. (c),(d) Degradation (red line) increases the number of molecules needed to achieve (c) $\psi(MIC_{PD}) = 0$ and (d) $\psi(MIC_{A_{50}}) = E_{max}/2$ compared to reversible binding only (black line). All parameters used for the simulations are listed in tab. S1.

term dynamics. This assumption is only valid if the net replication rate of a bacterial population of size N *at any time during the time-kill process* can be approximated by $\psi(A_0, N)$ that has been derived from the initial decline. This may not be the case because at later times the bacterial cells will have bound antimicrobial molecules, resulting in a state more susceptible to the antimicrobial effect. For these reasons, we asked if the extended PD function $\psi(A_0, N_{inoc})$ can be used to describe long term dynamics.

The multi-hit model enabled us to analyze if the short term population dynamics captured in the extended PD function predict long term dynamics. For this, we parameterized the extended PD function $\psi(A_0, N_{inoc})$ based on short term population dynamics from the multi-hit model. The multi-hit was then extended with a carrying capacity to account for density dependent reduction in growth and used to simulate long term population dynamics. The long term population dynamics from the multi-hit model were compared to predictions from a simple population model that relies on the extended PD function.

Fig. 4 (dashed lines) shows that the two approaches are in reasonable agreement assuming reversible binding only (basic model). Although the extended PD function underestimated the pharmacodynamic efficacy of the antimicrobial for low antimicrobial concentrations, the overall dynamical behavior of the system could be captured. In contrast, the dynamics based on the multi-hit model, in which enzymatic degradation was allowed, differed fundamentally from the dynamics predicted with the pharmacodynamic function at low bacterial inoculum (4, blue solid lines). Degradation decreased antimicrobials over time and the population was eventually able to increase in size. This effect was more pronounced at low bacterial inoculum sizes because at high bacterial inoculum sizes, the antimicrobial had already only little effect in the basic model. The extended PD function therefore captured simple bacterial population dynamics

(a)   (b)

Figure 4: Dynamics of bacterial population $N(t)$ within the first 24h after exposure to antimicrobials were simulated with (a) the multi-hit model and (b) a population model that incorporates $\psi(A_0, N_{inoc})$ (eq. 14). (a) We compared dynamics with (solid lines) and without extracellular proteins (dashed lines) as well as dynamics for low (blue) to high (orange) initial bacterial population sizes N_{inoc}. For the dynamics of $A(t)$ and $E(t)$, see fig. S6. The red area marks the time frame ($0h \leq t \leq 1h$) used to determine the net growth rate $\psi(A_0, N_{inoc})$. For a higher resolution of this time frame, but without carrying capacity, see figure 3a,3b). (b) We predicted long term dynamics based on $\psi(A_0, N_{inoc})$ with a simple population model (eq.14). All parameters used for the simulations are listed in tab. S1.

³⁹³ while only the mechanism-based multi-hit model could describe more complex

³⁹⁴ population dynamics.

³⁹⁵ Predicting the outcome of bacterial competition

³⁹⁶ We used the multi-hit model to ask how the inoculum effect influences competi-

³⁹⁷ tition and, more specifically, how the bacterial inoculum affects the competi-

³⁹⁸ tiveness of a recently mutated, more resistant strain. To answer this question,

³⁹⁹ we extended the multi-hit framework to two bacterial strains, strain S and R.

⁴⁰⁰ Strain R is more resistant to the antimicrobial than strain S due to a 10 $\times$

⁴⁰¹ lower attachment rate α. Note that production of extracellular enzymes is also

⁴⁰² a mechanism of bacterial resistance, but here we tried to avoid complex popu-

⁴⁰³ lation dynamics to be able to compare the results of the multi-hit model with

the dynamics based on the PD function. By extending the basic model by a strain R, we were able to simulate competition under complex pharmacokinetics (PKs).

We were interested if a resistant strain that is introduced in the system can take over the population within the simulation time. Strain R was not able to outcompete strain S at low antimicrobial concentrations (fig. 5a) due to an implemented cost of resistance. At intermediate concentrations, R took over the population (fig. 5b) and at high concentrations, both strains were killed (fig. 5c). These examples illustrate that the outcome of competition depended on antimicrobial dose given (A_0). We extended this analysis for a broad spectrum of input concentrations and bacterial inoculum sizes (fig. 5d). Here the range of antimicrobial concentrations for which the resistant strain can take over the population was dependent on the bacterial inoculum N_{inoc}. The higher the bacterial inoculum the smaller was the range of antimicrobial concentrations in which the resistant strain took over. These results were qualitatively reproduced with the extended PD function (fig. S7).

For comparison, we simulated the competition between the two strains in a model that does not display the inoculum effect. To parameterize such a model, we determined the PD function $\psi(A_0)$ for $N_{inoc} = 5 \times 10^5$. We assumed a wave-like pharmacokinetics. As expected, without the inoculum effect, the competitiveness of R did not depend on the inoculum (fig. 5e), and was overestimated, especially at high inoculum sizes (compare fig. 5e and fig. 5d).

Figure 5: The inoculum effect influenced competition outcomes. Antimicrobial doses A_0 changed the outcome of competition: (a) S won when exposed to low antimicrobial doses, (b) R won at intermediate antimicrobial doses and (c) bacteria went extinct at high antimicrobial doses. In an additional dimension, (d) the inoculum effect changes the outcome of competition (gray: S is more abundant, red: R is more abundant, white: extinction at the end of the simulation). (e) For comparison, we used the PD function to simulate competition without inoculum effect.

Discussion

The inoculum effect has been demonstrated experimentally for a number of bacterial strains and antimicrobials (*13, 15, 16, 19, 20, 24, 25, 27, 28*). In most cases the MIC (*13, 15, 20, 24, 25, 27*) was used to determine and quantify the inoculum effect. Udekwu *et al.* (*13*) demonstrated how to use MIC measurements to integrate the inoculum effect as change in MIC_{PD} in the PD function. However, they assumed that other PD parameters, among other things κ, were not affected by the inoculum effect.

In this work, we analyzed the inoculum effect based on mathematical modeling of bacterial population dynamics. We investigated whether the bacterial inoculum size has an impact on the efficacy of antimicrobials as measured by the PD function. A mechanism-based mathematical framework called multi-hit model (*31*) enabled us to explore the effect of increasing the inoculum size. Our findings are consistent with Abel zur Wiesch *et al.* (*12*): due to the kinetics of antimicrobials binding to bacterial cells, larger amounts of antimicrobials are required to attain the same efficacy when the bacterial inoculum size is increased.

In our analysis, the inoculum affected the PD parameters MIC_{PD}, A_{50}, and κ. Generally, the MIC_{PD}, A_{50}, and κ increased with increasing inoculum size. Because we calculated the slope parameter κ from MIC_{PD} and A_{50}, the effect of the inoculum on this parameter was a direct consequence of the effect on MIC_{PD} than A_{50}. We have previously shown that changes in PD parameters alter predictions about survival and resistance evolution of bacteria (*38*).

We quantified the inoculum effect as change in the PD parameters. An important insight in this context concerns a key parameter in our multi-hit modeling framework: the more targets are needed to be hit to kill a cell (z), the more pronounced was the inoculum effect on the PD parameters MIC_{PD},

A_{50}, and κ. This result suggest that the inoculum effect is stronger the more antimicrobial molecules are needed to kill a cell. In the case of antimicrobial peptides, this number varies from 80-800 in the case of protegrin (*36*) to $10^6 - 10^7$ in the case of PMAP-23 (*35*). With everything else being equal, we therefore expect a more pronounced inoculum effect for PMAP-23 than for protegrin.

In a next step, we determined the effect of enzymatic degradation in addition to the binding effects. The inoculum effect is often measured experimentally in systems with enzymes present that degrade antimicrobial molecules, for example β-lactamases (*16, 24, 48–50*). When allowing for enzymatic degradation in our simulations, the inoculum effect, quantified as increase in the pharmacodynamic parameters MIC_{PD} and A_{50}, was more pronounced. In these simulations, bacterial dynamics also increased in complexity, in particular at low bacterial concentrations. Importantly, however, our work shows that an inoculum effect can emerge without such enzymatic re-enforcement.

To quantitatively capture the effect of a range of antimicrobial concentrations on bacterial population growth, the pharmacodynamic function is commonly used. This function is parameterized to describe the short-term net replication rate of bacteria at one inoculum. When used to forecast the long-term behavior of bacterial populations under antimicrobial treatment, it is assumed that the pharmacodynamic function is also accurate for long-term dynamics. However, including enzymatic degradation of antimicrobials in our system revealed that the the complex long-term dynamics cannot be extrapolated from the short-term dynamics, and thus the extended PD function $\psi(A, N_{inoc})$ could not predict the bacterial population dynamics.

In our framework, the inoculum effect was caused solely by binding and degradation, but further mechanisms have been invoked to explain the inoculum effect. The efficacy of antimicrobials can change because of bacterial density

dependent change of abiotic factors. For example, pH (*29*), accessibility (*21*), susceptibility to antimicrobials (*30, 51*) can all change with bacterial density. Although we did not focus on these explanations, it is possible to extend the multi-hit framework to incorporate these explanations. These further mechanisms will exacerbate the inoculum effect that arises from the binding kinetics.

To investigate the influence of the inoculum effect on bacterial competition, we extended the basic multi-hit model by a resistant strain R. We assumed that the resistant strain is less prone to be hit by antimicrobials due to surface modifications (*21*). Simulation of bacterial competition revealed that the competitiveness of the resistant strain is overestimated when the inoculum effect is not considered. The range of antimicrobial concentration, in which R can take over corresponds to the mutant selection window (MSW) (*52, 53*). In our case, the MSW becomes smaller for higher bacterial inoculum sizes. This means that, while higher population sizes of S increase the probability of resistance emergence more S cells in the presence of antimicrobials also decreases the competitiveness of the emerged resistant strain by reducing antimicrobial efficacy. This unexpected feature of the inoculum effect might explain, at least in part, the puzzle of the sustained epidemiological co-existence of drug resistant and sensitive bacteria (*54*).

The absence of the inoculum effect has been reported repeatedly (*17, 55*). In our system, the inoculum effect was not perceivable in certain regions of the PD plane: for very small and very high antimicrobial numbers, the inoculum effect was absent. In addition, the inoculum effect on the MIC might be challenging to detect because of inherent uncertainties in determining the MIC by using doubling dilutions. As a concrete example, if, by increasing the inoculum, the "real" value of the MIC increases from 9 to 15, the measured MIC would be in both cases be 16. Furthermore, in many studies an arbitrary threshold of

an 8-fold increase of the MIC for a 100-fold higher bacterial inoculum is used to define the inoculum effect with MIC measurements (*9, 15, 16, 19, 20*), clouding inoculum effects of smaller extent.

It has been argued that the inoculum effect is an artifact (*17*). The core of this argument is that, to obtain the same pharmacodynamic efficacy, increasing the bacterial inoculum requires a proportionate increase in the amount of antimicrobials. Following this line of thought, keeping the ratio of antimicrobials to bacteria ($\frac{A_0}{N_{inoc}}$) constant should always result in the same pharmacodynamic efficacy. In our simulations, however, the ratio of antimicrobials to bacteria ($\frac{A_0}{N_{inoc}}$) does not predict efficacy. Rather, to obtain the same effect on a bacterial population of 100-fold the size of the original one we require less than 100-fold increase in antimicrobial molecules. For example, if we increase the bacterial inoculum 100-fold from 5×10^5 to 5×10^7, we only need a 9-fold increase in antimicrobial molecules to achieve the same pharmacodynamic efficacy and the ratio $\frac{A_0}{N_{inoc}}$ decreases from 80 to 7[1]. Thus, with increasing inoculum, each antimicrobial molecule becomes more effective with increasing inoculum.

In summary, we investigated the effect of the bacterial inoculum on the efficacy of antimicrobials in a simple conceptual model that describes the attached and detachment dynamics of antimicrobial molecules to target sites on the bacteria. We showed that the inoculum effect emerges from this simple dynamics. The quantitative aspect of this effect are surprisingly complex. We also showed that the inoculum effect influences the competitive dynamics between bacterial strains under treatment. We conclude that a comprehensive quantitative description of pharmacodynamics requires a detailed mechanistic population model and can not be captured by simple pharmacodynamic functions (*1, 13, 38, 47*).

---

[1] Calculations are based on eq. 20 and parameter values of slope m and intersection c are listed in tab. 1 (reversible binding, $n = z = 10$).

Authors contributions

R.R.R. and D.Y.B. initiated this project, conceived the model, analyzed the results and drafted the manuscript. D.Y.B. wrote the script in R.

Competing interests

We have no competing interests.

Funding

R.R.R. acknowledges the financial support of the ETH Zurich (ETH-41 15-2).

Acknowledgments

We thank Andrew Read for sparking our interest in the topic of inoculum effects, Pia Abel zur Wiesch, Sebastian Bonhoeffer, and Jens Rolff for reading the manuscript, and João Pires for interesting discussions about the project.

References

1. Regoes, R. R. *et al.* Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic Treatment Regimens. *Antimicrob. Agents Chemother.* **48,** 3670–3676 (2004).

2. Zhi, J., Nightingale, C. H. & Quintiliani, R. A Pharmacodynamic Model for the Activity of Antibiotics Against Microorganisms under Nonsaturable Conditions. *J. Pharm. Sci.* **75,** 1063–1067 (1986).

3. Nolting, A., Dalla Costa, T., Rand, K. H. & Derendorf, H. Pharma-cokinetic-

Pharmacodynamic Modeling of the Antibiotic effect of Piperacillin in Vitro.

Pharm. Res. **13,** 91–96 (1996).

4. Yu, G., Baeder, D. Y., Regoes, R. R. & Rolff, J. Combination Effects

of Antimicrobial Peptides. *Antimicrob. Agents Chemother.* AAC.02434–15

(2016).

5. Foerster, S., Unemo, M., Hathaway, L. J., Low, N. & Althaus, C. L. Time-

kill curve analysis and pharmacodynamic modelling for in vitro evaluation

of antimicrobials against Neisseria gonorrhoeae. *BMC microbiology* **16,** 216

(2016).

6. Wen, X. *et al.* Variation in fluoroquinolone pharmacodynamic parameter

values among isolates of two bacterial pathogens of bovine respiratory dis-

ease. *Scientific reports* **8,** 10553 (2018).

7. Ankomah, P. & Levin, B. R. Two-Drug Antimicrobial Chemotherapy:

A Mathematical Model and Experiments with *Mycobacterium marinum.*

PLoS Pathog. **8** (ed Fortune, S. M.) e1002487 (2012).

8. Mouton, J. W. & Vinks, A. A. Pharmacokinetic/pharmacodynamic mod-

elling of antibacterials in vitro and in vivo using bacterial growth and kill

kinetics. *Clinical pharmacokinetics* **44,** 201–210 (2005).

9. Wiegand, I., Hilpert, K. & Hancock, R. E. Agar and broth dilution methods

to determine the minimal inhibitory concentration (MIC) of antimicrobial

substances. *Nature protocols* **3,** 163 (2008).

10. CLSI. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria*

That Grow Aerobically; Approved Standard – Ninth Edition CLSI docu-

ment M07-A9 (Wayne, PA: Clinical and Laboratory Standards Institute,

2012).

11. ISO. *Clinical laboratory testing and in vitro diagnostic test systems – Susceptibility testing of infectious agents and evaluation of performance of antimicrobial susceptibility test devices – Part 1: Reference method for testing the in vitro activity of antimicrobial agents against rapidly growing aerobic bacteria involved in infectious diseases (Draft)* ISO/DIS 20776-1:2018(E) (Geneva, CH: International Organization for Standardization, 2018).

12. Abel zur Wiesch, P. *et al.* Classic reaction kinetics can explain complex patterns of antibiotic action. *Sci. Transl. Med.* **7,** 287ra73–287ra73 (2015).

13. Udekwu, K. I., Parrish, N., Ankomah, P., Baquero, F. & Levin, B. R. Functional relationship between bacterial cell density and the efficacy of antibiotics. *Journal of Antimicrobial Chemotherapy* **63,** 745–757 (2009).

14. Bhagunde, P. *et al.* Mathematical modeling to characterize the inoculum effect. *Antimicrobial agents and chemotherapy* **54,** 4739–4743 (2010).

15. Thomson, K. S. & Moland, E. S. Cefepime, piperacillin-tazobactam, and the inoculum effect in tests with extended-spectrum β-lactamase-producing Enterobacteriaceae. *Antimicrobial agents and chemotherapy* **45,** 3548–3554 (2001).

16. Adler, A., Chmelnitsky, I., Carmeli, Y., *et al.* Effect of resistance mechanisms on the inoculum effect of carbapenem in Klebsiella pneumoniae isolates with borderline carbapenem resistance. *Antimicrobial agents and chemotherapy* **59,** 5014–5017 (2015).

17. Craig, W. A., Bhavnani, S. M. & Ambrose, P. G. The inoculum effect: fact or artifact? *Diagnostic Microbiology and Infectious Disease* **50,** 229–230 (2004).

18. Brook, I. Inoculum effect. *Reviews of infectious diseases* **11,** 361–368 (1989).

19. Kang, C.-I. *et al.* Cefepime and the inoculum effect in tests with Klebsiella pneumoniae producing plasmid-mediated AmpC-type β-lactamase. *Journal of Antimicrobial Chemotherapy* **54,** 1130–1133 (2004).

20. Smith, K. P. & Kirby, J. E. The Inoculum Effect in the Era of Multidrug Resistance: Minor Differences in Inoculum Have Dramatic Effect on Minimal Inhibitory Concentration Determination. *Antimicrobial agents and chemotherapy,* AAC–00433 (2018).

21. Joo, H.-S., Fu, C.-I. & Otto, M. Bacterial strategies of resistance to antimicrobial peptides. *Phil. Trans. R. Soc. B* **371,** 20150292 (2016).

22. Ebbensgaard, A. *et al.* Comparative evaluation of the antimicrobial activity of different antimicrobial peptides against a range of pathogenic bacteria. *PLoS One* **10,** e0144611 (2015).

23. Sieprawska-Lupa, M. *et al.* Degradation of human antimicrobial peptide LL-37 by Staphylococcus aureus-derived proteinases. *Antimicrobial agents and chemotherapy* **48,** 4673–4679 (2004).

24. Saeki, M. *et al.* Inoculum effect of high concentrations of methicillin-susceptible Staphylococcus aureus on the efficacy of cefazolin and other beta-lactams. *Journal of infection and chemotherapy* **24,** 212–215 (2018).

25. Bedenić, B., Beader, N. & Žagar, Ž. Effect of inoculum size on the antibacterial activity of cefpirome and cefepime against Klebsiella pneumoniae strains producing SHV extended-spectrum β-lactamases. *Clinical microbiology and infection* **7,** 626–635 (2001).

26. Tan, C. *et al.* The inoculum effect and band-pass bacterial response to periodic antibiotic treatment. *Molecular systems biology* **8,** 617 (2012).

27. Chin, J. N., Rybak, M. J., Cheung, C. M. & Savage, P. B. Antimicrobial activities of ceragenins against clinical isolates of resistant Staphylococcus aureus. *Antimicrobial agents and chemotherapy* **51,** 1268–1273 (2007).

28. Savini, F. *et al.* Cell-density dependence of host-defense peptide activity and selectivity in the presence of host cells. *ACS chemical biology* **12,** 52–56 (2016).

29. Karslake, J., Maltas, J., Brumm, P. & Wood, K. B. Population density modulates drug inhibition and gives rise to potential bistability of treatment outcomes for bacterial infections. *PLoS computational biology* **12,** e1005098 (2016).

30. Bulitta, J. B. *et al.* Attenuation of colistin bactericidal activity by high inoculum of Pseudomonas aeruginosa characterized by a new mechanism-based population pharmacodynamic model. *Antimicrobial agents and chemotherapy* **54,** 2051–2062 (2010).

31. Hedges, A. J. An Examination of Single-hit and Multi-hit Hypotheses in Relation to the Possible Kinetics of Colicin Adsorption. *J. Theor. Biol.* **11,** 383–410 (1966).

32. Baeder, D. Y., Yu, G., Hozé, N., Rolff, J. & Regoes, R. R. Antimicrobial combinations: Bliss independence and Loewe additivity derived from mechanistic multi-hit models. *Phil. Trans. R. Soc. B* **371,** 20150294 (2016).

33. Magnus, C. & Regoes, R. R. Restricted occupancy models for neutralization of HIV virions and populations. *Journal of Theoretical Biology* **283,** 192–202 (2011).

34. McCallum, H., Barlow, N. & Hone, J. How should pathogen transmission be modelled? *Trends in ecology & evolution* **16,** 295–300 (2001).

35. Roversi, D. *et al.* How Many Antimicrobial Peptide Molecules Kill a Bacterium? The Case of PMAP-23. *ACS Chem. Biol.* **9,** 2003–2007 (2014).

36. Bolintineanu, D., Hazrati, E., Davis, H. T., Lehrer, R. I. & Kaznessis, Y. N. Antimicrobial mechanism of pore-forming protegrin peptides: 100 pores to kill E. coli. *Peptides* **31,** 1–8 (2010).

37. Mouton, J. W., Dudley, M. N., Cars, O., Derendorf, H. & Drusano, G. L. Standardization of pharmacokinetic/pharmacodynamic (PK/PD) terminology for anti-infective drugs: an update. *Journal of antimicrobial Chemotherapy* **55,** 601–607 (2005).

38. Yu, G., Baeder, D. Y., Regoes, R. R. & Rolff, J. Predicting drug resistance evolution: insights from antimicrobial peptides and antibiotics. *Proceedings. Biol. Sci.* **285,** 20172687 (2018).

39. R Core Team. *R: A Language and Environment for Statistical Computing* R Foundation for Statistical Computing (Vienna, Austria, 2017).

40. RStudio Team. *RStudio: Integrated Development Environment for R* RStudio, Inc. (Boston, MA, 2016).

41. Soetaert, K., Petzoldt, T. & Setzer, R. W. Solving Differential Equations in R: Package deSolve. *J. Stat. Softw.* **33,** 1–25 (2010).

42. Maechler, M. *et al. sfsmisc: Utilities from "Seminar fuer Statistik" ETH Zurich* R package version 1.1-0 (2016).

43. Neuwirth, E. *RColorBrewer: ColorBrewer Palettes* R package version 1.1-2 (2014).

44. Levin, B. R. & Rozen, D. E. Non-inherited antibiotic resistance. *Nature Reviews Microbiology* **4,** 556 (2006).

45. Ankomah, P. & Levin, B. R. Exploring the collaboration between antibiotics and the immune response in the treatment of acute, self-limiting infections. *Proceedings of the National Academy of Sciences,* 201400352 (2014).

46. Levin-Reisman, I. *et al.* Antibiotic tolerance facilitates the evolution of resistance. *Science,* eaaj2191 (2017).

47. Levin, B. R. & Udekwu, K. I. Population dynamics of antibiotic treatment: a mathematical model and hypotheses for time-kill and continuous-culture experiments. *Antimicrobial agents and chemotherapy* **54,** 3414–3426 (2010).

48. Burgess, D. S. & Hall II, R. G. In vitro killing of parenteral beta-lactams against standard and high inocula of extended-spectrum beta-lactamase and non-ESBL producing Klebsiella pneumoniae. *Diagnostic microbiology and infectious disease* **49,** 41–46 (2004).

49. Nannini, E. C. *et al.* Determination of an inoculum effect with various cephalosporins among clinical isolates of methicillin-susceptible Staphylococcus aureus. *Antimicrobial agents and chemotherapy* **54,** 2206–2208 (2010).

50. Wi, Y. M. *et al.* The cefazolin inoculum effect in methicillin-susceptible Staphylococcus aureus blood isolates: their association with dysfunctional accessory gene regulator (agr). *Diagnostic microbiology and infectious disease* **83,** 286–291 (2015).

51. Hirakawa, H. & Tomita, H. Interference of bacterial cell-to-cell communication: a new concept of antimicrobial chemotherapy breaks antibiotic resistance. *Frontiers in microbiology* **4,** 114 (2013).

52. Drlica, K. & Zhao, X. Mutant selection window hypothesis updated. *Clinical infectious diseases* **44,** 681–688 (2007).

53. Gullberg, E. *et al.* Selection of resistant bacteria at very low antibiotic concentrations. *PLoS pathogens* **7,** e1002158 (2011).

54. Lipsitch, M., Colijn, C., Cohen, T., Hanage, W. P. & Fraser, C. No coexistence for free: neutral null models for multistrain pathogens. *Epidemics* **1,** 2–13 (2009).

55. Mizunaga, S., Kamiyama, T., Fukuda, Y., Takahata, M. & Mitsuyama, J. Influence of inoculum size of Staphylococcus aureus and Pseudomonas aeruginosa on in vitro activities and in vivo efficacy of fluoroquinolones and carbapenems. *Journal of Antimicrobial Chemotherapy* **56,** 91–96 (2005).

711 Appendix

Table S1: Variables, indices and parameters of the multi-hit model. If not mentioned elsewhere, the parameter values used for the simulations are listed here. All rates are in $\frac{1}{h}$.

Variables	Definition
N_i	Number of bacteria that are hit by i antimicrobial molecules at time t.
N_{tot}	Total number of bacteria at time t, with $N_{tot} = \sum_{i=0}^{n} N_i$.
N_{inoc}	Number of bacteria at time $t = 0$, with $N_{inoc} = N_{tot}(t = 0)$.
A	Number of free antimicrobial molecules at time t, with $A_0 = A(t = 0)$.
E	Number of extracellular bacterial enzyme molecules at time t. We used $E(t = 0) = N_{inoc}p/(b_0 - d_0 - d_E)$.
$E.A$	Number of enzyme-antimicrobial complexes at time t, with $E.A(t = 0) = 0$.
S	Number of sensitive bacterial cells at time t.
R	Number of resistant bacterial cells at time t, with $R(t = 0) = 1$.
L	Number of targets for non-targeted binding at time t, with $L(t = 0) = 0$.

Indices	Definition
n	Total number of possible hits.
i,j	Number of hits, with $0 \leq i, j \leq n$. For the simulations, $n = 10$
z	Number of hits needed to kill a bacterial cell, with $1 \leq z \leq n$. If not mentioned, $z = 10$ is used.

Parameters	Definition	Value
t	time-point of the simulation	
b_i	Replication rate of bacterial cells in class N_i, with $i < z$.	1
b_z	Replication rate of bacterial cells in class N_i, with $i \geq z$.	0
d_i	Death rate of bacterial cells in class N_i, with $i < z$	0
d_z	Death rate of bacterial cells in class N_i, with $i \geq z$.	5
α_i	Hit rate of antimicrobial molecules with $0 \leq i \leq n$.	5×10^{-7}
μ_i	Detachment rate of antimicrobial molecules with $0 \leq i \leq n$.	0.1
r_1	Association rate of extracellular enzymes to antimicrobial molecules.	10^{-7}
r_2	Disassociation rate of $E.A$ complexes.	5×10^{-11}
r_3	Rate of catalysis.	5×10^{-4}
p	Production rate of bacterial enzymes.	50
d_E	Decay rate of bacterial enzymes.	0.1
α_d	Rate of non-targeted binding.	5×10^{-7}
l	Amount of intracellular targets of a bacterial cell.	20

Figure S1: Example of an estimation of PD parameters for $N_{inoc} = 0$. The estimation of PD parameters is based on the time-kill curves (middle panel) that are simulated with varying concentrations of A at time $t = 0$, A_0 (left panel), but always for the same inoculum concentration (middle panel). The slope was estimated for the time frame from $t_0 = 0$ to $t_1 = 1$h. Resulting, the parameter for $\psi(A_0)$ (right panel) are estimated for the given bacterial inoculum.

Figure S2: PD plane without inoculum effect due to fixed number of free antimicrobials. Here, we assumed $\frac{dA}{dt} = 0$.

Figure S3: PD parameters (a) MIC_{PD}, (b) A_{50}, and (c) κ for superfluous, non reversible and non-reversible and non-targeted binding compared to reversible binding ($z = 10$, as in fig. 2d- 2f). Note that all model variations shown here do not result in an change of the PD parameters E_{max}, ψ_{max}, and ψ_{min}. All parameters used for the simulations are listed in tab. S1.

Figure S4: Changing the bacterial inoculum does not change PD parameters (a) E_{max}, (b) ψ_{max}, and (c) ψ_{min}. Note that all model variations shown in fig. 2d- 2f result in no effect on these PD parameters. All parameters used for the simulations are listed in tab. S1.

Figure S5: Effect of degradation on PD parameters (a) ψ_{max}, (a) ψ_{min}, and κ. All parameters used for the simulations are listed in tab. S1.

<center>(a)</center><center>(b)</center>

Figure S6: Initial dynamics are differ fundamentally from long-term dynamics. The figures show the dynamics of (a) the amount of antimicrobial molecules ($A(t)$), (b) bacterial population size $N(t)$, and (c) extracellular proteins ($E(t)$) within the first $24h$ after exposure to antimicrobials. Here, we compared dynamics with extracellular proteins (solid lines) and dynamics of the basic model (dashed lines) as well as dynamics for a low (blue) and a high (orange) initial bacterial population size. In (b), we marked the time frame typically used to determine the net growth rate $\psi(A_0)$. All parameters used for the simulations are listed in tab. S1.

Figure S7: The effect of the bacterial inoculum on outcome of competition with the extended PD function (gray: S is more abundant, red: R is more abundant, white: extinction at the end of the simulation). All parameters used for the simulations are listed in tab. S1.