The pharmacodynamic inoculum effect from the perspective of bacterial population modeling

Desiree Y. Baeder and Roland R. Regoes[a]

Institute of Integrative Biology, ETH Zurich

[a]Author for correspondence: roland.regoes@env.ethz.ch

1 Abstract

The quantitative determination of the effects of antimicrobials is essential

for our understanding of pharmacodynamics and for their rational clinical

application. Common pharmacodynamic measures of antimicrobial efficacy,

such as the MIC and the pharmacodynamic function, fail to capture the

observed dependence of efficacy on the bacterial population size — a

phenomenon called inoculum effect. We assessed the relationship between

bacterial inoculum size and pharmacodynamic relationship and determined

the consequences of the inoculum effect on bacterial population dynamics

with a mathematical multi-hit model that explicitly describes the

interaction between antimicrobial molecules with their targets on the

bacterial cells. Our model showed that the inoculum effect can arise from

the binding dynamics of antimicrobial molecules to bacterial targets alone.

A pharmacodynamic function extended by the inoculum effect on its

parameters was able to predict the long-term population dynamics of

simple scenrios well. More complex scenarios, however, were only captured

with by the mechanistically more explicit multi-hit model. In simulations

with competing antimicrobial-susceptible and -resistant bacteria,

neglecting the inoculum effect led to an overestimation of the competitive

ability of the resistant strain. Our work underpins the importance of

including the inoculum effect into quantitative pharmacodynamic

frameworks, and provides approaches to accomplish that.

Introduction

Efficacy of antimicrobials against bacteria is commonly quantified using time-kill curves. The slopes of a log-linear function fitted to time-kill curves (Figure 1a) describe the change of the bacterial population over time and are a direct measure of the bacterial net growth rate. The net growth rate and its dependence on the antimicrobial concentration can be captured by the so-called pharmacodynamic (PD) function.[1,2,3] The parameters of this PD function, however, depend on the initial bacterial population size in the time-kill curve experiments.[4,5]

A more widespread, but less informative, way to express efficacy of antimicrobials is the MIC (Figure 1b). Although it is recognized that the MIC might change with bacterial inoculum size (see for example the MIC protocol by Wiegand *et al.*[6]), both the CLSI and the EUCAST recommendations to measure the MIC are based on a single standard bacterial inoculum of 5×10^5 cfu/mL.[7,8] This standardization of the inoculum yields an estimate of the MIC that cannot be extrapolated to other bacterial population sizes.

Changes in efficacy of antibacterial drugs caused by changing the initial bacterial inoculum size are summarized under the term *inoculum effect* (Figure 1). The inoculum effect was first described in the early 1940s,[9] and was observed for a variety of combinations of bacterial species and antimicrobials more recently.[4,10,11,12,13,14,15,16,17,18,19] In addition to its experimental establishment, the inoculum effect has been explored with mathematical models.[4,13,19,20,21,22,23,24] These experimental and theoretical

studies have put forward hypotheses explaining the inoculum effect that fall

into two broad categories. The first category attributes the inoculum effect

to the binding kinetics of antimicrobials: the higher the bacterial inoculum

the more antimicrobial molecules are bound, which leads to a larger

reduction of free antimicrobial molecules.[4,21,22] The second category invokes

the secretion enzymes at high bacterial inocula, e.g. β-

lactamases[9,10,14,15,16,25,26,27] and proteases.[28,29,30]

In this study, we used mathematical models as a tool to investigate these

competing hypotheses about the inoculum effect. Specifically, we use the so-

called multi-hit model[31] to address if the binding kinetics by itself already

leads to an inoculum effect, and to what extent it influences the PD

parameters. We have already used this model previously[32] and a similar,

albeit more antimicrobial-specific approach has been used to model classic

reaction kinetics of antibiotics.[21] Here we present an additional

implementation of this model by exploring the effects of the bacterial

inoculum conceptually. Additionally, we investigate how the inoculum effect

biases the competition between antimicrobial-susceptible and -resistant

strains.

65 Methods

66 Multi-hit model

We used the multi-hit model[21,31,32] that describes a bacterial population

consisting of N_{tot} cells. The key idea of this model is that targets on bacterial

cells are "hit", i.e. bound, by antimicrobial molecules, and a bacterial cell

dies if a threshold number of targets, which we call z, has been hit. The total

number of hits possible per bacterial cell is called n. Antimicrobial

molecules attach to and detach from a bacterial targets with the rates α_i and

μ_i, respectively. We denote the number of bacteria that have i antimicrobial

molecules adsorbed to cellular targets by N_i. Bacteria die with Bacteria die

with a death rate d_i Formally, the bacterial population in the multi-hit

model is structured according to how many antimicrobials have hit the cell

(Figure 2). For details of the multi-hit model, see supplement S1.

79 Variations in the Multi-hit framework

In the basic model (equation 1–3 in supplement S1), we assumed the case of

complete multihit,[31] with $z = n$, which maybe best describes antimicrobials

that need to hit a large fraction of the existing targets, for example AMPs.[33]

Moreover we assumed that the total amount of antimicrobials is stable and

that antimicrobial molecules are recycled back into the system when a cell

dies.

We considered variants of the multi-hit model by relaxing some of the

assumptions above: (i) not all targets need to be hit $z < n$, which leads to

non-effective binding, (ii) non-reversible binding, (iii) unspecific binding

(i.e. binding to non-targets) in combination with non-reversible binding,

leading to molecules attaching non-reversibly to targets and non-targets

alike, and (iv) enzymatic degradation. For details and model equations, see

Supplement S2.

Pharmacodynamic and Pharmacokinetic function

To investigate how the efficacy of antimicrobials changes with the inoculum in our multi-hit model simulations, we used the PD function[1,2,3] as a measure of antimicrobial efficacy. Specifically, we calculated the bacterial net growth rate ψ on basis of the antimicrobial concentration A at the start of the simulation ($A(t = 0) = A_0$), with

$$\psi(A_0) \quad = \quad \psi_{max} - e(A_0) \tag{1}$$

In this common conceptualization of PD effects, the net growth rate ψ is divided into the growth rate of the bacteria in absence of the antimicrobials ψ_{max} and an effect $e(A_0)$ that depends on the concentration of the antimicrobials. Note that we use the concentration A_0, because throughout the simulation, A varies due to binding and degradation.

The effect $e(A_0)$ is further parameterized by the concentration, MIC_{PD}, that leads to a zero net growth rate, $\psi(A_0 = MIC_{PD}) = 0$, the minimum net growth rate that can be obtained with antimicrobials $\psi_{min} = \psi(A_0 \to \infty)$, and a parameter that characterizes the steepness of the pharmacodynamic relationship, κ:

$$\psi(A_0) \quad = \quad \psi_{max} - \frac{(\psi_{max} - \psi_{min})(A_0/MIC_{PD})^\kappa}{(A_0/MIC_{PD})^\kappa - \psi_{min}/\psi_{max}} \tag{2}$$

Here, the MIC_{PD} approximates the MIC if the antimicrobial is concentration independent.[34]

113   To estimate the PD parameters, we simulated time-kill curves for which

114   we varied the antimicrobial concentration (Figure S1). In a first step, the net

115   growth rate was calculated based on the simulated time-kill curves using

116   the

117   relationship:

118
$$\psi(A) \;=\; \frac{log(N_{tot}(t_1)) - log(N_{tot}(t_0))}{t_1 - t_0} \tag{3}$$

The time window, t_1-t_0, is typically chosen to be on the order of

hours[1,32,35]**Error! Reference source not found.**,[36] to allow the measurement of

potentially steep declines of the bacterial population size at high

antimicrobial concentrations. For example, Tobramycin decreases the

bacterial population faster at high concentrations than Ticarcillin[35]. Thus, to

estimate the PD parameters, a shorter time window needs to be chosen for

Tobramycin than for Ticarcillin. Additionally, for wider time windows, more

complexities arise, such as persister populations that are detectable within a

few hours.[37,38,39,40] For the sake of simplicity, we set the time window that

we use to calculate the growth or decline of the bacterial population in our

simulations to 1 hour. The time window can be set to be narrower or wider

when, instead of aiming for a generic understanding of the inoculum effect,

a specific bug-drug combinations is being investigated.

The determined net growth rates were then used to estimate the PD

parameters. The steepness parameter κ was calculated (see supplement

S3). The estimates of the PD function parameters differed by inoculum. This

dependence of the PD parameters were captured by an extended PD

function, $\psi(A_0, N_{inoc})$. Specifically, the inoculum effect was quantified as slope

m of

the function

$$A_{0, \psi=y}(N_{inoc}) \quad = \quad mN_{inoc} + c, \qquad (4)$$

with $y = 0$ and $y = E_{max}/2$ in the case of $\psi(A_0 = MIC_{PD})$ and $\psi(A_0 = A_{50})$,

respectively.

Note that the pharmacodynamic function does not capture evolutionary

processes. However, evolved bacterial strains can be incorporated into this

pharmacodynamic modeling framework as subpopulations with their own

set of PD parameters.[36,41]

The pharmacokinetic function we used is described in Regoes *et al.*[1] In

short, the initial antimicrobial concentration A_0 is assumed to decay with a

rate k_A, and reset to A_0 every 8 hours.

Population model

151 To investigate the agreement between the bacterial population dynamics

152 predicted by the common parameterization of antimicrobial efficacy with

153 PD functions and those predicted by the more detailed multi-hit model, we

154 incorporated the extended PD function and the pharmacokinetic function

155 into a population model:

$$\frac{dN_{tot}}{dt} = \psi_{max}N\left(1 - \frac{N}{K}\right) - e(A_0, N_{inoc})N \tag{5}$$

157 Here, K is the carrying capacity. We used this framework before in.[42]

Competition

To describe competition between an antimicrobial-susceptible and an

antimicrobial resistant strain, we extended the basic multi-hit model with a

second strain. The number of susceptible and resistant cells are denoted by S

and R, respectively. In our simulations, strain R is assumed to have a 10 times

lower attachment rate, which could correspond to surface modification on

resistant bacterial cells — a common resistance mechanism against AMPs.[28]

For more details and model equations, see Supplement S4.

Implementation

167 The multi-hit model was implemented in R[43] (www.r-project.org; version

168 3.4.0). We solved the deterministic multi-hit model numerically with the

169 package deSolve.[44] For plotting, we used the following packages: sfsmisc[45]

170 and RColorBrewer.[46] The code is available upon request.

Results

We developed a mathematical model to describe the interaction between a

bacterial population and antimicrobials. The model is based on the multi-hit

concept of antimicrobial action,[31] according to which bacteria are killed as

soon as a certain number of antimicrobial molecules hit the cell. This

modeling framework describes the interaction between antimicrobial

molecules and bacterial cells in more detail than common population

biological models, and thus allowed us to investigate the determinants of

the inoculum effects and predict quantitatively how the inoculum affects the

pharmacodynamics.

Binding kinetics alone generates an inoculum effect

First, we investigated if an inoculum effect emerges simply as a result of

concentration-dependent interactions between bacteria and antimicrobials

and their binding kinetics. To that end, we used the basic multi-hit model

that assumes reversible binding of antimicrobial molecules to target sites

on

the bacterial cells.

We found that an inoculum effect emerged. In our simulations, a

bacterial population starting at a high inoculum grew over time, while a

smaller initial population exposed to the same amount of antimicrobials

declined (Figure 3a). The explanation of the effect can be found in the

concentration of free antimicrobial molecules that is sensitively affected by

the bacterial inoculum (Figure 3b), resulting in a lower efficacy if the

inoculum is high. Consistent with this explanation, the inoculum effect was

abrogated if the free antimicrobial concentration was assumed to be

constant independent of any binding processes (Figure S2). Thus, only the

binding of antimicrobial molecules — even though it was assumed

reversible — caused an inoculum

effect. Allowing for degradation of antimicrobials by extracellular enzymes

in our model exacerbated the inoculum effect (Figure 3) but was not

necessary.

202 Inhibitory concentrations increase linearly with inoculum,
203 but with a proportionality constant < 1

This inoculum effect can be quantitatively captured. Specifically, we studied

how the inoculum affects the parameters of the *pharmacodynamic function*

We considered the effect of the inoculum on the pharmacodynamic function

that describes relationship between the net growth rate of bacterial

population and the antimicrobial concentration (see Methods). Performing

in silico time kill experiments, we estimated the pharmacodynamic

functions for different inocula (Figure 4a) and found that the efficacy across

all antimicrobial concentrations is lower for larger inocula.

The most important efficacy parameter, the pharmacodynamic

parameter MIC_{PD} increases linearly with the inoculum (Figure 4b). However,

the proportionality constant is less than one, meaning that a doubling of the

inoculum does not lead to a doubling of number of antimicrobial molecules

required to halt bacterial population from growing, and thus does not lead

to a doubling of the MIC_{PD}. We found a similar linear increase of another

common efficacy measure, the A_{50} (see Methods for definition and Figure 4c),

which is interestingly also linearly affected by z.[32] We also found that these

two efficacy parameters increased linearly with the number of hits required

to kill a bacterial cell, z (Figure 4b,c) and that the increase was steeper when

we allowed for degradation (Figure S3). Furthermore, the steepness

parameter κ did not increase linearly, but saturated (Figure 4d). These

results hold for different modes of binding (Figure S4).

Formally, we found the following relationship between the efficacy
parameters and the inoculum N_{inoc}:

$$MIC_{PD}(N_{inoc}) \quad = \quad m_{MIC_{PD}}N_{inoc} + c_{MIC_{PD}} \tag{6}$$

$$A_{50}(N_{inoc}) \quad = \quad m_{A_{50}}N_{inoc} + c_{A_{50}} \tag{7}$$

227 Estimates for the intercept and slope parameters, c and m, of these linear

228 relationships are shown in Table 1. With these two parameters, we can also

229 describe how many more antimicrobial molecules we need to achieve the

230 same pharmacodynamic efficacy when we increase the bacterial inoculum

231 x-fold:

232
$$\frac{MIC_{PD}(xN_{inoc})}{MIC_{PD}(N_{inoc})} = \frac{mxN_{inoc} + c}{mN_{inoc} + c} \quad < \quad x \text{ for } c > 0 \tag{8}$$

The same relationship holds for the A_{50}, but, of course, with A_{50}-specific

estimates for c and m. Thus, the increase in antimicrobials required to

achieve the same effect on an x-fold increased inoculum is, according to our

model, less than x-fold and thus the proportionality constant is less than

one.

The parameters characterizing the net bacterial growth rate in absence

of antimicrobials is independent of the inoculum (Figure S5a). Also, the

maximum effect that can be achieved with an antimicrobial, E_{max}, and the

related minimum net growth rate, ψ_{min}, are independent of the inoculum
(Figure S5b-c). This is a consequence of the fact that these parameters are
not affected by the binding of antimicrobial molecules to targets on the
bacterial cells.[32]

The PD function $\psi(A0)$ has been used to predict the long-term bacterial
population dynamics from antimicrobial effect on the initial net growth
rate.[1,4,5,35,42,47,48,49] However, if we include the inoculum effects on MIC_{PD},
$A50$, and κ into the PD function, we found that this extended PD function
therefore captured only the short-term bacterial population dynamics while
only the mechanism-based multi-hit model could describe more complex
population dynamics (see supplement S5, Figure S6, Figure S7).

Inoculum effect alters the outcome of bacterial competition

To shed light on the effects of the inoculum on the competition between
bacterial strains, we extended the multi-hit model framework by an
additional resistant strain R. We assume that strain R carries a fitness cost
and is hence not able to outcompete strain S at low antimicrobial
concentrations (Figure 5a). At intermediate concentrations, R outcompetes
strain S (Figure 5b). Lastly, at high concentrations, both strains are killed
(Figure 5c). The antimicrobial concentrations that mark the transitions
between these three dynamical outcomes define the so-called *mutant*
selection window (MSW).[50,51]

To investigate how the inoculum affects the size of the MSW, we
conducted simulations across a range of inocula. In these simulations we
also included a time frame of 7 days and repeated doses of the
antimicrobial, to mimic antimicrobial usage in a treatment scenario. We
therefore use the term" treatment MSW". The difference between the MSW
and the treatment MSW is that the MSW is based on time-kill curves with a
single input of antimicrobials, while the treatment MSW is based on time-
kill curves in treatment regimes.

We find that the treatment MSW shrinks and even vanishes for
increasing inocula (Figure 5d). Qualitatively similar results hold for
simulations based on the extended PD function (Figure S8). Simulating
bacterial competition with a model that does not capture the inoculum
effect, in contrast, left the MSW unaffected (Figure 5e).

Discussion

In this study, we used a mathematical model to understand how an
inoculum effect can arise and to investigate how the bacterial inoculum
affects the pharmacodynamics of antimicrobials. With our model
simulations, we were able to derive predictions for the inoculum effect not
only on the MIC, but on multiple pharmacodynamic parameters. While the
insights from our study are generic and conceptual, they can be adapted to
any specific antimicrobial-bacterial pathogen combination. Our predictions
give rise to many experimentally testable hypotheses that we detail below.

Udekwu et al.[4] previously showed how the inoculum effect on the MIC
can be integrated into the PD function. We extended these efforts to the

additional pharmacodynamic parameters: the antibiotic concentration at

which 50% of the maximum effect is obtained, A_{50}, and the parameter that

describes the steepness of the pharmacodynamic relationship, κ. We found

that the inoculum affected the PD parameters MIC_{PD} and A_{50} linearly, and the

parameter κ non-linearly, resulting in an inoculum effect that does not

simply scale with the number of bacterial cells and antimicrobial molecules

in the system. For example, to obtain the same effect on a, say, 100-fold

increased bacterial population, we require less than 100-fold increase in the

number antimicrobial molecules. This finding directly contradicts the

argument that the inoculum effect is an artifact[27] arising from simple

scaling. We could show that to achieve the same antimicrobial effect for a

higher inoculum always requires a smaller increase in the number of

antimicrobial molecules than a simple scaling would suggest. How much

smaller depends on the specific parameters of the bacteria, antimicrobials,

and assay system. Interestingly, this means that, on a per-molecule level,

antimicrobials become more effective with increasing inoculum.

The mechanistic underpinning of the inoculum effect in our simulations

is that an increase in the bacterial inoculum results in higher depletion of

free antimicrobials. We call this the *numbers game*: Binding, even if it is

reversible, takes molecules out of the pool of free antimicrobial molecules,

which in turn means that there are fewer molecules available for binding in

the next time step. For larger bacterial inocula, the antimicrobial depletion

by binding is increased, resulting in the inoculum effect. Thus, for a given

antimicrobial-bacterial pathogen combination, we expect an inoculum effect

if the numbers game is at play. This numbers game results in an inoculum

effect has already been observed in mathematical modeling studies.[21,31]

Experimental studies that investigate the possibility and extent of an

inoculum effect arising from the numbers game alone[4,18,21], however, are

rare. Often, the inoculum effect is measured in systems in which additional

effects, e.g. non-targeted binding[52] or enzymatic

degradation[10,11,12,14,15,16,25,53,54,55], occur. These effects strongly increased the

complexity of the population dynamics in our simulations. We therefore

suggest conducting experimental studies that address the inoculum effect

first in systems that are governed only by a simple binding kinetics of

antimicrobials to their targets. These experiments should be conducted in a

regime of declining free antimicrobial molecule concentrations with increasing

bacterial inoculum sizes, because, in the numbers game scenario, this decline

causes the inoculum effect. Such studies will form the basis of a quantitative

understanding of the inoculum effect and will provide valuable baselines for

studies of systems with more complexity.

In addition to determining if the binding kinetics of antimicrobials alone

results in an inoculum effect, we suggest experiments that assess the more

qualitative prediction of the multi-hit modeling framework we presented

here. For example, we showed that the more targets are needed to be hit to

kill a cell (z), the more pronounced the inoculum affect was. In the case of

antimicrobial peptides, z varies from 80-800 in the case of protegrin[56] to 10^6

-10^7 in the case of PMAP-23.[33] With everything else being equal, we

therefore expect a more pronounced inoculum effect for PMAP-23 than for

protegrin. Similar predictions can be easily derived for other antimicrobials

if quantitative parameters characterizing their mechanisms of action are

known. Generally, the quantitative relationships between the mechanisms

of action of an antimicrobial, its pharmacodynamics, and the size of the

inoculum effect represent a promising area of future theoretical and

experimental studies.

The inoculum effect has been demonstrated experimentally for a number

of bacterial strains and antimicrobials. [4,10,11,12,14,15,16,17,18,19,21,53,54,57] In most

cases, the MIC was used to determine and quantify the inoculum

effect.[4,10,11,12,14,17] Our comprehensive analysis of the inoculum's effect on all

pharmacodynamic parameters showed that the steepness parameter κ

increases. Therefore, the inoculum effect is stronger for A_{50} than for MIC_{PD}

and determining the effect on A_{50} represent the most sensitive measure of

the inoculum effect. Based on this observation, we recommend to not only

quantify the MIC, but the whole pharmacodynamic relationship, and to use

OD measurements with caution because they do not allow the estimation of

the pharmacodynamic relationship for antibiotic concentrations above the

MIC_{PD}.

The absence of the inoculum effect has been reported repeatedly.[27,57]

Our results shed light on the conditions under which an inoculum effect is

expected to arise. First, in our simulations the inoculum effect did not

manifest itself in certain regions of the PD plane: for very small and very

high antimicrobial molecule numbers, the inoculum effect was absent.

Second, in our simulations, the inoculum effect essentially depends on a

decline in free antimicrobial concentrations. Therefore, we expect the

inoculum effect not to arise in any system, in which the antimicrobial

concentration does not decline with increasing bacterial inoculum. Third,

the establishment of the inoculum effect on the MIC is impaired because of

the uncertainties arising from the doubling of the antimicrobial

concentration that is required according to the standard protocol for MIC

determination. To give a specific example: according to the standard

protocol, "real" values of the MIC of 9 and 15 *mg/L* would be both estimated

as 16 *mg/L*; therefore, an inoculum effects increasing the MIC from 9 to 15

mg/L would not be detected. Lastly, in many studies an arbitrary threshold

of an 8-fold increase of the MIC for a 100-fold higher bacterial inoculum is

used to define the inoculum effect[6,10,14,15,16] clouding inoculum effects of

smaller extent. Rather than setting such a cutoff, we suggest determining

the MIC_{PD} or, even better, the A_{50} from adequately powered time-kill

experiments.

Most surprisingly, we found that the inoculum effect can have

unexpected and wide-ranging consequences for the evolutionary dynamics

of bacterial populations under drug treatment. Our simulation of bacterial

competition between drug-susceptible and -resistant strains revealed that

the competitiveness of drug-resistant strain is overestimated when the

inoculum effect is not considered. We found that, in our simulations, the

"treatment MSW" — an estimation of the MSW[50,51] under drug treatment —

shrinks for higher bacterial inoculum sizes. Although the treatment MSW is

an idealized concept, that does not capture all complexities of bacterial

evolution and competition, a negative effect of higher inocula on the
emergence of resistant strains should be observable in competition
experiments. In competition experiments between drug-susceptible and -
resistant bacterial strains, such as those presented in e.g.[51], our analysis
predicts the selection coefficients to decrease with inoculum. Hence the
inoculum effect is generally expected to impact bacterial survival and
evolution *in vitro*, within infected hosts, and potentially even on the
epidemiological level.

Acknowledgments

We thank Andrew Read for sparking our interest in the topic of inoculum effects, Pia Abel zur Wiesch, Sebastian Bonhoeffer, and Jens Rolff for reading the manuscript, Bruce Levin for giving us valuable feedback on our discussion, and João Pires for interesting discussions about the project. This work was supported by an ETH Grant (ETH-41 15-2, R.R.R.) and by the Volkswagen-Stiftung (Az 96 695, R.R.R.).

References

1 Regoes RR, Wiuff C, Zappala RM, *et al.* Pharmacodynamic Functions: a Multiparameter Approach to the Design of Antibiotic Treatment Regimens. *Antimicrob Agents Chemother* 2004; **10:** 3670–3676.

2 Zhi J, Nightingale CH, and Quintiliani R. A Pharmacodynamic Model for the Activity of Antibiotics Against Microorganisms under Nonsaturable Conditions. *J Pharm Sci* 1986; **11:** 1063–1067.

3 Nolting A, Dalla Costa T, Rand KH, *et al.* Pharma-cokinetic-Pharmacodynamic Modeling of the Antibiotic effect of Piperacillin in Vitro. *Pharm Res* 1996; **1:** 91–96.

4 Udekwu KI, Parrish N, Ankomah P, *et al.* Functional relationship between bacterial cell density and the efficacy of antibiotics. *J Antimicrob Chemother* 2009; **4:** 745–757.

5 Levin BR and Udekwu KI. Population dynamics of antibiotic treatment:

a mathematical model and hypotheses for time-kill and

continuousculture experiments. *Antimicrob Agents Chemother* 2010; **8:**

3414– 3426.

6 Wiegand I, Hilpert K, and Hancock RE. Agar and broth dilution meth-

ods to determine the minimal inhibitory concentration (MIC) of

antimicrobial substances. *Nat Protoc* 2008; **2:** 163.

7 CLSI. *Methods for Dilution Antimicrobial Susceptibility Tests for Bacteria*

That Grow Aerobically; Approved Standard – Ninth Edition. CLSI

document M07-A9. Wayne, PA: Clinical and Laboratory Standards In-

stitute, 2012.

8 ISO. *Clinical laboratory testing and in vitro diagnostic test systems –*

Susceptibility testing of infectious agents and evaluation of performance

of antimicrobial susceptibility test devices – Part 1: Reference method for

testing the in vitro activity of antimicrobial agents against rapidly

growing aerobic bacteria involved in infectious diseases (Draft). ISO/DIS

20776-1:2018(E). Geneva, CH: International Organization for

Standard-

ization, 2018.

9 Brook I. Inoculum effect. *Rev. Infect. Dis.* 1989; **3:** 361–368.

10   Thomson KS and Moland ES. Cefepime, piperacillin-tazobactam, and

the inoculum effect in tests with extended-spectrum β-lactamase-

producing Enterobacteriaceae. *Antimicrob Agents Chemother* 2001; **12:**

3548– 3554.

11   Saeki M, Shinagawa M, Yakuwa Y, *et al.* Inoculum effect of high concen-

trations of methicillin-susceptible Staphylococcus aureus on the

efficacy of cefazolin and other beta-lactams *J Antimicrob Chemother*

2018; **3:** 212–215.

12   Bedenić B, Beader N, Zagar Z. Effect of inoculum size on the

antibacterial activity of cefpirome and cefepime against *Klebsiella*

pneumoniae strains producing SHV extended-spectrum β-lactamases.

Clin Microbiol Infect 2001; **11:** 626–635.

13   Tan C, Smith RP, Srimani JK, *et al.* The inoculum effect and band-pass

bacterial response to periodic antibiotic treatment. *Mol Syst Biol* 2012;

1: 617.

14   Smith KP and Kirby JE. The Inoculum Effect in the Era of Multidrug

Resistance: Minor Differences in Inoculum Have Dramatic Effect on

Minimal Inhibitory Concentration Determination. *Antimicrob Agents*

Chemother 2018; **62:** e00433-18.

15 Kang CI, Pai H, Kim SH, *et al.* Cefepime and the inoculum effect in tests
with Klebsiella pneumoniae producing plasmid-mediated AmpCtype *β*-
lactamase. *J Antimicrob Chemother* 2004; **6:** 1130– 1133.

16 Adler A, Chmelnitsky I, Carmeli Y, *et al.* Effect of resistance mechanisms
on the inoculum effect of carbapenem in Klebsiella pneumoniae
isolates with borderline carbapenem resistance. *Antimicrob Agents*
Chemother 2015; **8:** 5014–5017.

17 Chin JN, Rybak MJ, Cheung CM, et al. Antimicrobial activities of
ceragenins against clinical isolates of resistant Staphylococcus aureus.
Antimicrob Agents Chemother 2007; **4:** 1268–1273.

18 Savini F, Luca V, Bocedi A, *et al.* Cell-density dependence of host-
defense peptide activity and selectivity in the presence of host cells.
ACS chem biol 2016; **1:** 52–56.

19 Loffredo MR, Savini F, Bobone S, *et al.* Inoculum effect of antimicrobial
peptides. *Proc. Natl. Acad. Sci.* 2021; **118.**

20 Karslake J, Maltas J, Brumm P, *et al.* Population density modulates drug
inhibition and gives rise to potential bistability of treatment outcomes
for bacterial infections. *PLoS comput biol* 2016; **10:** e1005098.

21 Abel zur Wiesch P, Abel S, Gkotzis S, *et al.* Classic reaction kinetics can
explain complex patterns of antibiotic action. *Sci Transl Med* 2015;
287: 287ra73–287ra73.

22   Bhagunde P, Chang KT, Singh R, *et al.* Mathematical modeling to

characterize the inoculum effect. *Antimicrob Agents Chemother* 2010;

11: 4739–4743.

23   Bulitta JB, Yang JC, Yohonn L, *et al.* Attenuation of colistin bactericidal

activity by high inoculum of Pseudomonas aeruginosa characterized by

a new mechanism-based population pharmacodynamic model.

Antimicrob Agents Chemother 2010; **5:** 2051–2062.

24   Greulich P, Scott M, Evans MR, *et al.* Growth-dependent bacterial

susceptibility to ribosome-targeting antibiotics. *Mol systbiol* 2015; **11:**

796.

25   Lenhard JR and Bulman ZP. Inoculum effect of β-lactam antibiotics. *J*

Antimicrob Chemother 2019; **10:** 2825–2843.

26   Rupp ME and Fey PD. Extended Spectrum β-Lactamase (ESBL)-

Producing Enterobacteriaceae. *Drugs* 2003; **4:** 353–365.

27   Craig WA, Bhavnani SM, and Ambrose PG. The inoculum effect: fact or

artifact? *Diagn Microbiol Infect Dis* 2004; **50:** 229–230.

28   Joo HS, Fu CI, and Otto M. Bacterial strategies of resistance to

antimicrobial peptides. *Phil Trans R Soc B* 2016; **1695:** 20150292.

29   Ebbensgaard A, Mordhorst H, Overgaard MT, *et al.* Comparative

evaluation of the antimicrobial activity of different antimicrobial

peptides against a range of pathogenic bacteria. *PLoS One* 2015; **12:**
e0144611.

30 Sieprawska-Lupa M, Mydel P, Krawczyk K, *et al.* Degradation of human
antimicrobial peptide LL-37 by Staphylococcus aureus-derived
proteinases. *Antimicrob Agents chemother* 2004; **12:** 4673–4679.

31 Hedges AJ. An Examination of Single-hit and Multi-hit Hypotheses in
Relation to the Possible Kinetics of Colicin Adsorption. *J Theor Biol*
1966; **3:** 383–410.

32 Baeder DY, Yu G, Hozé N, *et al.* Antimicrobial combinations: Bliss
independence and Loewe additivity derived from mechanistic multi-
hit models. *Phil Trans R Soc B* 2016; **1695:** 20150294.

33 Roversi D, Luca V, Aureli S, *et al.* How Many Antimicrobial Peptide
Molecules Kill a Bacterium? The Case of PMAP-23. *ACS Chem Biol* 2014;
9: 2003–2007.

34 Mouton JW and Vinks AA. Pharmacokinetic/pharmacodynamic
modelling of antibacterials in vitro and in vivo using bacterial growth
and kill kinetics. *Clin Pharmacokinet* 2005; **2:** 201–210.

35 Drusano, GL (2004). Antimicrobial pharmacodynamics: critical
interactions of 'bug and drug'. *Nat. Rev. Microbiol* 2004; **2:** 289-300.

36 Foerster S, Unemo M, Hathaway LJ, *et al.* Time-kill curve analysis and
pharmacodynamic modelling for in vitro evaluation of antimicrobials
against Neisseria gonorrhoeae*. BMC microbiol* 2016; **16:** 1-11.

37 Balaban NQ, Merrin J, Chait R, *et al*. Bacterial persistence as a
phenotypic switch. *Science* 2004*; **305:** 1622-1625.

38 Wiuff C, Zappala RM, Regoes, RR, *et al.* Phenotypic tolerance: antibiotic
enrichment   of   noninherited   resistance   in   bacterial
populations. *Antimicrob. Agents Chemother.* 2005; **49:** 1483-1494.

39 Balaban, NQ, Helaine S, Lewis K, *et al*. Definitions and guidelines for
research on antibiotic persistence. *Nat. Rev. Microbiol* 2019; **17:** 441-
448.

40 Rodríguez-Rojas A, Baeder DY, Johnston P, *et al.* Bacteria primed by
antimicrobial peptides develop tolerance and persist. *PLoS Pathog*
2021; **17:** e1009443.

41 El Shazely B, Yu G, Johnston PR, et al. Resistance evolution against
antimicrobial   peptides   in   Staphylococcus   aureus   alters
pharmacodynamics beyond the MIC. *Front microbiol* 2020; **11:** 103.

42 Yu G, Baeder DY, Regoes RR, *et al.* Predicting drug resistance evolution:
insights from antimicrobial peptides and antibiotics. *Proceedings Biol*
Sci 2018; **1874:** 20172687.

43   R Core Team. *R: A Language and Environment for Statistical Computing*.

R Foundation for Statistical Computing. Vienna, Austria, 2017.

44   Soetaert K, Petzoldt T, and Setzer RW. Solving Differential Equations in

R: Package deSolve. *J Stat Softw* 2010; **9:** 1–25.

45   Maechler M *et al. sfsmisc: Utilities from "Seminar fuer Statistik" ETH*

Zurich. R package version 1.1-0. 2016.

46   Neuwirth E. *RColorBrewer: ColorBrewer Palettes*. R package version

1.1- 2. 2014.

47   Levin BR and Rozen DE. Non-inherited antibiotic resistance. *Nat Rev*

Microbiol 2006; **7:** 556.

48   Ankomah P and Levin BR. Exploring the collaboration between

antibiotics and the immune response in the treatment of acute, self-

limiting infections. *Proc Natl Acad Sci* 2014; **111:** 8331-8338.

49   Levin-Reisman I, Ronin I, Gefen O, *et al.* Antibiotic tolerance facilitates

the evolution of resistance. *Science* 2017; **355:** 826-830.

50   Drlica K and Zhao X. Mutant selection window hypothesis updated. *Clin*

Infect Dis 2007; **5:** 681–688.

51   Gullberg E, Cao S, Berg OG, *et al.* Selection of resistant bacteria at very

low antibiotic concentrations. *PLoS pathog* 2011; **7:** e1002158.

52   Savini F, Loffredo M, Troiano C, *et al.* Binding of an antimicrobial

peptide to bacterial cells: Interaction with different species, strains and

552   cellular components. *Biochim Biophys Acta (BBA)-Biomembranes* 2020;

553   **1862:** 183291.

554 **53** Burgess DS and Hall II RG. In vitro killing of parenteral beta-lactams

555   against standard and high inocula of extended-spectrum beta-

556   lactamase and non-ESBL producing Klebsiella pneumoniae. *Diagn*

557   *Microbiol Infect Dis* 2004; **1:** 41–46.

558 **54** Nannini EC, Stryjewski ME, Singh KV, *et al.* Determination of an

559   inoculum effect with various cephalosporins among clinical isolates of

560   methicillin-susceptible Staphylococcus aureus. *Antimicrob Agents*

561   *Chemother* 2010; **5:** 2206–2208.

562 **55** Wi YM, Park YK, Moon C, *et al.* The cefazolin inoculum effect in

563   methicillin-susceptible Staphylococcus aureus blood isolates: their

564   association with dysfunctional accessory gene regulator (agr). *Diagn*

565   *Microbiol Infect Dis* 2015; **3:** 286–291.

566 **56** Bolintineanu D, Hazrati E, Davis HT, *et al.* Antimicrobial mechanism of

567   pore-forming protegrin peptides: 100 pores to kill E. coli. *Peptides*

568   2010; **1:** 1–8.

569 **57** Salas JR, Jaberi-Douraki M, Wen X, *et al.* Mathematical modeling of the

570   'inoculum effect': six applicable models and the MIC advancement

571   point concept. *FEMS microbiology letters* 2020; **5:** fnaa012.

58   Mizunaga S, Kamiyama T, Fukuda Y, *et al.* Influence of inoculum size of

Staphylococcus aureus and Pseudomonas aeruginosa on in vitro

activities and in vivo efficacy of fluoroquinolones and carbapenems. *J*

Antimicrob Chemother 2005; **1:** 91–96.

59   Magnus C and Regoes RR. Restricted occupancy models for

neutralization of HIV virions and populations. *J Theor Biol* 2011; **1:**

192–202.

Figures and Tables

Table 1: Results of fitting linear function $x = m_x N_{inoc} + c_x$ with $x = MIC_{PD}$ and $x = A_{50}$ to the simulated data. Reversible binding with $z = 1,2,10$ is also plotted in Figure 4b-d.

z	n	binding	MIC_{PD}		A_{50}	
			$c_{MIC_{PD}}$	$m_{MIC_{PD}}$	$c_{MIC_{PD}}$	$m_{A_{50}}$
1	1	reversible	2514516	0.167	8284221	0.258
2	2	reversible	6234471	0.728	14665506	0.812
10	10	reversible	37675418	6.508	61533236	6.577
10	39	reversible	37368958	6.672	61618874	6.902
10	10	non-reversible	36111621	9.081	62435986	9.944
10	10	non-reversible, non-targeted	57313826	9.416	111901448	11.08

588   (a)   (b)

Figure 1: The inoculum effect changes efficacy of antimicrobials. Two common
methods to quantify the efficacy of antimicrobials are (a) time-kill curves and (b)
MIC measurements. (a) Time-kill curves illustrate the change in population size
over time in presence of antimicrobials. The bacterial inoculum is the bacterial
population size N at $t0 = 0h$ ($N_{inoc} = N(t = 0)$). The slope of the curve results from
fitting a log-linear function to the population dynamics. Here, only the log-linear
curves and not individual measurements are shown. The slope of the time-kill
curve representing dynamics of bacteria starting at a high inoculum size (orange)
is shallower than dynamics of bacteria starting at a low inoculum (blue). In the
absence of an inoculum effect, the slope is independent of the inoculum (dashed
orange line). (b) Alternative to measuring time-kill curves, efficacy is typically
determined with the MIC. An inoculum effect is observed if increasing the
inoculum size from low (blue) to high (orange) increases the MIC.

Figure 2: Schematic representation of the basic multi-hit model with three classes
($n = 2$). At every time point the bacterial population N_{tot} can be divided in n
subpopulations based on the number of antimicrobials that have hit the bacterial
cells. A hit is the mathematical simplification of adsorption and subsequent
downstream processes connected to binding of one antimicrobial molecule to a
bacterial target site and is described with the rate α. The parameter μ is the

detachment of an antimicrobial from the bacterial target. We assumed that finding
the bacterial cell is the rate limiting step for an antimicrobial molecule.[59] This is
appropriate if the number of targets are in abundance, e.g. when antimicrobial
peptides adsorb to the cell membrane of bacterial cells. For antimicrobials that act
within the cell, a hit rate dependent on the number of free targets is more
appropriate (see[21]).

Each class N_i gains bacterial cells from the class N_{i-1} with the term $\alpha_{i-1}AN_{i-1}$ and a
bacterium dies with an increased death rate d_z if two antimicrobial molecules have
hit the cell, i.e. $z = 2$. Therefore, $N_{tot} = N_0 + N_1 + N_2$, with $N_2 = N_z$. A is the number of
free antimicrobial molecules, here indicated in blue, antimicrobials that have hit a
cell are indicated in pink. The rates of the ODE (see supplement S1 for equation)
are indicated with arrows and bacterial replication with dashed arrows.

623   (a)   (b)

Figure 3: Increasing the bacterial inoculum size from 5 ⬚ 10⁵ (blue) to 5 ⬚ 10⁷
(orange) changes the efficacy of the antimicrobial due to reduction of the number
of free antimicrobials. Simulations with reversible binding is shown with solid
lines, dashed lines indicate simulations with degradation. (a) The bacterial
inoculum *N(t)* is affected differently by antimicrobials depending on bacterial
inoculum size and binding (b). Time course of free antimicrobial molecules. The
range of the decline of free antimicrobials is consistent with other modelling
studies[19] , and *in vitro*[33] and *in vivo* observations[35]. Parameters not mentioned here
are listed in Table S1.

634
(a)

(b) (c) (d)

Figure 4: Inoculum size affects pharmacodynamics. (a) We determined the efficacy
for varying inoculum sizes (N_{inoc}) with z = 10, resulting in a PD plane (note that in
contrast to (b)-(d), the inoculum effect here is illustrated on a log_{10}-log_{10} scale). In
the PD plane, solid gray contour lines show combinations of antimicrobials and
inoculum sizes with the same net growth rate, both in the PD plane and projected
above the PD plot. MIC_{PD} and A_{50} are indicated as bold solid and dashed line,
respectively. For several inoculum sizes, we plotted the PD curve as dashed
contour lines along the antimicrobial molecule number axis and again projected
the contour lines, this time in front of the plot. The slopes of Fig. 3a (solid lines).
are marked as dots in the corresponding colors. In (b)-(d) we plotted the PD
parameters MIC_{PD}, A_{50}, and κ against the bacterial inoculum for varying number of
hits needed to kill a bacterial cell (z). The bold black line in (b) corresponds to the
bold solid line in (a) and the bold black line in (c) correspond the bold dashed lines
in (a). Parameters not mentioned here are listed in Table S1.

652
653   (a)   (b)   (c)

655   (d)   (e)

Figure 5: The inoculum effect influenced competition outcomes. Antimicrobial
doses A_0 changed the outcome of competition: (a) S won when exposed to low
antimicrobial doses, (b) R won at intermediate antimicrobial doses and (c) bacteria
went extinct at high antimicrobial doses. (d) In an additional dimension, the
inoculum effect changed the outcome of competition (gray: S is more abundant,
red: R is more abundant, white: extinction ($S,R < 1$) at the end of the simulation).
(e) For comparison, we used the PD function to simulate competition without
inoculum effect.