Faculty of Biology Medicine and Health
The University of Manchester
Oxford Road
Manchester   M13 9PT

www.manchester.ac.uk

The University of Manchester

8 February 2019

To the Editor(s),

Please find enclosed our manuscript entitled "**The essential role played by B cells in supporting protective immunity against *Trichuris muris* infection is dependent on host genetic background and is independent of antibody"** that we would like to be considered for publication in Plos Pathogens.

The importance of the B cell in resistance to gut dwelling nematode parasites has been debated at length. Our work will be of particular interest to the field because it reveals that B cells play an essential, but antibody independent, role in resistance to *Trichuris muris* but that this dependence varies with host genetic background.

Specifically, we demonstrate that:

(a) The essential role played by the B cell in the protective immune response varies with genetic background and the degree of Th cell polarisation of the host, with the B cell supporting the development and maintenance of protective Th2 immune responses in IFN-g-rich environments. Thus, if IFN-g is depleted from mouse strains that mount mixed Th1/Th2 response (eg C57BL/6), or the host strain is one that mounts dominant Th2 immune responses, the B cell becomes redundant in the protective immune response.

(b) The important role played by the B cell in promoting resistance to *T. muris* infection in mixed Th1/Th2 cytokine settings is in supporting the development and maintenance of the Th2 immune response and is not related to antibody production.

We look forward to your comments,

KJElce.

Kathryn Else and Rinal Sahputra, for the authors.

Kathryn Else,
Professor of Immunology
Faculty of Biology, Medicine and Health,
University of Manchester
Kathryn.j.else@manchester.ac.uk

The essential role played by B cells in supporting protective immunity against *Trichuris*

***muris* infection is dependent on host genetic background and is independent of antibody**

Rinal Sahputra[¥], Dominik Ruckerl, Kevin Couper, Werner Muller, Kathryn J Else[¥]

Lydia Becker Institute for Immunology, The University of Manchester

[¥]Joint corresponding authors

Email: Kathryn.j.else@manchester.ac.uk

Rinal.sahputra@manchester.ac.uk

Abstract

This study investigates the role of B cells in immunity to *Trichuris muris* (*T. muris*)

infection in two genetically distinct strains of mouse, using anti-CD20 monoclonal antibody

(mAb) (Genentech-clone 5D2) to deplete B cells. Data is presented for the mouse strains:

C57BL/6 and BALB/c, which mount mixed Th1/Th2, and highly polarised Th2 immune

responses to *T. muris*, respectively. C57BL/6 mice receiving anti-CD20 treatment prior to

and during, or anti-CD20 treatment that commenced two weeks post infection (p.i.), were

susceptible to *T. muris*. Parasite-specific IgG1 antibodies were absent and Th2 type cytokines

produced by mesenteric lymph nodes cells from mice receiving α-CD20 mAb treatment were

significantly lower than produced by cells from isotype control treated mice. T follicular

helper cells were also significantly reduced. Importantly, and in complete contrast, BALB/c

mice were still able to expel *T.muris* in the absence of B cells, revealing that the essential role

played by B cells in protective immunity was dependent on genetic background. To explore

whether the important role played by the B cell in the protective immune response of

C57BL/6 mice was in enabling strong Th2 responses in the presence of IFN-γ, IFN-γ was

blocked using anti-IFN-γ mAb post B cell depletion. Depleting IFN-γ, in the absence of B

cells restored worm expulsion in the absence of parasite-specific IgG1/IgG2c and partially

rescued the *T. muris* specific IL-13 response. Thus, our data suggest an important, antibody

independent role for B cells in supporting Th2 type immune responses in mixed IFN-γ-rich

Th1/Th2 immune response settings.

Author summary

How B cells contribute to protective immunity against parasitic nematodes remains

unclear, with their importance as accessory cells under-explored. This study reveals that, on

some genetic backgrounds, B cells are important for the expulsion of *T. muris* by acting as

accessory cells, supporting Th2 immune responses.

Introduction

Infecting over two billion people around the world, mostly in resource-limited countries, the ability of parasitic helminths to maintain long standing chronic infections makes them a major health care issue (1). *Trichuris trichiura* (*T. trichiura*) is one of the most common gastrointestinal nematodes, infecting approximately 465 million people worldwide (2), primarily children. In infected children, trichuriasis is strongly associated to malnutrition, growth stunting, and intellectual retardation; whereas in pregnant adults it is related to anaemia and low birth weight babies (3).

For decades, *T. muris* in the mouse has provided a useful and relevant model system with which to explore immunity to *T. trichiura* in man. Infection of mice with intestinal nematode parasite *T. muris* drives polarized T helper cell (Th) responses which associate with resistance (Th2) or susceptibility (Th1). However, the key cellular contributions which support Th2 cell polarization during *T. muris* infection are still not well understood. One of the cells thought to be important is the B cell. It is well established that B cell function is not only related to antibody production, but B cells can also act as antigen presenting cells (APCs) due to the expression of MHC class II molecules and several co-stimulators, including CD40, CD80, and CD86 on its surface (4-6), and as accessory cells, acting as a cellular source of multiple cytokines (7).

B cells are well placed to act as accessory cells as they are able to influence CD4 T cells polarisation. For example, previous studies have shown that CD4 T cells produce significantly more IL-4 when stimulated with antigen presented by B cells compared to macrophages (8). In addition, B cells have been shown to produce either Th1 or Th2 type cytokines *in vivo* depending on the type of parasite. Thus, for the Th1 driving parasite *Toxoplasma gondii*, B cells produce IFN-γ and IL-12p40 (9); whilst in the context of the Th2 polarising parasite *Heligmosomoides polygyrus* (*H. polygyrus*), B cells produce IL-4 and IL-2

(10, 11). Recent studies reported that IL-4 producing B cells during early infection of

Schistosoma mansoni (*S. mansoni*) are critical for Th2 polarisation, protecting the host

against *S. mansoni*-induced pathology (12). B cells can also influence T cell polarisation

towards Th2 response by providing co-stimulatory molecules. Linton *et al*., (13) showed that

co-stimulation via OX40L expressed by B cells is essential for T cell polarisation towards

Th2 cells. Likewise, both CD80 and CD86 co-stimulatory molecules expressed by B cells, as

well as other APC, are required for the Th2 response during *H. polygyrus* infection (14).

Recently, it was suggested that B cells, together with T cells and DCs are required not

only to optimize the development of Th2 type responses, but also to maintain T follicular

helper (T$_{FH}$) and memory Th2 cells development following *H. polygyrus* infection (15). Thus,

Leon *et al.* (15) showed that the deletion of CXCR5 on either DC or CD4 T cells, during *H.*

polygyrus-infection of C57BL/6 mice, impaired the development of T$_{FH}$ and Th2 cells by

impairing the migration of CXCR5+ cells towards the B cell follicle in response to CXCL13.

Interestingly, B cell depletion also impaired Th2 responses (15).

Previous studies suggested that B cells and antibody are not important in mediating

resistance following a primary *T. muris* infection (16-18). Since then, many different studies

have been performed to show the importance of CD4 T cells in mediating resistance against

T. muris (19-21). By contrast, the role of B cells in immunity to *T. muris* remains largely

unexplored. *T. muris*-infected μMT mice on a C57BL6 background develop Th-1 type

responses, resulting in the susceptibility to *T. muris* infection (22). These susceptible *T.*

muris-infected μMT C57BL6 mice produced high IFN-γ, without any Th2 cytokines

production. Furthermore, when these mice were treated with B cells from naive C57BL6 or

with anti-IL-12 antibody, resistance to infection was restored (22). This suggests that B cells

are important in either inhibiting Th1 development or supporting Th2 type immune

responses. However, given the importance of B cells in the development of lymph nodes and

tissue organisation (23, 24), data from μMT mice is difficult to interpret.

This study therefore investigates the role of B cells and antibodies in immunity to *T.*

muris infection using anti-CD20 monoclonal antibody (mAb) to deplete B cells from mice of

two distinct genetic backgrounds, C57BL/6 and BALB/c. The benefit of using anti-CD20

mAb is that it allows depletion of CD19+ cells either prior to or post infection and avoids the

complicating consequences of B cell deficiency during embryonic development. We

demonstrate that B cells are important in the protective immune response to *T. muris*; that the

role played by the B cell is antibody-independent; and that the importance of the B cell varies

with genetic background of the host. Thus the B cell plays an essential role in supporting Th2

type immune responses only in mixed Th1/Th2 IFN-γ rich settings, as seen in C57BL/6 mice,

and is redundant in the highly Th2 polarised, IFN−γ-deficient environment of the BALB/c

mouse post *T. muris* infection.

Results

2.1 Depletion of B cells throughout infection requires two injections of anti-CD20 mAb

As a transmembrane calcium channel, CD20 is important in B cell activation, proliferation, and differentiation (25). The CD20 molecule is normally expressed on the surface of B cells during the late pre-B cell. Therefore, the injection of anti-CD20 mAb will deplete all B cells, except early pre-B cells and plasma cells (26). The vast majority of IgM$^+$ or B220$^+$ cells in the spleen, lymph nodes, peritoneal cavity, and blood express CD20, whilst in bone marrow the expression of CD20 increases with B cell maturation (27). Previous studies showed that anti-CD20 effectively depleted B cells either via complement-mediated lysis of target cells (28) or by inducing FcγR-mediated clearance (Ab-dependent cell mediated cytotoxicity) (29). Furthermore, a single dose of anti-CD20 treatment prior to lymphocytic choriomeningitis virus infection depleted B cell populations for up to 45 days (30). However, the current study showed that a single anti-CD20 injection failed to ablate B cells for the full duration of *T. muris* infection, with a clear CD19+ population re-emerging in the MLNs, spleen, and blood by day 35 p.i. (Fig. 1A&B). Therefore, a second anti-CD20 injection was administered at day 10 p.i. in order to maintain B cell depletion in the blood and secondary lymphoid organs throughout infection (Fig. 1A&B). CD19$^+$ cells were still present in the bone marrow after anti-CD20 treatment (Fig. 1B), however these CD19$^+$ cells were pro-B cells (Fig. 1C) as has been shown previously (26).

2.2 C57BL/6 mice treated with anti-CD20 mAb fail to expel *T. muris* by d35 p.i., correlates with an absence of class switched antibodies and a significant decrease in Th2 cytokines

To investigate whether B cells are important in resistance against *T. muris* during primary infection of C57BL/6 mice, mice were injected with anti-CD20 mAb 7 days prior to infection and 10 days post infection. Chronic infection, characterised by persisting adult stage

parasites from day 32 p. i. defines susceptibility (22). Thus, autopsies were performed beyond this time point. The experimental design is shown in Fig. 2A. Previous studies have shown that C57BL/6 mice take up to 35 days to completely expel the parasite (31). As shown in Fig. 2B, C57BL/6 depleted of B cells were significantly more susceptible to infection than control treated mice.

It has previously been shown that IgG1 antibodies are associated with resistance to *T. muris* infection, whilst IgG2c antibodies are related to susceptibility (32). Therefore, the levels of parasite specific antibodies in the sera of anti-CD20 treated mice and isotype control treated mice were compared. Mice depleted of B cells using anti-CD20 mAb failed to secrete IgG1 and IgG2c parasite specific antibodies, in contrast to control treated mice (Fig. 2C&D). Further, the levels of *T. muris* specific IgM antibodies in the sera of anti-CD20 treated mice were significantly lower than in the sera of the isotype control-treated mice (Fig. 2E). These effects of B cell depletion on antigen-specific IgG and IgM antibodies are consistent with reports in other model systems (33).

In common with other nematode parasites, hosts resistant to *T. muris* mount Th-2 immune responses characterised by the production of IL-4, IL-5, IL-9 and IL-13; in contrast, mice susceptible to *T. muris* mount a Th-1 type response, dominated by the release of IFN-γ (32, 34, 35). B cell depleted mice on a C57BL/6 background produced significantly lower Th2 cytokines, including IL-5, IL-9, and IL-13 compared to mice treated with the isotype control (Fig. 2F-H). Interestingly, the production of IFN-γ and IL-17 was similar in both groups (Fig. 2I&J) suggesting that B cells do not affect Th1 development but rather boost Th2 responses. These data are in keeping with the previous study by Leon, *et al*., which revealed that B cell depletion impaired the development Th2 in mice against *H. polygyrus* infection (15).

2.3 B cell depletion reduced the T_{FH} population and altered DC subsets in the MLN

The importance of B cells for the formation of T_{FH} has been shown previously in both

mouse (15) and man (36). In keeping with these studies, the MLN T_{FH} population of anti-

CD20 treated mice in C57BL/6 genetic background was significantly reduced compared to

isotype control treated mice (Fig. 3A-C). Gating strategy was shown in Supplementary Fig. 1.

In contrast, T_{FH} development in the spleen, distal to the site of infection, was not significantly

affected by B cell depletion (Supplementary Fig.2), although the lack of statistical

significance may be due to the high variability in the control mice.

Previous studies have suggested that $CD11b^+$ DCs in MLNs might be important for

promoting Th2 immune responses against *T. muris* infection (37). Because Th2 cytokines

were significantly reduced in the MLNs from anti-CD20 treated mice during *T. muris*

infection, DC subsets in MLNs were analysed (Fig. 3D-H). The gating strategy is shown in

Supplementary Fig. 3. Total cell number of $CD103^-CD11b^+$ DCs (Fig. 3F) and $CD103^+$

$CD11b^+$ DCs (Fig. 3H) were significantly reduced in MLNs of anti-CD20 treated mice

compared to isotype control treated mice.

2.4 B cell depletion from day 14 post infection also altered the worm burden

As depletion of B cells throughout the course of infection of C57BL/6 mice lead to

susceptibility to *T. muris*, we wondered whether B cell depletion from day 14 p.i. would also

alter the phenotype. Previous studies on the expulsion kinetics of *T. muris* have shown that

the onset of worm expulsion occurs after day 12 p.i., with T cell activation occurring after the

first 7 days (38). Therefore, B cells were depleted 2 weeks post *T. muris* infection to see if

this still impacted on the ability to expel the parasite. The experimental design is shown in

Fig. 4A. Surprisingly, the depletion of B cells from day 14 p.i. also impaired worm expulsion

with significantly more parasites present at day 35 p.i. than seen in control-treated mice (Fig.

4B). T$_{FH}$ cells and Th2 type cytokines, including IL-5, IL-9, and IL-13 were also significantly

reduced in the absence of B cells (Fig, 4C-H). Interestingly, TNF-α was significantly

increased in B cell depleted mice (Fig. 4I), whilst IFN-γ and IL-17A remained the same

between groups (Fig. 4J&K). These data suggest that the B cell plays an important role in

maintaining Th2 responses.

2.5 Anti-IFN-γ treatment partially restored IL-13 production in B cell depleted mice and

rescued worm expulsion

In order to investigate whether the impaired resistance to infection in the absence of B

cells was due to the reduced Th2 immune response or loss of parasite specific antibodies, B

cell depleted mice were injected with anti-IFN-γ. Anti-IFN-γ treatment is a common strategy

to promote resistance in susceptible mice (39). The experimental design for anti-IFN-γ

injection plus B cell depletion is shown in Fig. 5A. As shown in Fig. 5B, anti-IFN-γ treatment

restored resistance to *T. muris* in B cell depleted mice.

As expected, blocking IFN-γ in the absence of B cells did not rescue the *T. muris*

specific IgG1 response (Fig. 5C). T. muris specific IgG2c antibodies were not detected in the

sera of anti-IFN-γ treated mice with intact B cells (Fig. 5D), consistent with a highly

polarised Th2 immune response. In contrast, the isotype control mice produced both IgG1

and IgG2c antibodies (Fig. 5C&D). Consistent with our previous data, mice treated with anti-

CD20 mAb produced significantly lower Th2 cytokines, such as, IL-13, IL-4, IL-5 and IL-9

compared to isotype control treated mice (Fig. 5D-G) with the Th1 response unaffected (data

not shown). Importantly, blocking IFN-γ significantly increased IL-13 production in mice

depleted of B cells (Fig. 5D), although levels were still significantly lower than in isotype

control treated mice (Fig. 5D). Although no significant difference was noted in the

production of other Th2 type cytokines, there were trends towards increased *T. muris* specific

IL-4, IL-5, and IL-9 after anti-IFN-γ treatment in B cell depleted mice compared to mice

depleted of B cells but not IFN-γ (Fig. 5F-H). Taken together, these results show that in the

absence of IFN-γ, B cells and IgG1 parasite specific antibodies are not important for *T. muris*

expulsion on a C57BL/6 genetic background; thus without any competing IFN-γ partial

restoration of the Th2 immune response is sufficient to render animals resistant to infection

Consistent with data in Fig 4, B cell depletion reduced the T_{FH} population in MLNs

(Fig. 5I&J). T_{FH} cells have been associated with Th2 immunity due to their ability to secrete

IL-4, a Th2 signature cytokine (33, 40, 41). Given that IL-13 production persisted in B cell

depleted mice after anti-IFN-γ injection, we wondered whether the T_{FH} population was also

altered. Anti-IFN-γ treatment did not restore the T_{FH} population in MLNs of anti-CD20

treated mice at day 21 p.i. (Fig. 5I&J), however it remains a possibility that the few TfH

remaining were sufficient to induce the partial Th2 response.

2.6 B cells are essential in supporting Th2 immune responses against *T. muris* only in an IFN-

γ-rich environment.

In the current study we found that when C57BL/6 mice were depleted of IFN-γ, the

immune response became more highly polarised towards Th2 mice, evidenced by parasite-

specific IgG1 antibodies, in the absence of IgG2c. In this more polarised Th2 environment

the depletion of B cells did not prevent worm expulsion. These data suggest that the

important role played by B cells in supporting Th2 immune responses against *T. muris* is only

necessary in IFN-γ-rich environments. Therefore, we decided to deplete B cells in BALB/c

mice, which are naturally very resistant to *T. muris* and mount highly polarised Th2 immune

responses in the absence of IgG2a/c and IFN-γ (31). The experimental design is shown in

Fig. 6A. CD19+ cells in MLNs, spleen, and PerC of BALB/c mice were not detected after

anti-CD20 mAb injection (Fig. 6B). In complete contrast to B cell depleted C57BL/6 mice,

BALB/c mice were still able to expel the parasite in the absence of B cells (Fig. 6C). IgG1

and IgG2c parasite specific antibodies were undetectable in B cell depleted mice (Fig.

6D&E), and levels of *T. muris* specific IgM were significantly lower than in the sera of

isotype control treated mice (Fig. 6F). As shown previously (31, 39), isotype control treated

mice on BALB/c genetic background mounted a strong parasite specific IgG1 response but

did not secrete *T. muris*-specific IgG2a/c (Fig. 3K), indicative of a highly polarised Th2

immune response.

Collectively, our data reveals that the important role played by the B cell in promoting

resistance to *T. muris* infection in mixed Th1/Th2 cytokine settings is in supporting the

development and maintenance of the Th2 immune response and is not related to antibody

production. Importantly, the essential role played by the B cell in the protective immune

response varied with genetic background and the degree of Th cell polarisation of the host.

Thus, if IFN-γ is depleted from mixed Th1/Th2 settings, or the Th2 immune response is

dominant, the B cell becomes redundant in the protective immune response.

Discussion

T. muris infection drives different Th cell responses in different strains of mice (34). BALB/c, BALB/k, and NIH mice are very resistant to *T. muris* infection, expelling the worms by around day 18 post infection, while AKR mice are susceptible to infection, unable to expel the parasite and harbour patent chronic infections (42). C57BL6 and C57BL10 mice are also resistant to infection, but they expel the worms more slowly, between day 18 and 35 p.i. (43). Susceptible hosts mount predominantly Th-1 immune responses associated with the presence of IFN-γ and IL-12, whilst very resistant host strains mount strong Th-2 type polarised immune response characterised by cytokines such as IL-4, IL-5, IL-9 and IL-13, and very low levels of IFN-γ. The quality of the Th cell response in C57BL6 mice is less clearly polarised with the slower expulsion kinetic associated with a mixed Th1/Th2 phenotype and presence of IgG1 and IgG2c. Despite a good understanding of the Th cell response during *T. muris* infection, the key cellular contributions which support T helper polarization are still not well understood, nor how genetic background impinges on this.

This study aimed to investigate the role of B cells in immunity to *T. muris* during primary infection by using anti-CD20 mAb to deplete B cells. In order to understand whether the contribution of the B cell varied with the nature of the protective immune response, we used mice of two different genetic backgrounds: C57BL/6 and BALB/c. CD20 is specifically expressed on the surface of B cells from the pre-B cells stage to immature B cells, but then disappear when B cells differentiate to plasma cells (28). Adding antibody against CD20 inhibits the progression of B cells from the G1 phase into S/G2+ M stages (44), resulting in the inhibition of B cell differentiation, antibody production, and inducing B cell apoptosis (45). Using anti-CD20 mediated depletion of B cells, antigen specific class switched antibodies are ablated and IgM antibodies are significantly reduced (41, 46). These finding

are in keeping with the finding of current study and enable an assessment of the importance

of parasite-specific IgG1 antibodies in the expulsion of *T. muris*.

Our data reveal that antibody is not essential for the expulsion of *T. muris*, with both

B cell depleted BALB/c mice and B cell depleted C57BL/6 mice also depleted of IFN-γ, able

to clear infection. These data also exemplify that the mechanism of immunity to *T. muris* in

BALB/c mice is entirely B cell independent. However, our data do identify an important and

significant antibody-independent role for the B cell in promoting immunity to infection in

mouse strains, such as C57BL/6, which are not highly polarised towards a Th2 immune

response. Thus, the importance of the B cell in resistance to infection is dependent on genetic

background, and represents a key consideration when interpreting data from other

experimental models. Further, given that protective immunity was lost when B cells were

depleted both throughout the time course of infection and only after the first two weeks of

infection, our data suggest that the role played by the B cell is in supporting both the

development and maintenance of Th2 immune responses.

Mechanistically, B cell depletion correlated with a significant reduction in

CD4+CXCR5+PD-1^{high} cells. CD4+CXCR5+PD-1^{high} cells are recognised as T_{FH} and are

known to be able to produce the Th2-type signature cytokine; IL-4 (47). Thus, these data are

consistent with previous studies showing that B cells are important in regulating Th2 immune

responses (10, 14, 15, 48-51) and maintaining T_{FH} development (15, 52). Interestingly, T_{FH}

populations in both MLN and spleen of BALB/c mice were not affected after B cell

depletion. Further, the presence of T_{FH} has been associated with Th2-type immune responses

in *H. polygyrus* infection (15). Although in the current study, depletion of B cells in mice also

treated with anti- IFN-γ restored expulsion of *T. muris* without rescuing the T_{FH} response, the

few T_{FH} cells remaining may be sufficient to support the partially restored Th2 response.

Thus it remains possible that T$_{FH}$ cells are essential in building strong Th2 responses in the

absence of IFN-γ.

292   The importance of B cells in immunity to *T. muris* has previously been proposed

using μMT mice (22) on a C57BL/6 background which are susceptible to infection unless

Th1 responses are inhibited using anti-IL-12. These data are keeping with the current study

supporting a role for B cells as accessory cells promoting and maintaining Th2 responses

rather than antibody producers. Antibody independent expulsion of *T. muris* is also evidenced

by the fact that primed CD4+ T cell transfer to SCID mice is sufficient to support worm

expulsion in the absence of antibody (18).

299   More broadly, the role of B cells in immunity to gastro intestinal nematodes in

general has been debated at length (53-56). Proposed mechanisms include production of

antibody (57), promoting and maintaining of primary and memory Th2 cells (10, 14) via their

ability to produce cytokines, especially IL-4 (11, 53, 58), and/or by expressing co-stimulatory

molecules, including OX40L and CD40 (13, 59). A variety of experimental approaches have

been used including the use of transgenic B cell deficient mice (10, 22, 57, 60), FcγR

deficiency (61, 62), passive immunization (22, 63, 64) and maternal antibody transfer (54,

65); and a variety of conclusions have been drawn. IgG1 antibodies were shown to be

essential for worm expulsion against *H. polygyrus* infection based on previous studies using

AID mice, which retain a secretory IgM response, on C57BL/6 genetic background (57).

Antibody was shown to be essential for worm expulsion against *H. polygyrus* infection via

antibody-dependent cell-mediated cytotoxic (ADCC) mechanisms (66). However, ADCC

mechanisms do not play a critical role in immunity to *T. muris* infection as FcγR deficiency

mice are still able to expel the parasite (62).

313   More recently, two studies have strongly suggested a role for B cells in directing T

cell polarisation towards Th2 at the time of antigen presentation (15, 67). Leon *et al*., showed

that B cell depletion in *H. polygyrus*-infected C57BL/6 mice prevented the migration of DCs

into B cell areas and impaired the Th2 immune response (15). Lymphotoxin, produced by B

cells, is important in the control of CXCL13 expression (15), a chemokine expressed by

follicular dendritic cells (68) and marginal reticular cells in the peri and interfollicular regions

between B cell follicles (69) and that is essential to attract CXCR5-expressing cells towards

the B cell area (15). Thus by treating C57BL/6 mice with anti-CXCL13, Leon *et al.,* (15)

also revealed that CD4 T cells and DCs accumulated in the T cell area rather than in

perifollicular region and that IL-4+ Th2 cells in mesenteric lymph nodes (MLNs) were

significantly reduced. In support of a mechanism involving the movement of DCs and CD4 T

cells to the B cell area to allow CD4 T cells to be educated by B cells towards a Th2, recent

studies have revealed that transgenic mice in which CXCR5 depletion was specifically

restricted to DCs become susceptible to *T. muris* with very few DCs detected in the B cell

area compared to control mice (67).

In addition to the proposed role for B cells in educating the DC-CD4 T cell

conversation suggested by Leon *et al.,* (15) and Bradford *et al.,* (67), the importance of

CD11b+ DCs in MLNs for Th2 polarisation has been shown in previous studies, against both

T. muris (37) and *S. mansoni* (70) infection. In this context, in the current study we observed

reduced numbers of CD11b+ DCs in B cell depleted C57BL/6 mice and this may contribute

to the reduced Th2 immune response and increased susceptibility of anti-CD20 treated mice

to *T. muris* infection. Although previous studies have indicated the importance of B cells in

regulating the development of DC subsets in secondary lymphoid organs (50, 71, 72), further

studies are required to investigate how the absence of B cells reduced the number of CD11b+

DCs in the MLN.

Overall, we present a study showing that, irrespective of host genetic background,

antibody is not essential in the expulsion of *T. muris*. However, our data do identify an

essential role played by the B cell on certain genetic backgrounds where IFN-γ rich, mixed

Th1/Th2 cell responses are observed, and strongly suggest that the B cell is able to adopt

multiple functions in inducing and maintaining protective Th2 responses during gastro-

intestinal nematode infection.

Material and Methods

1. Animals

C57BL/6 and BALB/c mice were purchased from Envigo, UK and were maintained in ventilated cages in the Biological Services Facilities (BSF) of the University of Manchester. Mice were housed in the facility at least 7 days prior to experimentation and were infected at 6-8 weeks old with *T. muris* by oral gavage.

2. B cells depletion and infection

To assess the importance of B cells for worm expulsion in different strains, C57BL/6 and BALB/c mice were split into 2 groups of 4-5 mice: anti-CD20 mAb and isotype control treated mice. C57BL/6 or BALB/c mice were treated with 100μg in 200μl PBS of anti-CD20 mAb (5D2, Genentech) or isotype control (Rat IgG2a, Biolegend) i.v. injection via tail vein. Mice were infected with approximately 200 T. muris eggs at day 7 post injection. Mice were re-injected with anti-CD20 mab or isotype control at day 10 p.i. C57BL/6 mice were autopsied at day 35 p.i., whilst BALB/c mice were autopsied at day 42 p.i. For anti-IFN-γ, C57BL/6 mice were split into 4 groups of 5 mice: isotype control of anti-CD20 mAb + isotype control of anti-IFN-γ mAb (rat Ig), anti-CD20 mAb + isotype control of anti-IFN-γ mAb, isotype control of anti-CD20 mAb + anti-IFN-γ mAb, anti-CD20 mAb + anti-IFN-γ mAb. C57BL/6 mice were treated with 100μg in 200μl PBS of anti-CD20 mAb (5D2, Genentech) or isotype control (Rat IgG2a, Biolegend) i.v. injection via tail vein. Mice were infected by oral gavage with approximately 200 *T. muris* eggs at day 7 post injection. Mice were re-injected with anti-CD20 mab or isotype control at day 10 p.i. 1 mg of α-IFN-γ antibody or rat IgG1 (as a control) antibody was given at day 0, day 7, and day 14 p.i. Mice were euthanised by CO_2 followed by autopsy at day 21 or day 35 p.i..

3. *Trichuris muris* maintenance and the preparation of E/S proteins

All protocols to maintain the parasite and to prepare the E/S were as previously

described (73). Briefly, the parasite was passaged through SCID mice that are susceptible to

T. muris infection. SCIDs received a high dose of approximately 300 *T.muris* embryonated

eggs and at approximately day 35 p.i. the large intestine was collected to produce adult E/S.

Guts were collected and then longitudinally split open before washed in warmed 5x

penicillin/streptomycin in complete RMPI 1640 medium (500U/ml penicillin, 500μg/ml

streptomycin, 500ml RMPI 1640 medium). Adult worms were then carefully pulled from the

gut using fine forceps and transferred to a 6 well plate containing 4ml warmed 5x pen/strep in

RMPI 1640 medium. Plates were incubated in a moist humidity box for 4 hours at 37°C to

collect 4 hr E/S. For overnight E/S, adult worms were then split into 2 wells containing fresh

medium and incubated again in a humidity box at 37°C overnight. Supernatant from 4 hour

and 24 hour incubations was collected and centrifuged at 2000g for 15 minutes. *T. muris* eggs

from adult worms were resuspended in 40ml deionised water and filtered through a 100μm

nylon sieve before transferring to a cell culture flask. To allow embryonation, eggs were

stored in darkness for approximately 8 weeks and then stored at 4°C. SCID mice were

subsequently infected with a high dose infection to determine infectivity of each new batch of

eggs. Thus, larvae were counted at around d14 p.i. and the number of larvae counted

expressed as a % over the number of eggs given to determine the infectivity of the egg batch.

All E/S supernatant was filter sterilised through a 0.2μm syringe filter (Merk). E/S was

concentrated using an Amicon Ultra-15 centrifugal filter unit (Millipore) by spinning at

3000g for 15 minutes at 4°C. E/S was dialysed against PBS using Slide-A-Lyzer Dialysis

Cassettes, 3.500 MWCO (Thermo Science) at 4°C. The concentration of E/S was measured

using the Nanodrop 1000 spectrophotometer (Thermo Fisher Science) and aliquoted before

storing at -80°C.

For high dose *Trichuris muris* infection, approximately 3-4 ml of egg suspension was

transferred to a universal tube and topped up with deionised water before centrifuging for 15

minutes at 2000g. Pelleted eggs were washed with deionised water, resuspended and only

embryonated eggs were counted. Eggs were concentrated or diluted with deionised water,

depending on the egg count. For example, eggs with 100% infectivity, approximately 50 eggs

per 50μl were counted. Mice were then infected with 200μl by oral gavage to infect mice

with approximately 200 eggs.

4. Worm burden of *Trichuris muris*

During autopsy, the caecum and proximal colon were collected and stored at -20°C

before analysis. Before worm count, the intestine was thawed at room temperature and cut

longitudinally using blunt ended scissors and the epithelium was scraped using curved

forceps in a petri dish. Worms were counted blindly under a dissecting microscope (Leica).

5. Preparation of single cell suspensions for fluorescence activated cell sorting (FACS)

Mesenteric lymph nodes (MLNs), spleen, blood and bone marrow were collected and

prepared for FACS staining. Lymph nodes and spleen were squeezed through a 70μm nylon

cell strainer (Fisher Scientific) and cells were pelleted by centrifugation at 1500 rpm for 5

minutes. The supernatant was removed and the pelleted cells were resuspended in 500μl to 1

ml of Red Blood Cell Lysing Buffer Hybri-Max™ (Sigma-Aldrich) for 30 seconds to 1

minute before adding 10 ml 1xPBS. Cells were pelleted by centrifugation at 1500 rpm for 5

minutes and resuspended in 1 ml of complete RPMI 1640 medium. Cells were counted on a

CASY cell counter (Scharfe System).

Approximately 50μl of blood were placed into 1.5 ml eppendorf tube containing 50μl

0.5M EDTA and stored on ice before analysis. 500 μl of Red Blood Cell Lysing Buffer

Hybri-Max™ (Sigma-Aldrich) were added and samples were incubated for 5 minutes at room

temperature. 1 ml of 1xPBS were added and cells were pelleted by centrifugation at 1500 rpm

for 5 minutes. Red blood cell lysis process was repeated twice before cells were resuspended

in 1 ml of complete RPMI 1640 medium.

Femurs and tibias were collected at autopsy and after removing any remaining tissue

without damaging the bone integrity, the bone was placed on ice until ready to process. Bone

was transferred to 70% ethanol for 2-3 minutes and then rinse in 3 changes of 1xPBS. In a

petri dish, both ends of bone were cut and the bone was flushed gently with 1xPBS using a

3cc syringe and a 23 ga needle. To break up the clumps, the marrow was sucked up and was

gently pushed back. Single cells suspension was filtered through 100μm nylon cell strainer

(Fisher Scientific) and cells were pelleted by centrifugation at 1500 rpm for 5 minutes and

resuspended in 1 ml of 1xPBS. The supernatant was removed and the pelleted cells were

resuspended in 500μl of Red Blood Cell Lysing Buffer Hybri-Max™ (Sigma-Aldrich) for 1

minute before adding 10 ml 1xPBS. Cells were pelleted by centrifugation at 1500 rpm for 5

minutes and resuspended in 1 ml of complete RPMI 1640 medium.

6. Cell surface markers

Cells from MLNs, spleen, blood and bone marrow were stained for live dead (Zombie

UV, Biolegend) and Fc block (eBiosciences) prior to cell surface cellular markers staining.

Samples were read on a BD LSR Fortessa flow cytometer (BD Biosciences) and data was

analysed using FlowJo X (Tree Star, Inc).

Cell surface markers: anti-B220 (RA3-6B2); anti-CD19 (6D5); anti-CD3ε (17A2);

anti-CD4 (RM4.5); anti-CD8α (53-6.7); anti-CD279 (PD-1) (29F.1A12); anti-

CD185/CXCR5 (L138D7); anti-CD93 (AA4.1); anti-CD25 (3C7); and anti-CD117/c-kit

(2B8) were purchased from Biolegend. Anti-CD43 (S11); anti-CD103 (2E7); anti-CD11c

(N418); anti-CD317/PDCA-1 (927); anti CD11b (M1/70); anti-CD64 (X54-5/7.1); anti-I-A/I-

E (M5/114.15.2); and anti-CD23 (B3B4) were purchased from BD Biosciences. Anti-CD45

(30-F11); anti-ly6G (RB6-8C5); anti-NK.1 (PK136); anti-Ter119 (Ter-119) purchased from

eBiosciences.

7. Quantification of parasite specific IgG1, IgG2a/c and IgM

To detect IgG1, IgG2a/c and IgM specific *T. muris*, an enzyme linked immunosorbant

assay (ELISA) was completed. Blood was collected from mice at autopsy and serum was

isolated by centrifuging samples for 10 minutes at 15,000g at room temperature. 96 well

immunoGrade plates (BrandTech Scientific, Inc) were coated with 5μg/ml *T. muris* diluted in

0.05M carbonate/bicarbonate buffer and incubated on plates overnight at 4°C. Plates were

washed using a Skatron Scan Washer 500 (Molecular Devices, Norway) 3 times with 0.05%

Tween 20 (Sigma) in PBS (PBS-T). Non-specific binding were blocked with 100μl 3%

bovine serum albumin (BSA) (Melford Laboratories)/ PBS at 37°C for 45 minutes. Plates

were washed 3 times with PBS-T and 50μl of double diluted serum in PBS (1:20, 1:40, 1:80,

1:160, 1:320, 1:640, 1:1280, 1:2560) was added to plates and incubated for 60 minutes. Plates

were then washed 3 times with PBS-T and 50μl of either biotinylated rat anti-mouse IgG1

(1:500, BD Bioscience), rat anti-mouse IgG2a (1:1000, BD Bioscience) or rat anti-mouse

IgM (1:500, BD Bioscience) was added to wells and incubated for 60 minutes. C57BL/6 mice

do not make IgG2a, but IgG2c. However, the rat anti-mouse IgG2a antibody also recognises

mouse IgG2c, the IgG2a equivalent. Plates were then washed with PBS-T 3 times before the

plates were incubated with 75μl Streptavidin peroxidase (1:1000, Sigma) for 60 minutes.

Plates were washed 3 times with PBS-T before the colour were developed using 100μl 0.03%

hydrogen peroxidase activated ABTS (10% 2,2'azino 3-thyl benzthiazoline in 0.045M citrate

buffer). Plates were read at 450nm on a VersaMax Microplate reader (Molecular Devices).

8. Cytokine analysis

During autopsy, mesenteric lymph nodes were isolated and collected in complete

RPMI 1640 medium. The tissue were squeezed through a 70μm nylon cell strainer (Fisher

Scientific) and cells were pelleted by centrifugation at 1500 rpm for 5 minutes. The

supernatant was removed and the pelleted cells were resuspended in 500μl to 1 ml of Red

Blood Cell Lysing Buffer Hybri-Max™ (Sigma-Aldrich) for 30 seconds to 1 minute before

adding 10 ml 1xPBS. Cells were pelleted by centrifugation at 1500 rpm for 5 minutes and

resuspended in 1 ml of complete RPMI 1640 medium. Cells were counted on a CASY cell

counter (Scharfe System) and then diluted to a concentration of $1x10^7$ cells/ml. 100μl of cells

were plated out into a 96 well flat bottom plate. Cells were restimulated with 100μl 4 hr *T.*

muris E/S at 100μg/ml (to give a final concentration of E/S in the cell culture of 50μg/ml).

Cells were incubated for 48 hours at 37°C and 5% CO2. To collect the supernatant, plates

were centrifuged at 1400g for 5 minutes and the supernatant was stored at -20°C.

The cytokines IL-4, IL-5, IL-6, IL-9, IL-10, IL-17, IL-13, TNF and IFN-γ were

detected in supernatant by Cytometric bead assay (CBA). 12.5μl of supernatant from *T. muris*

E/S stimulated cells was added to a 96-well round bottom plate. A capture bead cocktail (BD

Bioscience), containing beads for each cytokine, was diluted in capture diluent (BD

Bioscience) and 12.5μl was added to each well before incubating on a rocker at room

temperature for 1 hour. 12.5μl of detection beads (BD Bioscience), diluted in detection

reagent (BD Bioscience) was added to each well and incubated again on a rocker at room

temperature for 1 hour. Plates were washed and resuspended in 70μl wash buffer (BD

Bioscience). Cytokines were measured on a MACSQuant Analyser (Miltenyi Biotec) and

analysed using the FCAP array software in reference to a standard curve.

9. Statistical analysis

Statistical analysis was performed using Prism4 (Graph-Pad software Inc., La Jolla,

CA). The significant differences between two groups (P<0.05) were analysed with the t-test

or Mann Whitney test, depend on the n-size and the distribution of samples. For multiple

groups, the significant differences were analysed by Anova.

10. Ethics statement

All experiments were approved by The University of Manchester Local Ethical

Review Committee and were performed in accordance with the UK Home Office Animals

(Scientific Procedures) Act 1986, under the home Office project licence number 70/8127

Acknowledgment

This study was funded by the Indonesian endowment fund for education. We would like to

thank Rebecca Dookie for helping us with i.v. injection, Genentech (San Francisco, CA) for

their gift of purified anti-CD20 mAb (clone-5D2), and Gareth Howell from Flow cytometry

facility at the University of Manchester.

Conflict of interests

The authors declare no commercial or financial conflict of interest

    Figure legend

    Fig. 1. Two injections of anti CD20-mab are required to ensure B cells are depleted

    occurred throughout infection. C57BL/6 mice were treated with anti CD20 mAb or Rat

    IgG2a isotype control at 100 μg in 200 μl PBS by i.v. injection via tail vein. Mice were

    infected with approximately 150 *T. muris* eggs at day 7 post injection. Some mice were only

    injected once with anti CD20/ rat IgG2a isotype control, whilst some mice were re-injected

    i.v. with anti CD20 mAb or Rat IgG2a at 100 μg in 200 μl PBS on day 10 p.i. Mice were

    autopsied at day 35 p.i. (A&B) CD19+ cells were assessed in MLNs, spleen, blood,

    peritoneal cavity (PerC), and bone marrow (BM) using flow cytometry. (C) Representative

    flow cytometric analysis of CD19+ cells in BM of anti-CD20 treated mice. Both immature

    and mature B cells are IgM+, whilst pro/pre-B cells are IgM-. Pro-B cells are defined as

    CD19+B220^{low}IgM-CD93+CD43+CD25-ckit+. Data shown is mean ± SEM, representative

    of 2 independent experiments for Rat IgG2a and α-CD20 2x, from 1 experiment for α-CD20

    1x, n=5, males. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, Anova.

    Fig. 2. B cell depleted C57BL6 mice were susceptible to *T. muris*, correlating with an

    absence of class switched antibodies and a significant decrease in Th2 cytokines.

    C57BL/6 mice were treated with anti CD20 mAb or Rat IgG2a isotype control at 100 μg in

    200 μl PBS by i.v. injection via tail vein prior to infection. Mice were infected with

    approximately 150 *T. muris* eggs. Mice were autopsied at day 35 p.i.. (A) Diagram of

    experimental design. (B) Worm burdens were assessed blindly after autopsy. (C-E) Sera were

    analysed using ELISA for parasite specific antibodies. (F-J) MLN cells were re-stimulated

    with parasite E/S antigen for 48 hours and cytokines in supernatants were determined using

    cytokine bead array (CBA). Data shows mean ± SEM, pooled from 3 independent

experiments (B&F-K), representative of 3 independent experiments (C-E), males, *p<0.05,

**p<0.01 ***p<0.001, Student's t-test.

Fig. 3. B cell depletion alters the T$_{FH}$ population and DC subsets in MLNs. C57BL/6 mice

were treated with anti CD20 mAb or Rat IgG2a isotype control at 100 µg in 200 µl PBS by

i.v. injection via the tail vein. Mice were infected with approximately 150 *T. muris* eggs at

day 7 post injection. Mice were re-injected with anti CD20 mAb or isotype control at 100 µg

in 200 µl PBS by i.v. injection via the tail vein at day 10 p.i.. Mice were autopsied on day 0,

day 10 and day 35 p.i. for T$_{FH}$ and on day 35 p.i. for DC subsets. (A) Gating on

CD4+CXCR5+PD-1^{high} to define T$_{FH}$ cells in MLNs. (B&C) Relative % and total cell

number of T$_{FH}$ in MLNs, respectively. (D-H) DC subsets in MLNs. (E&F) Relative

percentage and total cell number of CD103-CD11b+ DCs, respectively. (G&H) Relative

percentage and total cell number of CD103+CD11b+ DCs, respectively. Data shows mean ±

SEM, from 1 experiment, n=3-5, males, *p<0.05, **p<0.01, Mann Whitney test.

Fig. 4. B cell depletion from day 14 p.i. also leads to susceptibility to *T. muris* infection.

C57BL/6 were treated with anti CD20 mAb or Rat IgG2a isotype control at 100 µg in 200 µl

PBS by i.v. injection via the tail vein 2 weeks post infection (p.i.). Mice were re-injected with

anti CD20 mAb or isotype control at 100 µg in 200 µl PBS by i.v. injection via the tail vein

on day 24 p.i. Mice were autopsied on day 35 p.i. (A) Diagram of the experiment design for

B cells depletion 2 weeks post-infection. (B) Worm burdens were assessed blindly after

autopsy. (C) Gating on CD4+CXCR5+PD-1^{high} to define T$_{FH}$ cells in MLNs. (D&E) Relative

% and total cell number of T$_{FH}$ in MLNs, respectively. (F-K) MLN cells were re-stimulated

with parasite E/S antigen for 48 hours and cytokines in supernatants were determined using

cytokine bead array (CBA). Data show mean ± SEM from 1 experiment, n=5, males,

*p<0.05, **p<0.01, Mann Whitney test.

Fig. 5. Anti-IFN-γ treatment restored resistance to *T. muris* infection and partially

rescued the IL-13 response in B cell depleted mice in the absence of *T. muris* specific

IgG1 antibodies and without preserving T_{FH} population. C57BL/6 mice were treated with

anti CD20 mAb or isotype control at 100 μg in 200 μl PBS by i.v. injection via the tail vein.

Mice were infected with approximately 150 *T. muris* eggs at day 7 post injection. Mice were

re-injected with anti CD20 mAb or isotype control at 100 μg in 200 μl PBS i.v. injection via

tail vein on day 10 p.i.. 1 mg of α-IFN-γ antibody or Rat Ig (as a control) was given on day 0,

day 7, and day 14 p.i.. Mice were autopsied on day 21 p.i. and day 35 p.i.. (A) Diagram of the

experimental design. (B) Worm burdens were assessed blindly after autopsy at day 35 p.i..

(C&D) *T. muris* specific IgG1 and IgG2c antibodies in the sera by day 21 p.i., respectively.

(E-H) Cytokine analysis of re-stimulated MLN cells day 21 p.i.. (I) Gating on

CD4+CXCR5+PD-1^{high} to define T_{FH} cells in MLNs day 21 p.i.. (J) Total numbers of T_{FH}

cells in MLNs day 21 p.i.. Data shows mean ± SEM, from 1 experiment, n=5, males,

*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001, Anova.

Fig. 6. BALB/c mice were able to expel *T. muris* in the absence of B cells. BALB/c mice

were treated with anti CD20 mAb or Rat IgG2a isotype control at 100 μg in 200 μl PBS by

i.v. injection via tail vein prior to infection. Mice were infected with approximately 150 *T.*

muris eggs and were autopsied at day 42 p.i.. (A) Diagram of experimental design. (B)

CD19+ cells were assessed in MLNs, spleen, and peritoneal cavity (PerC) using flow

cytometry. (C) Worm burdens were assessed blindly after autopsy. (D-F) Sera were analysed

590 using ELISA for parasite specific antibodies. Data shows mean ± SEM, from 1 experiment,

591 n=5, males. *p<0.05, ***p<0.001 ****p<0.0001, Student's t-test.

1.   Hotez PJ, Brindley PJ, Bethony JM, King CH, Pearce EJ, Jacobson J. Helminth infections: the
great neglected tropical diseases. The Journal of clinical investigation. 2008;118(118 (4)):1311-21.
2.   Zhan B, Beaumier CM, Briggs N, Jones KM, Keegan BP, Bottazzi ME, et al. Advancing a
multivalent'Pan-anthelmintic'vaccine against soil-transmitted nematode infections. Expert review of
vaccines. 2014;13(3):321-31.
3.   Bethony J, Brooker S, Albonico M, Geiger SM, Loukas A, Diemert D, et al. Soil-transmitted
helminth infections: ascariasis, trichuriasis, and hookworm. The Lancet. 2006;367(9521):1521-32.
4.   Constant S, Schweitzer N, West J, Ranney P, Bottomly K. B lymphocytes can be competent
antigen-presenting cells for priming CD4+ T cells to protein antigens in vivo. The Journal of
Immunology. 1995;155(8):3734-41.
5.   Liu Y, Wu Y, Ramarathinam L, Guo Y, Huszar D, Trounstine M, et al. Gene-targeted B-
deficient mice reveal a critical role for B cells in the CD4 T cell response. International immunology.
1995;7(8):1353-62.
6.   Magee CN, Boenisch O, Najafian N. The role of costimulatory molecules in directing the
functional differentiation of alloreactive T helper cells. American Journal of Transplantation.
2012;12(10):2588-600.
7.   Lund FE, Garvy BA, Randall TD, Harris DP. Regulatory roles for cytokine-producing B cells in
infection and autoimmune disease. 2005.
8.   Williams M, Chang T, Burke S, Lichtman A, Abbas A. Activation of functionally distinct
subsets of CD4+ T lymphocytes. Research in immunology. 1991;142(1):23-8.
9.   Harris DP, Goodrich S, Gerth AJ, Peng SL, Lund FE. Regulation of IFN-γ production by B
effector 1 cells: essential roles for T-bet and the IFN-γ receptor. The Journal of Immunology.
2005;174(11):6781-90.
10.   Wojciechowski W, Harris DP, Sprague F, Mousseau B, Makris M, Kusser K, et al. Cytokine-
producing effector B cells regulate type 2 immunity to H. polygyrus. Immunity. 2009;30(3):421-33.
11.   Harris DP, Goodrich S, Mohrs K, Mohrs M, Lund FE. Cutting edge: the development of IL-4-
producing B cells (B effector 2 cells) is controlled by IL-4, IL-4 receptor α, and Th2 cells. The Journal
of Immunology. 2005;175(11):7103-7.
12.   Hurdayal R, Ndlovu HH, Revaz-Breton M, Parihar SP, Nono JK, Govender M, et al. IL-4–
producing B cells regulate T helper cell dichotomy in type 1-and type 2-controlled diseases.
Proceedings of the National Academy of Sciences. 2017;114(40):E8430-E9.
13.   Linton P-J, Bautista B, Biederman E, Bradley ES, Harbertson J, Kondrack RM, et al.
Costimulation via OX40L expressed by B cells is sufficient to determine the extent of primary CD4 cell
expansion and Th2 cytokine secretion in vivo. Journal of Experimental Medicine. 2003;197(7):875-
83.
14.   Liu Q, Liu Z, Rozo CT, Hamed HA, Alem F, Urban JF, et al. The role of B cells in the
development of CD4 effector T cells during a polarized Th2 immune response. The Journal of
Immunology. 2007;179(6):3821-30.
15.   León B, Ballesteros-Tato A, Browning JL, Dunn R, Randall TD, Lund FE. Regulation of TH2
development by CXCR5+ dendritic cells and lymphotoxin-expressing B cells. Nature immunology.
2012;13(7):681-90.
16.   Yoichi I. The absence of resistance in congenitally athymic nude mice toward infection with
the intestinal nematode, Trichuris muris: resistance restored by lymphoid cell transfer. International
journal for parasitology. 1991;21(1):65-9.
17.   KOYAMA K, TAMAUCHI H, ITO Y. The role of CD4+ and CD8+ T cells in protective immunity to
the murine nematode parasite Trichuris muris. Parasite immunology. 1995;17(3):161-5.
18.   Else KJ, Grencis RK. Antibody-independent effector mechanisms in resistance to the
intestinal nematode parasite Trichuris muris. Infection and immunity. 1996;64(8):2950-4.
19.   Betts C, Else K. Trichuris muris: CD4+ T cell-mediated protection in reconstituted SCID mice.
Parasitology. 2000;121(6):631-7.

20.   Little MC, Bell LV, Cliffe LJ, Else KJ. The characterization of intraepithelial lymphocytes,
lamina propria leukocytes, and isolated lymphoid follicles in the large intestine of mice infected with
the intestinal nematode parasite Trichuris muris. The Journal of Immunology. 2005;175(10):6713-22.
21.   Svensson M, Russell K, Mack M, Else KJ. CD4+ T-cell localization to the large intestinal
mucosa during Trichuris muris infection is mediated by Gαi-coupled receptors but is CCR6-and
CXCR3-independent. Immunology. 2010;129(2):257-67.
22.   Blackwell NM, Else KJ. B cells and antibodies are required for resistance to the parasitic
gastrointestinal nematode Trichuris muris. Infection and immunity. 2001;69(6):3860-8.
23.   Fu Y-X, Chaplin DD. Development and maturation of secondary lymphoid tissues. Annual
review of immunology. 1999;17(1):399-433.
24.   Golovkina TV, Shlomchik M, Hannum L, Chervonsky A. Organogenic role of B lymphocytes in
mucosal immunity. Science. 1999;286(5446):1965-8.
25.   Cheson BD, Leonard JP. Monoclonal antibody therapy for B-cell non-Hodgkin's lymphoma.
New England Journal of Medicine. 2008;359(6):613-26.
26.   Shahaf G, Zisman-Rozen S, Benhamou D, Melamed D, Mehr R. B Cell Development in the
Bone Marrow Is Regulated by Homeostatic Feedback Exerted by Mature B Cells. Frontiers in
immunology. 2016;7:77.
27.   Uchida J, Lee Y, Hasegawa M, Liang Y, Bradney A, Oliver JA, et al. Mouse CD20 expression
and function. International immunology. 2004;16(1):119-29.
28.   Uchida J, Hamaguchi Y, Oliver JA, Ravetch JV, Poe JC, Haas KM, et al. The innate
mononuclear phagocyte network depletes B lymphocytes through Fc receptor–dependent
mechanisms during anti-CD20 antibody immunotherapy. Journal of Experimental Medicine.
2004;199(12):1659-69.
29.   Chan AC, Carter PJ. Therapeutic antibodies for autoimmunity and inflammation. Nature
Reviews Immunology. 2010;10(5):301.
30.   Misumi I, Whitmire JK. B cell depletion curtails CD4+ T cell memory and reduces protection
against disseminating virus infection. The Journal of Immunology. 2014;192(4):1597-608.
31.   Little MC, Hurst RJ, Else KJ. Dynamic changes in macrophage activation and proliferation
during the development and resolution of intestinal inflammation. The Journal of Immunology.
2014:1400502.
32.   Bancroft AJ, McKenzie AN, Grencis RK. A critical role for IL-13 in resistance to intestinal
nematode infection. The Journal of Immunology. 1998;160(7):3453-61.
33.   Zaretsky AG, Taylor JJ, King IL, Marshall FA, Mohrs M, Pearce EJ. T follicular helper cells
differentiate from Th2 cells in response to helminth antigens. The Journal of experimental medicine.
2009;206(5):991-9.
34.   Else K, Grencis R. Cellular immune responses to the murine nematode parasite Trichuris
muris. I. Differential cytokine production during acute or chronic infection. Immunology.
1991;72(4):508.
35.   Bancroft AJ, Else KJ, Sypek JP, Grencis RK. Interleukin-12 promotes a chronic intestinal
nematode infection. European journal of immunology. 1997;27(4):866-70.
36.   Xu X, Shi Y, Cai Y, Zhang Q, Yang F, Chen H, et al. Inhibition of increased circulating Tfh cell by
anti-CD20 monoclonal antibody in patients with type 1 diabetes. PLoS One. 2013;8(11):e79858.
37.   Demiri M, Müller-Luda K, Agace WW, Svensson-Frej M. Distinct DC subsets regulate adaptive
Th1 and 2 responses during Trichuris muris infection. Parasite immunology. 2017;39(10):e12458.
38.   Cruickshank SM, Deschoolmeester ML, Svensson M, Howell G, Bazakou A, Logunova L, et al.
Rapid dendritic cell mobilization to the large intestinal epithelium is associated with resistance to
Trichuris muris infection. The Journal of Immunology. 2009;182(5):3055-62.
39.   Else K, Finkelman F, Maliszewski C, Grencis R. Cytokine-mediated regulation of chronic
intestinal helminth infection. The Journal of experimental medicine. 1994;179(1):347-51.
40.   King IL, Mohrs M. IL-4–producing CD4+ T cells in reactive lymph nodes during helminth
infection are T follicular helper cells. Journal of Experimental Medicine. 2009;206(5):1001-7.

41.   Sahoo A, Alekseev A, Tanaka K, Obertas L, Lerman B, Haymaker C, et al. Batf is important for
IL-4 expression in T follicular helper cells. Nature communications. 2015;6:7997.

42.   Else K, Wakelin D. The effects of H-2 and non-H-2 genes on the expulsion of the nematode
Trichuris muris from inbred and congenic mice. Parasitology. 1988;96(03):543-50.

43.   Else K, Entwistle G, Grencis R. Correlations between worm burden and markers of Th1 and
Th2 cell subset induction in an inbred strain of mouse infected with Trichuris muris. Parasite
immunology. 1993;15(10):595-600.

44.   Tedder T, Klejman G, Disteche C, Adler D, Schlossman S, Saito H. Cloning of a complementary
DNA encoding a new mouse B lymphocyte differentiation antigen, homologous to the human B1
(CD20) antigen, and localization of the gene to chromosome 19. The Journal of Immunology.
1988;141(12):4388-94.

45.   Tedder TF, Forsgren A, Boyd AW, Nadler LM, Schlossman SF. Antibodies reactive with the B1
molecule inhibit cell cycle progression but not activation of human B lymphocytes. European journal
of immunology. 1986;16(8):881-7.

46.   Ait-Oufella H, Herbin O, Bouaziz J-D, Binder CJ, Uyttenhove C, Laurans L, et al. B cell
depletion reduces the development of atherosclerosis in mice. Journal of Experimental Medicine.
2010;207(8):1579-87.

47.   Wang C, Hillsamer P, Kim CH. Phenotype, effector function, and tissue localization of PD-1-
expressing human follicular helper T cell subsets. BMC immunology. 2011;12(1):53.

48.   Crawford A, MacLeod M, Schumacher T, Corlett L, Gray D. Primary T cell expansion and
differentiation in vivo requires antigen presentation by B cells. The Journal of Immunology.
2006;176(6):3498-506.

49.   Hernandez HJ, Wang Y, Stadecker MJ. In infection with Schistosoma mansoni, B cells are
required for T helper type 2 cell responses but not for granuloma formation. The Journal of
Immunology. 1997;158(10):4832-7.

50.   Moulin V, Andris F, Thielemans K, Maliszewski C, Urbain J, Moser M. B lymphocytes regulate
dendritic cell (DC) function in vivo: increased interleukin 12 production by DCs from B cell–deficient
mice results in T helper cell type 1 deviation. Journal of Experimental Medicine. 2000;192(4):475-82.

51.   Elsegeiny W, Eddens T, Chen K, Kolls JK. Anti-CD20 antibody therapy and susceptibility to
Pneumocystis pneumonia. Infection and immunity. 2015;83(5):2043-52.

52.   Choi YS, Kageyama R, Eto D, Escobar TC, Johnston RJ, Monticelli L, et al. ICOS receptor
instructs T follicular helper cell versus effector cell differentiation via induction of the transcriptional
repressor Bcl6. Immunity. 2011;34(6):932-46.

53.   Johansson-Lindbom B, Borrebaeck CA. Germinal center B cells constitute a predominant
physiological source of IL-4: implication for Th2 development in vivo. The Journal of Immunology.
2002;168(7):3165-72.

54.   Appleton J, McGregor D. Characterization of the immune mediator of rapid expulsion of
Trichinella spiralis in suckling rats. Immunology. 1987;62(3):477.

55.   Harris N, Pleass R, Behnke J. Understanding the role of antibodies in murine infections with
Heligmosomoides (polygyrus) bakeri: 35 years ago, now and 35 years ahead. Parasite immunology.
2014;36(3):115-24.

56.   Pleass RJ, Behnke JM. B-cells get the T-cells but antibodies get the worms. Trends in
parasitology. 2009;25(10):443-6.

57.   McCoy KD, Stoel M, Stettler R, Merky P, Fink K, Senn BM, et al. Polyclonal and specific
antibodies mediate protective immunity against enteric helminth infection. Cell host & microbe.
2008;4(4):362-73.

58.   Lund FE. Cytokine-producing B lymphocytes—key regulators of immunity. Current opinion in
immunology. 2008;20(3):332-8.

59.   Lu P, Urban JF, Zhou X, Chen S, Madden K, Moorman M, et al. CD40-mediated stimulation
contributes to lymphocyte proliferation, antibody production, eosinophilia, and mastocytosis during

an in vivo type 2 response, but is not required for T cell IL-4 production. The Journal of Immunology.
1996;156(9):3327-33.
60. Liu Q, Kreider T, Bowdridge S, Liu Z, Song Y, Gaydo AG, et al. B cells have distinct roles in host
protection against different nematode parasites. The Journal of Immunology. 2010;184(9):5213-23.
61. Ligas JA, Kerepesi LA, Galioto AM, Lustigman S, Nolan TJ, Schad GA, et al. Specificity and
mechanism of immunoglobulin M (IgM)-and IgG-dependent protective immunity to larval
Strongyloides stercoralis in mice. Infection and immunity. 2003;71(12):6835-43.
62. BETTS CJ, ELSE KJ. Mast cells, eosinophils and antibody-mediated cellular cytotoxicity are not
critical in resistance to Trichuris muris. Parasite immunology. 1999;21(1):45-52.
63. Rajan B, Ramalingam T, Rajan TV. Critical Role for IgM in Host Protection in Experimental
Filarial Infection. The Journal of Immunology. 2005;175(3):1827-33.
64. Inaba T, Sato H, Kamiya H. Monoclonal IgA antibody-mediated expulsion of Trichinella from
the intestine of mice. Parasitology. 2003;126(6):591-8.
65. Harris N, Gause WC. To B or not to B: B cells and the Th2-type immune response to
helminths. Trends in immunology. 2011;32(2):80-8.
66. Harris N, Gause WC. To B or not to B: B cells and the Th2-type immune response to
helminths. Trends Immunol. 2011;32(2):80-8.
67. Bradford BM, Donaldson DS, Forman R, Else KJ, Mabbott NA. Increased susceptibility to oral
Trichuris muris infection in the specific absence of CXCR 5+ CD 11c+ cells. Parasite Immunology.
2018:e12566.
68. Tew JG, Wu J, Fakher M, Szakal AK, Qin D. Follicular dendritic cells: beyond the necessity of
T-cell help. Trends in immunology. 2001;22(7):361-7.
69. Katakai T, Suto H, Sugai M, Gonda H, Togawa A, Suematsu S, et al. Organizer-like reticular
stromal cell layer common to adult secondary lymphoid organs. The Journal of Immunology.
2008;181(9):6189-200.
70. Mayer JU, Demiri M, Agace WW, MacDonald AS, Svensson-Frej M, Milling SW. Different
populations of CD11b+ dendritic cells drive Th2 responses in the small intestine and colon. Nature
communications. 2017;8:15820.
71. Ziętara N, Łyszkiewicz M, Puchałka J, Pei G, Gutierrez MG, Lienenklaus S, et al.
Immunoglobulins drive terminal maturation of splenic dendritic cells. Proceedings of the National
Academy of Sciences. 2013;110(6):2282-7.
72. Sahputra R, Yam-Puc JC, Waisman A, Muller W, Else KJ. Evaluating the IgMi mouse as a novel
tool to study B cell biology. European journal of immunology. 2018.
73. Wakelin D. Acquired immunity to Trichuris muris in the albino laboratory mouse.
Parasitology. 1967;57(3):515-24.

A

B

C

Bone marrow-α-CD20 treated mice

Supplementary Fig.1. Flow cytometry gating strategy for T$_{FH}$. T$_{FH}$ cells are defined as CD4+CXCR5+PD-1^{high}

Supplementary Fig. 2. T_{FH} population in the spleen of anti-CD20 mAb treated mice. C57BL/6 mice were treated with anti CD20 mab or isotype control 100 μg in 200 μl PBS i.v. injection via tail vein. Mice were infected with approximately 150 *T. muris* eggs at day 7 post injection. Mice were re-injected with anti CD20 mab or isotype control 100 μg in 200 μl PBS i.v. injection via tail vein by day 10 p.i. Mice were autopsied by day 0, day 10 and day 35 p.i. Gating on CD4+CXCR5+PD-1^{high} to define T$_{FH}$ cells in spleen (A). (B&C) Relative % and total cell number of T$_{FH}$ in the spleen, respectively. Data show mean±SEM, from 1 experiment, males., *p<0.05, Mann whitney U test

Supplementary Fig.3. Flow cytometry gating strategy for DC subsets. DCs are defined as CD45+lineage-CD11c+MHCII+CD64-.Two main DCs populations: conventional DCs (cDCs) and plasmacytoid DCs (pDCs). Conventional DCs (CD45+lineage-CD11c+MHCII+CD64-PDCA1-) were divided into migratory DCs and resident DCs. Migratory DCs was consisted of 4 subpopulations: CD103+CD11b-, CD103+CD11b+, CD103-CD11b+ and CD103-CD11b-. Plasmacytoid DCs:CD45+lineage-CD11c+MHCII+CD64-PDCA1+

Supplementary Fig. 4. Tfh population in the MLNs and spleen of anti-CD20 mab treated mice on a Balb/c genetic background were not affected. Balb/c mice were treated with anti CD20 mab or isotype control 100 μg in 200 μl PBS i.v. injection via tail vein. Mice were infected with approximately 150 *T. muris* eggs at day 7 post injection. Mice were re-injected with anti CD20 mab or isotype control 100 μg in 200 μl PBS i.v. injection via tail vein by day 10 p.i. Mice were autopsied by day 0 and day 42 p.i. Gating on CD4+CXCR5+PD-1^high to define Tfh cells in MLNs (A) and spleen (B). (C&D) Relative % of Tfh in MLNs and spleen, respectively. (E&F) Total Tfh in MLNs and spleen, respectively. . Data show mean±SEM, from 1 experiment, males.