1 **"Self-inactivating" rabies viruses are susceptible to loss of their intended attenuating**
2 **modification**
3

4 Lei Jin[1], Makoto Matsuyama[1], Heather A. Sullivan[1], Mulangma Zhu[1], Thomas K. Lavin[1],
5 YuanYuan Hou[1], Nicholas E. Lea[1], Maxwell T. Pruner[1], María Lucía Dam Ferdínez[1], and Ian R.
6 Wickersham[1]*
7
8 [1]McGovern Institute for Brain Research, Massachusetts Institute of Technology, Cambridge, MA
9
10 *Corresponding author (wickersham@mit.edu).

11
12
13 **Competing interests statement**
14 I.R.W. is a consultant for Monosynaptix, LLC, advising on design of neuroscientific experiments.
15

ABSTRACT

Monosynaptic tracing using rabies virus is an important technique in neuroscience, allowing brain-wide labeling of neurons directly presynaptic to a targeted neuronal population. A 2017 article reported development of a noncytotoxic version – a major advance – based on attenuating the rabies virus by addition of a destabilization domain to the C-terminus of a viral protein. However, this modification did not appear to hinder the ability of the virus to spread between neurons. We analyzed two viruses provided by the authors and show here that both were mutants that had lost the intended modification, explaining the paper's paradoxical results. We then made a virus that actually did have the intended modification in at least the majority of virions and found that it did not spread efficiently under the conditions described in the original paper, namely, without an exogenous protease being expressed in order to remove the destabilization domain. We found that it did spread when the protease was supplied, although this also appeared to result in the deaths of most source cells by three weeks postinjection. We conclude that the new approach is not robust but that it could become a viable technique given further optimization and validation.

SIGNIFICANCE STATEMENT

Rabies virus, which spreads between synaptically-connected neurons, has been one of the primary tools used by neuroscientists to reveal the organization of the brain. A new modification to rabies virus was recently reported to allow the mapping of connected neurons without adverse effects on the cells' health, unlike earlier versions. Here we show that the conclusions of that study were probably incorrect and based on having used viruses that had lost the intended modification because of mutations. We also show that a rabies virus that does retain the intended modification does not spread efficiently between neurons under the conditions reported previously; however, it does spread between neurons under different conditions, suggesting that the approach may be successful if refined further.

INTRODUCTION

Viruses have become important tools for neuroscience (1-16), and "monosynaptic tracing" based on rabies virus (5) has become the primary method of labeling neurons directly presynaptic to some targeted group of neurons (17-20). Its core components are, first, selective infection of the targeted neuronal group with a recombinant rabies virus with a deleted gene (which in all work published to date is the "G" gene encoding the glycoprotein that coats the viral envelope) and, second, complementation of the deletion in the targeted starting neurons, by expression of the deleted gene in *trans*. With all of its gene products therefore present in the starting cells, the virus can fully replicate within them and spreads, as wild-type rabies virus does, to cells directly presynaptic to the initially-infected neurons. Assuming that G has not been provided in *trans* in these presynaptic cells too, the deletion-mutant ("ΔG", denoting the deletion of G) virus is unable to spread beyond them, resulting in labeling of just the neurons in the initially-targeted population and ones that are directly presynaptic to them (5).

A drawback of these ΔG (or "first-generation" (21)) rabies viruses is that they are cytotoxic (4, 21, 22), which has spurred several labs to develop less toxic versions. Reardon, Murray, and colleagues (22) showed that simply using ΔG rabies virus of a different parent strain — switching from the original SAD B19 strain to the more neuroinvasive CVS-N2c strain (23) — decreased toxicity and increased the efficiency of transneuronal spread. Our own group has taken a more drastic approach, introducing nontoxic "second-generation" rabies viruses from which both G and a second gene, "L", encoding the viral polymerase, have been deleted (21); when complemented

by expression of both deleted genes *in trans*, these double-deletion-mutant vectors spread to
presynaptic neurons, just as with the first-generation system (24).
Taking a quite different approach, Ciabatti et al. (25) introduced "self-inactivating rabies"
("SiR") viruses, which differed from simple first-generation (ΔG, SAD B19 strain) ones by the
addition of a destabilization domain to the C-terminus of one of the viral proteins, so that the
protein would be rapidly degraded soon after it was produced. Because the protein in question
(the nucleoprotein, encoded by the "N" gene) is essential for viral gene expression and replication,
its destabilization was intended to "silence" viral gene expression and prevent replication, making
the viruses nontoxic.
These SiR viruses were designed to be unable to replicate unless an exogenous protease
(tobacco etch virus protease, TEVP) was expressed in infected cells in order to remove the
destabilization (or "PEST" (26)) domain. However, the paper reported that they were able to
spread between neurons – which requires replication – just as efficiently as unmodified first-
generation viruses did, without the protease being provided at all.
We hypothesized that the viruses that were used for the reported transsynaptic tracing
experiments (25) were mutants with premature stop codons at or near the end of the native
nucleoprotein gene and before the sequence of the destabilization domain. Rhabdoviruses have
high mutation rates (27-32), and production of high-titer rabies virus stocks for *in vivo* injection
typically involves repeated passaging on complementing cell lines (33-35), which affords ample
opportunity for accumulation of mutants with a selective replication advantage.
Here we show that, in both of the two SiR virus samples that we analyzed, the great
majority of viral particles did have mutations in their genomes that caused the complete loss of
the intended C-terminal addition to the nucleoprotein, so that they were effectively just ordinary
first-generation ΔG rabies viral vectors. We also tested the SiR-CRE virus *in vivo* and found that
it was rapidly cytotoxic.
We also show that a ΔG virus that does have the intended modification to the
nucleoprotein does not spread efficiently *in vivo* in the absence of TEVP, but that it does spread
efficiently when TEVP is provided.
95
96
97 **RESULTS**
98
We analyzed samples of two viruses sent directly from the Tripodi lab to MIT two months
after their publication(25) and given directly to the Wickersham lab soon afterward, still frozen and
unopened, with express permission from Marco Tripodi by email on December 5, 2017:
"EnvA/SiR-CRE" (made from genome plasmid Addgene 99608, pSAD-F3-NPEST-iCRE-2A-
mCherryPEST) and "EnvA/SiR-FLPo" (made from genome plasmid Addgene 99609, pSAD-
F3-NPEST-FLPo-2A-mCherryPEST). Both had the SAD B19 strain of rabies virus as their parent
strain and had been packaged with the avian and sarcoma virus subgroup A envelope
glycoprotein ("EnvA") for targeted infection of cells expressing EnvA's receptor, TVA (5).
For comparison with the two SiR viruses, we made five control viruses in our own
laboratory: three first-generation vectors, RVΔG-4Cre (21), RVΔG-4FLPo (see Methods), and
RVΔG-4mCherry (36), and two second-generation vectors, RVΔGL-4Cre and RVΔGL-4FLPo
(21). All of these viruses are on the SAD B19 background, like the SiR viruses. For each of the
four recombinase-expressing viruses from our laboratory, we made one preparation packaged
with the EnvA envelope protein and one preparation packaged with the native rabies virus (SAD
B19 strain) glycoprotein (denoted as "B19G"); RVΔG-4mCherry (used only as a control for the
Sanger sequencing) was only packaged with the EnvA envelope protein.
115
116 **Sequencing of viral genomes: Sanger sequencing**
117

In order to directly test our hypothesis that the SiR viruses had developed premature stop
codons removing the PEST domain in a majority of viral particles, we sequenced the genomes of
a large number of individual viral particles using two different techniques.
First, we used ordinary Sanger sequencing to determine the sequence in the vicinity of
the end of the nucleoprotein gene for 50 to 51 individual viral particles of each of the two SiR
viruses and of a first-generation virus from our own laboratory, RVΔG-4mCherry (Figure 1 and
Supplementary File S1). We ensured the isolation of individual viral genomes by using a primer
with a random 8-base index for the reverse transcription step, so that the cDNA copy of each RNA
viral genome would have a unique index. Following the reverse transcription step, we amplified
the genomes by standard PCR, cloned the amplicons into a generic plasmid, transformed this
library into E.coli and sequenced plasmids purified from individual colonies.
As shown in Figure 1, the results confirmed our hypothesis that SiR viruses are prone to
loss of the 3' addition to the nucleoprotein gene. Specifically, in the SiR-CRE sample, 100% of
the 51 sequenced viral particles had lost the PEST domain. Fifty out of the 51 had the same point
mutation in the linker between the end of the native nucleoprotein gene and the TEVP cleavage
site, converting a glycine codon (GGA) to a stop codon (TGA) so that the only modification to the
C-terminus of the nucleoprotein was the addition of two amino acids (a glycine and a serine). The
one sequenced viral particle that did not have this point mutation had a single-base insertion in
the second-to-last codon of the native nucleoprotein gene, frameshifting the rest of the sequence
and resulting in 15 amino acids of nonsense followed by a stop codon before the start of the PEST
domain sequence.
In the SiR-FLPo sample, the population was more heterogeneous: out of 50 sequenced
viral particles, 18 had the same stop codon that was found in almost all genomes in the Cre
sample, while another 28 had a different stop codon three amino acids upstream, immediately at
the end of the native nucleoprotein gene (converting a serine codon (TCA) to a stop codon (TGA)).
Four viral particles had no mutations in the sequenced region. Thus 46/50 (92%) of the SiR-FLPo
viral particles sequenced had lost the PEST domain.
In contrast, in the first-generation virus from our own lab, RVΔG-4mCherry, none of the 50
viral particles sequenced had mutations in the sequenced region containing the end of the
nucleoprotein gene.
148
149   **Sequencing of viral genomes: Single-molecule, real-time (SMRT) sequencing**
150
As a second approach to analyzing the mutations present in the SiR viruses, we employed
a large-scale sequencing technology: single-molecule, real-time ("SMRT") sequencing, which
provides independent sequences of tens of thousands of individual single molecules in a sample
in parallel (Figure 2 and Supplementary File S2). The results from this advanced sequencing
method were quite consistent with the results from the Sanger sequencing presented above. As
with the sample preparation for Sanger sequencing, we included a random index (10 bases, in
this case) in the reverse transcription primer, so that again the cDNA copy of each RNA viral
genome molecule would be labeled with a unique index.
SMRT sequencing entails circularization of the DNA amplicons and multiple consecutive
passes around the resulting circular molecule, with the redundancy provided by this repeated
sequencing of each position increasing the signal to noise ratio and statistical significance of the
results. The numbers presented in Figure 2 and below use the default of including only clones
that had at least three reads of each base ("circular consensus sequence 3", or "CCS3" in
Supplementary File S2). Using the increasingly stringent criteria of requiring either five or eight
reads per base (CCS5 or CCS8) reduced the numbers of qualifying genomes in all cases and
changed the percentages slightly but gave very similar results overall. Because read accuracy for
SMRT sequencing is ≥98% for circular consensus sequencing with 3 passes (see
https://www.mscience.com.au/upload/pages/pacbio/technical-note---experimental-design-for-

targeted-sequencing.pdf), we used a conservative threshold of 2% frequency of any given point
mutation position in order to screen out false positives. Also to be very conservative, for Figure 2
we ignored all apparent frame shifts caused by insertions and deletions, because insertions in
particular are prone to false positives with SMRT sequencing (37). See Supplementary File S2
for details, including details of frameshifts due to insertions; Supplementary Files S3-S5 contain
the sequences of the PCR amplicons that would be expected based on published sequences of
the three viruses, but to summarize here:
As a control, we used a virus from our own laboratory, RVΔG-4Cre (21) (see Addgene
#98034 for reference sequence). Out of 17,978 sequenced genomes of this virus, we found no
mutations above threshold frequency at the end of N. We did find that 1,706 viral particles (9.49%)
had a nonsynonymous mutation (TCT (Ser) → ACT (Thr)) farther up in N at amino acid position
419 (31 amino acids upstream of the end of the 450-aa native protein). We do not know if this
mutation is functionally significant, although it is not present in CVS-N2c (38), HEP-Flury (39),
ERA (40), or Pasteur strains (Genbank GU992320), so these particles may effectively be N-
knockouts that were propagated by coinfection with virions with intact N (see Discussion for more
on such parasitic co-propagating mutants).
For the SiR-CRE virus, out of 22,205 viral genomes sequenced, 22,032 had the premature
stop codon (GGA -> TGA) in the linker between the native nucleoprotein gene and the TEVP
cleavage site sequence. In other words, even without including frameshifts, at least 99.22% of
the individual viral particles in the SiR-CRE sample were essentially first-generation ΔG vectors,
with the only modification of the nucleoprotein being an additional two amino acids at the C-
terminus.
For the SiR-FLPo virus, out of 17,086 viral genomes sequenced, 5,979 had the stop codon
(GGA -> TGA) in the linker, 8,624 had the stop codon (TCA -> TGA) at the end of N, and a further
28 had a different stop codon (TCA -> TAA) at the same position at the end of N. Of these, 305
viral particles had premature stop codons at both of these two positions, so that the total number
of viral particles with one or both stop codons immediately before the PEST domain was (8624 +
5979 + 28 - 305 = 14,326. In other words, again even without including frameshifts, at least
83.85% of the individual viral particles in the SiR-FLPo sample were essentially first-generation
ΔG vectors, with the only modification of the nucleoprotein being either two amino acids added
to, or one amino acid lost from, the C-terminus.
200
201 **Anti-nucleoprotein immunostaining**
202
We infected reporter cell lines with serial dilutions of the two EnvA-enveloped SiR viruses
as well as the eight recombinase-expressing ones from our own lab: ΔG vs. ΔGL, Cre vs. FLPo,
EnvA vs. B19G envelopes. Three days later, we immunostained the cells for rabies virus
nucleoprotein and imaged the cells with confocal microscopy.
As seen in Supplementary Figure S1, we found that the cells infected with the SiR viruses
looked very similar to those infected with the first-generation, ΔG viruses. Notably, the viral
nucleoprotein, which in the SiR viruses is intended to be destabilized and degrade rapidly in the
absence of TEVP, accumulated in the SiR-infected cells in clumpy distributions that looked very
similar to those in the cells infected with the first-generation, ΔG viruses. By contrast, the cells
infected with the second-generation, ΔGL viruses, which we have shown to be noncytotoxic (21),
did not show any such nucleoprotein accumulation, clumped or otherwise, only punctate labeling
presumably indicating isolated viral particles or post-infection uncoated viral particles
(ribonucleoprotein complexes) that are not replicating.
216
217 **Longitudinal two-photon imaging *in vivo***
218

To see whether the SiR viruses kill neurons in the brain, we conducted longitudinal two-
photon imaging *in vivo* of virus-labeled neurons in visual cortex of tdTomato reporter mice, as we
had done previously to demonstrate the nontoxicity of second-generation rabies virus (21) (Figure
3). Because the SiR viruses were EnvA-enveloped, we first injected a lentivirus expressing EnvA's
receptor TVA, then one week later we injected either SiR-CRE or one of two EnvA-enveloped
viruses made in our laboratory: the first-generation virus RVΔG-4Cre(EnvA) or the second-
generation virus RVΔGL-4Cre(EnvA). Beginning one week after rabies virus injection, we imaged
labeled neurons at the injection site every seven days for four weeks, so that we could track the
fate of individual neurons over time.
As we found in our previous work (21), our second-generation virus RVΔGL-4Cre did not
kill neurons to any appreciable degree: all but a tiny handful of the neurons labeled by this virus
at seven days after injection were still present three weeks later in all mice. Also as we have found
previously (21), our first-generation virus RVΔG-4Cre did kill neurons, but by no means all of them
(see the Discussion for possible reasons for this).
However, we found that the putatively nontoxic SiR-CRE caused a steep loss of neurons
much more pronounced than even our first-generation virus did. By 14 days after injection, 70%
of cells seen at seven days were dead; by 28 days, 81% were.
There is a possible confound from our use of the tdTomato reporter line Ai14 (which we
used primarily because we already had large numbers of mice of this line): because SiR-CRE is
actually "SiR-iCRE-2A-mCherryPEST", designed to coexpress mCherry (with an added C-
terminal PEST domain intended to destabilize it, as for the nucleoprotein) along with Cre, it is
conceivable that some of the SiR-CRE-labeled red cells at seven days were only expressing
mCherry and not tdTomato. If the destabilized mCherry were expressed only transiently, as
intended(25), and a significant fraction of SiR-CRE virions had mutations in the Cre gene so that
they did not express functioning Cre, then it is possible that some of the red cells seen at seven
days were labeled only with mCherry that stopped being visible by 14 days, so that it would only
look like those cells had died.
We viewed this alternative explanation as unlikely, because the designers of SiR-CRE
had injected it in an EYFP reporter line and found no cells labeled only with mCherry and not
EYFP at six days and nine days postinjection (see Figure S4 in Ciabatti et al. (25)). Nevertheless,
we addressed this potential objection in several ways.
First, we sequenced the transgene inserts (iCre-P2A-mCherryPEST) of 21 individual SiR-
CRE viral particles (see Supplementary File S6) and found that only two out of 21 had mutations
in the Cre gene, suggesting that there would not have been a large population of cells only labeled
by mCherry and not by tdTomato.
Second, we repeated some of the SiR-CRE injections and imaging in a different reporter
line: Ai35, expressing Arch-EGFP-ER2 after Cre recombination (41) (Jax 012735). Although we
found that the membrane-localized green fluorescence from the Arch-EGFP-ER2 fusion protein
was too dim and diffuse at seven days postinjection to be imaged clearly, we were able to obtain
clear images of a number of cells at 11 days postinjection. We found that 46% of them had
disappeared only three days later (see Supplementary Figure S2 and Supplementary Video S1),
and 86% had disappeared by 28 days postinjection, consistent with a rapid die-off. Furthermore,
we found that the red fluorescence in Ai35 mice, which was due only to the mCherry expressed
by the virus, was much dimmer than the red fluorescence in Ai14 mice at the same time point of
seven days postinjection and with the same imaging parameters (see Supplementary Figure S3):
the mean intensity was 45.86 (arbitrary units, or "a.u.") in Ai14 but only 16.29 a.u. in Ai35. This is
consistent with the published findings that tdTomato is a much brighter fluorophore than mCherry
(42), particularly with two-photon excitation (43), and it is also consistent with Ciabatti et al.'s
addition of a destabilization domain to mCherry's C-terminus. We therefore redid the counts of
labeled cells in our Ai14 datasets to include only cells with fluorescence at seven days of more
than 32.33 a.u., the midpoint of the mean intensities in Ai35 versus Ai14 mice, in order to exclude

neurons that might have been labeled with mCherry alone. As seen in Supplementary Figure S4,
restricting the analysis to the cells that were brightest at seven days (and therefore almost
certainly not labeled with just mCherry instead of either just tdTomato or a combination of both
mCherry and tdTomato) made no major difference: 70.0% of SiR-labeled neurons had
disappeared by 14 days, and 80.8% were gone by 21 days.
Although in theory it is possible that the disappearance of the infected cells could be due
to cessation of tdTomato or Arch-EGFP-ER2 expression rather than to the cells' deaths, because
of downregulation by rabies virus of host cell gene expression (44), we view this as highly unlikely.
Downregulation of host cell gene expression by rabies virus is neither total ("cells with high
expression of RbV transcripts retain sufficient transcriptional information for their classification
into a specific cell type." (44)) nor uniform (45); in practice, we saw no evidence of a decline in
reporter expression in the infected cells but in fact found the exact opposite. As can be seen in a
number of cells in Figures 3 and S4, the cells got brighter and brighter over time, unless they
abruptly disappeared. In our experience, including in this case, cells infected with rabies virus
increase in brightness until they die, often blebbing and coming apart into brightly labeled pieces,
regardless of whether the fluorophore is expressed from a reporter allele (as in this case) or
directly by the virus (see Chatterjee et al. 2018 for many more examples of this (21)).
287
288 **Construction and testing of a virus with an intact PEST domain**
289
We decided to directly test whether a rabies virus with an intact PEST domain fused to its
nucleoprotein can spread transsynaptically, with or without TEVP (Figure 4). Beginning with our
lab's first-generation virus RV∆G-4Cre (21), we constructed a "self-inactivating" version, "RV∆G-
NPEST-4Cre", by adding the coding sequence for the C-terminal addition from Ciabatti et al. (25)
to the 3' end of the nucleoprotein gene. To reduce the chance of the PEST domain being lost to
nonsense mutations during production of the virus, we made synonymous changes to five codons
near the junction of the end of the native nucleoprotein gene and the beginning of the addition,
so that those codons were no longer a single mutation away from being stop codons; apart from
these five synonymous changes, the nucleotide sequence of the addition was identical to that
used by Ciabatti et al. (25) (Figure 4A). We also made a matched "revertant" version, "RV∆G-
N*PEST-4Cre", with exactly the same sequence as RV∆G-NPEST-4Cre except with a stop codon
three codons into the linker, at the same location as the stop codon that we had found in the
overwhelming majority of viral particles in the original SiR-CRE. While the genomes of these two
new viruses therefore only differed from each other by one codon, at the protein level one virus
(RV∆G-NPEST-4Cre) had a full-length PEST domain on the end of its nucleoprotein, while the
other (RV∆G-N*PEST-4Cre) had only a two-amino-acid (Gly-Ser) addition to the end of its
nucleoprotein and otherwise was an ordinary first-generation ∆G virus.
Following production of high-titer EnvA-enveloped virus (see Methods), we confirmed that
the final stocks retained the intended 3' additions to the nucleoprotein gene by extracting the
genomic RNA and used Sanger sequencing on the genomes of 32 viral particles for each virus.
All of the sequenced clones for each of the two viruses had the respective intended sequences
(Figure 4, panel A; details in Supplementary File S7), except for one clone of RV∆G-NPEST-
4Cre(EnvA) with a point mutation in the transcriptional termination signal of the nucleoprotein
gene (from CATGAAAAAAA to CATAAAAAAAA), and two clones of RV∆G-N*PEST-4Cre(EnvA)
that had one synonymous mutation each, one in the nucleoprotein gene and the other in the PEST
domain after the introduced stop codon (and that was therefore irrelevant).
Having verified that the NPEST and N*PEST viruses retained their respective
modifications (at least in all 32 of the clones sequenced for each), we tested their ability to spread
transsynaptically *in vivo* in the absence of TEVP, using corticostriatal neurons as the starting cells
(Figure 4, panel B). We injected an AAV2-retro (11) expressing FLPo into the dorsolateral striatum
of three mice each of the tdTomato reporter line Ai14(46), with a FLP-dependent cocktail of two

helper viruses (AAV1) injected into the primary somatosensory cortex (barrel field) in the same
surgery. The helper virus cocktail was a FLP-dependent version of a combination we have
published elsewhere (47-49), with the sole modifications being the substitution of orthogonal FRT
sites for orthogonal lox ones, and the incorporation of a 2-bp frameshift to prevent the production
of a premature stop codon within the first FRT site, in the first helper virus (Supplementary Figure
S5 shows the results of pilot testing of this FLP-dependent helper virus). Corticostriatal neurons
coinfected with all three AAVs were therefore intended to express, from the first helper virus, TVA
(to allow infection by EnvA-enveloped RV), the tetracycline transactivator (tTA), and EGFP (to
mark expression of TVA and tTA), as well as, from the second helper virus, G and the blue
fluorophore mTagBFP2 (to mark G expression). One week after AAV injection, we injected either
the NPEST or the N*PEST virus at equalized titers, then perfused the mice either 7 days or 3
weeks after the rabies virus injections.
Figure 4 and Supplementary Figure S6 show the results of these injections. The
"revertant" N*PEST virus, which had only an additional two amino acids on the C-terminus of its
nucleoprotein, spread efficiently (bottom rows of images in panels C and D, and rightmost bar in
each pair in the charts in panel E). We counted tdTomato-labeled neurons in ipsilateral cortex
(regardless of cortical area) as well as in thalamus and in contralateral cortex (regardless of
cortical area). Note that, because we counted neurons only in every sixth 50 μm section (see
Methods), the numbers of labeled neurons in the entire brain of each mouse would be
approximately six times the numbers given below and in the figures. At 7 days after rabies virus
injection (Supplementary Figure S6), we found an average of 4751 tdTomato-labeled neurons in
ipsilateral cortex (all areas); we also found an average of 73 labeled neurons in contralateral
cortex and 321 in thalamus. At 3 weeks after rabies virus injection (Figure 4), a time point which
is unusually long for a first-generation virus (21) but which we included to match the duration used
by Ciabatti et al. (25), the N*PEST virus had spread to many more neurons: on average 20759 in
ipsilateral cortex, 472 in contralateral cortex, and 244 in thalamus. In order to ensure that all this
label was not due simply to leaky TVA expression or residual RVG-enveloped virus, we performed
control experiments in which we omitted the AAV expressing G and replaced it by an AAV
expressing TEVP. This virus, AAV1-TREtight-H2b-emiRFP670-TEVP, was of the same design as
the G-expressing AAV (AAV1-TREtight-mTagBFP2-B19G) but encoded TEVP (S219V mutant)
instead of G and an H2b-fused near-infrared fluorescent protein (emiRFP670 (50)) instead of
mTagBFP2. These control injections resulted in very little labeling in these input regions
(Supplementary Figure S8), suggesting that the large amount of label seen when G was included
indicated *bona fide* and efficient transsynaptic spread of the N*PEST virus, consistent with our
lab's prior experience with the parent virus RVΔG-4Cre(EnvA) as well as with other first-
generation RVΔG vectors.
In contrast, matched experiments using the NPEST virus, which had the intact PEST
domain fused to its nucleoprotein, showed much less transsynaptic spread without TEVP at either
time point examined (top rows of images in panels C and D, and leftmost bar in each pair in the
charts in panel E). At 7 days (Supplementary Figure S6), we found averages of 1024 labeled cells
in ipsilateral cortex (all areas), 2 in contralateral cortex, and 4 in thalamus (although these
numbers were much lower than the comparable numbers for the N*PEST virus, the differences
at this timepoint were not statistically significant due to high variance and low n (3 mice for each
condition). See Supplementary File S8 for all cell counts and results of statistical analyses, and
see Supplementary File S9 for a summary table of mean cell counts for all conditions. Single-
factor ANOVAs were used for all comparisons). At 3 weeks, we found averages of 1366 labeled
cells at in ipsilateral cortex, 6 in contralateral cortex, and 11 in thalamus; these numbers were
significantly lower than those obtained with the "revertant" N*PEST virus. Here again, this label
in input regions did not appear to be due simply to direct retrograde infection by residual RVG-
enveloped virus, because it was almost completely absent in control animals in which the G-

expressing AAV was omitted and replaced with one expressing TEVP (Supplementary Figure S8).

Finally, we tested the ability of the two viruses to spread when both TEVP and G were supplied (Figure 5 and Supplementary Figure S7). These experiments were done in exactly the same way as the experiments described above but with both the G-expressing and TEVP-expressing AAVs included in the helper virus mixture.

Whereas the inclusion of TEVP did not much change the results using the "revertant" N*PEST virus, it greatly increased the spread of the NPEST virus, with means at 7 days of 8802 labeled cells in ipsilateral cortex, 82 in contralateral cortex, and 191 in thalamus; at 3 weeks, there were on average 9652 in ipsilateral cortex, 197 in contralateral cortex, and 283 in thalamus. These numbers were all not statistically significantly different from the corresponding ones for the N*PEST virus. These results suggest that the inclusion of TEVP restored the ability of the NPEST virus to replicate and spread, as we assume that the original authors had initially intended.

Importantly, we also found indications that both viruses eventually killed most of the "source cells" in which they replicated (Supplementary Figure S9). For example, using the NPEST virus with TEVP provided, the average number of remaining "source cells" (defined in this case as cells that were labeled by the virus and expressed both G and TEVP) dropped from 29 at 7 days to 2 at 3 weeks (although here again the comparison was not quite statistically significant (p = 0.071820149) due to the high variance and low number (3) of mice per group).

DISCUSSION

Our transsynaptic tracing results using virus with an intact PEST domain contradict the earlier claim that such a virus can efficiently spread between neurons in the absence of TEVP(25). Our findings from sequencing the SiR viruses from the originating laboratory suggest that the reason for that claim was that the viruses used in the original study had lost the intended modification.

While it is possible that the escape of the two SiR virus samples from the modification intended to attenuate them was a fluke due to bad luck with those two batches, we view this as unlikely, for three reasons. First, the reported finding that viral replication and spread occurred in the absence of TEVP is difficult to understand in the absence of mutations but is easily explained if the viral preparations used for those experiments harbored the kind of mutations that we found in the two preparations to which we had access. Second, the two SiR virus samples that we analyzed had independently developed mutations causing loss of the intended C-terminal addition to the nucleoprotein; we know that the mutations were independent because the two samples were of different viruses so did not both derive from a single compromised parental stock. Third, the mutation profiles of the two viruses were very different: whereas the SiR-CRE sample had the same point mutation in nearly 100% of its viral particles, only a minority of the SiR-FLPo particles had that particular mutation, with the majority having a different point mutation three codons away that had the same result. This suggests that any of the many opportunities for removing the C-terminal addition — creation of a premature stop codon at any one of a number of sites, or a frameshift mutation anywhere in the vicinity — can be exploited by a given batch of virus, greatly increasing the probability of such mutants arising.

While it is clearly possible to make virus with the intended modification to the nucleoprotein in at least the majority of virions, because we have done so here (and the authors of the original paper also report doing so, in a recent preprint (51) in response to our own preprint of an earlier version of this paper), our findings suggest that the approach in its current form is vulnerable to being undermined by viral mutation. Consistent with this, the original authors' recent preprint reports loss of the 3' addition in a significant minority of virions within as few as four passages(51). Although a number of groups have made recombinant rabies viruses — as well as other rhabdoviruses and other nonsegmented negative-strand RNA viruses — encoding fusions of

exogenous proteins to viral proteins (52-62), most of these groups have found that the additions
significantly impaired function, and some have found that the viruses rapidly lost C-terminal
additions to viral proteins. For example, an attempt to make SAD B19 rabies virus with EGFP
fused to the C-terminus of the nucleoprotein was unsuccessful, suggesting that large C-terminal
additions make the nucleoprotein dysfunctional; the authors of that paper resorted instead to
making virus encoding the fusion protein in addition to the wild-type nucleoprotein (55). A
vesicular stomatitis virus with GFP fused to the C-terminus of the glycoprotein gene lost the
modification within a single passage of the virus because of a point mutation creating a premature
stop codon (52). Relatedly, a VSV with its glycoprotein gene replaced with that of a different virus
was found to quickly develop a premature stop codon causing loss of the last 21 amino acids of
the exogenous glycoprotein, conferring a marked replication advantage to the mutants bearing
the truncated version (63). Generalizing from these prior examples as well as our findings here,
we suggest that any attempt to attenuate a virus by addition to the C-terminus of a viral protein
will be vulnerable to loss of the modification, and that any such virus will therefore need to be
monitored very carefully.
It is unclear whether further improvements to the design of the viruses can be made to
make loss of the PEST domain less likely. Although the synonymous mutations that we made to
five codons near the junction of N and the 3' addition are a good start, there are numerous other
codons in the immediate vicinity that are also one point mutation away from being stop codons
but that do not have synonyms that are not. Furthermore, no such changes would protect against
frameshifts.
If the viruses used for the transsynaptic tracing experiments in the original paper were
actually *de facto* first-generation, ΔG viruses like the SiR samples that we analyzed, how could
the authors have found, in postmortem tissue, cells labeled by SiR-CRE that had survived for
weeks? The answer may simply be that, as we have shown in Chatterjee et al. (21) and again
here (Figure 3), and as the original authors also report in their new preprint (51), a preparation of
first-generation rabies viral vector expressing Cre can leave a large fraction of labeled cells alive
for at least months, in contrast to similar ones encoding tdTomato (21) or EGFP (4). Similarly,
Gomme et al. found long-term survival of some neurons following infection by a replication-
competent rabies virus expressing Cre (64)). Our results with the "revertant" control virus RVΔG-
N*PEST-4Cre (Figures 4 and 5, and Supplementary Figures S5-S7), showing thousands of
labeled neurons three weeks after rabies virus injection, are consistent with this as well.
One reason, in turn, why a preparation of a simple ΔG rabies virus encoding Cre can leave
many cells alive may be that not all the virions are in fact first-generation viral particles, because
of the high mutation rate that we have highlighted in this paper. We have shown in Chatterjee et
al. (21) that a second-generation (ΔGL) rabies virus, which has both its glycoprotein gene G and
its polymerase gene L deleted, leaves cells alive for the entire four months that we followed them.
However, any first-generation (ΔG) virus that contains a frameshift or point mutation knocking out
L will in practice be a ΔGL virus. In fact, a stop codon or frameshift mutation in any of several
other viral genes is likely to have a similar effect as one in L (and it might be that the Ser419Thr
mutation that we found in 9.49% of our RVΔG-4Cre virions is just such a knockout mutation of N).
Together with the high mutation rate of rabies virus, this means that, within every preparation of
first-generation rabies virus there is almost guaranteed to be a population of de facto second-
generation variants mixed in with the intended first-generation population and propagated in the
producer cells by complementation by the first-generation virions. Any rabies virus preparation
(whether made in the laboratory or occurring naturally) can be expected to contain a population
of such knockout (whether by substitution, frameshift, or deletion) mutants (related to the classic
phenomenon of "defective interfering particles", or mutants with a marked replication advantage
(65-67), and the higher the multiplicity of infection when passaging the virus, the higher the
proportion of such freeloading viral particles typically will be. This would not necessarily be noticed
in the case of a virus encoding a more common transgene product such as a fluorophore, because

the expression levels of these by the knockout mutants would be too low to label cells clearly (see
Figure 1 in Chatterjee et al. (21)). However, with Cre as the payload, any "second-generation"
particles would be able to label neurons but not kill them, because second-generation rabies viral
vectors do not kill cells for at least months (21). This explanation would predict that the percentage
of neurons surviving infection with a rabies virus encoding Cre will depend on the particular viral
preparation that is injected, with some having a greater fraction of knockout particles than others.
This could explain why the SiR-CRE virus sample killed cells faster than our own RVΔG-
4Cre (Figure 3). This analysis would also presumably apply to first-generation (ΔG) viruses
expressing FLPo: while we found that the FLPo-expressing version that we made did not leave
as many cells alive as the Cre-expressing version did (Supplementary Figure S10), that
preparation may simply have had fewer mutants with knockout of genes essential for replication.
Most interestingly, we found that virus with an intact PEST domain does spread efficiently
between neurons when TEVP is supplied. This suggests that such viruses could become the
basis for monosynaptic tracing systems with reduced toxicity after all. Our finding is
complementary to those of the original authors in their recent preprint (51) that an intact SiR virus
did not kill labeled neurons for five months: that study was of neurons only infected by the rabies
virus (i.e., without expression of G to complement the ΔG virus or expression of TEVP to remove
the C-terminal addition to the nucleoprotein); they did not show that their intact virus could spread
between neurons or that it is nontoxic as it does so. Conversely, we showed here that an intact
NPEST virus can spread between neurons, but we did not specifically investigate toxicity during
this process. It remains to be seen to what degree an NPEST or SiR virus that spreads
transsynaptically is cytotoxic, not only to transsynaptically-labeled cells but also to the starting
cells, which need to express both G and TEVP to allow replication and spread of the virus. This
is not a trivial point, as G is toxic when overexpressed (68), and the original authors' finding that
cultured cells rapidly lost TEVP activity (51) suggests that TEVP expressed at sufficient levels
may be toxic as well. Indeed, the much lower average numbers of source cells that we found at
3 weeks than at 7 days (Supplementary Figure S9) suggest that the NPEST virus in the presence
of TEVP may be just as cytotoxic to source cells as ordinary ΔG virus is.
However, there may be room for improvement: the TEVP expression levels that we
provided here are unlikely to happen to be optimal. Given that we have used the tTA/TRE system
to provide both TEVP and G in the source cells, the expression of both could be controlled with
doxycycline to result in a system that is less toxic to source cells (mirroring an approach we have
used with ΔGL viruses (24). Another consideration (perhaps minor) is the nine additional amino
acids that are unavoidably left on the C-terminus of the nucleoprotein after the rest of the addition
is removed by TEVP: while these additional amino acids may not much impair the function of the
protein, they are unlikely to help it.
It is also unclear to what degree a minority population of revertant mutants that do arise in
stocks of otherwise-intact virus would pose a problem for monosynaptic tracing studies. If, for
example, ~5% of the virions in a given preparation were revertant mutants (e.g., as the original
authors report obtaining after six passages in cells highly expressing TEVP (51)), one might
expect the same percentage of labeled presynaptic neurons to be labeled by those mutants and
therefore to experience the toxicity of infection by a first-generation, ΔG virus. However, because
the process of infecting the source cells, replicating within them (if provided with G), and spreading
to other cells is comparable to an additional passage in cell culture, the percentage of presynaptic
cells labeled by the revertant mutants could be higher, and would presumably depend on the level
of TEVP expression in each source cell.
Such a minority population of revertant mutants in our "intact" NPEST virus preparation
may explain the limited spread of this virus that we observed in the absence of TEVP. Although
all 32 of the clones that we checked using Sanger sequencing had the intended 3' modification to
the nucleoprotein gene (Supplementary Figure S7), it is entirely possible (perhaps even
inevitable) that revertant virions were present in lower numbers, possibly at a level sufficient to

support some replication of the mixed population of virions in cells expressing G without TEVP.
While we view this as the most likely explanation, showing this convincingly would be difficult,
even assuming that high-throughput sequencing (such as the SMRT sequencing we used to
analyze the originating lab's stocks) would reveal the presence of revertant mutants at some level.
Even *in situ* sequencing sequencing(69-71) of viral genomes in pre- and postsynaptic neurons in
mice injected with the "intact" NPEST virus might not be conclusive: because functional
nucleoprotein produced in postsynaptic cells by revertant virus would complement and package
"intact" NPEST-encoding virions, allowing them to spread, it would not be particularly surprising
to find that the majority of virions in presynaptic neurons were "intact", even if a minority population
of revertant mutants were necessary to allow any spread at all in the absence of TEVP. This is
speculation, however.
What seems clear from the results we have presented here is that reliance on addition to
a viral protein of a C-terminal domain that could be lost by mutation at any replication event is a
tenuous basis for rendering a virus nontoxic. At the very least, we would suggest that any such
attenuating addition be placed on the N terminus of the viral protein (if this turns out to be possible
without impairing the function or trafficking of the protein) so that it could not easily be mutated
away.
In summary, our results suggest that rabies virus with a PEST domain added to the C
terminus of its nucleoprotein is vulnerable to the loss of that modification and only spreads
efficiently between neurons if it is removed, whether by unintended mutation or by expression of
TEVP. Our finding that virus with a PEST domain that is intact in the majority of virions can spread
efficiently when TEVP is provided, as the designers had presumably originally intended, raises
the possibility that further optimization and validation could make the SiR approach a viable option
for monosynaptic tracing with reduced toxicity.

548
549
550 **SUMMARY OF METHODS (see Supplementary Methods for details)**
551
Cloning
The following novel plasmids were made using standard cloning techniques (see Supplementary
Methods): pLV-CAG-FLEX-BFP-(mCherry)' (Addgene 115234), pLV-CAG-F14F15S-BFP-
(mCherry)' (Addgene 115235), pLV-U-TVA950 (Addgene 115236), pRVΔG-4FLPo (Addgene
122050), pAAV-syn-F14F15S-splitTVA-EGFP-tTA_v2 (Addgene 183352), pB-CAG-TEVP-IRES-
mCherry (Addgene 174377), pAAV-syn-FLPo (Addgene 174378), pAAV-TREtight-H2b-
emiRFP670-TEVP (Addgene 174379), pRVΔG-NPEST-4Cre (Addgene 174380), pRVΔG-
N*PEST-4Cre (Addgene 174381), pCAG-hypBase. All of the above novel plasmids have been
deposited with Addgene and can be obtained from there, except for pCAG-hypBase, the
distribution of which is not permitted due to intellectual property constraints.
562
563 **Production of lentiviral and adeno-associated viral vectors**
564 Lentiviral vectors were made as described (72) but using a vesicular stomatitis virus envelope
565 expression plasmid pMD2.G for most vectors except for LV-U-TVA950(B19G), which was made
566 using the rabies virus envelope expression plasmid pCAG-B19GVSVGCD (72).
567
568 AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2, AAV1-TREtight-H2b-emiRFP670-P2A-TEVP, and
569 AAV2-retro-syn-FLPo were made in-house by standard techniques (see Supplementary
570 Methods).
571
572 AAV1-TREtight-mTagBFP2-B19G (which we have described previously(47, 48)), was packaged
573 as serotype 1 by Addgene (catalog # 100798-AAV1).
574

575
Production of titering cell lines
576
577 Reporter cell lines 293T-FLEX-BC and 293T-F14F15S-BC were made using lentiviral vectors
578 made from pLV-CAG-FLEX-BFP-(mCherry)' and pLV-CAG-F14F15S-BFP-(mCherry)', described
579 above. TVA-expressing versions, 293T-FLEX-BC-TVA and 293T-F14F15S-BC-TVA, were made
580 by infecting the above lines with LV-U-TVA950(VSVG) (described above).
581
582 **Production of TEVP-expressing cell line**
583 293T-TEVP was made by transfecting HEK 293T/17 cells with pCAG-hypBase and pB-CAG-
584 TEVP-IRES-mCherry, then sorting.
585
586 **Production and titering of rabies viruses**
587 RV∆G-4Cre, RV∆GL-4Cre, RV∆G-NPEST-4Cre, and RV∆G-N*PEST-4Cre were produced
588 mostly as described (21, 35) (see Supplementary Methods); titering was as described (33) but
589 using the 293T-FLEX-BC and 293T-F14F15S-BC lines used for B19G-enveloped viruses and the
590 293T-FLEX-BC-TVA and 293T-F14F15S-BC-TVA used for the EnvA-enveloped viruses.
591
Extraction of viral genomic RNA and preparation for Sanger sequencing
RNA viral genomes were extracted from virus samples using a Nucleospin RNA kit (Macherey-
Nagel, Germany), then converted to cDNA by RT-PCR (Agilent Technologies, USA) with a
barcoded primer. cDNA sequences were amplified using Platinum SuperFi Green Master Mix
(Invitrogen (Thermo Fisher), USA) and cloned into pEX-A (Eurofins Genomics, USA) using an In-
Fusion HD Cloning Kit (Takara Bio, Japan). Sequencing data was collected for over fifty clones
per sample.
599
Single-molecule, real-time (SMRT) sequencing
Double-stranded DNA samples for SMRT sequencing were prepared similarly to the above,
except that that the clones generated from each of the three virus samples were tagged with one
of the standard PacBio barcode sequences to allow identification of each clone's sample of origin
following multiplex sequencing. This was in addition to the random index (10 nucleotides in this
case) that was again included in the RT primers in order to uniquely tag each individual genome.
606
607 **Surgeries and virus injections for two-photon imaging**
608 All experimental procedures using mice were conducted according to NIH guidelines and were
609 approved by the MIT Committee for Animal Care (CAC). Mice were housed 1-4 per cage under a
610 normal light/dark cycle for all experiments.
611
612 Adult mice of Cre-dependent reporter strains Ai14 (73) (Jackson Laboratory #007908) or Ai35D
613 (41) (Jackson Laboratory # 012735) mice were injected in V1 with LV-U-TVA950(B19G), then
614 implanted with a glass window. Seven days later, windows were removed and one of the three
615 EnvA-enveloped rabies viral vectors (with equalized titers) was injected at the same coordinates,
616 then coverslips were reapplied.
617
618 *In vivo* **two-photon imaging and image analysis**
619 Beginning seven days after injection of each rabies virus and continuing every seven days up to
620 a maximum of four weeks following rabies virus injection, the injection sites were imaged on a
621 two-photon microscope. One field of view was chosen in each mouse in the area of maximal
622 fluorescent labelling. Cell counting was performed with the ImageJ Cell Counter plugin.
623
624 **Monosynaptic tracing experiments: surgeries and virus injections**

The three helper AAV1s were combined at final titers of 3.6E10 gc/ml for AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2 and 6.60E11 gc/ml for AAV1-TREtight-mTagBFP2-B19G and/or AAV1-TREtight-H2b-emiRFP670-P2A-TEVP. 250 nl of helper virus mixture was injected into layer 5 of barrel cortex of Ai14 mice; in the same surgery, 300 nl of AAV2-retro-syn-FLPo (1.16E13 gc/ml) was injected into dorsolateral striatum. 7 days after AAV injection, 300nl of RVΔG-NPEST-4Cre(EnvA) (1.86E9 iu/ml) or RVΔG-N*PEST-4Cre(EnvA) (diluted to 1.86E9 iu/ml) was injected in barrel cortex at the same site as the helper AAV mixtures.

Monosynaptic tracing experiments: perfusions and histology
12 days or 3 weeks after injection of rabies virus, mice were perfused; Brains were postfixed overnight and cut into 50 μm coronal sections on a vibrating microtome. Sections were immunostained as described(48) with a chicken anti-GFP (Aves Labs GFP-1020) 1:500 and donkey anti-chicken Alexa Fluor 488 (Jackson Immuno 703-545-155) 1:200.

Monosynaptic tracing experiments: cell counts and microscopy
tdTomato-labeled neurons in contralateral cortex and thalamus were counted manually with the Cell Counter plugin in ImageJ. Cells at the injection site were counted either manually or (when dense) using the Analyze Particle function in ImageJ. Only one of the six series of sections (i.e., every sixth section: see above) was counted for each mouse. Images for figures were taken on a confocal microscope (Zeiss, LSM 900).

ACKNOWLEDGEMENTS

We thank Ernesto Ciabatti and Marco Tripodi for sharing samples of EnvA/SiR-CRE and EnvA/SiR-FLPo and for comments on the manuscript. We thank Ed Callaway, Sean Whelan, Ayano Matsushima, and Kim Ritola for helpful discussion and Jun Zhuang, Soumya Chatterjee, and Ali Cetin for helpful discussion and sharing their own results with SiR viruses. We thank Stuart Levine, Noelani Kamelamela, and Huiming Ding of the MIT BioMicro Center for assistance with SMRT sequencing and bioinformatic data analysis and Sara Beach for helpful feedback on the manuscript. We thank Xin Fu for noticing the stop codon that I.R.W. had inadvertently introduced into the FLP-dependent helper virus plasmid pAAV-F14F15S-splitTVA-EGFP-tTA. Research reported in this publication was supported by the following BRAIN Initiative awards from the National Institute of Mental Health: U01MH106018 (Wickersham) RF1MH120017 (Wickersham), U01MH114829 (Dong), and U19MH114830 (Zeng).

 FIGURE LEGENDS

Figure 1: Single-molecule Sanger sequencing of individual viral genomes revealed that
most SiR virions had lost the intended C-terminal modification to the nucleoprotein.
(A) Schematic of the RT-PCR workflow. In the reverse transcription (RT) step, the RT primer,
containing a random 8-nucleotide sequence, anneals to the 3' rabies virus leader, adding a unique
random index to the 5' end of the cDNA molecule corresponding to each individual viral particle's
RNA genome. In the PCR step, the forward PCR primer anneals to the RT primer sequence and
the reverse PCR primer anneals within the viral phosphoprotein gene P. Both PCR primers have
15-base sequences homologous to those flanking an insertion site in a plasmid used for
sequencing, allowing the amplicons to be cloned into the plasmid using a seamless cloning
method before transformation into bacteria. The resulting plasmid library consists of plasmids
containing up to 4^8 different index sequences, allowing confirmation that the sequences of
plasmids purified from individual picked colonies correspond to the sequences of different
individual rabies viral particles' genomes.
(B) Representative Sanger sequencing data of the 8-bp index and the TEV-PEST sequence.
Mutations are highlighted in red.
(C) Mutation variants and their frequencies in each viral vector sample based on Sanger
sequencing data. No unmutated genomes were found in the SiR-CRE sample: 50 out of 51 had
a substitution creating an opal stop codon just before the TEV cleavage site, and the 51st genome
contained a frameshift which also removed the C-terminal addition. In the SiR-FLPo sample, only
4 out of 50 clones had an intact sequence of the C-terminal addition; the other 46 out of 50 had
one of two *de novo* stop codons at the end of N or the beginning of the TEV cleavage site. In the
sample of RVΔG-4mCherry, a virus from our laboratory included as a control to distinguish true
mutations on the rabies genomes from mutations due to the RT-PCR process, none of the 51
clones analyzed had mutations in the sequenced region.

A

B

C

Figure 2: Single-molecule, real-time (SMRT) sequencing of thousands of barcoded viral genomes confirmed that most SiR virions had lost the intended C-terminal modification to the nucleoprotein.

(A) Schematic of workflow for SMRT sequencing. An RT primer with a random 10-nucleotide sequence anneals to the leader sequence on the negative-sense single-stranded RNA genome. Forward and reverse PCR primers have distinct SMRT barcodes at their 5' ends for the three different virus samples. After RT-PCR, each amplicon library consists of amplicons each containing a SMRT barcode to identify the sample of origin as well as the 10-nucleotide index (i.e., with a potential diversity of 4^{10} different indices) to uniquely label the individual genome of origin. SMRT "dumbbell" adaptors are then ligated to the amplicons' ends, making circular templates which are then repeatedly traversed by a DNA polymerase, resulting in long polymerase reads each containing multiple reads of the positive and negative strands. The individual subreads for a given molecule are combined to form circular consensus sequence (CCS) reads.

(B) High-frequency (>2%) point mutations found in the rabies vector samples based on SMRT sequencing. Horizontal axis represents nucleotide position along the reference sequences (see text); vertical axis represents variant frequency. Total number of CCS3 reads (i.e., with at least 3 subreads for each position) are 22,205 for SiR-CRE, 17,086 reads for SiR-FLPo, and 17,978 reads for RVΔG-4Cre. The great majority of SiR-CRE and SiR-FLPo genomes have point mutations creating premature stop codons at or just after the end of N and before the C-terminal addition. The only frequent (>2%) mutation found in the control virus, RVΔG-4Cre, was a single amino acid substitution at position 419 in 9.49% of virions. Insertions and deletions are not shown here (see text).

(C) Summary of results. In the SiR virus samples, 99.22% of SiR-CRE virions and 83.85% of SiR-FLPo virions had point mutations creating premature stop codons that completely removed the intended C-terminal addition to the nucleoprotein, making them simply first-generation (ΔG) rabies viral vectors. This does not include any insertions or deletions causing frameshifts (see text), which would further increase the percentage of first-generation-type virions in these samples. In the RVΔG-4Cre sample, there were no premature stop codons at or near the end of the nucleoprotein gene.

Figure 3: Longitudinal two-photon imaging *in vivo* showed that SiR virus killed approximately 80% of infected neurons in vivo within 2-4 weeks.

A) Representative fields of view (FOVs) of visual cortical neurons labeled with RVΔG-4Cre (top row), RVΔGL-4Cre (middle row), or SiR-CRE (bottom row) in Ai14 mice (Cre-dependent expression of tdTomato). Images within each row are of the same FOV imaged at the four different time points in the same mouse. Circles indicate cells that are present at 7 days postinjection but no longer visible at a subsequent time point. Scale bar: 50 µm, applies to all images.

B-D) Numbers of cells present at week 1 that were still present in subsequent weeks. While very few cells labeled with RVΔGL-4Cre were lost, and RVΔG-4Cre killed a significant minority of cells, SiR-CRE killed the majority of labeled neurons within 14 days following injection. Each FOV is from a single mouse: n = 2 mice in B, 3 mice in C, and 4 mice in D.

E) Percentages of cells present at week 1 that were still present in subsequent imaging sessions. By 28 days postinjection, an average of only 20.5% of cells labeled by SiR-CRE remained.

A

Week 1 | Week 2 | Week 3 | Week 4

RVΔG-4Cre

RVΔGL-4Cre

SiR-CRE

B RVΔG-4Cre

Remaining week1 cells

C RVΔGL-4Cre

Cell count

D SiR-CRE

Week 1 2 3 4 | 1 2 3 4 | 1 2 3 4 | 1 2 3 4
FOV-1 | FOV-2 | FOV-3 | FOV-4

E

RVΔG-4Cre
RVΔGL-4Cre
SiR-CRE

Cell count (percentage of week1 cells)

Time (week)

Figure 4: In the absence of TEVP, rabies virus with an apparently intact PEST domain spreads much less than one without a PEST domain.

A) Sequences of the region of the junction between the nucleoprotein gene N and its 3' addition in SiR-CRE (intended sequence) as well as in our two new viruses RVΔG-NPEST-4Cre and RVΔG-N*PEST-4Cre. Apart from synonymous mutations we made in five codons (highlighted in green) to make them less likely to mutate to stop codons, RVΔG-NPEST-4Cre has exactly the same 3' addition to N as was intended for SiR. RVΔG-N*PEST-4Cre has exactly the same sequence as RVΔG-NPEST-4Cre apart from a stop codon that we deliberately introduced at the same location as the nonsense mutation that we had found in almost every SiR-CRE virion, so that the only modification to the native nucleoprotein is an additional two amino acids (Gly-Ser) on its C-terminus. Sanger sequencing of 32 viral particles of each virus confirmed that the intended additions were still present in the final stocks.

B) Diagram of virus injections for monosynaptic tracing experiments. An AAV2-retro expressing FLPo was injected in dorsolateral striatum, and a FLP-dependent helper virus combination was injected into barrel cortex to express G in corticostriatal cells. Rabies virus was injected seven days later at the same location in barrel cortex, and mice were perfused either 7 days or 21 days after rabies virus injection.

C-D) Results at 21 days after injection (see Supplementary Fig. S6 for similar results at 7 days). At 21 days after injection, the NPEST virus (top row), with an intact PEST domain, shows very limited spread, whereas the N*PEST virus (bottom row), without the PEST domain, has labeled many thousands of neurons in ipsilateral cortex (C) and also spread to many cells in contralateral cortex and thalamus (D) (see quantification in E). For all panels in this figure as well as Figure 5 and the related supplementary figures, the red channel shows tdTomato, reporting Cre expression from the rabies viruses; the green channel shows EGFP, coexpressed with tTA by the first helper virus; the blue channel shows mTagBFP2, coexpressed with G by the second helper virus. Scale bar: 200 μm, applies to all images.

E) Counts of labeled neurons in ipsilateral cortex (left), contralateral cortex (center), and thalamus (right) at 21 days for the two viruses. Each dot indicates the total number of labeled cells found in a given mouse brain when examining every sixth 50 μm section, so that the corresponding numbers in the entire brain would be roughly six times as high (see Methods). All differences in numbers of cells labeled by the two viruses are significant for all conditions (see Supplementary File S8 for all counts and statistical analyses).

A

	N	TEV protease cleavage site	PEST

SiR-CRE
K T Y S S D S G S G E N L Y F Q S G S G S H G F P
AAG ACA TAT TCT AGT GAC TCT GGT TCC GGT GAG AAC CTC TAC TTC CAA TCT GGT TCC GGT AGC CAT GGC TTC CCG

RVΔG-NPEST-4Cre (32/32) ("intact")
K T Y S S D S G S G E N L Y F Q S G S G S H G F P
AAG ACA TAT AGT AGT GAC AGT GGT TCC GGC GAG AAC CTC TAC TTC CAA AGT GGC TCC GGT AGC CAT GGC TTC CCG

RVΔG-N*PEST-4Cre (32/32) ("revertant")
K T Y S S D S G S *
AAG ACA TAT AGT AGT GAC AGT GGT TCC TGA GAG AAC CTC TAC TTC CAA AGT GGC TCC GGT AGC CAT GGC TTC CCG

B
Day -7:
AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2
AAV1-TREtight-mTagBFP2-B19G
in S1

Day -7:
AAV2-retro-synP-FLPo
in striatum

Day 0:
RVΔG-NPEST-4Cre(EnvA)
("intact")
or RVΔG-N*PEST-4Cre(EnvA)
("revertant") in S1

Day 7 or 21:
Perfuse

Ai14 mice

C

Without TEVP: 3 weeks

NPEST ("intact") / N*PEST ("revertant")

tdTomato (reports Cre) — EGFP (reports TVA & tTA) — mTagBFP2 (reports G) — Merged

200 μm

D

Without TEVP: 3 weeks

NPEST ("intact") / N*PEST ("revertant")

M1/M2 cortex — S2 cortex — Thalamus — Contralateral cortex

200 μm

E

Ipsilateral cortex — Contralateral cortex — Thalamus

tdTomato+ cells

NPEST ("intact") / N*PEST ("revertant")

Figure 5: In the presence of TEVP, rabies virus with an apparently intact PEST domain spreads to a similar degree as one without a PEST domain.

With the experimental design exactly the same as shown in Figure 4, Panel B but with a TEVP-expressing AAV included in the helper virus mixture (A), the NPEST virus did spread transsynaptically, to a similar degree as the N*PEST one.

B-D) Results at 21 days after injection (see Supplementary Fig. S7 for similar results at 7 days). 21 days after injection, both NPEST (top row) and N*PEST (bottom row) viruses have labeled many cells in ipsilateral cortex (B). Both viruses have also spread to contralateral cortex and thalamus (C).

D) Counts of tdTomato+ cells in ipsilateral cortex, contralateral cortex, and thalamus. Each dot indicates the total number of labeled cells found in a given mouse brain when examining every sixth 50 μm section, so that the corresponding numbers in the entire brain would be roughly six times as high (see Methods). Differences between numbers of cells labeled by the NPEST and N*PEST viruses are not significant (see Supplementary File S8 for all counts and statistical analyses). Note that the inclusion of TEVP reduced the spread of the N*PEST virus (cf. Figure 4; all counts and statistical comparisons are in Supplementary File S8), possibly because of competition for receptors or expression machinery in source cells.

A

Day -7:
AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2
AAV1-TREtight-mTagBFP2-B19G
AAV1-TREtight-TEVP
in S1

Day -7:
AAV2-retro-synP-FLPo
in striatum

Day 0:
RVΔG-NPEST-4Cre(EnvA)
("intact")
or RVΔG-N*PEST-4Cre(EnvA)
("revertant")
in S1

Day 7 or 21:
Perfuse

Ai14 mice

B

C

D

REFERENCES

1. G. Ugolini, H. G. Kuypers, P. L. Strick, Transneuronal transfer of herpes virus from peripheral nerves to cortex and brainstem. *Science* **243**, 89-91 (1989).
2. G. Ugolini, Specificity of rabies virus as a transneuronal tracer of motor networks: transfer from hypoglossal motoneurons to connected second-order and higher order central nervous system cell groups. *J Comp Neurol* **356**, 457-480 (1995).
3. R. M. Kelly, P. L. Strick, Rabies as a transneuronal tracer of circuits in the central nervous system. *J Neurosci Methods* **103**, 63-71 (2000).
4. I. R. Wickersham, S. Finke, K. K. Conzelmann, E. M. Callaway, Retrograde neuronal tracing with a deletion-mutant rabies virus. *Nature Methods* **4**, 47-49 (2007).
5. I. R. Wickersham *et al.*, Monosynaptic restriction of transsynaptic tracing from single, genetically targeted neurons. *Neuron* **53**, 639-647 (2007).
6. D. Atasoy, Y. Aponte, H. H. Su, S. M. Sternson, A FLEX switch targets Channelrhodopsin-2 to multiple cell types for imaging and long-range circuit mapping. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **28**, 7025-7030 (2008).
7. K. T. Beier *et al.*, Anterograde or retrograde transsynaptic labeling of CNS neurons with vesicular stomatitis virus vectors. *P Natl Acad Sci USA* 10.1073/pnas.1110854108 (2011).
8. L. Lo, D. J. Anderson, A Cre-dependent, anterograde transsynaptic viral tracer for mapping output pathways of genetically marked neurons. *Neuron* **72**, 938-950 (2011).
9. L. A. Schwarz *et al.*, Viral-genetic tracing of the input-output organization of a central noradrenaline circuit. *Nature* 10.1038/nature14600 (2015).
10. B. E. Deverman *et al.*, Cre-dependent selection yields AAV variants for widespread gene transfer to the adult brain. *Nat Biotechnol* **34**, 204-209 (2016).
11. D. G. Tervo *et al.*, A Designer AAV Variant Permits Efficient Retrograde Access to Projection Neurons. *Neuron* **92**, 372-382 (2016).
12. J. Dimidschstein *et al.*, A viral strategy for targeting and manipulating interneurons across vertebrate species. *Nat Neurosci* **19**, 1743-1749 (2016).
13. W. B. Zeng *et al.*, Anterograde monosynaptic transneuronal tracers derived from herpes simplex virus 1 strain H129. *Mol Neurodegener* **12**, 38 (2017).
14. L. Luo, E. M. Callaway, K. Svoboda, Genetic Dissection of Neural Circuits: A Decade of Progress. *Neuron* **98**, 865 (2018).
15. S. Ravindra Kumar *et al.*, Multiplexed Cre-dependent selection yields systemic AAVs for targeting distinct brain cell types. *Nat Methods* **17**, 541-550 (2020).
16. J. K. Mich *et al.*, Functional enhancer elements drive subclass-selective expression from mouse to primate neocortex. *Cell Rep* **34**, 108754 (2021).
17. V. Augustine *et al.*, Hierarchical neural architecture underlying thirst regulation. *Nature* **555**, 204-209 (2018).
18. J. Kohl *et al.*, Functional circuit architecture underlying parental behaviour. *Nature* **556**, 326-331 (2018).
19. D. A. Evans *et al.*, A synaptic threshold mechanism for computing escape decisions. *Nature* **558**, 590-594 (2018).
20. M. M. Kaelberer *et al.*, A gut-brain neural circuit for nutrient sensory transduction. *Science* **361** (2018).

822   21.   S. Chatterjee *et al.*, Nontoxic, double-deletion-mutant rabies viral vectors for retrograde
823   targeting of projection neurons. *Nat Neurosci* **21**, 638-646 (2018).
824   22.   T. R. Reardon *et al.*, Rabies Virus CVS-N2c(DeltaG) Strain Enhances Retrograde Synaptic
825   Transfer and Neuronal Viability. *Neuron* **89**, 711-724 (2016).
826   23.   K. Morimoto, D. C. Hooper, S. Spitsin, H. Koprowski, B. Dietzschold, Pathogenicity of
827   different rabies virus variants inversely correlates with apoptosis and rabies virus
828   glycoprotein expression in infected primary neuron cultures. *J Virol* **73**, 510-518 (1999).
829   24.   L. Jin *et al.*, Long-term labeling and imaging of synaptically-connected neuronal
830   networks in vivo using nontoxic, double-deletion-mutant rabies viruses.
831   *bioRxiv* 10.1101/2021.12.04.471186, 2021.2012.2004.471186 (2021).
832   25.   E. Ciabatti, A. Gonzalez-Rueda, L. Mariotti, F. Morgese, M. Tripodi, Life-Long Genetic and
833   Functional Access to Neural Circuits Using Self-Inactivating Rabies Virus. *Cell* **170**, 382-
834   392 e314 (2017).
835   26.   S. Rogers, R. Wells, M. Rechsteiner, Amino acid sequences common to rapidly degraded
836   proteins: the PEST hypothesis. *Science* **234**, 364-368 (1986).
837   27.   D. A. Steinhauer, J. J. Holland, Rapid evolution of RNA viruses. *Annu Rev Microbiol* **41**,
838   409-433 (1987).
839   28.   D. A. Steinhauer, J. C. de la Torre, J. J. Holland, High nucleotide substitution error
840   frequencies in clonal pools of vesicular stomatitis virus. *J Virol* **63**, 2063-2071 (1989).
841   29.   J. J. Holland, J. C. De La Torre, D. A. Steinhauer, RNA virus populations as quasispecies.
842   *Curr Top Microbiol Immunol* **176**, 1-20 (1992).
843   30.   E. C. Holmes, C. H. Woelk, R. Kassis, H. Bourhy, Genetic constraints and the adaptive
844   evolution of rabies virus in nature. *Virology* **292**, 247-257 (2002).
845   31.   G. M. Jenkins, A. Rambaut, O. G. Pybus, E. C. Holmes, Rates of molecular evolution in
846   RNA viruses: a quantitative phylogenetic analysis. *J Mol Evol* **54**, 156-165 (2002).
847   32.   M. Combe, R. Sanjuan, Variation in RNA virus mutation rates across host cells. *PLoS
848   Pathog* **10**, e1003855 (2014).
849   33.   I. R. Wickersham, H. A. Sullivan, H. S. Seung, Production of glycoprotein-deleted rabies
850   viruses for monosynaptic tracing and high-level gene expression in neurons. *Nature
851   protocols* **5**, 595-606 (2010).
852   34.   F. Osakada, E. M. Callaway, Design and generation of recombinant rabies virus vectors.
853   *Nat Protoc* **8**, 1583-1601 (2013).
854   35.   I. R. Wickersham, H. A. Sullivan, Rabies viral vectors for monosynaptic tracing and
855   targeted transgene expression in neurons. *Cold Spring Harb Protoc* **2015**, 375-385
856   (2015).
857   36.   A. P. Weible *et al.*, Transgenic targeting of recombinant rabies virus reveals
858   monosynaptic connectivity of specific neurons. *The Journal of neuroscience : the official
859   journal of the Society for Neuroscience* **30**, 16509-16513 (2010).
860   37.   M. O. Carneiro *et al.*, Pacific biosciences sequencing technology for genotyping and
861   variation discovery in human data. *BMC Genomics* **13**, 375 (2012).
862   38.   C. Wirblich, M. J. Schnell, Rabies virus (RV) glycoprotein expression levels are not critical
863   for pathogenicity of RV. *J Virol* **85**, 697-704 (2011).

39. L. Tao *et al.*, Molecular basis of neurovirulence of flury rabies virus vaccine strains:
importance of the polymerase and the glycoprotein R333Q mutation. *J Virol* **84**, 8926-
8936 (2010).
40. C. Prehaud, S. Lay, B. Dietzschold, M. Lafon, Glycoprotein of nonpathogenic rabies
viruses is a key determinant of human cell apoptosis. *J Virol* **77**, 10537-10547 (2003).
41. L. Madisen *et al.*, A toolbox of Cre-dependent optogenetic transgenic mice for light-
induced activation and silencing. *Nature neuroscience* **15**, 793-802 (2012).
42. N. C. Shaner *et al.*, Improved monomeric red, orange and yellow fluorescent proteins
derived from Discosoma sp. red fluorescent protein. *Nat Biotechnol* **22**, 1567-1572
(2004).
43. M. Drobizhev, N. S. Makarov, S. E. Tillo, T. E. Hughes, A. Rebane, Two-photon absorption
properties of fluorescent proteins. *Nat Methods* **8**, 393-399 (2011).
44. K. W. Huang, B. L. Sabatini, Single-Cell Analysis of Neuroinflammatory Responses
Following Intracranial Injection of G-Deleted Rabies Viruses. *Front Cell Neurosci* **14**, 65
(2020).
45. M. Prosniak, D. C. Hooper, B. Dietzschold, H. Koprowski, Effect of rabies virus infection
on gene expression in mouse brain. *Proc Natl Acad Sci U S A* **98**, 2758-2763 (2001).
46. L. Madisen *et al.*, Transgenic mice for intersectional targeting of neural sensors and
effectors with high specificity and performance. *Neuron* **85**, 942-958 (2015).
47. K. Liu *et al.*, Lhx6-positive GABA-releasing neurons of the zona incerta promote sleep.
Nature **548**, 582-587 (2017).
48. T. K. Lavin, L. Jin, I. R. Wickersham, Monosynaptic tracing: a step-by-step protocol. *J
Chem Neuroanat* 10.1016/j.jchemneu.2019.101661, 101661 (2019).
49. T. K. Lavin, L. Jin, N. E. Lea, I. R. Wickersham, Monosynaptic Tracing Success Depends
Critically on Helper Virus Concentrations. *Front Synaptic Neurosci* **12**, 6 (2020).
50. M. E. Matlashov *et al.*, A set of monomeric near-infrared fluorescent proteins for
multicolor imaging across scales. *Nat Commun* **11**, 239 (2020).
51. E. Ciabatti *et al.*, Genomic stability of Self-inactivating Rabies. *bioRxiv*
10.1101/2020.09.19.304683, 2020.2009.2019.304683 (2020).
52. K. P. Dalton, J. K. Rose, Vesicular stomatitis virus glycoprotein containing the entire
green fluorescent protein on its cytoplasmic domain is incorporated efficiently into virus
particles. *Virology* **279**, 414-421 (2001).
53. W. P. Duprex, F. M. Collins, B. K. Rima, Modulating the function of the measles virus
RNA-dependent RNA polymerase by insertion of green fluorescent protein into the open
reading frame. *J Virol* **76**, 7322-7328 (2002).
54. S. Finke, K. Brzozka, K. K. Conzelmann, Tracking fluorescence-labeled rabies virus:
enhanced green fluorescent protein-tagged phosphoprotein p supports virus gene
expression and formation of infectious particles. *J Virol* **78**, 12333-12343 (2004).
55. M. L. Koser *et al.*, Rabies virus nucleoprotein as a carrier for foreign antigens. *Proc Natl
Acad Sci U S A* **101**, 9405-9410 (2004).
56. D. D. Brown *et al.*, Rational attenuation of a morbillivirus by modulating the activity of
the RNA-dependent RNA polymerase. *J Virol* **79**, 14330-14338 (2005).
57. S. C. Das, D. Nayak, Y. Zhou, A. K. Pattnaik, Visualization of intracellular transport of
vesicular stomatitis virus nucleocapsids in living cells. *J Virol* **80**, 6368-6377 (2006).

908 58. Y. Klingen, K. K. Conzelmann, S. Finke, Double-labeled rabies virus: live tracking of
909   enveloped virus transport. *Journal of virology* **82**, 237-245 (2008).

910 59. S. C. Das, D. Panda, D. Nayak, A. K. Pattnaik, Biarsenical labeling of vesicular stomatitis
911   virus encoding tetracysteine-tagged m protein allows dynamic imaging of m protein and
912   virus uncoating in infected cells. *J Virol* **83**, 2611-2622 (2009).

913 60. A. C. Marriott, C. A. Hornsey, Reverse genetics system for Chandipura virus: tagging the
914   viral matrix protein with green fluorescent protein. *Virus Res* **160**, 166-172 (2011).

915 61. T. K. Soh, S. P. Whelan, Tracking the Fate of Genetically Distinct Vesicular Stomatitis
916   Virus Matrix Proteins Highlights the Role for Late Domains in Assembly. *J Virol* **89**,
917   11750-11760 (2015).

918 62. J. Nikolic, A. Civas, Z. Lama, C. Lagaudriere-Gesbert, D. Blondel, Rabies Virus Infection
919   Induces the Formation of Stress Granules Closely Connected to the Viral Factories. *PLoS*
920   *Pathog* **12**, e1005942 (2016).

921 63. J. B. Case *et al.*, Replication-competent vesicular stomatitis virus vaccine vector protects
922   against SARS-CoV-2-mediated pathogenesis. *bioRxiv* 10.1101/2020.07.09.196386 (2020).

923 64. E. A. Gomme, C. Wirblich, S. Addya, G. F. Rall, M. J. Schnell, Immune clearance of
924   attenuated rabies virus results in neuronal survival with altered gene expression. *PLoS*
925   *Pathog* **8**, e1002971 (2012).

926 65. T. J. Wiktor, B. Dietzschold, R. N. Leamnson, H. Koprowski, Induction and biological
927   properties of defective interfering particles of rabies virus. *J Virol* **21**, 626-635 (1977).

928 66. A. Kawai, S. Matsumoto, Interfering and noninterfering defective particles generated by
929   a rabies small plaque variant virus. *Virology* **76**, 60-71 (1977).

930 67. H. F. Clark, N. F. Parks, W. H. Wunner, Defective interfering particles of fixed rabies
931   viruses: lack of correlation with attenuation or auto-interference in mice. *J Gen Virol* **52**,
932   245-258 (1981).

933 68. M. Faber *et al.*, Overexpression of the rabies virus glycoprotein results in enhancement
934   of apoptosis and antiviral immune response. *J Virol* **76**, 3374-3381 (2002).

935 69. X. Chen, Y. C. Sun, G. M. Church, J. H. Lee, A. M. Zador, Efficient in situ barcode
936   sequencing using padlock probe-based BaristaSeq. *Nucleic Acids Res* **46**, e22 (2018).

937 70. X. Chen *et al.*, High-Throughput Mapping of Long-Range Neuronal Projection Using In
938   Situ Sequencing. *Cell* **179**, 772-786 e719 (2019).

939 71. Y. C. Sun *et al.*, Integrating barcoded neuroanatomy with spatial transcriptional profiling
940   enables identification of gene correlates of projections. *Nat Neurosci* **24**, 873-885
941   (2021).

942 72. I. R. Wickersham *et al.*, Lentiviral vectors for retrograde delivery of recombinases and
943   transactivators. *Cold Spring Harb Protoc* **2015**, 368-374 (2015).

944 73. L. Madisen *et al.*, A robust and high-throughput Cre reporting and characterization
945   system for the whole mouse brain. *Nature neuroscience* **13**, 133-140 (2010).

SUPPLEMENTARY METHODS

Cloning

Lentiviral transfer plasmids were made by cloning, into pCSC-SP-PW-GFP (1) (Addgene #12337), the following components:

the CAG promoter (2) and a Cre-dependent "FLEX" (3) construct consisting of pairs of orthogonal lox sites flanking a back-to-back fusion of the gene for mTagBFP2 (4) immediately followed by the reverse-complemented gene for mCherry (5), to make the Cre reporter construct pLV-CAG-FLEX-BFP-(mCherry)' (Addgene 115234);

the CAG promoter (2) and a Flp-dependent "FLEX" (3) construct consisting of pairs of orthogonal FRT sites (6) flanking a back-to-back fusion of the gene for mTagBFP2 (4) immediately followed by the reverse-complemented gene for mCherry (5), to make the Flp reporter construct pLV-CAG-F14F15S-BFP-(mCherry)' (Addgene 115235);

the ubiquitin C promoter from pUB-GFP (7) (Addgene 11155) and the long isoform of TVA (8) to make the TVA expression vector pLV-U-TVA950 (Addgene 115236).

The first-generation vector genome plasmid pRVΔG-4FLPo (Addgene 122050) was made by cloning the FLPo gene (9) into pRVΔG-4Cre.

pAAV-syn-F14F15S-splitTVA-EGFP-tTA_v2 (Addgene 183352) is a FLP-dependent version of the Cre-dependent helper virus genome plasmid pAAV-syn-FLEX-splitTVA-EGFP-tTA (Addgene 52473) with orthogonal FRT sites(6) instead of orthogonal lox sites, and with a 2-bp frameshift to avoid creation of a premature stop codon in the first FRT site. Note that this plasmid is a replacement for pAAV-syn-F14F15S-splitTVA-EGFP-tTA, which was used in an earlier preprint of this manuscript.

pCAG-hypBase was made by synthesizing the 1785-bp gene for an improved version of piggyBac transposase (10) and cloning it into the EcoRI and NotI sites of pCAG-GFP (7) (Addgene 11150).

pB-CAG-TEVP-IRES-mCherry (Addgene 174377) was made by cloning the CAG promoter (2), a mammalian codon-optimized version (11, 12) of the TEVP gene (S219V mutant) (13), the EMCV IRES (14), and the mCherry gene (5) into pB-CMV-MCS-EF1-Puro (System Biosciences #PB510B-1).

pAAV-syn-FLPo (Addgene 174378) was made by cloning the FLPo gene(9) into the EcoRI and AccIII sites of pAAV-syn-FLEX-EGFP-B19G (Addgene 59333).

pAAV-TREtight-H2b-emiRFP670-TEVP (Addgene 174379) was made by cloning an H2b-emiRFP670 fusion gene (15) (sequence from Addgene 136571 but with the internal kozak sequence replaced by a short GSG linker to prevent translation of fluorophore unfused to H2b), followed by a P2A sequence and the above-described TEVP gene, into the EcoRI and NheI sites of pAAV-TREtight-mTagBFP2-B19G (16) (Addgene 100799).

pRVΔG-NPEST-4Cre (Addgene 174380) was made by cloning a synthesized fragment containing the 3' addition from Ciabatti et al. '17, with synonymous changes to 5 codons (17) in the immediate vicinity of the junction between the nucleoprotein gene and the 3' addition so that more than a single point mutation would be required to convert them to stop codons, into the PmlI and BstI sites of pRVΔG-4Cre(18) (Addgene 98034).

pRVΔG-N*PEST-4Cre (Addgene 174381) was made identically to the above except that the glycine codon in position 453 was replaced with a stop codon (TGA).

All of the above novel plasmids have been deposited with Addgene, with the accession numbers given above, and can be purchased from there except for pCAG-hypBase, the distribution of which is not permitted due to intellectual property constraints.

Production of lentiviral and adeno-associated viral vectors

Lentiviral vectors were made by transfection of HEK-293T/17 cells (ATCC 11268) as described (19) but using the vesicular stomatitis virus envelope expression plasmid pMD2.G (Addgene 12259) for all vectors except for LV-U-TVA950(B19G), which was made using the rabies virus envelope expression plasmid pCAG-B19GVSVGCD (19). Lentiviral vectors expressing fluorophores were titered as described (20); titers of LV-U-TVA950(VSVG) and LV-U-TVA950(B19G) were assumed to be approximately the same as those of the fluorophore-expressing lentiviral vectors produced in parallel.

AAV1-TREtight-mTagBFP2-B19G (which we have described previously(16, 21)), was packaged as serotype 1 by Addgene (catalog # 100798-AAV1).

AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2, AAV1-TREtight-H2b-emiRFP670-P2A-TEVP and AAV2-retro-syn-FLPo were made by transfecting HEK 293T/17 cells with the respective genome plasmid along with pHelper (Cellbiolabs VPK-421) and either pAAV-RC1 (Cellbiolabs VPK-421) or "rAAV2-retro helper" (Addgene 81070)(22), using Xfect Transfection Reagent (Takara 631318) according to the manufacturer's protocol, with collection of supernatant (and replacement with fresh media) at 3 days after transfection and collection of supernatant as well as the transfected cells at 5 days after transfection. Virus was pelleted from supernatants using PEG 8000; cells were lysed by four freeze-thaw cycles. Pelleted virus and cell lysate were pooled and treated with benzonase, purified on an iodixanol gradient, then concentrated in an Amicon Ultra-15 centrifugal filter unit (Millipore Sigma UFC9100).

Production of titering cell lines
To make reporter cell lines, HEK-293T/17 cells were infected with either pLV-CAG-FLEX-BFP-(mCherry)' or pLV-CAG-F14F15S-BFP-(mCherry)' at a multiplicity of infection of 100 in one 24-well plate well each. Cells were expanded to 2x 15cm plates each, then sorted on a FACS Aria to retain the top 10% most brightly blue fluorescent cells. After sorting, cells were expanded again to produce the cell lines 293T-FLEX-BC and 293T-F14F15S-BC, reporter lines for Cre and FLPo activity, respectively. TVA-expressing versions of these two cell lines were made by infecting one 24-well plate well each with LV-U-TVA950(VSVG) at an MOI of approximately 100; these cells were expanded to produce the cell lines 293T-FLEX-BC-TVA and 293T-F14F15S-BC-TVA.

Production of TEVP-expressing cell line
The 293T-TEVP cell line was made by transfecting HEK 293T/17 (ATCC CRL-11268) cells with pCAG-hypBase and pB-CAG-TEVP-IRES-mCherry in a 15 cm plate (293T/17) or one well each of a 24 well plate (BHK-B19G2 and BHK-EnvA2) using Lipofectamine 2000 (Thermo Fisher 11668019) according to the manufacturer's instructions, then expanding and sorting the cells on a FACSAria (BD Biosciences) for the brightest 20% of red fluorescent cells, then expanding and freezing the sorted cells.

Production and titering of rabies viruses
RV∆G-4Cre and RV∆GL-4Cre were produced as described (18, 23), with EnvA-enveloped viruses made by using cells expressing EnvA instead of G for the last passage. Titering and infection of cell lines with serial dilutions of viruses was as described (24), with the 293T-TVA-FLEX-BC and 293T-TVA-F14F15S-BC lines used for B19G-enveloped viruses and the 293T-TVA-FLEX-BC and 293T-TVA-F14F15S-BC used for the EnvA-enveloped viruses. For the *in vivo* injections, the three EnvA-enveloped, Cre-encoding viruses were titered side by side, and the two higher-titer viruses were diluted so that the final titer of the injected stocks of all three viruses were approximately equal at 1.39E9 infectious units (i.u.) per milliliter.

RV∆G-4mCherry(EnvA) was produced and titered as described (18, 23, 24), with final titer of 1.7E10 i.u./mL on 293T-TVA800 cells and 1.60E6 i.u./mL on HEK-293T cells. RV∆G-4Cre(EnvA) used for Supplementary Figure S5 was produced and titered as described (18, 23, 24), with final titers of 8.58E9 i.u./mL on 293T-TVA800 cells (and 3.14E5 i.u./mL on HEK-293T cells) and was diluted to a titer of 4.49E9 i.u./mL before injection.

The two new rabies viruses RV∆G-NPEST-4Cre and RV∆G-N*PEST-4Cre were produced as described(18, 23) but with only one amplification passage (P1) between the rescue transfection step and the final passage on EnvA-expressing cells. For the NPEST version, the following additional modifications were made: for the rescue transfection, the new cell line 293T-TEVP (see above) was used instead of HEK 293T-17, and pB-CAG-TEVP-IRES-mCherry (10 µg per 15 cm plate) was included in the plasmid mix. Because pilot testing showed no clear advantage of the 293T-TEVP line over transfected 293Ts for passaging, the P1 passage was on HEK 293T/17 cells transfected with equal amounts of pCAG-B19G and pB-CAG-TEVP-IRES-mCherry

(32 μg of each plasmid per 15 cm plate). The final passage was on BHK-EnvA2 cells(24) transfected with pB-CAG-TEVP-IRES-mCherry. Lipofectamine 2000 (Thermo Fisher 11668019) was used for all transfections, following the manufacturer's protocol, except for the final passage of RVΔG-NPEST-4Cre, for which Lipofectamine 3000 (Thermo Fisher L3000-015) was used.

The final stocks of RVΔG-NPEST-4Cre(EnvA) and RVΔG-N*PEST-Cre(EnvA) were titered side by side as described(24) on 293T-FLEX-BC-TVA cells (see above) in two-fold dilution series and on 293T-FLEX-BC cells (see above) undiluted (for determination of the titer of contaminating non-EnvA-enveloped virus). Final titers of the viruses on the TVA-expressing cells were determined to be 1.18E9 i.u./mL for RVΔG-NPEST-4Cre(EnvA) and 2.41E10 i.u./mL (normalized) for RVΔG-N*PEST-4Cre(EnvA). Prior to injection *in vivo* (see below), the N*PEST virus was diluted 20.42-fold in DPBS to match the titer of the NPEST version. Final titers of the viruses on the non-TVA-expressing cells were determined to be 3.79E5 i.u./mL for RVΔG-NPEST-4Cre(EnvA) and 8.28E5 i.u./mL (normalized) for RVΔG-N*PEST-4Cre(EnvA).

Immunostaining and imaging of cultured cells
Reporter cells (see above) plated on coverslips coated in poly-L-lysine (Sigma) were infected with serial dilutions of RVΔG-4Cre and RVΔGL-4Cre as described (24). Three days after infection, cells were fixed with 2% paraformaldehyde, washed repeatedly with blocking/permeabilization buffer (0.1% Triton-X (Sigma) and 1% bovine serum albumin (Sigma) in PBS), then labeled with a blend of three FITC-conjugated anti-nucleoprotein monoclonal antibodies (Light Diagnostics Rabies DFA Reagent, EMD Millipore 5100) diluted 1:100 in blocking buffer for 30 minutes, followed by further washes in blocking buffer, then finally briefly rinsed with distilled water and air-dried before mounting the coverslips onto microscope slides with Prolong Diamond Antifade (Thermo P36970) mounting medium. Images of wells at comparable multiplicities of infection (~0.1) were collected on a Zeiss 710 confocal microscope.

Extraction of viral genomic RNA and preparation for Sanger sequencing
RNA viral genomes were extracted from two Tripodi lab (EnvA/SiR-CRE and EnvA/Sir-FLPo) and three Wickersham lab (RVΔG-4mCherry, RVΔG-NPEST-4Cre(EnvA), and RVΔG-N*PEST-Cre(EnvA)) rabies virus samples using a Nucleospin RNA kit (Macherey-Nagel, Germany) and treated with DNase I (37°C for 1 hour, followed by 70°C for 5 minutes). Extracted RNA genomes were converted to complementary DNA using an AccuScript PfuUltra II RT-PCR kit (Agilent Technologies, USA) at 42°C for 2 hours with the following barcoded (so that individual viral particles' genomes would be marked with distinct barcodes) primer annealing to the rabies virus leader sequence:
Adapter_N8_leader_fp: TCAGACGATGCGTCATGCNNNNNNNNNACGCTTAACAACCAGATC

cDNA sequences from the leader through the first half of the rabies virus P gene were amplified using Platinum SuperFi Green Master Mix (Invitrogen (Thermo Fisher), USA) with cycling conditions as follows: denaturation at 98°C for 30 seconds, followed by 25 cycles of amplification (denaturation at 98°C for 5 seconds and extension at 72°C for 75 seconds), with a final extension at 72°C for 5 minutes, using the following primers:
pEX_adapter_fp: CAGCTCAGACGATGCGTCATGC
Barcode2_P_rp: GCAGAGTCATGTATAGCTTCTTGAGCTCTCGGCCAG

The ~2kb PCR amplicons were extracted from an agarose gel, purified with Nucleospin Gel and PCR Clean-up (Macherey-Nagel, Germany), and cloned into pEX-A (Eurofins Genomics, USA) using an In-Fusion HD Cloning Kit (Takara Bio, Japan). The cloned plasmids were transformed into Stellar competent cells (Takara Bio, Japan), and 200 clones per rabies virus sample were isolated and purified for sequencing. For each clone, the index and the 3' end of the N gene were sequenced until sequencing data was collected for over fifty clones per sample: 51 clones from SiR-CRE(EnvA), 50 from SiR-FLPo(EnvA), and 51 from RVΔG-4mCherry(EnvA). Although viral samples may contain plasmid DNA, viral mRNA, and positive-sense anti-genomic RNA, this RT-PCR procedure can amplify only the negative-sense RNA genome: the reverse transcription primer is designed to anneal to the leader sequence of the negative-strand genome so that cDNA synthesis can start from the negative-sense RNA genome, with no other possible templates. Additionally, the PCR amplifies the cDNA, not any plasmids which were transfected into producer cell lines during viral vector production, because the forward PCR primer anneals to the primer used in the reverse transcription, rather than any viral sequence. This RT-PCR protocol ensures that only negative-sense RNA rabies viral genomes can be sequenced.

Sanger sequencing of transgenes in SiR viruses

The procedure for sequencing the transgene inserts was the same as above, but with the RT primer being Adaptor_N8_M_fp (see below), annealing to the M gene and again with a random 8-nucleotide index to tag each clone, and with PCR primers pEX_adaptor_fp (see above) and Barcode2_L_rp (see below), to amplify the sequences from the 3' end of the M gene to the 5' end of the L gene, covering the iCre-P2A-mCherryPEST (or FLPo-P2A-mCherryPEST) sequence.

Primers for RT and PCR for Sanger sequencing were as follows:
Adaptor_N8_M_fp:
TCAGACGATGCGTCATGCNNNNNNNNCAACTCCAACCCTTGGGAGCA
Barcode2_L_rp:
GCAGAGTCATGTATAGTTGGGGACAATGGGGGGTTCC

Sanger sequencing analysis of PEST region in SiR and control viruses
Alignment and mutation detection were performed using SnapGene 4.1.9 (GSL Biotech LLC, USA). Reference sequences of the viral samples used in this study were based on deposited plasmids in Addgene: pSAD-F3-NPEST-iCRE-2A-mCherryPEST (Addgene #99608), pSAD-F3-NPEST-FLPo-2A-mCherryPEST (Addgene #99609), and pRVΔG-4mCherry (Addgene #52488). Traces corresponding to indices and mutations listed in Figure 1 and Supplementary File S1 were also manually inspected and confirmed.

Single-molecule, real-time (SMRT) sequencing
Double-stranded DNA samples for SMRT sequencing were prepared similarly to the above, except that that the clones generated from each of the three virus samples were tagged with one of the standard PacBio barcode sequences to allow identification of each clone's sample of origin following multiplex sequencing (see https://www.pacb.com/wp-content/uploads/multiplex-target-enrichment-barcoded-multi-kilobase-fragments-probe-based-capture-technologies.pdf and https://github.com/PacificBiosciences/Bioinformatics-Training/wiki/Barcoding-with-SMRT-Analysis-2.3). This was in addition to the random index (10 nucleotides in this case) that was again included in the RT primers in order to uniquely tag each individual genome.

RNA viral genomes were extracted from two Tripodi lab (SiR-CRE and Sir-FLPo) and one Wickersham lab (RVΔG-4Cre (18); see Addgene #98034 for reference sequence) virus samples using a Nucleospin RNA kit (Macherey-Nagel, Germany) and treated with DNase I (37°C for 1 hour, followed by 70° for 5 minutes). Primers for RT and PCR are listed below. PCR cycling conditions were as follows: denaturation at 98°C for 30 seconds, followed by 20 cycles of amplification (denaturation at 98°C for 5 seconds and extension at 72°C for 75 seconds), with a final extension at 72°C for 5 minutes. This left each amplicon with a 16bp barcode at each of its ends that indicated which virus sample it was derived from, in addition to a 10-nt index sequence that was unique to each genome molecule.

Primers for RT and PCR for SMRT sequencing were as follows:
RVΔG-4Cre:
    RT:
        Barcode1_cagc_N10_leader_fp:
        TCAGACGATGCGTCATCAGCNNNNNNNNNNACGCTTAACAACCAGATC
    PCR:
        Barcode1_cagc_fp: TCAGACGATGCGTCAT-CAGC
        Barcode2_P_rp (see above)

SiR-CRE:
    RT:
        Barcode5_cagc_N10_leader_fp:
        ACACGCATGACACACTCAGCNNNNNNNNNNACGCTTAACAACCAGATC
    PCR:
        Barcode5_cagc_fp: ACACGCATGACACACT-CAGC
        Barcode3_P_rp: GAGTGCTACTCTAGTACTTCTTGAGCTCTCGGCCAG

SiR-FLPo:
    RT:

Barcode9_cagc_N10_leader_fp:
CTGCGTGCTCTACGACCAGCNNNNNNNNNNACGCTTAACAACCAGATC
PCR:
Barcode9_cagc_fp: CTGCGTGCTCTACGAC-CAGC
Barcode4_P_rp: CATGTACTGATACACACTTCTTGAGCTCTCGGCCAG

After the amplicons were extracted and purified from an agarose gel, the three were mixed together at 1:1:1 molar ratio. The amplicons' sizes were confirmed on the Fragment Analyzer (Agilent Technologies, USA), then hairpin loops were ligated to both ends of the mixed amplicons to make circular SMRTbell templates for Pacbio Sequel sequencing. SMRTbell library preparation used the PacBio Template Preparation Kit v1.0 and Sequel Chemistry v3. Samples were then sequenced on a PacBio Sequel system running Sequel System v.6.0 (Pacific Biosciences, USA), with a 10-hour movie time.

Bioinformatics for PacBio sequence analysis

For the ~2kb template, the DNA polymerase with a strand displacement function can circle around the template and hairpins multiple times; the consensus sequence of multiple passes yields a CCS (circular consensus sequence) read for each molecule. Raw sequences were initially processed using SMRT Link v.6.0 (Pacific Biosciences, USA). Sequences were filtered for a minimum of read length 10 bp, pass 3, and read score 65. 127,178 CCS reads were filtered through passes 3 and Q10; 89,188 CCS reads through passes 5 and Q20; 29,924 CCS reads through passes 8 and Q30. Downstream bioinformatics analysis was performed using BLASR V5.3.2 for the alignment, bcftools v.1.6 for variant calling. Mutations listed in Figure 2 and Supplementary File S2 were also manually inspected and confirmed using Integrative Genomics Viewer 2.3.32 (software.broadinstitute.org/software/igv/). Analysis steps included the following: 1. Exclude CCS reads under 1000 bases, which may have been derived from non-specific reverse transcription or PCR reactions. 2. Classify the CCS reads to the three samples, according to the PacBio barcodes on the 5' ends. 3. For any CCS reads that contain the same 10-nucleotide random index, select only one of them, to avoid double-counting of clones derived from the same cDNA molecule. 4. Align the reads to the corresponding reference sequence (see Supplementary Files S3-S5). 5. Count the number of mutations at each nucleotide position of the reference sequences.

Surgeries and virus injections for two-photon imaging

All experimental procedures using mice were conducted according to NIH guidelines and were approved by the MIT Committee for Animal Care (CAC). Mice were housed 1-4 per cage under a normal light/dark cycle for all experiments.

Adult (>9 weeks, male and female) Cre-dependent tdTomato reporter Ai14 (25) (Jackson Laboratory #007908) or Arch-EGFP reporter Ai35D (26) (Jackson Laboratory # 012735) mice were anesthetized with isoflurane (4% in oxygen) and ketamine/xylazine (100mg/kg and 10mg/kg respectively, i.p.). Mice were given buprenorphine (0.1 mg/kg s.q.) and meloxicam (2 mg/kg s.q.) as preemptive analgesics, as well as eye ointment (Puralube); the scalp was then shaved, depilated with Nair, and thoroughly rinsed before the mice were mounted on a stereotaxic instrument (Stoelting Co.) with a hand warmer (Heat Factory) underneath the animal to maintain body temperature. The scalp was then disinfected with povidone-iodine, and an incision was made at the appropriate location (see below).

A 3 mm craniotomy was opened over primary visual cortex (V1). 300 nl of LV-U-TVA950(B19G) (see above) was injected into V1 (-2.70 mm AP, 2.50 mm LM, -0.26 mm DV; AP and LM stereotaxic coordinates are with respect to bregma; DV coordinate is with respect to brain surface) using a custom injection apparatus comprised of a hydraulic manipulator (MO-10, Narishige) with headstage coupled via custom adaptors to a wire plunger advanced through pulled glass capillaries (Wiretrol II, Drummond) back-filled with mineral oil and front-filled with virus solution. Glass windows composed of a 3mm-diameter glass coverslip (Warner Instruments CS-3R) glued (Optical Adhesive 61, Norland Products) to a 5mm-diameter glass coverslip (Warner Instruments CS-5R) were then affixed over the craniotomy with Metabond (Parkell). Seven days after injection of the lentiviral vector, the coverslips were removed and 300 nl of one of the three EnvA-enveloped rabies viral vectors (with equalized titers as described above) was injected at the same stereotaxic coordinates. Coverslips were reapplied and custom stainless steel headplates (eMachineShop) were affixed to the skulls around the windows.

In vivo two-photon imaging and image analysis
Beginning seven days after injection of each rabies virus and continuing every seven days up to a maximum of four weeks following rabies virus injection, the injection sites were imaged on a Prairie/Bruker Ultima IV In Vivo two-photon microscope driven by a Spectra Physics Mai-Tai Deep See laser with a mode locked Ti:sapphire laser emitting at a wavelength of 1020 nm for tdTomato and mCherry or 920 nm for EGFP. Mice were reanesthetized and mounted via their headplates to a custom frame, again with ointment applied to protect their eyes and with a handwarmer maintaining body temperature. One field of view was chosen in each mouse in the area of maximal fluorescent labelling. The imaging parameters were as follows: image size 512 X 512 pixels (282.6 μm x 282.6 μm), 0.782 Hz frame rate, dwell time 4.0 μs, 2x optical zoom, Z-stack step size 1 μm. Image acquisition was controlled with Prairie View 5.4 software. Laser power exiting the 20x water-immersion objective (Zeiss, W plan-apochromat, NA 1.0) varied between 20 and 65 mW depending on focal plane depth (Pockel cell value was automatically increased from 450 at the top section of each stack to 750 at the bottom section). For the example images of labeled cells, maximum intensity projections (stacks of 150-400 μm) were made with Fiji software. Cell counting was performed with the ImageJ Cell Counter plugin. When doing cell counting, week 1 tdTomato labelled cells were defined as a reference; remaining week 1 cells were the same cells at later time point that align with week 1 reference cells but the not-visible cells at week 1 (the dead cells). Plots of cell counts were made with Origin 7.0 software (OriginLab, Northampton, MA). For the thresholded version of this analysis (Supplementary Figure S4), in order to exclude cells that could possibly have been labeled only with mCherry in the SiR-CRE group, only cells with fluorescence intensity greater than the average of the mean red fluorescence intensities of cells imaged in Ai35 versus Ai14 mice at the same laser power at 1020 nm at 7 days postinjection (32.33 a.u.) were included in the population of cells tracked from 7 days onward.

Monosynaptic tracing experiments: Sanger sequencing of viral genomic RNA
Sanger sequencing of RVΔG-NPEST-4Cre(EnvA) and RVΔG-N*PEST-4Cre(EnvA) was as described above but with the following modifications. The following barcoded primer was used for RT-PCR.
Adaptor-UMI-N_fp_57:
ACACTCTTTCCCTACACGACGCTCTTCCGATCTNNNNNNNNNNNNNNNNNNNNNAGAAGTCCGGAGGCTGTTTAT
cDNA sequences from the nucleoprotein gene through the first half of the rabies virus P gene were amplified using Platinum SuperFi Green Master Mix (Invitrogen (Thermo Fisher), USA) with cycling conditions as follows: denaturation at 98°C for 30 seconds, followed by 21 cycles of amplification (denaturation at 98°C for 5 seconds, annealing at 60°C for 10 seconds and extension at 72°C for 21 seconds), with a final extension at 72°C for 5 minutes, using the following primers:
i5-anchor_CTAGCGCT_fp_56:
AATGATACGGCGACCACCGAGATCTACACCTAGCGCTACACTCTTTCCCTACACGAC
P_Sanger_rp_56:
CAAGCAGAAGACGGCACGATTTTCCATCATCCAGGTG

The 700bp PCR amplicons for RVdG-NPEST-4Cre (EnvA) or RVΔG-N*PEST-4Cre (EnvA) virus were cloned into pEX-A (Eurofins Genomics, USA) using an In-Fusion HD Cloning Kit (Takara Bio, Japan) as above. The plasmids were transformed into Stellar competent cells (Takara Bio, Japan), and 32 clones from each rabies virus sample were isolated and purified for sequencing.

Monosynaptic tracing experiments: surgeries and virus injections
For pilot testing of AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2 virus (Supplementary Fig. S5), 200 nl of AAV2-retro-syn-FLPo (1.16E13 g.c/mL) was injected into dorsolateral striatum (AP = +0.74 mm w.r.t. bregma, LM = 2.25 mm w.r.t. bregma, DV = -2.20 mm w.r.t the brain surface) (except when omitted for no-Flpo controls); in the same surgery, 250 nl of helper virus mixture (AAV1-syn-F14F15S-sTpEptTA_v2 (diluted to 7.88E10 g.c./mL) mixed with AAV1-TREtight-mTagBFP2-B19G (diluted to 6.50E11) in a 50/50 ratio by volume) was injected into layer 5 of barrel cortex (AP -1.55 mm w.r.t. bregma, LM 3.00 mm w.r.t. bregma, DV -0.75 mm w.r.t the brain surface) in Ai14 (het) mice. 7 days after AAV injection, 250nl of RVΔG-Cre (EnvA) (8.58E9 i.u./mL, diluted to 4.49E09 i.u./mL) was injected at the same cortical injection site as the helper viruses.

For comparisons of NPEST and N*PEST viruses, the three helper AAVs were combined at final titers of 3.6E10 genome count ("g.c.")/mL for AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2, 6.60E11 g.c./mL for AAV1-TREtight-mTagBFP2-B19G (when included) and AAV1-TREtight-H2b-emiRFP670-P2A-TEVP (when included) in DPBS

(Fisher, 14-190-250). 250 nl of helper virus mixture was injected into layer 5 of barrel cortex (AP -1.55 mm w.r.t. bregma, LM 3.00 mm w.r.t. bregma, -DV 0.75 mm w.r.t. brain surface) of Ai14 mice; in the same surgery, 300 nl of AAV2-retro-syn-FLPo (1.16E13 g.c./mL) was injected into dorsolateral striatum (AP 0.74 mm w.r.t. bregma, LM 2.25 mm w.r.t. bregma, DV -2.30 mm w.r.t the brain surface). 7 days after AAV injection, 300nl of RVΔG-NPEST-4Cre(EnvA) (1.86E9 i.u./mL) or RVΔG-N*PEST-4Cre(EnvA) (diluted 12.96-fold in DPBS from 2.41E10 i.u./mL to 1.86E9 i.u./mL) was injected in barrel cortex at the same site as the helper AAV mixtures.

Monosynaptic tracing experiments: perfusions and histology
12 days or 3 weeks (depending on experiment; see main text) after injection of rabies virus, mice were transcardially perfused with 4% paraformaldehyde in phosphate-buffered saline. Brains were postfixed overnight in 4% paraformaldehyde in PBS on a shaker at 4°C and cut into 50 µm coronal sections on a vibrating microtome (Leica, VT-1000S). Sections were collected anterior to posteriorly into 6 tubes containing cryoprotectant. Collection goes on for 15 rounds so that each tube contains a sixth of the collected tissue (15 sections in each tube). Sections were immunostained as described(21) with a chicken anti-GFP primary antibody (Aves Labs GFP-1020) 1:500 and donkey anti-chicken Alexa Fluor 488 secondary antibody (Jackson Immuno 703-545-155) 1:200. Sections were mounted with Prolong Diamond Antifade mounting medium (Thermo Fisher P36970).

Monosynaptic tracing experiments: cell counts and microscopy
Coronal sections between 1.2mm and -3.3mm relative to bregma were examined under an epifluorescence microscope (Zeiss, Imager.Z2). When necessary due to high density of labeled cells, images were taken with the same microscope for cell counting. tdTomato-labeled neurons in contralateral cortex and thalamus were counted manually with the Cell Counter plugin in ImageJ. For cells at the injection site, when tdTomato expressing cells were few and sparse (usually less than 100 per section), cells coexpressing mTagBFP2 or H2b-emiRFP670 alongside were counted manually adding separate labels to each and then looking for overlapping cells. When tdTomato expressing cells were dense, tdTomato labeled cells were first counted using the Analyze Particle function in ImageJ (size in micron^2: 20-400; circularity: 0.20-1.00). The outline of these cells was then merged on top of images of mTagBFP2 and H2b-emiRFP670 labeled cells for the counting of the overlapping cells. Only one of the six series of sections (i.e., every sixth section: see above) was counted for each mouse.

Images for figures were taken on a confocal microscope (Zeiss, LSM 900). So that the confocal images of brain tissue included in the figures in this paper are representative of each group, the images were taken after the counts were conducted (see above), in each case using the mouse with the middle number of labeled neurons in that group (i.e., neither the highest nor the lowest in the group of 3 mice used for each condition).

REFERENCES FOR SUPPLEMENTARY METHODS

1. Marr RA*, et al.* (2004) Neprilysin regulates amyloid Beta peptide levels. *J Mol Neurosci* 22(1-2):5-11.
2. Niwa H, Yamamura K, & Miyazaki J (1991) Efficient selection for high-expression transfectants with a novel eukaryotic vector. *Gene* 108(2):193-199.
3. Atasoy D, Aponte Y, Su HH, & Sternson SM (2008) A FLEX switch targets Channelrhodopsin-2 to multiple cell types for imaging and long-range circuit mapping. *The Journal of neuroscience : the official journal of the Society for Neuroscience* 28(28):7025-7030.
4. Subach OM, Cranfill PJ, Davidson MW, & Verkhusha VV (2011) An enhanced monomeric blue fluorescent protein with the high chemical stability of the chromophore. *PLoS One* 6(12):e28674.
5. Shaner NC*, et al.* (2004) Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma sp. red fluorescent protein. *Nat Biotechnol* 22(12):1567-1572.
6. Turan S, Kuehle J, Schambach A, Baum C, & Bode J (2010) Multiplexing RMCE: versatile extensions of the Flp-recombinase-mediated cassette-exchange technology. *J Mol Biol* 402(1):52-69.
7. Matsuda T & Cepko CL (2004) Electroporation and RNA interference in the rodent retina in vivo and in vitro. *P Natl Acad Sci USA* 101(1):16-22.
8. Bates P, Young JA, & Varmus HE (1993) A receptor for subgroup A Rous sarcoma virus is related to the low density lipoprotein receptor. *Cell* 74(6):1043-1051.
9. Raymond CS & Soriano P (2007) High-efficiency FLP and PhiC31 site-specific recombination in mammalian cells. *PLoS One* 2(1):e162.

10. Yusa K, Zhou L, Li MA, Bradley A, & Craig NL (2011) A hyperactive piggyBac transposase for mammalian applications. *Proc Natl Acad Sci U S A* 108(4):1531-1536.

11. Gray DC, Mahrus S, & Wells JA (2010) Activation of specific apoptotic caspases with an engineered small-molecule-activated protease. *Cell* 142(4):637-646.

12. Yang CF*, et al.* (2013) Sexually dimorphic neurons in the ventromedial hypothalamus govern mating in both sexes and aggression in males. *Cell* 153(4):896-909.

13. Kapust RB*, et al.* (2001) Tobacco etch virus protease: mechanism of autolysis and rational design of stable mutants with wild-type catalytic proficiency. *Protein Eng* 14(12):993-1000.

14. Gallardo HF, Tan C, & Sadelain M (1997) The internal ribosomal entry site of the encephalomyocarditis virus enables reliable coexpression of two transgenes in human primary T lymphocytes. *Gene Ther* 4(10):1115-1119.

15. Matlashov ME*, et al.* (2020) A set of monomeric near-infrared fluorescent proteins for multicolor imaging across scales. *Nat Commun* 11(1):239.

16. Liu K*, et al.* (2017) Lhx6-positive GABA-releasing neurons of the zona incerta promote sleep. *Nature* 548(7669):582-587.

17. Ciabatti E, Gonzalez-Rueda A, Mariotti L, Morgese F, & Tripodi M (2017) Life-Long Genetic and Functional Access to Neural Circuits Using Self-Inactivating Rabies Virus. *Cell* 170(2):382-392 e314.

18. Chatterjee S*, et al.* (2018) Nontoxic, double-deletion-mutant rabies viral vectors for retrograde targeting of projection neurons. *Nat Neurosci* 21(4):638-646.

19. Wickersham IR*, et al.* (2015) Lentiviral vectors for retrograde delivery of recombinases and transactivators. *Cold Spring Harb Protoc* 2015(4):368-374.

20. Sullivan HA & Wickersham IR (2015) Concentration and purification of rabies viral and lentiviral vectors. *Cold Spring Harb Protoc* 2015(4):386-391.

21. Lavin TK, Jin L, & Wickersham IR (2019) Monosynaptic tracing: a step-by-step protocol. *J Chem Neuroanat*:101661.

22. Tervo DG*, et al.* (2016) A Designer AAV Variant Permits Efficient Retrograde Access to Projection Neurons. *Neuron* 92(2):372-382.

23. Wickersham IR & Sullivan HA (2015) Rabies viral vectors for monosynaptic tracing and targeted transgene expression in neurons. *Cold Spring Harb Protoc* 2015(4):375-385.

24. Wickersham IR, Sullivan HA, & Seung HS (2010) Production of glycoprotein-deleted rabies viruses for monosynaptic tracing and high-level gene expression in neurons. *Nature protocols* 5(3):595-606.

25. Madisen L*, et al.* (2010) A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. *Nature neuroscience* 13(1):133-140.

26. Madisen L*, et al.* (2012) A toolbox of Cre-dependent optogenetic transgenic mice for light-induced activation and silencing. *Nature neuroscience* 15(5):793-802.

Supplementary File S1: Sanger sequencing data of all clones shown in Figure 1.
51 clones derived from SiR-CRE, 50 from SiR-FLPo, and 51 from RVΔG-4mCherry are identified by their unique indices. All of the indices as well as the sequences corresponding to the 3' end of the nucleoprotein gene are shown.

Supplementary File S1

Random 8-nucleotide index

N gene, tabacco etch virus protease cleavage site, PEST, transcription stop/polyA signal, mutation

---

Gly453* in N-TEV-PEST of SiR-CRE (50/51 clones)

TCTCGGAA, TTATGGGC, CCTGCCGA, CGTCGTTG, ACTCTAGT, GATCCGGT, ATTATGTA, TATTATCC, ATGCTAGA, GTTGTTTC, GAGTCGAC, GTCGACCT, CAAGAGTA, CGAAGCCA, GCGTGATA, CGTCCAGA, CTTGGTCG, GGGGTTTT, GTCCATAC, GTTTAATA, CAAAATCC, GTGGAGGC, CGGAAGGT, AGGACGGG, CCTCGGGA, GCATTTGC, TGGTTCGG, ATTTAGTC, ATGGGTAA, CTACGGGG, CCTGGTAC, GCATTTCA, AGTGTAGG, CGCATAGA, GTCACAAT, GTATGCTT, TCGGGTTT, ATGTTGTC, CAGCCGTA, GAGTGGCC, GGCCCATT, GGATTTCC, GTACAAGC, GAAGATAT, AAACAGAT, AAGGCCTT, CGTTCGTC, AAAGACTA, ACTGAAAG, TGGGGATA

ATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATTCGAGTGACTCAGGTTCCTGAGAGA ACCTCTACTTCCAATCGGGATCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCT GCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGT AAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA

Asp449Glufs*16 in N-TEV-PEST of SiR-CRE (1/51 clones)

AGGCTAAC

ATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATTCGAGTGAACTCAGGTTCCTGAGAG AACCTCTACTTCCAATCGGGATCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGC TGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTG TAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA

No mutations in sequenced region of N-TEV-PEST of SiR-FLPo (4/50 clones)

TCATTTAT, AAGTCGAA, TGAATACA, TATACACG

ATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATTCGAGTGACTCAGGTTCCGGAGAGA ACCTCTACTTCCAATCGGGATCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCT GCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGT AAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA

Gly453* in N-TEV-PEST of SiR-FLPo (18/50 clones)

GAGGCCGC, CTTATCAC, TCAGTTTT, CTTCAGCA, TTACTTTT, AAAATAAG, CCATTTCT, GTCACATT, GCATACGG, AGTATTAA, CCGGGTTG, GGTAAAAT, CAGGGAGA, CATTTTTC, TGACAAAT, TCGTATAT, TAGGGATT, GATCCCCG

ATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATTCGAGTGACTCAGGTTCCTGAGAGA ACCTCTACTTCCAATCGGGATCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCT GCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGT AAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA

Ser450* in N-TEV-PEST of SiR-FLPo (28/50 clones)

TACTAAAT, TTCAACTA, TAATAGGC, CGCGGGTC, AGGGTGCC, TTGCTGAC, CGGCCCTT, TACAGCGA, GAGTAGGA, GTAAACGC, CTATGGGT, TTGATTGA, AGGATTCT, GAGGTTTG, TACTTAGC, CCTGTAGT, GGGGTGGA, TTAAGTTG, CTCAATGA, CAGATCTG, GTCTGGTA, GCGTTGGG, TAAACGAG, TACTTGAG, CGGGGTGT, CGATCCCG, TGGGTGTC, GCCACGAT

ATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATTCGAGTGACTGAGGTTCCGGAGAGA ACCTCTACTTCCAATCGGGATCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCT GCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGT AAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA

No mutations in sequenced region of RVΔG-4mCherry (51/51 clones)

CGCCCGCG, GCCAATCT, GGCGGAAT, TGTGGAAC, GGAGGGAT, CGTAGTGT, AGAATCTC, TGTCTGGC, GCCTTTTA, TGGAATCT, TTGTAATG, AGGCCTGT, CGGATATA, AATCCAAA, AAGGAAAA, GAACTCAT, GCCGCTTC, CTTTGCCG, CGCAACCT, TCGGGCAT, GGGCGACT, AACTGGAT, TGCGTCGG, TCAAAGCG, CGAGGCCC, ATGTAGGA, AGATACGT, TGGCTTCG, ATTTTCTA, TCGTCCGG, TCGGAGCG, CTGTTATA, TACCGTTC, CGAATGTC, CCCTCTTT, TGAGTGGG, TGTAAATA, AAGGAGTC, ATTACCCT, TCCCTGAC, TGGTAAAA, ACTATCTC, CCGGAAGC, CTTGGAGG, GCAACAAT, ATTACCGT, AATAGCAG, ACCATAGT, GGGTTGGT, CATTTATT, GGGTACAC

ATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATTCGAGTGACTCATAAGAAGTTGAATA ACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA

Supplementary File S2: Summary tables of SMRT sequencing data.
These tables show all mutations occurring at positions mutated at greater than 2% frequency in the three virus samples analyzed. Position numbers in these tables refer to the sequences in the three Genbank files below (Supplementary Files S3-S5).

SINGLE-MOLECULE, REAL-TIME SEQUENCING RESULTS

Frameshifts and insertions

Position numbers in this file refer to the reference sequences included as Supplementary Files S4-S5. A "frameshift" is included in Tables 1a-2b if the number of deleted bases in positions 1439-1492 (the vicinity of the junction of the end of the N gene and the intended 3' addition) is not an integer multiple of 3, with insertions ignored. "Any error" includes either the apparent frameshifts, or the new TAA/TAG/TGA stop codons, or both, with insertions ignored. The number of "frameshifts" increases considerably if insertion mutations are included in the calculation, indicating that there is a much higher insertion rate as compared to that of deletion; however, previous studies have found that spurious insertions are high with SMRT (see main text), so we ignore insertions in this paper apart from summarizing the data below.

SiR-CRE		CCS3	CCS5	CCS8
Position	**Mutation**	**CCS3**	**CCS5**	**CCS8**
# Sequences		22205	866	239
243	**GAT (reference)**	20273	784	209
	GCT	3	0	0
	GGT	5	0	0
	GTT	1856	79	29
	DEL	68	3	1
1111	**GAA (reference)**	19807	760	197
	GAC	1	0	0
	GAG	2396	105	42
	GAT	0	0	0
	DEL	1	1	0
1457	AGA	6	0	0
	CGA	1	0	0
	GGA (reference)	140	5	1
	TGA	22032	858	237
	DEL	26	3	1
[1439, 1492]	Frame shift (#DELs not an integer multiple of 3, Insertion ignored)	507	13	3
[1439, 1492]	Any error (insertion ignored)	22104	864	239
% mutation in [1439, 1492]		100%	100%	100%

Table 1a. All mutations in the SiR-CRE sample at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8), as well as frameshift mutations found in the C-terminal region of N.

SiR-CRE		CCS3	CCS5	CC8
Position	**Mutation**	**CCS3**	**CCS5**	**CC8**
[1439, 1492]	Number of frame shifts due to DEL (number of DELs not an integer multiple of 3, insertion ignored)	507	13	3
[1439, 1492]	Number of frame shifts due to either DEL or INS or both (Sequence length not an integer multiple of 3)	2504	89	14
[1439, 1492]	Any error (insertion included)	22174	865	239

Table 1b. Frameshift mutations in the C-terminal region of N in the SiR-CRE sample at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8).

SiR-FLPo		CCS3	CCS5	CCS8
Position	**Mutation**	**CCS3**	**CCS5**	**CCS8**
#Sequences		17086	695	210
1449	TAA	28	1	0
	TCA (reference)	8405	333	94
	TGA	8624	360	115
	TTA	3	0	0
	DEL	26	1	1
1457	AGA	3	0	0
	CGA	0	0	0
	GGA (reference)	11088	462	139
	TGA	5979	233	71
	DEL	16	0	0
[1439, 1492]	Frameshifts (#DELs not an integer multiple of 3, Insertion ignored)	448	11	3
[1439, 1492]	Any error (insertion ignored)	14444	589	180
% of mutation		**85%**	**85%**	**86%**

Table 2a. All mutations in the SiR-FLPo sample at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8), as well as frameshift mutations found in the C-terminal region of N.

SiR-FLPo		CCS3	CCS5	CCS8
Position	Mutation	**CCS3**	**CCS5**	**CCS8**
[1439, 1492]	Number of frame shifts due to DEL (number of DELs not an integer multiple of 3, insertion ignored)	448	11	3
[1439, 1492]	Number of frame shifts due to either DEL or INS or both (Sequence length not an integer multiple of 3)	3562	121	24
[1439, 1492]	Any error (insertion included)	15818	642	196

Table 2b. Frameshift mutations in the C-terminal region of N in the SiR-FLPo sample at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8).

RV∆G-4Cre		CCS3	CCS5	CC8
Position	**Mutation**	**CCS3**	**CCS5**	**CC8**
# Sequences		17978	757	254
1355	ACT	1706	84	28
	CCT	3	0	0
	GCT	1	1	1
	TCT (reference)	16139	667	224
	DEL	129	5	1
Total				

Table 3. All mutations in the SiR-FLPo sample at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8).

Tables of all mutations above 2% frequency threshold

Table 4a to 4c list all single-nucleotide substitutions and deletions at positions mutated at >2% threshold frequency. The percentage of mutations is calculated based on the total number of single nucleotide and deletion mutations divided by the total number of reads aligned, when insertion mutations are ignored. Deletion mutations dominate in the medium-frequency range between 2% and 5%.

	Reference	Position	A	C	G	T	DEL	Un-mutated	Mutated (SNP/DEL)	% Mutation (SNP/DEL)
CCS3	G	1457	6	1	140	22032	26	140	22065	99.4
	A	1111	19807	1	2396	0	1	19807	2398	10.8
	A	243	20273	3	5	1856	68	20273	1932	8.7
	T	615	4	0	2	21229	970	21229	976	4.4
	G	1124	5	0	21356	7	837	21356	849	3.8
	A	81	21358	9	2	0	836	21358	847	3.8
	A	31	21381	0	0	0	824	21381	824	3.7
	T	713	1	2	4	21487	711	21487	718	3.2
	T	338	0	0	9	21498	698	21498	707	3.2
	A	1665	21544	1	4	0	656	21544	661	3.0
	A	53	21606	0	0	1	598	21606	599	2.7
	C	411	8	21683	0	1	513	21683	522	2.4
	A	838	21718	1	0	1	485	21718	487	2.2
CCS5	G	1457	0	0	5	858	3	5	861	99.4
	A	1111	760	0	105	0	1	760	106	12.2
	A	243	784	0	0	79	3	784	82	9.5
	A	31	828	0	0	0	38	828	38	4.4
	G	1124	0	0	831	0	35	831	35	4.0
	T	615	0	0	0	834	32	834	32	3.7
	A	81	837	1	1	0	27	837	29	3.3
	T	338	0	0	1	840	25	840	26	3.0
	T	713	0	0	1	841	24	841	25	2.9
	C	411	0	843	0	0	23	843	23	2.7
	A	53	846	0	0	0	20	846	20	2.3
CCS8	G	1457	0	0	1	237	1	1	238	99.6
	A	1111	197	0	42	0	0	197	42	17.6
	A	243	209	0	0	29	1	209	30	12.6
	A	31	228	0	0	0	11	228	11	4.6
	T	615	0	0	0	232	7	232	7	2.9
	G	1124	0	0	232	0	7	232	7	2.9
	C	411	0	233	0	0	6	233	6	2.5

Table 4a. SiR-CRE: substitutions and deletions at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8).

	Reference	Position	A	C	G	T	DEL	Un-mutated	Mutated (SNP/DEL)	% Mutation (SNP/DEL)
CCS3	C	1449	28	8405	8624	3	26	8405	8681	50.8
	G	1457	3	0	11088	5979	16	11088	5998	35.1
	T	615	4	0	0	16322	760	16322	764	4.5
	A	81	16410	4	2	5	665	16410	676	4.0
	G	1124	0	0	16410	3	673	16410	676	4.0
	T	713	1	0	2	16500	583	16500	586	3.4
	A	1665	16536	1	1	0	548	16536	550	3.2
	A	31	16547	0	0	0	539	16547	539	3.2
	T	338	0	1	4	16582	499	16582	504	2.9
	A	53	16672	2	2	0	410	16672	414	2.4
	C	411	6	16683	1	0	396	16683	403	2.4
	A	838	16696	0	0	0	390	16696	390	2.3
	G	208	301	0	16739	44	2	16739	347	2.0
CCS5	C	1449	1	333	360	0	1	333	362	52.1
	G	1457	0	0	462	233	0	462	233	33.5
	A	81	665	0	0	0	30	665	30	4.3
	T	615	1	0	0	668	26	668	27	3.9
	A	1665	669	0	1	0	25	669	26	3.7
	T	713	0	0	0	671	24	671	24	3.5
	T	338	0	0	0	673	22	673	22	3.2
	C	411	0	676	0	0	19	676	19	2.7
	G	1124	0	0	676	0	19	676	19	2.7
	A	838	679	0	0	0	16	679	16	2.3
	G	346	0	0	680	0	15	680	15	2.2
	A	31	681	0	0	0	14	681	14	2.0
CCS8	C	1449	0	94	115	0	1	94	116	55.2
	G	1457	0	0	139	71	0	139	71	33.8
	T	713	0	0	0	203	7	203	7	3.3
	A	53	205	0	0	0	5	205	5	2.4
	A	81	205	0	0	0	5	205	5	2.4

Table 4b. SiR-FLPo: substitutions and deletions at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8).
.

	Reference	Position	A	C	G	T	DEL	Un-mutated	Mutated (SNP/DEL)	% Mutation (SNP/DEL)
CCS3	T	1355	1706	3	1	16139	129	16139	1839	10.2
	A	31	16732	0	0	0	1246	16732	1246	6.9
	T	615	2	0	1	17266	709	17266	712	4.0
	A	81	17300	4	2	1	671	17300	678	3.8
	G	1124	0	0	17379	4	595	17379	599	3.3
	T	713	0	1	1	17469	507	17469	509	2.8
	T	338	1	0	2	17503	472	17503	475	2.6
	A	1506	17520	0	0	0	458	17520	458	2.5
	A	53	17579	1	2	1	395	17579	399	2.2
CCS5	T	1355	84	0	1	667	5	667	90	11.9
	A	31	703	0	0	0	54	703	54	7.1
	A	81	717	0	0	0	40	717	40	5.3
	T	615	0	0	0	729	28	729	28	3.7
	G	1124	0	0	738	0	19	738	19	2.5
	A	53	739	0	0	0	18	739	18	2.4
	T	713	0	0	0	740	17	740	17	2.2
	A	1506	740	0	0	0	17	740	17	2.2
CCS8	T	1355	28	0	1	224	1	224	30	11.8
	A	31	241	0	0	0	13	241	13	5.1
	T	615	0	0	0	248	6	248	6	2.4

Table 4c. RVΔG-4Cre: substitutions and deletions at positions mutated at >2% frequency at all stringencies (CCS3, CCS5, CCS8).

Supplementary File S3: SiR-CRE amplicon reference sequence.
This Genbank-format file contains the expected (i.e., based on the published sequence: Addgene #99608) sequence of amplicons obtained from SiR-CRE for SMRT sequencing.

```
LOCUS           Exported                 2183 bp ds-DNA      linear   SYN
07-JAN-2019
DEFINITION  synthetic linear DNA.
ACCESSION   .
VERSION     .
KEYWORDS    .
SOURCE      synthetic DNA construct
  ORGANISM  synthetic DNA construct
REFERENCE   1  (bases 1 to 2183)
  AUTHORS   Trial User
  TITLE     Direct Submission
  JOURNAL   Exported Jan 7, 2019 from SnapGene 4.1.9
            http://www.snapgene.com
FEATURES             Location/Qualifiers
     source          1..2183
                     /organism="PCR product"
                     /mol_type="other DNA"
     primer_bind     1..48
                     /label=Barcode5_cagc_N10_leader_fp
     primer_bind     1..20
                     /label=Barcode5_cagc_fp
     gap             21..30
                     /estimated_length=10
     5'UTR           31..100
                     /label=RV leader
                     /note="SPBN leader"
     CDS             101..1450
                     /codon_start=1
                     /locus_tag="N"
                     /label=N
                     /note="N"
                     /
translation="MDADKIVFKVNNQVVSLKPEIIVDQYEYKYPAIKDLKKPCITLGK

APDLNKAYKSVLSGMSAAKLNPDDVCSYLAAAMQFFEGTCPEDWTSYGIVIARKGDKIT

PGSLVEIKRTDVEGNWALTGGMELTRDPTVPEHASLVGLLLSLYRLSKISGQNTGNYKT

NIADRIEQIFETAPFVKIVEHHTLMTTHKMCANWSTIPNFRFLAGTYDMFFSRIEHLYS

AIRVGTVVTAYEDCSGLVSFTGFIKQINLTAREAILYFFHKNFEEEIRRMFEPGQETAV

PHSYFIHFRSLGLSGKSPYSSNAVGHVFNLIHFVGCYMGQVRSLNATVIAACAPHEMSV

LGGYLGEEFFGKGTFERRFFRDEKELQEYEAAELTKTDVALADDGTVNSDDEDYFSGET

RSPEAVYTRIMMNGGRLKRSHIRRYVSVSSNHQARPNSFAEFLNKTYSSDS"
     CDS             1460..1480
                     /codon_start=1
                     /locus_tag="TEV site"
```

```
                              /label=TEV site
                              /note="TEV site"
                              /translation="ENLYFQS"
     CDS                      1490..1609
                              /codon_start=1
                              /locus_tag="ECFP destabilized by fusion to
residues 422–461
                              of mouse ornithine decarboxylase, giving an in
vivo(1)"
                              /product="ECFP destabilized by fusion to residues
422–461
                              of mouse ornithine decarboxylase, giving an in
vivo
                              half–life of ~2 hours"
                              /label=ECFP destabilized by fusion to residues
422–461
                              /label=ECFP destabilized by fusion to residues
422–461...
                              /label=ECFP destabilized by fusion to residues
422–461 of
                              mouse ornithine decarboxylase, giving an in
vivo(1)
                              /note="destabilization domain"
                              /note="mammalian codon–optimized"
                              /
translation="SHGFPPEVEEQDDGTLPMSCAQESGMDRHPAACASARINV"
     CDS                      1703..2167
                              /codon_start=1
                              /locus_tag="P"
                              /label=P
                              /note="P"
                              /
translation="MSKIFVNPSAIRAGLADLEMAEETVDLINRNIEDNQAHLQGEPIE

VDNLPEDMGRLHLDDGKSPNHGEIAKVGEGKYREDFQMDEGEDPSFLFQSYLENVGVQI

VRQMRSGERFLKIWSQTVEEIISYVAVNFPNPPGKSSEDKSTQTTGRELKK"
     primer_bind    complement(2148..2183)
                              /label=Barcode3_P_rp
ORIGIN
        1 acacgcatga cacactcagc nnnnnnnnnn acgcttaaca accagatcaa
agaaaaaaca
       61 gacattgtca attgcaaagc aaaaatgtaa cacccctaca atggatgccg
acaagattgt
      121 attcaaagtc aataatcagg tggtctcttt gaagcctgag attatcgtgg
atcaatatga
      181 gtacaagtac cctgccatca aagatttgaa aaagccctgt ataaccctag
gaaaggctcc
      241 cgatttaaat aaagcataca agtcagtttt gtcaggcatg agcgccgcca
aacttaatcc
```

```
 301 tgacgatgta tgttcctatt tggcagcggc aatgcagttt tttgagggga
catgtccgga
 361 agactggacc agctatggaa ttgtgattgc acgaaaagga gataagatca
ccccaggttc
 421 tctggtggag ataaaacgta ctgatgtaga agggaattgg gctctgacag
gaggcatgga
 481 actgacaaga gaccccactg tccctgagca tgcgtcctta gtcggtcttc
tcttgagtct
 541 gtataggttg agcaaaatat ccgggcaaaa cactggtaac tataagacaa
acattgcaga
 601 caggatagag cagatttttg agacagcccc ttttgttaaa atcgtggaac
accatactct
 661 aatgacaact cacaaaatgt gtgctaattg gagtactata ccaaacttca
gatttttggc
 721 cggaacctat gacatgtttt tctcccggat tgagcatcta tattcagcaa
tcagagtggg
 781 cacagttgtc actgcttatg aagactgttc aggactggta tcatttactg
ggttcataaa
 841 acaaatcaat ctcaccgcta gagaggcaat actatatttc ttccacaaga
actttgagga
 901 agagataaga agaatgtttg agccagggca ggagacagct gttcctcact
cttatttcat
 961 ccacttccgt tcactaggct tgagtgggaa atctccttat tcatcaaatg
ctgttggtca
1021 cgtgttcaat ctcattcact ttgtaggatg ctatatgggt caagtcagat
ccctaaatgc
1081 aacggttatt gctgcatgtg ctcctcatga aatgtctgtt ctagggggct
atctgggaga
1141 ggaattcttc gggaaaggga catttgaaag aagattcttc agagatgaga
aagaacttca
1201 agaatacgag gcggctgaac tgacaaagac tgacgtagca ctggcagatg
atggaactgt
1261 caactctgac gacgaggact acttttcagg tgaaaccaga agtccggagg
ctgtttatac
1321 tcgaatcatg atgaatggag gtcgactaaa gagatctcac atacggagat
atgtctcagt
1381 cagttccaat catcaagccc gtccaaactc attcgccgag tttctaaaca
agacatattc
1441 gagtgactca ggttccggag agaacctcta cttccaatcg ggatccggta
gccatggctt
1501 cccgccggag gtggaggagc aggatgatgg cacgctgccc atgtcttgtg
cccaggagag
1561 cgggatggac cgtcaccctg cagcctgtgc ttctgctagg atcaatgtgt
aagaagttga
1621 ataacaaaat gccggaaatc tacggattgt gtatatccat catgaaaaaa
actaacaccc
1681 ctcctttcga accatcccaa acatgagcaa gatctttgtc aatcctagtg
ctattagagc
1741 cggtctggcc gatcttgaga tggctgaaga aactgttgat ctgatcaata
gaaatatcga
```

```
     1801 agacaatcag gctcatctcc aaggggaacc catagaggtg gacaatctcc
ctgaggatat
     1861 ggggcgactt cacctggatg atggaaaatc gcccaaccat ggtgagatag
ccaaggtggg
     1921 agaaggcaag tatcgagagg actttcagat ggatgaagga gaggatccta
gcttcctgtt
     1981 ccagtcatac ctggaaaatg ttggagtcca aatagtcaga caaatgaggt
caggagagag
     2041 atttctcaag atatggtcac agaccgtaga agagattata tcctatgtcg
cggtcaactt
     2101 tcccaaccct ccaggaaagt cttcagagga taaatcaacc cagactactg
gccgagagct
     2161 caagaagtac tagagtagca ctc
//
```

Supplementary File S4: SiR-FLPo amplicon reference sequence.
This Genbank-format file contains the expected (i.e., based on the published sequence: Addgene # 99609) sequence of amplicons obtained from SiR-FLPo for SMRT sequencing.

```
LOCUS           Exported                 2183 bp ds-DNA     linear   SYN
07-JAN-2019
DEFINITION  synthetic linear DNA.
ACCESSION   .
VERSION     .
KEYWORDS    .
SOURCE      synthetic DNA construct
  ORGANISM  synthetic DNA construct
REFERENCE   1  (bases 1 to 2183)
  AUTHORS   Trial User
  TITLE     Direct Submission
  JOURNAL   Exported Jan 7, 2019 from SnapGene 4.1.9
            http://www.snapgene.com
FEATURES             Location/Qualifiers
     source          1..2183
                     /organism="synthetic DNA construct"
                     /mol_type="other DNA"
     primer_bind     1..48
                     /label=Barcode9_cagc_N10_leader_fp
     primer_bind     1..20
                     /label=Barcode9_cagc_fp
     gap             21..30
                     /estimated_length=10
     5'UTR           31..100
                     /label=RV leader
                     /note="SPBN leader"
     misc_feature    89..97
                     /label=Transcription start signal (N)
     CDS             101..1450
                     /codon_start=1
                     /locus_tag="N"
                     /label=N
                     /note="N"
                     /
translation="MDADKIVFKVNNQVVSLKPEIIVDQYEYKYPAIKDLKKPCITLGK

APDLNKAYKSVLSGMSAAKLNPDDVCSYLAAAMQFFEGTCPEDWTSYGIVIARKGDKIT

PGSLVEIKRTDVEGNWALTGGMELTRDPTVPEHASLVGLLLSLYRLSKISGQNTGNYKT

NIADRIEQIFETAPFVKIVEHHTLMTTHKMCANWSTIPNFRFLAGTYDMFFSRIEHLYS

AIRVGTVVTAYEDCSGLVSFTGFIKQINLTAREAILYFFHKNFEEEIRRMFEPGQETAV

PHSYFIHFRSLGLSGKSPYSSNAVGHVFNLIHFVGCYMGQVRSLNATVIAACAPHEMSV

LGGYLGEEFFGKGTFERRFFRDEKELQEYEAAELTKTDVALADDGTVNSDDEDYFSGET

RSPEAVYTRIMMNGGRLKRSHIRRYVSVSSNHQARPNSFAEFLNKTYSSDS"
     CDS             1460..1480
```

```
                        /codon_start=1
                        /product="tobacco etch virus (TEV) protease
recognition and
                        cleavage site"
                        /label=TEV site
                        /translation="ENLYFQS"
     CDS                1490..1609
                        /codon_start=1
                        /locus_tag="ECFP destabilized by fusion to
residues 422–461
                        of mouse ornithine decarboxylase, giving an in
vivo(1)"
                        /product="ECFP destabilized by fusion to residues
422–461
                        of mouse ornithine decarboxylase, giving an in
vivo
                        half-life of ~2 hours"
                        /label=PEST
                        /label=ECFP destabilized by fusion to residues
422–461 of
                        mouse ornithine decarboxylase, giving an in
vivo(1)
                        /note="destabilization domain"
                        /note="mammalian codon-optimized"
                        /
translation="SHGFPPEVEEQDDGTLPMSCAQESGMDRHPAACASARINV"
     misc_feature       1661..1671
                        /label=Transcription stop/pA signal (N/P & P/M))
     misc_feature       1674..1682
                        /label=Transcription start signal (P)
     misc_feature       1703..2167
                        /locus_tag="P"
                        /label=P
                        /note="P"
     primer_bind        complement(2148..2183)
                        /label=Barcode4_P_rp
ORIGIN
        1 ctgcgtgctc tacgaccagc nnnnnnnnnn acgcttaaca accagatcaa
agaaaaaaca
       61 gacattgtca attgcaaagc aaaaatgtaa cacccctaca atggatgccg
acaagattgt
      121 attcaaagtc aataatcagg tggtctcttt gaagcctgag attatcgtgg
atcaatatga
      181 gtacaagtac cctgccatca agatttgaa aaagccctgt ataaccctag
gaaaggctcc
      241 cgatttaaat aaagcataca agtcagtttt gtcaggcatg agcgccgcca
aacttaatcc
      301 tgacgatgta tgttcctatt tggcagcggc aatgcagttt tttgagggga
catgtccgga
      361 agactggacc agctatggaa ttgtgattgc acgaaaagga gataagatca
```

```
        ccccaggttc
    421 tctggtggag ataaaacgta ctgatgtaga agggaattgg gctctgacag
        gaggcatgga
    481 actgacaaga gaccccactg tccctgagca tgcgtcctta gtcggtcttc
        tcttgagtct
    541 gtataggttg agcaaaatat ccgggcaaaa cactggtaac tataagacaa
        acattgcaga
    601 caggatagag cagattttg  agacagcccc ttttgttaaa atcgtggaac
        accatactct
    661 aatgacaact cacaaaatgt gtgctaattg gagtactata ccaaacttca
        gatttttggc
    721 cggaacctat gacatgtttt tctcccggat tgagcatcta tattcagcaa
        tcagagtggg
    781 cacagttgtc actgcttatg aagactgttc aggactggta tcatttactg
        ggttcataaa
    841 acaaatcaat ctcaccgcta gagaggcaat actatatttc ttccacaaga
        actttgagga
    901 agagataaga agaatgtttg agccagggca ggagacagct gttcctcact
        cttatttcat
    961 ccacttccgt tcactaggct tgagtgggaa atctccttat tcatcaaatg
        ctgttggtca
   1021 cgtgttcaat ctcattcact ttgtaggatg ctatatgggt caagtcagat
        ccctaaatgc
   1081 aacggttatt gctgcatgtg ctcctcatga aatgtctgtt ctagggggct
        atctgggaga
   1141 ggaattcttc gggaaaggga catttgaaag aagattcttc agagatgaga
        aagaacttca
   1201 agaatacgag gcggctgaac tgacaaagac tgacgtagca ctggcagatg
        atggaactgt
   1261 caactctgac gacgaggact acttttcagg tgaaaccaga agtccggagg
        ctgtttatac
   1321 tcgaatcatg atgaatggag gtcgactaaa gagatctcac atacggagat
        atgtctcagt
   1381 cagttccaat catcaagccc gtccaaactc attcgccgag tttctaaaca
        agacatattc
   1441 gagtgactca ggttccggag agaacctcta cttccaatcg ggatccggta
        gccatggctt
   1501 cccgccggag gtggaggagc aggatgatgg cacgctgccc atgtcttgtg
        cccaggagag
   1561 cgggatggac cgtcaccctg cagcctgtgc ttctgctagg atcaatgtgt
        aagaagttga
   1621 ataacaaaat gccggaaatc tacggattgt gtatatccat catgaaaaaa
        actaacaccc
   1681 ctcctttcga accatcccaa acatgagcaa gatctttgtc aatcctagtg
        ctattagagc
   1741 cggtctggcc gatcttgaga tggctgaaga aactgttgat ctgatcaata
        gaaatatcga
   1801 agacaatcag gctcatctcc aaggggaacc catagaggtg gacaatctcc
        ctgaggatat
   1861 ggggcgactt cacctggatg atggaaaatc gcccaaccat ggtgagatag
```

```
ccaaggtggg
     1921 agaaggcaag tatcgagagg actttcagat ggatgaagga gaggatccta
gcttcctgtt
     1981 ccagtcatac ctggaaaatg ttggagtcca aatagtcaga caaatgaggt
caggagagag
     2041 atttctcaag atatggtcac agaccgtaga agagattata tcctatgtcg
cggtcaactt
     2101 tcccaaccct ccaggaaagt cttcagagga taaatcaacc cagactactg
gccgagagct
     2161 caagaagtgt gtatcagtac atg
//
```

Supplementary File S5: RV∆G-4Cre amplicon reference sequence.
This Genbank-format file contains the expected (i.e., based on the published sequence: see Addgene #98034) sequence of amplicons obtained from RV∆G-4Cre for SMRT sequencing.

```
LOCUS       Exported                2024 bp ds-DNA     linear   SYN
07-JAN-2019
DEFINITION  synthetic linear DNA.
ACCESSION   .
VERSION     .
KEYWORDS    .
SOURCE      synthetic DNA construct
  ORGANISM  synthetic DNA construct
REFERENCE   1  (bases 1 to 2024)
  AUTHORS   Trial User
  TITLE     Direct Submission
  JOURNAL   Exported Jan 7, 2019 from SnapGene 4.1.9
            http://www.snapgene.com
FEATURES             Location/Qualifiers
     source          1..2024
                     /organism="synthetic DNA construct"
                     /mol_type="other DNA"
     primer_bind     1..48
                     /label=Barcode1_cagc_N10_leader_fp
     primer_bind     1..20
                     /label=Barcode1_cagc_fp
     gap             21..30
                     /estimated_length=10
     5'UTR           31..100
                     /label=RV leader
                     /note="SPBN leader"
     misc_feature    89..97
                     /locus_tag="Transcriptional start (N)"
                     /label=Transcriptional start (N)
     CDS             101..1450
                     /codon_start=1
                     /locus_tag="N"
                     /label=N
                     /note="N"
                     /
translation="MDADKIVFKVNNQVVSLKPEIIVDQYEYKYPAIKDLKKPCITLGK

APDLNKAYKSVLSGMSAAKLNPDDVCSYLAAAMQFFEGTCPEDWTSYGIVIARKGDKIT

PGSLVEIKRTDVEGNWALTGGMELTRDPTVPEHASLVGLLLSLYRLSKISGQNTGNYKT

NIADRIEQIFETAPFVKIVEHHTLMTTHKMCANWSTIPNFRFLAGTYDMFFSRIEHLYS

AIRVGTVVTAYEDCSGLVSFTGFIKQINLTAREAILYFFHKNFEEEIRRMFEPGQETAV

PHSYFIHFRSLGLSGKSPYSSNAVGHVFNLIHFVGCYMGQVRSLNATVIAACAPHEMSV

LGGYLGEEFFGKGTFERRFFRDEKELQEYEAAELTKTDVALADDGTVNSDDEDYFSGET

RSPEAVYTRIMMNGGRLKRSHIRRYVSVSSNHQARPNSFAEFLNKTYSSDS"
```

```
     misc_feature     1502..1512
                      /label=Transcription stop/pA signal (N/P & P/M))
     misc_feature     1515..1523
                      /locus_tag="Transcriptional start (P)"
                      /label=Transcriptional start (P)
     misc_feature     1544..2008
                      /locus_tag="P"
                      /label=P
                      /note="P"
     primer_bind      complement(1989..2024)
                      /label=Barcode2_P_rp
ORIGIN
        1 tcagacgatg cgtcatcagc nnnnnnnnnn acgcttaaca accagatcaa
agaaaaaaca
       61 gacattgtca attgcaaagc aaaaatgtaa caccccctaca atggatgccg
acaagattgt
      121 attcaaagtc aataatcagg tggtctcttt gaagcctgag attatcgtgg
atcaatatga
      181 gtacaagtac cctgccatca aagatttgaa aaagccctgt ataaccctag
gaaaggctcc
      241 cgatttaaat aaagcataca agtcagtttt gtcaggcatg agcgccgcca
aacttaatcc
      301 tgacgatgta tgttcctatt tggcagcggc aatgcagttt tttgagggga
catgtccgga
      361 agactggacc agctatggaa ttgtgattgc acgaaaagga gataagatca
ccccaggttc
      421 tctggtggag ataaaacgta ctgatgtaga agggaattgg gctctgacag
gaggcatgga
      481 actgacaaga gaccccactg tccctgagca tgcgtcctta gtcggtcttc
tcttgagtct
      541 gtataggttg agcaaaatat ccgggcaaaa cactggtaac tataagacaa
acattgcaga
      601 caggatagag cagattttg agacagcccc ttttgttaaa atcgtggaac
accatactct
      661 aatgacaact cacaaaatgt gtgctaattg gagtactata ccaaacttca
gatttttggc
      721 cggaacctat gacatgtttt tctcccggat tgagcatcta tattcagcaa
tcagagtggg
      781 cacagttgtc actgcttatg aagactgttc aggactggta tcatttactg
ggttcataaa
      841 acaaatcaat ctcaccgcta gagaggcaat actatatttc ttccacaaga
actttgagga
      901 agagataaga agaatgtttg agccagggca ggagacagct gttcctcact
cttatttcat
      961 ccacttccgt tcactaggct tgagtgggaa atctccttat tcatcaaatg
ctgttggtca
     1021 cgtgttcaat ctcattcact ttgtaggatg ctatatgggt caagtcagat
ccctaaatgc
     1081 aacggttatt gctgcatgtg ctcctcatga aatgtctgtt ctaggggct
atctgggaga
```

```
1141 ggaattcttc gggaaaggga catttgaaag aagattcttc agagatgaga
aagaacttca
1201 agaatacgag gcggctgaac tgacaaagac tgacgtagca ctggcagatg
atggaactgt
1261 caactctgac gacgaggact acttttcagg tgaaaccaga agtccggagg
ctgtttatac
1321 tcgaatcatg atgaatggag gtcgactaaa gagatctcac atacggagat
atgtctcagt
1381 cagttccaat catcaagccc gtccaaactc attcgccgag tttctaaaca
agacatattc
1441 gagtgactca taagaagttg aataacaaaa tgccggaaat ctacggattg
tgtatatcca
1501 tcatgaaaaa aactaacacc cctcctttcg aaccatccca aacatgagca
agatctttgt
1561 caatcctagt gctattagag ccggtctggc cgatcttgag atggctgaag
aaactgttga
1621 tctgatcaat agaaatatcg aagacaatca ggctcatctc caaggggaac
ccatagaggt
1681 ggacaatctc cctgaggata tggggcgact tcacctggat gatggaaaat
cgcccaacca
1741 tggtgagata gccaaggtgg gagaaggcaa gtatcgagag gactttcaga
tggatgaagg
1801 agaggatcct agcttcctgt tccagtcata cctggaaaat gttggagtcc
aaatagtcag
1861 acaaatgagg tcaggagaga gatttctcaa gatatggtca cagaccgtag
aagagattat
1921 atcctatgtc gcggtcaact ttcccaaccc tccaggaaag tcttcagagg
ataaatcaac
1981 ccagactact ggccgagagc tcaagaagct atacatgact ctgc
//
```

Supplementary Figure S1: SiR viruses appeared to cause expression of viral nucleoprotein at levels similar to those of first-generation ΔG viruses.

(A-D) Reporter cells infected with first-generation, ΔG viruses show characteristic bright, clumpy anti-nucleoprotein staining (green), indicating high nucleoprotein expression and active viral replication. Red is mCherry expression, reporting expression of Cre or FLPo; blue is mTagBFP2, constitutively expressed by these reporter cell lines.

(E-H) Reporter cells infected with second-generation, ΔGL viruses show only punctate staining for nucleoprotein, indicating isolated individual viral particles or ribonucleoprotein complexes; these viruses do not replicate intracellularly (21). Reporter cassette activation takes longer from the lower recombinase expression levels of these viruses, so mCherry expression is dimmer than in cells infected with ΔG viruses at the same time point.

(I-J) Reporter cells infected with SiR viruses show clumps of nucleoprotein and rapid reporter expression indicating high expression of recombinases, similarly to cells infected with ΔG viruses. Scale bar: 100 μm, applies to all panels.

First-generation (△G) **Second-generation (△GL)** **SiR**

Cre

A RVΔG-4Cre(B19G)   C RVΔG-4Cre(EnvA)   E RVΔGL-4Cre(B19G)   G RVΔGL-4Cre(EnvA)   I SiR-CRE(EnvA)

FLPo

B RVΔG-4FLPo(B19G)   D RVΔG-4FLPo(EnvA)   F RVΔGL-4FLPo(B19G)   H RVΔGL-4FLPo(EnvA)   J SiR-FLPo(EnvA)

Supplementary File S6: >90% of SiR-CRE viral particles with the mCherry gene intact also have the Cre gene intact, suggesting that most of the SiR-CRE-infected cells that disappear over time in tdTomato reporter mice are dying rather than simply stopping expression of mCherry.

We sequenced the transgene inserts for 21 individual SiR-CRE clones (see Methods). 19 out of 21 had no mutations in the Cre gene, and two had one point mutation each (Ala88Val and Arg189Ile). All 21 had an intact mCherry gene. The lack of a large proportion of Cre-knockout mutants is one indication that the majority of red fluorescent neurons in SiR-CRE-injected Ai14 (tdTomato reporter) mice are not labeled only with mCherry, providing evidence that their disappearance is equivalent to their death.

Random 8-nucleotide index
Cre gene, P2A, mCherry gene, PEST, and transcription stop/polyA signal, transcription start signal, L gene, mutation

No mutations in transgene insert of SiR-CRE (19/21 clones)

ACTTTCTA, AGCCCAGT, TTATCAGG, AGGGCGTT, AGTGGGCA, GGGCCTTT, TCCCGCAA, GGTCGGGG, GGCCTATA, TGTCTCTC, CCTGTCGG, TAGCACAC, GTATGTGG, TAAGGCCT, CTCACTAT, TTTAGTCC, AACCCGAC, AGCGGCGC, AGCGGGGG
CAACTCCAACCCTTGGGAGCAATATAACAAAAAACATGTTATGGTGCCATTAAACCGCTGCATTTCATCAAAGTCA AGTTGATTACCTTTACATTTTGATCCTCTTGGATGTGAAAAAAACTATTAACATCCCTCAAAGGACCTGCAGGTACG CGGCCGCGGTACCGCCACCATGGTGCCCAAGAAGAAGAGGAAAGTCTCCAACCTGCTGACTGTGCACCAAAAC CTGCCTGCCCTCCCTGTGGATGCCACCTCTGATGAAGTCAGGAAGAACCTGATGGACATGTTCAGGGACAGGCA GGCCTTCTCTGAACACACCTGGAAGATGCTCCTGTCTGTGTGCAGATCCTGGGCTGCCTGGTGCAAGCTGAACA ACAGGAAATGGTTCCCTGCTGAACCTGAGGATGTGAGGGACTACCTCCTGTACCTGCAAGCCAGAGGCCTGGC TGTGAAGACCATCCAACAGCACCTGGGCCAGCTCAACATGCTGCACAGGAGATCTGGCCTGCCTCGCCCTTCT GACTCCAATGCTGTGTCCCTGGTGATGAGGAGAATCAGAAAGGAGAATGTGGATGCTGGGGAGAGAGCCAAGC AGGCCCTGGCCTTTGAACGCACTGACTTTGACCAAGTCAGATCCCTGATGGAGAACTCTGACAGATGCCAGGAC ATCAGGAACCTGGCCTTCCTGGGCATTGCCTACAACACCCTGCTGCGCATTGCCGAAATTGCCAGAATCAGAGT GAAGGACATCTCCCGCACCGATGGTGGGAGAATGCTGATCCACATTGGCAGGACCAAGACCCTGGTGTCCACA GCTGGTGTGGAGAAGGCCCTGTCCCTGGGGGTTACCAAGCTGGTGGAGAGATGGATCTCTGTGTCTGGTGTGG CTGATGACCCCAACAACTACCTGTTCTGCCGGGTCAGAAAGAATGGTGTGGCTGCCCCTTCTGCCACCTCCCAA CTGTCCACCCGGGCCCTGGAAGGGATCTTTGAGGCCACCCACCGCCTGATCTATGGTGCCAAGGATGACTCTG GGCAGAGATACCTGGCCTGGTCTGGCCACTCTGCCAGAGTGGGTGCTGCCAGGGACATGGCCAGGGCTGGTG TGTCCATCCCTGAAATCATGCAGGCTGGTGGCTGGACCAATGTGAACATTGTGATGAACTACATCAGAAACCTGG ACTCTGAGACTGGGGCCATGGTGAGGCTGCTCGAGGATGGGGACGGCAGTGGAGGATCCGGAGCCACGAACT TCTCTCTGTTAAAGCAAGCAGGAGACGTGGAAGAAAACCCCGGTCCTACCGGTGTGAGCAAGGGCGAGGAGGA TAACATGGCCATCATCAAGGAGTTCATGCGCTTCAAGGTGCACATGGAGGGCTCCGTGAACGGCCACGAGTTCG AGATCGAGGGCGAGGGCGAGGGCCGCCCCTACGAGGGCACCCAGACCGCCAAGCTGAAGGTGACCAAGGGT GGCCCCCTGCCCTTCGCCTGGGACATCCTGTCCCCTCAGTTCATGTACGGCTCCAAGGCCTACGTGAAGCACC CGGCCGACATCCCCGACTACTTGAAGCTGTCCTTCCCCGAGGGCTTCAAGTGGGAGCGCGTGATGAACTTCGA GGACGGCGGCGTGGTGACCGTGACCCAGGACTCCTCCCTGCAGGACGGCGAGTTCATCTACAAGGTGAAGCT GCGCGGCACCAACTTCCCCTCCGACGGCCCCGTAATGCAGAAGAAGACCATGGGCTGGGAGGCCTCCTCCGA GCGGATGTACCCCGAGGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCTGAAGGACGGCGGCCA CTACGACGCTGAGGTCAAGACCACCTACAAGGCCAAGAAGCCCGTGCAGCTGCCCCGGCGCCTACAACGTCAAC ATCAAGTTGGACATCACCTCCCACAACGAGGACTACACCATCGTGGAACAGTACGAACGCGCCGAGGGCCGCC ACTCCACCGGCGGCATGGACGAGCTGTACAAGGGATATCTCAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCA GGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCT GCTAGGATCAATGTGTGACTCGAGGGCGCGCCTACCCGCGGTAGCTTTTCAGTCGAGAAAAAAACATTAGATCA GAAGAACAACTGGCAACACTTCTCAACCTGAGACTTACTTCAAGATGCTCGATCCTGGAGAGGTCTATGATGACC CTATTGACCCAATCGAGTTAGAGGCTGAACCCAGAGGAACCCCCATTGTCCCCAAC

Ala88Val mutation in iCre gene of SiR-CRE (1/21 clones)

AGCTGGTT
CAACTCCAACCCTTGGGAGCAATATAACAAAAAACATGTTATGGTGCCATTAAACCGCTGCATTTCATCAAAGTCA AGTTGATTACCTTTACATTTTGATCCTCTTGGATGTGAAAAAAACTATTAACATCCCTCAAAGGACCTGCAGGTACG CGGCCGCGGTACCGCCACCATGGTGCCCAAGAAGAAGAGGAAAGTCTCCAACCTGCTGACTGTGCACCAAAAC CTGCCTGCCCTCCCTGTGGATGCCACCTCTGATGAAGTCAGGAAGAACCTGATGGACATGTTCAGGGACAGGCA GGCCTTCTCTGAACACACCTGGAAGATGCTCCTGTCTGTGTGCAGATCCTGGGCTGCCTGGTGCAAGCTGAACA ACAGGAAATGGTTCCCTGCTGAACCTGAGGATGTGAGGGACTACCTCCTGTACCTGCAAGTCAGAGGCCTGGCT GTGAAGACCATCCAACAGCACCTGGGCCAGCTCAACATGCTGCACAGGAGATCTGGCCTGCCTCGCCCTTCTG ACTCCAATGCTGTGTCCCTGGTGATGAGGAGAATCAGAAAGGAGAATGTGGATGCTGGGGAGAGAGCCAAGCA GGCCCTGGCCTTTGAACGCACTGACTTTGACCAAGTCAGATCCCTGATGGAGAACTCTGACAGATGCCAGGACA TCAGGAACCTGGCCTTCCTGGGCATTGCCTACAACACCCTGCTGCGCATTGCCGAAATTGCCAGAATCAGAGTG AAGGACATCTCCCGCACCGATGGTGGGAGAATGCTGATCCACATTGGCAGGACCAAGACCCTGGTGTCCACAG CTGGTGTGGAGAAGGCCCTGTCCCTGGGGGTTACCAAGCTGGTGGAGAGATGGATCTCTGTGTCTGGTGGC TGATGACCCCAACAACTACCTGTTCTGCCGGGTCAGAAAGAATGGTGTGGCTGCCCCTTCTGCCACCTCCCAAC TGTCCACCCGGGCCCTGGAAGGGATCTTTGAGGCCACCCACCGCCTGATCTATGGTGCCAAGGATGACTCTGG GCAGAGATACCTGGCCTGGTCTGGCCACTCTGCCAGAGTGGGTGCTGCCAGGGACATGGCCAGGGCTGGTGT GTCCATCCCTGAAATCATGCAGGCTGGTGGCTGGACCAATGTGAACATTGTGATGAACTACATCAGAAACCTGGA

```
CTCTGAGACTGGGGCCATGGTGAGGCTGCTCGAGGATGGGGACGGCAGTGGAGGATCCGGAGCCACGAACTT
CTCTCTGTTAAAGCAAGCAGGAGACGTGGAAGAAAACCCCGGTCCTACCGGTGTGAGCAAGGGCGAGGAGGAT
AACATGGCCATCATCAAGGAGTTCATGCGCTTCAAGGTGCACATGGAGGGCTCCGTGAACGGCCACGAGTTCGA
GATCGAGGGCGAGGGCGAGGGCCGCCCCTACGAGGGCACCCAGACCGCCAAGCTGAAGGTGACCAAGGGGTG
GCCCCCTGCCCTTCGCCTGGGACATCCTGTCCCCTCAGTTCATGTACGGCTCCAAGGCCTACGTGAAGCACCC
CGCCGACATCCCCGACTACTTGAAGCTGTCCTTCCCCGAGGGCTTCAAGTGGGAGCGCGTGATGAACTTCGAG
GACGGCGGCGTGGTGACCGTGACCCAGGACTCCTCCCTGCAGGACGGCGAGTTCATCTACAAGGTGAAGCTG
CGCGGCACCAACTTCCCCTCCGACGGCCCCGTAATGCAGAAGAAGACCATGGGCTGGGAGGCCTCCTCCGAG
CGGATGTACCCCGAGGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCTGAAGGACGGCGGCCAC
TACGACGCTGAGGTCAAGACCACCTACAAGGCCAAGAAGCCCGTGCAGCTGCCCGGCGCCTACAACGTCAACA
TCAAGTTGGACATCACCTCCCACAACGAGGACTACACCATCGTGGAACAGTACGAACGCGCCGAGGGCCGCCA
CTCCACCGGCGGCATGGACGAGCTGTACAAGGGATATCTCAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAG
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTG
CTAGGATCAATGTGTGACTCGAGGGCGCGCCTACCCGCGGTAGCTTTTCAGTCGAGAAAAAAACATTAGATCAGA
AGAACAACTGGCAACACTTCTCAACCTGAGACTTACTTCAAGATGCTCGATCCTGGAGAGGTCTATGATGACCCT
ATTGACCCAATCGAGTTAGAGGCTGAACCCAGAGGAACCCCCATTGTCCCCAAC
```

Arg189Ile mutation in iCre gene of SiR-CRE

```
AGCGGGGT
```

```
CAACTCCAACCCTTGGGAGCAATATAACAAAAAACATGTTATGGTGCCATTAAACCGCTGCATTTCATCAAAGTCA
AGTTGATTACCTTTACATTTTGATCCTCTTGGATGTGAAAAAAACTATTAACATCCCTCAAAGGACCTGCAGGTACG
CGGCCGCGGTACCGCCACCATGGTGCCCAAGAAGAAGAGGAAAGTCTCCAACCTGCTGACTGTGCACCAAAAC
CTGCCTGCCCTCCCTGTGGATGCCACCTCTGATGAAGTCAGGAAGAACCTGATGGACATGTTCAGGGACAGGCA
GGCCTTCTCTGAACACACCTGGAAGATGCTCCTGTCTGTGTGCAGATCCTGGGCTGCCTGGTGCAAGCTGAACA
ACAGGAAATGGTTCCCTGCTGAACCTGAGGATGTGAGGGACTACCTCCTGTACCTGCAAGCCAGAGGCCTGGC
TGTGAAGACCATCCAACAGCACCTGGGCCAGCTCAACATGCTGCACAGGAGATCTGGCCTGCCTCGCCCTTCT
GACTCCAATGCTGTGTCCCTGGTGATGAGGAGAATCAGAAAGGAGAATGTGGATGCTGGGGAGAGAGCCAAGC
AGGCCCTGGCCTTTGAACGCACTGACTTTGACCAAGTCAGATCCCTGATGGAGAACTCTGACAGATGCCAGGAC
ATCAGGAACCTGGCCTTCCTGGGCATTGCCTACAACACCCTGCTGCGCATTGCCGAAATTGCCAGAATCATAGTG
AAGGACATCTCCCGCACCGATGGTGGGAGAATGCTGATCCACATTGGCAGGACCAAGACCCTGGTGTCCACAG
CTGGTGTGGAGAAGGCCCTGTCCCTGGGGGTTACCAAGCTGGTGGAGATGGATCTCTGTGTCTGGTGTGGC
TGATGACCCCAACAACTACCTGTTCTGCCGGGTCAGAAAGAATGGTGTGGCTGCCCCTTCTGCCACCTCCCAAC
TGTCCACCCGGGCCCTGGAAGGGATCTTTGAGGCCACCCACCGCCTGATCTATGGTGCCAAGGATGACTCTGG
GCAGAGATACCTGGCCTGGTCTGGCCACTCTGCCAGAGTGGGTGCTGCCAGGGACATGGCCAGGGCTGGTGT
GTCCATCCCTGAAATCATGCAGGCTGGTGGCTGGACCAATGTGAACATTGTGATGAACTACATCAGAAACCTGGA
CTCTGAGACTGGGGCCATGGTGAGGCTGCTCGAGGATGGGGACGGCAGTGGAGGATCCGGAGCCACGAACTT
CTCTCTGTTAAAGCAAGCAGGAGACGTGGAAGAAAACCCCGGTCCTACCGGTGTGAGCAAGGGCGAGGAGGAT
AACATGGCCATCATCAAGGAGTTCATGCGCTTCAAGGTGCACATGGAGGGCTCCGTGAACGGCCACGAGTTCGA
GATCGAGGGCGAGGGCGAGGGCCGCCCCTACGAGGGCACCCAGACCGCCAAGCTGAAGGTGACCAAGGGGTG
GCCCCCTGCCCTTCGCCTGGGACATCCTGTCCCCTCAGTTCATGTACGGCTCCAAGGCCTACGTGAAGCACCC
CGCCGACATCCCCGACTACTTGAAGCTGTCCTTCCCCGAGGGCTTCAAGTGGGAGCGCGTGATGAACTTCGAG
GACGGCGGCGTGGTGACCGTGACCCAGGACTCCTCCCTGCAGGACGGCGAGTTCATCTACAAGGTGAAGCTG
CGCGGCACCAACTTCCCCTCCGACGGCCCCGTAATGCAGAAGAAGACCATGGGCTGGGAGGCCTCCTCCGAG
CGGATGTACCCCGAGGACGGCGCCCTGAAGGGCGAGATCAAGCAGAGGCTGAAGCTGAAGGACGGCGGCCAC
TACGACGCTGAGGTCAAGACCACCTACAAGGCCAAGAAGCCCGTGCAGCTGCCCGGCGCCTACAACGTCAACA
TCAAGTTGGACATCACCTCCCACAACGAGGACTACACCATCGTGGAACAGTACGAACGCGCCGAGGGCCGCCA
CTCCACCGGCGGCATGGACGAGCTGTACAAGGGATATCTCAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAG
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTG
CTAGGATCAATGTGTGACTCGAGGGCGCGCCTACCCGCGGTAGCTTTTCAGTCGAGAAAAAAACATTAGATCAGA
AGAACAACTGGCAACACTTCTCAACCTGAGACTTACTTCAAGATGCTCGATCCTGGAGAGGTCTATGATGACCCT
ATTGACCCAATCGAGTTAGAGGCTGAACCCAGAGGAACCCCCATTGTCCCCAAC
```

Supplementary Figure S2: 86% of SiR-CRE-labeled neurons in Arch-EGFP-ER2 reporter mice disappeared between 11 days and 28 days after injection.

A) Maximum intensity projections of the two-photon FOV shown in Supplementary Video S1 of visual cortical neurons labeled with SiR-CRE in an Ai35 mouse, 11-28 days postinjection. Images are from the same FOV at four different time points. All cells clearly visible on day 11 are circled. In this example, 18 out of 19 cells (red circles) disappeared by a subsequent imaging session. Only one cell (white circle) is still visible on day 28. Numbers below four of the circles mark the cells for which intensity profiles are shown in panel B. Scale bar: 50 μm, applies to all images. See Supplementary Video S1 for a clearer view of the same cells.

B) Green fluorescence intensity versus depth for the four representative neurons numbered in panel A at the four different time points, showing disappearance of three of them over time.

C) Fraction of cells visibly EGFP-labeled at day 11 still visible at later time points, from four different FOVs in two Ai35 mice. Connected sets of markers indicate cells from the same FOV. 86% of SiR-CRE-labeled neurons had disappeared by 4 weeks postinjection.

Supplementary Video S1 *(separate file)*: **Video of 95% of SiR-CRE-labeled neurons in an Arch-EGFP-ER2 reporter mouse disappearing between 11 days and 28 days postinjection.** Two-photon image stacks of a single FOV of visual cortical neurons in an Ai35 mouse imaged at four different time points; time in the video represents depth of focus. Large blobs are glia. 18 out of the 19 neurons visibly labeled with Arch-EGFP-ER2 at 11 days following injection of SiR-Cre are no longer visible 17 days later. White circles indicate cells present at both 11 days and all subsequent imaging sessions; red circles indicate cells present at 11 days but gone by 28 days.

Supplementary Figure S3: mCherry fluorescence from SiR-CRE is much dimmer than tdTomato fluorescence in Ai14 mice, suggesting that the disappearance of the brighter cells in SiR-CRE-injected Ai14 mice indicates their death.

A) Representative images of red fluorescence in SiR-CRE-labeled cells in Ai14 (Cre-dependent expression of tdTomato, top row) and Ai35 (Cre-dependent expression of Arch-EGFP-ER2, bottom row). The three images for each mouse line are from 3 different mice of each line, imaged 7 days following SiR-CRE injection (see Methods), all with the same laser intensity and wavelength (1020 nm). Red fluorescence due only to mCherry (i.e., in Ai35 mice) is obviously much dimmer than that due to tdTomato (i.e., in Ai14 mice). Scale bar: 50 µm, applies to all images.

B) Intensity of red fluorescence of SiR-CRE-labeled cells in Ai14 (left) and Ai35 (right) mice. Data point indicate intensity of individual cells in arbitrary units at the same laser and microscope settings (see Methods). Box plots indicate median, 25th–75th percentiles (boxes), and full range (whiskers) of intensities for each mouse. The average of the mean red fluorescent intensity in each mouse was 48.97 in Ai14 and 15.69 in Ai35 ($p=0.00283 < 0.01$, one-way ANOVA); the midpoint of these means, 32.33, was used as the cutoff for the reanalysis of the data in Ai14 mice to exclude neurons that could have been labeled with mCherry alone.

Supplementary Figure S4: 81% of SiR-CRE-labeled neurons in tdTomato reporter mice disappear within 2-4 weeks, even excluding dimmer cells that might have only been labeled with mCherry.

A) Same representative fields of view as in Figure 3 but with circles now marking only cells with intensity at 7 days of greater than 32.33 a.u. (see text and Supplementary Figure S3) that are no longer visible at a subsequent time point. Scale bar: 50 µm, applies to all images.

B-D) Numbers of cells above threshold fluorescence intensity at week 1 that were still present in subsequent weeks. The conclusions from Figure 3 are unchanged: few cells labeled with RV∆GL-4Cre were lost, RV∆G-4Cre killed a significant minority of cells, and SiR-CRE killed the majority of labeled neurons within two weeks following injection.

E) Percentages of cells above threshold at week 1 that were still present in subsequent imaging sessions. By 28 days postinjection, an average of only 19.2% of suprathreshold SiR-CRE-labeled cells remained.

Supplementary File S7: Sequencing results for RV∆G-NPEST-4Cre(EnvA) ("intact") and RV∆G-NPEST-4Cre(EnvA) ("revertant"), showing that the 3' additions to the nucleoprotein were as intended in the final stocks used for monosynaptic tracing experiments.

We sequenced the complete region of the 3' addition to the nucleoprotein region from 32 viral particles for each of the two high-titer, EnvA-enveloped preparations of RV∆G-NPEST-4Cre(EnvA) and RV∆G-N*PEST-4Cre(EnvA) that we made for our *in vivo* experiments. For RV∆G-NPEST-4Cre(EnvA), none of the 32 clones had any mutations in the 3' addition; one clone had a point mutation in the transcriptional termination signal of the nucleoprotein gene (from CATGAAAAAAA to CATAAAAAAAA). For RV∆G-N*PEST-4Cre(EnvA), 30 of the 32 clones had no mutations in the sequenced region; one clone had a synonymous mutation in the N gene (Asn436, from AAC to AAT), and one other clone had a synonymous mutation in the PEST region (Pro5, from CCG to CCA) after the stop codon.

Sample number	Mutation	UMI sequence
NPEST-1	None	ATGAGGGATCTTTAGTAGTT
NPEST-2	None	GTTCCCAGTGGGCAGCATTT
NPEST-3	None	GGACGACTACGCTGGACTTG
NPEST-4	None	CTGGAACTCCTAGTTTGCTT
NPEST-5	None	CGTCGTGATCACTAGAGTCA
NPEST-6	None	TAAGGGTGGTTTTTTTTTA
NPEST-7	None	TTCAGCGTTTCGAACCCGAA
NPEST-8	None	GTTATAGATGATGTTATGTA
NPEST-9	None	ATGAGTGCGTTCTAAATTGT
NPEST-10	None	GGGGAGATACGGCAACGTCG
NPEST-11	None	ATTTCAGAATGTCGTGGTTA
NPEST-12	None	TTCCAACTAAATAAGGTACCA
NPEST-13	None	GTTGGGTTATTTTAAACATT
NPEST-14	None	TTTCTTGATGCAGGTTTGCC
NPEST-15	None	GAGAATCAAGACATATAGCC
NPEST-16	None	TGAGGACGGACTCTGATCCT
NPEST-17	None	GGTCTTCTTCTTCTTATTCT
NPEST-18	In the N transcriptional terminator from CAT**G**AAAAAAA to CAT**A**AAAAAAA	TAGCAACCGTTTTTAGTACG
NPEST-19	None	GTTTTTCGTCATTATGTTGC
NPEST-20	None	TCACTCGGGTAGATACATTG
NPEST-21	None	TTCTTGCAAAGTATTGAACC
NPEST-22	None	GTTCTGGGGGGTTCTTATTA
NPEST-23	None	GATTAACTCTAGCTTGCCCC
NPEST-24	None	TATGACTCTTCTGTTTCATG
NPEST-25	None	CTCTCTGTTCTCTTGGCAAT
NPEST-26	None	TTACCTTATTGGGTTTGGTT
NPEST-27	None	CAAAGATGGGATCACAGTAG
NPEST-28	None	TAGGTTCCTCCTGTGGTATC
NPEST-29	None	CCAACGATACAGTTGTGGGG
NPEST-30	None	CGATATGATATAAGAATCTT
NPEST-31	None	TCATAATGCGAGGTATGTCC
NPEST-32	None	TGCCTAGGTGTCGGGTTTGT

Sanger sequencing results (594nt from viral genome)

```
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGTTCCGGCGAGA
```

```
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
GATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCAATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAA(
```

CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG
CTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGATCTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGG

```
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
AACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
```

Sample number	Mutation	UMI sequence
N*PEST-1	None	TTTTCCATTTGTCCCAAATC
N*PEST-2	None	TAACCTCTTATATATTTGCC
N*PEST-3	None	GAAGGTACCTTACATAGTGT
N*PEST-4	Synonymous mutation in the N gene position Asn436 from AA**C** to AA**T**	TCAATTTCTTGTAATTTACT
N*PEST-5	None	GTAATTCATATAAAATATCG
N*PEST-6	None	CTTGTTCGGCCACGCAAGGG
N*PEST-7	None	TATATTGTACTTATCCCTCA
N*PEST-8	None	AGAAACTGCGCAGTCCGGCC
N*PEST-9	None	TCATCGTTTTGCCGGTAATT
N*PEST-10	None	TCCTCACTGCTTGTTTTCTG
N*PEST-11	None	CGACTTTAATATGTTTAAGG
N*PEST-12	None	TTACATCGTTATAAATATCG
N*PEST-13	None	CACCATAACGATTTTTTTG
N*PEST-14	None	CCTTTAACTATGCGCCACTC
N*PEST-15	None	AATTTCGTAGAAATCATATT
N*PEST-16	None	GTTTCATAGCTCGGGAGTTT
N*PEST-17	Synonymous mutation in the PEST region position Pro5 from CC**G** to CC**A**	GTTCCTCAAGTTATTATTAA
N*PEST-18	None	CCCGATTGGTCTGATGGTCA
N*PEST-19	None	CAACAGTAAGGAAAGTTGGC
N*PEST-20	None	AGTAGGCATTTCTTGTTATT
N*PEST-21	None	TTATTCGTTCTCGGTAAGAG
N*PEST-22	None	TTGAACATTAAACTCGCACC
N*PEST-23	None	GTATGAAGTCGTGTAAGTTT
N*PEST-24	None	ATAATGTGATGTAGATAACT
N*PEST-25	None	TACCATAAGATTTTTATTTA
N*PEST-26	None	TCTTAGTACGATTTCGATCT
N*PEST-27	None	ATATTCATTGCAGATTCGAA
N*PEST-28	None	AGTGAGGCTTTTACATTCTC
N*PEST-29	None	GTATTATGACTTACTTAAGT
N*PEST-30	None	GCATGATCGTGAGAAGTAAC
N*PEST-31	None	CTGAACCAGGTCAGTCAACC
N*PEST-32	None	AATAAGTAAATTTAACGTTT

Sanger sequencing results (594nt from viral genome)

```
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAATTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
AGAAGTCCGGAGGCTGTTTATACTCGAATCATGATGAATGGAGGTCGACTAAAGAGATCTCACATACGGAGATATGTCTCAGTCAGTTCCAATCATCAAGCCCGTCCAAACTCATTCGCCGAGTTTCTAAACAAGACATATAGTAGTGACAGTGGT
```

```
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTCCCACCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
TCCTGAGAGAACCTCTACTTCCAAAGTGGCTCCGGTAGCCATGGCTTCCCGCCGGAGGTGGAGGAGCAGGATGATGGCACGCTGCCCATGTCTTGTGCCCAGGAGAGCGGGATGGACCGTCACCCTGCAGCCTGTGCTTCTGCTAGGATCA
```

```
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
ATGTGTAAGAAGTTGAATAACAAAATGCCGGAAATCTACGGATTGTGTATATCCATCATGAAAAAAACTAACACCCCTCCTTTCGAACCATCCCAAACATGAGCAAGATCTTTGTCAATCCTAGTGCTATTAGAGCCGGTCTGGCCGAT
```

`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG
`CTTGAGATGGCTGAAGAAACTGTTGATCTGATCAATAGAAATATCGAAGACAATCAGGCTCATCTCCAAGGGGAACCCATAGAGGTGGACAATCTCCCTGAGGATATGGGGCGACTTCACCTGGATGATGGAAAATCG

Supplementary File S8: Counts and statistical analyses of labeled neurons in transsynaptic tracing experiments.

Sheet 1: Detailed counts of tdTomato-labeled neurons found in every (6th) brain section for all mice, including cell numbers from ipsilateral cortex, contralateral cortex, and thalamus, as well as counts of source cells for each animal;

Sheet 2: Total counts of labeled cells for each mouse in each condition, as well as the ratios of tdTomato+ cells to source cells and the average numbers for each condition;

Sheet 3: P-values of comparisons using single-factor ANOVAs.

Condition	Mouse number	Section number	Ipsilateral cortex tdTomato	Ipsilateral cortex tdTomato + mTagBFP2	Ipsilateral cortex tdTomato + mTagBFP2 + H2b-emiRFP670	ipsilateral tdTomato + H2b-emiRFP670 (for No-G controls only)	contralateral	thalamus
G without TEVP for N*PEST, 7 days	22022401LJ	1	236	0			2	0
G without TEVP for N*PEST, 7 days	22022401LJ	2	158	0			3	0
G without TEVP for N*PEST, 7 days	22022401LJ	3	187	0			0	0
G without TEVP for N*PEST, 7 days	22022401LJ	4	233	0			1	0
G without TEVP for N*PEST, 7 days	22022401LJ	5	221	0			1	0
G without TEVP for N*PEST, 7 days	22022401LJ	6	279	0			5	0
G without TEVP for N*PEST, 7 days	22022401LJ	7	603	0			3	3
G without TEVP for N*PEST, 7 days	22022401LJ	8	1446	8			16	39
G without TEVP for N*PEST, 7 days	22022401LJ	9	1076	22			22	76
G without TEVP for N*PEST, 7 days	22022401LJ	10	1323	19			28	61
G without TEVP for N*PEST, 7 days	22022401LJ	11	762	0			20	203
G without TEVP for N*PEST, 7 days	22022401LJ	12	499	0			19	249
G without TEVP for N*PEST, 7 days	22022401LJ	13	283	0			23	44
G without TEVP for N*PEST, 7 days	22022401LJ	14	183	0			6	10
G without TEVP for N*PEST, 7 days	22022401LJ	15	283	0			0	11
G without TEVP for N*PEST, 7 days	22022401LJ	total:	7772	49			149	696
G without TEVP for N*PEST, 7 days	22022402LJ	1	24	0			0	0
G without TEVP for N*PEST, 7 days	22022402LJ	2	36	0			0	0
G without TEVP for N*PEST, 7 days	22022402LJ	3	25	0			1	0
G without TEVP for N*PEST, 7 days	22022402LJ	4	36	0			1	0
G without TEVP for N*PEST, 7 days	22022402LJ	5	51	0			0	0
G without TEVP for N*PEST, 7 days	22022402LJ	6	57	0			1	0
G without TEVP for N*PEST, 7 days	22022402LJ	7	80	0			2	0
G without TEVP for N*PEST, 7 days	22022402LJ	8	515	0			3	14
G without TEVP for N*PEST, 7 days	22022402LJ	9	632	13			12	8
G without TEVP for N*PEST, 7 days	22022402LJ	10	447	0			4	9
G without TEVP for N*PEST, 7 days	22022402LJ	11	222	0			3	14
G without TEVP for N*PEST, 7 days	22022402LJ	12	229	0			1	23
G without TEVP for N*PEST, 7 days	22022402LJ	13	53	0			0	6
G without TEVP for N*PEST, 7 days	22022402LJ	14	22	0			0	0
G without TEVP for N*PEST, 7 days	22022402LJ	15	18	0			0	0
G without TEVP for N*PEST, 7 days	22022402LJ	total:	2447	13			28	74
G without TEVP for N*PEST, 7 days	22022403LJ	1	20	0			1	0
G without TEVP for N*PEST, 7 days	22022403LJ	2	75	0			1	0
G without TEVP for N*PEST, 7 days	22022403LJ	3	37	0			0	0
G without TEVP for N*PEST, 7 days	22022403LJ	4	49	0			3	0
G without TEVP for N*PEST, 7 days	22022403LJ	5	91	0			8	0
G without TEVP for N*PEST, 7 days	22022403LJ	6	156	0			8	0
G without TEVP for N*PEST, 7 days	22022403LJ	7	296	0			8	1
G without TEVP for N*PEST, 7 days	22022403LJ	8	621	4			1	48
G without TEVP for N*PEST, 7 days	22022403LJ	9	1193	11			4	33
G without TEVP for N*PEST, 7 days	22022403LJ	10	979	5			5	38
G without TEVP for N*PEST, 7 days	22022403LJ	11	291	0			2	36
G without TEVP for N*PEST, 7 days	22022403LJ	12	95	0			1	36
G without TEVP for N*PEST, 7 days	22022403LJ	13	65	0			0	0
G without TEVP for N*PEST, 7 days	22022403LJ	14	39	0			0	0
G without TEVP for N*PEST, 7 days	22022403LJ	15	27	0			0	0
G without TEVP for N*PEST, 7 days	22022403LJ	total:	4034	20			42	192
G with TEVP for N*PEST, 7 days	22022501LJ	1	9	0	0		0	0
G with TEVP for N*PEST, 7 days	22022501LJ	2	9	0	0		0	0
G with TEVP for N*PEST, 7 days	22022501LJ	3	38	0	0		1	0
G with TEVP for N*PEST, 7 days	22022501LJ	4	31	0	0		0	0
G with TEVP for N*PEST, 7 days	22022501LJ	5	29	0	0		0	0
G with TEVP for N*PEST, 7 days	22022501LJ	6	36	0	0		0	0
G with TEVP for N*PEST, 7 days	22022501LJ	7	71	0	0		1	0
G with TEVP for N*PEST, 7 days	22022501LJ	8	177	0	0		4	0
G with TEVP for N*PEST, 7 days	22022501LJ	9	933	8	8		4	26
G with TEVP for N*PEST, 7 days	22022501LJ	10	936	6	6		2	7
G with TEVP for N*PEST, 7 days	22022501LJ	11	858	7	7		3	70
G with TEVP for N*PEST, 7 days	22022501LJ	12	432	1	1		1	26
G with TEVP for N*PEST, 7 days	22022501LJ	13	42	0	0		0	39
G with TEVP for N*PEST, 7 days	22022501LJ	14	23	0	0		0	0
G with TEVP for N*PEST, 7 days	22022501LJ	15	18	0	0		0	0
G with TEVP for N*PEST, 7 days	22022501LJ	total:	3642	22	22		16	168
G with TEVP for N*PEST, 7 days	22022502LJ	1	34	0	0		2	0
G with TEVP for N*PEST, 7 days	22022502LJ	2	52	0	0		1	0
G with TEVP for N*PEST, 7 days	22022502LJ	3	61	0	0		1	0
G with TEVP for N*PEST, 7 days	22022502LJ	4	132	0	0		1	0
G with TEVP for N*PEST, 7 days	22022502LJ	5	117	0	0		5	0
G with TEVP for N*PEST, 7 days	22022502LJ	6	193	0	0		6	0
G with TEVP for N*PEST, 7 days	22022502LJ	7	336	0	0		3	0
G with TEVP for N*PEST, 7 days	22022502LJ	8	610	1	1		8	31
G with TEVP for N*PEST, 7 days	22022502LJ	9	597	13	13		7	23
G with TEVP for N*PEST, 7 days	22022502LJ	10	539	14	14		4	29
G with TEVP for N*PEST, 7 days	22022502LJ	11	598	21	21		9	60
G with TEVP for N*PEST, 7 days	22022502LJ	12	788	1	1		5	95
G with TEVP for N*PEST, 7 days	22022502LJ	13	227	0	0		6	0
G with TEVP for N*PEST, 7 days	22022502LJ	14	110	0	0		1	0
G with TEVP for N*PEST, 7 days	22022502LJ	15	171	0	0		0	0
G with TEVP for N*PEST, 7 days	22022502LJ	total:	4565	50	50		59	238
G with TEVP for N*PEST, 7 days	22022503LJ	1	14	0	0		0	0
G with TEVP for N*PEST, 7 days	22022503LJ	2	60	0	0		1	0
G with TEVP for N*PEST, 7 days	22022503LJ	3	147	0	0		1	0
G with TEVP for N*PEST, 7 days	22022503LJ	4	88	0	0		2	0
G with TEVP for N*PEST, 7 days	22022503LJ	5	112	0	0		3	0
G with TEVP for N*PEST, 7 days	22022503LJ	6	64	0	0		1	0
G with TEVP for N*PEST, 7 days	22022503LJ	7	178	0	0		3	0
G with TEVP for N*PEST, 7 days	22022503LJ	8	337	0	0		9	0
G with TEVP for N*PEST, 7 days	22022503LJ	9	663	7	7		7	61
G with TEVP for N*PEST, 7 days	22022503LJ	10	1931	8	8		5	33
G with TEVP for N*PEST, 7 days	22022503LJ	11	1496	11	11		5	71
G with TEVP for N*PEST, 7 days	22022503LJ	12	595	2	2		1	70
G with TEVP for N*PEST, 7 days	22022503LJ	13	254	0	0		0	77
G with TEVP for N*PEST, 7 days	22022503LJ	14	241	0	0		1	69
G with TEVP for N*PEST, 7 days	22022503LJ	15	186	0	0		1	15
G with TEVP for N*PEST, 7 days	22022503LJ	total:	6366	28	28		40	396
No-G control with TEVP for N*PEST, 7 days	22022601LJ	1	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	2	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	3	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	4	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	5	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	6	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	7	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	8	6			2	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	9	75			30	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	10	114			44	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	11	100			40	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	12	63			22	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	13	2			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	14	13			6	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	15	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022601LJ	total:	373			144	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	1	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	2	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	3	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	4	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	5	1			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	6	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	7	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	8	2			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	9	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	10	6			1	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	11	50			29	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	12	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	13	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	14	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	15	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022602LJ	total:	59			30	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	1	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	2	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	3	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	4	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	5	1			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	6	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	7	12			6	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	8	140			30	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	9	75			26	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	10	14			9	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	11	1			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	12	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	13	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	14	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	15	0			0	0	0
No-G control with TEVP for N*PEST, 7 days	22022603LJ	total:	243			71	0	0
G without TEVP for NPEST, 7 days	22022801LJ	1	5	0			0	0
G without TEVP for NPEST, 7 days	22022801LJ	2	9	0			0	0
G without TEVP for NPEST, 7 days	22022801LJ	3	12	0			1	0
G without TEVP for NPEST, 7 days	22022801LJ	4	4	0			0	0
G without TEVP for NPEST, 7 days	22022801LJ	5	14	0			0	0
G without TEVP for NPEST, 7 days	22022801LJ	6	37	0			0	0

Label	ID	Week	C1	C2	C3	ColA	ColB
G without TEVP for NPEST, 7 days	22022801LJ	7	212	11		0	0
G without TEVP for NPEST, 7 days	22022801LJ	8	627	84		4	0
G without TEVP for NPEST, 7 days	22022801LJ	9	130	0		0	0
G without TEVP for NPEST, 7 days	22022801LJ	10	479	41		0	1
G without TEVP for NPEST, 7 days	22022801LJ	11	16	0		0	0
G without TEVP for NPEST, 7 days	22022801LJ	12	6	0		0	1
G without TEVP for NPEST, 7 days	22022801LJ	13	0	0		0	0
G without TEVP for NPEST, 7 days	22022801LJ	14	0	0		0	0
G without TEVP for NPEST, 7 days	22022801LJ	15	0	0		0	0
G without TEVP for NPEST, 7 days	22022801LJ	total:	1551	136		5	2
G without TEVP for NPEST, 7 days	22022802LJ	1	4	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	2	2	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	3	7	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	4	1	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	5	8	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	6	38	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	7	232	13		0	0
G without TEVP for NPEST, 7 days	22022802LJ	8	551	87		0	0
G without TEVP for NPEST, 7 days	22022802LJ	9	225	12		0	1
G without TEVP for NPEST, 7 days	22022802LJ	10	38	0		0	5
G without TEVP for NPEST, 7 days	22022802LJ	11	5	0		1	2
G without TEVP for NPEST, 7 days	22022802LJ	12	9	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	13	1	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	14	1	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	15	1	0		0	0
G without TEVP for NPEST, 7 days	22022802LJ	total:	1123	112		1	8
G without TEVP for NPEST, 7 days	22031508LJ	1	1	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	2	0	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	3	0	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	4	0	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	5	0	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	6	0	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	7	5	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	8	3	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	9	23	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	10	302	12		0	0
G without TEVP for NPEST, 7 days	22031508LJ	11	61	3		0	2
G without TEVP for NPEST, 7 days	22031508LJ	12	1	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	13	1	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	14	0	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	15	0	0		0	0
G without TEVP for NPEST, 7 days	22031508LJ	total:	397	15		0	2
G with TEVP for NPEST rabies: 7 days	22030101LJ	1	133	0	0	2	0
G with TEVP for NPEST rabies: 7 days	22030101LJ	2	155	0	0	0	0
G with TEVP for NPEST rabies: 7 days	22030101LJ	3	195	0	0	2	0
G with TEVP for NPEST rabies: 7 days	22030101LJ	4	492	0	0	3	0
G with TEVP for NPEST rabies: 7 days	22030101LJ	5	939	0	0	3	0
G with TEVP for NPEST rabies: 7 days	22030101LJ	6	841	5	5	8	0
G with TEVP for NPEST rabies: 7 days	22030101LJ	7	1653	21	21	7	28
G with TEVP for NPEST rabies: 7 days	22030101LJ	8	1661	16	16	19	29
G with TEVP for NPEST rabies: 7 days	22030101LJ	9	2414	8	8	47	53
G with TEVP for NPEST rabies: 7 days	22030101LJ	10	556	0	0	15	52
G with TEVP for NPEST rabies: 7 days	22030101LJ	11	508	0	0	2	16
G with TEVP for NPEST rabies: 7 days	22030101LJ	12	246	0	0	0	10
G with TEVP for NPEST rabies: 7 days	22030101LJ	13	98	0	0	0	4
G with TEVP for NPEST rabies: 7 days	22030101LJ	14	52	0	0	0	14
G with TEVP for NPEST rabies: 7 days	22030101LJ	15	44	0	0	0	12
G with TEVP for NPEST rabies: 7 days	22030101LJ	total:	9987	50	50	108	218
G with TEVP for NPEST rabies: 7 days	22030102LJ	1	220	0	0	1	0
G with TEVP for NPEST rabies: 7 days	22030102LJ	2	357	0	0	2	0
G with TEVP for NPEST rabies: 7 days	22030102LJ	3	318	0	0	5	0
G with TEVP for NPEST rabies: 7 days	22030102LJ	4	627	0	0	8	0
G with TEVP for NPEST rabies: 7 days	22030102LJ	5	702	1	1	18	0
G with TEVP for NPEST rabies: 7 days	22030102LJ	6	1349	4	4	12	0
G with TEVP for NPEST rabies: 7 days	22030102LJ	7	1815	4	4	10	0
G with TEVP for NPEST rabies: 7 days	22030102LJ	8	2239	14	14	21	45
G with TEVP for NPEST rabies: 7 days	22030102LJ	9	1425	2	2	9	33
G with TEVP for NPEST rabies: 7 days	22030102LJ	10	859	0	0	7	53
G with TEVP for NPEST rabies: 7 days	22030102LJ	11	327	0	0	0	53
G with TEVP for NPEST rabies: 7 days	22030102LJ	12	258	0	0	1	14
G with TEVP for NPEST rabies: 7 days	22030102LJ	13	73	0	0	0	17
G with TEVP for NPEST rabies: 7 days	22030102LJ	14	43	0	0	1	2
G with TEVP for NPEST rabies: 7 days	22030102LJ	15	18	0	0	0	5
G with TEVP for NPEST rabies: 7 days	22030102LJ	total:	10630	25	25	95	222
G with TEVP for NPEST rabies: 7 days	22030103LJ	1	18	0	0	0	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	2	34	0	0	0	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	3	79	0	0	1	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	4	69	0	0	0	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	5	100	0	0	1	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	6	102	0	0	0	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	7	449	0	0	5	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	8	682	1	1	10	0
G with TEVP for NPEST rabies: 7 days	22030103LJ	9	1093	7	7	8	29
G with TEVP for NPEST rabies: 7 days	22030103LJ	10	1412	4	4	15	18
G with TEVP for NPEST rabies: 7 days	22030103LJ	11	1023	1	1	0	45
G with TEVP for NPEST rabies: 7 days	22030103LJ	12	546	0	0	2	31
G with TEVP for NPEST rabies: 7 days	22030103LJ	13	96	0	0	0	5
G with TEVP for NPEST rabies: 7 days	22030103LJ	14	58	0	0	0	4
G with TEVP for NPEST rabies: 7 days	22030103LJ	15	27	0	0	0	1
G with TEVP for NPEST rabies: 7 days	22030103LJ	total:	5788	13	13	42	133
No-G control with TEVP for NPEST rabies, 7 days	22031407LJ	10	1		0	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031407LJ	11	21		6	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031407LJ	12	1		0	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031407LJ	total:	23		6	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031408LJ	8	8		0	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031408LJ	9	66		10	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031408LJ	10	12		6	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031408LJ	11	12		3	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031408LJ	total:	98		19	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031507LJ	9	18		4	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031507LJ	10	43		12	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031507LJ	11	7		0	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031507LJ	12	1		0	0	0
No-G control with TEVP for NPEST rabies, 7 days	22031507LJ	total:	69		16	0	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	1	340	0		15	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	2	227	0		12	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	3	260	0		14	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	4	437	0		22	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	5	500	0		24	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	6	655	0		18	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	7	426	0		14	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	8	617	0		24	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	9	437	0		29	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	10	1527	0		38	1
G without TEVP for N*PEST, 3 weeks	22022404LJ	11	1951	1		37	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	12	2539	2		63	32
G without TEVP for N*PEST, 3 weeks	22022404LJ	13	2393	9		58	79
G without TEVP for N*PEST, 3 weeks	22022404LJ	14	2542	4		42	33
G without TEVP for N*PEST, 3 weeks	22022404LJ	15	3203	5		61	30
G without TEVP for N*PEST, 3 weeks	22022404LJ	16	2911	0		40	23
G without TEVP for N*PEST, 3 weeks	22022404LJ	17	865	0		15	66
G without TEVP for N*PEST, 3 weeks	22022404LJ	18	353	0		6	44
G without TEVP for N*PEST, 3 weeks	22022404LJ	19	218	0		7	33
G without TEVP for N*PEST, 3 weeks	22022404LJ	20	377	0		3	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	21	124	0		3	0
G without TEVP for N*PEST, 3 weeks	22022404LJ	total:	22902	21		545	341
G without TEVP for N*PEST, 3 weeks	22022405LJ	1	174	0		6	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	2	161	0		10	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	3	210	0		7	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	4	285	0		7	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	5	220	0		18	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	6	203	0		20	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	7	243	0		9	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	8	282	0		6	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	9	255	0		16	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	10	370	0		15	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	11	438	0		28	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	12	1130	0		28	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	13	1601	0		39	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	14	2712	1		55	5
G without TEVP for N*PEST, 3 weeks	22022405LJ	15	2906	2		83	16
G without TEVP for N*PEST, 3 weeks	22022405LJ	16	4065	4		135	90
G without TEVP for N*PEST, 3 weeks	22022405LJ	17	3543	0		88	10
G without TEVP for N*PEST, 3 weeks	22022405LJ	18	3275	0		77	38
G without TEVP for N*PEST, 3 weeks	22022405LJ	19	1714	0		31	54
G without TEVP for N*PEST, 3 weeks	22022405LJ	20	804	0		21	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	21	381	0		4	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	22	233	0		1	0
G without TEVP for N*PEST, 3 weeks	22022405LJ	total:	25205	7		704	213
G without TEVP for N*PEST, 3 weeks	22022406LJ	1	35	0		4	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	2	55	0		0	0

Description	ID	Idx	Count	A	B	C	D	E
G without TEVP for N*PEST, 3 weeks	22022406LJ	3	44	0			2	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	4	265	0			2	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	5	110	0			7	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	6	142	0			8	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	7	124	0			3	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	8	264	0			5	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	9	157	0			10	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	10	104	0			3	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	11	253	0			3	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	12	358	0			2	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	13	500	0			4	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	14	1929	0			3	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	15	2209	0			13	6
G without TEVP for N*PEST, 3 weeks	22022406LJ	16	2633	0			21	47
G without TEVP for N*PEST, 3 weeks	22022406LJ	17	3166	0			26	45
G without TEVP for N*PEST, 3 weeks	22022406LJ	18	1108	0			24	54
G without TEVP for N*PEST, 3 weeks	22022406LJ	19	345	0			17	25
G without TEVP for N*PEST, 3 weeks	22022406LJ	20	232	0			6	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	21	92	0			5	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	22	44	0			0	0
G without TEVP for N*PEST, 3 weeks	22022406LJ	total:	14169	0			168	177
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	1	168	0	0		7	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	2	224	0	0		7	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	3	210	0	0		9	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	4	208	0	0		12	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	5	209	0	0		9	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	6	846	0	0		5	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	7	1673	0	0		13	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	8	2465	0	0		19	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	9	2248	4	4		23	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	10	2581	0	0		22	6
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	11	2578	0	0		16	15
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	12	1277	0	0		15	18
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	13	379	0	0		5	36
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	14	210	0	0		0	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	15	366	0	0		0	0
G with TEVP for N*PEST rabies, 3 weeks	22022504LJ	total:	15642	4	4		162	75
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	1	154	0	0		11	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	2	262	0	0		18	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	3	347	0	0		16	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	4	373	0	0		18	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	5	391	0	0		20	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	6	395	0	0		23	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	7	623	0	0		24	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	8	1607	1	1		24	0
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	9	3515	2	2		69	11
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	10	4278	2	2		70	62
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	11	2667	2	2		110	33
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	12	2405	0	0		149	51
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	13	1120	0	0		159	63
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	14	611	0	0		109	39
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	15	439	0	0		38	15
G with TEVP for N*PEST rabies, 3 weeks	22022505LJ	total:	19187	7	7		858	274
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	1	97	0	0		6	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	2	113	0	0		6	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	3	224	0	0		9	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	4	213	0	0		4	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	5	181	0	0		14	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	6	128	0	0		10	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	7	149	0	0		17	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	8	235	0	0		26	0
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	9	644	0	0		20	1
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	10	856	0	0		38	15
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	11	1600	0	0		48	43
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	12	1343	0	0		31	25
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	13	657	0	0		13	39
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	14	316	0	0		4	38
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	15	301	0	0		6	23
G with TEVP for N*PEST rabies, 3 weeks	22022506LJ	total:	7057	0	0		252	184
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	1	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	2	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	3	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	4	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	5	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	6	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	7	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	8	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	9	5			2	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	10	12			6	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	11	3			2	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	12	1			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	13	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	14	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	15	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022604LJ	total:	21			10	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	1	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	2	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	3	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	4	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	5	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	6	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	7	1			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	8	3			2	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	9	4			2	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	10	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	11	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	12	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	13	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	14	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	15	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022605LJ	total:	8			4	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	1	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	2	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	3	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	4	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	5	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	6	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	7	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	8	4			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	9	9			3	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	10	3			2	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	11	1			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	12	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	13	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	14	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	15	0			0	0	0
No-G control with TEVP for N*PEST, 3 weeks	22022606LJ	total:	17			5	0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	1	12	0			0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	2	7	0			1	0
G without TEVP for NPEST, 3 weeks	22022803LJ	3	6	0			0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	4	10	0			0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	5	14	0			0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	6	85	0			0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	7	474	0			2	0
G without TEVP for NPEST, 3 weeks	22022803LJ	8	694	20			1	2
G without TEVP for NPEST, 3 weeks	22022803LJ	9	427	13			3	3
G without TEVP for NPEST, 3 weeks	22022803LJ	10	70	1			2	4
G without TEVP for NPEST, 3 weeks	22022803LJ	11	25	0			0	3
G without TEVP for NPEST, 3 weeks	22022803LJ	12	16	0			0	2
G without TEVP for NPEST, 3 weeks	22022803LJ	13	11	0			0	1
G without TEVP for NPEST, 3 weeks	22022803LJ	14	6	0			0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	15	1	0			0	0
G without TEVP for NPEST, 3 weeks	22022803LJ	total:	1858	34			9	16
G without TEVP for NPEST, 3 weeks	22022804LJ	1	3	0			0	0
G without TEVP for NPEST, 3 weeks	22022804LJ	2	10	0			1	0
G without TEVP for NPEST, 3 weeks	22022804LJ	3	18	0			0	0
G without TEVP for NPEST, 3 weeks	22022804LJ	4	5	0			0	0
G without TEVP for NPEST, 3 weeks	22022804LJ	5	10	0			1	0
G without TEVP for NPEST, 3 weeks	22022804LJ	6	16	0			0	0
G without TEVP for NPEST, 3 weeks	22022804LJ	7	36	0			0	0
G without TEVP for NPEST, 3 weeks	22022804LJ	8	168	0			2	0
G without TEVP for NPEST, 3 weeks	22022804LJ	9	491	27			2	1
G without TEVP for NPEST, 3 weeks	22022804LJ	10	405	47			0	4
G without TEVP for NPEST, 3 weeks	22022804LJ	11	433	9			0	6
G without TEVP for NPEST, 3 weeks	22022804LJ	12	74	0			0	2
G without TEVP for NPEST, 3 weeks	22022804LJ	13	14	0			0	1
G without TEVP for NPEST, 3 weeks	22022804LJ	14	28	0			0	0
G without TEVP for NPEST, 3 weeks	22022804LJ	15	5	0			0	1
G without TEVP for NPEST, 3 weeks	22022804LJ	total:	1716	83			6	15
G without TEVP for NPEST, 3 weeks	22022805LJ	1	2	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	2	8	0			0	0

Sample	ID	#	V1	V2	V3	V4	V5	V6
G without TEVP for NPEST, 3 weeks	22022805LJ	3	4	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	4	1	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	5	3	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	6	2	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	7	16	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	8	53	1			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	9	231	14			0	2
G without TEVP for NPEST, 3 weeks	22022805LJ	10	147	7			1	0
G without TEVP for NPEST, 3 weeks	22022805LJ	11	31	0			1	1
G without TEVP for NPEST, 3 weeks	22022805LJ	12	13	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	13	10	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	14	1	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	15	1	0			0	0
G without TEVP for NPEST, 3 weeks	22022805LJ	total:	523	22			2	3
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	1	197	0	0		4	0
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	2	285	0	0		5	0
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	3	189	0	0		2	0
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	4	221	0	0		9	0
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	5	216	0	0		13	0
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	6	251	0	0		23	0
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	7	380	0	0		26	1
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	8	766	0	0		46	15
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	9	1653	0	0		36	33
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	10	1955	0	0		31	20
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	11	1633	1	1		12	30
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	12	1104	0	0		3	38
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	13	545	0	0		4	59
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	14	397	0	0		2	49
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	15	252	0	0		0	16
G with TEVP for NPEST rabies: 3 weeks	22030104LJ	total:	10044	1	1		216	261
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	1	59	0	0		3	0
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	2	66	0	0		1	0
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	3	70	0	0		0	0
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	4	65	0	0		0	0
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	5	85	0	0		4	0
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	6	388	0	0		7	0
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	7	519	0	0		5	1
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	8	1200	1	1		7	9
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	9	1395	0	0		8	5
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	10	561	1	0		7	11
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	11	155	0	0		3	10
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	12	73	0	0		1	15
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	13	61	0	0		1	15
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	14	25	0	0		0	4
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	15	15	0	0		0	4
G with TEVP for NPEST rabies: 3 weeks	22030105LJ	total:	4737	2	1		47	74
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	1	386	0	0		15	0
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	2	598	0	0		7	0
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	3	753	0	0		18	0
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	4	1487	0	0		38	0
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	5	1577	1	1		24	0
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	6	1904	3	3		52	0
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	7	2688	1	1		57	45
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	8	2273	0	0		42	79
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	9	1701	0	0		40	61
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	10	448	0	0		16	61
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	11	157	0	0		11	87
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	12	130	0	0		8	72
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	13	36	0	0		1	62
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	14	25	0	0		0	29
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	15	13	0	0		0	17
G with TEVP for NPEST rabies: 3 weeks	22030106LJ	total:	14176	5	5		329	513
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	1	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	2	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	3	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	4	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	5	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	6	1			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	7	2			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	8	3			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	9	8			0	4	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	10	1			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	11	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	12	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	13	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	14	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	15	0			0	0	1
No-G control with TEVP for NPEST rabies, 3 weeks	22030202LJ	total:	15			0	4	1
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	1	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	2	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	3	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	4	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	5	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	6	1			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	7	7			3	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	8	4			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	9	2			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	10	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	11	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	12	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	13	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	14	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	15	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030203LJ	total:	14			3	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	1	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	2	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	3	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	4	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	5	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	6	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	7	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	8	9			5	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	9	3			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	10	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	11	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	12	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	13	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	14	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	15	0			0	0	0
No-G control with TEVP for NPEST rabies, 3 weeks	22030206LJ	total:	12			5	0	0

Ipsilateral cortex comparison

Ipsilateral cortex comparison	p-value
7 days w/o TEVP NPEST vs N*PEST	0.082086041
3 weeks w/o TEVP NPEST vs N*PEST	**0.004609673**
7 days with TEVP NPEST vs N*PEST	0.083091849
3 weeks with TEVP NPEST vs N*PEST	0.394341672
7 days NPEST with TEVP vs N*PEST w/o TEVP	0.138062424
3 weeks NPEST with TEVP vs N*PEST w/o TEVP	0.062362974
7 days NPEST without TEVP vs no G controls	0.159442998
3 weeks NPEST without TEVP vs no G controls	0.694330355
7 days no G control: NPEST vs N*PEST	0.159442998
3 weeks no G control: NPEST vs N*PEST	0.694330355

Ratio of tdTomato+ cells to source cells	p-value
7 days ipsilateral cortex with TEVP	0.09061624
3 weeks ipsilateral cortex with TEVP	0.62501912
7 days ipsilateral cortex NPEST w/ TEVP vs N*PEST w/o TEVP	0.094649458
3 weeks ipsilateral cortex NPEST w/ TEVP vs N*PEST w/o TEVP	0.472690245
7 days ipsilateral cortex without TEVP	**0.000266134**
3 weeks ipsilateral cortex without TEVP	0.089916919

Ratio of tdTomato+ cells to No-G mean (tdTomato + H2b-emiRFP670)	p-value
7 days ipsilateral cortex with TEVP	**0.006338478**
3 weeks ipsilateral cortex with TEVP	0.059221584
7 days ipsilateral cortex without TEVP	0.624238875
3 weeks ipsilateral cortex without TEVP	**0.039909542**

Source cells comparison	p-value
7 days without TEVP: NPEST vs N*PEST	0.193032609
3 weeks without TEVP: NPEST vs N*PEST	0.13286474
7 days with TEVP: NPEST vs N*PEST	0.786754292
3 weeks with TEVP: NPEST vs N*PEST	0.693196503
7 days vs 3 weeks: NPEST without TEVP	0.374876009
7 days vs 3 weeks: N*PEST without TEVP	0.227281063
7 days vs 3 weeks: NPEST with TEVP	0.071820149
7 days vs 3 weeks: N*PEST with TEVP	**0.027514001**

Contralateral Comparison	P-value
7 days w/o TEVP NPEST vs N*PEST	0.13696066
3 weeks w/o TEVP NPEST vs N*PEST	**0.04256234**
7 days with TEVP NPEST vs N*PEST	0.14163061
3 weeks with TEVP NPEST vs N*PEST	0.38647589
7 days NPEST with TEVP vs N*PEST w/o TEVP	0.85084588
3 weeks NPEST with TEVP vs N*PEST w/o TEVP	0.1988985
7 days NPEST without TEVP vs no G controls	all 0
3 weeks NPEST without TEVP vs no G controls	0.37390097
7 days no G control: NPEST vs N*PEST	#DIV/0!
3 weeks no G control: NPEST vs N*PEST	0.37390097

Ratio of tdTomato+ cells to source cells	p-value
7 days contralateral cortex with TEVP	**0.02023616**
3 weeks contralateral cortex with TEVP	0.81969833
7 days contralateral cortex NPEST w/ TEVP vs N*PEST w/o TEVP	0.32919389
3 weeks contralateral cortex NPEST w/ TEVP vs N*PEST w/o TEVP	0.66461936
7 days contralateral cortex without TEVP	**0.0013778**
3 weeks contralateral cortex without TEVP	0.10795063

Ratio of tdTomato+ cells to No-G mean (tdTomato + H2b-emiRFP670)	p-value
7 days contralateral cortex with TEVP	**0.02069633**
3 weeks contralateral cortex with TEVP	0.2323568
7 days contralateral cortex without TEVP	0.19501029
3 weeks contralateral cortex without TEVP	0.05130947

Thalamus Comparison	P-value
7 days w/o TEVP NPEST vs N*PEST	0.17221776
3 weeks w/o TEVP NPEST vs N*PEST	**0.00964446**
7 days with TEVP NPEST vs N*PEST	0.35715947
3 weeks with TEVP NPEST vs N*PEST	0.49379422
7 days NPEST with TEVP vs N*PEST w/o TEVP	0.53834668
3 weeks NPEST with TEVP vs N*PEST w/o TEVP	0.78940038
7 days NPEST without TEVP vs no G controls	all 0
3 weeks NPEST without TEVP vs no G controls	0.37390097
7 days no G control: NPEST vs N*PEST	#DIV/0!
3 weeks no G control: NPEST vs N*PEST	0.37390097

Ratio of tdTomato+ cells to source cells	p-value
7 days thalamus with TEVP	**0.01984607**
3 weeks thalamus with TEVP	0.31169121
7 days thalamus NPEST w/ TEVP vs N*PEST w/o TEVP	0.72341044
3 weeks thalamus NPEST w/ TEVP vs N*PEST w/o TEVP	0.29010953
7 days thalamus without TEVP	0.20553389
3 weeks thalamus without TEVP	**0.02228686**

Ratio of tdTomato+ cells to No-G mean (tdTomato + H2b-emiRFP670)	p-value
7 days thalamus with TEVP	**0.00935808**
3 weeks thalamus with TEVP	0.11846795
7 days thalamus without TEVP	0.19535066
3 weeks thalamus without TEVP	**0.03404937**

Supplementary File S9: Summary table of mean tdTomato+ cell counts for all conditions.
Mean numbers of tdTomato-positive cells in all three quantified brain regions (ipsilateral cortex, contralateral cortex, and thalamus) for each condition in the *in vivo* experiments using the NPEST and N*PEST viruses (cf. Figures 4 & 5 and Supplementary File S8). Numbers are average totals of labeled cells found in a given mouse brain when examining every sixth 50 μm section (see Methods).

Virus	Survival time	G provided?	TEVP provided?	Ipsilateral cortex	Contralateral cortex	Ipsilateral thalamus
N*PEST	7 days	yes	no	4751.0	73.0	320.7
N*PEST	7 days	yes	yes	4857.7	38.3	267.3
N*PEST	7 days	no	yes	225.0	0.0	0.0
NPEST	7 days	yes	no	1023.7	2.0	4.0
NPEST	7 days	yes	yes	8801.7	81.7	191.0
NPEST	7 days	no	yes	63.3	0.0	0.0
N*PEST	21 days	yes	no	20758.7	472.3	243.7
N*PEST	21 days	yes	yes	13962.0	424.0	177.7
N*PEST	21 days	no	yes	15.3	0.0	0.0
NPEST	21 days	yes	no	1365.7	5.7	11.3
NPEST	21 days	yes	yes	9652.3	197.3	282.7
NPEST	21 days	no	yes	13.7	1.3	0.3

Supplementary Figure S5: Pilot testing of Flp-dependent helper virus AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2.

A) Schematic of experimental design to test new Flp-dependent helper virus AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2, which was designed without the error discovered in AAV1-syn-F14F15S-splitTVA-EGFP-tTA, used in the initial preprint of this paper. The new helper virus as well as AAV1-TREtight-BFP2-B19G were injected in S1, with AAV2-retro-syn-Flpo injected in striatum (or not, for no-Flpo controls), of two Ai14 mice, in order to target corticostriatal neurons. The first-generation rabies virus RVΔG-Cre(EnvA) was injected one week after helper virus injection, and mice were perfused one week later.

B-C) Representative confocal images of injection site (B) and input regions (C) with FLPo virus in each of the two mice.

D-E) Representative confocal images of injection site (D) and input regions (E) without FLPo virus in each of the two mice. Far fewer cells are labeled at the injection sites; very cells are present in input regions.

A

Day -7:
AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2
AAV1-TREtight-mTagBFP2-B19G
in S1

Day 0:
RVΔG-4Cre(EnvA)
in S1

Day -7:
AAV2-retro-synP-FLPo
in striatum

Day 7:
Perfuse

Ai14 mice

B

Injection site: with FLPo

| tdTomato (reports Cre) | EGFP (reports TVA & tTA) | mTagBFP2 (reports G) | Merged |

Mouse 1
Mouse 2

C

Input regions: with FLPo

| M1/M2 cortex | S2 cortex | Thalamus | Contralateral cortex |

Mouse 1
Mouse 2

D

Injection site: without FLPo

| tdTomato (reports Cre) | EGFP (reports TVA & tTA) | mTagBFP2 (reports G) | Merged |

Mouse 1
Mouse 2

E

Input regions: without FLPo

| M1/M2 cortex | S2 cortex | Thalamus | Contralateral cortex |

Mouse 1
Mouse 2

Supplementary Figure S6: *In vivo* **comparison of NPEST and N*PEST viruses without TEVP: results at 7 days (cf. Figure 4).**

A) Schematic of experiment.

B-D) Results at 7 days after injection are similar to those at 21 days (see Fig. 4) but not statistically significant due to high variance.

B) Representative images of injection sites and surrounding ipsilateral cortex.

C) Representative images of input regions.

D) Counts of labeled neurons in ipsilateral cortex (left), contralateral cortex (center), and thalamus (right) at 7 days for the two viruses. Each dot indicates the total number of labeled cells found in a given mouse brain when examining every sixth 50 µm section (see Methods). Although the numbers of labeled cells are much higher in every case for the N*PEST virus than for the NPEST one, the comparisons are not statistically significant (see Supplementary File S8 for all counts and statistical analyses).

A

Day -7:
AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2
AAV1-TREtight-mTagBFP2-B19G
in S1

Day 0:
RVΔG-NPEST-4Cre(EnvA)
("intact")
or RVΔG-N*PEST-4Cre(EnvA)
("revertant")
in S1

Day -7:
AAV2-retro-synP-FLPo
in striatum

Day 7:
Perfuse

Ai14 mice

B

| | tdTomato (reports Cre) | EGFP (reports TVA & tTA) | mTagBFP2 (reports G) | Merged |

Without TEVP: 7 days — NPEST ("intact") / N*PEST ("revertant")

200 µm

C

| | M1/M2 cortex | S2 cortex | Thalamus | Contralateral cortex |

Without TEVP: 7 days — NPEST ("intact") / N*PEST ("revertant")

200 µm

D

Ipsilateral cortex — tdTomato+ cells — NPEST ("intact") / N*PEST ("revertant") — ns

Contralateral cortex — tdTomato+ cells — NPEST ("intact") / N*PEST ("revertant") — ns

Thalamus — tdTomato+ cells — NPEST ("intact") / N*PEST ("revertant") — ns

Supplementary Figure S7: *In vivo* comparison of NPEST and N*PEST viruses with TEVP: results at 7 days (cf. Figure 5).
A) Schematic of experiment.
B-D) Results at 7 days after injection are similar to those at 21 days (see Fig. 5).
B) Representative images of injection sites and surrounding ipsilateral cortex.
C) Representative images of input regions.
D) Counts of labeled neurons in ipsilateral cortex (left), contralateral cortex (center), and thalamus (right) at 7 days for the two viruses. Each dot indicates the total number of labeled cells found in a given mouse brain when examining every sixth 50 μm section (see Methods).
At this timepoint, the average numbers of labeled neurons in ipsilateral and contralateral cortex are higher for the NPEST virus, whereas the average number of labeled thalamic neurons is higher for the N*PEST virus, although again the comparisons are not statistically significant (see Supplementary File S8 for all counts and statistical analyses).

A

Day -7:
AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2
AAV1-TREtight-mTagBFP2-B19G
AAV1-TREtight-TEVP
in S1

Day -7:
AAV2-retro-synP-FLPo
in striatum

Day 0:
RVΔG-NPEST-4Cre(EnvA)
("intact")
or RVΔG-N*PEST-4Cre(EnvA)
("revertant")
in S1

Day 7:
Perfuse

Ai14 mice

B

| H2b-emiRFP670 (reports TEVP) | tdTomato (reports Cre) | EGFP (reports TVA & tTA) | mTagBFP2 (reports G) | Merged |

With TEVP: 7 days — NPEST ("intact"), N*PEST ("revertant")

Scale bar: 200 µm

C

| M1/M2 cortex | S2 cortex | Thalamus | Contralateral cortex | S1 cortex |

With TEVP: 7 days — NPEST ("intact"), N*PEST ("revertant")

Scale bar: 200 µm

D

Ipsilateral cortex

tdTomato+ cells

NPEST TEVP+ / N*PEST TEVP+

ns

Contralateral cortex

tdTomato+ cells

NPEST TEVP+ / N*PEST TEVP+

ns

Thalamus

tdTomato+ cells

NPEST TEVP+ / N*PEST TEVP+

ns

Supplementary Figure S8: Results for control experiments without G.
A) Schematic of experiment. Design is the same as for the transsynaptic tracing experiments with the helper virus expressing TEVP included, except that in these experiments the G-encoding helper virus, AAV1-TREtight-BFP2-B19G, was omitted.
B-C) Images of injection sites using NPEST (top) and N*PEST (bottom) viruses, with survival times of 7 days (B) or 3 weeks (C). Scale bar: 200 μm, applies to all images in each panel.
D) Counts of tdTomato-labeled neurons in ipsilateral cortex, contralateral cortex, and thalamus. Few cells were found in input regions with either virus at either survival time, suggesting that the extensive label found there in experiments in which G was included was primarily due to *bona fide* transneuronal spread rather than direct retrograde infection by injected rabies virus..

A

Day -7:
AAV1-syn-F14F15S-splitTVA-EGFP-tTA_v2
AAV1-TREtight-TEVP
in S1

Day -7:
AAV2-retro-synP-FLPo
in striatum

Day 0:
RVΔG-NPEST-4Cre(EnvA)
("intact")
or RVΔG-N*PEST-4Cre(EnvA)
("revertant")
in S1

Day 7 or 21:
Perfuse

Ai14 mice

B

H2b-emiRFP670 (reports TEVP)	tdTomato (reports Cre)	EGFP (reports TVA & tTA)	mTagBFP2 (reports G)	Merged

No G control: 7 days

NPEST ("intact")

N*PEST ("revertant")

200 µm

C

H2b-emiRFP670 (reports TEVP)	tdTomato (reports Cre)	EGFP (reports TVA & tTA)	mTagBFP2 (reports G)	Merged

No G control: 3 weeks

NPEST ("intact")

N*PEST ("revertant")

200 µm

D

Supplementary Figure S9: Images and counts of source cells for all conditions.
A-D) Representative images of injection sites without TEVP (A-B) or with TEVP (C-D), with 7-day (A & C) or 3-week (B & D) survival time following rabies virus injection. Top rows: NPEST ("intact"); bottom rows: N*PEST ("revertant"). Scale bars: 50 µm, apply to all images in each panel.
E-H) Counts of "source cells" and ratios of input cells to source cells in the absence (E-F) or presence (G-H) of TEVP. Source cells are here defined as cells coexpressing tdTomato and mTagBFP2 when TEVP was omitted, and cells coexpressing tdTomato, mTagBFP2, and emiRFP670 when TEVP was supplied. Note that the average numbers of source cells at 3 weeks are in most cases much lower than at 7 days, for both viruses, with or without TEVP, possibly suggesting toxicity to source cells that results in the death of the majority. Because of this apparent toxicity to source cells, the ratios of input cells to source cells may be artificially high and therefore arguably not particularly meaningful, particularly at the later timepoint: for example, for N*PEST at 3 weeks both with and without TEVP, no source cells were found (E & G), so the ratios for that mouse would be infinite and are omitted from the graphs (F & H). We have therefore omitted significance comparisons for the 3-week timepoint from the graphs, although Supplementary File S8 contains all counts and statistical analyses. I-J) Same as graphs in panels F & G but using the mean number of tdTomato+/emiRFP670+ cells in the 7d G-/TEVP+ control condition for each virus (i.e., 225 for N*PEST, 63.3 for NPEST; cf. Supplementary Figure S8 and Supplementary File S8) as proxy for the unknown number of original source cells in each case.

Supplementary Figure S10: First-generation vector RV△G-4FLPo appears to be toxic to most cells, unlike the comparable first-generation vector RV△G-Cre.
Although we did not rigorously quantify the effect, our FLPo-encoding RV△G-4Flpo appears to kill neurons more quickly than does RV△G-4Cre (cf. Figure 3 and Chatterjee et al. (21)). In this example field of view, most neurons clearly visible at earlier time points have disappeared by 28 days postinjection, leaving degenerating cellular debris. See Discussion for possible reasons why a preparation of a first-generation vector encoding a recombinase may or may not preserve a large percentage of infected neurons. Scale bar: 50 μm, applies to all panels.