Title: Rare Microbial Taxa Emerge When Communities Collide: Freshwater and Marine

Microbiome Responses to Experimental Seawater Intrusion

Jennifer D. Rocca[1], Marie Simonin[1,2], Justin P. Wright[1,3], Alex Washburne[4], Emily Bernhardt[1,3]

[1]Department of Biology, Duke University, Durham, NC, United States

[2]IRD, Cirad, University of Montpellier, IPME, Montpellier, France

[3]Nicholas School of the Environment, Duke University, Durham, NC, United States

[4]Department of Microbiology and Immunology, Montana State University, Bozeman, MT, United States

Running Title: Rare Taxa Emerge during Community Coalescence

Keywords: 16S rRNA, biotic interactions, community coalescence, competition, environmental

filtering, mass effects, mutualism, rare biosphere, salt stress, phylofactorization.

Authorship

JR, MS, EB, and JW conceived of the project. JR and MS set up and harvested the incubation. JR

extracted and processed the sequence data, and MS and AW contributed to the data analysis. All

authors contributed to equally to writing the manuscript.

Conflict of Interest

The contributors declare no conflict(s) of interest.

Corresponding Author

Jennifer D. Rocca

Biology Department, Duke University

Durham, NC 27708

Email: jenny.rocca@gmail.com

Phone: 001-919-684-6556

Abstract

Whole microbial communities regularly merge with one another, often in tandem with their

environments, in a process called community coalescence. Such events allow us to address a central

question in ecology – what processes shape community assembly. We used a reciprocal transplant

and mixing experiment to directly and independently unravel the effects of environmental filtering and

biotic interactions on microbiome success when freshwater and marine communities coalesce. The

brackish treatment and community mixing resulted in strong convergence of microbiome structure and

function toward the marine. Brackish exposure imposed a 96% taxa loss from freshwater and 66%

loss from marine microbiomes, which was somewhat counterbalanced by the emergence of tolerant

rare taxa. Community mixing further resulted in 29% and 49% loss from biotic interactions between

freshwater and marine microbiomes, offset somewhat by mutualistically-assisted rare microbial taxa.

Our study emphasizes the importance of the rare biosphere as a critical component of community

resilience.

Main Text

<u>**INTRODUCTION**</u>

A fundamental goal in ecology is to determine the distribution and abundance of species and

the mechanisms controlling this distribution. This central objective is particularly challenging in

microbial ecology because of the immense diversity, brief lifespan, and microscopic size of

microorganisms. Many studies examine organismal distributions along environmental gradients to

shed light on the natural history of species (Whittaker 1965) and microbial ecologists also use this

concept to understand the natural history of microorganisms at biogeographical scales (Fierer and

Jackson 2006) and along elevational (Bryant *et al.* 2008; Yang *et al.* 2014; Siles and Margesin 2016),

precipitation (Angel *et al.* 2009; Hawkes *et al.* 2017) and salinity (Herlemann *et al.* 2011; Herlemann *et

al.* 2016) gradients. However, distinguishing between environmental tolerance or competitive abilities

for microorganisms and determining their fundamental niche is difficult to assess with just distribution

information. Determining the relative importance of environmental filtering and biotic interactions in

structuring extant communities is difficult as both processes operate concurrently (Vellend 2010;

Goberna *et al.* 2014; Cadotte and Tucker 2017).

Quantifying the influence of these assembly filters is unique for microbial communities, which

typically migrate in the aggregate and in tandem with their environment. Thus 'habitat patches' move

along with an entire assemblage of organisms. In metacommunity ecology, such collective exchanges

are referred to as mass effects (Leibold *et al.* 2004, Souffreau *et al.* 2014, Comte *et al.* 2017). When

previously distinct communities combine along with their respective environments, the reassembly of

the novel community is termed 'community coalescence' (Webb 1976, Livingston *et al.* 2013, Rillig *et*

al. 2015, Rillig & Mansour 2017). Microbial community coalescence occurs every time a leaf falls to

the ground, a soil particle is blown into a new landscape, or two bodies of water mix. Despite the

ubiquity of microbial community coalescence, the formal recognition of this concept is fairly recent in

microbial ecology (Rillig *et al.* 2017; Mansour *et al.* 2018). In fact, Mansour *et al.* (2018) suggest that

many experimental studies with microbial communities are unacknowledged community coalescence

experiments.

Community coalescence is not only an interesting phenomenon, it also provides immense

opportunities to leverage experiments to learn about <u>how</u> individual microorganisms respond and

microbial communities assemble in response to the merging event. To develop the natural history of

microorganisms, we need to understand how both environmental and biological filters ultimately

determine who persists and who does not when communities collide. To address this fundamental

ecological question, we introduce a new methodological framework (Fig. 1) to directly measure the

independent and combined effects of environmental and biotic filters (sensu Vellend 2010) that

structure the distribution and abundance of microbial taxa.

We used a microcosm experiment to examine how microbial communities reassemble

following the blending of media and microbial communities between freshwater and marine

environments to simulate seawater intrusion - a mixing event that occurs along all coastal margins.

Our experimentally created coalescence event (Fig. 1) allowed us to ask: Q1) how do microbial

communities from distinct environments respond to merging? Q2) what is the relative role of

environmental filtering and biotic interactions in structuring this newly united community? Q3) are more

closely related taxa more likely to respond similarly to each filter? and Q4) what are the resulting

consequences for microbial community function?

Seawater intrusion into freshwater systems is a prevalent feature of tidal environments and is

increasing in frequency and extent as drought, irrigation and climate change are all increasing the

inland and upland movement of seawater into freshwater habitats (Weston *et al.* 2006, Herbert *et al.*

2015). The multivariate chemical transition to brackish water is well studied, and salinity is a well-

documented environmental stressor (Lozupone and Knight 2007), imposing strong evolutionary

selection on organisms (Paver *et al.* 2018). When freshwaters come into contact and blend with

seawater, we observe dramatic decreases in organic carbon content due to complexation with salts,

increased pH, and nutrient availability along with increases in salinity (Craft *et al.* 2009, Barlow &

Reichard 2010, Ardón *et al.* 2016a). Dynamic pulses of salty, oligotrophic marine water mixing with

relatively mesotrophic freshwater habitats results in brackish conditions, novel to either endpoint

microbial community. The microbial consortia derived from either endpoint habitat are well adapted to

their respective environmental conditions (Canfora *et al.* 2014, Herlemann *et al.* 2016), and exposure

to brackish water imposes substantial stress on members of each endpoint community, particularly the

freshwater sediment microbial communities (Baldwin *et al.* 2006, Jackson and Vallaire 2009,

Neubauer *et al.* 2013,) and water column microbial taxa (Burke and Baird 1931, Nielsen *et al.* 2003,

Ewert and Deming 2013). While we expect freshwater microorganisms to more stressed by the salt

exposure of the brackish conditions (Edmonds *et al.* 2009, Herbert *et al.* 2015), little is known about

the outcome of blending these distinct communities. We expect a substantial turnover of each

microbiome in response to the strong environmental filter imposed by the brackish exposure (Paver *et*

al. 2018), but less is known about the community responses to novel biotic interactions that might

arise through community blending. Though marine microbial taxa are more stress tolerant to the wide

range of salinity, they may not fare well if the freshwater microbial taxa are better competitors (Grimes

et al. 1977). Do the existing communities have a great phenotypic plasticity or can the rare biosphere

aid in the resilience to drastic environmental changes and community introductions?

Our experimental design allows us to differentiate amongst four potential outcomes of

microbial community coalescence (Fig. 1, top panel). Beginning with the assumption that our starting

endmember communities are distinct, we expect that the community resulting from their exposure to

the mixed environment and to the community merger would follow one of four possible trajectories of

the final community composition (Fig. 1, top panel): A) intermediates between the two home conditions

due to species sorting hinging on the environment, B) stratified yet similar to their initial inocula due to

symmetric community resistance; or C) converge towards the assemblage observed in only one of the

two environments due to strong asymmetric environmental selection *or* asymmetric community

resistance. Alternatively, a final outcome (D) is that the coalescence conditions result in highly variable

emergent communities via stochastic responses. In the scenario where the communities converge

towards a single endmember (C), our methodological framework allows us to directly distinguish

between environmental and biotic controls in driving this response (Fig. 1, bottom panel).

Our integrative research approach allows us the opportunity to add a natural history

perspective to the study of microbial ecology. Habitat transplants will apply an environmental filter to

identify generalist vs specialist microbial taxa. Community coalescence allows us to identify strong vs.

weak competitors and common associations amongst taxa.

MATERIALS and METHODS

Field sample collection and aquatic endmember characterization

The two endmember sources for this microcosm experiment are located in coastal North Carolina,

USA (Table 1). The Freshwater Wetland site, hereafter "Freshwater" site, is a blackwater wetland

ecosystem located within the Timberlake Wetland Restoration Project in Tyrrell Co., NC,

35°53'46.4"N, 76°09'51.4"W (Ardón *et al.* 2016b), exposed to episodic or storm-triggered flows

following rain or coastal storm systems. In contrast, the Coastal Marine site, hereafter "Marine" site,

located at the northern end of the Cape Hatteras National Seashore (Dare Co., NC, 35°49'57.4"N,

75°33'25.7"W) with persistent water turbulence.

The Freshwater and Marine endmember sites were selected for their close proximity (64 km)

and even latitude, while maximizing the potential range of habitats involved in seawater intrusion. Both

sites are not significantly impacted by human activity, and the Freshwater site had no recent seawater

exposure. Additionally, the water samples were specifically collected at the end of a seasonal period

where recent seawater intrusion was least likely. We acknowledge that these sites do not represent

true potential locations of seawater intrusion, as the mixing occurs gradually across the 64 km

gradient. However, the sites are true endmembers, with past seawater intrusion coalescence events.

In June 2016, we collected 80 L of surface water from each site into sterile carboys, stored at the

average temperature (23 $^{\circ}$C) of the two sites at time of collection. Within 12 hours, each sample was

filtered through axenic 1 mm mesh to remove macroorganisms and debris. Each sample was

subsampled for salinity, pH, and dissolved organic carbon. After filtration, the two endmember water

samples were halved to generate: (a) the microbe-free environments and (b) microbial inocula (Fig. 1).

 Microcosm incubation setup

We set up a laboratory incubation, exposing three inocula isolated from each endmember samples

(Freshwater, Marine, and a 1:1 mixture of Freshwater-Marine, hereafter "Coalescence"), into three

axenic aquatic environments (Freshwater, Marine, and a 1:1 mixture of the endmember environments,

hereafter "Brackish"). We sterilized each water source by autoclaving at 121 ^{o}C / 20 PSI for 30 mins in

covered acid-washed glassware. After cooling to room temperature, the autoclaving was repeated

twice more to ensure that any microorganisms exiting dormancy were killed. Subsamples of each

starting environment were streaked on Luria-Bertani agar plates to confirm sterility.

Intact microbial communities were isolated by concentrating cells off the remainder of the

Freshwater and Marine samples. Two complementary concentrating methods were implemented to

minimize biases imposed by either method. Half from each environment was gently centrifuged at

5,000 RCF in 50 mL batches in round-bottom tubes to ensure maximal viability of the microbial

assemblages (Pembrey *et al.* 1999, Peterson *et al.* 2012). In parallel, the remaining water samples

were filtered over gamma-irradiated Pall Supor 0.2 µm nitrocellulose membranes (Millipore, New York,

NY) in small batches to minimize fouling on the filter. For each inoculum, the filter-collected and

centrifuged-collected microbiomes were combined by resuspending in sterilized home. The

Coalescence microbiome was made by mixing 1:1 subsamples of the Freshwater and Marine inocula.

The full experimental design consisted of 120 independent microcosms (850 mL inoculated

with 550 µl), with an experimental replication of five, along with positive/negative controls (Suppl. Fig.

1). Here we present a subset (n=25 microcosms) of the full incubation with five treatment conditions:

"Freshwater-Home": Freshwater microbiome in Freshwater environment, "Marine-Home": Marine

microbiome in Marine environment, "Freshwater-Brackish" and "Marine-Brackish": Freshwater (or

Marine) microbiome in Brackish environment; and "Brackish-Coalescence": a 1:1 blended microbiome

in a 1:1 mixture of the endmember environments. We set up the microcosms into sterile glass Mason

jars under UV-treated PCR-hood conditions. The positive control microcosms were used to correct for

any experimental artifacts from inoculum and environment preparations, which did not vary

significantly from the 'home' microcosms (Suppl. Fig. 2). Microcosms with sterile environments or

sterile water with no added inocula were included as negative controls to confirm axenic conditions,

and none had any detectable microbial growth or genetic material and will not be discussed further.

The incubation ran for seven days in an environmental growth chamber (23 ^{o}C, 13.5 hr diurnal light

regime, and PAR: 250-450 µmols m^{-2}s^{-1}) reflecting field conditions. Twice-daily, the microcosms were

re-randomized to minimize potential biases from differential light and temperature across the chamber,

and each microcosm was mixed by inverting.

Microbial function

Microbial extracellular enzyme potential activity (hereafter enzyme activity) was measured in each

microcosm at the end of the incubation. Enzyme activity for eight enzymes were measured from a 91-

mL homogenized subsample of each microcosm following a protocol developed by Bell *et al. (*2013),

here modified to handle water. The following enzymes were targeted with fluorescently labeled

substrates to capture potential N, P, C, and S degradation activity: α-1,4-glucosidase (AG), aryl-

sulfatase (AS), β-1,4-glucosidase (BG), β-D-1,4-cellobiosidase (CB), L-leucine aminopeptidase (LAP),

β-1,4-N-acetylglucosaminidase (NAG), alkaline phosphatase (PHOS), and β-D-xylosidase (XYL). After

three-hours at room temperature, the centrifuged supernatant for each sample was read (340/460 nm)

on a FLUOstar Optima spectrophotometer (BMG Labtech, Cary, NC, USA) in black optical 96-well

plates.

DNA extraction & bacterial community analysis

After the incubation, a 250 mL subsample of each microcosm was filtered over gamma-irradiated Pall

Supor 0.2 µm nitrocellulose membranes (Millipore, New York, NY), and the filtrate was centrifuged at

10,000 x g for one hour to pellet any remaining ultra-small microorganisms (Luef *et al.* 2015). The

pellet and filter were combined and processed for genomic DNA using a MoBio PowerWater DNA

Isolate Kit (MoBio, Vancouver, CA) modified by adding a heating step during cell lysis. Total genomic

DNA was fluorometrically measured (Quant-iT dsDNA Assay Kit, ThermoScientific, Waltham, MA),

and used as proxy for total microbial biomass (Baas *et al.* 2015, Nagler *et al.* 2018). The samples

were amplified, targeting the V4 hypervariable region of the bacterial 16S rRNA gene (515-F/806-R,

Caporaso *et al.* 2011), and sequenced with Illumina MiSeq (PE 150bp; V2 chemistry) at the

Environmental Sample Preparation and Sequencing Facility (ESPSF) at Argonne National Laboratory.

Raw sequences are deposited into the NCBI Sequence Read Archive (SRA): PRJNAXXXX.

ESPSF returned 25 million raw sequences, which we processed through Quantitative Insights

Into Microbial Ecology 2 (Qiime2) pipeline (Bolyen *et al.* 2018) to remove low quality reads and

putative chimera, and to denoise the sequences into exact sequence variants (SVs) with Dada2

(Callahan *et al.* 2017). We aligned the representative sequences and assigned taxonomy using the

Silva V132 (99%) curated reference alignment (Quast *et al.* 2013), and a phylogeny and improved

alignment were simultaneously generated using the Practical Alignment using SATé and TrAnsitivity

(PASTA) software (Mirarab *et al.* 2013). Eukaryotic, mitochondrial, and chloroplast contaminant

sequences were removed. The SV table was rarefied to lowest sequence depth (17,500 sequences),

and the final dataset contained 3389 unique SVs with 12,293,437 total reads.

Microbiome Structure & Function Profile

To bulk characterize the bacterial communities, alpha-diversity (Chao1) and Pielou's evenness were

calculated on each sample. A Bray-Curtis dissimilarity matrix was created for the community dataset

(and Euclidean for the enzyme dataset) to examine differences in community structure and in the

functional profile among treatments, and to visualize these shifts using non-metric multidimensional

scaling (NMDS). To test the hypothetical outcomes of community structure detailed in Fig. 1A, a non-

parametric multivariate analysis of variance (per-MANOVA, Anderson 2001) was implemented to

identify significant overall and between group shifts in community. The "adonis" function in the vegan

R package (Version 3.3.1, Oksanen *et al.* 2017) was used to implement the perMANOVA by

estimating correlation coefficients and corresponding p-values (permutations = 999) for the effect of

each treatment. Subsequent pairwise comparisons were performed using Adonis. The "procrustes"

function (Jackson 1995) in Vegan was used to perform least-squares orthogonal mapping to

determine correlations between two multivariate datasets. Here, procrustes (PROTEST) was used to

determine correlations between bacterial community structure and microbial enzyme activities profile.

Additionally, we visualized the responses of the most abundant microbial SVs (> 2%) using UPGMA

clustering (Sokal & Michener 1958) to the environmental and microbiome mixing treatments. This

subset of taxa was further identified for halotolerance using the LPSN database (Parte 2013) of known

microbiological traits. Univariate data that violated assumptions of normal distribution and

homoscedasticity were log-transformed, and assumptions were subsequently re-verified with Shapiro-

Wilk and Levene's tests and examination of Q-Q plots. We used one-way ANOVAs to test the

significance of the treatments on microbiome diversity and enzyme activity, then *post-hoc* pairwise

comparisons among the treatment combinations were performed using Tukey's Honestly Significant

Difference multiple means comparison.

Direct separation of contribution of local assembly mechanisms

To directly disentangle the influence of environmental filtering and novel biotic interactions, we

compared specific experimental treatments (Fig. 1, bottom panel). To examine the environmental filter

(*Part 1*), we compared 'home' microbiomes to the corresponding Brackish microbiomes (i.e.

Freshwater-Home vs. Freshwater-Brackish or Marine-Home vs Marine-Brackish). To examine the

impact of novel biotic interactions from community blending (*Part 2*), the Freshwater-Brackish and

Marine-Brackish microbiomes were compared to the Brackish-Coalescence microbiomes. A new

phylogenetic-based method – phylofactorization – was used to identify clades driving changes in

community composition (Washburne *et al* 2017, 2019), based on a Holm's sequentially rejective 5%

cutoff for the family-wise error rate. The resultant phylofactor objects are available in the

supplementary material.

RESULTS

Starting Conditions: Bacterial Community and Chemical Characterization.

Our two environments and initial microbial communities were distinct among the Freshwater and

Marine endmembers (Table 1). Other than water temperature at the time of field collection, all other

water chemistry properties varied substantially between our endmembers (Table 1). Electrical

conductivity was ~740-fold greater and pH was 3.8 units higher in the Marine sample, while the

Freshwater environment had higher concentrations of both nutrients and dissolved organic matter

(Table 1).

The Freshwater and Marine microbial communities were also distinct from one another.

Microbial biomass (Table 1: 749 vs. 218 ng DNA/µL), alpha diversity (Fig. 2A: 328 vs. 253 observed

SVs/microcosm) and evenness (Fig. 2A: 0.1 vs. 0.02) were significantly higher in the Freshwater

microbial community. The two endmember microbiomes had minimal overlap in their community

composition (Fig. 2B), with only 21 SVs (<1.4% of total SVs) representing 22% of the Fresh+Marine

summed biomass (Suppl. Fig. 3). The Freshwater community was dominated by the families:

Acetobacteraceae, Paracaedibacteraceae, Beijerinckiaceae, Burkholderiaceae; while the Marine

microbiome was dominated by the families: *Alteromonadaceae, Rhodobacteraceae, Saprospiraceae,*

Spirosomaceae, and *Vibrionaceae* (Fig. 3) with a single taxon – *Alteromonas* - dominating 9.7% of the

Marine microbiome. The two communities contrasted in their functional potential as well, exhibiting

distinct enzyme profiles, with the Freshwater microbiome producing significantly higher amounts in

seven of the eight enzymes analyzed (Fig. 2C, Suppl. Fig. 4).

<u>*Convergence towards Marine Bacterial Community During Coalescence*</u>

When the two axenic endmember water samples were combined to create our Brackish media, the

resulting chemical properties were intermediate to the two endmembers. Substantial buffering led to

the Brackish media having a pH closer to the endmember, but other chemical components, were

essentially the average of the two contributing media (Table 1). The blending of the two communities

into the Coalescence inoculum was also the average biomass of the two endmembers (Table 1).

Environmental Filter: Despite their higher diversity and biomass (Fig. 2A), the Freshwater microbial

taxa did not fare well when added to the Brackish media in the absence of Marine community blending

(i.e. Freshwater-Brackish treatment). Only 36 of the 967 total taxa initially sequenced from our

Freshwater-Home microbiomes persisted following this environmental filter into Brackish media,

although 169 taxa that were below our detection (i.e. rare taxa) in the initial inoculum were detected in

the Brackish media (Fig. 4). We are confident that these taxa represent increases in abundance from

the rare biosphere contained in the initial inoculum as we failed to detect any genomic DNA in our

negative controls. Marine microbial taxa were more tolerant of the transfer into Brackish media, with

199 of the original 588 taxa surviving. A large number of rare biosphere taxa emerged from Marine

inoculum under Brackish conditions. We detected 393 taxa in our Marine-Brackish treatments that

were not detected in the original Marine-Home microcosms. Consequently, the diversity of the Marine

inoculum in Brackish media was equal to the starting inoculum (592 vs. 588), while there was a

substantial loss of total richness for the Freshwater microbiome in Brackish media compared to the

diversity of its initial inocula (205 vs 967) (Fig. 4). The community composition of the Marine-Brackish

replicates was very similar to the Marine-Home, while the Freshwater-Brackish community shifted

significantly in composition towards the Marine-Home relative to its initial Freshwater-Home

composition (Fig. 2B). We detected more shared taxa between these environmentally filtered

communities, with 65 taxa overlapping between Freshwater-Brackish and Marine-Brackish. For the

low diversity Freshwater-Brackish replicates, these shared taxa represent more than 25% of the total

diversity. These increased taxa include the following families, which were below detection limit in

Freshwater-Home: *Alteromonadaceae*, Oceanospirillaceae, *Rhodobacteraceae*, and *Vibrionaceae*

(Fig. 4). The enzymatic profile of each community followed similar trends, with the Marine-Home,

Marine-Brackish and Freshwater-Brackish replicates having reduced enzyme activity (Suppl. Fig. 4)

and more similar enzyme profiles relative to the Freshwater-Home enzyme profile (Fig. 2C).

Biotic Filter: In contrast to the extreme loss of abundant taxa caused by environmental filtering, the

addition of interacting taxa from the two endmember communities into our Brackish media ('Brackish-

Coalescence') had a more limited effect on microbial richness. Of the 967 Freshwater taxa found in

the initial Freshwater-Home treatment, 37 were detected in the Brackish-Coalescence treatment. This

set overlapped entirely with the set of taxa that survived through the environmental filter, with the

exception of a single taxa that disappeared in the Freshwater-Brackish treatment but increased in

abundance in response to the addition of an interacting community assemblage. There were 145

Freshwater-derived taxa that survived the Brackish treatment but did not persist when in the presence

of the new interacting microbiome (lost in the Brackish-Coalescence treatment). Of the 588 Marine

taxa detected in the original Marine-Home treatment, 171 were detected in the Brackish-Coalescence

treatment. This set of Marine survivors overlapped considerably with the list of taxa that were tolerant

of the environmental filter (with 143 taxa found in both taxa lists). There were 28 Marine-derived taxa

who only persisted in Brackish media when also combined with the interacting microbiome (Brackish-

Coalescence), and there were 302 Marine taxa that could survive the environmental filter (were found

in Marine-Brackish treatments) but could not persist in the presence of the new blended microbiome

(lost in the Brackish-Coalescence treatment).

We introduced at least 1528 taxa from both endmember inocula into the Brackish-

Coalescence treatments (this is the sum of the distinct taxa derived from the two endmember

communities). Given the rare biosphere constituents detected in our environmentally filtered

treatments, we likely added a further 495 taxa, for a total taxa pool of >2000 microbial taxa. After

coalescence, we detected only 472 taxa in the Brackish-Coalescence treatment. While richness

declined roughly to the level of the Marine-Home, evenness was intermediate between the two

endmembers, reflecting a shift in the shape of the dominance diversity curve. Taxa loss was not

symmetric: only 37 of the original Freshwater microbiomes were detected, while 171 Marine-derived

microbial taxa survived. Ten of these 'surviving' taxa were in common, and all ten of these overlapping

taxa were members of the set of 21 taxa found in both original endmember inocula. One quarter (n=126) of the taxa detected in the Brackish-Coalescence treatment were not observed in the Marine-Brackish or Freshwater-Brackish treatments and thus we do not know from which endmember community they were derived. These rare biosphere constituents increased in abundance as a result of interactions between the endmember microbiomes.

The composition of the Brackish-Coalescence community overlapped almost entirely with the Marine endmember community (Fig. 2B). Both the Marine endmember, the Marine-Brackish and the Brackish-Coalescence communities were dominated by *Alteromonas* (~50% of relative abundance) (Fig. 3). Taxa that dominated the Freshwater microbiome were lost (Fig. 3, Fig. 5).). The enzymatic response followed a similar trend: the enzymatic profile for the Brackish-Coalescence microcosms was indistinguishable from the Marine-Brackish and slightly different from the Marine-Home enzyme profile, while significantly different from the enzymes of both the Freshwater-Home and Freshwater-Brackish treatments (Fig. 2C).

Phylogenetic response to Coalescence varies by End Member Community

Taxa that were lost and gained from the Marine microbiome in response to environmental filter and to the biotic filter were closely related (Fig. 5). A sensitive taxon lost due to filtering was typically replaced by an increase in abundance of a tolerant sister taxa in the Marine microbiomes. In contrast, whole classes and orders of the Freshwater microbial community were lost and gained as a result of these two assembly filters. The phylofactorization of microbiomes exposed to the Brackish media identified nine factors, forming six non-overlapping clades capturing a total of 548 species (average 91 species) showed significant changes in the probability of being detected in Brackish water. All but one of these factors show an increase probability of being present relative to other Freshwater microorganisms (Suppl. Fig. 5). In contrast, there were four overlapping Marine clades (from five identified factors) totally 255 species (average 64 species) showing significant changes in detection probability with Brackish exposure (Suppl. Fig. 5).

DISCUSSION

Environmental filters and biotic interactions were both important in determining which Freshwater and Marine microorganisms survived under Brackish common garden conditions. When transplanted from their home environments into Brackish media, both Freshwater and Marine communities lost >70% of

all detectable taxa in the initial inoculum and home conditions. This was somewhat counter balanced

by the emergence of the rare biosphere of each microbiome, accounting for 66% of the Marine

microbiome and 82% of the Freshwater microbiome in the Brackish environment. The rare biosphere

emergence stabilized Marine microbial richness to just over 100% of original richness. In contrast,

Freshwater microbiome richness in Brackish conditions was only 21% of original richness, despite rare

biosphere emergence. This awakening of the rare biosphere during Brackish exposure supports Paver

et al. (2018), who show that certain rare microbial taxa that may possess wider salinity tolerance

('crossing the salty divide') may also be uniquely adapted to proliferating in these new environmental

conditions. The biotic filter also imposed taxonomic shifts for both communities when added in

combination to the Brackish arena. Under this Brackish-Coalescence treatment, a further 70.7% of the

initial Freshwater inocula and 51% of the Marine inocula were not detected. Taken together, these

patterns of major loss of specialists and counter balancing of emerging rare taxa explain the

community convergence on the Marine microbiome.

There were significant differences in the taxonomic richness response of these two

communities to our coalescence experiment. While both endmember communities were significantly

different from their coalescent counterparts, only the Freshwater community had a significant reduction

in taxa richness as a result of environmental filtering (Fig. 2C). This suggests that the Marine

microbiome is, on the whole, more capable of dealing with the intermediate Brackish conditions for two

main reasons: resistance of at least a third of the community to a range of salinity and nutrients, and

resilience due to the high community buffering capacity of the Marine rare biosphere. The prevalence

of Marine-derived rare Brackish tolerant taxa and consequential convergence towards the Marine

microbiome is perhaps expected given the high physiological threshold of marine microbial taxa to a

wide range of salinity (del Giorgio and Bouvier 2002, Wu *et al.* 2006, Herlemann *et al.* 2010). As we

lose sensitive dominant taxa, we sample the rare biosphere more deeply. The rich 'microbial seed

bank' buffers fluctuations in species richness (Lennon and Jones 2011, Jousset *et al.* 2017, Wang *et*

al. 2017), which may help explain the limited functional change in this common garden experiment.

These two unique microbial communities show distinct phylogenetic responses to the

environmental filters imposed by being transplanted into Brackish media. For our Freshwater

community, whole clades turned over. We saw that several sensitive Freshwater clades were lost

while multiple tolerant Freshwater clades become abundant enough to detect. In contrast, for our

saltwater community the turnover was at a finer taxonomic resolution, with a loss and gain of sister

taxa within clades. The composition of experimental replicates was remarkably similar, indicating that

there are real differences in the ability of microbial taxa to survive transplant into altered salinities and

that community composition responses are predictable. Community composition converged under

Brackish conditions because for both endmember microbiomes there was a reservoir of tolerant rare

taxa which increased in abundance when exposed to the intermediate environmental condition. The

shift in composition towards the Marine microbiome resulted from a greater reservoir of tolerant taxa

within that community. The evidence for this is both the compositional shift revealed through ordination

and also the fact that the loss of initially detected taxa is not accompanied by a decline in species

richness for the Marine microbiome.

From a fundamental science perspective, the immense contribution of the rare biosphere to

the Brackish conditions is most fascinating and this rapid turnover of the community somewhat unique

to microorganisms. With <2% of detectable overlap among the original microbiomes, the response to

Brackish conditions converges the microbiomes to much higher taxonomic overlap, with the retention

of the original overlapping taxa and immense emergence of the rare biosphere of each endmember

community. This emergence of the microbial 'seedbank' and the dampened response to aggregate

microbial function is an excellent example of microbial community resilience, where the rare biosphere

plays a pivotal role in ecological rescue.

From an applied microbiology perspective and the natural history of microorganisms, learning

which taxa are lost to environmental and biotic filtering will be instructive and useful for microbial

engineering of 'wild' unculturable microbial taxa (Libby & Silver 2019). There are potential commercial

applications for this experimental setup. The consistency of microbial community composition

responses to experimental treatments suggests that this approach may prove quite useful in

strategically identifying sensitive and tolerant taxa along many different real environmental gradients

(Rocca *et al.* 2019). Over time, such information could lead to the development of microbial sensors, in

which microbial community composition could be used to draw inferences about environmental

conditions. By better understanding the resultant microbial community structure and function when

microbial worlds collide, we may be able to better understand and modulate microbial communities in

areas as wide as agricultural efficiency by microbial consortia (Busby et al 2017), bioremediation

(Baez-Rogelio et al 2016, Sierocinski *et al.* 2017) or biomedical microbial transplants (Gibbons *et al.*

2017).

Our experiment also raises many new fundamental questions about the environmental and

biotic processes that structure the microbiome. Because we observed compositional shifts at very fine

levels of taxonomic resolution (~sister taxa) for our Marine microbiome when exposed to Brackish

conditions, we speculate that biotic interactions are far more important within this community. Why this

might be is an interesting question for microbial ecology.

<u>CONCLUSIONS</u>

Our experimental approach to the study of microbial coalescence provides one of the first

demonstration to directly and separately compare the relative strength of environmental and biotic

filters in structuring intact microbial communities. In the case of mixing Fresh and Marine water, the

Brackish intermediate condition proved to be a very strong environmental filter that had a greater

effect on the Freshwater microbiome than its Marine counterpart. Applying this technique to other

gradients may reveal cases in which biotic interactions dominate. Collectively, the use of this approach

to detect the phylogenetic distribution of sensitivity and tolerance to various environmental gradients is

likely to help us rapidly advance our ecological understanding of why microbial taxa live where they

do.

<u>REFERENCES</u>

Anderson, M.J. (2001). A new method for non-parametric multivariate analysis of variance. *Austral*
436    *Ecol.*, **26**, 32-46.

437 Angel, R., Soares, M.I.M., Ungar, E.D. & Gillor, O. (2009). Biogeography of soil archea and bacteria
438    along a steep precipitation gradient. *ISME J.*, **4**, 553-563.

439 Ardón, M., Helton, A.M. & Bernhardt, E.S. (2016). Drought and saltwater incursion synergistically
440    reduce dissolved organic carbon export from coastal freshwater wetlands. *Biogeochem.*, **127**,
441    411–426.

442 Ardón, M., Morse, J.L., Doyle, M.W. & Bernhardt, E.S. (2016). The Water Quality Consequences of
443    Restoring Wetland Hydrology to a Large Agricultural Watershed in the Southeastern Coastal
444    Plain. *Ecosystems*, **13**, 1060-1078.

445 Baas, D., Stentiford, G.D., Littlewood, D.T.J. & Hartikainen, H. (2015). Diverse applications of
446    environmental DNA methods in parasitology. *Trends Parasitol.*, **31**, 499-513.

447 Baez-Rogelio, A., Morales-Garcia, Y.E., Quintero-Hernandez, V. & Munoz-Rojas, J. (2016). Next
448    generation of microbial inoculants for agriculture and bioremediation. *Micro Biotechnol.*, **10**,
449    19-21.

450 Baldwin, D.S., Rees, G.N., Mitchell, A.M., Watson, G. & Williams, J. (2006). The short-term effects of
451    salinization on anaerobic nutrient cycling and microbial community structure in sediment from
452    a freshwater wetland. *Wetlands*, **26**, 455-464.

453 Barlow, P.M. & Reichard, E.G. (2010). Saltwater intrusion in coastal regions of North
454    America. *Hydrogeol J.*, **18**, 247-260.

455 Bell, C.W., Fricks, B.E., Rocca, J.D., Steinweg, J.M., McMahon, S.K. & Wallenstein, M.D. (2013).
456    High-throughput fluorometric measurement of potential soil extracellular enzyme activities. *J*
457    *Vis Exp.*, **81**, 50961.

458 Bolyen, E., Rideout, J.R., Dillon, M.R., Bokulich, N.A., Abnet, C., Al-Ghalith, G.A. *et al.* (2018).
459    QIIME2: Reproducible, interactive, scalable, and extensible microbiome data science. *PeerJ.*,
460    e27295v1.

Bryant, J.A., Lamanna, C., Morlon, H., Kerkhoff, A.J., Enquist, B.J. & Green, J.L. (2008). Microbes on

mountainsides: Contrasting elevational patterns of bacterial and plant diversity. *Proc Natl*

Acad Sci USA, **105**, 11505-11511.

Burke, V. & Baird, L.A. (1931). Fate of fresh water bacteria in the sea. *J. Bacteriol.*, **21**, 287-298.

Busby, P.E., Soman, C., Wagner, M.R., Friesen, M.L., Kremer, J., Bennett, A. *et al.* (2017). Research

priorities for harnessing plant microbiomes in sustainable agriculture. *PLoS Biol.*, **15**,

e2001793.

Cadotte, M.W. & Tucker, C.M. (2017). Embracing the Nonindependence of the Environmental Filter: A

Reply to Responses. *Trends Ecol Evol.*, **32**, 886-887.

Cahill, J.F., Kembel, S.W., Lamb, E.G. & Keddy, PA. (2008). Does phylogenetic relatedness influence

the strength of competition among vascular plants? *Perspect Plant Ecol Syst.*, **10**, 41-50.

Callahan, B.J., McMurdie, P.J. & Holmes, S.P. (2017). Exact sequence variants should replace

operational taxonomic units in marker-gene data analysis. *ISME J.*, **11**, 2639-2643.

Canfora, L., Bacci, G., Pinzari, F., Lo Papa, G., Dazzi, C. & Benedetti, A. (2014). Salinity and bacterial

diversity: to what extent does the concentration of salt affect the bacterial community in a

saline soil? *PLoS One.*, **9**, e106662.

Caporaso, J.G., Lauber, C.L., Walters, W.A., Berg-Lyons, D., Lozupone, C.A., Turnbaugh, P.J. *et al.*

(2011). Global patterns of 16S rRNA diversity at a depth of millions of sequences per sample.

Proc Natl Acad Sci USA., **108**, 4516–4522.

Comte, J., Berga, M., Severin, I., Logue, J.B. & Lindstrom, E.S. (2017). Contribution of different

bacterial dispersal sources to lakes: Population and community effects in different seasons.

Env Microbiol, **19**, 2391–2404.

Craft, C., Clough, J., Ehman, J., Joye, S., Park, R., Pennings, S. *et al.* (2009). Forecasting the effects

of accelerated sea-level rise on tidal marsh ecosystem services. *Front Ecol Environ*, **7**, 73-78.

del Giorgio, P.A. & Bouvier, T.C. (2002). Linking the physiologic and phylogenetic successions in free-

living bacterial communities along an estuarine salinity gradient. *Limnol Oceanogr.*, **47**, 471-

486.

Edmonds, J.W., Weston, N.B., Joye, S.B., Mou, X. & Moran, M.A. (2009). Microbial community
response to seawater amendment in low-salinity tidal sediments. *Microb Ecol.*, **58**, 558–568.

Ewert, M. & Deming, J.W. (2013). Sea Ice Microorganisms: Environmental Constraints and
Extracellular Responses. *Biology*, **2**, 603-628.

Fierer, N. & Jackson, R.B. (2006). The diversity and biogeography of soil bacterial communities. *Proc*
Natl Acad Sci USA, **103**, 626-631.

Gibbons, S.M., Kearney, S.M., Smilie, C.S. & Alm, E.J. (2017). Two dynamic regimes in the human
gut microbiome. *PLOS Comput Biol.*, **13**, e1005364.

Goberna, M., Navarro-Cano, J.A., Valiente-Banuet, A., García, C. & Verdú, M. (2014). Abiotic stress
tolerance and competition-related traits underlie phylogenetic clustering in soil bacterial
communities. *Ecol Lett.*, **17**, 1191-1201.

Grime, J.P. (1977). Evidence for the existence of three primary strategies in plants and its relevance to
ecological and evolutionary theory. *Am. Nat.*, **111**, 1169-1194.

Hawkes, C.V., Waring, B.G., Rocca, J.D. & Kivlin, S.N. (2017). Historical climate controls soil
respiration responses to current soil moisture. *Proc Natl Acad Sci USA*, **114**, 6322-6327.

Herbert, E.R., Boon, P., Burgin, A.J., Neubauer, S.C., Franklin, R.B., Ardón, M. *et al.* (2015). A global
perspective on wetland salinization: ecological consequences of a growing threat to
freshwater wetlands. *Ecosphere.*, **6**, 1-43.

Herlemann, D.P.R., Labrenz, M., Jürgens, K., Bertilsson, S., Waniek, J.J. & Andersson, A.F. (2011).
Transitions in bacterial communities along the 2000 km salinity gradient of the Baltic Sea.
ISME J., **5**, 1571-1579.

Herlemann, D.P.R., Lundin, D., Andersson, A.F., Labrenz, M. and Jürgens, K. (2016). Phylogenetic
signals of salinity and season in bacterial community composition across the salinity gradient
of the Baltic Sea. *Front Microbiol.*, **7**.

Jackson, C.R. & Vallaire, S.C. (2009). Salinity and nutrients on microbial assemblages in Louisian
wetland sediments. *Wetlands*, **29**, 277-287.

Jackson, D.A. (1995). PROTEST: A PROcrustean randomization TEST of community environment
concordance. *Ecoscience*, **3**, 297-303.

Jousset, A., Bienhold, C., Chatzinotas, A., Gallien, L., Gobet, A., Kurm, V. *et al.* (2017). Where less
517   may be more: how the rare biosphere pulls ecosystems strings. *ISME J*, **11**, 853-862.

518   Lennon, J.T. & Jones S.E. (2011). Microbial seed banks: the ecological and evolutionary implications
519   of dormancy. *Nat Rev Microbiol*, **9**, 119-130.

520   Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F. *et al.* (2004).
521   The metacommunity concept: a framework for multi-scale community ecology. *Ecol Lett.*, **7**,
522   601-613.

523   Libby, E.A. & Silver, P.A. (2019). Harnessing undomesticated life. *Nat Rev Microbiol.*, **4**, 212-213.

524   Livingston, G., Jiang, Y., Fox, J.W. & Leibold, M.A. (2013). The dynamics of community assembly
525   under sudden mixing in experimental microcosms. *Ecology*, **94**, 2898–2906.

526   Lozupone, C.A. & Knight, R. (2017). Global patterns in bacterial diversity. *Proc Natl Acad Sci USA*,
527   **104**, 11436-11440.

528   Luef, B., Frischkorn, K.R., Wrighton, K.C., Holman, H.Y.N., Birarda, G., Thomas, B.C. *et al.* (2015).
529   Diverse uncultivated ultra-small bacterial cells in groundwater. *Nat Commun*, **6**, 6372.

530   Mansour, I., Heppel, C.M., Ryo, M. & Rillig, M.C. (2018). Application of the microbial community
531   coalescence concept to riverine networks. *Biol Rev.*, **93**, 1832-1845.

532   Mirarab, S., Nguyen, N., Guo, S., Wang, L., Kim, J. & Warnow, T. (2014). PASTA: Ultra-Large Multiple
533   Sequence Alignment for Nucleotide and Amino-Acid Sequences. *J Comp Biol*, **22**, 377-386.

534   Nagler, M., Podmirseg, S.M., Griffith, G.W., Insam, H. & Ascher-Jenull, J. (2018). The use of
535   extracellular DNA as a proxy for specific microbial activity. *Appl Microbiol Biotech.*, **102**, 2885-
536   2898.

537   Nielsen, D.L., Brock, M.A., Rees, G.N. & Baldwin, D.S. (2003). Salinity and bacterial diversity: to what
538   extent does the concentration of salt affect the bacterial community in a saline soil? *Aust J*
539   *Bot.*, **51**, 655-665.

540   Neubauer, S.C., Franklin, R.B. & Berrier, D.J. (2013). Saltwater intrusion into tidal freshwater marshes
541   alters the biogeochemical processing of organic carbon. *Biogeosci.*, **10**, 8171-8183.

Oksanen, J., Blanchet, F.G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D. *et al.* (2017). Vegan:

Community Ecology Package. R package version 2.4-3.

Parte, A.C. (2013). LPSN – list of prokaryotic names with standing in nomenclature. *Nucleic Acids*

Research, **42**, D613-D616.

Paver, S.F., Muratore, D.J., Newton, R.J. & Coleman, M.L. (2018). Re-evaluating the salty divide:

phylogenetic specificity of transitions between marine and freshwater systems. *mSystems.*, **3**,

e00232-18.

Pembrey, R.S., Marshall, K.C. & Schneider, R.P. (1999). Cell surface analysis techniques: What do

cell preparation protocols do to cell surface properties? *Appl Environ Microbiol*, **65**, 2877-

2894.

Peterson, B.W., Sharma, P.K., van der Mei, H.C. & Busscher, H.J. (2012). Bacterial cell surface

damage due to centrifugal compaction. *Appl Environ Microbiol*, **78**, 120-125.

Quast, C., Pruesse, E., Yilmaz, P., Gerken, J., Schweer, T. & Yarza, P. *et al.* (2013). The SILVA

ribosomal RNA gene database project: improved data processing and web-based tools.

Nucleic Acids Res., **41**, D590-D596.

R Development Core Team. (2010). R: a Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Rillig, M.C., Antonovics, J., Caruso, T., Lehmann, A., Powell, J.R., Veresoglou, S.D. *et al.* (2015).

Interchange of entire communities: microbial community coalescence. *Trends Ecol Evol.*, **8**,

470-476.

Rillig, M.C. & Mansour, I. (2017) Microbial Ecology: Community Coalescence Stirs Things Up. *Curr*

Biol., **27**, R1268-R1286.

Rocca, J.D., Simonin, M., Blaszczak, J.R., Ernakovich, J.G., Gibbons, S.M., Midani, F.S., *et al.* (2019).

The Microbiome Stress Project: Toward a Global Meta-Analysis of Environmental Stressors

and Their Effects on Microbial Communities. *Front Microbiol.*, **9**, 3272.

Sierocinski, P., Milferstedt, K., Bayer, F., Alston, M., Swarbreck, D., Hobbs, P.J. *et al.* (2017). The

most efficient microbial community dominates during community coalescence. *Curr Biol.*, **27**,

3390–3395.

570 Siles, J.A. & Margesin, R. (2016). Abundance and Diversity of Bacterial, Archaeal, and Fungal

571   Communities Along an Altitudinal Gradient in Alpine Forest Soils: What Are the Driving

572   Factors? *Microb Ecol.*, **72**, 207-220.

573 Sokal, R. & Michener, C. (1958). *A statistical method for evaluating systematic relationships*. Univ

574   Press of Kansas, 38: 1409-1438.

575 Souffreau, C., Pecceu, B., Denis, C., Rummens, K. & Meester, L.D. (2014). An experimental analysis

576   of species sorting and mass effects in freshwater bacterioplankton. *Freshwater Biol.*, **10**,

577   2081-2095.

578 Vellend, M. (2010). Conceptual synthesis in community ecology. *Q Rev Biol.*, **85**, 183-206.

579 Wang, Y., Hatt, J.K., Tsementzi, D., Rodriguez-R, L.M., Ruiz-Pérez, C.A., Weigand, M.R. *et al.* (2017).

580   Quantifying the Importance of the Rare Biosphere for Microbial Community Response to

581   Organic Pollutants in a Freshwater Ecosystem. *Appl Environ Microbiol.*, **83**, e03321.

582 Washburne, A.D., Silverman, J.D., Leff, J.W., Bennett, D.J., Mukherjee, S. *et al.* (2017). Phylogenetic

583   factorization of compositional data yields lineage-level associations in microbiome datasets.

584   *PeerJ*, **5**, e2969.

585 Washburne, A.D., Silverman, J.D., Morton, J.T., Becker, D., Crowley, D., Mkherjee, S. *et al.* (2019).

586   Phylofactorization – a graph partitioning algorithm to identify phylogenetic scales of ecological

587   data. *Ecol Monogr.*, in press (accepted).

588 Webb, S.D. (1976). Mammalian Faunal Dynamics of the Great American Interchange. *Paleobiology*, **2**,

589   220–234.

590 Weston, N.B., Dixon, R. & Joye, S.B. (2006). Microbial and geochemical ramifications of salinity

591   intrusion into tidal freshwater sediments. *J. Geophys Res: Biogeosci.*, **111**, G01009.

592 Whittaker, R.H. (1965). Dominance and diversity in land plant communities. *Science*, **147**, 250-260.

593 Wu, Q.L., Zwart, G., Schauer, M., Kamst-van Agterveld, M.P. & Hahn, M.W. (2006). Bacterioplankton

594   community composition along a salinity gradient of sixteen high-mountain lakes located on the

595   Tibetan Plateau, China. *Appl Environ Microbiol.*, **72**, 5478-5485.

596 Yang, Y., Gao, Y., Wang, S., Xu, D., Yu, H., Wu, L. *et al.* (2014). The microbial gene diversity along an

597   elevation gradient of the Tibetan grassland. *ISME J.*, **8**, 430-440.

TABLE & FIGURE LEGENDS

Figure 1. (I.) Potential outcomes of coalescing communities relative to "home" conditions.

Ellipsoids represent the theoretical variation in community structure or function among experimental

replicates (symbols). Communities may be symmetrically impacted by coalescence (A,B), constrained

primarily by environment (A), or resistant to environmental change (B); (C) asymmetrical impact may

be driven by either environment or community; or (D) coalescence may result in stochastic shifts in

community structure or function. **(II.) Conceptual framework to disentangle the influence of**

assembly processes: components needed for each filter (left), experimental setup (center), and

hypothetical outcomes (right). Top panel: assessing the impact of the environmental filter, where no

impact of altered environment (Ho), versus a shift in community structure (Ha), reveals

environmentally sensitive and generalist taxa. Bottom panel determines the impact of novel biotic

interactions, with no impact (Ho) where the sum of Part 1 communities equals that of the coalesced

community, versus new biotic interactions, where taxa are lost due to competition or gained due to

mutualistic interactions (Ha).

Table 1. Location, chemical and microbial bulk characterization of the two distinct aquatic endmember

habitats in Coastal North Carolina, USA: Freshwater and Marine environments and inocula; and the

chemical characterization of the experimentally mixed (1:1) Brackish environment and blended

inoculum.

Figure 2. Impact of coalescence on microbial community structure and function: (A) alpha

diversity of each treatment for SV richness (left) and Pielou's Evenness (right), (B,C) nonmetric

multidimensional scaling (NMDS) of (B) bacterial community structure; and (C) microbial extracellular

enzyme activity. The NMDS arrows indicate movement through the filters relative to endmember

control conditions, weighted by significance from Adonis (solid = significant change; dotted = NS). No

community or functional data is available for one Brackish-Coalescent sample that was damaged

during the incubation (n=4, instead of experiment wide n=5).

Figure 3. Responses of the most abundant bacteria to community coalescence. Heatmap of the most prevalent bacterial SVs (>0.5% relative abundance among all microcosms), vertically clustered by SV, with corresponding taxonomy (shaded by salt tolerance) and horizontally by microcosm, identified by Environment (top row symbols) and Inoculum (bottom row). Blue and green flanked regions identify the "home" conditions for Freshwater and Marine conditions. Ranking estimated using standard UPGMA methods based on Bray-Curtis pairwise dissimilarities in Vegan R package.

Figure 4. Distribution of community origin due to each assembly filter: microbial taxa lost and gained due to Brackish exposure (Part 1: Impacts of Environmental Filtering); and due to microbiome merging with the Brackish-Coalescence treatments (Part 2: Biotic Filtering In Newly Merged Community). The numbers inside or above each box represent the number of taxa (SVs) found among the microcosms of that treatment (n=5), with Freshwater in "white" and Marine in "Black" boxes. The numbers adjacent to the lower boxes represent the number of taxa detected in the original communities (solid) or detected as rare taxa emergence (striped), and the gray box represents rare emergence of unknown endmember origin.

Figure 5. Community phylogenetic response to environmental filtering and to novel biotic interactions for Freshwater (left) and Marine (right) microbiomes. The _inner ring_ shows the distribution of microbial taxa present in "home" conditions; _middle ring_ represents the distribution of microbial taxa surviving the Brackish environment; and the _outer ring_ displays the microorganisms of known origin surviving to Brackish-Coalescence. The gray boxes highlight phylogenetic regions where the Freshwater and Marine microbiomes exhibit distinct dispersion signals in response to the assembly filters. _The bacterial phyla are labeled inside the rings, with the following abbreviations 'Actino': Actinobacteria; 'Arm': Armatimonadetes; 'Planc': Planctomycetes; 'Verru': Verrucomicrobia; 'Ac': Acidobacteria; 'Cy': Cyanobacteria; 'D': Dependentiae._

SUPPLEMENTARY INFORMATION

Supplemental Figure 1. Laboratory Experimental Incubation Setup. (Top) Reciprocal manipulation of communities into 'home', 'away' and 'mixed' environments, with and without community mixing. (Bottom) Field controls of: (Left) Intact communities with two levels of filtration to assess microbes-only and microbes + small predators; (Middle) Sterile environments of same filtration; (Right) Water-only negative controls. Replication is five microcosms per treatment.

Supplemental Figure 2. Nonmetric multidimensional scaling (NMDS) of bacterial community structure, demonstrating the minimal impacts of inoculum preparations and autoclaving environments on microbial community structure relative to unaltered positive control microcosms.

Supplemental Figure 3. 16S rRNA-based phylogeny of bacteria in the starting inocula. The inner ring shows SVs in the Freshwater inocula (green) and outer ring represents the Marine inocula (blue). Black arrows indicate the 21 overlapping SVs.

Supplemental Figure 4. Microbial function of Coalescing Microbial Communities vs. End member Controls. Barcharts of potential extracellular enzyme activity: (A) Aryl-sulfatase, (B) 1,4 Beta-glucosidase, (C) 1,4 N-acetylglucosidase, (D) Alkaline Phosphotase, (E) 1,4 Alphaglucosidase, (F) 1,4 Cellobiohydrolase, (G) D-xylosidase, and (H) L-leucine-aminopeptidase. Compact letter display represents pairwise comparisons among each of the five treatments, dotted lines denote the average of the end-point controls (Freshwater and Marine), and error bars display ±SE of each enzyme.

Supplemental Figure 5. Phylofactorization results for Brackish exposure on the Freshwater (Left) and Marine (Right) microbiomes. Top panels show the non-overlapping clades identified as significantly factors based on a Holm's sequentially rejective 5% FWER cutoff, and bottom panels represent the corresponding phylogenetic location for each of these factors (6 in Freshwater, 4 in Marine).

I. Hypothetical outcomes of coalescing community structure or function relative to 'home' conditions.

A. Species Sorting

B. Community Resistance

Environment	Community
Env 1 ⬭	Comm 1 ○
New Env ⬭	Mixed ◗
Env 2 ⬬	Comm 2 ●
Conditions	
● Home	▲ Changed

C. Environment or Community Convergence

D. Stochasticity

NMDS 1

NMDS 1

II. Approach to *directly* parse the contribution of community assembly mechanisms in community coalescence

Part 1: Isolating Community Impacts of the Environmental Filter

Components	*Experimental Setup*	*Hypothetical Outcomes*

Environments

Community origin

Ho: No impact of environment on community structure

Ha: Taxa lost due to new environment

Env 1 Specialists

Generalists

Env 2 Specialists

Generalists

Part 2: Identifying Biotic Interactions in Coalescing Community

Components	*Experimental Setup*	*Hypothetical Outcomes*

Environments

Community origin

Mixed Community

Ho: No loss/gain of taxa from novel community interactions

Ha: Taxa lost/gained due to novel biotic interactions

(-) Biotic Interactions

Competitors

(+) Biotic Interactions

Final Community

Environment	Freshwater	Brackish	Marine
Endpoint Location	Timberlake Restoration	Lab mixed 50:50 by volume	Cape Hatteras NS
Latitude,	35°53'46.4"N,	—	35°49'57.4"N,
Longitude	76°09'51.4"W	—	75°33'25.7"W
Mixing	Episodic/Storm	—	Turbulent
Temp (June '16)	23-25 °C	—	23 °C
Chemistry of Environments			
pH	4.4 (± 0.05)	7.8 (± 0.01)	8.2 (± 0.01)
Salinity (PSU)	0.07 (± 0.01)	26.7 (± 0.9)	51.8 (± 0.4)
Dissolved Organic Carbon (mg/L)	33.9 (± 1.2)	21.7 (± 3.2)	3.6 (± 0.4)
Dissolved Inorganic Nitrogen (mg/L)	0.92 (± 0.01)	0.61 (± 0.05)	0.12 (± 0.03)
Microbial Inoculum Characterization			
Biomass (ng DNA/μL)	749.7	584.8	218.3

A.

SV Richness: 300, 200, 100, 0

Fresh Home — *c*

Brackish: Fresh *a*, Coal *b*, Marine *b*

Marine Home — *b*

Evenness: 0.100, 0.075, 0.050, 0.025

Fresh Home — *b*

Brackish: Fresh *a*, Coal *ab*, Marine *a*

Marine Home — *a*

B.

NMDS 2 (Community)

NMDS 1 (Community)

stress = 0.04

C.

NMDS 2 (Function)

NMDS 1 (Function)

stress = 0.07

Community: Fresh ○ Brackish ◔ Marine ●

Environment: Home ● Changed ▲

Freshwater Community

Proteobacteria
Actino
Arm
Planc
Verru
Ac
Bacteroidetes
Cy
D

Freshwater Home

Freshwater - Brackish

Brackish - Coalescence

Marine Community

Proteobacteria
Actino
Arm
Planc
Verru
Ac
Bacteroidetes
Cy
D

Marine Home

Marine - Brackish

Brackish - Coalescence