An integrative approach identifies direct targets of the late viral transcription complex and

an expanded promoter recognition motif in Kaposi's sarcoma-associated herpesvirus

Divya Nandakumar [a, b] and Britt Glaunsinger [a, b]

[a] Department of Plant & Microbial Biology, University of California Berkeley, CA, USA

[b] Howard Hughes Medical Institute, Berkeley, CA, USA

Corresponding Author: glaunsinger@berkeley.edu

Abstract

The structural proteins of DNA viruses are generally encoded by late genes, whose

expression relies on recruitment of the host transcriptional machinery only after the onset of viral

genome replication. β and γ-herpesviruses encode a unique six-member viral pre-initiation

complex (vPIC) for this purpose, although how the vPIC directs specific activation of late genes

remains largely unknown. The specificity underlying late transcription is particularly notable

given that late gene promoters are unusually small, with a modified TATA-box being the only

recognizable element. Here, we explored the basis for this specificity using an integrative

approach to evaluate vPIC-dependent gene expression combined with promoter occupancy

during Kaposi's sarcoma-associated herpesvirus (KSHV) infection. This approach distinguished

the direct and indirect targets of the vPIC, ultimately revealing a novel promoter motif critical for

KSHV vPIC binding. Additionally, we found that the KSHV vPIC component ORF24 is

required for efficient viral DNA replication. Together, these results identify an elusive element

that contributes to vPIC specificity and suggest novel links between KSHV DNA replication and

late transcription.

Author summary

Gene expression in DNA viruses often occurs in temporal waves, with expression of

essential structural proteins occurring late in infection, after viral genome replication has begun.

Strategies underlying expression of these viral late genes are often sophisticated; for example,

the β- and γ-herpesviruses encode a six-component viral complex that directs late gene

transcription, largely by unknown mechanisms. Here, we evaluated how this complex

specifically recognizes late promoters during infection with the oncogenic human γ-herpesvirus

Kaposi's sarcoma-associated herpesvirus (KSHV). We found that one of the components of the

late transcription complex was required for robust viral DNA replication, suggesting new links

between KSHV replication and transcription. Combined measurements of late gene expression

and promoter occupancy then revealed which KHSV genes are directly controlled by the late

gene transcription complex, leading to identification of a key new regulatory element in KSHV

late promoters. Together, these data help explain how the late gene transcription complex is able

to bind seemingly minimal promoters with high specificity, ensuring robust expression of viral

factors necessary for assembly of progeny virions.

Introduction

A conserved feature of double-stranded DNA viruses is that they possess a set of genes that

become strongly activated only following viral DNA replication (1). These "late genes"

generally encode factors involved in viral assembly that are essential for virion production. The

mechanisms underlying their transcriptional regulation are often non-canonical or distinct from

those involved in activating either cellular genes or early viral genes. For example, baculoviruses

encode a multisubunit, α-amanitin-resistant DNA-directed RNA polymerase dedicated

specifically to the activation of compact late promoters (2). Similarly, bacteriophage T7 encodes

its own late gene specific RNA polymerase that recognizes unique promoter elements to

transcribe late genes (3,4). Adenoviruses regulate late gene expression using a complex

combination of viral and cellular factors; beyond promoter activation, these also regulate

transcription termination and contribute to differential posttranscriptional processing of late viral

mRNA (5).

Several studies suggest that the expression of late genes is also uniquely regulated in β- and

γ-herpesviruses, which include human cytomegalovirus (HCMV), Epstein-Barr virus (EBV), and

Kaposi's sarcoma-associated herpesvirus (KSHV) (6). Unlike the eukaryotic viruses described

above, β and γ herpesviruses do not engage cellular TATA binding protein (TBP) during late

gene pre-initiation complex formation. Their core late gene promoters are unusually small (12-15

base pairs (bp)), with the only recognizable conserved element being a modified TATA box

(TATT) ~30 bp upstream of the transcription start site (TSS) (7–9). This is in contrast to

eukaryotic and most early herpesviral promoters, which have multiple promoter elements that are

important for promoter recognition and activation by RNA polymerase II (RNA pol II) (10).

Another notable feature is that activation of these minimal promoters requires a virally-encoded

six-protein complex called the viral Pre-Initiation Complex (vPIC) that coordinates mammalian

RNA pol II (11,12). The vPIC is presumed to be uniquely adapted to activate minimal promoters,

yet the mechanisms underlying its activity and specificity remain largely unknown.

The core component of the KSHV vPIC is ORF24 (BcRF1 in EBV and UL87 in HCMV), a

viral TBP mimic that displays putative structural similarity to cellular TBP (13–15). Unlike

cellular TBP, the viral TBP mimics have expanded functionality in that they bind RNA pol II in

addition to promoter DNA, making them unique among transcription factors that function in

eukaryotic cells. Beyond ORF24, the other essential components of the KSHV vPIC are ORFs

18, 30, 31, 34, and 66, but little is known about how each of these factors contributes to vPIC

activity (14,16–18). ORF34 interacts with multiple components of the vPIC, and is thus thought

to function as a scaffolding factor (12). The ORF18 homolog in HCMV (pUL79) has been

shown to function as an elongation factor, although whether it plays a similar role in KSHV and

EBV is unclear (19). Despite the paucity of functional information on the individual vPIC

components, protein-protein interaction mapping has provided insight into the overall complex

organization (12,18). Furthermore, point mutants that disrupt individual inter-subunit vPIC

interactions abrogate late gene transcription, highlighting the fact that vPIC complex integrity is

critical for function (12,20,21).

The mechanistic basis for promoter selectivity has also been challenging to grasp in the

context of the minimal, largely nondescript nature of late gene promoters. While a modified

TATA box with a TATT sequence is enriched in several late promoters of KSHV, MHV68, and

EBV (7–9,22), the ORF24 TBP mimic shows partial binding to a mutant promoter in which the

TATT sequence is modified to resemble a canonical TATA box (7). However, ORF24 does not

bind promoters that inherently have a canonical TATA box (7), which suggests that other DNA

elements surrounding the TATT sequence in late gene promoters contribute to ORF24 binding or

vPIC assembly on the DNA.

One possible reason that additional DNA elements involved in vPIC promoter recognition

have not been identified is that we do not yet know the subset of viral promoters under the direct

influence of the vPIC. Techniques such as RNA Sequencing (RNA Seq) and microarrays have

helped define the set of genes that are expressed with late kinetics and in a manner dependent on

viral DNA replication (23–26). Furthermore, viral mutagenesis and knockdown studies have

revealed viral genes whose expression is impaired in the absence of vPIC components

(17,18,22,27,28). However, these assays cannot distinguish direct vPIC targets from indirect

targets, such as genes regulated by factors that are themselves dependent on the vPIC. Inclusion

of such indirect targets in the pool of vPIC-regulated promoters could confound the identification

of conserved elements important for direct vPIC regulation.

Here, we integrated RNA Seq with chromatin immunoprecipitation high-throughput

sequencing (ChIP Seq) for vPIC components during KSHV infection to distinguish genes whose

expression is directly versus indirectly controlled by the vPIC. In doing so we made three key

findings that provide insight into late gene promoter activation in KSHV. First, we defined the

set of direct vPIC promoter targets within the KSHV genome and showed that other promoters

are indeed nonproductively bound or only indirectly regulated by the vPIC. Second, by

restricting our analysis to these direct targets, we revealed a novel promoter motif that

contributes significantly to vPIC binding and thus provides insight into features governing vPIC

specificity. Notably, this motif did not emerge when the indirect vPIC targets were included in

the analysis. Finally, we found that the KSHV TBP mimic ORF24 is required for efficient viral

DNA replication. This was unexpected given that the other studied vPIC components have been

shown to be dispensable for this process, and may have implications for the connection between

viral DNA replication and late gene transcription.

Results

KSHV ORF24 is essential for late gene expression and contributes to viral DNA replication

Three leucine residues in the N-terminal domain of KSHV ORF24 (amino acids 73-75) are

essential for its interaction with RNA Pol II and its ability to promote late gene transcription

(14). Given that most viral proteins are multifunctional, we engineered the ORF24$_{RLLLG->RAAAG}$

mutation (ORF24$_{RAAAG}$) into KSHV BAC16 to precisely evaluate the impact of the

transcriptional function of ORF24 during lytic infection. We also generated the corresponding

mutant rescue (ORF24$_{RAAAG}$-MR) to ensure that any observed phenotypes were not the effect of

secondary mutations in the BAC. Identical banding patterns from *Rsr*II digestion of WT and

mutant BAC16 DNAs verified that there were no gross rearrangements of the viral genome

during mutagenesis (Figure S1-lanes1-3). We established stable cell lines latently infected with

WT, ORF24$_{RAAAG}$, and ORF24$_{RAAAG}$-MR KSHV in the iSLK background, which contains a

doxycycline-inducible version of the KSHV lytic transactivator protein ORF50 (RTA) to enable

lytic reactivation (29). During lytic reactivation, these three viruses expressed similar levels of

the early protein ORF59, while the ORF24$_{RAAAG}$ mutant was selectively unable to express the

K8.1 late protein or produce infectious virions (Fig 1A, 1B). These findings are consistent with

previous complementation data (14), and confirm the importance of the ORF24-Pol II interaction

for late gene expression.

Surprisingly, we also observed ~6-fold defect in DNA replication in the ORF24$_{RAAAG}$ mutant

(Fig 1C), which was unexpected given that previous studies with mutations in ORF24 or other

components of the vPIC did not show this defect (12,14,16–18). This defect was rescued in the

ORF24$_{RAAAG}$-MR cell line, indicating that it was not due to secondary mutations in the BAC

(Fig. 1C). However, to independently confirm this observation, we engineered two additional

ORF24 viral mutants: the point mutant ORF24$_{R328A}$, which abrogates its essential interaction

with the ORF34 component of the vPIC (12), and a stop mutant (ORF24$_{Stop}$) that has a premature

stop codon in the ORF24 coding sequence. Inclusion of the ORF24$_{Stop}$ virus allowed us to

evaluate whether the point mutants were functioning in a dominant negative manner, for example

through nonproductive DNA binding. Finally, we also N-terminally tagged endogenous WT ORF24 with HA (HA-ORF24) to assess whether any alterations to this locus impacted viral DNA replication. Notably, we observed a similar ~5-6-fold defect in DNA replication in the ORF24$_{RAAAG}$, ORF24$_{R328A}$, and ORF24$_{Stop}$ viruses, which was restored to WT levels in the ORF24$_{RAAAG}$ and ORF24$_{R328A}$ MR viruses. Moreover, there was no DNA replication defect in the HA-ORF24 virus (Fig 1C), demonstrating that the defect in DNA replication is not a consequence of manipulating the ORF24 locus but is specific to disrupting its role in transcription.

Specific gene clusters show differential dependence on the ORF24-Pol II interaction

To more comprehensively analyze how the ORF24-Pol II interaction impacts gene expression during infection, we deep sequenced RNA from three replicates each of iSLK cells infected with the ORF24$_{RAAAG}$ or the ORF24$_{RAAAG}$-MR virus. Samples were sequenced at 24 and 48 h post lytic reactivation with doxycycline and sodium butyrate, as these times captured predominantly early (24 h) versus both early and late (48 h) viral gene expression (Figure S2). Given that overlapping transcripts are pervasive in the KSHV genome (22,30), we calculated the primary transcript counts as described in Bruce et al (30) to accurately quantify transcript levels.

Differential expression analysis of genes in infected cells containing WT ORF24 (ORF24$_{RAAAG}$ MR) confirmed that the majority of viral genes are upregulated between 24 and 48 h post reactivation (Figure 2A), and those that are most strongly upregulated are predominantly classified as late genes (shown in blue). Conversely, there was a generalized reduction in transcript levels at the 48 h time point in cells containing KSHV ORF24$_{RAAAG}$ compared to MR, with the most strongly downregulated genes being those classified as late genes (Figure 2B).

Combining the above ORF24$_{RAAAG}$ mutant and MR induction data into a heatmap confirmed

that in general, genes with later expression kinetics tended to be more severely impacted by the

ORF24$_{RAAAG}$ mutation compared to genes with early expression kinetics (Figure 2C). (The fold

change data was scaled to allow for easy visualization). However, not all genes fit this model.

Some genes categorized as early based on their induction pattern showed a moderate dependence

on the ORF24-Pol II interaction (e.g. ORFs 9, 40, 34, 41, 16 and 37). Similarly, a subset of genes

that show late kinetics were less dependent on the ORF24-Pol II interaction (e.g. ORFs 67A, 62,

K4.2A, 64, K7, 54, 67, 56, 66, K4.2 and vIRFs-2, 3 and 4). Classification as early or late and

dependence on the ORF24-Pol II interaction was determined using mean log$_2$ fold change.

Together, these data suggest that not all kinetic late genes are regulated by the vPIC and there

may be subset of early genes that are direct or indirect targets of the vPIC.

The TATTWAA motif is enriched in the promoters of genes whose expression is linked to

the ORF24-Pol II interaction

We then looked for motifs enriched in the promoters of genes most strongly

downregulated in the ORF24$_{RAAAG}$ mutant using Multiple EM for Motif Elicitation (MEME)

(31), a program that discovers un-gapped motifs that are enriched in a given set of sequences.

We defined the promoter sequence as 100 bp upstream from the transcription start site (TSS) or

300 bp upstream from the translation start site when the TSS was unknown. Analysis of

promoter sequences from 14 of the most strongly down-regulated genes (Log$_2$ fold change

greater than one standard deviation from the mean) showed that TATTWAA is the only motif

that is enriched in this set of sequences (Figure 2D).

We next monitored for the occurrence of this motif across all promoters in the viral

genome using Find Individual Motif Occurrences (FIMO) (32), which scans a set of sequences

for matches to motifs provided by the user. 23 of 83 viral promoters were found to have the

TATTWAA motif with a high score, of which 16 promoters had the motif ~30 bp upstream from

the TSS (Figure 4 and Table 1), consistent with the location of the canonical TATA box in

eukaryotic promoters. Of the remaining seven promoter sequences, the TSS of six of the

promoters is not known (shown in red in Table 1); hence, the exact position of the TATTWAA

motif from the TSS cannot be estimated. ORF25 is the only exception, where the TATTWAA

sequence is not present ~30 bp upstream from the TSS, although it does have a canonical TATA

box (TATAAAA) at that position.

Notably, a subset of the genes containing the TATTWAA motif showed less than average

repression by the ORF24$_{RAAAG}$ mutation (ORF55, ORF64, ORF66, ORF59, K4.1, ORF4, ORF8,

ORF58, ORF26 and ORF62), suggesting that the motif may not be sufficient to mark regulation

by the vPIC, which is consistent with a similar observation in MHV68 (7). Furthermore, several

genes (ORF28, ORF30, K15, ORF31 and ORF75) were strongly affected by the mutation (Log$_2$

fold change greater than one standard deviation from the mean) but lack the TATTWAA motif.

The number of genes in this set increases to 26 if we broaden the criteria to include all the genes

that show greater than average repression in the presence of the mutation. These genes could

either be down regulated by an indirect mechanism in the absence of the ORF24-Pol II

interaction or could be bound by the vPIC in a manner independent of the TATT motif.

Chromatin immunoprecipitation reveals vPIC occupancy across the KSHV genome

Our RNA Seq experiments identified KSHV genes whose expression is influenced by the

transcription regulatory function of ORF24. However, RNA Seq does not distinguish genes that

are directly bound by the vPIC from those that are indirect transcriptional targets (e.g. controlled

by other factors whose expression is induced by ORF24). We therefore sought to identify

promoters bound by the KSHV vPIC complex using ChIP Seq. Previous studies suggest that

ORF34 acts as a scaffolding factor that brings together all components of the vPIC (12)

including ORF24, which binds both RNA Pol II and TATT-containing DNA. We reasoned that

ORF34 promoter occupancy would therefore be a good proxy for vPIC assembly, and

additionally would avoid potential pitfalls associated with any non-productive or independent

interactions between ORF24 and DNA.

To facilitate ChIP Seq, we added an N-terminal HA tag to endogenous ORF34 in KSHV

BAC16 (HA-ORF34), and generated a stable, doxycycline-inducible iSLK cell line containing

this tagged virus as previously described. HA-ORF34 produced WT levels of infectious virions

in a supernatant transfer assay (Figure S3A), indicating that the tag does not disrupt ORF34

activity. The integrity of the BAC DNA was verified by *Rsr*II digestion (Figure S1-Lane7). We

performed the ChIP Seq experiment in duplicate at 48 h post reactivation to ensure active late

gene expression. Reactivated iSLK cells with untagged BAC16 served as a negative control for

the immunoprecipitation. The individual ChIP peaks for the two replicates as well as the

respective inputs and controls are shown in Figure S3B and supplementary file 1. The ChIP Seq

data showed that HA-ORF34 bound several locations across the viral genome (Figure 3A) but

was not detectably associated with the host genome. A peak calling algorithm (HOMER) (33)

identified 25 common peaks in the viral genome across two replicates that were significantly

enriched in HA-ORF34 relative to untagged WT or input DNA (Supplementary File 1 and Figure

S3B). Notably, 20 of the 25 peaks were found upstream of ORFs, with ~80% of the peaks

centered less than 200 bp from the closest TSS (Supplementary File 1). Given the link between

DNA replication and late transcription in herpesviruses, it was particularly interesting that two of

the most prominent peaks are upstream of the two minimal lytic origins of replication in the

KSHV genome (Figure 3A). One of the peaks could not be uniquely assigned to a single ORF

and was not considered further. Of the remaining 19 ORFs associated with peaks, 13 are

involved in packaging, capsid assembly (or form part of capsid itself), 4 are involved in immune

evasion and 2 in DNA replication (Table 2). While several of them have also been previously

classified as late genes (12/20), it is notable that 6 of them have been classified as early genes,

again suggesting that not all targets of the vPIC are late genes.

To confirm that the observed binding pattern was representative of vPIC assembly, we

independently validated several prominent ChIP Seq peaks by ChIP-qPCR from cells infected

with KSHV containing HA-tagged ORF24 (HA-ORF24). Similar to HA-ORF34, the HA-ORF24

virus produced WT levels of infectious virions (Figure S3A). Indeed, the HA-ORF24 ChIP-

qPCR data were consistent with the HA-ORF34 ChIP-Seq data (Figure 3B), with peaks showing

higher percent input values compared to control regions (ORF64 and ORF37 promoters).

A MEME analysis of the promoters associated with the peaks again showed that the only

motif enriched in these promoters is TATTWAARV (Figure 3C). 17 of the 19 promoters (except

ORF45 and ORF68) have this motif and in a majority of the sequences, it is present ~30 bp from

the TSS (Table 2). Two key points can be drawn from the ChIP seq data. First, several genes

whose promoters contain a TATTWAA motif do not show ORF34 binding (e.g. ORFs 17, 32,

22, 55, 58, 64 and 66). Second, while most promoters bound by ORF34 have a TATTWAA

motif, there are some exceptions (ORF45, ORF68 and ORF53). These observations indicate that

while the TATTWAA motif is enriched in promoters regulated by the vPIC, it is not sufficient to

ensure binding of this specialized transcription complex.

Thirteen KSHV genes are direct targets of the vPIC

Combining the RNA Seq and ChIP Seq data, we defined the direct targets of the vPIC as

genes that are both downregulated in the absence of the ORF24-Pol II interaction and show

ORF34 binding in the promoter region. By this measure, a total of 13 genes are direct targets of

the vPIC (Figure 4 and Table 2 – gray rows). Of the 13 direct targets, 11 code for structural

proteins or proteins involved in viral egress while two are tegument proteins involved in immune

evasion. Genes whose expression was impaired in the KSHV ORF24$_{RAAAG}$ mutant but lacked

ORF34 binding at their promoters are likely to be indirect targets of the vPIC (Figure 4A).

Finally, it is interesting to note that not all genes that have a ChIP peak are down regulated in the

RNA Seq data (e.g. ORFs 4, 6, K4.1, 59, 62 and 68). These could be false positive ChIP peaks

and/or have alternate mechanisms of regulation that ensure their transcription even in the

absence of the vPIC.

The direct vPIC targets reveal an expanded promoter motif

A defining feature of characterized late gene promoters is that they are minimalistic, with

no conserved motifs beyond the TATTWAA signature recognized by ORF24. Yet, our data

indicate that the TATTWAA motif is insufficient to confer activation by the late gene

transcription complex. We hypothesized that the inclusion of potentially indirect vPIC targets

into prior motif searches may have hindered identification of other regulatory sequences within

these promoters. Indeed, repeating the MEME analysis on the 13 direct targets revealed that an

additional 5 bp motif (RVNYS) downstream of the TATTWAA motif was enriched in these

promoter sequences (Figure 4B).

To explore the contribution of the 5 bp region downstream of the TATTWAA sequence

for regulation by the vPIC, we used a plasmid-based promoter activation assay in which

expression of a luciferase reporter was driven by either the K8.1 late or ORF57 early promoter

(Figure 5A). The KSHV left lytic origin of replication (ori-lyt) was included on the plasmids, as

data from MHV68, KSHV and EBV indicate that the presence of ori-lyt in *cis* significantly

boosts late gene promoter activation during infection (7,9,34). Each plasmid was transfected

together with a plasmid constitutively expressing renilla luciferase (to normalize for transfection

efficiency) into lytically reactivated KSHV-positive iSLK cells and the luciferase signal was

measured 48 h post transfection and reactivation. As expected, the presence of ori-lyt enhanced

K8.1 promoter-driven luciferase expression by 15 – 20-fold (Figure 5B). Furthermore, unlike the

consistently robust ORF57 promoter activity, K8.1 promoter activity on the plasmid was

completely repressed in the presence of DNA replication inhibitor PAA (Figure 5C) or in cells

lytically infected with the $ORF24_{RAAAG}$ mutant (Figure 5D). Taken together, these data confirm

that the plasmid assay accurately recapitulates late gene promoter activity.

Notably, mutating the 5 bp RVNYS motif in the K8.1 promoter decreased transcription

by > 50% (Figure 5E). This decrease was comparable to mutating the T at position four to an A

in the TATTWAA sequence, which has been shown to be important (although not sufficient) for

recognition and binding by ORF24 (9). Replacement of the 5 bp RVNYS motif with either

synonymous (-S) or nonsynonymous (-NS) bases yielded a similar decrease in promoter activity

(Figure $5E-K8.1_{GGGAC->TACGA-S}$, $K8.1_{GGGAC->ACTCG-NS}$), suggesting a role for the overall

architecture of this motif rather than the primary sequence. In support of this hypothesis, a

preliminary analysis of four structural features of the expanded motif in the WT and mutant

predict that the average roll of the DNA in the expanded motif is lower in the mutant promoter

compared to the WT promoter (Figure S4). Furthermore, with the exception of GG89TT-NS,

systematic single and double base substitutions in the RVNYS motif were well tolerated, arguing

that this region may not be contributing to base specific contacts with the vPIC (Figure 5F). In

summary, while the expanded RVNYS motif appears important for transcriptional activation, we

hypothesize it may contribute to vPIC specificity by impacting the local topology of the

promoter DNA rather than through base specific contacts.

The expanded promoter motif is important for DNA binding by vPIC

The RVNYS motif could either contribute to vPIC binding to its target promoters, as is

the case for the TATTWAA sequence, or it could modulate downstream events involved in

transcription initiation at vPIC bound promoters. To distinguish these possibilities, we used ChIP

to measure HA-ORF34 occupancy on the luciferase plasmids containing WT or mutant K8.1

promoter in lytically reactivated HA-ORF34 iSLK cells. We engineered qPCR primers that

enabled distinction of the K8.1 promoter on the plasmid DNA from the endogenous viral locus

(Figure 6A).

Notably, both the K8.1$_{TATT->TATA}$ and K8.1$_{GGGAC->ACTCG}$ mutations similarly impaired

HA-ORF34 binding to the plasmid promoter DNA (Figure 6B- left panel). The ChIP signal was

specific for HA-ORF34, as we observed only a background signal when the assay was performed

in the WT iSLK cells containing an untagged ORF34 locus. Furthermore, there was comparable

vPIC assembly at the genomic K8.1 locus in cells transfected with WT or the mutant plasmids,

confirming similar immunoprecipitation efficiencies across samples (Figure 6B - right panel).

Together, these data indicate that the expanded promoter motif is important for vPIC assembly

on the promoter DNA.

Discussion
324
325
326   Late gene expression is uniquely regulated in KSHV by a set of viral proteins collectively

known as the vPIC. In this study, we generated a comprehensive list of direct targets of the vPIC

by combining transcriptome and genome-wide binding data of the complex. Among the direct

targets, we identified an expanded promoter recognition motif that contributes to stable binding

of the vPIC on the promoter. This is particularly useful to understanding alternate mechanisms of

transcription initiation by RNA pol II because of the unique ability of the vPIC to directly recruit

RNA Pol II to highly compact promoters.

333   Several key points emerge from independent analysis of the RNA Seq and ChIP Seq data.

First, the previously identified TATTWAA motif is insufficient to mark binding and regulation

by the vPIC, as demonstrated by the multiple ORFs that have the motif in their promoter (23/83

ORFs) but are not repressed in the ORF24$_{RAAAG}$ mutant (10/23 ORFs) or bound by ORF34 (7/23

ORFs). Second, the TATTWAA motif is not a pre-requisite for regulation or binding by the

vPIC. For example, although ORF53 expression is dependent on the ORF24-Pol II interaction

and its promoter is bound by ORF34, its promoter does not have the conserved TATTWAA

sequence but a TATATAA sequence in its place. Third, several genes that exhibit late kinetics

escape repression in the mutant, indicating that not all late genes are direct or indirect targets of

vPIC. It is notable that several of the kinetic late genes that escape repression are involved in

immune regulation, consistent with a similar observation in EBV (28). In particular, EBV BPLF1

is a late gene whose expression is independent of the vPIC, and we observe that its homolog

ORF64 also shows a similar pattern in KSHV. Finally, while there are several early genes that

are indirect targets of the vPIC, only 1 of 13 direct targets exhibited early kinetics (ORF45). It is

notable that ORF45 is a tegument protein, which might explain the requirement for its continued

regulation at later times of infection.

Our integrative analysis revealed that a distinct 5 bp motif is integral to vPIC promoter

recognition. Rather than engaging in base specific contacts (as is the case for the TATTWAA

sequence) (14), we instead hypothesize that the importance of this motif lies in its shape or

structure. In this regard, a preliminary analysis of basic structural properties of the expanded

motif suggest that the average roll (representing the rotational flexibility) of the DNA in the

expanded motif is lower in the 5 bp mutant promoter compared to WT K8.1 promoter. Structure-

based recognition and a contribution of base pairs flanking the core binding site to binding

affinity is a common feature among transcription factors, although the specific structural

properties that influence binding differ for various classes of factors (35). A previous study of the

bZIP family of transcription factors identified a correlation between the roll of DNA sequences

that flank the core recognition element and binding affinity (35). Consistent with that, we

observed a defect in the assembly of the vPIC when the expanded motif was mutated, suggesting

a stabilizing role for the flanking sequence.

The location of the expanded motif is reminiscent of the downstream recognition element

of the general transcription factor TFIIB (BRE[d]) (36). Interestingly, the BRE[d] element also

shows low sequence conservation and does not require an active promoter sequence to conform

with the consensus sequence at all positions (36), as seen with the expanded motif. Interaction of

TFIIB with the BRE[d] element is thought to stabilize the TFIIB-TBP interaction on the promoter

DNA, leading to activation or repression of the promoter in a context-dependent manner. While

there is some evidence to suggest that TFIIB is present at late gene promoters (14), whether the

interaction between the vPIC and the expanded motif is mediated by TFIIB or any other factors

remains to be elucidated.

Notably, we found that KSHV DNA replication was reduced in the absence of functional

ORF24, including with an ORF24 null mutant or ORF24 point mutants that fail to bind either

RNA Pol II or the vPIC component ORF34. This was surprising, given that deletion or mutation

of several other vPIC components has been shown not to affect DNA replication (12,14,17,18).

Furthermore, we previously failed to observe a replication defect in an ORF24 null virus (14).

We hypothesize that the discrepancy for the ORF24 null mutant relates to the methods used to

establish the infected cell lines. In our earlier study, the viral BAC DNA was directly transfected

into iSLK cells, resulting in only a ~5-7-fold increase in genome copy number upon reactivation.

It is likely that the majority of viral DNA was present in a non-reactivatable state, providing a

level of background that significantly reduced the dynamic range for DNA replication

measurements. Here, the iSLK cells were established through co-culture with reactivated 293T

cells harboring the BAC DNA, i.e. through infection rather than transfection. This reproducibly

yields ~50-70-fold increases in genome copy number upon lytic reactivation, more closely

representing actual infection. That said, the role of ORF24 in viral DNA replication may be

unique to KSHV, as this link has not been observed upon depletion of its homologs in EBV and

MHV68 (28,37).

One possibility for how KSHV ORF24 might contribute to DNA replication is through its

indirect transcriptional targets. We observed that components of the helicase/primase complex

(ORFs 40, 41), which are involved in DNA replication, show more than average repression in the

absence of functional ORF24 (Figure 2). Importantly, these genes were not repressed in previous

studies using stop mutants of the vPIC components ORF18, ORF30 and ORF31 (17,18), raising

the possibility that ORF24 has additional gene regulatory functions outside the context of the

vPIC. For example, our observed binding of ORF24 and ORF34 at the lytic origins of replication

could facilitate DNA replication. Consistent with this hypothesis, two previous studies that

defined the boundaries of the lytic origin of replication observed a decrease in replicated DNA

upon deletion of the ORF24-ORF34 binding site identified here (38,39). In theory, this

interaction could also contribute to the link between DNA replication and late gene activation,

and thus exploring the functional consequences of origin binding will be an exciting area for

future study.

Methods

Cell lines

HEK293T cells were maintained in DMEM (Invitrogen) + 10% Fetal Bovine Serum

(FBS). iSLK BAC16 cells (29) were maintained in DMEM + 10% FBS and 0.5 mg/mL

hygromycin B. HEK293T cells stably expressing ORF24-3xFlag were maintained in DMEM

(Invitrogen) + 10% Fetal Bovine Serum (FBS) and 325 ug/ml of zeocin. iSLK-puro cells were

maintained in DMEM + 10% FBS.

Plasmids

A 100 bp fragment of the K8.1 promoter or ORF57 promoter was generated by PCR from

the BAC16 genome and cloned into the KpnI and HindIII sites of the pGL4.16 reporter plasmid

using Infusion cloning (Clontech). The left origin of replication was similarly generated by PCR

from the BAC16 genome and cloned into the NotI and BstXI sites of the K8.1Pr-pGL4.16

plasmid. The promoter mutants were generated by site-directed mutagenesis using KAPA HiFi

DNA polymerase using the primers shown in Supplementary table 1. To make the ORF24-

3xFlag pJLM1 plasmid for lentiviral transduction, the ORF24-3xFlag fragment was generated by

PCR using ORF24-3xFlag-pCDNA4 as the template and cloned in to pLJM1 plasmid digested

with AgeI and EcoRI. All primers used for cloning and mutagenesis are listed in Supplementary

table 1.

Generation of BAC mutants and establishing latent cell lines

The ORF24$_{RAAAG}$, ORF24$_{R328A}$ and ORF24$_{Stop}$ mutants, the corresponding mutant rescues

(MR) for ORF24$_{RAAAG}$, ORF24$_{R328A}$ and N terminal HA tagged ORF24 and ORF34 were

engineered using the scarless Red recombination system in BAC16 GS1783 *E. coli* (40) as

described previously. The modified BACs were purified using the NucleoBond BAC 100 kit

(Clontech).

To establish iSLK cell lines latently infected with the modified virus, HEK293T cells

were transfected with 5-10 ug of the modified BAC16 using linear polyethylenimine (PEI, MW

~25,000) at a 1:3 DNA:PEI ratio. The following day, the 293T cells were mixed 1:1 with iSLK-

puro cells (which contain inducible RTA but lack KSHV) and treated with 30 nM 12-O-

Tetradecanoylphorbol-13-acetate (TPA) and 300 nM sodium butyrate for 4 days to induce lytic

replication. Cells were then grown in selection media containing 300 ug/ml hygromycin B, 1

ug/ml puromycin and 250 ug/ml G418. The hygromycin concentration was gradually increased

to 500 ug/ml and then 1 mg/ ml until all the HEK293Ts cells died. To generate the ORF24$_{RAAAG}$,

ORF24$_{R328A}$, and ORF24$_{Stop}$ (which are defective in infectious virion production), HEK293T

cells that stably express ORF24-3xFlag were used for the initial transfection and co-culture.

Supernatant transfer assay

For supernatant transfer assays to assess virion production, BAC16-containing iSLK cells were plated at a density of 1×10^6 cells in 10 cm dishes and treated with 1 ug/mL doxycycline and 1 mM sodium butyrate the following day. 72 h post induction, the supernatant was filtered through a 0.45 uM syringe filter and 2 ml of the supernatant was mixed with 1×10^6 freshly trypsinized HEK293T cells in a 6 well plate and centrifuged at 1,500x*g* for 2 h at 37˚C. The following day, cells were trypsinized, fixed in 4% paraformaldehyde and the percentage of cells expressing GFP was determined by flow cytometry (BD Csampler, BD Biosciences).

Viral genome replication

To measure viral genome replication, iSLK-BAC16 cells were reactivated for 72 h as described above. The cells were then scraped into the media and the media + cells were digested with proteinase K (80μg/mL) (Promega) in 5x proteinase K digestion buffer (50mM Tris-HCl pH 7.4, 500mM NaCl, 5mM EDTA, 2.5% SDS) overnight at 55˚C. The gDNA was isolated using Zymo Quick gDNA Miniprep Kit according to the manufacturer's instructions. Quantitative PCR (qPCR) was performed on the isolated DNA using iTaq Universal SYBR Green Supermix on a QuantStudio3 Real-Time PCR machine. DNA levels were quantified using relative standard curves with primers specific for KSHV ORF59 promoter and human CPSF6 promoter (Supplementary table 1). The relative genome numbers were normalized to CPSF6 to account for loading differences and to uninduced samples to account for differences in starting genome copy number.

RT-qPCR

To look at viral transcript levels, total RNA was isolated from reactivated iSLK cells at the indicated time points using the Zymo Direct-Zol RNA kit following the manufacturer's protocol. The purified RNA was treated with DNase and cDNAs were synthesized using AMV reverse transcriptase (Promega). The cDNAs were directly used for qPCR analysis using the iTaq Universal SYBR Green Supermix. The qPCR signal for each ORF was normalized to 18s rRNA.

Western blotting

Cells were lysed in 1x lysis buffer (50mM Tris-HCl pH 7.5, 150mM NaCl, 0.5% NP-40, cOmplete EDTA-free Protease Inhibitors [Roche]) by rotating for 30 min at 4˚C. The lysate was clarified by centrifugation at 21,000 x g for 15 min at 4˚C. The supernatant was quantified by Bradford assay, and equivalent amounts of each sample were resolved by SDS-PAGE and western blotted with rabbit polyclonal antibodies against vinculin (Abcam, 1:1000), ORF59, or K8.1. The ORF59 and K8.1 antibodies were generated by injecting rabbits with purified MBP-ORF59 or MBP-K8.1 (gifts from Denise Whitby (41)) (Pocono Rabbit Farm and Laboratory).

RNA Seq sample preparation and analysis

Total RNA was isolated from reactivated iSLK cells at indicated time points using Zymo Direct-Zol RNA kit following the manufacturer's protocol. The samples were ribo-depleted and stranded libraries were prepped by the QB3 Berkeley core sequencing facility. Multiplex sequencing was performed using the Illumina HiSeq 4000 to generate 100 bp paired end reads. The quality of the raw reads was checked using FastQC (42) and adapters were trimmed using

TrimGalore (43) using default parameters. The trimmed reads were aligned to both the viral

(NC_009333.1) and host genomes (hg19) using STAR aligner (ver2.5.3a) (44). The aligned

reads were counted using htseq-counts (45) and differential expression was assessed using

DESeq2 (46).

To compensate for the problem of quantitative assessment of transcripts due to the

overlapping transcripts in the KSHV genome, the primary counts of the transcripts were

calculated as recommended in Bruce et al (30) with the following small modifications. Length

correction was not applied to allow for differential expression analysis by DESeq2. Transcripts

for which the counts became negative in all 3 replicates when calculating primary counts were

removed before running DESeq2. Bar plots showing Log_2 Fold Change were generated using R.

The heatmap was generated using the heatmap.2 package in R. The fold change values were

scaled by column before plotting to ease visualization of the data.

Chromatin Immunoprecipitation (ChIP)

ChIP was performed on 3*15-cm plates of iSLK-cells (WT KSHV, $KSHV_{HA-ORF34}$ and

$KSHV_{ORF24-HA}$) reactivated for 48 h with 1 ug/ml dox and 1 mM sodium butyrate. Cells were

crosslinked in 2% formaldehyde for 15 min at room temperature, quenched in 0.125 M glycine,

and washed twice with ice-cold PBS. Crosslinked cell pellets were mixed with 1 ml ice-cold

ChIP lysis buffer (5mM PIPES pH 8.0, 85mM KCl, 0.5% NP-40) and incubated on ice for 10

min, whereupon the lysate was dounce homogenized to release nuclei and spun at 1700 g for 5

min at 4°C. Nuclei were then resuspended in 800 μl of nuclei lysis buffer (50mM Tris-HCl pH

8.0, 0.3% SDS, 10mM EDTA) and rotated for 10 min at 4°C followed by sonication using a

QSonica Ultrasonicator with a cup horn set to 25 amps for 20 min total (5 min on, 5 min off).

Chromatin was spun at 16,000 x *g* for 10 min at 4°C and the pellet was discarded. The chromatin

was precleared with protein A + protein G beads blocked with 5 ug/ml glycogen, 50 ug/ml BSA,

100 ug/ml Ecoli tRNA in 1ml dilution buffer (1.1% Triton-X-100, 1.2mM EDTA, 16.7mM Tris-

HCl pH 8, 167mM NaCl) for 2 hours.

350 µl of chromatin was diluted in ChIP dilution buffer (16.7 mM Tris-HCl pH 8.0, 1.1%

Triton X-100, 1.2 mM EDTA, 167 mM NaCl) to 500 µl and incubated with 10 µg anti-HA

antibody (Cell Signaling C29F4) overnight, whereupon samples were rotated with 20 µl pre-

blocked protein A + G beads (Thermofisher) for 2 h at 4°C. Beads were washed with low salt

immune complex (20 mM Tris pH 8.0, 1% Triton-x-100, 2 mM EDTA, 150 mM NaCl, 0.1%

SDS), high salt immune complex (20 mM Tris pH 8.0, 1% Triton-x-100, 2 mM EDTA, 500 mM

NaCl, 0.1% SDS), lithium chloride immune complex (10mM Tris pH 8.0, 0.25 M LiCl, 1% NP-

40, 1% Deoxycholic acid, 1 mM EDTA), and Tris-EDTA for 10 min each at 4°C with rotation.

DNA was eluted from the beads using 100 µl of elution buffer (150 mM NaCl, 50 µg/ml

Proteinase K) and incubated at 50 °C for 2 h, then 65 °C overnight. DNA was purified using the

Zymo PCR purification kit. For ChIP-qPCR, purified DNA was quantified by qPCR using iTaq

Universal SYBR Mastermix (BioRad) and the indicated primers (Supplementary table 1). Each

sample was normalized to its own input.

For ChIP-Seq, the concentration of the eluted DNA was measured using Qubit. 10-15 ng

of eluted DNA was used to generate libraries using the Accel-NGS 2S Plus DNA Library Kit

(Swift Biosciences). The libraries were sequenced on a Illumina HiSeq 4000 sequencer at the

QB3 Berkeley core facility to generate 100 bp single end reads.

ChIP Seq Analysis

The quality of the raw reads was checked using FastQC (42) and adapters were trimmed

using TrimGalore (43) using default parameters. The trimmed reads were aligned to both the

viral and host genomes (hg19) using Bowtie2 aligner using default parameters. The aligned reads

were filtered for uniquely mapped reads using SAMTools. The filtered reads were converted to

tdf format using IGV Tools and visualized on Integrated Genome Viewer (47,48). We used

HOMER (33) with the following minor adjustments to default parameters to call peaks for the

immunoprecipitated and input samples from cells that have HA tagged ORF34 and untagged

ORF34 (WT) as a control. The fold over input and local fold change were set to 1.3 as

recommended for smaller and denser viral genomes (personal communication with Chris Benner,

UCSD). The peak size and fragment length were also set at 200 and 150 respectively. From the

resulting peaks, only the ones that were present in both conditions (HA-ORF34 vs HA-Input,

HA-ORF34 vs WT) and both replicates were retained for further analysis. The common peaks

identified were verified visually on the Integrated Genome Viewer and were manually annotated.

To identify the direct targets, transcripts that had a Log_2 fold change lower than the mean

Log_2 fold change in the $ORF24_{RAAAG}$ mutant vs MR were considered to be down regulated in the

mutant. Although ORF8 and ORF26 were not included in the differential expression analysis,

they were considered to be downregulated because of negative values for their levels in the

mutant and not in the mutant rescue on calculation of primary transcript levels.

Promoter Analyses

We defined the promoter sequence as 100 base pairs (bp) upstream from the transcription

start site (TSS) or 300 bp upstream from the translation start site when the TSS was unknown.

The TSS data was obtained from Table 1 in (23) and a custom python script was used to extract

the sequences. These were then run through MEME (31) allowing for multiple occurrences of a

motif in a sequence and only searching the given strand to identify enriched motifs. The

identified motif was then used to search for similar motifs across all 83 promoter sequences from

the viral genome, again only allowing for searching the given strand.

Reporter assays

$5*10^5$ iSLK-BAC16 (WT or ORF24$_{RAAAG}$) cells were plated in a 6 well plate 24 h before

transfection and reactivation. The cells were reactivated with 1 μg/ml doxycycline and 1 mM

Sodium Butyrate and simultaneously transfected with 0.95 μg luciferase reporter plasmid and

0.05 μg renilla plasmid using polyjet transfection reagent. 48 h post transfection and reactivation,

the cells were rinsed twice with room temperature PBS, lysed by rocking for 15 min at room

temperature in 300 μl of Passive Lysis Buffer (Promega), and clarified by centrifuging at 14,000

g for 1 min. 20 μl of the clarified lysate was added in triplicate to a white chimney well

microplate (Greiner bio-one) to measure luminescence on a Tecan M1000 using a Dual

Luciferase Assay Kit (Promega). The firefly luminescence was normalized to the internal Renilla

luciferase control. The induced sample was then normalized to the uninduced (but transfected)

control to look at the fold induction during reactivation.

Transient ChIP-qPCR

$3.5x10^6$ iSLK-BAC16 were plated in a 15 cm dish 24 h before transfection and

reactivation. The cells were reactivated with 1 μg/ml doxycycline and 1 mM Sodium Butyrate

and simultaneously transfected with 8 μg of reporter plasmid using polyjet transfection reagent.

48 h post reactivation and transfection, cells were crosslinked and lysed as mentioned above.

Nuclei were sonicated using a Covaris S220 (Covaris) for 5 min with the recommended

sonication settings (Power: 140 W, Bursts per cycle: 200, Duty cycle: 5%). The chromatin was

pre-cleared as mentioned above and the concentration measured using a qubit. 25 μg of pre-

cleared chromatin was used in the IP and the following steps were same as mentioned in the

ChIP protocol above.

Acknowledgements

We thank all members of the Glaunsinger and Coscoy labs for their helpful suggestions.

In particular, we thank Angelica Casteñeda for providing the HEK293T cells stably expressing

ORF24-3xFlag and Chris Benner (UCSD) for suggestions on proper usage of HOMER for

organisms with small genomes. This work was supported by NIH R01 AI122528 to B.G., who is

also a Howard Hughes Medical Institute Investigator.

References

1.   Payne S. Chapter 28 - Introduction to DNA Viruses. In: Payne S, editor. Viruses

[Internet]. Academic Press; 2017. p. 231–6. Available from:

http://www.sciencedirect.com/science/article/pii/B9780128031094000283

2.   Grula MA, Buller PL, Weaver RF. α-Amanitin-Resistant Viral RNA Synthesis in Nuclei

Isolated from Nuclear Polyhedrosis Virus-Infected Heliothis zea

Larvae and Spodoptera frugiperda Cells. J Virol [Internet]. 1981

Jun 1;38(3):916. Available from: http://jvi.asm.org/content/38/3/916.abstract

3.   Kochetkov SN, Rusakova EE, Tunitskaya VL. Recent studies of T7 RNA polymerase

mechanism. FEBS Lett [Internet]. 1998;440(3):264–7. Available from:

http://www.sciencedirect.com/science/article/pii/S0014579398014847

4.   Rong M, He B, McAllister WT, Durbin RK. Promoter specificity determinants of T7

RNA polymerase. Proc Natl Acad Sci U S A [Internet]. The National Academy of

Sciences; 1998 Jan 20;95(2):515–9. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/9435223

5.   Young CSH. The Structure and Function of the Adenovirus Major Late Promoter. In:

Doerfler W, Böhm P, editors. Adenoviruses: Model and Vectors in Virus-Host

Interactions: Virion-Structure, Viral Replication and Host-Cell Interactions [Internet].

Berlin, Heidelberg: Springer Berlin Heidelberg; 2003. p. 213–49. Available from:

https://doi.org/10.1007/978-3-662-05597-7_8

6.   Gruffat H, Marchione R, Manet E. Herpesvirus Late Gene Expression: A Viral-Specific

Pre-initiation Complex Is Key. Front Microbiol [Internet]. Frontiers Media S.A.; 2016 Jun

6;7:869. Available from: https://www.ncbi.nlm.nih.gov/pubmed/27375590

7.   Wong-Ho E, Wu T-T, Davis ZH, Zhang B, Huang J, Gong H, et al. Unconventional

Sequence Requirement for Viral Late Gene Core Promoters of Murine Gammaherpesvirus

68. Frueh K, editor. J Virol [Internet]. 2014 Mar 15;88(6):3411. Available from:

http://jvi.asm.org/content/88/6/3411.abstract

8.   Serio TR, Cahill N, Prout ME, Miller G. A Functionally Distinct TATA Box Required for

Late Progression through the Epstein-Barr Virus Life Cycle. J Virol [Internet]. 1998 Oct

1;72(10):8338. Available from: http://jvi.asm.org/content/72/10/8338.abstract

9.   Tang S, Yamanegi K, Zheng Z-M. Requirement of a 12-Base-Pair TATT-Containing

Sequence and Viral Lytic DNA Replication in Activation of the Kaposi's Sarcoma-

Associated Herpesvirus K8.1 Late Promoter. J Virol [Internet]. 2004 Mar 1;78(5):2609.

Available from: http://jvi.asm.org/content/78/5/2609.abstract

10.   Haberle V, Stark A. Eukaryotic core promoters and the functional basis of transcription

initiation. Nat Rev Mol Cell Biol [Internet]. 2018;19(10):621–37. Available from:

https://doi.org/10.1038/s41580-018-0028-8

11.   Aubry V, Mure F, Mariamé B, Deschamps T, Wyrwicz LS, Manet E, et al. Epstein-Barr

Virus Late Gene Transcription Depends on the Assembly of a Virus-Specific Preinitiation

Complex. Longnecker RM, editor. J Virol [Internet]. 2014 Nov 1;88(21):12825. Available

from: http://jvi.asm.org/content/88/21/12825.abstract

12.   Davis ZH, Hesser CR, Park J, Glaunsinger BA. Interaction between ORF24 and ORF34 in

the Kaposi's Sarcoma-Associated Herpesvirus Late Gene Transcription Factor

Complex Is Essential for Viral Late Gene Expression. Sandri-Goldin RM, editor. J Virol

[Internet]. 2016 Jan 1;90(1):599. Available from:

http://jvi.asm.org/content/90/1/599.abstract

13.   Wyrwicz LS, Rychlewski L. Identification of Herpes TATT-binding protein. Antiviral

Res [Internet]. 2007;75(2):167–72. Available from:

http://www.sciencedirect.com/science/article/pii/S0166354207002367

14.   Davis ZH, Verschueren E, Jang GM, Kleffman K, Johnson JR, Park J, et al. Global

Mapping of Herpesvirus-Host Protein Complexes Reveals a Transcription Strategy for

Late Genes. Mol Cell [Internet]. 2015;57(2):349–60. Available from:

http://www.sciencedirect.com/science/article/pii/S1097276514009216

15.   Gruffat H, Kadjouf F, Mariamé B, Manet E. The Epstein-Barr Virus BcRF1 Gene Product

Is a TBP-Like Protein with an Essential Role in Late Gene Expression. J Virol [Internet].

2012 Jun 1;86(11):6023. Available from: http://jvi.asm.org/content/86/11/6023.abstract

16.   Nishimura M, Watanabe T, Yagi S, Yamanaka T, Fujimuro M. Kaposi's sarcoma-

645   associated herpesvirus ORF34 is essential for late gene expression and virus production.

646   Sci Rep [Internet]. 2017;7(1):329. Available from: https://doi.org/10.1038/s41598-017-

647   00401-7

648   17.   Gong D, Wu NC, Xie Y, Feng J, Tong L, Brulois KF, et al. Kaposi's Sarcoma-

649   Associated Herpesvirus ORF18 and ORF30 Are Essential for Late Gene Expression

650   during Lytic Replication. Frueh K, editor. J Virol [Internet]. 2014 Oct 1;88(19):11369.

651   Available from: http://jvi.asm.org/content/88/19/11369.abstract

652   18.   Brulois K, Wong L-Y, Lee H-R, Sivadas P, Ensser A, Feng P, et al. Association of

653   Kaposi's Sarcoma-Associated Herpesvirus ORF31 with ORF34 and ORF24 Is

654   Critical for Late Gene Expression. Longnecker RM, editor. J Virol [Internet]. 2015 Jun

655   1;89(11):6148. Available from: http://jvi.asm.org/content/89/11/6148.abstract

656   19.   Perng Y-C, Campbell JA, Lenschow DJ, Yu D. Human Cytomegalovirus pUL79 Is an

657   Elongation Factor of RNA Polymerase II for Viral Gene Transcription. PLOS Pathog

658   [Internet]. Public Library of Science; 2014 Aug 28;10(8):e1004350-. Available from:

659   https://doi.org/10.1371/journal.ppat.1004350

660   20.   Castañeda AF, Glaunsinger BA. The Interaction between ORF18 and ORF30 Is Required

661   for Late Gene Expression in Kaposi's Sarcoma-Associated Herpesvirus. Sandri-

662   Goldin RM, editor. J Virol [Internet]. 2019 Jan 1;93(1):e01488-18. Available from:

663   http://jvi.asm.org/content/93/1/e01488-18.abstract

664   21.   Pan D, Han T, Tang S, Xu W, Bao Q, Sun Y, et al. Murine Cytomegalovirus Protein

665   pM91 Interacts with pM79 and Is Critical for Viral Late Gene Expression. Sandri-Goldin

666   RM, editor. J Virol [Internet]. 2018 Sep 15;92(18):e00675-18. Available from:

667    http://jvi.asm.org/content/92/18/e00675-18.abstract

668 22.  Djavadian R, Hayes M, Johannsen E. CAGE-seq analysis of Epstein-Barr virus lytic gene

669    transcription: 3 kinetic classes from 2 mechanisms. PLOS Pathog [Internet]. Public

670    Library of Science; 2018 Jun 4;14(6):e1007114-. Available from:

671    https://doi.org/10.1371/journal.ppat.1007114

672 23.  Arias C, Weisburd B, Stern-Ginossar N, Mercier A, Madrid AS, Bellare P, et al. KSHV

673    2.0: A Comprehensive Annotation of the Kaposi's Sarcoma-Associated Herpesvirus

674    Genome Using Next-Generation Sequencing Reveals Novel Genomic and Functional

675    Features. PLOS Pathog [Internet]. Public Library of Science; 2014 Jan

676    16;10(1):e1003847-. Available from: https://doi.org/10.1371/journal.ppat.1003847

677 24.  Nakamura H, Lu M, Gwack Y, Souvlis J, Zeichner SL, Jung JU. Global Changes in

678    Kaposi's Sarcoma-Associated Virus Gene Expression Patterns following

679    Expression of a Tetracycline-Inducible Rta Transactivator. J Virol [Internet]. 2003 Apr

680    1;77(7):4205. Available from: http://jvi.asm.org/content/77/7/4205.abstract

681 25.  Jenner RG, Albà MM, Boshoff C, Kellam P. Kaposi's Sarcoma-Associated

682    Herpesvirus Latent and Lytic Gene Expression as Revealed by DNA Arrays. J Virol

683    [Internet]. 2001 Jan 15;75(2):891. Available from:

684    http://jvi.asm.org/content/75/2/891.abstract

685 26.  Paulose-Murphy M, Ha N-K, Xiang C, Chen Y, Gillim L, Yarchoan R, et al. Transcription

686    Program of Human Herpesvirus 8 (Kaposi's Sarcoma-Associated Herpesvirus). J

687    Virol [Internet]. 2001 May 15;75(10):4843. Available from:

688    http://jvi.asm.org/content/75/10/4843.abstract

689 27.  Arumugaswami V, Wu T-T, Martinez-Guzman D, Jia Q, Deng H, Reyes N, et al. ORF18

Is a Transfactor That Is Essential for Late Gene Transcription of a Gammaherpesvirus. J

Virol [Internet]. 2006 Oct 1;80(19):9730. Available from:

http://jvi.asm.org/content/80/19/9730.abstract

28.   McKenzie J, Lopez-Giraldez F, Delecluse H-J, Walsh A, El-Guindy A. The Epstein-Barr

Virus Immunoevasins BCRF1 and BPLF1 Are Expressed by a Mechanism Independent of

the Canonical Late Pre-initiation Complex. PLOS Pathog [Internet]. Public Library of

Science; 2016 Nov 17;12(11):e1006008-. Available from:

https://doi.org/10.1371/journal.ppat.1006008

29.   Brulois KF, Chang H, Lee AS-Y, Ensser A, Wong L-Y, Toth Z, et al. Construction and

Manipulation of a New Kaposi's Sarcoma-Associated Herpesvirus Bacterial

Artificial Chromosome Clone. J Virol [Internet]. 2012 Sep 15;86(18):9708. Available

from: http://jvi.asm.org/content/86/18/9708.abstract

30.   Bruce AG, Barcy S, DiMaio T, Gan E, Garrigues HJ, Lagunoff M, et al. Quantitative

Analysis of the KSHV Transcriptome Following Primary Infection of Blood and

Lymphatic Endothelial Cells. Pathog (Basel, Switzerland) [Internet]. MDPI; 2017 Mar

19;6(1):11. Available from: https://www.ncbi.nlm.nih.gov/pubmed/28335496

31.   Bailey L. T, Elkan C. Fitting a mixture model by expectation maximization to discover

motifs in biopolymers. In: Proceedings of the Second International Conference on

Intelligent Systems for Molecular Biology [Internet]. Menlo Park; 1994. p. 28–36.

Available from:

https://www.ncbi.nlm.nih.gov/pubmed/?term=Fitting+a+mixture+model+by+expectation+

maximization+to+discover+motifs+in+biopolymers

32.   Grant CE, Bailey TL, Noble WS. FIMO: scanning for occurrences of a given motif.

Bioinformatics [Internet]. 2011 Feb 16;27(7):1017–8. Available from:

https://dx.doi.org/10.1093/bioinformatics/btr064

33.   Heinz S, Benner C, Spann N, Bertolino E, Lin YC, Laslo P, et al. Simple Combinations of

Lineage-Determining Transcription Factors Prime cis-Regulatory Elements Required for

Macrophage and B Cell Identities. Mol Cell [Internet]. 2010;38(4):576–89. Available

from: http://www.sciencedirect.com/science/article/pii/S1097276510003667

34.   Djavadian R, Chiu Y-F, Johannsen E. An Epstein-Barr Virus-Encoded Protein Complex

Requires an Origin of Lytic Replication In Cis to Mediate Late Gene Transcription. PLOS

Pathog [Internet]. Public Library of Science; 2016 Jun 27;12(6):e1005718-. Available

from: https://doi.org/10.1371/journal.ppat.1005718

35.   Yella VR, Bhimsaria D, Ghoshdastidar D, Rodríguez-Martínez JA, Ansari AZ, Bansal M.

Flexibility and structure of flanking DNA impact transcription factor affinity for its core

motif. Nucleic Acids Res [Internet]. 2018/11/05. Oxford University Press; 2018 Dec

14;46(22):11883–97. Available from: https://www.ncbi.nlm.nih.gov/pubmed/30395339

36.   Deng W, Roberts SGE. A core promoter element downstream of the TATA box that is

recognized by TFIIB. Genes Dev [Internet]. Cold Spring Harbor Laboratory Press; 2005

Oct 15;19(20):2418–23. Available from: https://www.ncbi.nlm.nih.gov/pubmed/16230532

37.   Wong E, Wu T-T, Reyes N, Deng H, Sun R. Murine Gammaherpesvirus 68 Open Reading

Frame 24 Is Required for Late Gene Expression after DNA Replication. J Virol [Internet].

2007 Jun 15;81(12):6761. Available from: http://jvi.asm.org/content/81/12/6761.abstract

38.   Lin CL, Li H, Wang Y, Zhu FX, Kudchodkar S, Yuan Y. Kaposi's sarcoma-associated

herpesvirus lytic origin (ori-Lyt)-dependent DNA replication: identification of the ori-Lyt

and association of K8 bZip protein with the origin. J Virol [Internet]. American Society

for Microbiology; 2003 May;77(10):5578–88. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/12719550

39. AuCoin DP, Colletti KS, Xu Y, Cei SA, Pari GS. Kaposi's Sarcoma-Associated

Herpesvirus (Human Herpesvirus 8) Contains Two Functional Lytic Origins of DNA

Replication. J Virol [Internet]. 2002 Aug 1;76(15):7890. Available from:

http://jvi.asm.org/content/76/15/7890.abstract

40. Gardner MR, Glaunsinger BA. Kaposi's Sarcoma-Associated Herpesvirus ORF68

Is a DNA Binding Protein Required for Viral Genome Cleavage and Packaging. Sandri-

Goldin RM, editor. J Virol [Internet]. 2018 Aug 15;92(16):e00840-18. Available from:

http://jvi.asm.org/content/92/16/e00840-18.abstract

41. Labo N, Miley W, Marshall V, Gillette W, Esposito D, Bess M, et al. Heterogeneity and

Breadth of Host Antibody Response to KSHV Infection Demonstrated by Systematic

Analysis of the KSHV Proteome. PLOS Pathog [Internet]. Public Library of Science;

2014 Mar 27;10(3):e1004046. Available from:

https://doi.org/10.1371/journal.ppat.1004046

42. Andrews S. FastQC A Quality Control tool for High Throughput Sequence Data

[Internet]. Available from: http://www.bioinformatics.babraham.ac.uk/projects/fastqc/

43. Trim Galore - http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/. Available

from: http://www.bioinformatics.babraham.ac.uk/projects/trim_galore/

44. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. Bioinformatics [Internet]. 2012/10/25. Oxford University

Press; 2013 Jan;29(1):15–21. Available from:

https://www.ncbi.nlm.nih.gov/pubmed/23104886

45.   Pyl PT, Anders S, Huber W. HTSeq—a Python framework to work with high-throughput

760   sequencing data. Bioinformatics [Internet]. 2014 Sep 25;31(2):166–9. Available from:

761   https://dx.doi.org/10.1093/bioinformatics/btu638

762   46.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for

763   RNA-seq data with DESeq2. Genome Biol [Internet]. 2014;15(12):550. Available from:

764   https://doi.org/10.1186/s13059-014-0550-8

765   47.   Robinson JT, Thorvaldsdóttir H, Winckler W, Guttman M, Lander ES, Getz G, et al.

766   Integrative genomics viewer. Nat Biotechnol [Internet]. Nature Publishing Group, a

767   division of Macmillan Publishers Limited. All Rights Reserved.; 2011 Jan 10;29:24.

768   Available from: https://doi.org/10.1038/nbt.1754

769   48.   Thorvaldsdóttir H, Robinson JT, Mesirov JP. Integrative Genomics Viewer (IGV): high-

770   performance genomics data visualization and exploration. Brief Bioinform [Internet].

771   2012 Apr 19;14(2):178–92. Available from: https://dx.doi.org/10.1093/bib/bbs017

772   49.   Zhou T, Yang L, Lu Y, Dror I, Dantas Machado AC, Ghane T, et al. DNAshape: a method

773   for the high-throughput prediction of DNA structural features on a genomic scale. Nucleic

774   Acids Res [Internet]. 2013/05/22. Oxford University Press; 2013 Jul;41(Web Server

775   issue):W56–62. Available from: https://www.ncbi.nlm.nih.gov/pubmed/23703209

Figure Legends

Fig 1: ORF24-RNA pol II interaction is essential for late gene expression and contributes to

viral DNA replication

iSLK cells harboring WT BAC16, the ORF24$_{RAAAG}$ mutant or the ORF24$_{RAAAG}$ mutant rescue

(MR) were reactivated with doxycycline and sodium butyrate for 72 hours. (A) Western blot

showing expression of an early protein (ORF59) and a late protein (K8.1) in WT, ORF24$_{RAAAG}$

and ORF24$_{RAAAG}$-MR. Vinculin was used as a loading control. (B) Progeny virion production

was measured by supernatant transfer on to 293T cells and quantified by flow cytometry. (C)

DNA replication of different mutants and MRs was measured by qPCR and normalized to the

respective uninduced sample. The fold induction of all mutants was normalized to the WT to get

the relative change in DNA replication compared to WT.

Fig 2:   Differential expression analysis of ORF24-Pol II interaction mutant and mutant

rescue

iSLK cells harboring ORF24$_{RAAAG}$ and ORF24$_{RAAAG}$ -MR were reactivated with doxycycline and

sodium butyrate for 24 and 48 h, whereupon total RNA was extracted and subjected to library

preparation and RNA sequencing. (A) Bar plot showing log$_2$ fold change of the viral genes in the

ORF24$_{RAAAG}$ -MR at 48 h relative to 24 h. Genes were color coded by kinetic class as defined in

(30). (B) Bar plot showing log$_2$ fold change of the viral genes in the mutant relative to the MR.

Genes were color coded by kinetic class as defined in (30). (C) Heatmap of the scaled log$_2$ fold

change of the genes induced in the MR between 24 to 48 h and genes repressed in the mutant

compared to MR at 48 h. Gene clusters with different kinetics and dependence on ORF24-Pol II

interaction are indicated. (D) Consensus motif identified by MEME analysis of promoters from

14 genes that showed greater than one standard deviation repression in the mutant relative to the

MR.

Fig 3: ChIP-Seq with HA-ORF34 shows multiple binding sites across the viral genome

(A) ChIP-Seq was performed using an anti-HA antibody to immunoprecipitate DNA prepared

from iSLK HA-ORF34 cells reactivated for 48 h with doxycycline and sodium butyrate. iSLK-

WT cells lacking the HA tag and input DNA were used as negative controls. The alignment files

were converted to the tiled data file (tdf) format and the ChIP background from the untagged

cells was subtracted and visualized on the Integrated Genome Viewer (IGV) (B) ChIP-qPCR was

performed using an anti-HA antibody to immunoprecipitate DNA prepared from iSLK HA-

ORF24 cells reactivated for 48 h with doxycycline and sodium butyrate. The DNA was

quantified by qPCR with primers specific to the promoter regions of the indicated genes. Data

were normalized to the level of input DNA and are presented as percentages of the input. Data

shown are from three biological replicates. (C) Consensus motif identified by MEME analysis of

promoters from 19 genes whose promoters had a ChIP peak.

Fig 4: Identifying direct targets of the vPIC

(A) Heatmap showing genes repressed in the $ORF24_{RAAAG}$ mutant compared to the MR adjacent

to the list of genes that show ORF34 binding at their promoter by ChIP-Seq analysis. Also

indicated are viral genes that were identified to have the TATTWAA motif by FIMO analysis.

The direct targets were identified by their combined dependence on the ORF24-Pol II interaction

and the presence of a ChIP signal at the promoter. (B) Consensus motif identified by MEME

analysis of promoters from 13 direct targets.

Fig 5: Motif identified from direct targets is important for transcription from the promoter

(A) Schematic of plasmid-based promoter activation assay. iSLK-WT or $ORF24_{RAAAG}$ mutant

cells reactivated with doxycycline and sodium butyrate were transfected with reporter plasmid

carrying the indicated promoter sequence upstream of the firefly luciferase gene and a control

renilla luciferase expressing plasmid. 48 h post transfection and reactivation, the cells were

assessed for luciferase activity. (B) Reactivated iSLK-WT cells were transfected with the K8.1

promoter reporter plasmid with or without the KSHV left lytic origin of replication cloned in *cis*

and assessed for luciferase activity. (C) Reactivated iSLK-WT cells were transfected with the

K8.1 and ORF57 promoter reporter plasmids in the presence or absence of DNA replication

inhibitor (PAA) and assessed for luciferase activity. (D) Reactivated iSLK-WT and iSLK-

ORF24$_{RAAAG}$ mutant cells were transfected with K8.1 or ORF57 promoter reporter plasmids and

assessed for luciferase activity. (E-F) iSLK-WT cells were transfected with plasmids containing

the WT K8.1 promoter or the indicated single or double point mutant promoters and assessed for

luciferase activity. The data for each mutant was normalized to the WT K8.1 promoter. All data

shown are average of 3-5 biological replicates.

Fig 6: Expanded motif is important for DNA binding by vPIC

(A) Schematic of the transient ChIP assay. iSLK-WT or iSLK-HAORF34 cells reactivated with

doxycycline and sodium butyrate were transfected with plasmid carrying the indicated promoter

sequence. 48 h post transfection and reactivation, cells were subjected to ChIP with an anti-HA

antibody, and the associated DNA was quantified using primers specific to the promoter on the

plasmid and the genomic K8.1 promoter locus. (B) iSLK-HAORF34 cells were transfected with

K8.1 promoter containing plasmid, mutant plasmids or ORF57 promoter plasmid (as a control).

iSLK-WT cells without the HA tag transfected with K8.1 plasmid served as a control for the IP.

The ChIP DNA was quantified using plasmid specific primers (left panel) and K8.1 promoter

specific primers (right panel), normalized to the level of input DNA and presented as percent

input. Data shown are an average of 3-6 biological replicates.

Fig S1: Validation of mutant BAC DNA integrity

The integrity of the various BAC mutants used in this study was verified by *Rsr*II digestion.

Fig S2: Kinetics of early and late gene transcripts in ORF24$_{RAAAG}$ mutant and MR

iSLK-ORF24$_{RAAAG}$ and MR were reactivated with doxycycline and sodium butyrate for the

indicated time points, whereupon total RNA was extracted, converted to cDNA, and quantified

by qPCR using primers specific to the CDS of K8.1 (late gene) or ORF57 (early gene). The fold

induction was measured relative to uninduced sample.

Fig S3: ChIP-Seq data with HA-ORF34

(A) iSLK cells harboring WT BAC16, HA-ORF34 BAC or HA-ORF24 BAC were reactivated

with doxycycline and sodium butyrate for 72 hours. Progeny virion production was measured by

supernatant transfer on to 293T cells and quantified by flow cytometry. (B) IGV image showing

ChIP-Seq data from iSLK-HA-ORF34 cells reactivated with doxycycline and sodium butyrate

for 48 h and immunoprecipitated with anti-HA antibody. Also shown are plots for control data

from untagged iSLK cells and input DNA for two biological replicates

Fig S4: Effect of mutations on DNA structural parameters

Four structural parameters for 40 bp of the WT and expanded motif mutants were calculated

using the DNAShape program in R (49). Average of each structural parameter for the mutated

base pairs and the two adjacent base pairs at either end (a total of 7 bp) was calculated for each

DNA sequence analyzed. The two adjacent base pairs were included in the averaging as some of

the structural parameters refer to a base step rather than a specific base. The average for each

mutant was normalized to the WT K8.1 promoter.

Figure 1

Figure S1

Figure S2

Figure 2

A.

24 h vs 48 h
ORF24$_{RAAAG}$-MR

B.

48 h
ORF24$_{RAAAG}$ vs ORF24$_{RAAAG}$-MR

- ■ Latent
- ■ Early/Delayed Early
- ■ Late
- ■ Unknown

C.

Viral genes

24 h vs 48 h
ORF24$_{RAAAG}$MR
(Kinetics)

ORF24$_{RAAAG}$
vs
ORF24$_{RAAAG}$MR

Early genes
Independent of ORF24

Delayed Early / Late genes
Independent of ORF24

Early/Delayed Early
Dependent on ORF24

Delayed Early / Late genes
Dependent on ORF24

Scaled Fold Change

D.

E-value: 3.4e-006

Figure 3

A

B

C

E-value: 1.3e-014

Figure S3

A

Supernatant transfer

B

Replicate 1

Replicate 2

Figure 4

A

B TATTWAARVNYS

E-value: 5.8e-009

Figure 5

Figure 6

A

B

Figure S4

Table 1. Occurrence of TATTWAA motif in KSHV

Promoter	Start	Stop	Strand	Score	p-value	q-value	Matched Sequence	Position
ORF17	68	74	+	11.8614	0.000114	0.143	TATTTAA	32
ORF62	68	74	+	11.8614	0.000114	0.143	TATTTAA	32
ORF55	69	75	+	11.8614	0.000114	0.143	TATTTAA	31
ORF39	70	76	+	11.8614	0.000114	0.143	TATTTAA	30
ORF8	71	77	+	11.8614	0.000114	0.143	TATTTAA	29
ORF52	71	77	+	11.8614	0.000114	0.143	TATTTAA	29
ORF17.5	71	77	+	11.8614	0.000114	0.143	TATTTAA	29
ORF59	71	77	+	11.8614	0.000114	0.143	TATTTAA	29
ORF58	71	77	+	11.8614	0.000114	0.143	TATTTAA	29
ORF19	187	193	+	11.8614	0.000114	0.143	TATTTAA	113
ORF27	197	203	+	11.8614	0.000114	0.143	TATTTAA	103
ORF32	243	249	+	11.8614	0.000114	0.143	TATTTAA	57
ORF25	11	17	+	11.198	0.000235	0.153	TATTAAA	89
K8.1	68	74	+	11.198	0.000235	0.153	TATTAAA	32
K4.1	68	74	+	11.198	0.000235	0.153	TATTAAA	32
ORF65	69	75	+	11.198	0.000235	0.153	TATTAAA	31
ORF33	70	76	+	11.198	0.000235	0.153	TATTAAA	30
ORF26	71	77	+	11.198	0.000235	0.153	TATTAAA	29
ORF38	71	77	+	11.198	0.000235	0.153	TATTAAA	29
ORF4	71	77	+	11.198	0.000235	0.153	TATTAAA	29
ORF64	145	151	+	11.198	0.000235	0.153	TATTAAA	155
ORF66	218	224	+	11.198	0.000235	0.153	TATTAAA	82
ORF22	229	235	+	11.198	0.000235	0.153	TATTAAA	71

ORFs shown in red do not have a known Transcription Start Site (TSS)

Table 2. Features of ChIP-Seq targets

ChIP targets	Function	Kinetics	Strand	Position of motif from TSS
K8.1	Glycoprotein	L	+	32
ORF52	Immune Evasion	L	-	29
ORF27	Glycoprotein	L	+	103
ORF53	Glycoprotein	L	-	29
ORF25	Major capsid protein	L	+	89 (29[#])
ORF62	Assembly/DNA maturation	L	-	32
ORF33	Capsid binding	L	+	30
ORF8	Glycoprotein	L	+	29
ORF17.5	Viral capsid assembly	L	-	29 (73[#])
ORF19	Capsid vertex		-	113
ORF26	Minor capsid protein	L	+	29
ORF45	Immune evasion (Tegument)	E	-	No motif
ORF6	Single strand DNA binding protein	E	+	89 (31[#])
K4.1	Homologs of cellular chemokines	E	-	32
ORF59	DNA polymerase processivity factor	E	-	29
ORF68	Packaging protein	E	+	No motif
ORF4	Immune evasion	E	+	29
ORF65	Minor capsid protein	L	-	31
ORF38	Viral maturation and egress (Tegument protein)	L	+	29

Second site with lower consensus to TATTWAARV motif
Direct targets are shown in gray

Supplementary Table 1:
List of all DNA sequences used in the study

Name	Sequence	Used for
K8.1Pr_pGL4.16_F	TGGCCTAACTGGCCGGTACCTGTGCCGTCGTCCGGGA	Cloning
K8.1Pr_pGL4.16_R	CCGGATTGCCAAGCTTAGGATTAAGGGATTAACTTCGGTCCC	Cloning
ORF57Pr_pGL4.16_F	TGGCCTAACTGGCCGGTACCAGTGTAACAATAATGTTCCCACGGC	Cloning
ORF57Pr_pGL4.16_R	CCGGATTGCCAAGCTTTTTTGGGAACCTGGCAGCCAGGTTA	Cloning
Orilyt_KSHV_F	GGTCTGACAGCGGCCGCCAGCTTATCCGGTGGCAATCT	Cloning
Orilyt_KSHV_R	AGCACTGGTAACCACTGCAGTGGGCACAGAGATATGTGACGTGACGCCCT	Cloning
K8.1_G8A_1	GGCAGCAATATTAAAAGGACCGAAGTTAATC	Mutagenesis
K8.1_G8A_2	GATTAACTTCGGTCCTTTTAATATTGCTGCC	Mutagenesis
K8.1_G8C_1	GGCAGCAATATTAAACGGACCGAAGTTAATC	Mutagenesis
K8.1_G8C_2	GATTAACTTCGGTCCGTTTAATATTGCTGCC	Mutagenesis
K8.1_G8T_1	GGCAGCAATATTAAATGGACCGAAGTTAATC	Mutagenesis
K8.1_G8T_2	GATTAACTTCGGTCCATTTAATATTGCTGCC	Mutagenesis
K8.1_G9C_1	GCAGCAATATTAAAGCGACCGAAGTTAATCC	Mutagenesis
K8.1_G9C_2	GGATTAACTTCGGTCGCTTTAATATTGCTGC	Mutagenesis
K8.1_G9A_1	GCAGCAATATTAAAGAGACCGAAGTTAATCC	Mutagenesis
K8.1_G9A_2	GGATTAACTTCGGTCTCTTTAATATTGCTGC	Mutagenesis
K8.1_G9T_1	GCAGCAATATTAAAGTGACCGAAGTTAATCC	Mutagenesis
K8.1_G9T_2	GGATTAACTTCGGTCACTTTAATATTGCTGC	Mutagenesis
K8.1_G10T_1	CAGCAATATTAAAGGTACCGAAGTTAATCCC	Mutagenesis
K8.1_G10T_2	GGGATTAACTTCGGTACCTTTAATATTGCTG	Mutagenesis
K8.1_A11C_1	AGCAATATTAAAGGGCCCGAAGTTAATCCCT	Mutagenesis
K8.1_A11C_2	AGGGATTAACTTCGGGCCCTTTAATATTGCT	Mutagenesis
K8.1_A11T_1	AGCAATATTAAAGGGTCCGAAGTTAATCCCT	Mutagenesis
K8.1_A11T_2	AGGGATTAACTTCGGACCCTTTAATATTGCT	Mutagenesis
K8.1_A11G_1	AGCAATATTAAAGGGGCCGAAGTTAATCCCT	Mutagenesis
K8.1_A11G_2	AGGGATTAACTTCGGCCCCTTTAATATTGCT	Mutagenesis
K8.1_C12G_1	GCAATATTAAAGGGAGCGAAGTTAATCCCTT	Mutagenesis
K8.1_C12G_2	AAGGGATTAACTTCGCTCCCTTTAATATTGC	Mutagenesis
K8.1_C12T_1	GCAATATTAAAGGGATCGAAGTTAATCCCTT	Mutagenesis
K8.1_C12T_2	AAGGGATTAACTTCGATCCCTTTAATATTGC	Mutagenesis
K8.1_GG89AC_1	GGCAGCAATATTAAAACGACCGAAGTTAATCCCTT	Mutagenesis
K8.1_GG89AC_2	AAGGGATTAACTTCGGTCGTTTTAATATTGCTGCC	Mutagenesis
K8.1_GG89CT_1	GGCAGCAATATTAAACTGACCGAAGTTAATCCCTT	Mutagenesis
K8.1_GG89CT_2	AAGGGATTAACTTCGGTCAGTTTAATATTGCTGCC	Mutagenesis
K8.1_GG89TT_1	GGCAGCAATATTAAATTGACCGAAGTTAATCCCTT	Mutagenesis
K8.1_GG89TT_2	AAGGGATTAACTTCGGTCAATTTAATATTGCTGCC	Mutagenesis

K8.1_AC1112CG_1	GGCAGCAATATTAAAGGGCGCGAAGTTAATCCCTTAATC	Mutagenesis
K8.1_AC1112CG_2	GATTAAGGGATTAACTTCGCGCCCTTTAATATTGCTGCC	Mutagenesis
K8.1_AC1112GT_1	GGCAGCAATATTAAAGGGGTCGAAGTTAATCCCTTAATC	Mutagenesis
K8.1_AC1112GT_2	GATTAAGGGATTAACTTCGACCCCTTTAATATTGCTGCC	Mutagenesis
K8.1_TATT->TATA_1	GTTATCCGGCAGCAATATAAAAGGGACCGAAGTTAATC	Mutagenesis
K8.1_TATT->TATA_2	GATTAACTTCGGTCCCTTTTATATTGCTGCCGGATAAC	Mutagenesis
K8.1_GGGAC->TACGA_1	GGCAGCAATATTAAATACGACGAAGTTAATCCCTT	Mutagenesis
K8.1_GGGAC->TACGA_2	AAGGGATTAACTTCGTCGTATTTAATATTGCTGCC	Mutagenesis
K8.1_GGGAC->ACTCG_1	GGCAGCAATATTAAAACTCGCGAAGTTAATCCCTT	Mutagenesis
K8.1_GGGAC->ACTCG_2	AAGGGATTAACTTCGCGAGTTTTAATATTGCTGCC	Mutagenesis
K8.1Pr_qPCR_F	GGGAGAACCATGCCAGACTTTG	ChIP-qPCR
K8.1Pr_qPCR_R	GCATAGGATTAGGAGCGCCAC	ChIP-qPCR
ORF59Pr_qPCR_F	AATCCACAGGCATGATTGC	ChIP-qPCR/ DNA replication
ORF59Pr_qPCR_R	CACACTTCCACCTCCCCTAA	ChIP-qPCR DNA replication
CPSF6Pr_qPCR_F	CGAGGAATGTCCCTGTTTGT	DNA replication
CPSF6Pr_qPCR_R	ACTGGCTGTCTCCTCTCAGC	DNA replication
K8.1CDS_qPCR_F	CCGTCGGTGTGTAGGGATAAAG	RT-qPCR
K8.1CDS_qPCR_R	GTCGTTGTAGTGGTGGCAGAAA	RT-qPCR
ORF57CDS_qPCR_F	TTTGACGAATCGAGGGACGACG	RT-qPCR
ORF57CDS_qPCR_R	GCAGTTGAGAACGACCTTGAGAT	RT-qPCR
ORF17.5Pr_qPCR_F	GCTGGCCTTTAAATACGCCG	ChIP-qPCR
ORF17.5Pr_qPCR_R	TTTCGTCAGTCCCCTGGAGA	ChIP-qPCR
OriLyt_L_qPCR_F	CGGCGGAGGTTGTCAATAGT	ChIP-qPCR
OriLyt_L_qPCR_R	GGGTCCTGTTCTGGGATAGC	ChIP-qPCR
OriLyt_R_qPCR_F	CCCTGGGCACGATAACCATA	ChIP-qPCR
OriLyt_R_qPCR_R	AGGGGCGCAAAACAATTGTG	ChIP-qPCR
ORF64Pr_qPCR_F	GCAGAAAGAACTTGAGCTCT	ChIP-qPCR
ORF64Pr_qPCR_R	AGGCAGACCGGGAGTACGTT	ChIP-qPCR
ORF37Pr_qPCR_F	GCTGGCGTTTCAGAAGCAGTG	ChIP-qPCR
ORF37Pr_qPCR_R	CCATCCAGGGTGTCAACCA	ChIP-qPCR