1 *Article*

2 The Patchy Distribution of Restriction-Modification
3 System Genes and the Conservation of Orphan
4 Methyltransferases in Halobacteria

Matthew S. Fullmer[1,2,$], Matthew Ouellette[1,$], Artemis S. Louyakis[1,$],   R. Thane Papke[1*], J. Peter
Gogarten[1*]

[1] Department of Molecular and Cell Biology, University of Connecticut, Storrs, CT, USA
[2] Present Address: Bioinformatics Institute, School of Biological Sciences, The University of Auckland,
Auckland, New Zealand
[$] These authors contributed equally
* Correspondence: thane@uconn.edu; gogarten@uconn.edu

Abstract: Restriction-modification (RM) systems in Bacteria are implicated in multiple biological
roles ranging from defense against parasitic genetic elements, to selfish addiction cassettes, and
barriers to gene transfer and lineage homogenization. In Bacteria, DNA-methylation without
cognate restriction also plays important roles in DNA replication, mismatch repair, protein
expression, and in biasing DNA uptake.   Little is known about archaeal RM systems and DNA
methylation. To elucidate further understanding for the role of RM systems and DNA methylation
in Archaea, we undertook a survey of the presence of RM system genes and related genes, including
orphan DNA methylases, in the halophilic archaeal class Halobacteria. Our results reveal that some
orphan DNA methyltransferase genes were highly conserved among lineages indicating an
important functional constraint, whereas RM systems demonstrated patchy patterns of presence
and absence. This irregular distribution is due to frequent horizontal gene transfer and gene loss, a
finding suggesting that the evolution and life cycle of RM systems may be best described as that of
a selfish genetic element. A putative target motif (CTAG) of one of the orphan methylases was
underrepresented in all of the analyzed genomes, whereas another motif (GATC) was
overrepresented in most of the haloarchaeal genomes, particularly in those that encoded the cognate
orphan methylase.

Keywords: HGT, Restriction, Methylation, Gene Transfer, Selfish Genes, Archaea, Haloarchaea,
DNA methylase, epigenetics

Funding: This work was supported through grants from the Binational Science Foundation (BSF
2013061); the National Science Foundation (NSF/MCB 1716046) within the BSF-NSF joint research
program; and NASA exobiology (NNX15AM09G, and 80NSSC18K1533).

34 1. Introduction

DNA methyltransferases (MTases) are enzymes which catalyze the addition of a methyl group
to a nucleotide base in a DNA molecule. These enzymes will methylate either adenine, producing
*N*6-methyladenine (6mA), or cytosine, producing either *N*4-methylcytosine (4mC) or *C*5-
methylcytosine (5mC), depending on the type of MTase enzyme [1]. DNA methyltransferases
typically consist of three types of protein domains: an S-adenosyl-L-methionine (AdoMet) binding
domain which obtains the methyl group from the co-factor AdoMet, a target recognition domain
(TRD) which binds the enzyme to the DNA strand at a short nucleotide sequence known as the
recognition sequence, and a catalytic domain which transfers the methyl group from AdoMet to a

nucleotide at the recognition sequence [2]. The order in which these domains occur in a MTase varies
and can be used to classify the enzymes into the subtypes of α, β, γ, δ, ε, and ζ MTases [3–5].
45
In bacteria and archaea, MTases are often components of restriction-modification (RM) systems,
in which an MTase works alongside a cognate restriction endonuclease (REase) that targets the same
recognition site. The REase will cleave the recognition site when it is unmethylated, but the DNA will
escape cutting when the site has been methylated by the MTase; this provides a self-recognition
system to the host where it differentiates between its own methylated DNA and that of unmethylated,
potentially harmful foreign DNA that is then digested by the host's REase [6–8]. RM systems have
also been described as addiction cassettes akin to toxin-antitoxin systems, in which post-
segregational killing occurs when the RM system is lost since the MTase activity degrades more
quickly than REase activity, resulting in digestion of the host genome at unmodified recognition sites
[9,10].   RM systems have been hypothesized to act as barriers to genetic exchange and drive
population diversification [11,12]. In *Escherichia coli*, for example, conjugational uptake of plasmids is
reduced by the RM system EcoKI when the plasmids contain EcoKI recognition sequences [13].
However, transferred DNA that is digested by a cell's restriction endonuclease can still effectively
recombine with the recipient's chromosomal DNA [7,14,15]; the effect of DNA digestion serves to
limit homologous recombinant DNA fragment size [16]. Restriction thus advantages its host by
decreasing transfer of large mobile genetic elements and infection with phage originating in
organisms without the cognate MTase [8], while also reducing linkage between beneficial and slightly
deleterious mutations [17].
64
There are four major types of RM systems which have been classified in bacteria and archaea
[18,19]. Type I RM systems consist of three types of subunits: REase (R) subunits, MTase (M) subunits,
and site specificity (S) subunits which contain two tandem TRDs. These subunits form pentamer
complexes of two R subunits, two M subunits, and one S subunit, and these complexes will either
fully methylate recognition sites which are modified on only one DNA strand (hemimethylated) or
cleave the DNA several bases upstream or downstream of recognition sites which are unmethylated
on both strands [20,21]. The MTases and REases of Type II RM systems have their own TRDs and
operate independently of each other, but each one targets the same recognition site [22]. There are
many different subclasses of Type II RM system enzymes, such as Type IIG enzymes which contain
both REase and MTase domains and are therefore capable of both methylation and endonuclease
activity [23]. Type III RM systems consist of REase (Res) and MTase (Mod) subunits which work
together as complexes, with the Mod subunit containing the TRD which recognizes asymmetric target
sequences [24]. Type IV RM systems are made up of only REases, but unlike in other RM systems,
these REases will target and cleave methylated recognition sites [20,25].
79
MTases can also exist in bacterial and archaeal hosts as orphan MTases, in which they occur
independently of cognate restriction enzymes and typically have important physiological functions
[26]. In *E. coli*, the orphan MTase Dam, an adenine MTase which targets the recognition sequence
GATC, is involved in regulating the timing of DNA replication by methylating the GATC sites
present at the origin of replication (*oriC*) [27]. The protein SeqA binds to hemimethylated GATC sites
at *oriC*, which prevents re-initiation of DNA replication at *oriC* after a new strand has been
synthesized [28,29]. Dam methylation is also important in DNA repair in *E. coli*, where the
methylation state of GATC sites is used by the methyl-directed mismatch repair (MMR) system to
identify the original DNA strand in order to make repairs to the newly-synthesized strand [30–32].
In *Cauldobacter crescentus*, the methylation of target sites in genes such as *ctrA* by orphan adenine
MTase CcrM helps regulate the cell cycle of the organism [33–35]. The importance of orphan MTases
in cellular processes is likely the reason why they are more widespread and conserved in bacteria
compared to MTases associated with RM systems [36,37].
93

MTases and RM systems have been well-studied in the bacteria, but less research has been
performed in archaea, with most studies focused on characterizing RM systems of thermophilic
species [38–42]. Recent research into the halophilic archaeal species *Haloferax volcanii* has
demonstrated a role for DNA methylation in DNA metabolism, and probably uptake: cells could not
grow on wild type *E. coli* DNA as a phosphorous source, whereas unmethylated *E. coli* was
metabolized completely [43,44].   In an effort to better understand this phenomenon, we
characterized the genomic methylation patterns (methylome) and MTases in the halophilic archaeal
species *Haloferax volcanii* [45,46]. However, the distribution of RM systems and MTases among the
archaea has not been extensively studied, and thus their life histories and impact on host evolution
are unclear.
104
To that end we surveyed the breadth of available genomes from public databases representing
the class Halobacteria, also known as the Haloarchaea, for RM system and MTase candidate genes.
We further sequenced additional genomes from the genus *Halorubrum* which provided an
opportunity to examine patterns among very closely related strains. Upon examining their patterns
of occurrence, we discovered orphan methyltransferases widely distributed throughout the
Haloarchaea.   In contrast, RM system candidate genes had a sparse and spotty distribution
indicating frequent gene transfer and loss.   Even individuals from the same species isolated from
the same environment and at the same time, differed in the RM system complement.

2. Materials and Methods

Search Approach. The starting data consists of 217 Halobacteria genomes from NCBI and 14 in-
house sequenced genomes (Supplementary Table S1**)**. We note that some of these genomes were
assembled from shotgun metagenome sequences and not from individual cultured strains. Genome
completion was determined through identification of 371 Halobacteriaceae marker genes using
CheckM v1.0.7 [47]. Queries for all restriction-methylation-specificity genes were obtained from the
Restriction Enzyme dataBASE (REBASE) website [48,49]. As methylation genes are classified by
function rather than by homology [48] the protein sequences of each category were clustered into
homologous groups (HGs) via the *uclust* function of the USEARCH v9.0.2132 package [50] at a 40
percent identity.   The resulting ~36,000 HGs were aligned with MUSCLE v3.8.31 [51]. HMMs were
then generated from the alignments using the *hmmbuild* function of HMMER3 v3.1b2 (hmmer.org).
The ORFs of the 217 genomes were searched against the profiles via the *hmmsearch* function of
HMMER3. Top hits were extracted and cross hits filtered with in-house Perl scripts. Steps were taken
to collapse and filter HGs. First, the hits were searched against the arCOG database [52] using BLAST
[53] to assign arCOG identifiers to the members of each group. Second the R package *igraph* v1.2.2
[54] was used to create a list of connected components from the arCOG identifications. All members
of a connected component were collapsed into a single collapsed HG (cHG).
130
Because REBASE is a database of all methylation-restriction-related activities there are many
members of the database outside our interest. At this point we made a manual curation of our cHGs
attempting to identify known functions that did not apply to our area of interest. Examples include
protein methylation enzymes, exonucleases, cell-division proteins, etc. The final tally of this
clustering and filtering yielded 1696 hits across 48 total candidate cHGs. arCOG annotations indicate
DNA methylase activity, restriction enzyme activity, or specificity module activity as part of an RM
system for 26 cHGs. The remaining 22 cHGs had predominant arCOG annotations matching other
functions that may reasonably be excluded from conservative RM system-specific analyses. For a
graphical representation of the search strategy see supplementary materials **Figure S1.** The putative
Type IV methyl-directed restriction enzyme gene, *mrr*, which is known to be present in *Haloferax*
volcanii had not been assembled into an cHG.   We assembled a cluster of *mrr* homologs and
determined their presence and absence using Mrr from *Haloferax volcanii* DS2 (accession:
ADE02322.1) as query in BLASTP searches against each genome (E-value cut-off 10[-10]).
144

Reference Phylogeny. A reference tree was created using the full complement of ribosomal
proteins. The ribosomal protein set for *Halorubrum lacusprofundi* ATCC 49239 was obtained from the
BioCyc website [55]. Each protein orf was used as the query in a BLAST [53] search against each
genome. Hits for each gene were aligned with MUSCLE v3.8.31 [51] and then concatenated with in-
house scripting. The concatenated alignment was subjected to maximum likelihood phylogenetic
inference in the IQ-TREE v1.6.1 suite with ultrafast bootstrapping and automated model selection
[56,57]. The final model selection was LG+F+R9.
152
F81 Presence-Absence Phylogeny. It is desirable to use maximum-likelihood methodology
rather than simple distance measures. To realize this, the matrix was converted to an A/T alignment
by replacing each present with an "A" and absent with a "T." This allowed use of an F81 model with
empirical base frequencies. This confines the base parameters to only A and T while allowing all of
the other advantages of an ML approach. IQ-TREE was employed to infer the tree with 100 bootstraps
[57].
159
Horizontal Gene Transfer Detection. Gene trees for each of the cHGs were inferred using
RAxML v8.2.11 [58] under PROTCATLG models with 100 bootstraps. The gene trees were then
improved by resolving their poorly supported in nodes to match the species tree using TreeFix-DTL
[59]. Optimized gene tree rootings were inferred with the OptRoot function of Ranger-DTL.
Reconciliation costs for each gene tree were computed against the reference tree using Ranger-DTL
2.0 (http://compbio.engr.uconn.edu/software/RANGER-DTL/) [60] with default DTL costs. One-
hundred reconciliations, each using a different random seed, were calculated for each cHG. After
aggregating these with the AggregateRanger function of Ranger-DTL the results were summarized
and each prediction and any transfer inferred in 51% or greater of cases was counted as a transfer
event.
170
Data Analysis and Presentation: The presence-absence matrix of cHGs was plotted as a
heatmap onto the reference phylogeny using the *gheatmap* function of the R Bioconductor package
ggtree v1.14.4 [61,62].   The rarefaction curve was generated with the *specaccum* function of the
vegan v2.5-3 package in R [63] and number of genomes per homologous group was plotted with
ggplot2 v3.1.0 [64]. Spearman correlations and significances between the presence-absence of cHGs
was   calculated   with   the   *rcorr*   function   of   the   *hmisc*   v4.1-1   package   in   R
(http://biostat.mc.vanderbilt.edu/wiki/Main/Hmisc). A significance cutoff of $p < 0.05$ was used with
a Bonferroni correction. All comparisons failing this criterion were set to correlation = 0. These data
were plotted into a correlogram via the *corrplot* function of the R package *corrplot* v0.84.   To compare
the Phylogeny calculated from Presence-Absence data to the ribosomal protein reference, the
bootstrap support set of the presence-absence phylogeny was mapped onto the ribosomal protein
reference tree using the *plotBS* function in *phangorn* v2.4.0 [65]. Support values equal to or greater
than 10% are displayed.   To compare phylogenies using Internode Certainty, scores were
calculated using the IC/TC score calculation algorithm implemented in RAxML v8.2.11 [58,66].
185
186   **Synteny.** Genomes were searched for location of cHGs. Proximity was used to determine
187   synteny of groups of cHGs frequently identified on the same genomes.
188
189   **Presence-Absence PCoA.** Jaccard distances between presence-absence of taxa were calculated
190   using the *distance* function of the R package *philentropy* v0.2.0 [67]. The PCoA was generated using
191   the *wcmdscale* function in *vegan* v2.5-3 [63]. The two best sampled genera, *Halorubrum* (orange) and
192   *Haloferax* (red), are colored distinctively.
193
194   **Recognition Site Assignment.** To determine the most likely recognition sites, each member of
195   each cHG was searched against the REBASE Gold Standard set using BLASTp. The REBASE gold
196   standard set was chosen over the individual gene sets on account of it having a much higher density

197 of recognition site annotation. This simplifies the need to search for secondary hits to find predicted
198 target sites. After applying an e-value cut-off of 1E-20, the top hit was assigned to each ORF.
199
200   **CTAG and GATC** motifs were counted with an inhouse perl script available at the Gogarten-
201 lab's GitHub [68].
202
203   **Gene Ontology.** Sets of GO terms were identified for each cHG using Blast2GO [69].
204 Annotations were checked against the UniProt database [70] using arCOG identifiers.

3. Results

RM-system gene distribution

Analysis of 217 haloarchaeal genomes and metagenome assembled genomes yielded 48 total
candidate collapsed homologous groups (cHGs) of RM-system components.   Out of these 48 cHGs,
26 had arCOG annotation suggesting DNA methylase activity, restriction enzyme activity, or
specificity module activity as part of an RM system. We detected 22 weaker candidates with
predominant arCOG annotations matching other functions (**Table 1**). Our analysis shows that nearly
all of the cHGs are found more than once. (**Figure 1A**). Indeed, 16 families are found in 20 or more
genomes each (>9 %), and this frequency steadily increases culminating in five families being
conserved in greater than 80 genomes each (>37 %) with one cHG being in ~80 % of all Haloarchaea
surveyed. Though these genes appear frequently in taxa across the haloarchaeal class, the majority of
each candidate RM system cHG is present in fewer than half the genomes, - the second most
abundantly recovered cHG is found in only ~47 % of all taxa surveyed.   We note that the cHGs with
wide distribution are annotated as MTases without an identifiable co-evolving restriction
endonuclease: Group U DNA_methylase-022; W dam_methylase-031; Y dcm_methylase-044; and AT
Uncharacterized-032 (members of this cHG are also annotated as methylation subunit and N6-
Adenine MTase). Rarefaction analysis indicates about 50 % of the genomes assayed contain seven
dominant cHGs, and that all taxa on average are represented by half of the cHGs (**Figure 1B**).
Together, the separate analyses indicate extensive gene gain and loss of RM-system genes.   In
contrast, orphan MTases in cHG U and W, and to a lesser extent Y (Figure 2) have a wider distribution
in some genera (see below for further discussion).
226
The phylogeny of the class Halobacteria inferred from concatenated ribosomal proteins (**Figure
2)** was largely comparable to prior work [71], and with a taxonomy based on concatenations of
conserved proteins [72,73].   For instance, in our phylogeny the *Halorubracaea* group with the
Haloferacaceae recapitulating the order *Haloferacales,* and the families, *Halobacteriaceae, Haloarculaceae*
and *Halococcaceae* group within the order *Halobacteriales.*   Our genome survey in search of RM-
system genes encompassed a broad taxonomic sampling, and it explores in depth the genus
Halorubrum because it is a highly speciated genus, and because the existence of many genomes from
the same species allows within species distribution assessment.
235
Comparison of the phylogeny in **Figure 2** to the heatmap giving the presence/absence of RM
system cHG candidates demonstrates that the cHG distribution is highly variable (**Figure 2)**.   The
one glaring exception is cHG U, a DNA methylase found in 174 of the 217 genomes analyzed.   Since
it is not coupled with a restriction enzyme of equal abundance, it is assumed to be an orphan MTase.
The MTase from *Hfx. volcanii* (gene HVO_0794), which recognizes the CTAG motif [45] is a member
of this cHG. Though U is widely distributed, within the genus *Halorubrum* it is only found in ~37.5 %
(21/56) of the genomes.   While U's phylogenetic profile is compatible with vertical inheritance over
much of the phylogeny, the presence absence data also indicate a few gene transfer and loss events
within *Halorubrum.*   cHG U is present in *Hrr. tebenquichense* DSM14210, *Hrr. hochstenium*
ATCC700873, *Hrr.* sp. AJ767, and in strains from the related species *Hrr. distributum, Hrr. arcis, Hrr.*
litoreum and *Hrr. terrestre* suggesting an acquisition in the ancestor of this group.

247
Instead of U, another orphan MTase is abundantly present in *Halorubrum* spp., cHG W. It was
found in ~95 % of all *Halorubrum* strains, with three exceptions - an assembled genome from
metagenome sequence data, and two from incomplete draft genomes of the species *Halorubrum*
ezzemoulense. Interestingly, when U is present in a *Halorubrum* sp. genome, so too is W **(Figure 2)**.
In a complementary fashion, analysis of W outside of the *Halorubrum* shows that it is found patchily
distributed throughout the rest of the class Halobacteria (~20 % -32/158), and always as a second
orphan MTase with cHG U. When the members of cHG W were used to search the uniprot database,
the significant matches included the *E. coli* Dam MTase, a very well-characterized GATC MTase,
which provides strong evidence that this cHG is a GATC orphan MTase family. The presence and
absence of cHG U and W in completely sequenced genomes is given in Table S3, together with the
frequency of the CTAG and GATC motifs in the main chromosome.
259
The rest of the RM cHGs are much more patchily distributed (**Figure 2**). For instance, the cHGs
that make up columns A-G represent different gene families within the Type I RM system
classification; two MTases (A,B), three REases (C,D,E), and two site specificity units (SSUs) (F,G).
Throughout the Haloarchaea, cHGs from columns A, E and F, representing an MTase, an REase, and
an SSU respectively, are found co-occurring 35 times. In a subset of genomes studied for synteny,
A, E, and F are encoded next to one another in *Natrinema gari, Halorhabdus utahensis, Halorubrum*
SD690R, *Halorubrum ezzemoulense* G37, and *Haloorientalis* IM1011 (**Figure 3**). These genes probably
represent a single transcriptional unit of genes working together for restriction and modification
purposes. Since the Type I RM system is a five-component system, the likely stoichiometry is 2:2:1.
These three cHGs co-occur four times within the species *Halorubrum ezzemoulense*, and two of these
cHGs (A and E) co-occur an additional three more times, suggesting either a loss of the SSU, or an
incomplete genome sequence for those strains. If it is due to incomplete sequencing, then 7/16 (43
%) of the *Hrr. ezzemoulense* genomes have this set of co-occurring genes, while half do not have an
identified Type I system. This is particularly stunning since strains FB21, Ec15, G37 and Ga2p were
all cultivated at the same time from the same sample, a hypersaline lake in Iran. Furthermore, one
strain, Ga36, has a different identified Type I RM system composed of substituted cHGs A and E with
B and D, respectively, while maintaining the same SSU. This suggests the same DNA motif may be
recognized by the different cHGs and that these cHGs are therefore functionally interchangeable.
Members of cHGs B, F, and D were found as likely co-transcribed units in *Halococcus salifodinae,*
Natronolimnobius aegyptiacus, Halorubrum kocurii, Haloarcula amylolytica (**Figure 3**). In *Halorubrum*
DL, and *Halovivax ruber* XH70, genomes that contained members from cHGs A, B, D, E, and F these
genes were not found in a single unit, suggesting that they do not form a single RM system.
Together, these analyses suggest this Type I RM system has a wide but sporadic distribution, that
this RM system is not required for individual survival, and that functional substitutions occur for
cHGs.
285
Type II RM systems contain an MTase and an REase that target the same motif but do not require
an associated SSU because each enzyme has its own TRD. The Type II RM system cHGs are in
columns H-L for the MTases, and M-P for the REases. Memberships to the Type II MTase cHGs are
far more numerous in the Haloarchaea than their REase counterpart, as might be expected when
witnessing decaying RM systems through the loss of the REase. The opposite result, more REases is
a more difficult scenario because an unmethylated host genome would be subject to restriction by the
remaining cognate REase (e.g., addiction cassettes). There are 14 "orphan" Type II REases in **Figure**
2, but their cognate MTase's absence could be explained by incomplete genome sequence data.
294
Type III RM systems have been identified in cHGs Q (MTase) and R and S (REases). Type III
MTases and REases (cHGs Q and R) co-occur almost exclusively in the species *Halorubrum*
ezzemoulense, our most highly represented taxon. Furthermore, these Type III RM systems are highly
restricted in their distribution to that species, with cHGs co-occuring only twice more throughout the

Haloarchaea, and with a different REase cHG (S); once in *Halorubrum arcis*, and another in
Halobacterium D1. Orphan MTases occurred twice in cHG Q. Of particular interest is that closely
related strains also cultivated from Lake Bidgol in Iran but which are in a different but closely related
Halorubrum species (e.g., Ea8, IB24, Hd13, Ea1, Eb13) do not have a Type III RM system, implying
though exposed to the same halophilic viruses, they do not rely on this system for avoiding virus
infection.
305
Mrr is a Type IV REase that was suggested to cleave methylated GATC sites [74,75]. Mrr
homologs are identified in most *Haloferax* sp., they have a sporadic distribution among other
Haloferacaceae and in the *Halobacteriaceae,* and are absent in the *Natrialbaceae* (**Figure 2**). cHGs Z-AV
are not sufficiently characterized to pinpoint of their role in DNA RM systems or as MTase. These
cHGs likely include homing endonucleases or enzymes modifying nucleotides in RNA molecules;
however, their function as orphan MTases or restriction endonucleases can, at present, not be
excluded.
313

Horizontal Gene Transfer explains patchy distribution

The patchy appearance of RM system candidates was further investigated by plotting the
Jaccard distance of the presence-absence data against the alignment distance of the reference tree
(supplementary **Figure S2)**. If the presence-absence data followed vertical descent one would expect
the best-fit line to move from the origin with a strong positive slope. Instead, the best fit line is close
to horizontal with an r-squared value of 0.0047, indicating negligible relationship between the overall
genome phylogeny and RM system complement per genome. The presence-absence clustering
patterns were visualized by plotting a principle coordinate analysis **(**supplementary **Figure S3)**. The
high degree of overlap between the ranges of the three groups illustrates that there are few RM
system genes unique to a given group and a large amount of overlap in repertoires.
324

To further evaluate the lack of long-term vertical descent for RM system genes, a phylogeny was
inferred from the presence-absence pattern of cHGs. The resultant tree (**Figure S4)** is largely in
disagreement with the reference phylogeny. The bootstrap support set from the presence-absence
phylogeny was mapped onto the ribosomal topology (**Figure S5)**. The resulting support values
demonstrate an extremely small degree of agreement between the two methods. The few areas where
there is even 10% support are near the tips of the ribosomal phylogeny and correspond to parts of
established groups, such as *Haloferax*, *Natronobacterium*, and *Halorubrum*.   Internode Certainty (IC)
scores are another way to compare phylogenies. An average IC score of 1 represents complete
agreement between the two phylogneies, and score of -1 complete disagreement.   The average IC
scores for the reference tree using the support set from the F81 tree was -0.509, illustrating that the
presence absence data do not support the topology of the reference phylogeny.
336

The patchy distribution of the RM system candidate genes and their lack of conformity to the
reference phylogeny suggests frequent horizontal gene transfer combined with gene loss events as
the most probable explanation for the observed data. To quantify the amount of transfer, the TreeFix-
Ranger pipeline was employed. TreeFix-DTL resolves poorly supported areas of gene trees to better
match the concatenated ribosomal protein gene tree used as reference. Ranger-DTL resolves optimal
gene tree rooting against the species tree and then computes a reconciliation estimating the number
of duplications, transfers, and losses that best explains the data (**Table 2**). For almost every cHG with
four or more taxa, our analysis infers several HGT events. Only cHG R, a putative Type III restriction
enzyme found only in a group of closely related *Halorubrum ezzemoulense strains,* has not been inferred
to undergo at least one transfer event.
347

RM systems usually function as cooperative units [48,76,77]. It stands to reason that some of the
RM system candidates may be transferred as units, maintaining their cognate functionality. This

possibility was examined by a correlation analysis. A spearman correlation was made between all
pairs of cHGs. Those with a significant result at a Bonferroni-corrected $p < 0.05$ were plotted in a
correlogram (**Figure 4)**. As illustrated in **Figure 3**, cHGs with significant similar phylogenetic profiles
often are near to one another in the genomes.
354

3.2. Figures and Tables

Figure 1: Distribution of collapsed Homologous Group (cHG) among haloarchaeal genomes. (A) the
number of genomes present in each collapsed Homologous Group (cHG). No cHG contains a
representative from every genome used in this study. With the exception of one cHG, all contain
members from fewer than half of the genomes.   The cHGs are ordered by number of genomes they
contain. (B) rarefaction plot of the number of genomes represented as cHGs accumulate.   95%
confidence interval is shown in shaded blue area and yellow box whisker plots give the number of
taxa from random subsamples (permutations = 100) over 48 gene families.

Figure 2: Presence-absence matrix of the 48 candidate RMS cHGs plotted against the reference phylogeny. For most cHGs the pattern of presence-absence does not match the reference phylogeny (compare supplementary Figures S2-S5) RMS-candidate cHGs are loosely ordered by system type and with the ambiguously assigned RM candidates at the end. Above is a key relating the column names to the majority functional annotation.

372
Figure 3. Gene maps for syntenic clusters of gene families (A) EFA and (B) BFD found in a subset of
organisms identified to the right of each map.   Genes are colored by gene families with Type I
methylases (AB) in greys, Type I restriction endonucleases (DE) in blues, and Type I site specificity
unit (F) in green.

Figure 4. Heatmap of co-occurrence between the 48 RMS-candidate cHGs. Positive correlation
indicates the cHGs co-occur while negative indicates that the presence of one means the other will not
be present. Significance level is p < 0.05 with a Bonferroni correction applied for multiple tests. Blue
indicates significant positive correlation; red indicates a significant negative correlation.

Table 1. Collapsed homologous group descriptions. $^\$$

Alpha code	Numerical code	Annotated arCOG Function$^{\$\$}$	arCOG number
A	cHG 021	T I M	arCOG02632
B	cHG 024	T I M	arCOG05282
C	cHG 018	T I R	arCOG00880
D	cHG 034	T I R	arCOG00879
E	cHG 045	T I R	arCOG00878
F	cHG 006	T I S	arCOG02626
G	cHG 025	T I S	arCOG02628
H	cHG_036	probable_T_II_M	arCOG00890
I	cHG 001	T II M	arCOG02635
J	cHG 003	T II M	arCOG02634
K	cHG_011	T_II_M	arCOG04814
L	cHG 033	T II M	arCOG03521
M	cHG 007	T II R	arCOG11279
N	cHG 013	T II R	arCOG11717
O	cHG 023	T II R	arCOG03779
P	cHG 029	T II R	arCOG08993
Q	cHG 042	Adenine DNA methylase probable T III M	arCOG00108
R	cHG 008	T III R	arCOG06887
S	cHG 009	T III R probable	arCOG07494
T	cHG 014	Adenine DNA methylase	arCOG00889
U	cHG 022	DNA methylase	arCOG00115
V	cHG 027	DNA methylase	arCOG00129
W	cHG 031	dam methylase	arCOG03416
X	cHG 035	probable RMS M	arCOG08990
Y	cHG 044	dcm methylase	arCOG04157
Z	cHG 048	Adenine DNA methylase	arCOG02636
AA	cHG_010	RNA_methylase	arCOG00910
AB	cHG 040	SAM-methylase	arCOG01792
AC	cHG 012	RestrictionEndonuclease	arCOG05724
AD	cHG_038	PredictedRestrictionEndonuclease	arCOG06431
AE	cHG 015	HNH endonuclease	arCOG07787
AF	cHG 019	Endonuclease	arCOG02782
AG	cHG 020	Endonuclease	arCOG02781
AH	cHG 004	HNH endonuclease	arCOG09398
AI	cHG 037	HNH nuclease	arCOG05223
AJ	cHG 039	HNH nuclease	arCOG03898
AK	cHG 041	HNH nuclease	arCOG08099
AL	cHG 046	MBF1	arCOG01863
AM	cHG 028	CBS domain	arCOG00608
AN	cHG 005	MarR	arCOG03182
AO	cHG 030	ParB-like nuclease	arCOG01875
AP	cHG 016	GVPC	arCOG06392
AQ	cHG 002	ASCH domain RNA binding	arCOG01734
AR	cHG_017	Uncharacterized	arCOG10082
AS	cHG 026	Uncharacterized	arCOG13171
AT	cHG 032	Uncharacterized	arCOG08946
AU	cHG_043	Uncharacterized	arCOG08856
AV	cHG_047	Uncharacterized	arCOG04588

383    $^\$$: A listing of associated Gene Ontology terms and gene family descriptions is available in
384   supplementary Table S2
385    $^{\$\$}$: T_I and T_II denote type I and type II restriction enzyme, respectively. M, R, S denote
386   the methylase, restriction endonuclease, and specificity subunits, respectively.

Table 2. Important traits of cHGs with four or more ORFs.

Alpha (numeric) cHG	No. of taxa	No. of transfers[a]	Function[b]	Predicted Recognition sites[c]	Frequency[e]
I (001)	16	9	T_II_M	GAAGGC	31%
				GGRCA	31%
J (003)	38	21	T_II_M	CANCATC	53%
				TAGGAG	21%
AH (004)	12	4	HNH_endonuclease	GGCGCC	89%
				GATC	11%
F (006)	61	44	T_I_S	GGAYNNNNNNTGG	24%
				CAGNNNNNNTGCT	16%
R (008)	14	0	T_III_R	NA[d]	100%
AA (010)	55	15	RNA_methylase	ATTAAT	33%
K (011)	137	97	T_II_M	GCAAGG	49%
				GKAAYG	28%
AC (012)	8	5	Restriction Endonuclease	GCGAA	29%
				CAACNNNNNTC	29%
				CTGGAG	29%
T (014)	130	93	Adenine_DNA_methylase	GCAGG	45%
				AAGCTT	32%
AE (015)	21	13	HNH_endonuclease	GGCGCC	70%
				YSCNS	15%
AP (016)	12	6	GVPC	CANCATC	83%
C (018)	7	4	T_I_R	AACNNNNNNGTGC	73%
				CTANNNNNNRTTC	27%
AF (019)	4	3	Endonuclease	NA[d]	100%
A (021)	88	58	T_I_M	GGAYNNNNNNTGG	37%
				GTCANNNNNNRTCA	12%
				CTCGAG	9%
U (022)	290	120	DNA_methylase	CTAG	59%
				CATTC	14%
				CCCGGG	7%
O (023)	37	28	T_II_R	NA[d]	100%
B (024)	16	8	T_I_M	GAGNNNNNVTGAC	75%
				GACNNNNNNRTAC	19%
G (025)	4	2	T_I_S	GAGNNNNRTAA	75%
				GAGNNNNNTAC	25%
V (027)	5	1	DNA_methylase	CATTC	100%
AO (030)	4	2	ParB-like_nuclease	GATC	75%
				CTAG	25%
W (031)	153	70	dam_methylase	GATC	70%
				AB / SAAM	22%

AT (032)	116	60	Uncharacterized	GCAAGG	43%
				GKAAYG	26%
				GGTTAG	14%
L (033)	66	38	T_II_M-033	CAARCA	40%
				CTGAAG	36%
D (034)	16	11	T_I_R-034	GCANNNNRTTA	69%
				GGCANNNNNTTC	19%
X (035)	19	9	probable_RMS_M	GGGAC	83%
H (036)	38	24	probable_T_II_M	CCWGG	42%
				CCSGG	18%
				GTAC	16%
AI (037)	6	4	HNH_nuclease	NA[d]	100%
AJ (039)	5	4	HNH_nuclease	GGCGCC	100%
AK (041)	6	4	HNH_nuclease	NA[d]	100%
Q (042)	21	8	Adenine_DNA_methylase	RGTAAT	71%
			probable_T_III_M	NA[d]	19%
Y (044)	179	110	dcm_methylase	CGGCCG	24%
				GTCGAC	13%
				ACGT	11%
E (045)	58	42	T_I_R	CCCNNNNNRTTGY	63%
				GCANNNNNRTTA	28%
Z (048)	54	35	Adenine_DNA_methylase	CCRGAG	36%
				GTMKAC	30%

[a] Number of estimated horizontal gene transfer events

[b] T_I and T_II denote type I and type II restriction enzyme, respectively. M, R, S denote the methylase, restriction endonuclease, and specificity subunits, respectively.

[c] Top predicted recognition sites

[d] No predicted recognition site

[e] Frequency of predictions within the cHG

4. Discussion

A striking result of our study is the irregular distribution of the RM system gene candidates
throughout not just the haloarchaeal class, but also within its orders, genera, species, and even
communities and populations. The patchy distribution is almost certainly the result of frequent HGT
and gene loss. RM system genes are well known for their susceptibility to HGT and loss, and their
presence almost never define a clade or an environmental source (e.g., [36,78]). Frequent acquisition
of RM system genes through HGT is illustrated by their sporadic distribution.   For example,
Halorubrum genomes encode many candidate RM system cHGs that are absent from the remainder
of the Halobacteria (e.g., cHG M, R, S, AC, AG, AM). Only one of these (cHG R) is found in more than
3 genomes, a Type III restriction protein found in 14 of 57 *Halorubrum* genomes. Mrr homologs have
a sporadic distribution among *Haloferacaceae* and <u>Halobacteriaceae</u> and are absent in *Natrialbaceae*
(**Figure 2**). Gene loss undoubtedly contributed to the sparse cHGs distribution; however, without
invoking frequent gene transfer, many independent and parallel gene losses need to be postulated.
We also observed that a number haloarchaeal species possess multiple Type I subunit genes, allowing
for functional substitution of the different subunits in the RM system. The existence of multiple Type
I subunits has also been observed in *Helicobacter pylori*, in which 4 different SSU loci are used by the
organism's Type I system to target different recognition sequences; these SSUs can even exchange
TRDs, resulting in variation in the methylome of *H. pylori*   [79–81].   In our results, however, we
observed multiple MTase and REase subunits alongside a single SSU, suggesting the functional
substitution of the subunits in these haloarchaeal organisms does not result in variation in detected
recognition sequences.

Mrr is a Type IV REase that cleaves methylated target sites. Studies have demonstrated that this
gene reduces transformation efficiency of GATC-methylated plasmids in *H. volcanii*, and that deletion
of the *mrr* gene increases transformation efficiency on GATC-methylated plasmids, suggesting that
this Type IV REase can target GATC-methylated sites for cleavage [74,75].   However, we find no
anticorrelation between the presence of Mrr homologs and members of cHG W, which is homologous
to the *E. coli* Dam MTase, a very well-characterized GATC MTase (**Figures 2 and 4**).   This suggests
that some members of cHG W or the Mrr homologs either are dysfunctional of have a site specificity
different from the GATC motif.

It seems counterintuitive that RM systems are not more conserved as cellular countermeasures
against commonly occurring viruses.   It may be that cells do not require extensive protection via
RM systems, because they use multiple defensive systems some of which might be more effective.
For example, another well-known defense against viruses is the CRISPR-Cas system [82]. CRISPR
recognizes short (~40bp) regions of invading DNA that the host has been exposed to previously and
degrades it. While it can be very useful against virus infection, our prior work indicated that CRISPR-
Cas was also sporadically distributed within communities of closely related haloarchaeal species [83]
indicating they are not required for surviving virus infection.

Both the RM and CRISPR-Cas systems are only important countermeasures after external
fortifications have failed to prevent a virus from infiltrating, and therefore their limited distributions
also indicate that the cell's primary defense would be in preventing virus infection altogether, which
is accomplished by different mechanisms. By altering surfaces via glycosylation cells can avoid virus
predation prior to infection. In *Haloferax* species there are two pathways which control glycosylation
of external features. One is relatively conserved and could have functions other than virus avoidance,
while the other is highly variable and shows hallmarks of having genes mobilized by horizontal
transfer [84]. At least one halovirus has been found to require glycosylation by its host in order to
infect properly [85]. Comparison of genomes and metagenomes from hypersaline environments
showed widespread evidence for distinct "genomic" islands in closely related halophiles [86] that
contain a unique mixture of LPS and other genes that contribute to altering the cell's surface structure

and virus docking opportunities. Thus selective pressure on post infection, cytosolic and nucleic
acids-based virus defenses is eased, allowing them to be lost randomly in populations.
443
A major consideration in understanding RM system diversity is that viruses, or other infiltrating
selfish genetic elements, might gain access to the host's methylation after a successful infection that
was not stopped by the restriction system.   Indeed, haloviruses are known to encode DNA
methyltransferases in their genomes (e.g., see [87]). In this case, RM systems having a limited within
population distribution would then be an effective defense for that part of the population possessing
a different RM system.   Under this scenario, a large and diverse pool of mobilized RM system genes
could offer a stronger defense for the population as a whole. A single successful infection would no
longer endanger the entire group of potential hosts.
452
Group selection may be invoked to explain the within population diversity of RM systems; a
sparse distribution of RM systems may provide a potential benefit to the population as a whole,
because a virus cannot easily infect all members of the population.   However, often gene level
selection is a more appropriate alternative to group selection [88,89].   Under a gene centered
explanation, RM systems are considered as selfish addiction cassettes that may be of little benefit to
its carrier.   While RM systems may be difficult to delete as a whole, stepwise deletion, that begins
with inactivation of the REase activity can lead to their loss from a lineage.   Their long-term survival
thus may be a balance of gain through gene transfer, persistence through addiction, and gene loss.
This gene centered explanation is supported by a study from [36], which examined the distribution
of MTase genes in ~1000 bacterial genomes. They observed, similar to our results in the Halobacteria,
that MTases associated with RM systems are poorly conserved, whereas orphan MTases share
conservation patterns similar to average genes. They also demonstrated that many RM-associated
and orphan MTases are horizontally acquired, and that a number of orphan MTases in bacterial
genomes neighbor degraded REase genes, suggesting that they are the product of degraded RM
systems that have lost functional REases [36].   Similarly, Kong et al. [78] studying genome content
variation in *Neisseria meningitidis* found an irregular distribution of RM systems, suggesting that these
systems do not form an effective barrier to homologous recombination within the species. Kong et al.
also observed that the RM systems themselves had been frequently transferred within the species.
We conclude that RM genes in bacteria as well as archaea appear to undergo significant horizontal
transfer and are not well-conserved. Only when these genes pick up additional functions, do parts of
these systems persist for longer periods of time, as exemplified in the distribution of orphan MTases.
However, the transition from RM system MTase to orphan MTase is an infrequent event.   A study
of 43 pan-genomes by Oliveira et al. [90] suggests that orphan MTases occur more frequently from
transfer via large mobile genetic elements (MGEs) such as plasmids and phages rather than arise *de*
novo from RM degradation.   The distribution of orphan methylase cHG U and W, and their likely
target motifs, CTAG and GATC, respectively suggests different biological functions for these two
methylases. The widespread conservation of the CTAG MTase family cHG U supports the findings
of Blow *et al.* [37] who identified a well-conserved CTAG orphan MTase family in the Halobacteria.
Similar to other bacterial and archaeal genomes [91], the CTAG motif, the likely target for methylases
in cHG U, is underrepresented in all haloarchaeal genomes (see table S3).   The low frequency of
occurrence, only about once per 4000 nucleotides, suggests that this motif and the cognate orphan
methylase are not significantly involved in facilitating mismatch repair.   The underrepresented
CTAG motif was found to be less underrepresented near rRNA genes [91] and on plasmids; the
CTAG motif also is a known target sequence for some IS elements [92]; and it may be involved in
repressor binding, where the CTAG motif was found to be associated with kinks in the DNA when
bound to the repressor [93,94]   Interestingly, CTAG and GATC motifs are absent, or
underrepresented in several haloarchaeal viruses [87,95,96].   Both, the presence of the cHG U
methylase and the underrepresentation of the CTAG motif, appear to be maintained by selection;
however, at present the reasons for the underrepresentation of the motif in chromosomal DNA, and
the role that the methylation of this motif may play remain open questions.

5. Conclusions

495   RM systems have a sporadic distribution in haloarchaea, even within species and populations.   In
496   contrast, orphan methylases are more persistent in lineages, and the targeted motifs are under
497   selection for lower (in case of CTAG) or higher (in case of GATC) than expected frequency.   In case
498   of the GATC motif, the cognate orphan MTase was found only in genomes where this motif occurs
499   with high frequency.

Supplementary Materials: The following are available online at www.mdpi.com/xxx/s1,

Figure S1. Workflow of RMS-candidate gene search strategy.

Figure S2. Plot of alignment distance as a function of presence-absence distance.

Figure S3. PCoA plot of the distances between the RMS presence-absence profiles of the 217 analyzed Halobacterial genomes.

Figure S4. Maximum-likelihood phylogeny of cHG presence-absence matrix.

Figure S5. Bootstrap support values of the presence-absence phylogeny mapped onto the ribosomal protein reference tree.

Table S1. Basic statistics for Halobacteriaceae complete and draft genomes

Table S2. Gene Ontology (GO) terms for each collapsed homologous group

Table S3. Distribution of orphan methylases cHGs U and W and frequency of their putative recognition motifs in completely sequenced halobacterial chromosomes.

Author Contributions: Conceptualization, Thane R. Papke and Johann Peter Gogarten; Data curation, Matthew S. Fullmer, Matthew Ouellette and Artemis S. Louyakis; Formal analysis, Matthew S. Fullmer, Matthew Ouellette and Artemis S. Louyakis; Funding acquisition, Thane R. Papke and Johann Peter Gogarten; Investigation, Matthew S. Fullmer and Matthew Ouellette; Methodology, Matthew S. Fullmer, Artemis S. Louyakis and Johann Peter Gogarten; Project administration, Thane R. Papke and Johann Peter Gogarten; Software, Matthew S. Fullmer and Artemis S. Louyakis; Supervision, Thane R. Papke and Johann Peter Gogarten; Validation, Matthew S. Fullmer; Visualization, Matthew S. Fullmer and Artemis S. Louyakis; Writing – original draft, Matthew S. Fullmer and Johann Peter Gogarten; Writing – review & editing, Matthew Ouellette, Artemis S. Louyakis, Thane R. Papke and Johann Peter Gogarten.

Funding: This work was supported through grants from the Binational Science Foundation (BSF 2013061); the National Science Foundation (NSF/MCB 1716046) within the BSF-NSF joint research program; and NASA exobiology (NNX15AM09G, and 80NSSC18K1533).

Acknowledgments: The Computational Biology Core, Institute for Systems Genomics, University of Connecticut provided computational resources. The authors thank an anonymous reviewer for pointing out the presence of Mrr in *Haloferax volcanii*.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Korlach, J.; Turner, S. W. Going beyond five bases in DNA sequencing. *Curr. Opin. Struct. Biol.* **2012**, *22*, 251–261, doi:10.1016/j.sbi.2012.04.002.

2. Bheemanaik, S.; Reddy, Y. V. R.; Rao, D. N. Structure, function and mechanism of exocyclic DNA methyltransferases. *Biochem. J.* **2006**, *399*, 177–190, doi:10.1042/BJ20060854.

3. Malone, T.; Blumenthal, R. M.; Cheng, X. Structure-guided Analysis Reveals Nine Sequence Motifs Conserved among DNA Amino-methyl-transferases, and Suggests a Catalytic Mechanism for these Enzymes. *J. Mol. Biol.* **1995**, *253*, 618–632, doi:10.1006/jmbi.1995.0577.

4. Bujnicki, J. M.; Radlinska, M. Molecular evolution of DNA-(cytosine-N4) methyltransferases: evidence for their polyphyletic origin. *Nucleic Acids Res.* **1999**, *27*, 4501–9.

5. Bujnicki, J. M. Sequence permutations in the molecular evolution of DNA methyltransferases. *BMC Evol. Biol.* **2002**, *2*, 3.

6. Tock, M. R.; Dryden, D. T. The biology of restriction and anti-restriction. *Curr. Opin. Microbiol.* **2005**, *8*, 466–472, doi:10.1016/j.mib.2005.06.003.

7. Vasu, K.; Nagaraja, V. Diverse functions of restriction-modification systems in addition to cellular defense. *Microbiol. Mol. Biol. Rev.* **2013**, *77*, 53–72, doi:10.1128/MMBR.00044-12.

8. Pleška, M.; Qian, L.; Okura, R.; Bergmiller, T.; Wakamoto, Y.; Kussell, E.; Guet, C. C. Bacterial Autoimmunity Due to a Restriction-Modification System. *Curr. Biol. CB* **2016**, *26*, 404–409, doi:10.1016/j.cub.2015.12.041.

9. Ohno, S.; Handa, N.; Watanabe-Matsui, M.; Takahashi, N.; Kobayashi, I. Maintenance Forced by a Restriction-Modification System Can Be Modulated by a Region in Its Modification Enzyme Not Essential for Methyltransferase Activity. *J. Bacteriol.* **2008**, *190*, 2039–2049, doi:10.1128/JB.01319-07.

10. Kobayashi, I. Behavior of restriction-modification systems as selfish mobile elements and their impact on genome evolution. *Nucleic Acids Res.* **2001**, *29*, 3742–56.

11. Budroni, S.; Siena, E.; Hotopp, J. C. D.; Seib, K. L.; Serruto, D.; Nofroni, C.; Comanducci, M.; Riley, D. R.; Daugherty, S. C.; Angiuoli, S. V.; Covacci, A.; Pizza, M.; Rappuoli, R.; Moxon, E. R.; Tettelin, H.; Medini, D. Neisseria meningitidis is structured in clades associated with restriction modification systems that modulate homologous recombination. *Proc. Natl. Acad. Sci.* **2011**, *108*, 4494–4499, doi:10.1073/pnas.1019751108.

12. Erwin, A. L.; Sandstedt, S. A.; Bonthuis, P. J.; Geelhood, J. L.; Nelson, K. L.; Unrath, W. C. T.; Diggle, M. A.; Theodore, M. J.; Pleatman, C. R.; Mothershed, E. A.; Sacchi, C. T.; Mayer, L. W.; Gilsdorf, J. R.; Smith, A. L. Analysis of Genetic Relatedness of Haemophilus influenzae Isolates by Multilocus Sequence Typing. *J. Bacteriol.* **2008**, *190*, 1473–1483, doi:10.1128/JB.01207-07.

13. Roer, L.; Aarestrup, F. M.; Hasman, H. The EcoKI Type I Restriction-Modification System in Escherichia coli Affects but Is Not an Absolute Barrier for Conjugation. *J. Bacteriol.* **2015**, *197*, 337–342, doi:10.1128/JB.02418-14.

14. McKane, M.; Milkman, R. Transduction, restriction and recombination patterns in Escherichia coli. *Genetics* **1995**, *139*, 35–43.

15. Chang, S.; Cohen, S. N. In vivo site-specific genetic recombination promoted by the EcoRI restriction

endonuclease. *Proc. Natl. Acad. Sci. U. S. A.* **1977**, *74*, 4811–5.

16.   Lin, E. A.; Zhang, X.-S.; Levine, S. M.; Gill, S. R.; Falush, D.; Blaser, M. J. Natural Transformation of

Helicobacter pylori Involves the Integration of Short DNA Fragments Interrupted by Gaps of Variable

Size. *PLoS Pathog.* **2009**, *5*, e1000337, doi:10.1371/journal.ppat.1000337.

17.   Muller, H. J. The relation of recombination to mutational advance. *Mutat. Res.* **1964**, *1*, 2–9.

18.   Ershova, A. S.; Rusinov, I. S.; Spirin, S. A.; Karyagina, A. S.; Alexeevski, A. V. Role of restriction-

modification systems in prokaryotic evolution and ecology. *Biochem.* **2015**, *80*, 1373–1386,

doi:10.1134/S0006297915100193.

19.   Roberts, R. J.; Belfort, M.; Bestor, T.; Bhagwat, A. S.; Bickle, T. A.; Bitinaite, J.; Blumenthal, R. M.;

Degtyarev, S. K.; Dryden, D. T. F.; Dybvig, K.; Firman, K.; Gromova, E. S.; Gumport, R. I.; Halford, S. E.;

Hattman, S.; Heitman, J.; Hornby, D. P.; Janulaitis, A.; Jeltsch, A.; Josephsen, J.; Kiss, A.; Klaenhammer,

T. R.; Kobayashi, I.; Kong, H.; Krüger, D. H.; Lacks, S.; Marinus, M. G.; Miyahara, M.; Morgan, R. D.;

Murray, N. E.; Nagaraja, V.; Piekarowicz, A.; Pingoud, A.; Raleigh, E.; Rao, D. N.; Reich, N.; Repin, V.

E.; Selker, E. U.; Shaw, P.-C.; Stein, D. C.; Stoddard, B. L.; Szybalski, W.; Trautner, T. A.; Van Etten, J. L.;

Vitor, J. M. B.; Wilson, G. G.; Xu, S. A nomenclature for restriction enzymes, DNA methyltransferases,

homing endonucleases and their genes. *Nucleic Acids Res.* **2003**, *31*, 1805–12.

20.   Loenen, W. A. M.; Dryden, D. T. F.; Raleigh, E. A.; Wilson, G. G. Type I restriction enzymes and their

relatives. *Nucleic Acids Res.* **2014**, *42*, 20–44, doi:10.1093/nar/gkt847.

21.   Liu, Y.-P.; Tang, Q.; Zhang, J.-Z.; Tian, L.-F.; Gao, P.; Yan, X.-X. Structural basis underlying complex

assembly and conformational transition of the type I R-M system. *Proc. Natl. Acad. Sci.* **2017**, *114*, 11151–

11156, doi:10.1073/pnas.1711754114.

22.   Pingoud, A.; Wilson, G. G.; Wende, W. Type II restriction endonucleases—a historical perspective and

more. *Nucleic Acids Res.* **2014**, *42*, 7489–7527, doi:10.1093/nar/gku447.

23.   Morgan, R. D.; Bhatia, T. K.; Lovasco, L.; Davis, T. B. MmeI: a minimal Type II restriction-modification

system that only modifies one DNA strand for host protection. *Nucleic Acids Res.* **2008**, *36*, 6558–6570,

doi:10.1093/nar/gkn711.

24.   Rao, D. N.; Dryden, D. T. F.; Bheemanaik, S. Type III restriction-modification enzymes: a historical

perspective. *Nucleic Acids Res.* **2014**, *42*, 45–55, doi:10.1093/nar/gkt616.

25.   Czapinska, H.; Kowalska, M.; Zagorskaitė, E.; Manakova, E.; Slyvka, A.; Xu, S.; Siksnys, V.; Sasnauskas,

G.; Bochtler, M. Activity and structure of EcoKMcrA. *Nucleic Acids Res.* **2018**, *46*, 9829–9841,

doi:10.1093/nar/gky731.

26.   Adhikari, S.; Curtis, P. D. DNA methyltransferases and epigenetic regulation in bacteria. *FEMS*

Microbiol. Rev. **2016**, *40*, 575–591, doi:10.1093/femsre/fuw023.

27.   Messer, W.; Bellekes, U.; Lother, H. Effect of dam methylation on the activity of the E. coli replication

origin, oriC. *EMBO J.* **1985**, *4*, 1327–32.

28.   Kang, S.; Lee, H.; Han, J. S.; Hwang, D. S. Interaction of SeqA and Dam methylase on the hemimethylated

origin of Escherichia coli chromosomal DNA replication. *J. Biol. Chem.* **1999**, *274*, 11463–8.

29.   Sanchez-Romero, M. A.; Busby, S. J. W.; Dyer, N. P.; Ott, S.; Millard, A. D.; Grainger, D. C. Dynamic

Distribution of SeqA Protein across the Chromosome of Escherichia coli K-12. *MBio* **2010**, *1*,

doi:10.1128/mBio.00012-10.

30.   Welsh, K. M.; Lu, A. L.; Clark, S.; Modrich, P. Isolation and characterization of the Escherichia coli mutH

gene product. *J. Biol. Chem.* **1987**, *262*, 15624–9.

31.   Au, K. G.; Welsh, K.; Modrich, P. Initiation of methyl-directed mismatch repair. *J. Biol. Chem.* **1992**, *267*,
12142–8.

32.   Putnam, C. D. Evolution of the methyl directed mismatch repair system in Escherichia coli. *DNA Repair*
(Amst). **2016**, *38*, 32–41, doi:10.1016/j.dnarep.2015.11.016.

33.   Zweiger, G.; Marczynski, G.; Shapiro, L. A Caulobacter DNA Methyltransferase that Functions only in
the Predivisional Cell. *J. Mol. Biol.* **1994**, *235*, 472–485, doi:10.1006/jmbi.1994.1007.

34.   Domian, I. J.; Reisenauer, A.; Shapiro, L. Feedback control of a master bacterial cell-cycle regulator. *Proc.*
Natl. Acad. Sci. U. S. A. **1999**, *96*, 6648–53.

35.   Gonzalez, D.; Kozdon, J. B.; McAdams, H. H.; Shapiro, L.; Collier, J. The functions of DNA methylation
by CcrM in Caulobacter crescentus: a global approach. *Nucleic Acids Res.* **2014**, *42*, 3720–3735,
doi:10.1093/nar/gkt1352.

36.   Seshasayee, A. S. N.; Singh, P.; Krishna, S. Context-dependent conservation of DNA methyltransferases
in bacteria. *Nucleic Acids Res.* **2012**, *40*, 7066–7073, doi:10.1093/nar/gks390.

37.   Blow, M. J.; Clark, T. A.; Daum, C. G.; Deutschbauer, A. M.; Fomenkov, A.; Fries, R.; Froula, J.; Kang, D.
D.; Malmstrom, R. R.; Morgan, R. D.; Posfai, J.; Singh, K.; Visel, A.; Wetmore, K.; Zhao, Z.; Rubin, E. M.;
Korlach, J.; Pennacchio, L. A.; Roberts, R. J. The Epigenomic Landscape of Prokaryotes. *PLOS Genet.*
2016, *12*, e1005854, doi:10.1371/journal.pgen.1005854.

38.   Nölling, J.; de Vos, W. M. Identification of the CTAG-recognizing restriction-modification systems
MthZI and MthFI from Methanobacterium thermoformicicum and characterization of the plasmid-
encoded mthZIM gene. *Nucleic Acids Res.* **1992**, *20*, 5047–52.

39.   Grogan, D. W. Cytosine methylation by the SuaI restriction-modification system: implications for genetic
fidelity in a hyperthermophilic archaeon. *J. Bacteriol.* **2003**, *185*, 4657–61.

40.   Ishikawa, K.; Watanabe, M.; Kuroita, T.; Uchiyama, I.; Bujnicki, J. M.; Kawakami, B.; Tanokura, M.;
Kobayashi, I. Discovery of a novel restriction endonuclease by genome comparison and application of a
wheat-germ-based cell-free translation assay: PabI (5′-GTA/C) from the hyperthermophilic archaeon
Pyrococcus abyssi. *Nucleic Acids Res.* **2005**, *33*, e112–e112, doi:10.1093/nar/gni113.

41.   Watanabe, M.; Yuzawa, H.; Handa, N.; Kobayashi, I. Hyperthermophilic DNA Methyltransferase
M.PabI from the Archaeon Pyrococcus abyssi. *Appl. Environ. Microbiol.* **2006**, *72*, 5367–5375,
doi:10.1128/AEM.00433-06.

42.   Couturier, M.; Lindås, A.-C. The DNA Methylome of the Hyperthermoacidophilic Crenarchaeon
Sulfolobus acidocaldarius. *Front. Microbiol.* **2018**, *9*, 137, doi:10.3389/fmicb.2018.00137.

43.   Chimileski, S.; Dolas, K.; Naor, A.; Gophna, U.; Papke, R. T. Extracellular DNA metabolism in Haloferax
volcanii. *Front. Microbiol.* **2014**, *5*, 57, doi:10.3389/fmicb.2014.00057.

44.   Zerulla, K.; Chimileski, S.; Näther, D.; Gophna, U.; Papke, R. T.; Soppa, J. DNA as a Phosphate Storage
Polymer and the Alternative Advantages of Polyploidy for Growth or Survival. *PLoS One* **2014**, *9*, e94819,
doi:10.1371/journal.pone.0094819.

45.   Ouellette, M.; Gogarten, J.; Lajoie, J.; Makkay, A.; Papke, R. Characterizing the DNA Methyltransferases
of Haloferax volcanii via Bioinformatics, Gene Deletion, and SMRT Sequencing. *Genes (Basel).* **2018**, *9*,
129, doi:10.3390/genes9030129.

46.   Ouellette, M.; Jackson, L.; Chimileski, S.; Papke, R. T. Genome-wide DNA methylation analysis of

Haloferax volcanii H26 and identification of DNA methyltransferase related PD-(D/E)XK nuclease family protein HVO_A0006. *Front. Microbiol.* **2015**, *6*, 251, doi:10.3389/fmicb.2015.00251.

47. Parks, D. H.; Imelfort, M.; Skennerton, C. T.; Hugenholtz, P.; Tyson, G. W. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res.* **2015**, gr-186072.

48. Roberts, R. J.; Macelis, D. REBASE—restriction enzymes and methylases. *Nucleic Acids Res.* **2001**, *29*, 268–269, doi:10.1093/nar/29.1.268.

49. Roberts, R. J.; Vincze, T.; Posfai, J.; Macelis, D. REBASE—a database for DNA restriction and modification: enzymes, genes and genomes. *Nucleic Acids Res.* **2015**, *43*, D298–D299, doi:10.1093/nar/gku1046.

50. Edgar, R. C. Search and clustering orders of magnitude faster than BLAST. *Bioinformatics* **2010**, *26*, 2460–2461, doi:10.1093/bioinformatics/btq461.

51. Edgar, R. C. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucl. Acids Res.* **2004**, *32*, 1792–1797.

52. Makarova, K. S.; Wolf, Y. I.; Koonin, E. V. Archaeal Clusters of Orthologous Genes (arCOGs): An Update and Application for Analysis of Shared Features between Thermococcales, Methanococcales, and Methanobacteriales. *Life (Basel, Switzerland)* **2015**, *5*, 818–840, doi:10.3390/life5010818.

53. Altschul, S. F.; Madden, T. L.; Schäffer, A. A.; Zhang, J.; Zhang, Z.; Miller, W.; Lipman, D. J. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acids Res.* **1997**, *25*, 3389–3402, doi:10.1093/nar/25.17.3389.

54. Csardi, G.; Nepusz, T. The igraph software package for complex network research. *InterJournal* **2006**, *Complex Sy*, 1695.

55. Caspi, R.; Altman, T.; Dale, J. M.; Dreher, K.; Fulcher, C. A.; Gilham, F.; Kaipa, P.; Karthikeyan, A. S.; Kothari, A.; Krummenacker, M.; Latendresse, M.; Mueller, L. A.; Paley, S.; Popescu, L.; Pujar, A.; Shearer, A. G.; Zhang, P.; Karp, P. D. The MetaCyc database of metabolic pathways and enzymes and the BioCyc collection of pathway/genome databases. *Nucleic Acids Res.* **2010**, *38*, D473–D479, doi:10.1093/nar/gkp875.

56. Hoang, D. T.; Chernomor, O.; von Haeseler, A.; Minh, B. Q.; Le, S. V. UFBoot2: Improving the Ultrafast Bootstrap Approximation. *Mol. Biol. Evol.*, doi:10.1093/molbev/msx281.

57. Nguyen, L.-T.; Schmidt, H. A.; von Haeseler, A.; Minh, B. Q. IQ-TREE: A Fast and Effective Stochastic Algorithm for Estimating Maximum-Likelihood Phylogenies. *Mol. Biol. Evol.* **2015**, *32*, 268–274, doi:10.1093/molbev/msu300.

58. Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **2014**, *30*, 1312–1313, doi:10.1093/bioinformatics/btu033.

59. Bansal, M. S.; Wu, Y.-C.; Alm, E. J.; Kellis, M. Improved gene tree error correction in the presence of horizontal gene transfer. *Bioinformatics* **2015**, *31*, 1211–1218, doi:10.1093/bioinformatics/btu806.

60. Bansal, M. S.; Alm, E. J.; Kellis, M. Efficient algorithms for the reconciliation problem with gene duplication, horizontal transfer and loss. *Bioinformatics* **2012**, *28*, i283–i291, doi:10.1093/bioinformatics/bts225.

61. Yu, G.; Smith, D. K.; Zhu, H.; Guan, Y.; Lam, T. T.-Y. ggtree : an r package for visualization and annotation of phylogenetic trees with their covariates and other associated data. *Methods Ecol. Evol.* **2017**,

8, 28–36, doi:10.1111/2041-210X.12628.

62.   Yu, G.; Lam, T. T.-Y.; Zhu, H.; Guan, Y. Two Methods for Mapping and Visualizing Associated Data on
Phylogeny Using Ggtree. *Mol. Biol. Evol.* **2018**, *35*, 3041–3043, doi:10.1093/molbev/msy194.

63.   Dixon, P. VEGAN, a package of R functions for community ecology. *J. Veg. Sci.* **2003**, *14*, 927–930,
doi:10.1111/j.1654-1103.2003.tb02228.x.

64.   Wickham, H. *ggplot2: elegant graphics for data analysis*; Springer, 2016; ISBN 3319242776.

65.   Schliep, K. P. phangorn: phylogenetic analysis in R. *Bioinformatics* **2011**, *27*, 592–3,
doi:10.1093/bioinformatics/btq706.

66.   Salichos, L.; Rokas, A. Inferring ancient divergences requires genes with strong phylogenetic signals.
Nature **2013**, *497*, 327–331, doi:10.1038/nature12130.

67.   Drost, H. Philentropy: Information Theory and Distance Quantification with R. *J. Open Source Softw.*
2018, *3*.

68.   Gogarten, J. P. Perl script to measure frequency and distribution of CTAG and GATC motifs in DNA
Available online: https://github.com/Gogarten-Lab/CTAG-GATC-frequencies (accessed on Jan 12, 2019).

69.   Conesa, A.; Götz, S.; García-Gómez, J. M.; Terol, J.; Talón, M.; Robles, M. Blast2GO: a universal tool for
annotation, visualization and analysis in functional genomics research. *Bioinformatics* **2005**, *21*, 3674–
3676.

70.   Consortium, U. UniProt: the universal protein knowledgebase. *Nucleic Acids Res.* **2016**, *45*, D158–D169.

71.   Soucy, S. M.; Fullmer, M. S.; Thane Papke, R.; Gogarten, J. P. Inteins as indicators of gene flow in the
halobacteria. *Front. Microbiol.* **2014**, *5*, 1–14, doi:10.3389/fmicb.2014.00299.

72.   Gupta, R. S.; Naushad, S.; Fabros, R.; Adeolu, M. A phylogenomic reappraisal of family-level divisions
within the class Halobacteria: proposal to divide the order Halobacteriales into the families
Halobacteriaceae, Haloarculaceae fam. nov., and Halococcaceae fam. nov., and the order Haloferacales
into th. *Antonie Van Leeuwenhoek* **2016**, *109*, 565–587, doi:10.1007/s10482-016-0660-2.

73.   Gupta, R. S.; Naushad, S.; Baker, S. Phylogenomic analyses and molecular signatures for the class
Halobacteria and its two major clades: a proposal for division of the class Halobacteria into an emended
order Halobacteriales and two new orders, Haloferacales ord. nov. and Natrialbales ord. n. *Int. J. Syst.*
Evol. Microbiol. **2015**, *65*, 1050–69, doi:10.1099/ijs.0.070136-0.

74.   Allers, T.; Barak, S.; Liddell, S.; Wardell, K.; Mevarech, M. Improved strains and plasmid vectors for
conditional overexpression of His-tagged proteins in Haloferax volcanii. *Appl. Environ. Microbiol.* **2010**,
76, 1759–1769.

75.   Holmes, M. L.; Nuttall, S. D.; Dyall-Smith, M. L. Construction and use of halobacterial shuttle vectors
and further studies on Haloferax DNA gyrase. *J. Bacteriol.* **1991**, *173*, 3807–3813.

76.   Ohno, S.; Handa, N.; Watanabe-Matsui, M.; Takahashi, N.; Kobayashi, I. Maintenance Forced by a
Restriction-Modification System Can Be Modulated by a Region in Its Modification Enzyme Not
Essential for Methyltransferase Activity. *J. Bacteriol.* **2008**, *190*, 2039–2049, doi:10.1128/JB.01319-07.

77.   Roberts, R. J.; Belfort, M.; Bestor, T.; Bhagwat, A. S.; Bickle, T. A.; Bitinaite, J.; Blumenthal, R. M.;
Degtyarev, S. K.; Dryden, D. T. F.; Dybvig, K.; Firman, K.; Gromova, E. S.; Gumport, R. I.; Halford, S. E.;
Hattman, S.; Heitman, J.; Hornby, D. P.; Janulaitis, A.; Jeltsch, A.; Josephsen, J.; Kiss, A.; Klaenhammer,
T. R.; Kobayashi, I.; Kong, H.; Krüger, D. H.; Lacks, S.; Marinus, M. G.; Miyahara, M.; Morgan, R. D.;
Murray, N. E.; Nagaraja, V.; Piekarowicz, A.; Pingoud, A.; Raleigh, E.; Rao, D. N.; Reich, N.; Repin, V.

E.; Selker, E. U.; Shaw, P.-C.; Stein, D. C.; Stoddard, B. L.; Szybalski, W.; Trautner, T. A.; Etten, V.; L, J.; Vitor, J. M. B.; Wilson, G. G.; Xu, S. A nomenclature for restriction enzymes, DNA methyltransferases, homing endonucleases and their genes. *Nucleic Acids Res.* **2003**, *31*, 1805–1812, doi:10.1093/nar/gkg274.

78.   Kong, Y.; Ma, J. H.; Warren, K.; Tsang, R. S. W.; Low, D. E.; Jamieson, F. B.; Alexander, D. C.; Hao, W. Homologous recombination drives both sequence diversity and gene content variation in Neisseria meningitidis. *Genome Biol. Evol.* **2013**, *5*, 1611–27, doi:10.1093/gbe/evt116.

79.   Furuta, Y.; Namba-Fukuyo, H.; Shibata, T. F.; Nishiyama, T.; Shigenobu, S.; Suzuki, Y.; Sugano, S.; Hasebe, M.; Kobayashi, I. Methylome Diversification through Changes in DNA Methyltransferase Sequence Specificity. *PLoS Genet.* **2014**, *10*, e1004272, doi:10.1371/journal.pgen.1004272.

80.   Furuta, Y.; Kobayashi, I. Mobility of DNA sequence recognition domains in DNA methyltransferases suggests epigenetics-driven adaptive evolution. *Mob. Genet. Elements* **2012**, *2*, 292–296, doi:10.4161/mge.23371.

81.   Furuta, Y.; Kawai, M.; Uchiyama, I.; Kobayashi, I. Domain movement within a gene: a novel evolutionary mechanism for protein diversification. *PLoS One* **2011**, *6*, e18819, doi:10.1371/journal.pone.0018819.

82.   Gophna, U.; Brodt, A. CRISPR/Cas systems in archaea. *Mob. Genet. Elements* **2012**, *2*, 63–64, doi:10.4161/mge.19907.

83.   Fullmer, M. S.; Soucy, S. M.; Swithers, K. S.; Makkay, A. M.; Wheeler, R.; Ventosa, A.; Gogarten, J. P.; Papke, R. T. Population and genomic analysis of the genus Halorubrum. *Front. Microbiol.* **2014**, *5*, 140, doi:10.3389/fmicb.2014.00140.

84.   Shalev, Y.; Soucy, S.; Papke, R.; Gogarten, J.; Eichler, J.; Gophna, U. Comparative Analysis of Surface Layer Glycoproteins and Genes Involved in Protein Glycosylation in the Genus Haloferax. *Genes (Basel).* **2018**, *9*, 172, doi:10.3390/genes9030172.

85.   Kandiba, L.; Aitio, O.; Helin, J.; Guan, Z.; Permi, P.; Bamford, D. H.; Eichler, J.; Roine, E. Diversity in prokaryotic glycosylation: an archaeal-derived N-linked glycan contains legionaminic acid. *Mol. Microbiol.* **2012**, *84*, 578–593, doi:10.1111/j.1365-2958.2012.08045.x.

86.   Rodriguez-Valera, F.; Martin-Cuadrado, A.-B.; Rodriguez-Brito, B.; Pašić, L.; Thingstad, T. F.; Rohwer, F.; Mira, A. Explaining microbial population genomics through phage predation. *Nat. Rev. Microbiol.* **2009**, *7*, 828–836, doi:10.1038/nrmicro2235.

87.   Tang, S.-L.; Nuttall, S.; Ngui, K.; Fisher, C.; Lopez, P.; Dyall-Smith, M. HF2: a double-stranded DNA tailed haloarchaeal virus with a mosaic genome. *Mol. Microbiol.* **2002**, *44*, 283–96.

88.   Olendzenski, L.; Gogarten, J. P. Evolution of genes and organisms: the tree/web of life in light of horizontal gene transfer. *Ann N Y Acad Sci* **2009**, *1178*, 137–145, doi:NYAS4998 [pii]10.1111/j.1749-6632.2009.04998.x.

89.   Naor, A.; Altman-Price, N.; Soucy, S. M.; Green, A. G.; Mitiagin, Y.; Turgeman-Grott, I.; Davidovich, N.; Gogarten, J. P.; Gophna, U. Impact of a homing intein on recombination frequency and organismal fitness. *Proc. Natl. Acad. Sci. U. S. A.* **2016**, *113*, E4654-61, doi:10.1073/pnas.1606416113.

90.   Oliveira, P. H.; Touchon, M.; Rocha, E. P. C. The interplay of restriction-modification systems with mobile genetic elements and their prokaryotic hosts. *Nucleic Acids Res.* **2014**, *42*, 10618–31, doi:10.1093/nar/gku734.

91.   Karlin, S.; Mrázek, J.; Campbell, A. M. Compositional biases of bacterial genomes and evolutionary

implications. *J. Bacteriol.* **1997**, *179*, 3899–913.

92. Fournier, P.; Paulus, F.; Otten, L. IS870 requires a 5′-CTAG-3′ target sequence to generate the stop codon for its large ORF1. *J. Bacteriol.* **1993**, *175*, 3151–60.

93. Otwinowski, Z.; Schevitz, R. W.; Zhang, R.-G.; Lawson, C. L.; Joachimiak, A.; Marmorstein, R. Q.; Luisi, B. F.; Sigler, P. B. Crystal structure of trp represser/operator complex at atomic resolution. *Nature* **1988**, *335*, 321–329, doi:10.1038/335321a0.

94. Burge, C.; Campbell, A. M.; Karlin, S. Over- and under-representation of short oligonucleotides in DNA sequences. *Proc. Natl. Acad. Sci. U. S. A.* **1992**, *89*, 1358–62.

95. Bath, C.; Cukalac, T.; Porter, K.; Dyall-Smith, M. L. His1 and His2 are distantly related, spindle-shaped haloviruses belonging to the novel virus group, Salterprovirus. *Virology* **2006**, *350*, 228–39, doi:10.1016/j.virol.2006.02.005.

96. Porter, K.; Tang, S.-L.; Chen, C.-P.; Chiang, P.-W.; Hong, M.-J.; Dyall-Smith, M. PH1: an archaeovirus of Haloarcula hispanica related to SH1 and HHIV-2. *Archaea* **2013**, *2013*, 456318, doi:10.1155/2013/456318.