Modeling ribosome dwell times and relationships with tRNA loading and codon usage in mammals

Cédric Gobet [1,2], Benjamin Weger[1,2], Julien Marquis [2,3], Eva Martin[2], Frédéric Gachon *[1,2,4], Felix Naef*[1]

[1] Institute of Bioengineering, School of Life Sciences, Ecole Polytechnique Fédérale de Lausanne, Lausanne CH-1015, Switzerland

[2] Nestlé Research, CH-1015 Lausanne, Switzerland

[3] Current address: Lausanne Genomic Technologies Facility, Université de Lausanne, Lausanne CH-1015, Switzerland

[4] Current address: Institute for Molecular Bioscience, The University of Queensland, St. Lucia, Brisbane, QLD 4072, Australia

⋆ Corresponding authors

Abstract

Protein translation depends on mRNA-specific initiation, elongation and termination rates. While the regulation of ribosome elongation is well studied in bacteria and yeast, less is known in higher eukaryotes. Here, we combined ribosome and tRNA profiling to investigate the relations between ribosome elongation rates, (aminoacyl-) tRNA levels, and codon usage in mammals. We modeled codon-specific ribosome dwell times and translation fluxes from ribosome profiling, considering pair-interactions between ribosome sites. In mouse liver, the model revealed site and codon specific dwell times, as well as codon pair-interactions clustering by amino acids. While translation fluxes varied significantly across diurnal time and feeding regimen, codon dwell times were highly stable and conserved in human. Profiling of tRNA levels correlated with codon usage and several tRNAs were lowly aminoacylated, which was conserved in fasted mice. Finally, codons with lowly aminoacylated tRNAs and high codon usage relative to tRNA abundance exhibited long dwell times. Together, these analyses started to reveal some of the complex dependencies between ribosome dwell times, tRNA loading, and codon usage in mammals.

Introduction

Translation regulation dynamically controls gene expression in processes such as development, the cell cycle, circadian rhythms, and response to stress [1]. At least three steps underlie protein translation: translation initiation, often thought to be rate limiting, elongation, and termination[2]. Recently, however, elongation has emerged as an important layer to fine-tune gene expression (reviewed in [3]). Indeed, variations in elongation rates may influence mRNA stability [4, 5, 6], nascent protein folding [7], and even feedback on the initiation rates [8]. For example, recent studies showed that alteration of ribosome elongation rates in cancer cells influences their proliferation and invasion capabilities [9, 10, 11].

While the links between translation elongation and gene expression are increasingly studied, the mechanisms influencing ribosome elongation rates are poorly understood, notably in higher eukaryotes. In unicellular organisms, codon-specific elongation rates are well explained by cognate tRNA concentrations [12]. This is also reflected evolutionarily, since highly expressed genes are enriched for fast codons with high concentrations of tRNAs and favorable codon-anticodon interactions [13]. However, this concept has been challenged since pioneering work in *E.coli* showed that ribosomes move at different speeds on the codons GAA and GAG [14]. These codons are decoded by the same tRNA, raising the possibility that elongation rate is not only determined by the concentration of tRNAs.

More recently, the development of ribosome profiling (RP) shed new light on the regulation of translation elongation and revealed supplementary layers of complexity [15]. Indeed, the possibility to capture the positions of translating ribosomes on mRNAs [16] fostered the development of quantitative models providing genome-wide insights on key features regulating ribosome speed [17, 18, 19]. For instance, the properties of amino acids (AA) [20], (aminoacyl-) tRNA availability [21, 22, 23], tRNA modifications [24, 25, 26], secondary structures of mRNAs [27, 28, 29], folding of the nascent chain [30], pairs of codons [31, 32], and sterical interactions with the ribosome exit tunnel [33], were shown to influence the local density of ribosomes on

transcripts. While RP studies have brought new knowledge on translation elongation, these were performed mostly in unicellular organisms and have led to divergent results as highlighted in several meta-analyses [17, 34]. One reason is that ribosome footprints are sensitive to biases from differences in protocols [35, 36], drug usage [37, 38, 39], library preparations [19], and data analysis pipelines [40]. Consequently, the reported correlations between ribosome dwell times (DTs), tRNA abundances, and codon usage frequency and bias [41], show inconsistencies. In addition, while codon usage can be precisely estimated, it is still difficult to measure tRNA concentrations. Indeed, tRNAs exhibit a high degree of modifications and complex secondary structures, which alter cDNA synthesis and biases quantification by high-throughput sequencing [42]. Thus, improved methods have been proposed to circumvent these barriers [43, 44, 45, 9] and to quantify as well tRNA aminoacylation levels [46].

Here, to better establish the determinants of ribosome elongation in higher eukaryotes, we combined modeling of RP data, codon usage analysis, and (aminoacyl-) tRNA profiling in mouse liver. In particular, we built a genome-wide statistical model that allowed us to identify key features predicting ribosome densities along transcripts, notably site-specific codon DTs and importantly ribosome site interactions. In yeast, our analysis was able to confirm previously measured slow inter-site codon pairs. In mouse liver, we found a wider dynamic range of codon- and AA-specific ribosome DTs. Our model was also able to identify codon pairs contributing significantly to the DTs, notably involving the ribosome P and A sites. The identified patterns of codon and codon-pair specific DTs in the liver were remarkably stable along the feeding/fasting cycle, and even under conditions of prolonged fasting. Moreover, meta-analysis in mammals revealed conserved DT patterns between mouse tissues and in human, which were however significantly distinct from those in yeast. Finally, we extended a recent tRNA profiling method [9] to quantify (aminoacyl-) tRNA levels in liver of *Ad Libitum* (AL) and fasted (FA) mice. tRNA levels correlated with codon usage and several tRNAs were lowly aminoacylated, which was conserved in FA mice. These data, together with codon usage properties, allowed us to explain

some codon-specificity in the estimated ribosome DTs.

Results

Modeling ribosome fluxes and codon-specific DTs including ribosome inter-site interactions

RP read counts along transcripts typically show large variations with high and low densities of ribosomes. To obtain accurate ribosome DTs in higher eukaryotes, we explicitly modeled RP read counts on a genome-wide scale. Specifically, we assumed steady-state, no ribosome drop-off, and low density of ribosomes per transcript (no traffic jams), such that the probability of finding a ribosome at a specific position on an mRNA is proportional to the translation initiation rate times ribosome DTs. As introduced, these DTs depend on the codons translated in the E, P, and A sites, as well as surrounding sequences (Fig.S1A). The total translation flux is then equal to the translation flux per mRNA multiplied by the number of mRNA copies. Specifically, we developed a generalized linear model (GLM), which models the expected read counts as gene specific fluxes (gene covariates) multiplied by ribosome DTs (codon covariates), and uses appropriate noise models (Methods). We modeled DTs additively (in log) using site-specific contributions of the three E, P, and A sites and their surrounding codons, as well as possible pair interactions between the E, P, and A sites, noted E:P, P:A, or E:A. (Fig.S1A-B). The GLM used the 61 sense codon alphabet, and we considered positions around the ribosome spanning 120 nucleotides around the E site (Fig.S1B). The same model is also applied on RNA-seq experiments (when available) to normalize the fluxes per mRNA and attenuate possible technical biases. While previous models allow to faithfully reproduce RP signal [17, 34, 28, 18, 19], our approach is different as it allows to fit globally, for each condition, all model parameters (single/pairs DTs and gene specific fluxes) directly from the raw reads counts at every position on the coding sequences (CDSs) (Fig.S1C-E) (Methods). Thus, our model allows to extract

unknown features regulating translation elongation, in addition to faithfully capturing raw RP signals (see below).

In yeast ribosome DTs anti-correlate with codon usage and display inter-site interactions

To validate our model, we analyzed two published RP datasets in *Saccharomyces cerevisiae* [23, 47]. One set was under normal (WT) [47] conditions and one treated with 3-amino-1,2,4-triazol (3-AT), which inhibits the histidine (His) biosynthesis pathway [23], thereby reducing loaded His-tRNAs. Both datasets used cycloheximide (CHX) only in the lysis buffer. Our model faithfully reproduced raw RP read counts along the transcripts as 1000 and 3000 genes had a correlation coefficient larger than 0.5 between the measured and predicted read counts, respectively, in WT and 3-AT conditions (Fig. S2A-B). Such correlation coefficients are within the range of reported values using previous methods [17, 34, 28, 18, 19].

Focusing on the model parameters describing ribosome elongation rates, we found that codon-specific DTs in WT exhibited about a four-fold range at each of the three E, P, and A sites (Fig. 1A-B, left). While codons for proline (Pro) and arginine (Arg) showed long DTs in all three sites, and were slowest in the P site, codons for isoleucine (Ile), leucine (Leu), and valine (Val) were fastest in the A site (Fig. 1A-B, left).

As expected, we found that the shortage of His in the 3-AT condition resulted in lengthened DTs in the P and A sites for both His codons (CAC and CAT) (Fig. 1A-B, right). Interestingly, outside the E, P, and A sites, DTs showed a dependency on His codons at around 30 nucleotides (positions 11 and 12) downstream of the P-site (Fig. S2C), likely reflecting queued ribosomes (disomes) behind His codons[23]. Also, some Arg codons showed slightly longer DTs in the upstream sequence, highlighting possible interactions of this positively charged AA with the ribosome exit tunnel (Fig. S2C).

Moreover, the DTs also displayed signatures of technical biases. Notably, the high variation in

DTs at position -4, coinciding with the most 5' nucleotide of the insert, was previously shown to reflect biases in library preparation (Fig. S2C) (reviewed in [48]). To further validate the biological relevance of our DTs, we compared ribosome DTs in WT condition with codon usage weighted by mRNA translation levels (wCU), to take into account condition-specific demands in codons. Interestingly, we found high negative correlations ($R^2 = 0.565$ and $R^2 = 0.495$) between the wCU and the DTs at the P and A sites (Fig. 1C). This observation suggests an evolutionary pressure to enrich for fast codons in highly expressed genes, and conversely.

In addition to the site-specific DTs, we probed whether pairs of codons in the ribosome sites synergize by adding interaction terms (E:P, P:A, and E:A) to the modelled DTs (Fig. S2D). We compared these predicted DTs with a GFP-reporter experiment in yeast probing for pairs of codons that inhibit translation [32]. Indeed, the experimentally determined inhibitory pairs (EDIPs) exhibited long predicted DTs at the E-P and P-A sites (Fig. 1D). While for these pairs the summed DTs from the individual sites were already long, the interaction terms clearly improved the match with experiments (Fig. 1E-F). Interestingly, though the E:P and P:A interactions were not correlated overall, EDIPs stood out as showing strong DT contributions in both E:P and P:A (Fig. 1G). Globally the P:A interaction matrix was sparse and not highly structured, but revealed large values and spread for the pairs involving codons for Arg or Pro (Fig S2D). Moreover, the total DTs for the 61^2 codon pairs, which were computed by including the single sites and interactions, summed for the E-P and P-A sites, revealed Arg in most of the top 50 slowest pairs, making this AA potent at decreasing translation elongation rate. Conversely, Val was contained in most of the fastest pairs. Thus, modeling RP data can identify subtle properties of ribosome DTs, such as codon-specific and inter-site contributions, signatures of sequences outside of the E, P, and A sites, and library biases.

Figure 1: In yeast, ribosome DT anticorrelate with codon usage and display site interactions
Panels for the single-site DTs are retrieved from the fit with the P:A interaction. A. DTs (log2, mean centered per site) for the 61 sense codons in the two conditions (WT/left, 3-AT/right) for the E, P, and A sites. Codons are colored according to AA. DTs with $p >= 0.05$ are not shown. Relatively fast and slow interactions are shown respectively in dark red and dark blue. B. Heatmap representation of panel A). Here, DTs with $p >= 0.05$ are set to zero. C. wCU correlates with the codon DTs for the E, P, and A sites. Black line: linear fit. D. Sum of DTs ($\log_2(2^{(E+P+E:P)} + 2^{(P+A+P:A)})$, added in linear scale) for all pairs of codons vs. the GFP signal in [32]. Red: pairs described as inhibitory. E. DTs (log2) for codon pairs. Total DTs (P+A+P:A, *i.e.* including the interactions P:A) vs. the individual contributions alone (P+A). Red: Pairs described as inhibitory. F. DTs (log2) for codon pairs. Total DTs (E+P+E:P, *i.e.* including the interactions E:P) vs. the individual contributions alone (E+P). Red: Pairs described as inhibitory. G. P:A (log2) interactions versus E:P (log2) interactions for all codon pairs.

Site-specific and inter-site codon DTs (for A and P sites) in mouse liver cluster by amino acids

In general, determinants of translation elongation are less studied in mammals. We and others have previously shown that feeding/fasting cycles can regulate translation initiation in mouse liver [49], via well described mechanisms, notably through mTOR and GCN2 related nutrient sensor pathways (reviewed in [50]). Here, we aimed to extend this analysis to the level of DTs, in particular to asses whether likely perturbed AA pools during low nutrient availability can alter codon specific DTs. Therefore, we applied the above model to our previously 84 RP samples harvested in WT and circadian clock deficient mice (Bmal1 KO) every 2-4 h around the 24 h day, including four biological replicates. As for yeast, our model faithfully reproduced raw RP read counts along transcripts, specifically, more than 1000 genes showed a correlation coefficient larger than 0.5 between the measured and predicted reads counts (Fig. S3A-B). Remarkably, DTs were very stable accross all samples, showing high biological reproducibility and no time or circadian clock-dependent changes (Fig. S3C-E). Therefore, for the following analyses, we averaged the DTs over all the 84 samples.

Global patterns of DTs for the E, P, and A sites showed biologically significant codon and AA-specificity (Fig.S3C). DTs were strikingly different than in yeast and exhibited a larger dynamic range (Fig. 2A-B). In mouse liver, the P and A sites revealed nearly 10-fold range in DTs between the fastest and slowest codons, while DTs in the E site were more tightly distributed (Fig. 2A-B). While DTs in the P and A site were overall more strongly correlated to each other than with the E site, DTs also showed clear site-specificity (Fig. 2A-B). For instance, the four codons for Gly had long DTs in the P site, however, the Gly GGT codon was among the fastest in the E and A sites, while the GGA codon was markedly slow in the A site. Also, all three Ile codons had long DTs in the A site but a very short DT in the P site (Fig. 2A-B). For the negatively charged glutamate (Glu) and aspartate (Asp), all their codons showed long DTs in the P and A sites (Fig. 2A-B). Considering a larger window around the ribosome revealed that

P and A sites, followed by the E site, contributed the largest effects (Fig. S4A). Upstream and downstream sequences outside the (-4,+6) interval did not contribute (Fig. S4A), while codons in the vicinity of the ribosome (-3, -2, -1 and +3, +4, +5) exhibited significant and correlated variations in DTs on both sides. The detected signals at the -4 and +6 positions reflect known ligation biases during the library preparation [48] .

Inter-site codon pair DTs revealed a significant influence on translation elongation in mouse liver, with P:A interactions showing the widest dynamic range, followed by E:P and E:A (Fig. S4C). Note that the interaction matrices are not symmetric, showing codons or AA specificity at the respective ribosome sites (Fig. 2C and S4B). Intriguingly, the P:A matrix highlighted a striking clustering by AA for the A site (Fig. 2C), while E:P interactions clustered by AA in the P site (Fig. S4B). This suggests that P:A inter-site DTs are determined by AAs through their influence on the peptide bond formation. The clustering by AA was corroborated by a model selection analysis on the 84 samples, where the alphabet for the DT regression coefficients was taken as either the 20 natural AA, or the 61 sense codons (Fig. S4D). While the preferred alphabet was overall that of the codons, the model with AA coefficients at the A site for the P:A interaction was preferred to all the other models (Fig. S4D). In the case of the E:P interaction, the AA alphabet in the P site was considered as the best model. Overall, models including the site interactions were preferred to the reduced models, emphasizing the importance of inter-site interactions in determining ribosome DTs in mouse liver. In fact, the codon-pair interactions contributed to the total DTs in the P and A site by a factor larger than 2 for about one hundred pairs (Fig. 2D). The P:A matrix revealed strong positive interactions (ones lengthening the DTs) for pairs of bulky (Pro, tryptophan (Trp) and phenylalanine (Phe)) or achiral (Gly) (Fig. 2C and D) AAs. As in yeast, Arg codons were part of the slowest pairs (Fig. 2C and D). Surprisingly, the known stalling pair Pro-Pro showed the largest negative interaction (Fig. 2C and D), possibly related to eIF5A activity [51]. The E:P matrix was more complex: pairs involving the AA Gly, Asp, Asparagine (Asn), and Pro in the P site lengthened the total E-P

pair DTs (Fig. S4B).

Unlike in yeast (Fig. 1C), ribosome DTs did not correlate with codon usage in mouse liver (Fig. 2E), arguing for different evolutionary pressure on translation efficiency.

Ribosome DTs in liver are stable under prolonged fasting

The above analysis showed highly robust DTs between liver samples collected during the normal feeding (night) / fasting (day) cycle (Figure S3C-E). To probe whether ribosome DTs are sensitive to longer periods of fasting, we performed new RP experiments in mice fed either AL or fasted for up to 30 hours (FA) (Fig. 3A). Since the enrichment in ribosome footprints can be sensitive to RP protocols [52], we here used a small RNA-Seq protocol with random adapters (UMI) to reduce possible ligation biases and PCR duplicates. Moreover, as ribosome dynamics and DTs are affected in RP experiments with CHX in yeast [37, 53], we tested conditions without CHX in the lysis buffer.

First, we validated the effect of prolonged fasting by analyzing differential RP signals between AL and FA. Genes related to the Peroxisome Proliferator-Activated Receptor α (PPARα) pathway and to fatty acids oxidation were upregulated in FA, presumably to provide the energy needs (Fig. 3D). On the contrary, genes related to lipid biosynthesis were downregulated in FA (Fig. 3D), suggesting that animals switched from glucose to fatty acid metabolism in FA, as already described [54]. Moreover, *Mat1a*, *Asl* and *Got1* related to AA biosynthesis were upregulated in FA (Fig. 3D), presumably in response to perturbed AA homeostasis. The RP data in the new conditions showed a typical tri-repeat nucleotide pattern (Fig. 3B), confirming the presence of *bona fide* translating ribosomes in the FA samples, as well as in samples without CHX (NOCHX). In addition, the modified library preparation improved the goodness of the fits. Indeed, correlation coefficients were higher than described above (Figs. S3A-B). More than 3000 and 1000 genes had a correlation coefficient larger than 0.5 in AL and FA conditions, respectively (Fig S5A-B).

Figure 2: **Site-specific and inter-site codon DTs (for A and P sites) in mouse liver cluster by AA**
Site-specific DTs are retrieved from the fit with the P:A interaction. A. DTs (log2, mean centered per site) for
the E, P, and A sites averaged over the 84 samples in mouse liver. Codons are colored according to AA. DTs
with $p >= 0.05$ are set to zero. Fast and slow DTs (relative to the mean) are shown in dark red and dark blue,
respectively. B. Heatmap representation of panel A). DTs with $p >= 0.05$ are set to zero. C. Interaction matrix
for the pairs P:A (log2). Codons are colored according to AA. Codons in both sites are hierarchically clustered
(euclidean distance matrix, complete linkage algorithm). Fast and slow interactions are shown in dark red and
dark blue, respectively (colorbar). D. DTs (log2) for codon pairs. Total DTs (P+A+P:A, *i.e.* including the P:A
interactions) vs. the individual contributions alone (P+A). Pairs with interactions > 1 or < -1 are annotated
and colored, respectively, in blue and red. E. wCU does not correlate with codon DTs for the E, P, and A sites.
Black line: linear fit.

Surprisingly, codon DTs in the AL and FA mice showed no difference in both the CHX or NOCHX conditions (Figs. 3C and S5C), including the inter-site codon DTs (Fig. S5D). Moreover, these DTs were highly correlated with those described above for the 84 samples (Fig. S6C-D). Nevertheless, the FA samples showed a reduced dynamic range in the DTs, presumably due to variability in RP signal quality (Fig. S5A) related to the global decrease in translation levels.

Finally, we probed whether the perturbed metabolic state in FA might lead to differential codon usage [22]. Strikingly, when considering the wCU bias in WT and FA animals, we found that most of the codons with a G or C nucleotide at the third position (GC3) were enriched in up-regulated transcripts in FA, while codons with an A or T nucleotide were underrepresented (Fig. S5E).

Meta-analysis reveals conserved ribosome DTs in mammals

To further compare the estimated ribosome DTs, we analyzed published RP datasets in mouse liver (H: Howard *et al.*) [55], mouse kidney (CS: Castelo-Szekely *et al.*) [56], and in the human liver cell line Huh7 (L: Lintner *et al.*) [57]. Despite differences in library preparation protocols (Methods), codon DTs at the A site were highly correlated between the mammalian datasets ($0.48 < r < 0.96$), including in different tissues (kidney) and human cells (Fig. S6C, E). However, the mammalian DTs were markedly different from those in yeast [17] . Note that the relative contributions to the DTs from the E,P, and A sites vs. surrounding sites, in particular at positions -4 and +6, differed significantly depending on the protocols (Fig. S6A-B). Similar potential biases have been reported previously [52]. Here, we found that protocols without cDNA circularization showed highest signals in the P and A sites, presumably reflecting ribosome dynamics more faithfully (Fig. S6A-B). Moreover, this was also reflected in the inter-site codon pair DTs (P:A), which were more consistent across experiments (Fig. S6D) with that protocol.

Together, this meta-analysis highlighted how different library preparations lead to damped

RP signals in the A and P sites for some protocols, and showed that the codon DT patterns are conserved between mouse tissues and mammalian species.

(Aminoacyl-) tRNA profiles are conserved in fed and fasted mice

We next asked whether the estimated DTs can be linked with tRNA abundances or loading levels, which is poorly studied in higher eukaryotes [41]. The chemical modifications and secondary structure of tRNAs render them difficult to quantify. A recent hybridization method combined with sequencing, which controls specificity using left/right probes and a stringent ligation step, allows to bypass the cDNA synthesis to quantify tRNA levels [9]. To measure tRNA abundances and assess possible links with ribosome DTs in mouse liver, we adapted and optimized this method to target all annotated mouse tRNAs (Fig. S7A). Moreover, we quantified the fraction of (aminoacyl-) tRNAs using sodium periodate [58], which depletes unloaded tRNAs by selective biotinylation of 3'-ends (Fig. S7A). This way, we aimed to quantify the tRNA pools available for elongation in the ribosome A site.

tRNA molecules are encoded by a large number of genes. Therefore, we designed 303 DNA probe pairs (left and right) to target all mouse tRNA sequences from the *GtRNAdb* database [59]. Our modified protocol yielded a high proportion of specific ligations between left and right probes, showing target specificity for tRNAs (Fig. S7B). Indeed, mapping of the sequencing reads to all possible combinations (303^2) of left and right probes showed that more than 75% of ligated products belonged to tRNA genes of the same codon (Fig. S7B), and even 95% were from probe pairs that could be assigned to specific codons with high confidence (Fig. S7B) (Methods). We performed further experiments to validate the specificity and evaluated the efficiency of DNA ligases (Fig. S7C-D).

We measured tRNA abundance on mouse livers from the same samples as those used for the RP. Specifically, we quantified the total tRNA (control, NaCl) and the (aminoacyl-) tRNA (periodate, $NaIO_4$) abundances from the same pieces of liver in two replicates in the AL and

Figure 3: **Ribosome DTs are not affected in fasted mice and without CHX**
Individual DTs are retrieved from the fit with the P:A interaction. A. Livers from mice fed *AL* or FA were harvested at ZT4, ZT12, and ZT18. RP was performed without CHX in the lysis buffer. B. Normalized fractions of reads of length 32 around the start and stop codons in a window of 100 nucleotides genome-wide. Dark blue: AL/NOCHX; dark red: FA/NOCHX. C. DTs (log2, mean centered per site, heatmap) for the E, P, and A sites in AL/NOCHX and FA/NOCHX. Codons are ordered by AA. Side bars: log2 color scale. D. Differential expression of RP signals between AL and FA (Methods). Benjamini-Hochberg adjusted p-values (-log10) plotted against averaged log2 fold change between FA and AL. Genes with FDR < 1% and absolute log2 fold change > 1 are annotated and colored. Blue: genes in the KEGG "PPARα signal pathway"; Green: KEGG and GO term "lipid biosynthesis"; Orange: KEGG "AA biosynthesis".

FA conditions, at three different times during the day (ZT04, ZT12, ZT18) (Fig. 4A, Methods). tRNA abundances were highly reproducible (Fig. S7E), exhibited a large dynamic range (Fig. 4B), and were significantly correlated with PolIII ChIP-Seq data in mouse liver [60] (Fig. S7F). tRNA levels for AA encoded by four synonymous codons ("4-codon" box) were represented by one dominant highly expressed isoacceptor with a T at the wobble position 34 (*e.g.* TGC/Ala, TGG/Pro, TCC/Gly, TAC/Val, TGT/Thr) (Fig. S8A). The tRNA levels showed only small variations over the biological conditions, except for mitochondrial tRNAs (Fig. S8B). Strikingly though, principal component analysis (PCA) on the four conditions (*i.e.* NaCl/AL, NaCl/FA, NaIO$_4$/AL and NaIO$_4$/FA) exhibited a clear separation between the control and periodate conditions (Fig. 4C), indicating differential loading of the tRNAs. Surprisingly, the AL and FA samples were indistinguishable in the total tRNA and (aminoacyl-) tRNA conditions (Fig. 4C-D), indicating no imbalance of tRNA charging in FA. However, some codons for Asn, Asp, and Ile were lowly aminoacylated, independently of the feeding regime (Figs. 4D, S8C). Intriguingly, these same codons were found to exhibit the slowest DTs in the A site (Fig. 2).

Relationship between (aminoacyl-) tRNA levels, codon usage, and DTs

To substantiate this observation, we investigated whether variations of codon DTs and wCU could be explained by the available tRNA pools. In lower organisms, it is known that codon usage frequency and tRNA pools co-adapted to fine-tune translation elongation. However, whether this relation holds in mammals is debated, mainly due to the lack of good proxies for tRNA levels. Moreover, codon usage differences seem to be driven by mutational bias such as GC content [61]. Here, we discovered a significant correlation between wCU and our measured tRNA levels in mouse liver (Fig. 5A), extending previous work using POLIII loadings on tRNA genes as proxies [61]. Our analysis also highlighted codons with high or low demand (codon usage) compared to the supply (tRNA levels), as quantified by the codon balance [62] (Fig. 5A).

Figure 4: **(Aminoacyl-) tRNA profiling in AL fed and FA mice**
A. Mice fed AL or FA were sacrificed at ZT4 (fasting duration: 16 h), ZT12 (24 h) and ZT18 (30 h) and livers were harvested. Each sample was treated with NaCl and sodium periodate (NaIO$_4$). B. Cumulative fraction of reads for each tRNA, ordered by abundances. Anticodons and AAs are indicated for the 50 first codons. Blue: four-codon box AA; red: mitochondrial tRNAs. C. PCA of the tRNA abundances (log2 UMI). PC1 and PC2 explain 42.2% and 21.3% of the variance, respectively, and separate NaCl from NaIO$_4$ treatment. NaCl (circle), NaIO$_4$ (square), AL (blue), FA (red), replicate 1 (black), replicate 2 (grey). ZT is shown beside the points. D. Ratio of tRNA abundances (log2 fold change, averaged over the time points) between the NaCl and NaIO$_4$ for AL fed vs. FA mice (significant changes, $p < 0.05$ in red). No tRNA showed a significant difference between AL and FA (*i.e.* fell out of the diagonal).

DTs in the A site did not exhibit a simple correlation with tRNA abundances (Fig. 5B), nor with the codon balance (Fig. 5C). However, some codons clearly stood out. In particular, the slow DTs for Glu codons (Fig. 5C) may well result from their low codon balance, hence limiting tRNA availability at the A site. Similarly, while DTs in the A site were overall poorly correlated with tRNA aminoacylation levels, codons for Asp, Asn, and Ile, which had particularly lowly charged tRNAs, coincided with some of the slowest DTs (Fig. 5D). We therefore included several effects in a linear model, which uncovered that a linear combination of tRNA aminoacylation levels and codon balance captures a significant portion of variation in the A site DTs, particularly the long DTs for Glu, Asp, Asn, and Ile codons (Fig. 5E).

Discussion

We extensively modeled RP datasets and uncovered codon-specific and inter-site DTs determining ribosome elongation rates in mammals. These DTs were highly stable across all conditions tested. In parallel, we quantified (aminoacyl-) tRNA levels in mouse liver and identified several features regulating ribosome elongation, such as aminoacylation levels and the balance between tRNA levels and codon usage.

In yeast, our model accurately inferred codon-specific DTs and highlighted mainly Arg and Pro as slow in the A and P sites. These AAs are known for their inefficient peptide formation and sterical interactions with the ribosome [63, 64]. A significant negative correlation was observed between DTs and wCU, reflecting natural selection for fast codons in highly translated genes. While this relationship has been described [65, 34], the found correlation is, to our knowledge, the highest reported. Moreover, our analysis confirmed recently identified EDIPs [32], and deciphered their synergistic effect in addition to the site-specific contributions. We showed that the EDIPs had slow DTs both on the E:P and P:A positions, highlighting potentially inefficient

Figure 5: **Relationship between (aminoacyl-) tRNA levels, codon usage and DT**
A. Significant correlation ($R^2 = 0.22, p = 1.2e - 4$) between wCU frequency (log2) and normalized total tRNA (NaCl) read count for each codon (log2, UMI, averaged over the AL samples). The gray line shows the first principal component (PC). Orthogonal distance to the PC reflects the balance between tRNA supply and demand. Codons with positive (resp. negative) balance are colored blue (resp. red). Codons are assigned to their canonical tRNAs (dark-grey) or to wobble tRNAs (light-grey) where appropriate (Methods). B. DTs in the A site (log2) *vs.* tRNA levels (NaCl, log2) averaged over the AL samples ($R^2 = 0.01$, p-value= $3.9e - 1$). C. Correlation between tRNA levels/codon usage balance and DT in the A site (log2) averaged over the AL samples. R^2 and p-value are reported for the linear regression ($R^2 = 0.02$, p-value= $2.9e - 1$). D. Correlation between tRNA uncharged levels (NaCl/$NaIO_4$, log2) and DTs in the A site (log2) averaged over the AL samples. R^2 and p-value are reported for the linear regression ($R^2 = 0.03$, p-value= $1.7e - 1$). Positively correlated codons are annotated by their one-letter AA. E. Significant correlation($R^2 = 0.17$, p-value=$2.15e - 02$) between estimated and predicted DTs in the A site. Prediction uses a linear model with the balance and uncharged levels, in linear scale, as explanatory variables. In C-E, annotations refer to one-letter AA.

translocation of the pair due to wobble base-pairing or other mechanisms [32].

In mouse liver, DTs differed significantly from yeast, showing a larger spread and higher complexity. Remarkably, DTs were very similar between different tissues, species, and RP protocols. Moreover, the DTs were consistent with a peptide motif enriched in stalled ribosome sites in mouse embryonic stem cells (mESCs) [66].

We found that the smallest and achiral AA Gly exhibited very long DTs (in the A and P site) with different magnitudes between the isoacceptors and tissues (*i.e.* liver and kidney). Interestingly, in bacteria, Gly codons are slow, although this effect is still difficult to separate from Shine-Dalgarno (SD) dependent stalling [67] or protocol artifacts [68], and is therefore debated [69]. As mammals do not use a SD mechanism, our result support an alternative hypothesis, such as slow codon-anticodon pairing [70] or inefficient peptide bond formation. Pioneering work in *E. coli* suggested that Gly-tRNAs adopt a particular conformation due to the U nucleotide in position 32 and that unmodified U_{34} on $tRNA_{UCC}^{Gly}$ could decode the four Gly codons (a pairing known as superwobbling [71]), but with low efficiency (reviewed in [70]). While this mechanism was shown in unicellular organisms, our tRNA profiling found $tRNA_{UCC}^{Gly}$ as the major Gly isoacceptors and one of the most abundant tRNAs in mouse liver. Further work may indicate whether superwobbling occurs in the liver.

The DTs for the acidic AAs (Asp and Glu) were among the slowest. Glu showed a particularly low balance of tRNA levels/codon usage and Asp tRNA was lowly charged. This could lead to a shortage of tRNA availability and ribosome stalling [21]. As their codons share the same first two bases, competition with near-cognate tRNAs [25], or pairing inefficiency due to the wobble mechanism could also explain the long DTs. Indeed, slower elongation would allow higher precision in codon-anticodon discrimination [72].

Ile codon DTs were slow in the A site while fast in the P site. Remarkably, the isomeric Leu codons were the fastest in the P and A sites, highlighting a structure-independent mechanism. Indeed, we showed that Ile-tRNAs were lowly aminoacylated, reducing Ile availability on the A site, but other explanations are possible. For instance, since Ile is decoded by three differ-ent codons, a suitable pairing mechanism such as inosine or other U34 modifications could be used to avoid pairing of the fourth near-cognate codon (Met) and therefore increase the DTs [73].

One of our main results concerned the contributions of codon-pair interactions between ribo-some sites towards DTs, mainly at the P and A site. At these positions, the ribosome catalyzes the peptide bond formation between (aminoacyl-) tRNA in the A site and peptide-tRNA bound to the P site. Our analysis revealed that the identity of the AA in the A site (acceptor), and not the codon, was the best descriptor of those codon pair interactions. Pairs including bulky AAs or Gly in the A site were slow, highlighting their potential inefficiency in peptide bond formation. Interestingly the DT for Pro-Pro pairs, known to inefficiently form peptide bonds [74], was markedly reduced by the interaction. This observation probably shows the role of eIF5A in resolving this stalling motif. On the other hand, Gly, Asp, and Glu, which were slow in our analysis, were shown by others to require eIF5A for their efficient translation [75, 76].

Other features not included in the model, and which are independent of the codon identity, might regulate ribosome elongation. A high number of liver proteins are secreted and thereby translated by ribosomes bound to the endoplasmic reticulum (ER) via the interaction of signal recognition particles with the nascent peptide chain. These interactions are known to stall the ribosomes, however, as these appear to be codon independent, we did not detect them in our analysis [28]. In addition, chaperone proteins interacting with the nascent peptide can influence co-translational folding and subsequent ribosome density on mRNAs [77]. The model could also

be further extended by including RNA secondary structure and modifications, pseudo-knots acting as ribosome roadblocks, and slippery sequences inducing frame shifting [28, 18, 78, 79].

While we found a striking correlation between DTs and wCU in yeast, the same did not hold in mammals. This suggests that biased codon usage in mammals reflects more complex evolutionary forces, such as mutation driven GC bias [61]. Nevertheless, the measured tRNA abundances showed signatures of adaptation, since tRNA levels correlated with the wCU. These correlations extended previous results at the transcription level or in highly expressed genes [61, 80]. Related to this, one still open challenge is to assign tRNAs to their corresponding codons, due to the extended wobble base pairing rules related to tRNA modifications.

Surprisingly, tRNA loading was unaffected by prolonged fasting. Several studies in cell lines showed that decreasing AAs in culture media leads to decreased (aminoacyl-) tRNA availability and therefore increases ribosome stalling [58, 21]. Moreover, others have shown that codon optimality contributes to differential mRNA translation in response to starvation [22]. While we did not observe this, probably due to the *in vivo* state, GC3 bias (*i.e.* GC bias at position N3 in codons) was significantly different between genes translated in AL and FA mice or also between night and day conditions (not shown). Genes with high GC3 content have been shown to provide more targets for methylation than those with low GC3 and to be enriched in stress responsive genes [81]. Oxidative stress occurs during fasting/day in mouse and correlates with GC3 content. Nevertheless, the reason of the higher GC3 level in FA compared to AL still needs to be identified.

Like (aminoacyl-) tRNA levels, DTs were unchanged between AL and FA. We can hypothesize that after more than 30 hours of starvation, mice compensate the lack of AAs by a large global decrease of translation initiation through mTORC1/GCN2 [82], making tRNA availabil-

ity and translation elongation non limiting. Moreover, since RP signals, DTs, and tRNAs were measured in relative and not absolute amounts, we cannot exclude a total decrease of translation elongation rate, aminoacylation, or tRNA levels.

In conclusion, ribosome DTs, codon usage, tRNA levels, and translation elongation in mammals do not obey simple relationships. Nevertheless, although a global understanding is still missing, we were able to link both tRNA/codon usage balance and aminoacylation levels with anomalously slow DTs in the P and A site of the ribosome. Probing different ribosome states (*e.g.* free A site) using RP combined with different drugs [68] or improving the quantification of (aminoacyl-) tRNA through nucleotide modification removal [43] will lead to better understanding of the determinants of translation elongation. Finally, more work is needed to understand the consequences of changes in ribosome elongation rates for mRNA stability and nascent protein folding.

Methods

Inference of DT and translation fluxes

Preprocessing of RP data

RP from yeast, mouse, and human were respectively mapped on the sacCer3, mm10 and Hg38 genomes using STAR [83] with parameters `-seedSearchStartLmax 15`. Genomes indexes were built using Ensembl transcripts annotations. Adapters were retrieved for the different datasets and input as parameter for STAR. In the case of NEXTFlex library, fastqs files were parsed and duplicated sequences (UMI and insert) are removed. Sequences were trimmed for adapters using fastx_clipper with parameters `-Q33 -a TGGAATTCTCGGGTGCCAAGG -l 11` and UMIs are removed (4 nucleotides on both sides). Then, the fastq files are mapped using STAR with options `-seedSearchStartLmax 15`. The subsequence BAM files were sorted and indexed.

Read counting on the CDSs

For each protein coding transcript with a CDS larger than 120 nucleotides, reads with zero mismatches, unique mapping (nM:i:0 & NH:i:1) and a length between 25 and 40 nucleotides were retrieved using `samtools view` in the respective region. E site position was defined, for each read size, in function of the frame on the CDS and pileup plots at the start codon. From the E site position, the sequence in the window [-60,+60] nts was reported and incremented by one at each new observation. Sequences with a window spanning the start or stop codon were removed.

Filtering

A sliding window of 120 nucleotides moving 3 by 3 on the CDS of protein coding genes were computed and the respective sequences were reported (Figure S1D). This set of sequences is used as a reference and their respective number of counts is set to zero. Every time a read occurs at one of these sequences, we incremented the count by one (Fig.S1D) . Genes with less than 5% of positions covered or less than 5 positions observed were discarded. Genes with less than 100 counts were removed. Sequences containing a stop codon (TAG, TGA, or TAA) or non-unique in the coding genome were discarded. Depending on the sample coverage, we monitored about 5000 genes in mammals.

Generalized linear model

We used a generalized linear model for the observed RP read counts at the different positions on the gene CDS. The read counts at a specific codon position i corresponding to the ribosome E site on the CDS of a gene g in sample s were modeled as a negative binomial (NB) with mean μ_{igs} and dispersion parameter θ_s. θ_s was taken as a sample specific parameter and was empirically estimated from the variance-mean relationship for a NB (Eq.1). Specifically, for each gene we used adjacent codon pairs occurring more than once, the rationale being that the counts

observed on the multiple instances behave as replicates (i.e. the counts are sampled from the

same NB). For those pairs of codons, we computed the respective mean and variance of counts

and since we assumed that θ_s only depends on the sample, we then estimated θ_s globally (from

all pairs on all genes) by linear regression using Eq.1 (Fig.S1 E) . The reason to consider counts

on pairs of codons was to take into account a minimal contextual dependence of the counts,

while keeping the counts not too sparse.

$$Y_{ig} \sim NB\left(\mu_{ig}, \theta\right)$$

$$\mathrm{E}[Y_{ig}] = \mu_{ig} \tag{1}$$

$$\mathrm{Var}\left(Y\right) = \mu + \frac{\mu^2}{\theta}$$

460 where we omitted the sample index for clarity. The model for each sample is as follows (omitting

461 the sample index) :

$$h(\mu_{i,j}) = \underbrace{f_g}_{\text{gene}} + \left(\sum_{k=-20}^{20} \underbrace{\tau_{k,c(i+k)}^{(1)}}_{\text{single}}\right) + \underbrace{\tau_{c(i+s1),c(i+s2)}^{(2)}}_{\text{pairs}} + \text{offset(library size and/or RNA-Seq)}$$

$$\text{with}(s_1, s_2) \in \begin{cases} (0,1)\text{for the E:P fit} \\ (0,2)\text{for the E:A fit} \\ (1,2)\text{for the P:A fit} \end{cases} \tag{2}$$

where the offset term(s) makes the gene fluxes normalized by library size. In addition (when

specified in the text), to have gene translation fluxes normalized per mRNA abundance, we fitted

the same model to RNA-seq data (interpreting the DT as sequencing dependent efficiencies),

and predicted from it the gene expected specific RNA abundance for use as an offset term.

$c(i) \in \{AAA, AAC, AAG, ..., TTT\}$. $h(x) = \log(x)$ the natural link function for count data.

$\tau_{k,c(k)}^{(1)}$ denote the individual codon DTs for the 61 sense codons in log scale at position k, with

k the relative position to the ribosome E site. $\tau^{(2)}_{c(i+s1),c(i+s2)}$ denote the inter-site codon pair

DTs for the 61^2 pairs of sense codons in log scale at positions (s_1, s_2) relative to the ribosome

E site. These codon pairs are modeled for the sites E:A, E:P, and P:A. f_g is the gene-specific

translation flux in log scale.

The fit was performed using *glm4()* function from the R package *MatrixModels* with the noise

family *negative.binomial(θ_s)* from the *MASS* package and with sparse design matrix option.

Sequencing library size is used as an offset. RNA-Seq data is fitted (when available) and read

counts are predicted at every positions and used as an offset.

Since this problem does not have full rank, we set for the fit : $\tau^{(1)}_{k,AAA} = 0$ for $\forall k$ and $\tau^{(2)}_{(AAA,.)} = 0$,

$\tau^{(2)}_{(.,AAA)} = 0$, $\tau^{(2)}_{(AAT,AAT)} = 0$. To present the results, we then chose the more natural convention

(zero average): $\sum_c \tau^{(1)}_{k,c} = 0$ for all k, $\sum_c \tau^{(2)}_{c,c'} = 0$ for all c', and $\sum_{c'} \tau^{(2)}_{c,c'} = 0$ for all c and shifted

the gene fluxes accordingly. In all figures showing DTs, point estimates as returned by *glm4* are

plotted. The reproducibility of those point estimate accross biological samples is shown in Fig.

S3C.

Differential expression in AL vs. FA

Two outlier samples (ZT12/FA/CHX and ZT04/FA/NOCHX) were excluded for the differential

expression analysis and DT modelling . Statistics were computed using EdgeR [84] comparing

a model including factors for time, feeding, and drug conditions against a model without the

feeding term.

Animals experiments

Animal studies were approved by the local ethics committee, and all protocols were approved

by the Service Vétérinaire Cantonal (Lausanne, Switzerland) under license VD3613. 8 weeks

old male C57BL6/J mice (Charles River Laboratory) are kept under diurnal lighting conditions

(12-h light, 12-h dark) at a temperature of 21 °C +/- 2 °C. After a complete night of fasting,

the mice were kept without access to food for an additional period of up to 24 hours. During this time period animals were sacrificed every 8 hours starting at ZT4. Control animals were kept on *ad libitum* feeding regimen.

Ribosome profiling

Samples preparation for RP was performed as described in [49] except for the conditions without cycloheximide (CHX) in which fresh livers were directly lysed in ice-cold lysis buffer without CHX and directly flash-frozen in liquid nitrogen. To limit possible bias due to footprint size selection related to different conformations of the ribosome [67] [36], a larger band was cut on the TBE-gel. Libraries were generated using NEXTflex Small RNA Sequencing Kit v3 (bioo scientific) following the manufacturer's protocol. Samples were pooled based on the Illumina indices used. Denatured pools were spiked with 5% PhiX and clustered (loading equivalent to 3 samples per lane) onto a rapid single-end v2 flowcells at a concentration of 8pM. Libraries were sequenced on a HiSeq 2500 (Illumina) for 50 cycles.

(Aminoacyl-) tRNA profiling

The tRNA profiling protocol was adapted and modified from [9]. We tested the initial protocol [9] on mouse liver samples but the results showed a high proportion of unspecific ligations between the left and right probes from distinct tRNAs. We solved this issue by inverting the order of two steps in the protocol: we performed the pull-down and cleaning on magnetic beads before the splint ligation between the two DNA probes on the tRNA (Fig. S7A). Oxidation of 3'-tRNA by periodate was adapted from [58]. All the steps were performed under cold and acidic conditions to avoid deacylation of the tRNAs before Na periodate oxidation.

Probe Design

DNA probes were designed to target all the annotated mouse tRNAs from `http://gtrnadb.ucsc.edu/`. The database contains tRNA gene predictions by tRNAscan-SE [59]. tRNA se-

quences for *Mus musculus* (GRCm38/mm10) were downloaded and spliced *in silico*. The sequences were split in the middle of the anticodon in order to design left and right probes. After reverse complementation of the sequences, overhangs (for PCR primer binding) and unique molecular identifiers (UMIs, 2x6N) were added (right-probe adapter:

5'-GCACCCGAGAATTCCANNNNNNTGG-3, left-probe adapter:

5'-NNNNNNGATCGTCGGACTGTAGAACTC-3'). Left probes were ordered with a 5'-phosphate to allow ligation with the right probe upon annealing with the corresponding tRNA. The random nucleotides were ordered as «high fidelity wobble »to ensure homogeneous representation of the four bases in the UMI and to avoid bias. DNA probes were ordered at MicroSynth AG (Switzerland).

tRNA extraction and oxidation

50-100 μg of frozen mouse liver tissues were weighted under cold conditions. Beating beads were added and the samples were homogenized in 350 μl of cold Qiazol (Qiagen) lysis reagent in a TissueLyser (Qiagen) for 2 x 2 min at 20 Hz. Tubes were left 5 min at room temperature. 140 μl of $CHCl_3$ was added and homogenates were shaken vigorously followed by centrifugation at 4°C for 15 min (12'000 x g). The upper aqueous phase was carefully removed and 1 volume (350 μl) of buffered phenol (Phenol:chloroform:isoamyl alcool, 25:24:1, pH 4.9) was added. Samples were mixed and centrifuged for 15 minutes at 4°C (12'000 x g). Upper phase (300 μl) was supplemented with 1 volume (300 μl) of cold isopropanol, precipitated 30 minutes at 4°C and then centrifuged for 15 minutes at 4°C (12'000 x g). RNA pellets were dried at room temperature and re-suspended in 500 μl of Sodium Acetate buffer pH 4.9 (0.2M). Samples were split in two tubes (2 x 250μl) for sodium periodate oxidation (NaIO$_4$) or control (NaCl) treatment. 50 μl of NaCl (0.3M) or NaIO$_4$ (0.3M) was added and samples were incubated for 30 minutes at room temperature. The reaction was then supplemented with 300 μl Ethanol (70%) and loaded on a miRNeasy column (Qiagen). tRNA were extracted following the miRNA easy protocol from

Qiagen. 390 μl Ethanol (100%) was added to the flow through and loaded on a MinElute column (Qiagen). Columns were washed following the manufacturer's protocol and RNAs were eluted in 15 μl RNase-free H_2O.

Deacylation

Purified tRNAs (14 μl) supplemented with 6 μl of Tris-HCl (pH 8) were deacylated by heating at 40°C for 35 minutes. Reaction was stopped by the addition of 30 μl NaAcetate (0.3 M). RNAs were purified using RNA Clean & Concentrator -5 kit (Zymo) according to manufacturer's instructions and eluted in 15 μl RNase-free H_2O.

3'-tRNAs biotinylation

3'-tRNAs biotinylation was adapted from Pierce RNA 3'-End Biotinylation Kit (Thermo Fisher). Deacylated tRNAs were denaturated in 25% DMSO at 85°C for 5 minutes and directly chilled on ice. Biotinylation was performed in a 90 μl reaction with 6 U of T4 ssRNA Ligase (NEB), 4 μl Biotinylated Cytidine (Thermo Fisher, 1mM), 2 U RNase inhibitor, 9 μl RNase Buffer (NEB), 9 μl ATP (NEB, 10mM), 40 μl PEG 800 (50%), and 20 μl denaturated RNAs. The reaction was performed overnight at 16°C. Biotinylated tRNAs were cleaned using RNA Clean & Concentrator -5 kit (Zymo) according to manufacturer's instructions and eluted in 20 μl $H_2$0.

Probes hybridization

DNA probes were synthesized by *Microsynth AG* and resuspended at a 100 μM concentration. The 606 probes were then mixed at an equimolar ratio (0.15 μM each) and aliquoted for further usage. Hybridization of probes was performed in a 300 μl$-$reaction with 45 μl probes mastermix, 30 μl hybridization buffer 5x (500μM EGTA, 25mM NaCl, 50mM Tris$-$HCl), 205 μl RNase-free water and 20 μl tRNAs. After a 15 minutes denaturation at 95 °C, the mixture was slowly cooled down to 55 °C (0.2 °C/second) and incubated for 30 minutes.

Beads purification

200 μl of Dynabeads MyOne Streptavidin C1 (Thermo Fisher) were washed following manu-facturer's instructions for RNA usage. 250 μl of beads, re-suspended in washing buffer 2x (10 mM Tris−HCl, 1 mM EDTA, 2M NaCl), were incubated with $300ul$ of the resulting RNA-DNA hybridization reaction for 40 minutes with gentle rotation. Beads were washed/magnetized three times with 1ml of washing buffer (1x) and re-suspended in 300 μl H_2O.

RNA-DNA hybrid ligation

Bead purified DNA-RNA hybrid on beads were ligated at the anticodon nick by a combination of SplintR and T4 DNA ligases (NEB) to minimize ligation efficiency bias. 300 μl of DNA-RNA hybrids were splint-ligated with 2.5 U of SplintR DNA ligase and 30μl of SplintR DNA ligase buffer (10X, NEB) for 1 hours at 25 °C. Then, 10 U of T4 DNA ligase (NEB) and 33μl of T4 DNA ligase buffer (10x, NEB) were added. Ligation was performed overnight at 16 °C.

RNA digestion

Beads were magnetized and washed once with washing buffer (1X) to remove any remaining ligases. Next, beads were re-suspended in 10μl H_2O. 2 U of RNase A (Thermo Fisher) and 10 U of RNase H (NEB) with RNase H buffer (10X) (NEB) were added and digestion was performed for 30 minutes at 37 °C . Elution buffer (5X) was added for a final concentration of 50 mM tris pH8, 10 mM EDTA, 1% SDS and samples were incubated at 65 °C for 30 minutes with intermittent shaking to retrieve ligated DNA probes. Beads were magnetized and supernatant extracted. DNA ligated probes were purified using DNA Clean & Concentrator -5 kit (Zymo) according to manufacturer's instructions and eluted in 20 μl RNase-free H_2O.

qPCR for quality control and relative concentration estimation

The relative concentration of the resulting DNA ligated-probes was assessed by quantitative PCR (qPCR) using the LightCycler 480 SYBR Green I Master kit (Roche). 3.5 μl of $\frac{1}{10}$ diluted samples was used to assemble a 10 μl reaction and run on a Light Cycler 480 II (Roche) with the primers CAGAGTTCTACAGTCCGACGAT and TTGGCACCCGAGAATTCCA (matching each probe's ends) at a final concentration of 0.3 μM. Cycling conditions consisted of an initial denaturation step of 5 min at 95 °C followed by 40 cycles of 30 s at 95 °C and 45 sat 45 °C. The Cp obtained were used to calculate the optimal number of PCR cycles amplification required for library amplification, as described previously [49]. The number of required cycles were between 13 and 17 depending on the experiments and samples. The quality of the ligation was assessed on a Bioanalyzer small RNA chip (Agilent Technologies).

PCR amplification

The PCR was designed following Illumina's recommendation taking advantage of the indexed oligos from the TruSeq small RNA kit. A 50 μl-reaction was assembled with Kapa Polymerase and 15 ul of DNA ligated probes, and run for the optimal number of PCR cycles calculated as described above.

Library postprocessing

Amplified libraries were purified with 100 μl AMPure XP beads (Beckman) and eluted in 20 μl resuspension buffer (10 mM Tris, pH8.0). Libraries were quantified with Picogreen (Life Technology) and usually yield 50-400 ng DNA. The libraries size patterns were checked using a fragment analyzer (Agilent).

Library Sequencing

An equimolar library pool was prepared from the individual libraries based on the concentrations

calculated from Picogreen data. Pools were denaturated with NaOH and neutralized with HT1

buffer (Illumina) to reach a final concentration of 20pM. Pools were spiked with 10 % PhiX and

clustered (loading equivalent to 12 samples per lane) onto a rapid single-end flow cell v2 at a

final concentration of 7pM. Sequencing was performed on a HiSeq 2500 (Illumina) in rapid mode

for 130 cycles.

Data Analysis

To assess the fidelity of left/right probe ligation and efficiency of hybridization, a fasta file with

all the possible combinations between left and right probes was created. In case two tRNA genes

share the same left/right probes, they were grouped and annotated accordingly. It led us to a

total of 68526 sequences. Genome index was generated with STAR with options

`-runMode genomeGenerate -genomeSAindexNbases 3`. Fastq files were trimmed for adapters.

Sequencing reads were aligned against the index with parameters:

`-outFilterScoreMinOverLread 0 -outFilterMatchNminOverLread 0`

`-winAnchorMultimapNmax 1000 -outFilterMismatchNmax`

`-clip3pNbases 6 -clip5pNbases 6`

`-outFilterMultimapNmax 50 -outSAMattributes NM nM NH NM -alignIntronMax 1`

`-alignEndsType EndToEnd -seedSearchStartLmax 20 -seedMultimapNmax 100000`.

For each combination, the number of counts were computed and corrected for PCR duplicates

using both unique molecular identifiers sequences. Reads larger than 60 nucleotides with less

than 3 mismatches and less than 5 insertions compared to the reference were retained. Combina-

tions with less than 10 reads were discarded. Reads mapping to combinations of probes coming

from different codons were reassigned in function of the newly ligated sequence. Abundances of tRNA coding for the same codon were summed up and normalized by library size using *edgeR* R package [84]. Because tRNA moieties (the two halves of the tRNA) have very similar sequences, and since the specificity of the hybrid DNA-RNA around the anticodon is important for the ligation, we used the sequence around the anticodon to reassign the ambiguous combinations.

Data Availability and specificity

Sequencing data of this study have been submitted to Gene Expression Omnibus (GEO) under accession number GSE126384: `https://www.ncbi.nlm.nih.gov/geo/query/acc.cgi?acc=GSE126384`). Datasets and GEO references: Atger Liver (GSE73553, n=84), Howard Liver (SRR826795, SRR826796, SRR826797, n=3), Huh Linter (SRR5227294, SRR5227295, SRR5227296, SRR5227303, SRR5227304, SRR5227305, n=6), Jan Yeast (SRR1562907, SRR1562909, SRR1562911, SRR1562913, n = 4), Kidney Castelo (GSE81283, n=24), Guydosh Yeast (SRR1042865, SRR1042866, SRR1042867, n= 1).

Acknowledgments

We thank Nicolas Bonhoure and Stefan Morgenthaler for useful discussions. Research in the Naef laboratory was supported by the EPFL. Some computations were performed on the Vital-It computing platform.

Conflict of interest statement

CG, BW, EM,and FG were employees of Nestle Institute of Health Sciences SA, CH-1015 Lausanne, Switzerland.

Supplementary material

Table 1. Inferred codon-specific DTs

Table 2. Inferred inter-site codon pair DTs

Table 3. Ribosome profiling in AL and FA mice w/o CHX

Table 4. (aminoacyl-) tRNA profiling in AL and FA mice

Code availability

Snakefile to model DTs and flux from ribosome profiling data will be made publicly available on github upon publication.

References

1. Gobet, C. & Naef, F. Ribosome profiling and dynamic regulation of translation in mammals. *Current Opinion in Genetics and Development* **43.** ISSN: 18790380. doi:10.1016/j.gde.2017.03.005 (2017).

2. Hinnebusch, A. G. & Lorsch, J. R. The mechanism of eukaryotic translation initiation: new insights and challenges. *Cold Spring Harbor perspectives in biology* **4,** a011544. ISSN: 1943-0264 (Oct. 2012).

3. Brule, C. E. & Grayhack, E. J. Synonymous Codons: Choose Wisely for Expression. *Trends in Genetics* **33,** 283–297. ISSN: 01689525 (Apr. 2017).

4. Presnyak, V. *et al.* Codon Optimality Is a Major Determinant of mRNA Stability. *Cell* **160,** 1111–1124. ISSN: 00928674 (Mar. 2015).

5. Radhakrishnan, A. *et al.* The DEAD-Box Protein Dhh1p Couples mRNA Decay and Translation by Monitoring Codon Optimality. *Cell* **0,** 1497–1506. ISSN: 00928674 (2016).

6. Pelechano, V., Wei, W. & Steinmetz, L. M. Widespread Co-translational RNA Decay Reveals Ribosome Dynamics. *Cell* **161,** 1400–1412. ISSN: 00928674 (June 2015).

7. Yu, C.-H. *et al.* Codon Usage Influences the Local Rate of Translation Elongation to Regulate Co-translational Protein Folding. *Molecular Cell* **59,** 744–754. ISSN: 10972765 (Sept. 2015).

8. Chu, D., Kazana, E., Bellanger, N., Singh, T., Tuite, M. F. & von der Haar, T. Translation elongation can control translation initiation on eukaryotic mRNAs. *The EMBO Journal* **33,** 21–34. ISSN: 02614189 (Jan. 2014).

9. Goodarzi, H., Nguyen, H. C., Zhang, S., Dill, B. D., Molina, H. & Tavazoie, S. F. Modulated Expression of Specific tRNAs Drives Gene Expression and Cancer Progression. *Cell* **165,** 1416–1427. ISSN: 00928674 (June 2016).

10. Loayza-Puch, F. *et al.* Tumour-specific proline vulnerability uncovered by differential ribosome codon reading. *Nature* **530,** 490–494. ISSN: 0028-0836 (Feb. 2016).

11. Rapino, F. *et al.* Codon-specific translation reprogramming promotes resistance to targeted therapy. *Nature* **558,** 605–609. ISSN: 14764687 (2018).

12. Varenne, S., Buc, J., Lloubes, R. & Lazdunski, C. Translation is a non-uniform process. *Journal of Molecular Biology* **180,** 549–576. ISSN: 00222836 (Dec. 1984).

13. Quax, T. E., Claassens, N. J., Söll, D. & van der Oost, J. Codon Bias as a Means to Fine-Tune Gene Expression. *Molecular Cell* **59,** 149–161. ISSN: 10972765 (July 2015).

14. Sørensen, M. A. & Pedersen, S. Absolute in vivo translation rates of individual codons in Escherichia coli. The two glutamic acid codons GAA and GAG are translated with a threefold difference in rate. *Journal of Molecular Biology* **222,** 265–280. ISSN: 00222836 (1991).

15. Ingolia, N. T., Lareau, L. F. & Weissman, J. S. Ribosome Profiling of Mouse Embryonic Stem Cells Reveals the Complexity and Dynamics of Mammalian Proteomes. *Cell* **147,** 789–802. ISSN: 00928674 (Nov. 2011).

16. Ingolia, N. T. Ribosome Footprint Profiling of Translation throughout the Genome. *Cell* **165,** 22–33. ISSN: 00928674 (Mar. 2016).

17. O'Connor, P. B. F., Andreev, D. E. & Baranov, P. V. Comparative survey of the relative impact of mRNA features on local ribosome profiling read density. *Nature Communications* **7,** 12915. ISSN: 2041-1723 (Oct. 2016).

18. Fang, H., Huang, Y. F., Radhakrishnan, A., Siepel, A., Lyon, G. J. & Schatz, M. C. Scikit-ribo Enables Accurate Estimation and Robust Modeling of Translation Dynamics at Codon Resolution. *Cell Systems.* ISSN: 24054720. doi:10.1016/j.cels.2017.12.007 (2018).

19. Tunney, R., McGlincy, N. J., Graham, M. E., Naddaf, N., Pachter, L. & Lareau, L. F. Accurate design of translational output by a neural network model of ribosome distribution. *Nature Structural and Molecular Biology* **25,** 577–582. ISSN: 15459985 (2018).

20. Charneski, C. A. & Hurst, L. D. Positively charged residues are the major determinants of ribosomal velocity. *PLoS biology* **11,** e1001508. ISSN: 1545-7885 (2013).

21. Darnell, A. M., Subramaniam, A. R. & O'Shea, E. K. Translational Control through Differential Ribosome Pausing during Amino Acid Limitation in Mammalian Cells. *Molecular Cell* **71,** 229–243. ISSN: 10974164 (2018).

22. Saikia, M., Wang, X., Mao, Y., Wan, J., Pan, T. & Qian, S.-B. Codon optimality controls differential mRNA translation during amino acid starvation. doi:`10.1261/rna.058180.116`.

23. Guydosh, N. R. & Green, R. Dom34 Rescues Ribosomes in 3 Untranslated Regions. *Cell* **156,** 950–962 (2014).

24. Nedialkova, D. D. & Leidel, S. A. Optimization of Codon Translation Rates via tRNA Modifications Maintains Proteome Integrity. *Cell* **161,** 1606–1618. ISSN: 00928674 (June 2015).

25. Tuorto, F. *et al.* The tRNA methyltransferase Dnmt2 is required for accurate polypeptide synthesis during haematopoiesis. *The EMBO Journal* **34,** 2350–2362 (2015).

26. Chou, H.-J., Donnard, E., Gustafsson, H. T., Garber, M. & Rando, O. J. Transcriptome-wide Analysis of Roles for tRNA Modifications in Translational Regulation. *Molecular Cell* **68,** 978–992. ISSN: 10972765 (Dec. 2017).

27. Pop, C. *et al.* Causal signals between codon bias , mRNA structure , and the efficiency of translation and elongation. *Molecular systems biology,* 1–16 (2014).

28. Zhang, S., Hu, H., Zhou, J., He, X., Jiang, T. & Zeng, J. Analysis of Ribosome Stalling and Translation Elongation Dynamics by Deep Learning. *Cell Systems* **5,** 212–220. ISSN: 24054720 (Sept. 2017).

29. Gorochowski, T. E., Ignatova, Z., Bovenberg, R. a. L. & Roubos, J. A. Trade-offs between tRNA abundance and mRNA secondary structure support smoothing of translation elongation rate. *Nucleic Acids Research* **43,** 3022–3032. ISSN: 0305-1048 (Mar. 2015).

30. Döring, K. *et al.* Profiling Ssb-Nascent Chain Interactions Reveals Principles of Hsp70-Assisted Folding. *Cell* **170,** 298–311. ISSN: 10974172 (2017).

31. Harigaya, Y. & Parker, R. The link between adjacent codon pairs and mRNA stability. *BMC Genomics* **18,** 364. ISSN: 1471-2164 (Dec. 2017).

32. Gamble, C. E., Brule, C. E., Dean, K. M., Fields, S. & Grayhack, E. J. Adjacent Codons Act in Concert to Modulate Translation Efficiency in Yeast. *Cell* **166,** 679–690. ISSN: 00928674 (July 2016).

33. Dao Duc, K. & Song, Y. S. The impact of ribosomal interference, codon usage, and exit tunnel interactions on translation elongation rate variation. *PLOS Genetics* **14** (ed Tuller, T.) e1007166. ISSN: 1553-7404 (Jan. 2018).

34. Gardin, J., Yeasmin, R., Yurovsky, A., Cai, Y., Skiena, S. & Futcher, B. Measurement of average decoding rates of the 61 sense codons in vivo. *eLife* **3.** ISSN: 2050-084X. doi:`10.7554/eLife.03735`. `https://elifesciences.org/articles/03735` (Oct. 2014).

35. Aeschimann, F., Xiong, J., Arnold, A., Dieterich, C. & Großhans, H. Transcriptome-wide measurement of ribosomal occupancy by ribosome profiling. *Methods* **85,** 75–89. ISSN: 10462023 (Sept. 2015).

36. Lareau, L. F., Hite, D. H., Hogan, G. J. & Brown, P. O. Distinct stages of the translation elongation cycle revealed by sequencing ribosome-protected mRNA fragments. *eLife* **3,** e01257. ISSN: 2050-084X (May 2014).

37. Duncan, C. D. S. & Mata, J. Effects of cycloheximide on the interpretation of ribosome profiling experiments in Schizosaccharomyces pombe. *Scientific Reports.* ISSN: 2045-2322. doi:`10.1038/s41598-017-10650-1` (2017).

38. Gerashchenko, M. V., Lobanov, A. V. & Gladyshev, V. N. Genome-wide ribosome profiling reveals complex translational regulation in response to oxidative stress. *Proceedings of the National Academy of Sciences* **109,** 17394–17399. ISSN: 0027-8424 (Oct. 2012).

39. Hussmann, J. a., Patchett, S., Johnson, A., Sawyer, S. & Press, W. H. Understanding Biases in Ribosome Profiling Experiments Reveals Signatures of Translation Dynamics in Yeast. *PLOS Genetics* **11** (ed Snyder, M.) e1005732. ISSN: 1553-7404 (Dec. 2015).

40. Bartholomäus, A., Del Campo, C. & Ignatova, Z. Mapping the non-standardized biases of ribosome profiling. *Biological Chemistry* **397.** ISSN: 1437-4315. doi:`10.1515/hsz-2015-0197`. `https://www.degruyter.com/view/j/bchm.2016.397.issue-1/hsz-2015-0197/hsz-2015-0197.xml` (Jan. 2016).

41. Quax, T. E., Claassens, N. J., Söll, D. & van der Oost, J. Codon Bias as a Means to Fine-Tune Gene Expression. *Molecular Cell* **59,** 149–161. ISSN: 10972765 (July 2015).

42. Orioli, A. tRNA biology in the omics era: Stress signalling dynamics and cancer progression. *BioEssays.* ISSN: 15211878. doi:`10.1002/bies.201600158` (2017).

43. Zheng, G. *et al.* Efficient and quantitative high-throughput tRNA sequencing. *Nature Methods* **12,** 835–837. ISSN: 1548-7091 (Sept. 2015).

44. Gogakos, T., Brown, M., Garzia, A., Meyer, C., Hafner, M. & Tuschl, T. Characterizing Expression and Processing of Precursor and Mature Human tRNAs by Hydro-tRNAseq and PAR-CLIP. *Cell Reports* **20,** 1463–1475. ISSN: 22111247 (Aug. 2017).

45. Cozen, A. E., Quartley, E., Holmes, A. D., Hrabeta-Robinson, E., Phizicky, E. M. & Lowe, T. M. ARM-seq: AlkB-facilitated RNA methylation sequencing reveals a complex landscape of modified tRNA fragments. *Nature Methods* **12,** 879–884. ISSN: 1548-7091 (Sept. 2015).

46. Evans, M. E., Clark, W. C., Zheng, G. & Pan, T. Determination of tRNA aminoacylation levels by high-throughput sequencing. *Nucleic Acids Research* **45,** e133–e133. ISSN: 0305-1048 (Aug. 2017).

47. Jan, C. H., Williams, C. C. & Weissman, J. S. Principles of ER cotranslational translocation revealed by proximity-specific ribosome profiling. *Science* **346,** 1257521–1257521. ISSN: 0036-8075 (Nov. 2014).

48. Ingolia, N. T., Hussmann, J. A. & Weissman, J. S. Ribosome Profiling: Global Views of Translation. *Cold Spring Harbor Perspectives in Biology,* a032698. ISSN: 1943-0264 (July 2018).

49. Atger, F. *et al.* Circadian and feeding rhythms differentially affect rhythmic mRNA transcription and translation in mouse liver. *Proceedings of the National Academy of Sciences* **112,** E6579–E6588. ISSN: 0027-8424 (Nov. 2015).

50. González, A. & Hall, M. N. Nutrient sensing and TOR signaling in yeast and mammals. *The EMBO Journal* **36,** 397–408. ISSN: 0261-4189 (Feb. 2017).

51. Gutierrez, E. *et al.* eIF5A Promotes Translation of Polyproline Motifs. *Molecular Cell* **51,** 35–45. ISSN: 10972765 (July 2013).

52. Lecanda, A. *et al.* Dual randomization of oligonucleotides to reduce the bias in ribosome-profiling libraries. *Methods* **107,** 89–97. ISSN: 10959130 (2016).

53. Gerashchenko, M. V. & Gladyshev, V. N. Translation inhibitors cause abnormalities in ribosome profiling experiments. *Nucleic Acids Research.* ISSN: 13624962. doi:`10.1093/nar/gku671` (2014).

54. Zhang, F., Xu, X., Zhou, B., He, Z. & Zhai, Q. Gene Expression Profile Change and Associated Physiological and Pathological Effects in Mouse Liver Induced by Fasting and Refeeding. *PLoS ONE* **6** (ed Androulakis, I. P.) e27553. ISSN: 1932-6203 (Nov. 2011).

55. Howard, M. T., Carlson, B. a., Anderson, C. B. & Hatfield, D. L. Translational redefinition of UGA codons is regulated by selenium availability. *Journal of Biological Chemistry* **288,** 19401–19413. ISSN: 00219258 (2013).

56. Castelo-Szekely, V., Arpat, A. B., Janich, P. & Gatfield, D. Translational contributions to tissue specificity in rhythmic and constitutive gene expression. *Genome Biology* **18.** ISSN: 1474760X. doi:`10.1186/s13059-017-1222-2` (2017).

57. Lintner, N. G. *et al.* Selective stalling of human translation through small-molecule engagement of the ribosome nascent chain. *PLOS Biology* **15** (ed Khosla, C.) e2001882. ISSN: 1545-7885 (Mar. 2017).

58. Dittmar, K. A., Sørensen, M. A., Elf, J., Ehrenberg, M. & Pan, T. Selective charging of tRNA isoacceptors induced by amino-acid starvation. *EMBO Reports.* ISSN: 1469221X. doi:`10.1038/sj.embor.7400341` (2005).

59. Lowe, T. M. & Chan, P. P. tRNAscan-SE On-line: integrating search and context for analysis of transfer RNA genes. *Nucleic acids research.* ISSN: 13624962. doi:`10.1093/nar/gkw413` (2016).

60. Canella, D. *et al.* A multiplicity of factors contributes to selective RNA polymerase III occupancy of a subset of RNA polymerase III genes in mouse liver. *Genome Research.* ISSN: 10889051. doi:`10.1101/gr.130286.111` (2012).

61. Rudolph, K. L. M. *et al.* Codon-Driven Translational Efficiency Is Stable across Diverse Mammalian Cell States. *PLoS Genetics* **12,** 1–23. ISSN: 15537404 (2016).

62. Pechmann, S. & Frydman, J. Evolutionary conservation of codon optimality reveals hidden signatures of cotranslational folding. *Nature Structural & Molecular Biology* **20,** 237–243. ISSN: 1545-9993 (Feb. 2013).

63. Woolstenhulme, C. J., Guydosh, N. R., Green, R. & Buskirk, A. R. High-Precision Analysis of Translational Pausing by Ribosome Profiling in Bacteria Lacking EFP. *Cell Reports* **11,** 13–21. ISSN: 22111247 (Apr. 2015).

64. Chevance, F. F. V. & Hughes, K. T. Case for the genetic code as a triplet of triplets. *Proceedings of the National Academy of Sciences* **114,** 4745–4750. ISSN: 0027-8424 (May 2017).

65. Kanaya, S., Yamada, Y., Kudo, Y. & Ikemura, T. Studies of codon usage and tRNA genes of 18 unicellular organisms and quantification of Bacillus subtilis tRNAs: Gene expression

level and species-specific diversity of codon usage based on multivariate analysis. *Gene.* ISSN: 03781119. doi:`10.1016/S0378-1119(99)00225-5` (1999).

66. Ingolia, N. T. Ribosome profiling: new views of translation, from single codons to genome scale. *Nature Reviews Genetics* **15,** 205–213. ISSN: 1471-0056 (Mar. 2014).

67. Mohammad, F., Woolstenhulme, C. J., Green, R. & Buskirk, A. R. Clarifying the Translational Pausing Landscape in Bacteria by Ribosome Profiling. *Cell Reports* **14,** 686–694. ISSN: 22111247 (Feb. 2016).

68. Wu, C. C.-C., Zinshteyn, B., Wehner, K. A. & Green, R. High-Resolution Ribosome Profiling Defines Discrete Ribosome Elongation States and Translational Regulation during Cellular Stress. *Molecular Cell* **0,** 1–12. ISSN: 10972765 (Jan. 2019).

69. Rodnina, M. V. The ribosome in action: Tuning of translational efficiency and protein folding. *Protein Science* **25,** 1390–1406. ISSN: 09618368 (Aug. 2016).

70. Agris, P. F., Eruysal, E. R., Narendran, A., Väre, V. Y. P., Vangaveti, S. & Ranganathan, S. V. Celebrating wobble decoding: Half a century and still much is new. *RNA Biology* **15,** 537–553. ISSN: 1547-6286 (May 2018).

71. Alkatib, S. *et al.* The Contributions of Wobbling and Superwobbling to the Reading of the Genetic Code. *PLoS Genetics* **8** (ed Maraia, R. J.) e1003076. ISSN: 1553-7404 (Nov. 2012).

72. Yang, J. R., Chen, X. & Zhang, J. Codon-by-Codon Modulation of Translational Speed and Accuracy Via mRNA Folding. *PLoS Biology.* ISSN: 15457885. doi:`10.1371/journal.pbio.1001910` (2014).

73. Stadler, M. & Fire, A. Wobble base-pairing slows in vivo translation elongation in metazoans. *RNA* **17,** 2063–2073. ISSN: 1355-8382 (Dec. 2011).

74. Pavlov, M. Y., Watts, R. E., Tan, Z., Cornish, V. W., Ehrenberg, M. & Forster, A. C. Slow peptide bond formation by proline and other N-alkylamino acids in translation. *Proceedings of the National Academy of Sciences* **106,** 50–54. ISSN: 0027-8424 (2009).

75. Lubas, M. *et al.* eIF5A is required for autophagy by mediating ATG3 translation. doi:`10.15252/embr.201846072`.

76. Pelechano, V. & Alepuz, P. eIF5A facilitates translation termination globally and promotes the elongation of many non polyproline-specific tripeptide sequences. *Nucleic Acids Research* **45,** 7326–7338. ISSN: 0305-1048 (July 2017).

77. Shalgi, R., Hurt, J. A., Krykbaeva, I., Taipale, M., Lindquist, S. & Burge, C. B. Widespread Regulation of Translation by Elongation Pausing in Heat Shock. *Molecular Cell* **49,** 439–452. ISSN: 10972765 (Feb. 2013).

78. Somogyi, P., Jenner, A. J., Brierley, I. & Inglis, S. C. Ribosomal pausing during translation of an RNA pseudoknot. *Molecular and cellular biology* **13,** 6931–40. ISSN: 0270-7306 (1993).

79. Tholstrup, J., Oddershede, L. B. & Sørensen, M. A. MRNA pseudoknot structures can act as ribosomal roadblocks. *Nucleic Acids Research.* ISSN: 03051048. doi:`10.1093/nar/gkr686` (2012).

80. Dittmar, K. A., Goodenbour, J. M. & Pan, T. Tissue-specific differences in human transfer RNA expression. *PLoS genetics* **2,** e221. ISSN: 1553-7404 (Dec. 2006).

81. Tatarinova, T. V., Alexandrov, N. N., Bouck, J. B. & Feldmann, K. A. GC3biology in corn, rice, sorghum and other grasses. *BMC Genomics* **11.** ISSN: 14712164. doi:`10.1186/1471-2164-11-308` (2010).

82. Wolfson, R. L. & Sabatini, D. M. The Dawn of the Age of Amino Acid Sensors for the mTORC1 Pathway. *Cell Metabolism* **26,** 301–309. ISSN: 19327420 (Aug. 2017).

83. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29,** 15–21. ISSN: 13674803 (2013).

84. Robinson, M. D., Mccarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *BIOINFORMATICS APPLICATIONS NOTE* **26,** 139–140 (2010).

Figure S1: **Modeling ribosome fluxes and codon-specific DT including ribosome inter-site interactions**

A. Ribosome elongation: Snapshot of two different mRNA species (m_1, m_2) translated with two different fluxes (F_1, F_2). Zoomed ribosome shows that numerous factors regulate ribosome DTs: 40 codons (+20,-20) around the E site are taken into account in the model to alleviate possible library biases, exit tunnel interactions, and influence of upstream/downstream sequences. Codon-pair interactions between the three sites (E, P, A) are also modeled. The ribosome densities on the mRNAs are estimated by RP, and modeled as genes fluxes multiplied by DTs. B. Site- and codon- specific DTs are visualized in a heatmap, relative to the position mean. The matrix of inter-sites interactions (E:P, P:A, and E:A) shows all possible combinations of codon pairs. All regression coefficients (gene fluxes and DTs) are inferred genome-wide. C. Bioinformatics pipeline: Sequencing reads are trimmed and mapped to the genome. Reads are first selected based on their size and E sites are assigned for each read. All annotated CDSs are parsed with a step of 3 nucleotides and number of reads are reported at each position. Genes with insufficient total read counts and read densities are removed, as well as regions with non-unique mappability. Dispersion parameters for negative binomial distributions are estimated for each sample and the GLM is fitted with a sparse design matrix and negative binomial (NB) noise model. DTs are centered (in log2 scale) and represented as shown in Fig. S1B. D. Construction of the data matrix for the GLM. Example of a gene CDS with two different positions (dark blue and dark red) covered by 56 and 2 reads, respectively. The assigned E, P, and A sites are shown. The CDS is parsed 3-by-3 and a matrix is designed with the corresponding position-dependent codons. E. Mean and variance of measured counts for pairs of codons occurring multiple times on a gene. The green line shows a *Poisson* regime with the variance equal to the mean. The red line represents the estimated fit for a negative binomial distribution (Methods). The dispersion parameter is estimated from these fits and used to parameterize the NB used in the GLM, independently for each sample.

Figure S2: **In yeast ribosome DT anticorrelate with codon usage and display site interactions**
Panels for the site-specific DTs are retrieved from the fit with the P:A interaction. A. Gene-specific Pearson correlation coefficient (ρ) between measure and fitted read counts. Number of genes with ρ larger than x-axis values are depicted. WT (green) and 3-AT (blue). B. Measured (black) and fitted (red) read counts for three representative genes. C. Heatmap representation of the DTs (log2, mean centered per site) in a window of 40 codons around the E site. Codons are ordered by AA and colored accordingly at the top of the heatmap. DTs with p >= 0.05 are not shown(set to zero). D. Interaction matrix for the pairs P:A (log2). Codons are colored according to AA. Codons in both sites are hierarchically clustered based on the euclidean distance matrix and a complete linkage algorithm. Fast and slow interactions are shown respectively in dark red and dark blue (colorbar). DTs with p-value >= 0.05 are set to zero.

Figure S3: **Sample specific DTs and correlations**
A. Gene-specific Pearson correlation coefficient (ρ) between measured and fitted read counts for 4 AL samples [49] . Number of genes with ρ larger than x-axis values are depicted. B. Measured (black) and fitted (red) read counts for three representative genes in one AL sample. C. Mean DTs and standard deviation over the 84 samples for the E, P, and A sites. D. DTs (log2, mean centered per site) at the E, P, and A sites for the 84 samples in the three conditions WT AL (WT ad libitum), WT RF (WT night-restricted feeding), and KO RF (*Bmal1* KO night-restricted feeding). E. Inter-sample Spearman correlation coefficients for site-specific DTs (single) and inter-site codon DTs for the interactions P:A, E:P, and E:A.

Figure S4: **Extended analysis of the translation elongation landscape in mouse liver**

Site-specific DTs are retrieved from the fit with the P:A interaction. A. Heatmap representation of the DTs (log2, mean centered per site) in a window of 40 codons around the E site. Codons are ordered by AA and colored accordingly. DTs with $p >= 0.05$ are not shown (set to zero). B. Interaction matrix for E:P pairs (log2). Codons are colored according to AA. Codons in both sites are hierarchically clustered based on the euclidean distance matrix and a complete linkage algorithm. Relatively fast and slow interactions are shown respectively in dark red and dark blue. DTs with p-value $>= 0.05$ are not shown (set to zero). C. DT (log2) distributions and boxplots for the three interaction terms P:A, E:P and E:A of the 84 samples in mouse liver. D. Differences of Bayesian Information Criterion (ΔBIC) between the model shown and the best model. (ΔBIC) is computed for each sample and proposed model, in which the alphabet for the DT covariates was taken as either the 20 natural AAs or the 61 sense codons. A lowercase 'a' on the right of an uppercase letter indicates that the AA alphabet was used for this position.

Figure S5: **Ribosome DTs are not affected in fasted mice and with CHX**
A. Gene-specific Pearson correlation coefficient (ρ) between measured and fitted read counts. Number of genes with ρ larger than x-axis values are depicted. WT (blue) and FA (green) B. Measured (black) and fitted (red) read counts for three representative genes in one AL sample. C. DTs in the E (top), P (middle) and A site (bottom) for the CHX vs. NOCHX conditions (left) and AL vs. FA conditions (right). Standard deviation is depicted. D. Pearson correlation coefficient between the different conditions for the interaction terms (log2) E:P, P:A, and E:A. E. Codon usage bias is computed for each gene up- or down-regulated in fasted animals and averaged. The difference in codon usage bias is computed between the FA and AL conditions. Codons are colored accordingly to their nucleotide at the third position (G-C in blue and A-T in red).

Figure S6: **Meta-analysis revealed conserved DT patterns and technical biases**
Analysis of published RP datasets in yeast (Yeast Jan) [47], mouse liver (Liver Atger, Liver Howard, Liver Gobet (this paper)) [49, 55], mouse kidney (Kidney Castelo) [56] and in a human hepatocyte cell line (Huh7 Lintner) [57]. DTs were inferred for each sample and averaged by condition. A. Spread of the DTs (max-min, log2) at every positions in a window of 40 codons around the E site for studies using small RNA library protocols. Colors show the different datasets. B. Same as (A) for studies using "cDNA circularization" library protocols. C. Correlation for the A site DTs between the different datasets (Pearson coefficient is color coded). D. Same as (C) for P:A interaction. E. DTs in the A site for the Liver Howard, Kidney Castelo and Huh7 Linter datasets vs. DTs in the A site from the RP data in this paper (Liver Gobet, AL and FA).

Figure S7: **(aminoacyl-) tRNA profiling in AL and FA mice (bis)**
A. tRNA profiling protocol. tRNAs were extracted in acidic conditions and uncharged tRNAs were hydrolyzed at 3'-end upon periodate treatment. tRNAs were treated with NaCl in control conditions. Then, tRNAs were deacylated and biotinylated at their 3'-end. Mix of left and right DNA probes were hybridized to the tRNA pools and pulled-down on magnetic beads through biotin-streptavidin interactions. Nicks in the anticodon between the left and right probes were ligated. tRNAs were degraded and DNA probes sequenced after amplification. B. Reads were mapped on every combination of left and right probes. Fraction of reads corresponding to left-right probe combinations belonging to the same codon or AA is reported for the 24 samples. The same measure is computed after reassignment of the probe combinations (Methods). C. tRNA abundances (log2) at the codon level for the control vs. altered conditions in which probes related to tRNAs coding for Pro and Gly were removed. D. tRNA abundances (log2) at the codon level for experiments with T4 or SplintR DNA ligases. Significant differences are shown in red. E. tRNA abundances (log2) at the probe level between biological replicates in the NaCl/AL condition at ZT04. F. Correlation between tRNA abundances in control AL vs. RNA polymerase III (POL3) ChIP-Seq signal quantified on the tRNAs gene loci. Data were extracted from the supplementary table of ref. [60].

Figure S8: **(aminoacyl-) tRNA profiling in AL and FA mice (bis)**
A. Fraction of reads (UMI) per AA for the different isoacceptor tRNAs. Colors indicate the nucleotide in position 34 on the tRNA. B. Heatmaps of the normalized tRNA read counts (log2) at the codon level for the 24 samples. tRNAs are ordered with hierarchical clustering. Dark blue (high), dark red (low). C. Expression levels for three tRNA coding respectively for Asn, Asp, and Ile, showing 5-8 fold differences between control and periodate-treated conditions.