Age-structured vectorial capacity in a ZIKV-*Ae. aegypti* model system

E. Handly Mayton[1], A. Ryan Tramonte[1], Rebecca C. Christofferson*[1,2]

[1]Department of Pathobiological Sciences, School of Veterinary Medicine, Department of

Pathobiological Sciences, Baton Rouge, USA

[2]Center for Computation and Technology, Louisiana State University, Baton Rouge,

USA

ABSTRACT

The transmission of vector-borne diseases is a dynamic process dependent on a

multitude of factors. A better understanding of the temporal factors that modify

transmission is needed. Here, we examined the impacts of age at the time of infectious

bloodmeal on the transmission potential of Zika virus (ZIKV) by *Aedes aegypti.* To do

this, we applied three treatments: treatment Z, where mosquitoes were exposed to ZIKV

at 5 days post-emergence; treatment M.Z, where mosquitoes were given a non-

infectious bloodmeal at 5 days post-emergence, followed by a ZIKV-infectious

bloodmeal at 12 days post-emergence; and treatment S.Z, where mosquitoes were

exposed to ZIKV at 12 days post-emergence. We then observed the impact of these

treatments on mosquito mortality, vector competence, and biting rate where we

introduce a modification of the human landing rate to measure "laboratory bloodmeal

landing rate" in the laboratory. Mosquitoes from treatment Z survived significantly longer

than treatments M.Z and S.Z, though this difference was modest. There were no

statistically significant differences observed in either vector competence or in biting

rates among the three treatment groups when assessed as functions of days post

exposure. When results were put in the context of mosquito age, we found a significant

age effect in both vector competence and biting rate. Subsequently, we combined the

life traits and viral kinetics results and modified the vectorial capacity equation in a way

that addresses these age-dependent interactions, providing insight into the transmission

potential of arboviruses as a function of mosquito age. Our principal finding is that

mosquito age at the time of the infectious blood meal affects the subsequent

transmission potential of the arbovirus and offers a means by which to quantify this

effect in our modified vectorial capacity equation.

Introduction

ZIka virus (ZIKV) is an arthropod-borne virus belonging to the family *Flaviviridae*.

ZIKV was first isolated from a primate in Uganda in 1947, and shortly thereafter an

Aedes mosquito in 1948 [1]. The primary vector in subsequent outbreaks is believed to

be *Ae. aegypti* [2]. The virus emerged in the Americas in 2015 leading to an

unprecedented outbreak, resulting in the WHO declaring ZIKV a public health

emergency [3]. ZIKV rapidly spread across South America and there was short-lived

autochthonous transmission in the Southern United States [4]. Unlike previous

outbreaks, however, ZIKV has been associated with congenital malformations and

autoimmune problems [1]. Because there are no antiviral treatments or vaccines

available, prevention relies on vector control and understanding the temporal dynamics

of virus-mosquito interactions.

Specifically, understanding the temporal nature of transmission and the

heterogeneities inherent in such is key for precision in estimates of risk [5-9]. The most

widely utilized measure of transmission potential is vectorial capacity (VC) given in the

equation below.

$$VC = \frac{ma^2bp^N}{-\ln(p)}$$

where 'm' is the density of mosquitoes, 'a' is the biting rate, 'b' is the vector competence

of the mosquito for a particular virus, 'N" is the extrinsic incubation period, and 'p' is the

daily probability of mosquito survival [10].

Vector competence is the ability of a mosquito to acquire and ultimately transmit

a virus. In order for the mosquito to become infectious and transmit a virus, several

events must occur. First, the mosquito must acquire an infectious bloodmeal and the

virus must establish infection in the midgut. Next, the virus escapes the midgut into the

hemocoel and hemolymph of the mosquito circulatory system, eventually making it to

the salivary glands. Shortly after the virus establishes infection in the salivary glands,

the mosquito becomes competent to transmit the virus to vertebrate hosts [5, 10]. The

time it takes for this process to occur is referred to as the extrinsic incubation period

(EIP). Not all mosquitoes that are exposed will be able to transmit, and the EIP is not a

constant. EIP has often been inferred from static sampling done at days 7 and 14 dpi,

which does not capture the dynamism of the process of vector competence [11-14]. The

EIP_{50} is the time it takes for 50% of the exposed mosquito population to become

infectious and offers a standardized means of reporting transmission potential of

arboviruses and other vector-borne diseases [15, 16].

Additionally, mosquito mortality is a critical factor in the spread of arboviruses like

ZIKV. Shorter lifespans may result in reductions in viral transmission as the mosquito

may not survive the EIP [15, 17-19]. However, in the traditional vectorial capacity

equation, the average probability of daily survival is used to determine the probability

that a mosquito will live through the EIP, though the EIP should be thought of in the

context of the age of the mosquito at the time of infectious bloodmeal [20, 21]. That is,

the length of the EIP has differing consequences for a mosquito that receives an

infectious bloodmeal at age 10 days old versus one that acquires virus at 20 days old.

Further, the biting rate of the mosquito is the most critical factor in determining

transmission potential. Biting rate is difficult to quantify and is not often measured in

laboratory studies [22, 23]. More often, biting rates are estimated by proxy. Human

landing rates are the gold standard as they measure the number of female mosquitoes

that land on a human during a given time [24]. While laboratory bloodmeal landing rates

appear to have limited field relevance, it may be able to inform models which can then

be compared to field data to potentially form suppositions about the makeup of the

effective density of transmitting mosquitoes compared to the overall density of

mosquitoes.

Ae. aegypti feed more than once and the timing at which an individual will

acquire an infectious bloodmeal is not uniform [25]. Thus, the age at which an infectious

bloodmeal occurs will determine the likelihood that the virus will get back out of the

mosquito and transmission will continue. Studies have addressed some of the issues of

age-structure in vectorial capacity, and this work builds on those studies [7, 17, 18, 21].

We were interested to know whether timing of the mosquito-infecting bloodmeal would

affect the transmission potential of ZIKV. To that end, we investigated age-specific life

traits of *Ae. aegypti* mosquitoes – mortality, biting rate, and vector competence for ZIKV

– and altered the vectorial capacity equation to account for these interactions relative to

infectious bloodmeal timing. As many of the parameters of vectorial capacity vary

across viral strain, genotype, and species, as well as mosquito species and conspecific

populations, this model will have utility to investigate differential age-dependence of

vector-pathogen interactions for determination of transmission potential.

Materials and Methods

Virus and mosquitoes

ZIKV strain PRVABC59 (ZIKV-PRV), originally isolated from a human patient in

Puerto Rico in 2015, was provided by Dr. Barbara Johnson at the US Centers for

Disease Control and Prevention. Prior to use, it was passaged three times in Vero cells

and cultured as in [26]. Supernatant was collected 3 days post-inoculation and titer

determined as previously described [26]. Titer of ZIKV was verified by qRT-PCR at

approximately 8×10^7 plaque forming units (pfu)/mL, matched across all exposure

experiments. Virus used for mosquito exposure was never frozen. Colony *Aedes*

aegypti (Rockefeller) were provided by Dr. Daniel Swale of the LSU Entomology

Department. Mosquitoes were maintained at constant conditions as in [27] with 16:8

light/dark periods and at 28°C constant temperature. Sucrose solution was removed 24

hours before experiments.

Blood-feeding and oral exposure of Ae. aegypti

At approximately 5 days post-emergence, mosquitoes were either exposed to an

infectious bloodmeal at a 1:2 virus supernatant-to-blood ratio as in [27] or to a mock

infected bloodmeal containing uninfected cell culture media at the same ratio as in [27].

Mosquitoes were allowed to feed via the Hemotek (Discovery Labs, UK) membrane

feeding system for 45 minutes, after which mosquitoes were cold anesthetized and

blood-fed females were placed into clean cartons until further treatment. Each treatment

was repeated three times. The three treatment groups were: ZIKV-infectious bloodmeal

given at 5 days post emergence (group 'Z'), mock bloodmeal at 5 days post emergence

followed by a ZIKV-infectious bloodmeal at approximately 12 days post emergence

(group 'M.Z'), and a last group that got only a single ZIKV-infectious bloodmeal at 12

days post emergence (group 'S.Z'). To prevent confounding, we starved all groups 24

hours prior to days 5 and 12 post-emergence (regardless if that group received a

bloodmeal) and all groups were cold anesthetized and moved to a new carton at days 5

and 12 post-emergence (again, regardless of whether they got a bloodmeal). Thus all

cohorts were treated in the exact same way with the exception of treatment conditions.

The experimental design is depicted in Figure 1.

Figure 1. Illustration of treatment design.
Three treatments were applied to assess our hypothesis. Treatment Z (ZIKV) received an infectious bloodmeal at 5 days post-emergence (dpe); Treatment M.Z (mock-ZIKV) received a mock infectious bloodmeal at 5 days dpe, followed by an infectious bloodmeal at 12 dpe; and Treatment S.Z (sugar-ZIKV) received an infectious bloodmeal at 12 dpe.

Infection kinetics of ZIKV

For vector competence studies, mosquitoes were sampled at 5, 8, and 11 days

post-exposure to test for infection and dissemination. Mosquito legs and bodies were

put into separate tubes containing 900 uL BA-1 diluent media and nickel plated BBs.

Samples were then homogenized twice at 25 Hz for 3 minutes using a Qiagen

Tissuelyzer. RNA was extracted and tested for the presence of viral RNA (see below).

Each treatment was replicated three times, and data are averages of all three replicates

per treatment.

Transmission and determination of EIP50

In order to test for transmission, an additional 10 mosquitoes per ZIKV treatment

were force-salivated. ZIKV-exposed mosquitoes were immobilized on ice before

removing legs and wings. Mosquitoes were then placed on double-sided tape, and the

proboscis of each mosquito was placed into a pipette tip containing 35 uL FBS with 3

mmol/L ATP for 30 minutes, as previously described in [28]. Contents of the pipette tip

were ejected into 100 uL BA-1 diluent and stored before testing (see below).

The EIP$_{50}$ was calculated by fitting a logistic regression (SSlogis function in R) to

the days post infection (DPI) and obtaining the value of the function at 50%.

$$T(t_{DPI}) = \frac{s}{1 + exp^{-\frac{DPI-x}{k}}}$$

Where s is the asymptote, x is the value of x at the inflection point, and k is the scale

parameter.

Mortality

Mortality studies were performed for the same three treatments as those used for

vector competence (Z, M.Z, and S.Z) as well as appropriate negative controls where a

mock bloodmeal was used in place of infectious bloodmeals. The three negative

controls were: 1) a mock bloodmeal at 5 days post-emergence (M) to correspond to the

Z treatment, 2) a mock bloodmeal at 5 days post-emergence, followed by another mock

bloodmeal at 12 days post-emergence (M.M) to correspond to the M.Z treatment, and 3)

a mock bloodmeal at 10 days post-emergence (S.M) to correspond to the S.Z

treatment. Another treatment was created where the mosquitoes were cold

anesthetized and sorted at approximately 5 days and 12 days post-emergence, but

were never blood fed (S). All treatments were cold anesthetized at 5 days, regardless of

whether they were offered a bloodmeal so that all mosquitoes experienced the same

treatment and mosquitoes were kept at relatively similar density/carton with an average

of 47 mosquitoes/carton (range 36-58). Each mortality treatment was repeated a total of

three times and data are averages of the three replicates.

Mosquitoes were checked daily and, when applicable, dead mosquitoes counted

for 23 days post initial treatment. For the purposes of this illustrative study, we took the

mortality study out to 23 days post exposure, or 28 days old (approximately 1 month),

as this has been shown to be the upper limit of believed survival of *Ae. aegypti* and is

similar to the range used in Tesla, et al. [28, 29]*.* For treatments where mosquitoes were

offered a later bloodmeal (M.M, S.M, M.Z, and S.Z), only mosquitoes that took one or

both bloodmeals were included in the mortality analyses. During the mortality study,

dead mosquitoes of ZIKV-exposed treatments were removed from the carton,

processed, and tested for infection and dissemination. Studies have previously shown

that viral RNA detection from dead mosquitoes was successful [30]. At 28 days old,

remaining live mosquitoes were taken down, counted, processed, and tested for

infection and dissemination.

Daily mortality estimates relative to age ($M(t_{age})$) were then predicted fitting an

asymptomatic regression (SSasymp function) using a non-linear least squares method

where:

$$M\left(t_{age}\right) = s + (r - s) * exp^{-exp^{c}*(-age)}$$

202   where s is the asymptote, r is the zero response parameter, and c is the logarithmic rate

203   constant.

 Laboratory blood-meal landing rates

Ten to twelve mosquitoes per day post exposure from each ZIKV-infection

treatments (Z, M.Z, and S.Z) were placed in individual, clear plastic canisters (Bioquip)

twenty-four hours before being provided a bloodmeal via membrane feeder using 1 mL

discs with 800 μL of blood (Hemotek, Discovery Labs, UK). Blood was kept at constant

temperature of 37°C and placed on top of the canisters. Herein we devised a method of

measuring a proxy biting rate similar to human landing rates, "laboratory bloodmeal

landing rate" [23]. This was assessed in two ways. First, mosquitoes were observed

through the clear canister for their general position in the canister and second, the disc

was removed to determine if they were on or near the mesh at the top of the canister. In

all cases, these two methods of observation matched. That is, if a mosquito was

observed to be at the blood meal prior to disc removal (looking through the canister),

she did not move to the bottom of the canister upon disc removal.

A mosquito was assessed as "landed" and recorded as "1" if the female was at

the top of the canister or "not landed" and recorded as "0" if she was at the bottom of

the canister. This observation was done at 1, 20, and 45 minutes post placement of the

disc and the disc was replaced between observation time points.

The sum of "landings" was taken for each mosquito as the individual frequency of

biting. The daily biting rate relative to age ($B(t_{age})$) was then predicted using the same

asymptotic regression model as described above.

$$B\left(t_{age}\right) = s + (r - s) * exp^{-exp^{c}*(-age)}$$

Additionally, we calculated a probability of biting where the frequency of biting

from above was characterized binomially as either having bitten at least once of the

three observations (coded as "1") or not at all (coded as "0"). The probability of biting as

a function of age ($Z(t_{age})$) was determined by fitting the proportion of mosquitoes that

landed or fed at least once a day using a self-starting non-linear least squares

regression as above (SSasymp).

$$Z(t_{age}) = s + (r - s) * exp^{-exp^c*(-age)}$$

For the ZIKV treatment, biting behavior was assessed at days post emergence

10, 13, 16 (to represent 5, 8, and 11 days post exposure) as well as 17, 20, 23, and 28

days post emergence to match age with the M.Z and S.Z treatments. (For clarity, days

17, 20, and 23 in M.Z and S.Z treatments correspond to days 5, 8, 11 post-exposure).

The blood that individual mosquitoes were exposed to were collected and serum

was separated by centrifugation at 6,000 rpm for 4 minutes at 4 degrees. Serum was

tested for the presence of ZIKV via RNA extraction and qRT-PCR.

Viral detection

RNA extractions performed using the MagMax-96 kit on a King Fisher nucleic

acid extraction instrument according to the manufacturer's instructions as in [26]. Viral

RNA was then detected via qRT-PCR using the SuperScript III One-Step RT-PCR

system with Platinum Taq on a Roche Lightcycler 480. Protocols and primers used were

previously described [31]. Samples with suspected positive Cp values above 35 were

inoculated onto confluent 6-well plates of Vero cells, rocked for 45 minutes, and

supernatant collected at 5 days post inoculation. For abdomen, leg, and forced

salivation samples, plating was performed with M199 Medium containing 2% FBS and

2% PSF. For serum samples, plating was performed with M199 Medium containing 2%

248 FBS and 4% PSF. Wells containing serum samples were also washed after rocking.

249 RNA was extracted and tested for the presence of viral RNA and tested as described.

250 Upon second testing, if the Cp value was again greater than 35, they were considered

251 negative.

253 *Statistics*

254  All statistics and subsequent graphics were performed and generated using R

255 version 3.4.3. To test for differences in mortality rates among treatments, Kaplan-Meier

256 survival analyses were conducted and the average time to death (TTD) estimated.

257 Differences among the treatment groups for infection, dissemination, and transmission

258 rates were tested by a chi-square test for multiple proportions on a day-by-day basis.

259 Biting rates were tested using a non-parametric analysis of variance (Kruskal-Wallis) for

260 treatment over each day post exposure.

262 *Vectorial capacity as a function of timing of infectious bloodmeal*

263  Using the data generated for blood feeding as a proxy for biting rate (see

264 Results), the mortality data to estimate daily probabilities of survival, and our

265 transmission data from forced salivation to predict the EIP_{50} of ZIKV in our mosquito

266 colony, we altered the vectorial capacity equation as follows to estimate vectorial

267 capacity as a function of when the mosquito acquires an infectious bloodmeal. In

268 addition, we add the parameter of probability of biting as a function of age (z).

$$VC_{age} = \frac{m(z_{age}a_{age} * z_i a_i)b \prod_1^i p_i}{-\ln(p_{age})}$$

Here we derive an age-structured vectorial capcity (VC_{age}) with the average time

to transmission of EIP_{50}. We then allow VC_{age} to be customized based on the timing of

the infectious bloodmeal taken by the mosquito.

To account for age-dependent mortality, we assign two different mortalities. The

traditional calculation of p^N calculated the probability of a mosquito living through to an

age of N, the EIP. In the context of an age-dependent vectorial capacity framework, we

can calculate a more precise probability of surviving based on the day the mosquito bit

(and obtained an infection) as p_i where *i* is the age at the time of a mosquito-infecting

bloodmeal + EIP_{50}. Allowing for the daily survival correction (denominator), p_{age} is the

probability of daily survival at the age of the mosquito acquiring a ZIKV infection. Where

p_{age} is 100%, we substituted 99% in order to satisfy non-zero denominator. Similarly, we

derive an age-dependent biting rate. Whereas a^2 – as the average biting rate – is

squared to account for the bite at which point a mosquito acquires an infection and the

second bite when the mosquito transmits this infection. Putting this in the context of

age-dependence, a^2 becomes the product of a_{age} (the biting rate at the age of infection

acquisition) and a_i (the biting rate at *i*). In addition, we introduced an additional

parameter to determine if accounting for the probability of biting per day is impactful.

This was parameterized as z_{age} and z_i to account for the probability of biting at times *age*

and *i*, as defined above.

Results

Mortality of *Aedes aegypti* is affected by exposure status and timing of bloodmeal

Pairwise comparisons of the ZIKV-infection treatments determined that the Z

group had a significantly longer TTD when compared to M.Z and S.Z (Figure 2). The

average time to death for the Z group was 25.9 days, 25.3 days for the M.Z group, and

24.5 days for the S.Z group. When the appropriate unexposed controls were compared

to each infection treatment group, there was a significant difference between the S.Z

and S.M comparison only, with an estimated difference in TTD of two days

(Supplemental Table 1). Of interest, the non-blood fed sugar-only controls died

significantly faster than any of the other treatments with an average TTD of 19.6 days

(Supplemental Table 1).

Figure 2. Survival curves of female *Ae. aegypti* by treatment.
Each line represents the combined data from three replicates per treatment: Zika (Z,
blue line), Mock-ZIKV (M.Z, red line), and sugar-ZIKA (S.Z, yellow line). Average time to
death of treatment Z was significantly longer than treatments M.Z and S.Z.
306

To determine if there was a generalized effect of exposure on mortality we

compared all groups with a ZIKV exposure to those without, excepting the group that

received no bloodmeal at all which was removed from the analysis. There was no

significant difference between ZIKV-exposed mosquitoes (groups Z, M.Z, and S.Z) and

the non-exposed groups (p>.05). When we predicted the daily survival rates, we pooled

all three ZIKV treatments together (Z, M.Z, S.Z) and obtained the parameter estimates

for the non-linear fit (Supplemental Figure 1). Though there was a significant difference

between Z and the M.Z and S.Z groups, the difference was relatively modest (0.6 and

1.4 days, respectively). The infection and dissemination rates of the dead mosquitoes

are given in Supplemental Table 2.

<u>ZIKV Infection Kinetics and Transmission Potential</u>

When we tested for differences in the infection and dissemination rates of *Ae.*

aegypti after ZIKV exposure, we found that the kinetics were virtually the same across

all three treatments when assessed as a function of days post exposure (dpe) (Figure

3A and3C). When we tested the proportion of infected and disseminated mosquitoes

per day among treatments, there were no significant differences noted (Supplemental

Table 3). Figure 3B and 3D show the same data as a function of age, rather than dpe.

Figure 3. Infection and dissemination rates per treatments.
Each line represents the combined data from three replicates per treatment.
Abdomens/legs sampled on days 5, 8, and 11 post-exposure. Rates of
infection/dissemination examined in the context of days post-exposure (A)(C) and
mosquito age (B)(D). No significant differences were noted between treatments at any
days post-exposure.

EIP_{50} and interaction with mosquito age

We first compared the proportion of transmitting mosquitoes at shared days post

exposure (5, 8, and 11 dpe). There was no transmission from any of the groups at days

5 and 8 post exposure. At day 11 post exposure, there was no significant difference in

transmission between the three treatments (p>0.05). Based on our findings with

infection and dissemination, we wanted to examine transmission in the context of age.

We compared the proportion of transmitting mosquitoes at shared ages (17, 20, 23, and

28 days post-emergence). At ages 17 and 20, only 10% of mosquitoes from Z showed

transmission, with no signs of transmission for the other two treatments. At age 23, we

found a significant difference among the three treatments (p<0.05). Z had the highest

proportion of transmission (60%), while S.Z had low transmission (10%) and M.Z had no

signs of transmission. Significant differences were also found among the three

treatments at age 28 (p<0.05), with no signs of transmission for M.Z or S.Z, and 60%

transmission for Z. This is summarized in Table 1.

We determined the EIP_{50}, or the time needed for 50% of the mosquitoes to

become infectious, for the Z-group only, as this was the only treatment to reach 50%

transmission rates. The EIP_{50} was determined to be 18.8 days post-exposure. This

translates to 23.8 days old in our system.

Table 1: Transmission rates for each day post exposure (DPE) and corresponding
mosquito age for each of the three treatment groups.

Treatment	DPE (Age)	% Transmission (n)
Z	5 (10)	0 (10)
	8 (13)	0 (10)
	11 (16)	10 (10)
	12 (17)	10 (10)
	15 (20)	10 (10)
	18 (23)	60 (10)
	23 (28)	60 (10)
M.Z	5 (17)	0 (10)
	8 (20)	0 (10)
	11 (23)	0 (10)
	16 (28)	0 (10)

S.Z	5 (17)	0 (10)
	8 (20)	0 (10)
	11 (23)	10 (10)
	16 (28)	0 (10)

360   <u>Biting Rates</u>

361   Using a modification of human landing rates which we term "bloodmeal landing

rate", we were able to estimate the biting rates as a function of age. There was no

difference among the three treatments with regard to biting rates across days post

exposure (p>.05) (Figure 4A). When we re-center the data and test for differences

among the groups as a function of age, there were no significant differences (p>.05)

(Figure 4B). We then combined all three treatments into a single exposed group and

determined whether infection or dissemination status affected the biting rate. There

were no significant differences in biting rates between infected mosquitoes and

uninfected mosquitoes, as well as those with a disseminated infection and those without

at each day post exposure (p>.05), though an effect of infection has been shown for

DENV-2 and DENV-3 [32-34]. Again combining all the treatments into a single group

due to no statistical differences, we determined that there was a day-to-day difference in

biting rate by age (p<.0001), indicating that age does affect the biting rate of *Ae.*

aegypti. The non-linear least squares fit is shown in Figure 4C, with a lower asymptote

of 0.

378
379 **Figure 4. Biting rates per treatment. (**A) Estimated biting rate per each treatment as a
380 function of day post ZIKV exposure. (B) Estimated biting rates per treatment as a
381 function of mosquito age. (C) Collapsed biting rates as a function of age across all three
382 treatments (dots) and daily biting rate predictions from fitted model (red line).
383

In addition, the proportion of mosquitoes that fed per day post emergence was

not significantly different among the treatment groups (p>.05). The individual per-day

biting proportions and the results of the non-linear least squares fit of probability of

biting as a function of age are given in Supplemental Figure 2.

<u>Vectorial Capacity as a function of age at the time of mosquito-infectious bloodmeal</u>

When we calculated VC_{age} with all average values (and assuming a constant

density of 1), we determined that the VC_{age} of ZIKV in our colony mosquitoes was

0.3119338 indicating a low likelihood of outbreak. However, when we calculated VC_{50}

as a function of the timing of the infectious bloodmeal, only days 1-10 post emergence

resulted in VC_{age} values greater than 1, indicative of outbreak potential [20]. The range

of VC_{age} over these 10 days was 66 at its height (day 1 post emergence) and 1.06 at

day 10 post emergence. This is shown in Figure 5, where we demarcate the window of

opportunity for biting based on age.

**Figure 5. Age-structured vectorial capacity (VC_{age}) based on mosquito age at time
of infectious bloodmeal.** The extrinsic incubation period 50 (EIP_{50}), or the time it takes
for 50% of the population to be transmitting, was used to calculate the VC_{age}. Red
shading shows the time range post-emergence in which mosquitoes must bite in order
to survive the EIP_{50} (days 1-10).

Discussion

Heterogeneity in vector-borne disease transmission systems has been shown to alter predictions of disease spread [5, 10, 27]. Temperature, for example, has been shown to be a critical modifier of vector competence and mosquito life traits [28, 35, 36]. Variable infectiousness of the human and the subsequent acquisition of virus by a mosquito has also been shown to have different estimates of transmission compared to average measures of virus acquisition by the vector [37] We demonstrate here that the interactions of several life traits and the viral kinetics of ZIKV can be parameterized in an age-structured framework.

Further, we demonstrate that this framework yields drastically different results relative to a measure of VC_{age} that uses averages for parameterization. At first glance, an EIP_{50} of 18.8 days would suggest that ZIKV transmission is an inefficient process, considering that laboratory mosquitoes likely live longer than populations in the field. However, by taking into account predicted daily survival and biting rates based on our data, we show that there is transmission potential ($VC_{age} \geq 1$) when the infectious bloodmeal is obtained within the first 10 days post emergence. We introduce the inclusion of the probability of biting per day to compliment biting rates. When we excluded the probability of feeding per day, the novel parameter introduced here, the range of VC_{50} increased to 71.3-1.6 over 10 days post emergence. This is likely due to the high values of predicted biting probabilities in the first 10 days post emergence. However, more data is needed to fully define the patterns of biting behaviors in *Ae. aegypti*, especially at early time points.

Of course, our system represents an artificial one where colony mosquitoes have

a higher inclination to take an artificial bloodmeal, which is based entirely on heat

seeking cues and does not account for other olfactory and chemosensory attractants,

and may have altered vector competence than field populations [38-40]. It has been

shown, however, that heat alone is sufficient to attract mosquitoes and at least initiate

host seeking behaviors [41]. However, the main purpose of our study was to

demonstrate, critically, that the timing of an infectious bloodmeal, coupled with

heterogeneity in mosquito life-traits and the interactions with viral kinetics alter

estimates of VC_{age}.

We show that mosquito age at the time of the infectious bite affects the

subsequent transmission potential of the arbovirus, and offers a means by which to

quantify this effect in our modified vectorial capacity equation. Indeed, if we alter the

EIP_{50} to 8 days, as some data suggests CHIKV to have [42], the VC_{age} reaches 203 if

the infectious bite occurs on the first day post emergence and there VC_{age} <1 at 18 days

post emergence. However, field infection rates remain quite low in surveillance settings

for several *Aedes*-borne viruses, which suggests that the number of mosquitoes that fall

within this window are quite small [43-45]. This could be due to several reasons. First,

the effective density of mosquitoes, which we did not address in this effort, may not be a

straightforward measure. Entomological indices such as the house index (HI), the

container index (CI), or the Breteau index (BI) are all indirect measures of density (how

many juveniles/houses (HI), how many containers are infected with juvenile stages (CI),

or breeding containers/100 houses (BI)) and do not directly assess actual contact and

have not been good corollaries of disease [46, 47]. In addition, these measures often

occur outside or adjacent to houses, while transmission of *Aedes*-borne viruses has

been shown to occur at the household level [48]. This suggests that a small subset of

mosquitoes may be responsible for most of the transmission, and could necessitate

further adaptation of the vectorial capacity equation to account for epidemiological

relevant measures of mosquito density.

Our study uses a laboratory system to investigate the consequences of

alternative parameterization of the vectorial capacity equation. We show that several

parameters are age-dependent and offer inclusion of a new parameter of biting

probability in addition to the daily biting rate. The influence of individual age-dependent

parameters should be carefully studied in the context of field-caught material and

outbreak relevant strains in order to fully appreciate the impact and necessity of these

modifications.

Acknowledgements: We'd like to thank all the members of the Christofferson Lab

(Chrissy Smith and Fausta Dutuze) and honorary member, Claire Birkenheuer, for

emotional support, insights, and exceptional cheerleading. Go Team. In addition, we'd

like to acknowledge Peggy, our mascot mosquito. May she rest in peace.

Funding: This work was supported by NIH/NIGMS grant R01GM122077.

References:

1.   Musso D, Gubler DJ. Zika Virus. Clin Microbiol Rev. 2016;29(3):487-524. Epub
2016/04/01. doi: 10.1128/CMR.00072-15. PubMed PMID: 27029595; PubMed Central PMCID:
PMCPMC4861986.

2.   Ciota AT, Bialosuknia SM, Zink SD, Brecher M, Ehrbar DJ, Morrissette MN, et al. Effects
of Zika Virus Strain and Aedes Mosquito Species on Vector Competence. Emerg Infect Dis.
2017;23(7):1110-7. Epub 2017/04/22. doi: 10.3201/eid2307.161633. PubMed PMID: 28430564;
PubMed Central PMCID: PMCPMC5512477.
3.   Gulland A. Zika virus is a global public health emergency, declares WHO. BMJ.
2016;352:i657. doi: 10.1136/bmj.i657. PubMed PMID: 26839247.
4.   Baud D, Gubler DJ, Schaub B, Lanteri MC, Musso D. An update on Zika virus infection.
Lancet. 2017;390(10107):2099-109. doi: 10.1016/S0140-6736(17)31450-2. PubMed PMID:
28647173.
5.   Valderrama A, Diaz Y, Lopez-Verges S. Interaction of Flavivirus with their mosquito
vectors and their impact on the human health in the Americas. Biochem Biophys Res Commun.
2017;492(4):541-7. Epub 2017/05/14. doi: 10.1016/j.bbrc.2017.05.050. PubMed PMID:
28499872.
6.   Dye C. The analysis of parasite transmission by bloodsucking insects. Annu Rev Entomol.
1992;37:1-19. Epub 1992/01/01. doi: 10.1146/annurev.en.37.010192.000245. PubMed PMID:
1539935.
7.   Maciel-de-Freitas R, Koella JC, Lourenco-de-Oliveira R. Lower survival rate, longevity and
fecundity of Aedes aegypti (Diptera: Culicidae) females orally challenged with dengue virus
serotype 2. Trans R Soc Trop Med Hyg. 2011;105(8):452-8. doi: 10.1016/j.trstmh.2011.05.006.
PubMed PMID: 21700303.
8.   Ruckert C, Ebel GD. How Do Virus-Mosquito Interactions Lead to Viral Emergence?
Trends Parasitol. 2018;34(4):310-21. doi: 10.1016/j.pt.2017.12.004. PubMed PMID: 29305089;
PubMed Central PMCID: PMCPMC5879000.
9.   Petersen MT, Silveira IDD, Tatila-Ferreira A, David MR, Chouin-Carneiro T, Van den
Wouwer L, et al. The impact of the age of first blood meal and Zika virus infection on Aedes
aegypti egg production and longevity. PLoS One. 2018;13(7):e0200766. doi:
10.1371/journal.pone.0200766. PubMed PMID: 30048481; PubMed Central PMCID:
PMCPMC6062029.
10.   Kramer LD, Ciota AT. Dissecting vectorial capacity for mosquito-borne viruses. Curr Opin
Virol. 2015;15:112-8. Epub 2015/11/17. doi: 10.1016/j.coviro.2015.10.003. PubMed PMID:
26569343; PubMed Central PMCID: PMCPMC4688158.
11.   Huang YS, Lyons AC, Hsu WW, Park SL, Higgs S, Vanlandingham DL. Differential
outcomes of Zika virus infection in Aedes aegypti orally challenged with infectious blood meals
and infectious protein meals. PLoS One. 2017;12(8):e0182386. Epub 2017/08/11. doi:
10.1371/journal.pone.0182386. PubMed PMID: 28796799; PubMed Central PMCID:
PMCPMC5552158.
12.   Chouin-Carneiro T, Vega-Rua A, Vazeille M, Yebakima A, Girod R, Goindin D, et al.
Differential Susceptibilities of Aedes aegypti and Aedes albopictus from the Americas to Zika
Virus. PLoS Negl Trop Dis. 2016;10(3):e0004543. Epub 2016/03/05. doi:
10.1371/journal.pntd.0004543. PubMed PMID: 26938868; PubMed Central PMCID:
PMCPMC4777396.
13.   Roundy CM, Azar SR, Rossi SL, Huang JH, Leal G, Yun R, et al. Variation in Aedes aegypti
Mosquito Competence for Zika Virus Transmission. Emerg Infect Dis. 2017;23(4):625-32. Epub

2017/03/14. doi: 10.3201/eid2304.161484. PubMed PMID: 28287375; PubMed Central PMCID:
PMCPMC5367433.
14.   Costa-da-Silva AL, Ioshino RS, Araujo HR, Kojin BB, Zanotto PM, Oliveira DB, et al.
Laboratory strains of Aedes aegypti are competent to Brazilian Zika virus. PLoS One.
2017;12(2):e0171951. Epub 2017/02/12. doi: 10.1371/journal.pone.0171951. PubMed PMID:
28187183; PubMed Central PMCID: PMCPMC5302382.
15.   Christofferson RC, Mores CN, Wearing HJ. Bridging the Gap Between Experimental Data
and Model Parameterization for Chikungunya Virus Transmission Predictions. J Infect Dis.
2016;214(suppl 5):S466-S70. Epub 2016/12/07. doi: 10.1093/infdis/jiw283. PubMed PMID:
27920175; PubMed Central PMCID: PMCPMC5137240.
16.   Ohm JR, Baldini F, Barreaux P, Lefevre T, Lynch PA, Suh E, et al. Rethinking the extrinsic
incubation period of malaria parasites. Parasit Vectors. 2018;11(1):178. Epub 2018/03/14. doi:
10.1186/s13071-018-2761-4. PubMed PMID: 29530073; PubMed Central PMCID:
PMCPMC5848458.
17.   Bellan SE. The importance of age dependent mortality and the extrinsic incubation
period in models of mosquito-borne disease transmission and control. PLoS One.
2010;5(4):e10165. Epub 2010/04/21. doi: 10.1371/journal.pone.0010165. PubMed PMID:
20405010; PubMed Central PMCID: PMCPMC2854142.
18.   Styer LM, Carey JR, Wang JL, Scott TW. Mosquitoes do senesce: departure from the
paradigm of constant mortality. Am J Trop Med Hyg. 2007;76(1):111-7. Epub 2007/01/27.
PubMed PMID: 17255238; PubMed Central PMCID: PMCPMC2408870.
19.   Maciel-de-Freitas R, Codeco CT, Lourenco-de-Oliveira R. Daily survival rates and
dispersal of Aedes aegypti females in Rio de Janeiro, Brazil. Am J Trop Med Hyg.
2007;76(4):659-65. PubMed PMID: 17426166.
20.   Novoseltsev VN, Michalski AI, Novoseltseva JA, Yashin AI, Carey JR, Ellis AM. An age-
structured extension to the vectorial capacity model. PLoS One. 2012;7(6):e39479. Epub
2012/06/23. doi: 10.1371/journal.pone.0039479. PubMed PMID: 22724022; PubMed Central
PMCID: PMCPMC3378582.
21.   Christofferson RC, Mores CN. Potential for Extrinsic Incubation Temperature to Alter
Interplay Between Transmission Potential and Mortality of Dengue-Infected Aedes aegypti.
Environ Health Insights. 2016;10:119-23. Epub 2016/08/02. doi: 10.4137/EHI.S38345. PubMed
PMID: 27478382; PubMed Central PMCID: PMCPMC4961054.
22.   Focks DA. A Review of Entomological Sampling Methods and Indicators for Dengue
Vectors. World Health Organization, 2003.
23.   Farjana T, Tuno N. Multiple blood feeding and host-seeking behavior in Aedes aegypti
and Aedes albopictus (Diptera: Culicidae). J Med Entomol. 2013;50(4):838-46. PubMed PMID:
23926783.
24.   Tangena JA, Thammavong P, Hiscox A, Lindsay SW, Brey PT. The Human-Baited Double
Net Trap: An Alternative to Human Landing Catches for Collecting Outdoor Biting Mosquitoes in
Lao PDR. PLoS One. 2015;10(9):e0138735. doi: 10.1371/journal.pone.0138735. PubMed PMID:
26381896; PubMed Central PMCID: PMCPMC4575072.
25.   Harrington LC, Fleisher A, Ruiz-Moreno D, Vermeylen F, Wa CV, Poulson RL, et al.
Heterogeneous feeding patterns of the dengue vector, Aedes aegypti, on individual human
hosts in rural Thailand. PLoS Negl Trop Dis. 2014;8(8):e3048. doi:

10.1371/journal.pntd.0003048. PubMed PMID: 25102306; PubMed Central PMCID:
PMCPMC4125296.
26.   Kawiecki AB, Mayton EH, Dutuze MF, Goupil BA, Langohr IM, Del Piero F, et al. Tissue
tropisms, infection kinetics, histologic lesions, and antibody response of the MR766 strain of
Zika virus in a murine model. Virol J. 2017;14(1):82. Epub 2017/04/20. doi: 10.1186/s12985-
017-0749-x. PubMed PMID: 28420392; PubMed Central PMCID: PMCPMC5395720.
27.   Christofferson RC, Mores CN. Estimating the magnitude and direction of altered
arbovirus transmission due to viral phenotype. PLoS One. 2011;6(1):e16298. Epub 2011/02/08.
doi: 10.1371/journal.pone.0016298. PubMed PMID: 21298018; PubMed Central PMCID:
PMCPMC3029343.
28.   Tesla B, Demakovsky LR, Mordecai EA, Ryan SJ, Bonds MH, Ngonghala CN, et al.
Temperature drives Zika virus transmission: evidence from empirical and mathematical models.
Proc Biol Sci. 2018;285(1884). Epub 2018/08/17. doi: 10.1098/rspb.2018.0795. PubMed PMID:
30111605; PubMed Central PMCID: PMCPMC6111177.
29.   Goindin D, Delannay C, Ramdini C, Gustave J, Fouque F. Parity and longevity of Aedes
aegypti according to temperatures in controlled conditions and consequences on dengue
transmission risks. PLoS One. 2015;10(8):e0135489. doi: 10.1371/journal.pone.0135489.
PubMed PMID: 26258684; PubMed Central PMCID: PMCPMC4530937.
30.   Burkhalter KL, Savage HM. Detection of Zika Virus in Desiccated Mosquitoes by Real-
Time Reverse Transcription PCR and Plaque Assay. Emerg Infect Dis. 2017;23(4):680-1. Epub
2017/01/12. doi: 10.3201/eid2304.161772. PubMed PMID: 28075325; PubMed Central PMCID:
PMCPMC5367413.
31.   Faye O, Faye O, Diallo D, Diallo M, Weidmann M, Sall AA. Quantitative real-time PCR
detection of Zika virus and evaluation with field-caught mosquitoes. Virol J. 2013;10:311. Epub
2013/10/24. doi: 10.1186/1743-422X-10-311. PubMed PMID: 24148652; PubMed Central
PMCID: PMCPMC4016539.
32.   Sylvestre G, Gandini M, Maciel-de-Freitas R. Age-dependent effects of oral infection
with dengue virus on Aedes aegypti (Diptera: Culicidae) feeding behavior, survival, oviposition
success and fecundity. PLoS One. 2013;8(3):e59933. Epub 2013/04/05. doi:
10.1371/journal.pone.0059933. PubMed PMID: 23555838; PubMed Central PMCID:
PMCPMC3612067.
33.   Maciel-de-Freitas R, Sylvestre G, Gandini M, Koella JC. The influence of dengue virus
serotype-2 infection on Aedes aegypti (Diptera: Culicidae) motivation and avidity to blood feed.
PLoS One. 2013;8(6):e65252. Epub 2013/06/12. doi: 10.1371/journal.pone.0065252. PubMed
PMID: 23755202; PubMed Central PMCID: PMCPMC3670916.
34.   Platt KB, Linthicum KJ, Myint KS, Innis BL, Lerdthusnee K, Vaughn DW. Impact of dengue
virus infection on feeding behavior of Aedes aegypti. Am J Trop Med Hyg. 1997;57(2):119-25.
Epub 1997/08/01. PubMed PMID: 9288801.
35.   Brady OJ, Johansson MA, Guerra CA, Bhatt S, Golding N, Pigott DM, et al. Modelling
adult Aedes aegypti and Aedes albopictus survival at different temperatures in laboratory and
field settings. Parasit Vectors. 2013;6:351. Epub 2013/12/18. doi: 10.1186/1756-3305-6-351.
PubMed PMID: 24330720; PubMed Central PMCID: PMCPMC3867219.

36.   Alto BW, Wiggins K, Eastmond B, Ortiz S, Zirbel K, Lounibos LP. Diurnal Temperature
Range and Chikungunya Virus Infection in Invasive Mosquito Vectors. J Med Entomol.
2018;55(1):217-24. Epub 2017/10/19. doi: 10.1093/jme/tjx182. PubMed PMID: 29040730.
37.   Tesla B, Demakovsky LR, Packiam HS, Mordecai EA, Rodriguez AD, Bonds MH, et al.
Estimating the effects of variation in viremia on mosquito susceptibility, infectiousness, and R0
of Zika in Aedes aegypti. PLoS Negl Trop Dis. 2018;12(8):e0006733. Epub 2018/08/23. doi:
10.1371/journal.pntd.0006733. PubMed PMID: 30133450; PubMed Central PMCID:
PMCPMC6122838.
38.   Friend WG, Smith JJ. Factors affecting feeding by bloodsucking insects. Annu Rev
Entomol. 1977;22:309-31. doi: 10.1146/annurev.en.22.010177.001521. PubMed PMID: 319741.
39.   Saratha R, Mathew N. Development of a mosquito attractant blend of small molecules
against host-seeking Aedes aegypti. Parasitol Res. 2016;115(4):1529-36. doi: 10.1007/s00436-
015-4886-0. PubMed PMID: 26693718.
40.   Majeed S, Hill SR, Dekker T, Ignell R. Detection and perception of generic host volatiles
by mosquitoes: responses to CO2 constrains host-seeking behaviour. R Soc Open Sci.
2017;4(5):170189. doi: 10.1098/rsos.170189. PubMed PMID: 28573028; PubMed Central
PMCID: PMCPMC5451829.
41.   Zermoglio PF, Robuchon E, Leonardi MS, Chandre F, Lazzari CR. What does heat tell a
mosquito? Characterization of the orientation behaviour of Aedes aegypti towards heat
sources. J Insect Physiol. 2017;100:9-14. doi: 10.1016/j.jinsphys.2017.04.010. PubMed PMID:
28479122.
42.   Ledermann JP, Borland EM, Powers AM. Minimum infectious dose for chikungunya virus
in Aedes aegypti and Ae. albopictus mosquitoes. Rev Panam Salud Publica. 2017;41:e65.
PubMed PMID: 28902278.
43.   Chen CF, Shu PY, Teng HJ, Su CL, Wu JW, Wang JH, et al. Screening of dengue virus in
field-caught Aedes aegypti and Aedes albopictus (Diptera: Culicidae) by one-step SYBR green-
based reverse transcriptase-polymerase chain reaction assay during 2004-2007 in Southern
Taiwan. Vector Borne Zoonotic Dis. 2010;10(10):1017-25. doi: 10.1089/vbz.2008.0069. PubMed
PMID: 21128850.
44.   Chow VT, Chan YC, Yong R, Lee KM, Lim LK, Chung YK, et al. Monitoring of dengue
viruses in field-caught Aedes aegypti and Aedes albopictus mosquitoes by a type-specific
polymerase chain reaction and cycle sequencing. Am J Trop Med Hyg. 1998;58(5):578-86.
PubMed PMID: 9598444.
45.   Chung YK, Pang FY. Dengue virus infection rate in field populations of female Aedes
aegypti and Aedes albopictus in Singapore. Trop Med Int Health. 2002;7(4):322-30. PubMed
PMID: 11952948.
46.   Bowman LR, Runge-Ranzinger S, McCall PJ. Assessing the relationship between vector
indices and dengue transmission: a systematic review of the evidence. PLoS Negl Trop Dis.
2014;8(5):e2848. doi: 10.1371/journal.pntd.0002848. PubMed PMID: 24810901; PubMed
Central PMCID: PMCPMC4014441.
47.   Yalwala S, Clark J, Oullo D, Ngonga D, Abuom D, Wanja E, et al. Comparative efficacy of
existing surveillance tools for Aedes aegypti in Western Kenya. J Vector Ecol. 2015;40(2):301-7.
doi: 10.1111/jvec.12168. PubMed PMID: 26611965.

48. Stoddard ST, Forshey BM, Morrison AC, Paz-Soldan VA, Vazquez-Prokopec GM, Astete H,
et al. House-to-house human movement drives dengue virus transmission. Proc Natl Acad Sci U
S A. 2013;110(3):994-9. doi: 10.1073/pnas.1213349110. PubMed PMID: 23277539; PubMed
Central PMCID: PMCPMC3549073.

S1 Figure. Observed and predicted daily probabilities of survival. Combined
observed daily survival rates of all three treatments (dots) and the predicted daily
survival rates (red curve).
659
660 **S2 Figure. Observed and predicted probabilities of daily biting.** The observed daily
661 biting rates (dots) from the "blood-meal" landing experiments and the fitted predictions
662 (red curve) with a force lower asymptote of 0.
663
**S1 Table. Average time to death and sample size for ZIKV-infection treatments
and unexposed controls used in the mortality study.** Exposed groups were the
three treatments exposed to ZIKV: Z -- ZIKV at 5 days post exposure (dpe); M.Z – mock
bloodmeal at 5 dpe and ZIKV bloodmeal at 12 dpe; S.Z – only a ZIKV bloodmeal at 12
dpe. Unexposed groups were not exposed to ZIKV but matched for bloodmeal uptake:
M – mock bloodmeal at 5 dpe; M.M – mock bloodmeals at 5 and 12 dpe; S.M – mock
bloodmeal at 12 dpe; and S – no bloodmeal, sugar only. Average times to death (TTD)
and total sample sizes per treatment are given below.
672
**S2 Table. Daily infection and dissemination proportions of dead or censored
mosquitoes from the three ZIKV-infection treatments of the mortality study.** Each
day when a mosquito was observed to have died, that mosquito was tested for
presence of ZIKV RNA in the abdomen (infection) and legs (disseminated infection).
The proportion of daily dead mosquitoes that were infected and/or disseminated are
given below, as well as the total number tested (n).
679
**S3 Table. Infection and dissemination rates for each day post-exposure (DPE) and
corresponding mosquito age for each of the three treatments.** Percent infection
was determined by the proportion of infected abdomens over total exposed; and percent
dissemination was determined as the proportion of infected legs over total exposed.
Three replicates were performed for each treatment, but proportions and sample sizes
(n) are combined from all three replicates.
686
687 **S4 Table. Modeled fits of parameters, the type of model, and the parameter
688 values.** For each parameter where predictions were made from experimental data, the
689 type of model (and R self-starting function) and parameter estimates for each fit are
690 given below.
691