1 **Adaptation and latitudinal gradients in species interactions: nest predation in**

2 **birds**

3   Benjamin G. Freeman[1,2*], Micah N. Scholer[1,2], Mannfred M. A. Boehm[1,3], Julian

4   Heavyside[1,2], Dolph Schluter[1,2]

6   [1] Biodiversity Research Centre, University of British Columbia

7   [2] Department of Zoology, University of British Columbia

8   [3] Department of Botany, University of British Columbia

9   *Corresponding author: freeman@zoology.ubc.ca. 604.822.2387. 2212 Main Mall,

10 Vancouver, BC V6T1Z4, Canada

Abstract

The "biotic interactions" hypothesis proposes that strong species interactions in the

tropics drive faster divergence in the tropics. However, support for the idea that

interactions are stronger in the tropics is mixed. Here we propose an explanation for

why observed interaction strengths might often be similar across latitudes. We

suggest populations might adapt to latitudinal differences in species interaction

regimes, which can have the effect of flattening observed latitudinal gradients in

interaction rates. To investigate this idea, we examine a canonical example of a strong

tropical biotic interaction—nest predation rates in land birds. Surprisingly, we find

that daily rates of nest predation vary minimally with latitude. We then consider the

possibility that evolved life history differences across latitudes contribute to the flat

gradient. We focus on the duration of the nesting period, which is longer in the tropics

than in the temperate zone and thought to represent a component of adaptation to

tropical nest predators. Greater nesting period duration is known to be associated with

lower daily predation rates. When we statistically control for nesting period duration,

we recover a pattern where daily rates of nest predation are highest in the tropics. The

implication of this analysis is that the evolution of longer nesting periods in tropical

species (and shorter nesting periods in the temperate zone) has helped to flatten a

"baseline" latitudinal gradient in daily rates of nest predation. More generally, we

propose that adaptation to latitudinal differences in biotic interactions can flatten

latitudinal gradients in interaction strength.

Keywords: biotic interaction, latitudinal gradient, latitudinal diversity gradient, nest

predation, predation, species interactions

Since Darwin, ecologists have suggested that biotic interactions increase in strength

towards the Equator (Darwin 1859; Wallace 1869; Dobzhansky 1950; MacArthur

1972). Dobzhansky (1950) expressed this viewpoint by arguing "Where physical

conditions are easy, interrelationships between species become the paramount

adaptive problem…This is probably the case in most tropical communities." Strong

biotic interactions in the tropics are hypothesized to generate strong selection that, in

turn, leads to faster rates of evolution and speciation in the tropics (Schemske 2009).

If so, strong biotic interactions in the tropics may explain in part why there are far

more species at low latitudes than in the temperate zone (Schemske 2009).

The biotic interactions hypothesis has inspired a growing number of studies

that test the prediction that total biotic interactions are indeed strongest in the tropics

(Schemske et al. 2009; Moles and Ollerton 2016). Two principal approaches use

measurements of the rates of biotic interactions to assess interaction strength across

latitudes. The first is to place a standardized artificial model at many sites, such as an

artificial prey type, and measure the frequency of biotic interactions, such as predator

attack rates, experienced by this model (Roslin et al. 2017; Hargreaves et al. 2019).

The second is to measure the frequency of biotic interactions that wild populations

actually experience, repeating this at a large number of sites across latitudes to

account for variability (e.g., Kubelka et al. 2018). A fundamental difference between

these two approaches is that standardized models are evolutionarily naïve and

measure rates of biotic interaction experienced by populations that have not adapted

to local interactions. By contrast, wild studies can measure rates of biotic interaction

experienced by populations that have had the opportunity to adapt to local species

interactions. The possibility exists that adaptation in response to interacting species

could reduce, or perhaps even eliminate, any latitudinal gradient in interaction rates.

This might explain why several studies using naïve models report higher interaction

rates at low latitudes (e.g., Roslin et al. 2017; Hargreaves et al. 2019) whereas many

observational studies of wild populations report latitudinal gradients in interaction

rates that are more flat (reviewed by Moles and Ollerton 2016).

Here we study rates of nest predation experienced by land birds, which are

widely thought to be greatest at low latitudes (Skutch 1985; Robinson et al. 2000;

Schemske et al. 2009; Mckinnon et al. 2010; Remeš et al. 2012; DeGregorio et al.

2016; Kubelka et al. 2018; but see Martin et al. 2017). The drivers of purported high

nest predation in the tropics are not certain, but one possible cause is the apparently

higher diversity of nest predators in the tropics, which include tropical predator

species that appear to prey almost exclusively upon eggs (DeGregorio et al. 2016;

Menezes and Marini 2017). To test these ideas, we compiled observational data on

daily rates of nest survival in wild Western Hemisphere birds from the published

literature. We use daily rates of nest survival to estimate daily rates of nest predation,

as the bulk of nest failure is due to predation and this is consistent across latitudes

(see methods). Tropical birds must deal with a distinct and diverse community of nest

predators, and we explore whether tropical birds have evolved life history traits that

reduce the realized rates of nest predation they experience in nature. We then

statistically control for one divergent life history trait to estimate a "baseline"

latitudinal gradient in rates of nest predation, and compare this to the observed

("derived") latitudinal gradient in rates of nest predation. Our synthesis thus

investigates not only latitudinal patterns in the intensity of a biotic interaction, but

also the interplay between ecological interaction and evolutionary consequence across

latitudinal gradients.

Methods

Assembling nest predation data

We searched the peer-reviewed literature to find studies that have measured nest

predation for land bird populations breeding in continental North, Central or South

America. We focus on the Western Hemisphere because nest predation data for

tropical birds outside of the Americas (e.g., from the Asian and African tropics) is

scarce (for data from Australia see Remeš et al. 2012). We included only studies of

real nests because we were interested in biotic interactions experienced by

populations in nature and because predation on artificial nests is poorly correlated

with predation on real nests (King et al. 1999; Burke et al. 2004; Moore and Robinson

2004; Robinson et al. 2005). Following previous studies investigating latitudinal

trends in nest predation, we restricted our analysis to land birds that are not cavity

nesters—primarily passerines, but also a small number of doves, hummingbirds, and

other non-passerines (Robinson et al. 2000; Remeš et al. 2012; Martin et al. 2017).

Several previous studies have analyzed nest predation data from the Americas (Nice

1957; Skutch 1985; Kulesza 1990; Conway and Martin 2000; Robinson et al. 2000;

Boyle et al. 2016; Martin et al. 2017). These syntheses are valuable summaries of

relevant studies, but did not always present the complete set of data that we were

interested in for each study. Hence, we extracted data from the original publications in

all cases. We only included studies that reported data based on at least 10 nests. When

studies found that nest predation differed between "natural" and "disturbed" habitats

(typically differently sized forest fragments), we used data from the "natural" habitat.

Last, we did not include data from studies in which authors explicitly stated that nest

failure was due to human activities such as mowing (e.g., birds nesting in hay fields),

or, in one case, when "much of the predation is known to have been by young boys"

(Peterson and Young 1950).

We located additional studies by conducting a Web of Science search in

February 2018 with the keywords "Nest predation" OR "Breeding ecology" OR "Nest

success" AND "bird". Because there is a latitudinal gradient in data availability (more

data in the temperate zone, less in the tropics), we expended additional effort to more

exhaustively search for tropical studies. Specifically, we: (1) conducted additional

country-specific Web of Science searches for each nation in Central and South

America, with keywords "Nest predation" OR "Nest success" AND "Country Name",

where "Country Name" was the name of each individual Central and South American

country; (2) examined the entire publication records of scientists who have

extensively studied Neotropical bird breeding biology; and (3) followed citation webs

to search for additional relevant studies from the tropics and southern temperate zone.

For each species from each study that met our criteria described above, we

extracted the following information: (1) Species name; (2) Sample size of nests; (3)

Nest success, presented either as fledging success (often termed "apparent success",

an estimate of the percentage of nests that successfully fledge young) or as daily

survival rate (the probability that an egg or nestling survives from one day to the

next); (4) Latitude and longitude; and (5) Clutch size, incubation and nestling periods,

extracted from the paper itself or from Handbook of the Birds of the World Alive (del

Hoyo et al. 2018).

Our final dataset included nest predation data for 515 unique species-site

combinations (from 244 studies and representing 314 species). The bulk of this

dataset comes from studies conducted between ~40° South and ~50° North. The

majority of studies came from the Northern Hemisphere temperate zone (269 unique

species-site combinations), but the tropics (absolute latitude < 23.4; 187 unique

species-site combinations) and Southern Hemisphere temperate zone (59 unique

species-site combinations) were also well represented (Fig 1). This dataset included

369 estimates of fledging success and 267 estimates of daily survival rate (see Fig

S1).

Nest predation vs. nest failure

Nest predation is difficult to study directly, as it is difficult to identify the cause of

nest failure. Predators are responsible for the large majority of nest mortality in most

places (Remeš et al. 2012; Martin et al. 2017), In our dataset, around one quarter of

studies (66 out of 244) attempted to identify causes of nest failure. Most nest failure

was attributed to predation (73%; N = 112 unique species-site combinations), a figure

remarkably similar to that previously reported for Australian birds (72%; Remeš et al.

2012). Further, the percentage of nest failures due to predation was similar across

latitudes (slope estimate for latitude in a univariate linear model with percentage of

nest failure attributed to predation as the response variable and latitude as a predictor

variable = -0.00083 ± 0.00074, p = 0.27). Hence, we follow previous analyses and

classify all nest failures as due to predation (Martin et al. 2017). This approach likely

slightly overestimates nest predation rates. However, it does not appear to introduce

any bias, and offers the advantage of greater clarity, as our focus in this paper is on

predators and nest predation.

Latitudinal variation in daily rates of nest predation

We assessed whether daily rates of nest predation (i.e., predation of the bird egg or

nestling) vary as a function of latitude by fitting mixed effect meta-analytic models

using the "metafor" package (Viechtbauer 2010) in R (R Development Core Team

2017). Data for this analysis were 267 estimates of daily survival rate, which we used

to calculate daily predation rate (132 from the north temperate zone, 90 from the

tropics, and 45 from the south temperate zone). The meta-analytic models weight

individual estimates by the inverse of their squared standard errors, and incorporate

the estimated variance among the study-specific effect sizes. We fit three models that

correspond to different biological hypotheses: (1) no latitudinal gradient in predation

(intercept-only model); (2) a linear latitudinal gradient in predation (single slope term,

equal slopes in both Northern and Southern hemispheres); and (3) a breakpoint linear

model wherein predation differs between the tropics and temperate zone (zero slope

within the tropics, ≤ 23.4 degrees absolute latitude, equal slopes for temperate

latitudes in both Northern and Southern hemispheres). We then compared model fits

using AIC. After determining the best-fit model, we fit an expanded version of the

best-fit model to test the influence of non-independence among effect sizes. This

model included species as a random effect, and we incorporated phylogenetic

relationships between species by specifying shared phylogenetic branch length in the

variance-covariance matrix. Branch lengths were measured from a majority rules

consensus tree calculated from 1000 phylogenies with the "Hackett" backbone pruned

to our study taxa and downloaded from birdtree.org (Jetz et al. 2012). We did not

include phylogenetic relationships in any of our initial models of latitudinal variation

in interaction rate, because we do not believe our question requires a phylogenetic

approach (e.g., in principle we could test for latitudinal variation in rates of nest

predation even if there was only one species with a broad latitudinal distribution in

the dataset). We then incorporated phylogenetic relationships into secondary models

to probe one potential explanation of our results, that latitudinal differences in nest

predation are related to phylogenetic differences between tropical and temperate taxa.

Latitudinal variation in nesting period duration

We found minimal variation in daily predation rates among latitudes (see Results). To

explore causes of this surprising result, we examined latitudinal patterns in nesting

period duration. We calculated nesting period length as the number of days from the

laying of the first egg until fledging, assuming that species begin incubating upon

laying the final egg. We first plotted species' nesting period durations vs. latitude

(using data on nesting period for 292 species from our dataset for which we could

calculated nesting period; 110 from the north temperate zone, 142 from the tropics,

and 40 from the south temperate zone), and observed clear latitudinal patterns in

nesting period duration, with longer nesting periods in the tropics (see Results). We

therefore tested the evolutionary association between latitude and nesting period

duration by fitting a phylogenetic generalized least squares (PGLS) regression using

the "ape" package (Paradis et al. 2004). The response variable in this model was

nesting period duration, absolute value of latitude was a fixed effect, Pagel's λ was

estimated using maximum likelihood, and we used a majority rules consensus tree

from 1000 phylogenies with the "Hackett" backbone downloaded from birdtree.org

(Jetz et al. 2012). In both reality and our dataset, oscine passerines predominate in the

temperate zone whereas suboscines are more common in the tropics. To explore

whether our overall results are mirrored within these groups, we repeated the PGLS

for oscines only ($N = 175$ species) and for suboscines only ($N = 86$ species).

To quantify the relationship between nesting period duration and daily

predation rates, we fit a meta-analytic model to predict daily predation rate using the

"metafor" package. Data for this analysis came from 175 species for which we had

both estimates of daily survival rate (used to calculate daily predation rate) and

information on nesting period length (132 from the north temperate zone, 84 from the

tropics, and 36 from the south temperate zone). We first fit a model that estimated

different slopes and intercepts for different latitudinal zones (predictor variables =

nesting period duration, latitudinal zone [north temperate, tropical, south temperate],

and an interaction between nesting period duration and latitudinal zone). We next fit a

model where latitudinal zones had different intercepts but the same slope (i.e., without

the interaction term between nesting period duration and latitudinal zone), and

compared fit of the "different slopes" and "same slopes" models using the "anova"

function in R. Last, we fit an expanded version of the "same slopes" model that

included species as a random effect and incorporated phylogeny by specifying

phylogenetic branch length as the variance-covariance matrix. Again, branch lengths

were measured from a majority rules consensus tree calculated from 1000

phylogenies with the "Hackett" backbone", pruned to our study taxa and downloaded

from birdtree.org (Jetz et al. 2012)

Latitudinal variation in fledging success

We examined latitudinal patterns in fledging success by repeating the three meta-

analytic models described above for latitudinal patterns in daily predation rate, but

with fledging success as the response variable. Data for this analysis were 369

estimates of fledging success (187 from the north temperate zone, 152 from the

tropics, and 30 from the south temperate zone). Upon plotting the data for the best-fit

model (breakpoint regression with equal slopes in Northern and Southern temperate

zones), we noticed that this model provided a poor fit to data from the Southern

temperate zone. We therefore fit an additional breakpoint regression model where we

allowed slopes to differ between Northern and Southern temperate zones, and

compared model fits using AIC. Last, we fit an expanded version of the best-fit model

that included species as a random effect and specified phylogenetic branch lengths as

the variance-covariance matrix (calculated as previously described).

Results

Latitudinal variation in daily rates of nest predation is minimal

We found evidence that daily rates of nest predation in land birds are similar across

latitudes within the Western Hemisphere. The most strongly supported model of daily

nest survival fit a line with equal daily predation rate (~ 0.04) across the entire

latitudinal gradient (Fig 2, Table 1). Alternative models that fit symmetric, non-zero

slopes to the relationship between latitude and daily survival rates were less well

supported (ΔAIC ~ 7; Table 1). These less supported models were similar to the best-

fit model in that they estimated slopes that were nearly flat (Table S1), with estimated

daily nest predation at 45° North only slightly lower (by ~ 0.004) than at the equator.

Our results did not change when including phylogenetic relationships and species

identity in our model (Table S2).

Nesting period length is a key covariate

We next conducted a series of exploratory analyses to investigate our finding of

minimal latitudinal variation in daily predation rate. We first examined patterns in a

conspicuous life history difference between latitudes, the length of the nesting period

(Martin 2002; Chalfoun and Martin 2007). In our dataset, tropical nesting periods

average approximately 15% longer than in the Northern temperate zone (~ 32 days vs

~ 28 days; Fig 3). This difference reflects a repeated pattern of evolution across a

diversity of avian lineages. Latitude is negatively related to nesting period in a

phylogenetic generalized least squares regression model (p < 0.0001, Table S3), and

patterns are similar when analyzing only oscines (temperate origin, predominate in

the temperate zone) or suboscines (tropical origin, predominate in the tropics; Table

S6). An equal slopes model estimated that a 10-day increase in nesting period is

associated with a 1.9% decrease in daily predation rates within each latitudinal zone

(Table S4).

We found that daily rates of nest predation are indeed greater in the tropics when

we statistically controlled for length of nesting period (North temperate zone vs.

tropics; p < 0.0001 in "same slopes" model to predict daily predation rate given

latitudinal zone and nesting period; Fig 4, Table S4). This "same slopes" model was

marginally better supported than a model that fit different slopes to different

latitudinal zones (p = 0.092). When we controlled for nesting period length in the

"different slopes" model, daily rates of nest predation were also greater in the tropics

(comparing estimates for North temperate zone vs. tropics for the mean nesting period

of 29 days, $p < 0.001$; Table S5, Fig S2). Again, this result is robust to potential

sources of non-independence in our dataset (Table S6).

Fledging success is lower in the tropics

The observation that nesting periods are longer in the tropics, while daily predation

rates are similar across latitudes, implies that fledging success (the probability that a

given nest survives to fledging) is lower in the tropics compared to the temperate

zone. Indeed, we found strong evidence that fledging success of nests is highest in the

Northern temperate zone (Fig 5, Table 2). A breakpoint regression estimated ~33% of

nests successfully fledge nestlings within the tropics while ~ 50% of nests

successfully fledge nestlings at 45° North (Table S7). Plotting this data revealed that

this breakpoint regression model poorly fit data from the Southern temperate zone. A

breakpoint regression model that fit different slopes to the Southern and Northern

temperate zones was better supported (Table 2); this model estimated that ~34% of

nests successfully fledge nestlings within both the tropics *and* the Southern temperate

zone, while ~ 53% of nests successfully fledge nestlings at 45° N, a result that is

robust to sources of non-independence in our data (Table S8). Although these

differences in nest predation between the tropics and temperate zone seem consistent

with the biotic interactions hypothesis, fledging success is integrated over a different

nesting duration in the tropics than the temperate zone, and so is not an accurate

measure of predation rate.

Discussion

The "biotic interactions" hypothesis proposes that biotic interactions are stronger in

the tropics, and that these strong interactions lead to faster evolution (and ultimately

speciation) in the tropics. Inspired by these ideas, we ask whether adaptation may

alter the link between ecological interaction and latitude. Our general hypothesis is

that adaptation to different communities of interacting species at different latitudes

can alter geographic patterns in interaction strength (see also Rasmann and Agrawal

2011; Chen et al. 2017). Hence, even if "baseline" interactions experienced by

evolutionarily naïve populations are generally stronger in the tropics, evolved

adaptations to different interaction regimes might flatten observed latitudinal

gradients in interaction strength. We propose that this key role for adaptation may

partially explain why some studies measuring interaction rates using "naïve"

standardized models show higher interaction rates in the tropics (e.g., Roslin et al.

2017; Hargreaves et al. 2019) while studies on wild "adapted" populations often show

minimal variation between interaction rate and latitude (reviewed by Moles and

Ollerton 2016).

Our data are consistent with a potential role for adaptation in influencing

latitudinal patterns in interaction rates. We report that there is little or no latitudinal

gradient in daily rates of nest predation. At the same time, there is a latitudinal

gradient in daily rates of nest predation when statistically controlling for nesting

period duration. Reconciling these results requires explicit consideration of the fact

that birds at different latitudes possess different life history adaptations that mitigate

baseline differences in nest predation rates. Tropical birds have longer nesting

periods, and longer nesting periods are generally associated with lower daily

predation rates. The implication is that if birds at all latitudes had the same nesting

period length, then tropical birds would suffer higher rates of nest predation than

temperate zone birds. We view this scenario—where we statistically control for

latitudinal divergence in a life history trait—as a measurement of "baseline"

interaction strength. However, in reality, birds are not equivalent across latitudes.

Instead, tropical and temperate species differ in life history adaptations including

nesting period length. When taking consideration of different adaptations along the

gradient, tropical birds no longer suffer higher rates of nest predation, and hence the

latitudinal gradient in "derived" interaction strength is more flat. A prediction arising

from this interpretation is that a standardized study of predation on artificial nests

(which are "evolutionarily naïve") across latitudes would find higher rates of nest

predation in the tropics. The only such standardized study we are aware of was

conducted at multiple latitudes within the Arctic. Consistent with our expectation, this

study reported higher predation rates on artificial nests at lower latitudes (Mckinnon

et al. 2010).

Our interpretation of latitudinal patterns in nest predation assumes that longer

nesting periods in tropical birds represent in part an adaptation to tropical predator

assemblages. Substantial evidence suggests that predation can be an important driver

of latitudinal differences in avian life history traits (Ghalambor and Martin 2001;

Martin 2002; Martin 2015). Tropical predator assemblages could select for long

nesting periods in tropical birds relative to temperate birds either directly or

indirectly. For example, high baseline predation risk in the tropics could directly

select for reduced rates of parental care. Less parental care per day at the nest is often

associated with longer nesting periods (Chalfoun and Martin 2007; Martin et al. 2007;

but see Tieleman et al. 2004), and the tendency for birds to reduce their rates of

activity at the nest when confronted with high predation risk is especially strong in the

tropics (Matysioková and Remeš 2018). Perhaps more plausibly, tropical predator

assemblages could indirectly select for long nesting periods in tropical birds via its

correlation with a suite of life history traits. For example, high predation could

directly select for increased investment in adult survival, with longer nesting periods

as an indirect effect (e.g., by a trade-off between adult investment in survival vs.

incubation/provision rates; Martin et al. 2015). The exact mechanism by which

predator-driven selection (among many sources of selection on life history traits)

favors longer nesting periods in tropical birds is not crucial to our analysis. Our

general idea is that adaptation to gradients in baseline interaction regimes can alter

observed rates of interaction. Applied to the bird nest predation data we present here,

our assumption is simply that variation in nesting period duration across latitudes

represents a component of adaptation to changing predator communities.

Comparison with previous studies of nest predation across latitudes

We report that natural rates of nest predation experienced by land birds are similar

along a latitudinal gradient in the Americas stretching from ~ 40° South to ~ 60°

North. This finding surprised us, because nest predation has been considered an

archetypical example in support of the more general hypothesis that biotic interactions

are stronger at low latitudes (Schemske 2009). Here we show that, at least for

Western Hemisphere land birds, the long-held view that nest predation is higher in the

tropics appears to be derived from observations of reduced fledging success of

tropical birds. However, latitudinal differences in fledging success are a consequence

of evolved differences between birds at different latitudes in a life history trait

(nesting period) rather than of differences in interaction rates. We focus on daily

predation rates because analyzing latitudinal patterns in a biotic interaction requires a

common currency for measuring interaction strength (Bulla et al. 2018). Whether the

pattern we document for the Western Hemisphere applies to the Eastern Hemisphere

remains uncertain; for example, rates of nest predation are indeed higher in the tropics

in Australia (Remeš et al. 2012; note that this study found much greater longitudinal

variation in nest predation than latitudinal variation).

Conclusions

The biotic interactions hypothesis invokes strong biotic interactions in the tropics to

explain in part the high species richness of the tropics. However, whether interactions

are stronger in the tropics remains hotly debated. On one hand, one review reported

general support for stronger biotic interactions in the tropics (Schemske et al. 2009).

On the other hand, a different review suggested the notion that interactions are

stronger in the tropics to be a "zombie idea"—an idea that is not supported by

empirical data but that refuses to die (Moles and Ollerton 2016). We propose that

adaptation that alters baseline interaction rates may in part reconcile these opposing

viewpoints (note that methodological differences are another factor responsible for

variation in the outcomes of observational studies). From this perspective, we expect

interactions to be stronger in the tropics when studies use standardized evolutionarily

naïve models to measure it (e.g., clay caterpillars or sunflower seeds; Roslin et al.

2017; Hargreaves et al. 2019), but not necessarily when studies measure interaction

strengths experienced by wild evolutionarily "adapted" populations.

More broadly, we argue that documenting geographic patterns in interaction

strength is only part of the story. The biotic interactions hypothesis is intriguing

precisely because it links biotic interactions to rates of divergence to explain

geographic patterns in species richness. We propose that variation in biotic

interactions across latitudes might exert strong selection for differential adaptations

along the gradient, and that these adaptations may reduce or flatten latitudinal

gradients in interaction strength. If so, adaptations could potentially flatten any

latitudinal gradient in evolutionary rates as well, though this remains untested. We

advocate for an increased focus on the evolutionary consequences of variation in the

strength of biotic interactions (Benkman 2013). Moving forward, we suggest that the

most rigorous tests of the biotic interaction hypothesis will consider how both biotic

interaction strength and evolutionary rates of trait divergence vary across latitude.

Acknowledgements and Funding

This research was supported by postdoctoral fellowships from the Biodiversity

Research Centre and Banting Canada (#379958) to BGF. None of our funders had

any influence on the content of the submitted manuscript, and none of our funders

required approval of the final manuscript to be published. Comments from the

Schluter lab group and Ralf Yorque greatly improved this manuscript.

References

Benkman, C. W. 2013. Biotic interaction strength and the intensity of selection.

Ecology Letters 16:1054–1060.

Boyle, A. W., B. K. Sandercock, and K. Martin. 2016. Patterns and drivers of

intraspecific variation in avian life history along elevational gradients: A

meta-analysis. Biological Reviews 91:469–482.

Bulla, M., K. Reneerkens, E. Weiser, A. Sokolov, A. Taylor, B. Sittler, B. McCaffery,

et al. 2018. Comment on "Global pattern of nest predation is disrupted by

climate change in shorebirds." Science 362:eaaw8529.

Burke, D. M., K. Elliott, L. Moore, W. Dunford, E. Nol, J. Phillips, S. Holmes, et al.

2004. Patterns of Nest Predation on Artificial and Natural Nests in Forests.

Conservation Biology 18:381–388.

Chalfoun, A. D., and T. E. Martin. 2007. Latitudinal variation in avian incubation

attentiveness and a test of the food limitation hypothesis. Animal Behaviour

73:579–585.

Chen, S. C., F. A. Hemmings, F. Chen, and A. T. Moles. 2017. Plants do not suffer

greater losses to seed predation towards the tropics. Global Ecology and

Biogeography 26:1283–1291.

Conway, C. J., and T. E. Martin. 2000. Evolution of Passerine Incubation Behavior :

Influence of Food , Temperature , and Nest Predation. Evolution 54:670–685.

Darwin, C. 1859. On the origin of species by means of natural selection. J. Murray,

London,.

DeGregorio, B. A., S. J. Chiavacci, T. J. Benson, J. H. Sperry, and P. J. Weatherhead.

2016. Nest Predators of North American Birds: Continental Patterns and

Implications. BioScience 66:655–665.

del Hoyo, J., A. Elliott, J. Sargatal, D. Christie, and de J. E. 2018. Handbook of the

Birds of the World Alive.

Dobzhansky, T. 1950. Evolution in the tropics. American Scientist 38:209–221.

Ghalambor, C. K., and T. E. Martin. 2001. Fecundity-survival trade-offs and parental

risk-taking in birds. Science 292:494–497.

Hargreaves, A. L., E. Suárez, K. Mehltreter, I. Myers-smith, S. E. Vanderplank, H. L.

Slinn, Y. L. Vargas-rodriguez, et al. 2019. Seed predation increases from the

Arctic to the Equator and from high to low elevations. Science Advances

5:eaau4403.

Jetz, W., G. H. Thomas, J. B. Joy, K. Hartmann, and A. O. Mooers. 2012. The global

diversity of birds in space and time. Nature 491:444–448.

King, D. I., R. M. DeGraaf, C. R. Griffin, and T. J. Maier. 1999. Do Predation Rates

on Artificial Nests Accurately Reflect Predation Rates on Natural Bird Nests?

Journal of Field Ornithology 70:257–262.

Kubelka, V., M. Šálek, P. Tomkovich, Z. Végvári, R. Freckleton, and T. Székely.

2018. Global pattern of nest predation is disrupted by climate change in

shorebirds. Science 362:680–683.

Kulesza, G. 1990. An analysis of clutch size in New World passerine birds. Ibis

132:407–422.

MacArthur, R. H. 1972. Geographical ecology: patterns in the distribution of species.

Princeton University Press, Princeton.

Martin, T. E. 2002. A new view of avian life-history evolution tested on an incubation

paradox. Proceedings of the Royal Society B: Biological Sciences 269:309–

316.

Martin, T. E. 2015. Age-related mortality explains life history strategies of tropical

and temperate songbirds. Science 349:966–970.

Martin, T. E., S. K. Auer, R. D. Bassar, A. M. Niklison, and P. Lloyd. 2007.

Geographic variation in avian incubation periods and parental influences on

embryonic temperature. Evolution 61:2558–2569.

Martin, T. E., A. J. Boyce, K. Fierro-Calderón, A. E. Mitchell, C. E. Armstad, J. C.

Mouton, and E. E. Bin Soudi. 2017. Enclosed nests may provide greater

thermal than nest predation benefits compared with open nests across

latitudes. Functional Ecology 31:1231–1240.

Martin, T. E., J. C. Oteyza, A. J. Boyce, P. Lloyd, and R. Ton. 2015. Adult mortality

probability and nest predation rates explain parental effort in warming eggs

with consequences for embryonic development time. American Naturalist

186:223–236.

Matysioková, B., and V. Remeš. 2018. Evolution of parental activity at the nest is

shaped by the risk of nest predation and ambient temperature across bird

species. Evolution 72:2214–2224.

Mckinnon, L., P. A. Smith, E. Nol, J. L. Martin, F. I. Doyle, K. F. Abraham, H. G.

Gilchrist, et al. 2010. Lower Predation Risk for Migratory birds. Science

327:6–8.

Menezes, J. C. T., and M. Â. Marini. 2017. Predators of bird nests in the Neotropics: a

review. Journal of Field Ornithology 88:99–114.

Moles, A. T., and J. Ollerton. 2016. Is the notion that species interactions are stronger

and more specialized in the tropics a zombie idea? Biotropica 48:141–145.

Moore, R. P., and W. D. Robinson. 2004. Artificial Bird Nests , External Validity,

and Bias in Ecological Field Studies. Ecology 85:1562–1567.

Nice, M. 1957. Nesting Success in Altricial Birds. The Auk 74:305–321.

Paradis, E., J. Claude, and K. Strimmer. 2004. APE: analyses of phylogenetics and

evolution in R language. Bioinformatics 20:289–290.

Peterson, A., and H. Young. 1950. A nesting study of the Bronzed Grackle. The Auk

67:466–476.

R Development Core Team. 2017. R: A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria.

Rasmann, S., and A. A. Agrawal. 2011. Latitudinal patterns in plant defense:

Evolution of cardenolides, their toxicity and induction following herbivory.

Ecology Letters 14:476–483.

Remeš, V., B. Matysioková, and A. Cockburn. 2012. Long-term and large-scale

analyses of nest predation patterns in Australian songbirds and a global

comparison of nest predation rates. Journal of Avian Biology 43:435–444.

Robinson, W. D., T. R. Robinson, S. K. Robinson, and J. D. Brawn. 2000. Nesting

success of understory forest birds in central Panama. Journal of Avian Biology

31:151–164.

Robinson, W. D., J. N. Styrsky, and J. D. Brawn. 2005. Are artificial bird nests

effective surrogates for estimating predation on real bird nests? A test with

tropical birds. The Auk 122:843–852.

Roslin, T., B. Hardwick, V. Novotny, W. K. Petry, N. R. Andrew, A. Asmus, I. C.

Barrio, et al. 2017. Latitudinal gradients: Higher predation risk for insect prey

522   at low latitudes and elevations. Science 356:742–744.

523 Schemske, D. 2009. Biotic interactions and speciation in the tropics. (Butlin R, Bridle

524   J, & Schluter D, eds.)Speciation and patterns of … 219–240.

525 Schemske, D. W., G. G. Mittelbach, H. V Cornell, J. M. Sobel, and K. Roy. 2009. Is

526   there a latitudinal gradient in the importance of biotic interactions? Annual

527   Review of Ecology, Evolution, and Systematics 40:245–269.

528 Skutch, A. F. 1985. Clutch size, nesting success, and predation on nests of

529   Neotropical birds, reviewed. Ornithological Monographs 36:575–594.

530 Tieleman, B. I., J. B. Williams, and R. E. Ricklefs. 2004. Nest attentiveness and egg

531   temperature do not explain the variation in incubation periods in tropical birds.

532   Functional Ecology 18:571–577.

533 Viechtbauer, W. 2010. Conducting meta-analyses in R with the metafor package.

534   Journal of Statistical Software 36:1–48.

535 Wallace, A. R. 1869. The Malay Archipelago. Cosimo Classics.

Figure 1. Map of studies measuring nest predation experienced by land birds in the

Americas (dataset = 244 studies). Many studies report data for multiple species from

the same site, illustrated by the size of the circle. The Tropics of Cancer and

Capricorn (at 23.4° N and S, respectively) delimit the tropics, and are illustrated with

dashed lines.

Figure 2. The latitudinal gradient in daily predation rates for land birds in the

Americas (N = 267). Predictions from the best-fit metafor model are plotted as a

dashed line—this simple model fit a constant value for daily predation rates across

latitudes. For comparison, the loess fit, which used the same weights for each data

point as the best-fit model, and incorporates the same estimated variance among the

study-specific effect sizes, is plotted in blue with shaded 95% confidence intervals.

Daily predation rates are estimated from daily survival rates reported in the literature.

Figure 3. The latitudinal gradient in nesting period length for land birds in the

Americas (N = 292 species). The dashed line illustrates the loess trendline, with 95%

confidence intervals shaded in gray. Nesting periods average ~ 32 days within the

tropics and Southern temperate zone but ~28 days within the Northern temperate

zone. Nesting period is calculated as the number of days from the laying of the first

egg to fledging.

Figure 4. The relationship between daily predation rate and nesting period duration

among latitudinal zones. Daily predation rate decreases with nesting period length in

each latitudinal zone, but with different intercepts. Predictions from a metafor model

are plotted as solid lines; values from individual studies are plotted in colors

corresponding to their latitudinal zone (N = 252). Daily predation rates are estimated

from daily survival rates reported in the literature; nesting period is calculated as the

number of days from the laying of the first egg to fledging.

Figure 5. The latitudinal gradient in fledging success for land birds in the Americas

(N = 367). Predictions from the best-fit model in metafor—a breakpoint regression

that fit a flat line within the tropics and different slopes in the Northern and Southern

Hemisphere temperate zones—, are plotted as a dashed line. For comparison, the

loess fit, which used the same weights for each data point as the best-fit model, and

incorporated the same estimated variance among the study-specific effect sizes, is

plotted in blue with shaded 95% confidence intervals.

582 Table 1. Model comparison of metafor models with daily predation rate as the

583 response variable. The best-fit model fit a constant value of daily predation rate across

584 the entire latitudinal expanse of the dataset.

Model	ΔAIC
Constant (daily predation rates do not vary with latitude)	--
Linear regression	6.89
Breakpoint regression (breakpoint = 23.4°)	7.54

587 Table 2. Model comparison of metafor models with fledging success as the response

588 variable. The best-fit model was a breakpoint regression that fit one slope within the

589 tropics and unequal slopes in the Northern and Southern temperate zones.

Model	ΔAIC
Breakpoint regression, unequal slopes (breakpoint = 23.4°)	--
Breakpoint regression, equal slopes (breakpoint = 23.4°)	4.68
Linear regression	21.29
Constant (fledging success does not vary with latitude)	66.35

Figure S1. Map of studies measuring daily rates of nest survival for land birds in the

Americas (dataset = 110 studies). We used these data to estimates daily rates of nest

predation, as the bulk of nest failure across latitudes is due to predation. Many studies

report data for multiple species from the same site, illustrated by the size of the circle.

The Tropics of Cancer and Capricorn (at 23.4° N and S, respectively) delimit the

tropics, and are illustrated with dashed lines.

Figure S2. The relationship between daily predation rate and nesting period duration

among latitudinal zones (different slopes model). Daily predation rate decreases with

nesting period length, but with different intercepts and slopes for different latitudinal

zones. Predictions from a metafor model are plotted as solid lines; values from

individual studies are plotted in colors corresponding to their latitudinal zone (N =

252). Daily predation rates are estimated from daily survival rates reported in the

literature; nesting period is calculated as the number of days from the laying of the

first egg to fledging.

620   Table S1. Parameter estimates for fixed effects from metafor models with daily

621   predation rate as the response variable. Slope estimates for the linear regression are

622   for straight lines outward from the equator towards higher latitudes; slope estimates

623   for breakpoint regression are for straight lines outward from the tropics (23.4 degrees

624   latitude) towards higher latitudes.

Model	Intercept ± SE	Slope ± SE
Intercept-only (constant)	0.042 ± 0.0015	NA
Linear regression	0.044 ± 0.0039	-0.0001 ± 0.0001
Breakpoint regression (breakpoint = 23.4°)	0.041 ± 0.0024	0.0001 ± 0.0002

627   Table S2. Model summary of intercept-only metafor model with daily predation rate

628   as the response variable and species and phylogeny as random effects.

Effect	Estimate
Intercept (fixed)	0.042 (se = 0.0083)
Species (random)	0.0000 (sqrt = 0.0067)
Phylogeny (random)	0.0003 (sqrt = 0.016)

635   Table S3. Parameter estimates for fixed effects from phylogenetic generalized least

636   squares regression models with nesting period as the response variable. Pagel's λ for

637   these model was 0.99 (entire dataset), 1.00 (oscines) and 0.94 (suboscines).

Model	Parameter	Estimate ± SE
Entire dataset	Intercept	27.29 ± 2.90
	Absolute value (latitude)	-0.040 ± 0.015
Oscines only	Intercept	34.92 ± 2.86
	Absolute value (latitude)	-0.026 ± 0.016
Suboscines only	Intercept	35.45 ± 2.99
	Absolute value (latitude)	-0.065 ± 0.039

643   Table S4. Parameter estimates from a metafor model with daily predation rate as the

644   response variable. This is the "equal slopes" model where different latitudinal zones

645   have the same slope. The reference category is the North temperate zone; parameter

646   estimates for the South temperate zone and tropics represent deviations from this

647   reference category.

Parameter	Estimate ± SE
Intercept	0.091 ± 0.0084
Nesting period	-0.0019 ± 0.0003
South temperate zone	0.0049 ± 0.0047
Tropics	0.015 ± 0.0037

649 Table S5. Parameter estimates from a metafor model with daily predation rate as the

650 response variable. This is the "different slopes" model where different latitudinal

651 zones have different slopes. The reference category is the North temperate zone;

652 parameter estimates for "Nesting Period: South temperate zone" and "Nesting Period:

653 Tropics" represent deviations from this reference category.

Parameter	Estimate ± SE
Intercept	0.10 ± 0.012
Nesting period	-0.0025 ± 0.00043
South temperate zone	-0.038 ± 0.035
Tropics	-0.024 ± 0.019
Nesting period: South temperate zone	0.0015 ± 0.0012
Nesting period: Tropics	0.0013 ± 0.00063

Table S6. Parameter estimates from a metafor model with daily predation rate as the

response variable and study, species and phylogeny as random effects. This is the

"equal slopes" model where different latitudinal zones have the same slope. The

reference category for the intercept is the North temperate zone; parameter estimates

for the South temperate zone and tropics represent deviations from this reference

category.

Parameter	Estimate ± SE
Intercept	0.091 ± 0.0084
Nesting period	-0.0019 ± 0.00030
South temperate zone	0.0050 ± 0.0047
Tropics	0.015 ± 0.0037
Species (random)	0.0 (sqrt = 0.0064)
Phylogeny (random)	0.0 (sqrt = 0.0)

Table S7. Parameter estimates for fixed effects from metafor models with fledging

success as the response variable. Slope estimates for the linear regression are for

straight lines outward from the equator towards higher latitudes; slope estimates for

breakpoint regression are for straight lines outward from the tropics (23.4 degrees

latitude) towards higher latitudes. Breakpoint regressions fit either equal slopes (same

slope value in both Northern and Southern temperate zones) or unequal slopes

(different slopes in Northern vs. Southern temperate zones).

Model	Intercept ± SE	Slope ± SE
Intercept-only (constant)	0.41 ± 0.010	NA
Linear regression	0.29 ± 0.019	0.0043 ± 0.0006
Breakpoint regression, equal slopes (breakpoint = 23.4°)	0.33 ± 0.013	0.0080 ± 0.001
Breakpoint regression, unequal slopes (breakpoint = 23.4°)	0.34 ± 0.013	-0.0008 ± 0.0032 (S Hemisphere) 0.0089 ± 0.0031 (N Hemisphere)

677   Table S8. Model summary of unequal slopes breakpoint regression metafor model

678   with fledging success as the response variable, and species and phylogeny included as

679   random effects.

Effect	Estimate
Intercept (fixed)	0.34 (se = 0.013)
Slope (fixed)	-0.0008 ± 0.0032 (S Hemisphere)
	0.0089 ± 0.0031 (N Hemisphere)
Species (random)	0.026 (sqrt = 0.16)
Phylogeny (random)	0 (sqrt = 0.0000)