Adaptation and latitudinal gradients in species interactions: nest predation in

birds

Benjamin G. Freeman[1,2*], Micah N. Scholer[1,2], Mannfred M. A. Boehm[1,3], Julian

Heavyside[1,2], Dolph Schluter[1,2]

[1] Biodiversity Research Centre, University of British Columbia

[2] Department of Zoology, University of British Columbia

[3] Department of Botany, University of British Columbia

*Corresponding author: freeman@zoology.ubc.ca. 604.822.2387. 2212 Main Mall,

Vancouver, BC V6T1Z4, Canada

Abstract

The "biotic interactions" hypothesis—that stronger interspecific interactions in the

tropics drive faster evolution and speciation, giving rise to the latitudinal diversity

gradient—has inspired many tests of whether certain biotic interactions are indeed

stronger in the tropics. However, the possibility that populations have adapted to

latitudinal differences in species interactions, potentially blunting effects on

evolutionary rates, has been largely ignored. Here we examine latitudinal patterns in

nest predation in Western Hemisphere birds, a canonical example of a strong tropical

biotic interaction. Surprisingly, daily rates of nest survival experienced by land birds

vary minimally with latitude. Only when controlling for length of nesting period do

tropical birds suffer higher predation than temperate zone birds. Tropical birds have

evolved particularly long nesting periods, and long nesting periods are generally

associated with reduced predation rates. Hence, long nesting periods may be a

component of adaptation by tropical birds to a diverse community of nest predators.

More generally, we suggest that adaptation to strong biotic interactions can alter

observed latitudinal gradients in interaction strength, potentially equalizing

evolutionary rates among latitudes. Moving forward, tests of the biotic interactions

hypothesis should consider measuring both latitudinal patterns in interaction strength

and evolutionary responses to these interactions.

Keywords: biotic interaction, latitudinal gradient, latitudinal diversity gradient, nest

predation, predation, species interactions

Since Darwin, ecologists have suggested that biotic interactions increase in strength

towards the Equator (Darwin 1859; Dobzhansky 1950; MacArthur 1972; Wallace

2007). Dobzhansky (1950) expressed this viewpoint by arguing "Where physical

conditions are easy, interrelationships between species become the paramount

adaptive problem…This is probably the case in most tropical communities." Strong

biotic interactions in the tropics are hypothesized to generate strong selection that, in

turn, leads to faster rates of evolution and speciation (Schemske 2009). If so, strong

tropical biotic interactions may explain in part why there are far more species at low

latitudes than in the temperate zone (the "biotic interactions hypothesis; Schemske

2009).

The biotic interactions hypothesis has inspired a growing number of studies

that test the prediction that total biotic interactions are indeed strongest in the tropics

(Schemske *et al.* 2009; Moles & Ollerton 2016). There are two principal approaches

to measuring rates of biotic interactions. First, one can place a standardized model at

many sites, and measure the frequency of biotic interactions experienced by this

model (e.g., Roslin *et al.* 2017; Hargreaves *et al.* 2019). Second, one can measure the

intensity of biotic interactions that wild local populations actually experience,

repeating this across a large number of sites to account for variability (e.g., Kubelka *et*

al. 2018). A fundamental difference between these two approaches is that

standardized model studies attempt to measure the biotic interaction as experienced

by populations that have not adapted to local interactions, whereas wild studies

measure biotic interactions as experienced by populations that have had the

opportunity to adapt to local interactions (i.e., "naïve" vs. "adapted"). Hence,

standardized model studies neglect the possibility that adaptation in response to

interaction strength and the perpetrators of these interactions could reduce, or perhaps

even eliminate, any gradient. If so, adaptations could blunt any differential effects of

interactions on evolutionary rates in the tropics vs. the temperate zone.

Here, we examine the biotic interactions hypothesis by studying nest predation

in land birds. Nest predation experienced by birds is widely held to be most intense at

lower latitudes (Skutch 1985; Robinson *et al.* 2000; Schemske *et al.* 2009; Mckinnon

et al. 2010; Remeš *et al.* 2012; DeGregorio *et al.* 2016; Kubelka *et al.* 2018), but see

(Martin *et al.* 2017). High nest predation in the tropics is thought to be related to the

fact that the tropics are home to both more species and more types of nest predators

than the temperate zone, including some tropical species that appear to prey almost

exclusively upon eggs (DeGregorio *et al.* 2016; Menezes & Marini 2017). Here, we

test latitudinal patterns in nest predation using a newly compiled dataset of estimates

of daily rates of nest survival experienced by land bird populations across the Western

Hemisphere. Our study measures interactions experienced by populations in nature

(i.e., "adapted" interactions). We therefore explore whether tropical birds, which must

deal with a distinct and diverse community of nest predators, might have evolved

adaptations that reduce the rates of nest predation they experience in nature.

Specifically, we examine the possibility that nesting period length is a component of

adaptation to the higher predator diversity in the tropics. Our synthesis thus

investigates not only latitudinal patterns in the intensity of a biotic interaction, but

also the interplay between ecological interaction and evolutionary consequence across

latitudinal gradients.

Methods

Assembling nest predation data

We searched the peer-reviewed literature to find studies that have measured nest

predation for land bird populations breeding in continental North, Central or South

America. We focus on the Western Hemisphere because nest predation data for

tropical birds outside of the Americas (e.g., the Asian and African tropics) is scarce

(but see Remeš *et al.* 2012 for an analysis of Australian birds). We included only

studies of real nests because we were interested in biotic interactions experienced by

populations in nature, as predation on artificial nests is poorly correlated with

predation on real nests (King *et al.* 1999; Burke *et al.* 2004; Moore & Robinson 2004;

Robinson *et al.* 2005). Following previous studies investigating latitudinal trends in

nest predation, we restricted our analysis to land birds that are not cavity nesters—

primarily passerines, but also a small number of doves, hummingbirds, and other non-

passerines (Robinson *et al.* 2000; Remeš *et al.* 2012; Martin *et al.* 2017). Several

previous studies have analyzed nest predation data from the Americas (Nice 1957;

Skutch 1985; Kulesza 1990; Conway & Martin 2000; Robinson *et al.* 2000; Boyle *et*

al. 2016; Martin *et al.* 2017). These syntheses are valuable summaries of relevant

studies, but did not always present the complete set of data that we were interested in

for each study. Hence, we extracted data from the original publications in all cases.

We only considered studies that reported the fate of at least 10 nests, and did not

include studies where authors explicitly described that nest failure was due to human

activities such as mowing (e.g., birds nesting in hay fields), or, in one case, when

"much of the predation is known to have been by young boys" (Peterson & Young

1950).

We located additional studies by conducting a Web of Science search in

February 2018 with the keywords "Nest predation" OR "Breeding ecology" OR "Nest

success" AND "bird". Because there is a latitudinal gradient in data availability (more

data in the temperate zone, less in the tropics), we expended additional effort to more

exhaustively search for tropical studies. Specifically, we: (1) conducted additional

country-specific Web of Science searches for each nation in Central and South

America, with keywords "Nest predation" OR "Nest success" AND "Country Name",

where "Country Name" was the name of each individual Central and South American

country; (2) examined the entire publication records of scientists who have

extensively studied Neotropical bird breeding biology; and (3) followed citation webs

to search for additional relevant studies from the tropics and southern temperate zone.

For each species from each study that met our criteria described above, we

extracted the following information: (1) Species name; (2) Sample size of nests; (3)

Nest success, presented either as fledging success (often termed "apparent success",

an estimate of the percentage of nests that successfully fledge young) or as daily

survival rate (the probability that an egg or nestling survives from one day to the

next); (4) Latitude and longitude; (5) Elevation (meters above sea level, extracted

using the Google Maps Platform); and (6) Clutch size, incubation and nestling

periods, extracted from the paper itself or from Handbook of the Birds of the World

Alive (del Hoyo *et al.* 2018).

Our final dataset included nest predation data for 515 unique species-site

combinations (from 244 studies and representing 314 species). The bulk of this

dataset comes from studies conducted between ~40° South and ~50° North. The

majority of studies come from the Northern Hemisphere temperate zone (269 unique

species-site combinations), but the tropics (187 unique species-site combinations) and

Southern Hemisphere temperate zone (59 unique species-site combinations) are also

well represented (Fig 1). This dataset includes 369 estimates of fledging success and

267 estimates of daily survival rate (see Fig S1).

Quantifying nest predation vs. nest failure

Here we are interested in daily rates of nest predation, but researchers typically report

daily rates of nest survival and do not parse out causes causes of nest failure.

Predators are responsible for the large majority of nest mortality in most places

(Remeš *et al.* 2012), and previous syntheses have simply classified all nest failures as

due to predation (Martin *et al.* 2017). In our dataset, around one quarter of studies (66

out of 244) attempted to identify causes of nest failure. Most nest failure was indeed

attributed to predation (73%; $N = 112$ unique species-site combinations), a figure

remarkably similar to that previously reported for Australian birds (72%; Remeš *et al.*

2012). Further, the percentage of nest failures due to predation was similar across

latitudes (slope estimate for latitude in a linear model predicting the percentage of

nest failure attributed to predation = -0.00083 ± 0.00074, $p = 0.27$). Hence, analyzing

a dataset of studies that report nest success appears to be appropriate for our goal of

analyzing latitudinal patterns in nest predation.

Latitudinal variation in nest predation

We first assessed whether daily rates of nest survival vary as a function of latitude by

fitting mixed effect meta-analytic models using the "metafor" package (Viechtbauer

2010) in R (R Development Core Team 2017). These models weight individual

estimates by the inverse of their squared standard errors, and incorporate the

estimated variance among the study-specific effect sizes. We fit three models that

correspond to different biological hypotheses: (1) no latitudinal gradient in predation

(intercept-only); (2) a linear latitudinal gradient in predation (single slope term, equal

slopes in both Northern and Southern hemispheres); and (3) a breakpoint linear model

wherein predation differs categorically between the tropics and temperate zone (zero

slope within the tropics, ≤ 23.4 degrees absolute latitude, equal slopes for temperate

latitudes in both Northern and Southern hemispheres). We then compared model fits

using AIC. After determining the best-fit model, we fit an expanded version of the

best-fit model to test the influence of non-independence among effect sizes. This

model included species as a random effect and incorporated phylogeny into this

model by specifying phylogenetic branch length as the variance-covariance matrix.

Branch lengths were measured from a majority rules consensus tree calculated from

1000 phylogenies pruned to our study taxa and downloaded from birdtree.org

("Hackett" backbone; Jetz *et al.* 2012).

Latitudinal variation in nesting period duration

Despite longstanding support for the idea that nest predation is more intense in the

tropics, we found minimal variation in daily survival rates with latitude (see Results).

To explore this surprising result, we examined latitudinal patterns in nesting period

duration. We calculated nesting period as the number of days from the laying of the

first egg until fledging, assuming that species begin incubating upon laying the final

egg. We first plotted species' nesting period durations vs. latitude (N = 292 species).

Upon observing clear latitudinal patterns in nesting period duration, with longer

nesting periods in the tropics (see Results), we tested the evolutionary association

between latitude and nesting period duration by fitting a phylogenetic generalized

least squares (PGLS) regression using the "ape" package (Paradis *et al.* 2004). The

response variable in this model was nesting period duration, absolute value of latitude

was a fixed effect, Pagel's λ was estimated using maximum likelihood, and we used .

a majority rules consensus tree from 1000 phylogenies downloaded from birdtree.org

("Hackett" backbone; Jetz *et al.* 2012). In both reality and our dataset, oscine

passerines predominate in the temperate zone while suboscines are more common in

the tropics. To explore whether our overall results are mirrored within these groups,

we repeated the PGLS for oscines only (*N* = 175 species) and for suboscines only (*N*

= 86 species).

To explore the relationship between nesting period duration and daily survival

rates, we fit a meta-analytic model to predict daily survival rate using the "metafor"

package. We first fit a model that estimated different slopes and intercepts for

different latitudinal zones (predictor variables = nesting period duration, latitudinal

zone [north temperate, tropical, south temperate], and an interaction between nesting

period duration and latitudinal zone). We next fit a model where latitudinal zones had

different intercepts but the same slope (i.e., without the interaction term between

nesting period duration and latitudinal zone), and compared fit of the "different

slopes" and "same slopes" models using the "anova" function. Last, we fit an

expanded version of the "same slopes" model that included species as a random effect

and incorporated phylogeny by specifying phylogenetic branch length as the variance-

covariance matrix. Again, branch lengths were measured from a majority rules

consensus tree calculated from 1000 phylogenies pruned to our study taxa and

downloaded from birdtree.org ("Hackett" backbone; Jetz *et al.* 2012).

Latitudinal variation in fledging success

We examined latitudinal patterns in fledging success by repeating the three meta-

analytic models described above for latitudinal patterns in daily survival rate, but with

fledging success as the response variable. Upon plotting the data for the best-fit model

(breakpoint regression with equal slopes in Northern and Southern temperate zones),

we noticed that this model provided a poor fit to data from the Southern temperate

zone. We therefore fit an additional breakpoint regression model where we allowed

slopes to differ between Northern and Southern temperate zones, and compared

model fits using AIC. Last, we fit an expanded version of the best-fit model that

included species as a random effect and specified phylogenetic branch lengths as the

variance-covariance matrix (calculated as previously described).

Results

Latitudinal variation in nest predation

We found evidence that daily rates of nest predation in land birds are similar across

latitudes within the Western Hemisphere. The most supported model of daily nest

survival fit a line with equal daily survival rate (0.96) across the entire latitudinal

gradient (Fig 2, Table 1). Alternative models that fit symmetric, non-zero slopes to

the relationship between latitude and daily survival rates were less well supported

(ΔAIC ~ 7; Table 1). These less supported models were similar to the best-fit model in

that they estimated slopes that were nearly flat (Table S1), with estimated daily nest

survival at 60° North only slightly greater (by ~ 0.006) than at the equator. Our results

did not change when including phylogenetic relationships and species identity in our

model (Table S2), indicating that our findings are robust to sources of non-

independence in our data.

Thus, unlike in tests that have used standardized models to measure predation

in other systems (e.g., Mckinnon *et al.* 2010; Roslin *et al.* 2017; Hargreaves *et al.*

2019), we conclude that natural rates of nest predation experienced by land birds in

the Americas are similar across latitudes. A possible explanation for this contrast

between real nests and experimental nests is that adaptation to contrasting predation

regimes has equalized observed rates in wild birds across latitudes.

Latitudinal variation in nesting period duration

Birds exhibit many adaptations to deter nest predators. We investigated the role of

one specific trait that varies considerably between the tropics and the North temperate

zone—the length of the nesting period (Fig 3). Eggs and chicks remain longer in the

nest in the tropics compared to the North temperate zone (Martin 2002; Chalfoun &

Martin 2007); in our dataset, tropical nesting periods average approximately 15%

longer than in the North temperate zone (~ 32 days vs ~ 28 days; Fig 3). This

difference reflects repeated patterns of evolution across a diversity of avian

lineages—latitude is negatively related to nesting period in a phylogenetic generalized

least squares regression model ($p < 0.0001$, Table S3), and patterns are similar when

analyzing only oscines (temperate origin, predominate in the temperate zone) or

suboscines (tropical origin, predominate in the tropics; Table S3).

Is nesting period length an adaptation to predation?

We hypothesize that longer nesting periods in the tropics are in part an adaptation to

tropical predation regimes. In this view, longer nesting periods in the tropics lead to

lower daily mortality levels on tropical nests than would be expected if nestling

period length remained unchanged. Supporting this idea, when we compare species of

similar nesting duration, daily survival rates are lower in the tropics than in the

temperate zone. This comparison is simplest when we compare intercepts from the

equal slopes model (North temperate zone vs. tropics $p < 0.0001$; Fig 4, Table S4),

which was marginally better supported than a model that fit different slopes to

different latitudinal zones ($p = 0.092$), and is also present in the different slopes model

(comparing estimates for North temperate zone vs. tropics for the mean nesting period

of 29 days, $p < 0.001$; Table S5, Fig S2). Again, this result is robust to sources of non-

independence in our dataset (Table S6).

Nesting period duration is generally positively related to daily survival rate

(Fig 4). For example, the equal slopes model estimated that a 10-day increase in

nesting period is associated with a 1.9% increase in daily survival rates within each

latitudinal zone (Table S4). The suspected reason is that a longer nesting period is

associated with fewer visits to the nest per day by attending parents, reducing rates of

detection by nest predators that cue on parental activity to locate and depredate the

nest (Matysioková & Remeš 2018).

Latitudinal variation in fledging success

The observation that nesting periods are longer in the tropics, while daily survival

rates are similar across latitudes, implies that fledging success (~ daily survival rate *

number of days in nesting period) is lower in the tropics compared to the temperate

zone. Indeed, we found strong evidence that fledging success of nests is highest in the

Northern temperate zone (Fig 5, Tables 2), and a breakpoint regression estimated

~33% of nests successfully fledge nestlings within the tropics while ~ 62% of nests

successfully fledge nestlings at 60° (Table S7). Plotting this data revealed that this

breakpoint regression model poorly fit data from the Southern temperate zone.

Indeed, a breakpoint regression model that fit different slopes to the Southern and

Northern temperate zones was better supported (Table 2); this model estimated that

~34% of nests successfully fledge nestlings within both the tropics *and* the Southern

temperate zone, while ~ 66% of nests successfully fledge nestlings at 60° N, a result

that is robust to sources of non-independence in our data (Table S8). Fledging success

in the tropics vs. the Northern temperate zone is based on mortality accumulated over

different time spans. Hence, our fledging success results are not a demonstration of

higher overall predation in the tropics vs. the Northern temperate zone.

Discussion

We report that natural rates of nest predation experienced by land birds are similar

along a latitudinal gradient in the Americas stretching from ~ 40° South to ~ 60°

North. This result is surprising. Nest predation has long been known to be intense in

the tropics (Skutch 1985) such that nest predation is an archetypical example

illustrating the more general pattern that biotic interactions are stronger at low

latitudes (Schemske 2009). Moreover, recent well-designed globally distributed

experiments have shown that predation rates on standardized models (clay

caterpillars, sunflower seeds) are indeed higher in the tropics (Roslin *et al.* 2017;

Hargreaves *et al.* 2019). Why do clay caterpillars and sunflower seeds get gobbled up

rapidly in the tropics, while tropical birds appear to suffer nest predation at similar

rates to temperate zone birds? We propose that adaptation may explain this

discrepancy, and hypothesize that adaptation to local predation regimes may generate

relatively flat latitudinal gradients in predation experienced by wild populations.

The biotic interactions hypothesis proposes that strong biotic interactions in

the tropics generate strong selection that results in faster evolution in the tropics. Here

we argue that strong biotic interactions in the tropics do not invariably lead to faster

rates of evolution (and ultimately speciation) in the tropics. Instead, strong selection

may lead to adaptations that reduce experienced interaction strength (for the link

between interaction strength and strength of selection see Benkman 2013). For

example, caterpillars living within tropical forests may respond to high rates of

predation by adapting anti-predator traits that reduce the rates at which they are eaten

in nature. It seems noteworthy to us that caterpillars in tropical forests—at least those

noticed by an entomologically-naïve ornithologist— tend not to be green inchworms

but instead are often covered with spikes, brightly colored and presumably poisonous,

or exquisitely camouflaged (BGF pers. obs). The idea that strong negative interactions

drives adaptations that have consequences for the interaction is pervasive in

evolutionary ecology, but has seldom been considered in the context of latitudinal

gradients in interaction strength (but see Rasmann & Agrawal 2011; Chen *et al.*

2017).

Here we hypothesize that tropical birds have adapted to the greater diversity of

nest predators in the tropics, and that these adaptations in part explain the observation

that the latitudinal gradient in daily rates of nest survival is relatively flat.

Specifically, we hypothesize that one component of tropical bird adaptation against

the diversity of tropical nest predators is the evolution of longer nesting periods.

Supporting this hypothesis, when we compare species with similar nesting periods,

rates of nest predation are higher on average in the tropics than in the temperate zone.

However, nesting periods are not equivalent between the tropics and temperate zone.

Instead, tropical birds have repeatedly evolved nesting periods that are longer than

those of their relatives that live in the temperate zone. Thus, we hypothesize that

tropical birds have achieved higher daily survival rates in part by evolving longer

nesting periods (i.e., moving to the right and upwards along the regression line for

daily survival vs. nesting period in Fig 4). We note that this link between daily

survival rate and nesting period duration is based on observational data, so

unmeasured correlated traits (e.g., adult survival; Martin 2002) may also be involved.

Further, many factors likely exert selection on nesting period duration, and the

evolutionary association we document between tropical birds and long nesting periods

could reflect a combination of tropical birds evolving prolonged nesting periods (as

described above) and temperate zone birds evolving shorter nesting periods (e.g., if

shorter breeding seasons in the temperate zone select for faster nesting periods).

High predation risk could potentially drive tropical populations to evolve

longer nesting periods via reductions in rates of parental care that lead to longer

nesting periods. Previous studies provide support for these correlations. When

confronted with high predation risk, there is a general pattern for birds to reduce their

rates of activity at the nest (nest attentiveness), and this correlation appears to be

especially strong in tropical environments (Matysioková & Remeš 2018). In addition,

less parental care per day at the nest is associated with longer nesting periods

(Chalfoun & Martin 2007; Martin et al. 2007) but see (Tieleman et al. 2004). Future

studies that include data on nest attendance (and other relevant breeding biology and

life history traits) are needed to more directly test these relationships.

Comparison with previous studies of latitudinal gradients in nest predation

Tropical birds have long been thought to suffer greater nest predation than temperate

zone birds (e.g., Robinson *et al.* 2000; Schemske *et al.* 2009; Kubelka *et al.* 2018),

but not all previous studies have supported this idea (e.g., Martin *et al.* 2017). Here

we show that one important factor generating disparate results is the choice of metric

of nest predation, as in temperate-tropical comparisons, daily rates vs. fledging

success are measured over non-equivalent time periods. Our finding that fledging

success is much lower in the tropics compared to the North temperate zone is in

agreement with previous analyses of fledging success (Robinson *et al.* 2000;

Schemske *et al.* 2009). However, this pattern is deceptive because daily mortality

rates are in fact similar across latitudes. We suggest that lower fledging success in the

tropics results mainly because tropical birds accumulate mortality over a greater

number of days in the tropics than in the temperate zone. In addition, we find that

fledging success in temperate South America is similar to that within the tropics (and

that nesting periods are similarly long in the tropics and Southern temperate zone),

consistent with previous global studies highlighting that temperate-tropical

comparisons in bird breeding biology may differ between Northern and Southern

Hemispheres (Martin *et al.* 2007, 2017). However, this finding should be treated as

provisional given the small amount of data from latitudes > 35° S in our analysis.

The degree to which our findings can be extrapolated to other regions remains

uncertain. A previous global study found little latitudinal variation in daily rate of nest

predation (Martin *et al.* 2017), consistent with our results. In contrast, rates of nest

predation are indeed higher in the tropics in Australia (Remeš *et al.* 2012; this study

also found much greater longitudinal variation in nest predation than latitudinal

variation). Here we show that for the Western Hemisphere, the long-held view that

nest predation is higher in the tropics is mainly derived from the reduced fledging

success of tropical birds (Skutch 1985; Schemske 2009), which is more a

consequence of a longer nesting period than a lower daily survival rate. Since they are

measured over different time periods on average, fledging success rates in the tropics

vs. the Northern temperate zone do not adequately measure overall survival rates in

these two latitudinal zones.

Conclusions

The biotic interactions hypothesis invokes strong biotic interactions in the tropics to

explain in part the origin of high tropical species richness. This intriguing proposal

has spawned intense interest in documenting latitudinal clines in interaction strength.

We add to this literature by showing evidence that a latitudinal gradient in rates of

nest predation suffered by land birds in the Americas is weak or absent. Yet

documenting gradients in interaction strength is only part of the story. The idea that

species can adapt to strong interactions, potentially blunting their effects, has been

comparatively ignored. Here we present evidence that the lack of a latitudinal gradient

in nest predation suffered by birds may be in part a result of adaptation on the part of

tropical birds. We suggest this is an example of how an adaptation to strong negative

biotic interactions can flatten the observed latitudinal gradient in interaction strength.

More broadly, we advocate for an increased focus on the evolutionary consequences

of biotic interactions.

Acknowledgements

This research was supported by postdoctoral fellowships from the Biodiversity

Research Centre and Banting Canada (#379958) to BGF. None of our funders had

any influence on the content of the submitted manuscript, and none of our funders

required approval of the final manuscript to be published. Comments from the

Schluter lab group and Ralf Yorque greatly improved this manuscript.

References

Benkman, C.W. (2013). Biotic interaction strength and the intensity of selection.

Ecol. Lett., 16, 1054–1060.

Boyle, A.W., Sandercock, B.K. & Martin, K. (2016). Patterns and drivers of

intraspecific variation in avian life history along elevational gradients: A meta-

analysis. *Biol. Rev.*, 91, 469–482.

Burke, D.M., Elliott, K., Moore, L., Dunford, W., Nol, E., Phillips, J., *et al.* (2004).

Patterns of Nest Predation on Artificial and Natural Nests in Forests. *Conserv.*

Biol., 18, 381–388.

Chalfoun, A.D. & Martin, T.E. (2007). Latitudinal variation in avian incubation

attentiveness and a test of the food limitation hypothesis. *Anim. Behav.*, 73, 579–

585.

Chen, S.C., Hemmings, F.A., Chen, F. & Moles, A.T. (2017). Plants do not suffer

greater losses to seed predation towards the tropics. *Glob. Ecol. Biogeogr.*, 26,

1283–1291.

Conway, C.J. & Martin, T.E. (2000). Evolution of Passerine Incubation Behavior：

Influence of Food, Temperature, and Nest Predation. *Evolution*, 54, 670–685.

Darwin, C. (1859). *On the origin of species by means of natural selection*. J. Murray,

London.

DeGregorio, B.A., Chiavacci, S.J., Benson, T.J., Sperry, J.H. & Weatherhead, P.J.

(2016). Nest Predators of North American Birds: Continental Patterns and

Implications. *Bioscience*, 66, 655–665.

Dobzhansky, T. (1950). Evolution in the tropics. *Am. Sci.*, 38, 209–221.

Hargreaves, A.L., Suárez, E., Mehltreter, K., Myers-smith, I., Vanderplank, S.E.,

Slinn, H.L., *et al.* (2019). Seed predation increases from the Arctic to the Equator

and from high to low elevations. *Sci. Adv.*, 5, eaau4403.

del Hoyo, J., Elliott, A., Sargatal, J., Christie, D. & E, de J. (2018). *Handbook of the*

Birds of the World Alive.

Jetz, W., Thomas, G.H., Joy, J.B., Hartmann, K. & Mooers, A.O. (2012). The global

diversity of birds in space and time. *Nature*, 491, 444–448.

King, D.I., DeGraaf, R.M., Griffin, C.R. & Maier, T.J. (1999). Do Predation Rates on

Artificial Nests Accurately Reflect Predation Rates on Natural Bird Nests? *J. F.*

Ornithol., 70, 257–262.

Kubelka, V., Šálek, M., Tomkovich, P., Végvári, Z., Freckleton, R. & Székely, T.

(2018). Global pattern of nest predation is disrupted by climate change in

shorebirds. *Science*, 362, 680–683.

Kulesza, G. (1990). An analysis of clutch size in New World passerine birds. *Ibis*,

132, 407–422.

MacArthur, R.H. (1972). *Geographical ecology: patterns in the distribution of*

species. Princeton University Press, Princeton.

Martin, T.E. (2002). A new view of avian life-history evolution tested on an

incubation paradox. *Proc. R. Soc. B Biol. Sci.*, 269, 309–316.

Martin, T.E., Auer, S.K., Bassar, R.D., Niklison, A.M. & Lloyd, P. (2007).

Geographic variation in avian incubation periods and parental influences on

embryonic temperature. *Evolution*, 61, 2558–2569.

Martin, T.E., Boyce, A.J., Fierro-Calderón, K., Mitchell, A.E., Armstad, C.E.,

Mouton, J.C., *et al.* (2017). Enclosed nests may provide greater thermal than nest

predation benefits compared with open nests across latitudes. *Funct. Ecol.*, 31,

1231–1240.

Matysioková, B. & Remeš, V. (2018). Evolution of parental activity at the nest is

shaped by the risk of nest predation and ambient temperature across bird species.

Evolution, 72, 2214–2224.

Mckinnon, L., Smith, P.A., Nol, E., Martin, J.L., Doyle, F.I., Abraham, K.F., *et al.*

(2010). Lower Predation Risk for Migratory birds. *Science*, 327, 6–8.

Menezes, J.C.T. & Marini, M.Â. (2017). Predators of bird nests in the Neotropics: a

review. *J. F. Ornithol.*, 88, 99–114.

Moles, A.T. & Ollerton, J. (2016). Is the notion that species interactions are stronger

and more specialized in the tropics a zombie idea? *Biotropica*, 48, 141–145.

Moore, R.P. & Robinson, W.D. (2004). Artificial Bird Nests, External Validity, and

Bias in Ecological Field Studies. *Ecology*, 85, 1562–1567.

Nice, M. (1957). Nesting Success in Altricial Birds. *Auk*, 74, 305–321.

Paradis, E., Claude, J. & Strimmer, K. (2004). APE: analyses of phylogenetics and

evolution in R language. *Bioinformatics*, 20, 289–290.

Peterson, A. & Young, H. (1950). A nesting study of the Bronzed Grackle. *Auk*, 67,

466–476.

R Development Core Team. (2017). R: A language and environment for statistical

computing.

Rasmann, S. & Agrawal, A.A. (2011). Latitudinal patterns in plant defense: evolution

of cardenolides, their toxicity and induction following herbivory. *Ecol. Lett.*, 14,

476-483.

Remeš, V., Matysioková, B. & Cockburn, A. (2012). Long-term and large-scale

analyses of nest predation patterns in Australian songbirds and a global

comparison of nest predation rates. *J. Avian Biol.*, 43, 435–444.

Robinson, W.D., Robinson, T.R., Robinson, S.K. & Brawn, J.D. (2000). Nesting

Success of Understory Forest Birds in Central Panama. *J. Avian Biol.*, 31, 151–164.

Robinson, W.D., Styrsky, J.N. & Brawn, J.D. (2005). Are artificial bird nests effective surrogates for estimating predation on real bird nests? A test with tropical birds. *Auk*, 122, 843–852.

Roslin, T., Hardwick, B., Novotny, V., Petry, W.K., Andrew, N.R., Asmus, A., *et al.* (2017). Higher predation risk for insect prey at low latitudes and elevations. *Science*, 356.

Schemske, D. (2009). Biotic interactions and speciation in the tropics. *Speciation and patterns of diversity*, pages 219–240.

Schemske, D.W., Mittelbach, G.G., Cornell, H. V, Sobel, J.M. & Roy, K. (2009). Is there a latitudinal gradient in the importance of biotic interactions? *Annu. Rev. Ecol. Evol. Syst.*, 40, 245–269.

Skutch, A.F. (1985). Clutch size, nesting success, and predation on nests of Neotropical birds, reviewed. *Ornithol. Monogr.*, 36, 575–594.

Tieleman, B.I., Williams, J.B. & Ricklefs, R.E. (2004). Nest attentiveness and egg temperature do not explain the variation in incubation periods in tropical birds. *Funct. Ecol.*, 18, 571–577.

Viechtbauer, W. (2010). Conducting meta-analyses in R with the metafor package. *J. Stat. Softw.*, 36, 1–48.

Wallace, A.R. (1869). *The Malay Archipelago*.

523   Figure 1. Map of studies measuring nest predation experienced by land birds in the

524   Americas (dataset = 244 studies). Many studies report data for multiple species from

525   the same site, illustrated by the size of the circle. The Tropics of Cancer and

526   Capricorn (at 23.4° N and S, respectively) delimit the tropics, and are illustrated with

527   dashed lines.

Figure 2. The latitudinal gradient in daily survival rates for land birds in the Americas

(N = 267). Predictions from the best-fit metafor model are plotted as a dashed line—

this simple model fit a constant value for daily survival rates across latitudes. For

comparison, the loess fit, which used the same weights for each data point as the best-

fit model, and incorporates the same estimated variance among the study-specific

effect sizes, is plotted in blue with shaded 95% confidence intervals.

Figure 3. The latitudinal gradient in nesting period length for land birds in the

Americas (N = 292 species). The dashed line illustrates the loess trendline, with 95%

confidence intervals shaded in gray. Nesting periods average ~ 32 days within the

tropics and Southern temperate zone but ~28 days within the Northern temperate

zone.

Figure 4. The relationship between daily survival rate and nesting period duration

among latitudinal zones. Daily survival rate increases with nesting period length in

each latitudinal zone, but with different intercepts. Predictions from a metafor model

are plotted as solid lines; values from individual studies are plotted in colors

corresponding to their latitudinal zone (N = 252).

Figure 5. The latitudinal gradient in fledging success for land birds in the Americas

(N = 367). Predictions from the best-fit model in metafor—a breakpoint regression

that fit a flat line within the tropics and different slopes in the Northern and Southern

Hemisphere temperate zones—, are plotted as a dashed line. For comparison, the

loess fit, which used the same weights for each data point as the best-fit model, and

incorporated the same estimated variance among the study-specific effect sizes, is

plotted in blue with shaded 95% confidence intervals.

561   Table 1. Model comparison of metafor models with daily survival rate as the response

562   variable. The best-fit model fit a constant value of daily survival rate across the entire

563   latitudinal expanse of the dataset.

Model	ΔAIC
Constant (daily survival rates do not vary with latitude)	--
Linear regression	6.89
Breakpoint regression (breakpoint = 23.4°)	7.54

566   Table 2. Model comparison of metafor models with fledging success as the response

567   variable. The best-fit model was a breakpoint regression that fit one slope within the

568   tropics and unequal slopes in the Northern and Southern temperate zones.

Model	ΔAIC
Breakpoint regression, unequal slopes (breakpoint = 23.4°)	--
Breakpoint regression, equal slopes (breakpoint = 23.4°)	4.68
Linear regression	21.29
Constant (fledging success does not vary with latitude)	66.35

Figure S1. Map of studies measuring daily rates of nest survival for land birds in the

Americas (dataset = 110 studies). Many studies report data for multiple species from

the same site, illustrated by the size of the circle. The Tropics of Cancer and

Capricorn (at 23.4° N and S, respectively) delimit the tropics, and are illustrated with

dashed lines.

Figure S2. The relationship between daily survival rate and nesting period duration

among latitudinal zones (different slopes model). Daily survival rate increases with

nesting period length, but with different intercepts and slopes for different latitudinal

zones. Predictions from a metafor model are plotted as solid lines; values from

individual studies are plotted in colors corresponding to their latitudinal zone (N =

252).

598   Table S1. Parameter estimates for fixed effects from metafor models with daily

599   survival rate as the response variable. Slope estimates for the linear regression are for

600   straight lines outward from the equator towards higher latitudes; slope estimates for

601   breakpoint regression are for straight lines outward from the tropics (23.4 degrees

602   latitude) towards higher latitudes.

Model	Intercept ± SE	Slope ± SE
Intercept-only (constant)	0.96 ± 0.0015	NA
Linear regression	0.96 ± 0.0039	0.0001 ± 0.0001
Breakpoint regression (breakpoint = 23.4°)	0.96 ± 0.0024	-0.0001 ± 0.0002

605   Table S2. Model summary of intercept-only metafor model with daily survival rate as

606   the response variable and species and phylogeny as random effects.

Effect	Estimate
Intercept (fixed)	0.96 (se = 0.0083)
Species (random)	0.0000 (sqrt = 0.0067)
Phylogeny (random)	0.0003 (sqrt = 0.016)

613   Table S3. Parameter estimates for fixed effects from phylogenetic generalized least

614   squares regression models with nesting period as the response variable. Pagel's λ for

615   these model was 0.99 (entire dataset), 1.00 (oscines) and 0.94 (suboscines).

Model	Parameter	Estimate $\pm$ SE
Entire dataset	Intercept	27.29 ± 2.90
	Absolute value (latitude)	-0.040 ± 0.015
Oscines only	Intercept	34.92 ± 2.86
	Absolute value (latitude)	-0.026 ± 0.016
Suboscines only	Intercept	35.45 ± 2.99
	Absolute value (latitude)	-0.065 ± 0.039

619   Table S4. Parameter estimates from a metafor model with daily survival rate as the

620   response variable. This is the "equal slopes" model where different latitudinal zones

621   have the same slope. The reference category is the North temperate zone; parameter

622   estimates for the South temperate zone and tropics represent deviations from this

623   reference category.

Parameter	Estimate $\pm$ SE
Intercept	0.91 ± 0.0084
Nesting period	0.0019 ± 0.0003
South temperate zone	-0.0049 ± 0.0047
Tropics	-0.015 ± 0.0037

627   Table S5. Parameter estimates from a metafor model with daily survival rate as the

628   response variable. This is the "different slopes" model where different latitudinal

629   zones have different slopes. The reference category is the North temperate zone;

630   parameter estimates for "Nesting Period: South temperate zone" and "Nesting Period:

631   Tropics" represent deviations from this reference category.

Parameter	Estimate ± SE
Intercept	0.89 ± 0.012
Nesting period	0.0025 ± 0.00043
South temperate zone	0.038 ± 0.035
Tropics	0.024 ± 0.019
Nesting period: South temperate zone	-0.0015 ± 0.0012
Nesting period: Tropics	-0.0013 ± 0.00063

Table S6. Parameter estimates from a metafor model with daily survival rate as the

response variable and study, species and phylogeny as random effects, based on data

from studies that reported daily survival rates. This is the "equal slopes" model where

different latitudinal zones have the same slope. The reference category for the

intercept is the North temperate zone; parameter estimates for the South temperate

zone and tropics represent deviations from this reference category.

Parameter	Estimate ± SE
Intercept	0.91 ± 0.0084
Nesting period	0.0019 ± 0.00030
South temperate zone	-0.0050 ± 0.0047
Tropics	-0.015 ± 0.0037
Species (random)	0.0 (sqrt = 0.0064)
Phylogeny (random)	0.0 (sqrt = 0.0)

Table S7. Parameter estimates for fixed effects from metafor models with fledging

success as the response variable. Slope estimates for the linear regression are for

straight lines outward from the equator towards higher latitudes; slope estimates for

breakpoint regression are for straight lines outward from the tropics (23.4 degrees

latitude) towards higher latitudes. Breakpoint regressions fit either equal slopes (same

slope value in both Northern and Southern temperate zones) or unequal slopes

(different slopes in Northern vs. Southern temperate zones).

Model	Intercept ± SE	Slope ± SE
Intercept-only (constant)	0.41 ± 0.010	NA
Linear regression	0.29 ± 0.019	0.0043 ± 0.0006
Breakpoint regression, equal slopes (breakpoint = 23.4°)	0.33 ± 0.013	0.0080 ± 0.001
Breakpoint regression, unequal slopes (breakpoint = 23.4°)	0.34 ± 0.013	-0.0008 ± 0.0032 (S Hemisphere) 0.0089 ± 0.0031 (N Hemisphere)

681   Table S8. Model summary of unequal slopes breakpoint regression metafor model

682   with fledging success as the response variable, and species and phylogeny included as

683   random effects.

Effect	Estimate
Intercept (fixed)	0.34 (se = 0.013)
Slope (fixed)	-0.0008 ± 0.0032 (S Hemisphere)
	0.0089 ± 0.0031 (N Hemisphere)
Species (random)	0.026 (sqrt = 0.16)
Phylogeny (random)	0 (sqrt = 0.0000)