Single AAV-mediated scarless genome editing in dysfunctional retinal neurons

mediates robust visual restoration in mice

Short title: Single AAV scarless genome editing

Koji M Nishiguchi,[1,2][†][*] Kosuke Fujita,[3][†] Shota Katayama,[1] Toru Nakazawa[1,2,3*]

[1] Department of Advanced Ophthalmic Medicine, Tohoku University Graduate School of

Medicine, Sendai 980-8574, Japan

[2] Department of Ophthalmology, Tohoku University Graduate School of Medicine, Sendai

980-8574, Japan

[3] Department of Ophthalmic Imaging and Information Analytics, Tohoku University Graduate

School of Medicine, Sendai 980-8574, Japan

[†] These authors contributed equally to the work.

[*] Corresponding authors. Tel: +81-22-717-7294

Koji M Nishiguchi, E-mail: nishiguchi@oph.med.tohoku.ac.jp

Toru Nakazawa, E-mail: ntoru@oph.med.tohoku.ac.jp

The most versatile treatment for inherited disorders is to precisely replace a

mutated sequence with its wildtype counterpart, thereby _"normalizing"_ the genome.

We developed a single AAV platform that allows this in retinal neurons with

combined CRISPR-Cas9 and micro-homology-mediated end-joining. In blind mice,

the platform rescued ~10% of the retinal neurons, resulting in an incredible

~10,000-fold improvement in light sensitivity, equivalent to the restoration

mediated by conventional gene augmentation therapy.

Gene supplementation of wildtype copies of functionally defective genes (gene

augmentation) mediated by adeno associated virus (AAV) have shown remarkable success

in clinical trials for inherited neural disorders. [1,2] However, its application is restricted to

limited genes within the capacity (~5,000 bps) of AAV.[3] CRISPR-Cas9-mediated allele

knock-out based on error-prone non-homologous end joining (NHEJ) can correct gain-of-

function mutations with AAV.[4-7] However, many loss-of-function mutations in larger genes

still require local replacement with a wildtype sequence (scarless genome editing).

Here, we used micro-homology-mediated end-joining (MMEJ) to integrate a donor

template with micro-homology arms of 20 bp[8] as the basis for a new single-AAV platform

for scarless genome editing. First, we generated mutants of the preexisting retina-specific

promoters and conducted *in vivo* AAV reporter assays (Fig. S1a-c, Table S1). The smallest

promoter that maintained neural retina-specific transcription was a 93-bp mutant GRK1

promoter that allowed reporter expression in 65.5% of the photoreceptors (Fig. S1b-e). This

was used to drive SaCas9 (3.2 Kb) expression.

We tested our single-AAV vector platform in $Gnat1^{IRD2/IRD2}$/$Pde6c^{cpfl1/cpfl1}$ mice

(double-mutant mice); the *Gnat1* and *Pde6c* defects caused functional lack of rod and cone

photoreceptors,[9] respectively. The mice were essentially blind from birth and thus allowed

clear delineation of therapeutic effects.[10] We used our platform to correct *IRD2* mutations in

Gnat1; these mutations constitute a homozygous 59-bp deletion in intron 4, preventing

protein expression in the rod photoreceptors[11], which comprise ~75% of the murine retinal

cells.[12] Six guide RNAs (gRNAs) designed flanking the mutation were assessed with a T7

Endonuclease 1 (T7E1) assay (Fig. S2). A gRNA pair (1+4) that excised the mutation most

efficiently was selected. The constructed single-AAV vector (Fig. 1a) was then injected sub-

retinally in 6-month-old double-mutant mice. Five weeks later, histology showed scattered

GNAT1-positive photoreceptors, indicating successful genome editing (Fig. 1b). T7E1

assay (Fig. 1c) and RT-PCR (Fig. 1d) of the whole retina revealed absolute editing

efficacies of 8.6% and 12.7% among the rod photoceptors, respectively. Furthermore,

electroretinogram (ERG), which reflects the number of functional photoreceptors, revealed

6-Hz flicker responses, averaging 11.4% of the control mice (Fig. 1e). These results are

consistent with a ~10% rescue of the rod photoreceptors by genome editing.

Next, light sensitivity in the visual cortex was assessed with flash visually evoked

potentials (fVEPs). Surprisingly, cortical responses contralateral to the treated eye revealed

a ~10,000-fold (range: 1,000 – 100,000-fold) improvement in light sensitivity, equivalent to

gene augmentation in ~70% of the photoreceptors (Fig. 1f and g, S1e, S3).[10] This was

reflected by changes in light-induced behavior (fear conditioning, Fig. 1h). Furthermore,

contralateral cortical responses to phase-reversal gratings, i.e., the pattern VEP (pVEP),

were detected in higher resolution gratings with larger amplitudes following the treatment,

indicating improved visual acuity (Fig. 1i). Threshold assessment by measuring optokinetic

responses to rotating gratings showed visual acuity restoration in the treated mice to 59.1%

of the control mice. (Fig. 1j).

On-target analysis of retinal genomic DNA with sequencing revealed an absolute

indel rate of 36.4% (Table S3), somewhat higher than previous reports[13,14]. Three sites

predicted to have the highest off-target probabilities for each gRNA showed no detectable

mutation events in the retinal genomic DNA (Fig. S4) by the T7E1 assay. Further

sequencing of these sites found no mutations (Fig. S4).

We also used our AAV vector to treat 2-month-old *Gnat1[IRD2/IRD2]* mice, which retain

cone function and serve as a model of human retinal dystrophy.[15] In the early course of the

disease, patients suffer from severe loss of light sensitivity with preserved visual acuity. [15] A

histological analysis showed scattered GNAT1-postive photoreceptors in the treated mice

(Fig. 2a). T7E1 assay and RT-PCR measured absolute genome editing efficacy at 7.5%

and 7.2%, respectively (Fig. S5a,b). fVEP showed a ~1,000-fold increase in light sensitivity

(Fig. 2b). This was confirmed behaviorally with fear conditioning (Fig. 2c). However, the

improvement in retinal function could not be isolated from preexisting cone function with

ERG and visual acuity remained unchanged with pVEP, or OKR (Fig. S5c-e). These results

show that our platform had therapeutic effects also in a human disease model.

This study shows that scarless genome editing allows a striking improvement in

light sensitivity, the major biomarker adopted in the seminal clinical trials for retinal

dystrophy.[2] Moreover, the effect was equivalent to that of conventional gene augmentation

therapy[10], a platform that is already in clinical use,[16] and was achieved with a single AAV

vector. This is a major step forward from the dual vector system used in previous studies on

donor-based *in vivo* genome editing in post-mitotic cells, which generally yielded an

absolute editing efficiency of less than ~5%[13,14,17,18], compared to ~10% shown herein.

Additional important advantages of scarless genome editing include physiological and

uniform expression of the target gene, neither of which is possible with gene augmentation.

This may be critically important to avoid dose-related toxicity caused by both gene over-

expression[19] and under-expression[20], especially during long-term treatment in humans.

As a proof of concept, our platform for single AAV-mediated scarless genome

editing demonstrates amelioration of key features of dysfunctional neurons. This paves the

way to treat inherited neural disorders caused by loss-of-function mutations in larger genes,

for which conventional gene augmentation therapy or NHEJ-based genome editing

strategies are not currently feasible.

Reference

1.   Mendell, J.R.*, et al. N. Engl. J. Med.* **377**, 1713-1722 (2017).

2.   Cideciyan, A.V. *Prog. Retin. Eye Res.* **29**, 398-427 (2010).

3.   Wu, Z., Yang, H. & Colosi, P. *Mol. Ther.* **18**, 80-86 (2010).

4.   Tsai, Y.T.*, et al. Ophthalmology* **125**, 1421-1430 (2018).

5.   Yang, S.*, et al. J. Clin. Invest.* **127**, 2719-2724 (2017).

6.   Gaj, T.*, et al. Science advances* **3**, eaar3952 (2017).

7.   Li, P.*, et al. The CRISPR Journal* **1**, 55-64 (2018).

8.   Nakade, S.*, et al. Nature communications* **5**, 5560 (2014).

9.   Lamb, T.D. *Prog. Retin. Eye Res.* **36**, 52-119 (2013).

10.   Nishiguchi, K.M.*, et al. Mol. Ther.* **26**, 2397-2406 (2018).

11.   Miyamoto, M.*, et al. Exp. Eye Res.* **90**, 63-69 (2010).

12.   Jeon, C.J., Strettoi, E. & Masland, R.H. *J. Neurosci.* **18**, 8936-8946 (1998).

13.   Yin, H.*, et al. Nat. Biotechnol.* **34**, 328 (2016).

14.   Yang, Y.*, et al. Nat. Biotechnol.* **34**, 334 (2016).

15.   Carrigan, M.*, et al. Br. J. Ophthalmol.* **100**, 495-500 (2016).

16.   Kaltenboeck, A. & Bach, P.B. *JAMA* (2018).

17.   Suzuki, K.*, et al. Nature* **540**, 144-149 (2016).

18.   Yin, H.*, et al. Nat. Biotechnol.* **32**, 551 (2014).

19.   Tan, E.*, et al. Invest. Ophthalmol. Vis. Sci.* **42**, 589-600 (2001).

20.   Verbakel, SK., *et al. Invest. Ophthalmol. Vis. Sci.* (2019) *in press*.

Fig 1. *In vivo* scarless genome editing and visual restoration in blind mice

124 **a.** Design of the all-in-one vector with an enlarged map of the donor sequence. ITR,

125 inverted terminal repeat; NLS, nuclear localizing signal; pA, ploy A; PAM, protospacer

126 adjacent motif. **b.** GNAT1-positive photoreceptors (arrowhead) following genome editing in

127 a retinal section (top) and a flat mount (bottom). Scale bar: 20 μm. **c.** T7E1 assay.

128 Representative image (left) and stats (right, N = 4). Arrows: size of cleaved fragments (also

129 indicated below). **d.** RT-PCR (N = 4). **e.** 6-Hz flicker ERGs in treated and untreated eyes of

130 the same $Pde6c^{cpfl1/cpfl1}Gnat1^{IRD2/IRD2}$ mice (N = 6) and control $Pde6c^{cpfl1/cpfl1}$ mice (N = 5). **f.**

131 fVEPs recorded from contralateral visual cortices of treated and untreated eyes of

132 $Pde6c^{cpfl1/cpfl1}Gnat1^{IRD2/IRD2}$ mice (N = 9) and control $Pde6c^{cpfl1/cpfl1}$ mice (N = 5). **g.**

133 Thresholds of fVEP. $Pde6c^{cpfl1/cpfl1}Gnat1^{IRD2/IRD2}$ mice were treated either with genome

134 editing (N = 9) or $GNAT1$ over-expression (OE, N = 6) or left untreated (N =9). **h.** Fear

135 conditioning test. Freezing time before (Baseline) and during (Stimulus) presentation of

136 fear-conditioned light cue (N = 6 each). **i.** pVEPs from treated (N = 9) and untreated (N = 5)

137 $Pde6c^{cpfl1/cpfl1}Gnat1^{IRD2/IRD2}$ mice. **j.** Optokinetic response in the treated (N = 10) and

138 untreated (N = 4) $Pde6c^{cpfl1/cpfl1}Gnat1^{IRD2/IRD2}$ mice and untreated $Pde6c^{cpfl1/cpfl1}$ mice (N =

139 6). Data represent the mean ± S.E.M.; *P < 0.05; Tx, treated; NoTx untreated; OE, over-

140 expression; nd, non-detectable; ns, not significant.

Fig 2. In vivo scarless genome editing in a mouse model of retinal degeneration

a. GNAT1-positive photoreceptors (arrowhead) following treatment of $Gnat1^{IRD2/IRD2}$ mice

shown in a retinal section (top) and a flatmount (bottom). Scale bar: 20 μm. **b.** fVEPs

recorded from contralateral visual cortices in treated and untreated eyes of the same mice.

(N = 7). **c.** Fear conditioning test, showing freezing time before (Baseline) and during

(Stimulus) presentation of fear-conditioned light cue. Treated (Tx, N = 7) and untreated

(NoTx, N = 6) $Gnat1^{IRD2/IRD2}$ mice and CL57B6 mice (B6, N = 6). Data represent the mean ±

S.E.M.; *P < 0.05; ONL, outer nuclear layer. ns, not significant.

Acknowledgements

We thank Ms Misane Uchiike for her help with the experiments. This work was supported in

part by the Japan Agency for Medical Research and Development (KMN,

18ek0109213h0002). The manuscript was edited by a professional English editing service

(Mr. Tim Hilts).

Author contributions

KMN conceived and designed the experiments. KMN and KF performed the experiments and

analyzed the data. SK helped in vitro experiments and gave critical advice. KMN and TN

wrote the manuscript and obtained the funding.

Competing financial interests

KMN, KF, and TN are listed as inventors in a patent application related to this work. The

Departments of Advanced Ophthalmic Medicine and Ophthalmic Imaging and Information

Analytics are endowed departments, supported by an unrestricted grant from Senju

Pharmaceutical Co. (Osaka, Japan) and Topcon Co. Ltd. (Tokyo, Japan), respectively.

These funders had no role in the study design, data collection and analysis, decision to

169   publish, or preparation of the manuscript.

171   **Data availability**

172   The datasets generated during and/or analyzed during the current study are available from

173   the corresponding author on reasonable request.