Drosophila **p38 MAPK Interacts with BAG-3/starvin to Regulate Age-dependent**

Protein Homeostasis

Sarah M. Ryan[2,*], Michael Almassey[1,*], Amelia M. Burch[3], Gia Ngo[2], Scott Barbee[2],

Nathan T. Mortimer[1], Subhabrata Sanyal[3,4] and Alysia D. Vrailas-Mortimer[1-3]

1 Illinois State University, School of Biological Sciences, Normal, IL, USA

2 University of Denver, Department of Biological Sciences, Denver, CO, USA

3 Emory University, Department of Cell Biology, Atlanta, GA, USA

4 Calico, San Francisco, CA, USA

Running Title: p38K Regulates Age-Dependent Proteostasis

Key words/phrases: p38 MAPK, BAG-3/starvin, protein aggregation, aging, Lamin

Corresponding author:

Alysia D. Vrailas-Mortimer

Illinois State University

School of Biological Sciences

Science Laboratory Building

Normal, IL 61790

Phone: 309-348-8574

Fax: 309-348-3722

e-mail: admorti@ilstu.edu

Summary

As organisms age, they often accumulate protein aggregates that are thought to be

toxic, potentially leading to age-related diseases. This accumulation of protein

aggregates is partially attributed to a failure to maintain protein homeostasis. A variety of

genetic factors have been linked to longevity, but how these factors also contribute to

protein homeostasis is not completely understood. In order to understand the

relationship between aging and protein aggregation, we tested how a gene that

regulates lifespan and age-dependent locomotor behaviors, p38 MAPK (p38Kb),

influences protein homeostasis as an organism ages. We find that p38Kb regulates age-

dependent protein aggregation through an interaction with the Chaperone-Assisted

Selective Autophagy complex. Furthermore, we have identified Lamin as an age-

dependent target of p38Kb and the Chaperone-Assisted Selective Autophagy complex.

Introduction

Protein turnover is critical for maintaining tissue health as many proteins become

damaged or misfolded during normal tissue functions. Therefore, the cell utilizes a

variety of protein quality control mechanisms to refold or degrade these damaged

proteins, including the ubiquitin proteasome system and macroautophagy. During aging,

protein quality control mechanisms become less efficient leading to the accumulation of

damaged or misfolded proteins that begin to form protein aggregates[1]. It has been

hypothesized that these aggregates are toxic and may lead to the deleterious

phenotypes associated with normal aging, such as impaired tissue function[1].

Furthermore, decreased protein aggregation has been associated with longevity. For

example, over-expression of Foxo leads to an increased lifespan but also a concordant

decrease in protein aggregation in *C. elegans*, *Drosophila,* and mice [2–6], suggesting that

lifespan and protein aggregation are tightly linked processes. However, the molecular

mechanisms that underlie the relationship between aging and protein homeostasis have

not been fully characterized.

One pathway that has been linked to both aging and protein homeostasis is the

stress response p38 MAPK (p38K) pathway. In mammals, there are four p38K genes (α,

β, γ, and δ), and p38Kα has been linked to both the inhibition [7,8] and induction [9,10] of

macroautophagy, in particular in response to oxidative stress [11,12]. In addition, p38Kα

has been linked to regulating macroautophagy in cellular senescence [13–15]. However,

how p38K signaling may contribute to protein homeostasis in response to natural aging

is not well understood. The fruit fly *Drosophila melanogaster* has two canonical p38K

genes (p38Ka and p38Kb), and we have previously reported that p38Kb acts in the adult

musculature to regulate aging. We found that over-expression of p38Kb leads to

increased lifespan while loss of p38Kb results in a short lifespan and age-dependent

locomotor behavior defects [16]. In addition, oxidatively damaged proteins accumulate in

the muscle of p38Kb mutants with age [16], and loss of p38Kb leads to increased

polyubiquitination of insoluble proteins and alterations in oxidative stress dependent

translation [17], suggesting that these oxidatively damaged proteins may be aggregating in

p38Kb mutants. Furthermore, p38Kb has been shown in a *Drosophila* cell culture system

to pull down with the HspB8 homologue CG14207 [18], which plays a role in the muscle by

regulating protein homeostasis as a part of a protein quality control mechanism called

BAG-3 Mediated Selective Autophagy pathway or the Chaperone-Assisted Selective

Autophagy (CASA) complex in flies [19,20]. The CASA complex also includes the

chaperone Hsc70 and the co-chaperone BAG-3 (starvin in flies). BAG-3/starvin binds to

specific damaged or misfolded protein substrates and brings them to the CASA complex

where it binds to Hsc70 and HspB8. Those substrates that cannot be refolded by the

complex are polyubiquitinated and targeted to the autophagosome through a handover

between BAG-3/stv and the autophagy adaptor protein p62 (ref(2)p in flies), and

subsequently degraded through the autophagosome-lysosome system [21–25].

Here, we report that p38Kb regulates age-dependent muscle protein

homeostasis through an interaction with the CASA complex. We find that p38Kb acts as

an intermediary between BAG-3/starvin and p62/ref(2)p in targeting damaged or

misfolded proteins for degradation. This interaction is not only important for maintaining

protein homeostasis but also for lifespan extension. In addition, we find that the

Drosophila homologue of the Hutchinson-Gilford progeria protein, Lamin A/C, is a target

for p38Kb and CASA complex mediated protein turnover, suggesting that the p38Kb

aging phenotypes may be a result of impaired Lamin degradation.

Materials and Methods

Genotypes

UAS-p38Kb wt, UAS-p38Kb Kinase Dead, p38Kb$^{\Delta 45}$, p38Kb^{Ex41}, w^{1118}, Mef2-GAL4 and

MHC-GAL4 were as described in [16]. The p38Kb^{Ex41} is a precise excision allele and

serves as a genetic background control for p38Kb$^{\Delta 45}$ deletion mutation.

UAS-stv RNAi 34408 (w^{1118}; P{GD10796}v34408) and UAS-stv RNAi 34409 (w^{1118};

P{GD10796}v34409/TM3) are described in [26] and are from the Vienna Drosophila

Resource Center.

stv-GFP trap (w[11118];; Pbac{754.P.F3v30} stv[+CPTI 002824]), HspB8-GFP trap (w[1118]

PBac{810.P.FSVS-2}CG14207[CPTI004445]), Hsc70-4 GFP trap (*w[1118] PBac{544.SVS-*

1}N[CPTI002347]) are from the Kyoto Stock Center.

w[1118]; P{y[+mDint2] w[BR.E.BR]=SUPor-P}ref(2)P[KG00926], w[1118];;+mDint2 EY4969 stvEP,

Lam[A25] pr[1], and w[1118]; P{w[+mC]=UAS-Lam.GFP}3-3 were obtained from the

Bloomington Drosophila Stock Center.

All fly stocks were backcrossed into the w[1118] background and isogenized for 10

generations. All stocks were reared at 25ºC in a 12hr:12hr light:dark cycle on standard

fly food media.

Immunofluorescence

Adult flies were fixed in 4% paraformaldehyde for 48hrs at 4°C. Indirect flight muscles

were dissected in 1X PBS, permeabilized in 1X PBS 0.15% Triton-X 100, and blocked in

NGS + 0.15% Triton-X 100. Samples were incubated in primary antibody at 4°C

overnight, washed in 1X PBS 0.15% Triton-X 100, and incubated in secondary antibody

at room temperature for 2hrs. Samples were mounted in Vectashield mounting medium

(Vectorlabs) and visualized using a laser scanning confocal microscope. Antibodies:

rabbit anti-GFP 1:400 (Invitrogen), mouse anti-FLAG M2 1:1000 (Sigma), rabbit anti-stv

1:1000 (gift of Jrög Höhfeld), rat anti-α actinin 1:100 (Abcam), rabbit anti-ubiquitin

linkage-specific K63 anti-mouse 1:200 (Abcam), IgG- Alexa Fluor 488 1:200 (Life

Technologies), anti-mouse IgG- Alexa Fluor 568 1:500 (Life Technologies), anti-rabbit

IgG- Alexa Fluor 488 1:500 (Life Technologies) and Rhodamine Phalloidin 1:2000.

Protein aggregate analysis

Indirect flight muscles were prepared as described above from 9 individual flies per

genotype. Protein aggregates were identified using mouse anti-polyubiquitin 1:1000

(Enzo Life Sciences). Three muscles from each individual fly were imaged as z-series

and flattened into a single image as a max projection using confocal microscopy for a

total of 27 muscles per genotype. Images were analyzed using Image J "Analyze

Particles" function with a diameter of 100 pixels set for the minimum aggregate size.

Aggregate number and size were analyzed using ANOVA followed by Tukey's HSD

using the R package "multcomp" to generate significance groups with each letter group

being significantly different with a p value of ≤ 0.05. Within genotype/across time point

analyses were performed using the Welch two sample t-test in R.

Lifespan

For lifespan experiments using the UAS-p38Kb^{wt}, ref(2)p -/+, p38Kb$^{\Delta45/\Delta45}$, the stv EP

(stv^{wt}) lines, and their respective controls, virgin females were kept on standard

molasses *Drosophila* media. Due to a change in lab food, the stv RNAi 34408 and stv

RNAi 34409 lifespan experiments (with their respective controls) were performed on the

standard Bloomington *Drosophila* media. Virgin flies were collected and reared at 25°C

in a 12hour:12hour light:dark cycle. Flies were put on new food twice a week. The

number of dead animals was scored daily. Lifespan was analyzed using a log rank test

to compare genotypes with censored data on all genotypes and then on all pairwise

comparisons using the R package "survival" with Benjamini and Hochberg correction

(false discovery rate < 0.05).

Co-immunoprecipitation

Flies were aged 1 week or 5 weeks. Forty thoraxes per genotype per condition were

homogenized in high salt buffer (0.5 M KCl, 35% glycerol,10 mM HEPES pH 7.0, 5 mM

MgCl$_2$, 0.5 mM EDTA pH 8.0, 0.1% NP40 25 mM NaF, 1 mM Na$_2$VO$_4$, 1 mM DTT,

Complete protease inhibitor). The lysate was flash frozen in liquid nitrogen and quickly

thawed at 37°C. Then lysates were rocked at 4°C for 30 minutes and centrifuged at

14,200 x g for 30 minutes at 4°C. The supernatant was transferred to equilibrated beads

anti-Flag (M2) agarose (Sigma) or anti-GFP agarose (Chromotek) and rocked for 2

hours at 4°C. Beads were collected using a magnetic bar and washed four times with IP

buffer (50 mM HEPES pH 7.0, 100 mM KCl, 0.4% NP40 1.5 mM MgCl$_2$, 5% glycerol, 25

mM Na, 1 mM Na$_2$VO$_4$, 1 mM EDTA, 1 mM DTT, Complete protease inhibitor). Lysates

were then analyzed by immunoblotting using rabbit anti- GFP 1:1000 (Invitrogen),

mouse anti-FLAG M2 1:1000 (Sigma), rabbit anti-phospho-p38 1:1000 (Cell Signaling

Technologies), goat anti total p38 1:1000 (Santa Cruz Biotechnology), rabbit anti-stv

1:10,000 (gift of Jrög Höhfeld) or mouse anti-Lamin 1:1000 (DHSB).

Immunoblotting

Wild type flies (w[1118]) were aged either for 3, 15, 30, and 45 days or for 1-5 weeks. Three

thoraxes were dissected and homogenized in 1x Laemmli buffer. Immunoblots were

performed as described in [16]. Membranes were developed using SuperSignal West

Femto kit (ThermoFisher) or Pierce ECL (ThermoFisher) and exposed on

autoradiography film. Antibodies used were: rabbit anti-GFP 1:1000 (Invitrogen), rabbit

anti-starvin 1:10,000 (gift of Jrög Höhfeld), mouse anti-actin 1:5,000,000 (Sigma), mouse

anti-FLAG M2 (Sigma), rabbit anti-alpha tubulin (Cell Signaling Technologies), mouse

anti-Lamin 1:100 (DHSB), rabbit anti-phospho Lamin A Ser22 1:1000 (Thermofisher),

mouse anti-beta tubulin (E-10) 1:5000 (Santa Cruz Biotechnology), mouse anti- HRP

1:20,000 (Jackson Labs), rabbit anti-HRP 1:40,000 (Jackson Labs). Densitometry was

performed using a minimum of three independent blots. For statistical analysis of protein

expression level, pixel density of the tested protein was normalized within sample to the

loading control. These values were then normalized to control to calculate fold change.

The fold change values were analyzed by Student's t-test or ANOVA [27] as appropriate.

Sucrose Gradient Fractionation

$p38Kb^{Ex41/Ex41}$ and $p38Kb^{\Delta45/\Delta45}$ flies were aged three weeks. 30 thoraxes per genotype

were dissected and homogenized in NP40 lysis buffer. Samples were centrifuged at

800xg for 10 minutes at 4°C. The supernatant was then transferred to a 15-50% sucrose

discontinuous gradient. Samples were then ultracentrifuged at 55,000 rpm (201,000xg at

r_{av}) for 20 hours at 4°C in a TLS-55 in a Beckman Coulter Optima TLX Ultracentrifuge.

200μl fractions were collected and the pellet was resuspended in an equal volume of

NP40 Lysis Buffer.

Stv localization

Immunohistochemistry on $p38Kb^{Ex41/Ex41}$ and $p38Kb^{\Delta45/\Delta45}$ Indirect flight muscles was

performed as described above. Confocal images from five individual flies per genoytpe

were analyzed for average pixel density using ImageJ in three different non-overlapping

locations on each muscle for a total of 15 measurements per genotype. Average pixel

density was analyzed by Student's t-test using R.

Results

p38Kb regulates age-dependent protein homeostasis.

p38Kb null mutant animals ($p38Kb^{\Delta45/\Delta45}$, a deletion of the p38Kb gene) exhibit age-

dependent locomotor behavior defects and have a 48% lifespan reduction. In addition,

biochemical analysis suggests that p38Kb null mutants have increased levels of

insoluble polyubiquitinated proteins in the thoracic musculature of aged animals as

observed by immunoblot analysis [17]. However, protein aggregate size and distribution

have not been visualized in the p38Kb mutants, nor is it known whether augmenting

p38Kb activity in muscles leads to a change in protein homeostasis. Therefore, we

analyzed how p38Kb expression influences protein aggregation. We find that p38Kb null

mutants have an increased number of protein aggregates in the adult indirect flight

muscle at 1 week and 3 weeks of age (Figure 1A-B and E-F, Table S1) and increased

aggregate size with age (Figure 1G-H, Table S1) as compared to a genetic background

control (p38Kb$^{Ex41/Ex41}$, a precise excision allele). Furthermore, transgenic inhibition of

p38Kb in the muscle using a dominant negative kinase dead construct (p38Kb^{KD}, [16]) also

results in a significant increase in aggregate number, however aggregate size was not

affected (Figure 1I-L and Table S2). Conversely, as both strong and moderate levels of

p38Kb over-expression lead to an increased lifespan (37% and 14% extension,

respectively) [16], we tested if p38Kb over-expression also affects protein aggregation. We

find that both strong over-expression of p38Kb in the adult muscle using the Mef2-GAL4

driver (Figure 1C-D and M-P, Table S3) and moderate over-expression of p38Kb using

the MHC-GAL4 driver (Figure 1Q-T and Table S4) leads to decreased protein aggregate

number and size throughout the lifespan. It has been hypothesized that protein

aggregate accumulation and increased size may be toxic, potentially explaining the

decreased lifespan and age-dependent locomotor abnormalities in p38Kb^{Δ45} null mutants

and the increased longevity in the p38Kb over-expression animals.

p38Kb mediates age-dependent phenotypes through autophagy

In order to determine what mechanism plays a role in the clearance of polyubiquitinated

protein aggregates, we first tested what type of ubiquitin linkage is present in the

aggregates in wild type flies. In particular, K63-linked ubiquitination has been shown to

facilitate the formation of aggregates [28–30] that are cleared through autophagy [30–32]. We

find that a subset of aggregates from aging flies contain K63-linked ubiquitinated

proteins (Figure 2A-C), suggesting that these muscle protein aggregates are degraded

through autophagy. Polyubiquitinated protein aggregates can be degraded through

selective autophagy in which the adaptor protein p62 (ref(2)p in flies) promotes the

packaging and delivery of polyubiquitinated proteins to the autophagosome [33]. If p38Kb

requires selective autophagy to mediate protein homeostasis, then loss of ref(2)p will

block the p38Kb mediated decreased aggregation phenotype. We find that loss of a

single copy of *ref(2)p* results in fewer aggregates (Figure 2D-E and Table S5), which

may reflect compensation by other protein clearance mechanisms with aging, especially

as homozygous *ref(2)p* mutants are viable. When a single copy of *ref(2)p* is removed in

the p38Kb over-expression background, this prevents the reduced protein aggregation

observed in the p38Kb over-expression animals at both young and old ages (Figure 2D-

E and Table S5). This dominant interaction suggests that *ref(2)p* acts downstream of

p38Kb to promote the degradation of protein aggregates.

p38Kb colocalizes with the CASA complex in the adult flight muscle.

In order to better understand the role of p38Kb in protein homeostasis, we first

determined where in the muscle p38Kb localizes and find that a FLAG-tagged p38Kb

(green) colocalizes with the Z-disc marker alpha-actinin and is also present at the M-line

(Figure 2F). The muscle Z-disc is an area of high protein turnover, and one protein

quality control mechanism that localizes to the Z-disc in mice and adult flies is the

Chaperone-Assisted Selective Autophagy (CASA) complex [19,34]. In adult *Drosophila*

muscle, the CASA complex has been reported to consist of the Hsc70 homologue,

Hsc70-4, the HspB8 homologue CG14207 and the BAG-3 homologue starvin (stv) [19].

We find that p38Kb colocalizes with all the CASA complex members at the Z-disc and

M-line (Figure 2G-I).

p38Kb physically interacts with the CASA complex in the adult flight muscle.

In order to further determine if p38Kb physically interacts with the CASA complex in the

adult muscle, we performed co-immunoprecipitation experiments. In *Drosophila* cell

culture p38Kb^{wt} was shown to co-immunoprecipitate with the HspB8 homologue

CG14207[18], therefore, we tested if this interaction also occurs in adult indirect flight

muscle. As the interaction between p38Kb and its targets may be transient, we utilized a

FLAG-tagged p38Kb kinase dead construct (UAS-p38Kb^{KD} Mef2-GAL4) that is able to

be activated and bind to a target but cannot phosphorylate it, leading to a delayed

release of the target[35]. We expressed p38Kb^{KD} in the adult flight muscle of a fly that also

endogenously expressed GFP tagged HspB8, in which GFP is spliced in as a new exon

of the endogenous gene. This results in a GFP fusion protein that is expressed in the

same pattern as endogenous HspB8 ([19] and Figure 2H'). We find that p38Kb^{KD} was able

to pull down HspB8 (Figure 2J) as compared to the HspB8-GFP control background. In

addition, expression of p38Kb^{KD} in a wildtype background was able to pull down un-

tagged endogenous stv (Figure 2K).   We then performed reverse IPs in which we

immunoprecipitated each endogenously GFP tagged CASA complex protein (Hsc70-4,

HspB8, or stv) and probed for FLAG tagged p38Kb^{KD} and found that p38Kb co-

immunoprecipitates with the CASA complex (Figure 2L-N) with stronger binding at

younger ages than older ages. Interestingly, we found no significant difference in the

expression levels of Hsc70-GFP, stv-GFP, and over-expression of p38Kb^{KD} with age

(Figure S1A-B, D) while levels of HspB8-GFP increased with age (Figure S1C). These

data suggest that the age-dependent interaction between p38Kb and the CASA complex

may be due to changes in the physical interactions between these proteins rather than a

reduction in protein levels.

To verify that endogenous p38Kb can interact with the CASA complex, we

performed co-immunoprecipitations with muscle lysates from flies expressing either

endogenously GFP tagged Hsc70-4, HspB8 or stv in a wild type background and probed

for endogenous activated p38K that is dual phosphorylated on the TGY motif. We find

that all three members of the CASA can co-immunoprecipitate with endogenous

phosphorylated p38K in the muscle (Figure 2O). These results suggest that p38Kb is

able to physically interact with the CASA complex.

p38Kb acts downstream of the CASA complex.

As p38Kb physically interacts with the CASA complex, we next tested if p38Kb is

acting upstream activity of the complex to promote the initial recognition of misfolded

proteins or the downstream activity of the complex to direct un-foldable proteins to the

autophagosome. As BAG-3/stv provides both the specificity to the CASA complex and is

involved in the handoff of damaged proteins to p62/ref(2)p for autosomal degradation [21–

25], we tested for genetic interactions between p38Kb and *stv*. *stv* null mutants have

impaired locomotor functions, muscle degeneration and early lethality [19,36]. Due to the

severity of these null phenotypes, we utilized *stv* RNAi lines to generate an allelic series

of *stv* loss of function in the muscle. We find that weak inhibition of stv (UAS-stv

RNAi[34408] MHC-GAL4) had no effect on protein aggregation (Table S6). However,

moderate inhibition of stv (UAS-stv RNAi[34409] MHC-GAL4) results in increased protein

aggregate number and size (Figure 3A-B and Table S7) and results in a decrease in

lifespan particularly in the first half of life as compared to outcrossed controls (Figure 3C

and Table S8). Strong inhibition of *stv* (UAS-stv RNAi[34408] Mef2-GAL4) leads to a

severely reduced lifespan of ~4 days on average (Figure 3D and Table S9).

If p38Kb acts to regulate the downstream activity of the CASA complex, then over-

expression of p38Kb should rescue these *stv* RNAi phenotypes. We find that over-

expression of p38Kb rescues both protein aggregate number and size in the *stv*

moderate inhibition background (Figure 3A-B and Table S7) and also rescues the

lifespan defect (Figure 3C and Table S9). Furthermore, p38Kb over-expression is able to

partially rescue the decreased lifespan exhibited by strong *stv* inhibition increasing

average lifespan from 4 days to 13 days (Figure 3D and Table S10). To further test if

p38Kb acts downstream of the CASA complex, we combined loss of p38Kb with CASA

complex over-expression. If p38Kb acts downstream of the CASA complex, then over-

expression of the CASA complex members in the p38Kb mutant background will not be

able to rescue the p38Kb mutant phenotypes. We find that over-expression of stv in the

p38Kb mutants not only fails to rescue the p38Kb mediated short lifespan defect but

reduces the lifespan further (Figure 3E and Table S11). This is particularly striking as

over-expression of stv in a wild type background does not significantly affect lifespan

(Figure 4H and Table S12) as compared to the outcrossed transgene control (p value =

0.195). This suggests that p38Kb may be a limiting factor in regulating the downstream

activity of the CASA complex in which it hands the poly-ubiquitinated misfolded proteins

to ref(2)p for degradation through the autophagosome.

p38Kb regulates the activity of the CASA complex in protein homeostasis.

If p38Kb is a limiting factor for stv function, then the combined over-expression of

p38Kb and stv may result in a further beneficial effect. We find that over-expression of

stv alone results in fewer aggregates at young and old ages (Figure 4A-B and Table

S13) and smaller aggregates with age (Figure 4D and Table S13) as compared to

outcrossed controls. However, co-over-expression of p38Kb and stv does not result in a

further reduction in aggregate number as compared to over-expression of p38Kb or stv

alone (Figure 4A and Table S13). By 5 weeks of age, co-over-expression of p38Kb and

stv flies have comparable aggregate number to controls (Figure 4B and Table S13).

Conversely, p38Kb and stv co-over-expression results in a reduction in aggregate size at

a young age compared to over-expression of stv alone (Figure 4C and Table S13, p

value < 0.001), suggesting that p38Kb is a limiting factor for stv function in regulating

aggregate size. Unlike with aggregate number, the aggregates remain significantly

smaller in size as the flies age in the combined over-expression background (Figure 4D

and Table S13). We also find that co-over-expression of p38Kb and stv leads to an

additional 5% increase in lifespan relative to p38Kb over-expression alone (Figure 4H

and Table S12). Interestingly, the co-over-expression animals show a very similar

lifespan to the p38Kb over-expression alone animals until ~day 50, when the p38Kb

alone animals begin to die at a faster rate (Figure 4H). One explanation is that co-over-

expression of p38Kb and stv provides beneficial effects in early adulthood that continue

throughout adulthood leading to increased lifespan despite the presence of protein

aggregates. Another possibility is that aggregate size is a reflection of aggregate

content. If the CASA complex can only target a subset of proteins for refolding or

degradation, then increasing CASA complex activity could lead to smaller aggregates as

the CASA targets are effectively degraded through an interaction with p38Kb and ref(2)p

rather than aggregating. Therefore, size and/or content may play a more important role

in determining lifespan as compared to overall aggregate number.

p38Kb is required for proper localization of stv in the muscle.

Since we find that p38Kb acts between stv and ref(2)p in regulating protein

aggregation, we hypothesize that p38Kb plays a role in stabilizing the transfer of

misfolded proteins primed for degradation from the CASA complex to ref(2)p. If so, then

over-expression of p38Kb would lead to increased efficiency in targeting misfolded

proteins to the autophagosome (Figure 1M-T), and inhibition of p38Kb would lead to an

increase in protein aggregation (Figure 1E-L). In addition, loss of p38Kb may lead to a

failure of stv to maintain its interaction with the CASA complex or to interact with ref(2)p.

Therefore, we tested if stv is still able to localize to the Z-disc in the absence of p38Kb.

We find that in p38Kb null mutants, stv is more diffuse throughout the muscle but is still

able to localize to the Z-disc and M-line in (Figure 4E-G), suggesting that its localization

is partially impaired in the absence of p38Kb. Furthermore, we find that the localization

of the CASA complex member HspB8 in the muscle is unaffected by loss of p38Kb

(Figure S2). These data indicate that p38Kb may play a role in stabilizing the interaction

between the CASA complex and stv that allows stv to direct misfolded proteins to

ref(2)p.

Lamin protein binds to the CASA complex and accumulates in stv RNAi and

p38Kb mutants.

In order to determine if p38Kb might be playing a role in the targeting of

misfolded proteins from stv to ref(2)p, we first needed to identify a protein target of both

stv and p38Kb. Therefore, we focused on proteins associated with Limb-Griddle

Muscular Dystrophy (LGMD), which is caused by mutations in either BAG-3 (stv) or

HspB8 [37–39]. In addition, p38K signaling has been implicated in LGMD [40–42]. *Drosophila*

have 19 orthologues of LGMD proteins, one of which is Lamin A/C (Lamin or Lamin Dm0

in flies). Lamin is of particular interest since mutations in Lamin A/C also result in the

accelerated aging disorder Hutchinson-Gilford progeria [43–45], and Lamin has been shown

to aggregate under oxidative stress conditions [46]. Mutations in Lamin A/C are also

sufficient to induce Lamin aggregation and abnormal nuclear morphology in human cell

culture, *C. elegans*, and *Drosophila* systems [44,47–54]. Furthermore in *Drosophila*, Lamin

knockouts have similar phenotypes to p38Kb and/or stv mutants, such as reduced

locomotor function and increased activity of the Nrf-2/Keap-1 pathway [49,55,56].

If Lamin is a target of the CASA complex, then decreased activity of the CASA

complex should result in an accumulation of Lamin protein. We find that Lamin levels do

not change with age (Figure 5A), however, inhibition of stv results in a significant

increase in the total amount of Lamin protein regardless of age (Figure 5A-B, p value

0.028). Interestingly, loss of p38Kb leads to an age-dependent increase in Lamin protein

as compared to age matched controls (Figure 5C-D, p value = 0.006643). As the

association between p38Kb and the CASA seems to decline with age (Figure 2L-N),

these data suggest that p38Kb activity later in life is important for maintaining protein

homeostasis and lifespan.

To investigate if Lamin is accumulating in the protein aggregates, we performed

fractionation experiments in which we find that Lamin is predominantly found in fractions

5-7 in controls with the majority concentrated in faction 6 (Figure 5E). Interestingly, we

also find a high molecular weight from of Lamin mainly restricted to fraction 5 (Figure

5E). In addition, we also observe that both the main and high molecular weight Lamin

species are in the aggregate containing pellet (Figure 5E). We next tested how loss of

p38Kb affects the aggregation of Lamin and find that p38Kb mutants have decreased

Lamin in fractions 5-7 with a concurrent increase of both the main and high molecular

weight species of Lamin in the pellet (Figure 5E). This suggests that Lamin is a target of

p38Kb and that loss of p38Kb results in increased aggregation of Lamin.

Of interest is this high molecular weight form of Lamin, which is present in

fraction 5 and the aggregates (Figure 5E). One important protein domain in Lamin is the

CaaX box, which is farnesylated and required for Lamin localization to the inner nuclear

membrane[57–60]. To test if this might be a farnesylated form of Lamin, we utilized the

Lam^{A25} mutant which has a frameshift that results in the loss of the C-terminal CaaX

box[61]. We find that the high molecular weight form of Lamin is lost in the Lam^{A25} mutant

whereas the predominant 75kDa band of Lamin is still present (Figure 5F). These data

suggest that the high molecular weight Lamin we observe is the farnesylated form.

In addition to farnesylation, Lamin can also be phosphorylated by a variety of

kinases that can change the solubility of Lamin [54,62]. For example, in a *Drosophila* cell

culture system, a phospho-mimic mutation at Ser45 (the equivalent of Ser22 in humans)

results in Lamin aggregation in both the nucleus and cytoplasm [54]. This serine residue is

phosphorylated by the cd2 kinase [63], but also happens to be a potential p38K

phosphorylation site. Therefore, we tested if the levels of phospho- Ser45 Lamin are

altered in p38Kb mutants. We find that phospho-Lamin levels are not significantly altered

in the p38Kb mutants (Figure 5G-H p=0.09), suggesting that p38Kb is not required for

Ser45 phosphorylation. We next examined if phosphorylated Lamin is also present in the

aggregates. We find that in controls, phosphorylated Lamin is present in factions 5-7

with low amounts in the pellet (Figure 5I). We also observe a low molecular weight

phospho-Lamin form that appears in the pellet (Figure 5I). In the p38Kb mutants,

phospho-Lamin expression is reduced in fractions 6 and 7 and accumulates in the pellet

(Figure 5I). Furthermore, increasingly smaller low molecular weight species of phospho-

Lamin are present in the p38Kb$^{\Delta45}$ mutants including the pellet (Figure 5I), suggesting

that loss of p38Kb prevents the effective clearance of these Lamin cleavage forms.

Interestingly, loss of the Lamin CaaX box does not affect these lower molecular weight

forms, however, it does result in an increase in the phosphorylation of full-length Lamin

(Figure 5F). Additionally, we do not detect phosphorylation positive high molecular

weight forms of Lamin (Figure 5F), suggesting that farnesylated lamin is not

phosphorylated at Ser45. These data combined with our finding that full length Lamin is

phosphorylated, suggest that when Lamin is unphosphorylated it is prone to

farnesylation that may lead to aggregation.

To determine if Lamin is a direct target of p38Kb and the CASA complex, we

tested if Lamin can bind to the CASA complex. We immunoprecipitated endogenously

GFP tagged HspB8, Hsc70-4 and stv from adult *Drosophila* muscle and probed for

endogenous Lamin. We find that full-length Lamin co-immunoprecipitates with all three

members of the CASA complex (Figure 6A), but do not detect binding with either the

higher or lower molecular weight forms of Lamin found in the fractions (Figure 5E and I).

Furthermore, we expressed either a nuclear localization signal tagged GFP or Lamin

tagged with GFP in the muscle and found that endogenous stv co-immunoprecipitates

with Lamin (Figure 6B). These data suggest that the CASA complex is able to bind to

misfolded forms of full-length Lamin for targeting to the autophagosome.

As we hypothesize that p38Kb is interacting with the CASA complex at the point

in which the misfolded targets are handed over to ref(2)p for degradation, we tested if

p38Kb also binds to Lamin. We find that p38Kb[KD] expressed in the muscle co-

immunoprecipitates with both high and low molecular weight forms of Lamin (Figure 6C).

Interestingly, p38Kb physically interacts with the aggregate prone low molecular weight

species of Lamin as opposed to the predominant full-length form of Lamin (Figure 6C).

More striking is that p38Kb[KD] physically interacts with increasingly larger forms of Lamin

(Figure 6C), suggesting that p38Kb is interacting with the poly-ubiquitinated forms of

Lamin.

We have developed a model in which misfolded proteins are targeted by stv to

the CASA complex. Those proteins that cannot be refolded are then tagged with poly-

ubiquitin. These poly-ubiquitinated proteins are handed off to ref(2)p in a process

mediated by p38Kb. This model predicts that in p38Kb mutants these poly-ubiquitinated

proteins accumulate and then form aggregates. It additionally predicts that over-

expression of p38Kb would lead to increased efficiency of the poly-ubiquitinated proteins

being targeted to the autophagosome. This model would also predict that misfolded full-

length Lamin would bind to the CASA complex for refolding. If Lamin cannot be refolded

it is then poly-ubiquitinated and transferred to ref(2)p through an interaction with p38Kb.

Discussion

We find that the aging gene p38Kb regulates age-dependent protein homeostasis

through an interaction with the CASA complex and is acting at a step between the poly-

ubiquitination of a un-foldable target and its transfer to ref(2)p for autolysosomal

mediated degradation. Interestingly, we find that p38Kb is important for the proper

localization of stv to the Z-disc but does not affect the localization of other complex

members. This suggests that p38Kb may play a role in maintaining the interaction

between stv and the CASA complex which may be necessary for target transfer to

ref(2)p. stv has a conserved MAPK docking site as well several potential conserved

p38K phosphorylation sites. Furthermore, mammalian p62 has been shown to bind to

p38Kb in vitro through two domains [64], which are partially conserved in flies. Therefore,

one possibility is that p38Kb mediated phosphorylation of stv facilitates the localization of

a functional CASA complex to the Z-disc, where damaged proteins are rapidly turned

over. Another possibility is that p38Kb binds to stv and ref(2)p and that the

phosphorylation of stv is required for target hand-over to ref(2)p so that in the absence of

p38Kb, stv cannot transfer targets to ref(2)p. A consequence of this may be that stv and

the protein target together are released from the CASA complex, leading to

mislocalization of stv. As we don't detect a decrease in phospho-Lamin in the p38Kb

mutants or find that the p38Kb can pull-down phospho-Lamin, it is unlikely that p38Kb is

directly phosphorylating the target proteins as a part of the stv-ref(2)p hand-over

process.

How protein aggregation contributes to aging and disease has been an area of

great interest. One outstanding question is if protein aggregation is a consequence or

cause of aging. It has been hypothesized that protein aggregates accumulate with age

as the amount of damaged or misfolded proteins increase. However, it is not clear

whether or not these aggregating proteins are toxic leading to tissue dysfunction and a

disease state. Previous studies have found that long-lived fly strains such as over-

expression of Foxo or parkin result in reduced protein aggregate formation [4,65].

Therefore, we hypothesized that decreased protein aggregation would lead to improved

measures of healthspan, while increased protein aggregation would lead to reduced

measures of healthspan. As expected, we find that the short-lived p38Kb mutants, which

exhibit premature locomotor behavior defects [16], have large and numerous protein

aggregates. Similarly, inhibition of stv results in increased aggregation and decreased

lifespan. We also find that over-expression of p38Kb leads to decreased protein

aggregation and increased lifespan. However, over-expression of stv leads to decreased

aggregate number but does not extend lifespan. One possibility is that decreasing

aggregation alone is not sufficient to promote lifespan extension. Another possibility is

that aggregate number is not as critical as aggregate size and/or content, particularly

early in life, for lifespan extension. We find that stv over-expression only results in

decreased aggregate size in older animals, however, co-over-expression of p38Kb and

stv leads to reduced aggregate size at a young and old ages and leads to a further

increase in lifespan. Thus, the turnover of specific toxic protein or subset of proteins

early in life may lead to a reduction in the exposure to these toxic species and a lifespan

extension. We find that Lamin, which is mutated to an aggregation prone protein form in

Hutchinson-Gilford progeria [44,53,62,66], is a target of p38Kb and the CASA complex.

Interestingly, p38Kb and Lamin mutants share similar phenotypes such as age-

dependent locomotor impairments [16,55,56] and upregulation of the Nrf-2/Keap-1 pathway

[16,49]. As loss of p38Kb leads to an accumulation of Lamin in the aggregates, this may be

toxic to the cell, leading to impaired locomotor function, increased stress and decreased

lifespan. Thus, we have found that one aging gene (p38Kb) regulates a second,

unrelated aging gene (Lamin) via the CASA complex. These data suggest a new link

between aging pathways and how they may converge through the regulation of protein

homeostasis.

Acknowledgements

We would also like to acknowledge the Bloomington Drosophila Stock Center (NIH

P40OD018537), the Vienna Drosophila Resource Center, and Kyoto Stock Center for

providing fly stocks used in this study. A. Vrailas-Mortimer was funded by start-up funds

from the University of Denver, a Knoebel Center for the Study of Aging pilot grant, start-

up funds from Illinois State University and was supported by the National Institute of

Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health

under Award Number R15AR070505. The content is solely the responsibility of the

authors and does not necessarily represent the official views of the National Institutes of

Health. S. Sanyal was funded by startup funds from Emory University. S. Ryan was

supported by a Knoebel Center for the Study of Aging pilot grant to A. Vrailas-Mortimer

and S. Barbee, N. Mortimer was funded by start-up funds from Illinois State University.

The funders had no role in study design, data collection and analysis, decision to

publish, or preparation of the manuscript.

Figure Legends

Fig 1. p38Kb regulates age-dependent protein homeostasis.

Confocal micrographs of polyubiquitin positive protein aggregates in the adult indirect

flight muscle in **A)** a precise excision genetic background control p38Kb$^{Ex41/Ex41}$ and **B)**

p38Kb$^{\Delta45/\Delta45}$ null mutants at three weeks of age and in **C)** outcrossed Mef2-GAL4

controls (Mef2>w^{11118}) and **D)** UAS-p38Kb^{wt} Mef2-GAL4 (Mef2>p38Kb^{wt}) over-expression

animals at 5 weeks of age. Scale bar equals 6.2 μm. Box-Whisker plots of aggregate

number in p38Kb$^{\Delta45/\Delta45}$ mutants as compared to p38Kb$^{Ex41/Ex41}$ controls at **E)** 1 week and

F) 3 weeks of age and aggregate size at **G)** 1 week and **H)** 3 weeks of age. Aggregate

number in p38Kb^{KD} Mef2-GAL4 (Mef2>p38Kb^{KD}) and outcrossed Mef2-GAL4 controls at

I) 1 week and **J)** 5 weeks of age and aggregate size at **K)** 1 week and **L)** 5 weeks of

age. Aggregate number in strong p38Kb over-expression animals and outcrossed Mef2-

GAL4 controls at **M)** 1 week and **N)** 5 weeks of age and aggregate size at **O)** 1 week

and **P)** 5 weeks of age. Aggregate number in moderate p38Kb over-expression

(MHC>p38Kb^{wt}) animals and outcrossed MHC-GAL4 controls (MHC>w^{1118}) at **Q)** 1 week

and **R)** 5 weeks of age and aggregate size at **S)** 1 week and **T)** 5 weeks of age.

Fig 2. p38Kb regulates aging phenotypes through ref(2)p.

A-C) A subset of polyubiquitinated protein aggregates (green) in 3 week old

p38Kb$^{Ex4/Ex41}$ control muscle contain K63 ubiquitinated (magenta) proteins (white

arrows). **D-E)** Protein aggregate number in ref(2)p heterozygous mutant backgrounds at

D) 1 week and **E)** 5 weeks. Loss of a single copy of ref(2)p prevents the p38Kb mediated

reduced protein aggregation at 1 week and 5 weeks of age. Asterisks denote a p-value

of ≤0.001. **(F-I)** Localization of a FLAG-tagged p38Kb (green in **F-I** and **F''-I''**) in the

adult indirect flight muscle. **F)** FLAG-tagged p38Kb localizes to the Z-disc (arrows) as

exhibited by colocalization with the Z-disc protein alpha-actinin (magenta, **F'** and **F''**), as

well as the M-line (arrowheads). **G-I)** p38Kb colocalizes with each CASA complex

member (magenta, **G'-I'**) at the Z-disc. Over-expression of a FLAG tagged p38Kb in the

muscle in **G**) an endogenous Hsc70-GFP fusion protein background, **H**) an endogenous

HspB8-GFP fusion protein background, and **I)** a wildtype background with endogenous

levels of stv. **J-K)** Muscle lysates expressing control and Mef2-GAL4 UAS-p38Kb[KD-FLAG]

in either **J)** an endogenously expressed HspB8-GFP background or **K)** a wildtype

background were immunoprecipitated using anti-FLAG coated beads. **J)** Immunoblots

were probed with anti-GFP to detect the presence of HspB8 in the IP lysates or with **K)**

anti-stv to detect the presence of stv in the IP lysates. **L-N)** Muscle lysates expressing

p38Kb [KD-FLAG] in an endogenously expressed **L)** Hsc70-GFP, **M)** HspB8-GFP and **N)** stv-

GFP background were immunoprecipitated using anti-GFP coated beads. Immunoblots

were probed with anti-FLAG to detect the presence of p38Kb in the IP lysates. Note: stv

is tagged with both GFP and FLAG. **O)** Endogenous phospho-p38 co-

immunoprecipitates with Hsc70-4-GFP, HspB8-GFP, and stv-GFP. Muscle lysates were

immunoprecipitated with anti-GFP beads, and immunoblots were probed with anti-

phospho-p38K.

Fig 3. p38Kb genetically interacts with stv to regulate protein homeostasis and

lifespan. A) Protein aggregate number and **B)** size in the moderate stv knockdown

background using MHC-GAL4 at 5 weeks. Asterisks denote a p-value of ≤0.001. **C)**

Moderate over-expression of p38Kb (red line) results in an increased lifespan as

compared to the MHC-GAL4 controls and p38Kb transgene control (black line and gray

lines, respectively). Moderate knockdown of stv using the MHC-GAL4 results in a

decreased lifespan (pink line compared to yellow and black lines) and is rescued by

p38Kb over-expression (compare pink line to blue line). **D)** Strong over-expression of

p38Kb (red line) results in an increased lifespan as compared to the Mef2-GAL4 controls

and p38Kb transgene control (black line and gray lines, respectively). Strong knockdown

of stv using the Mef2-GAL4 results in a decreased lifespan (pink line compared to yellow

and black lines) and is partially rescued by p38Kb over-expression (compare pink line to

blue line). **E)** Over-expression of stv in the p38Kb mutant background results in a further

reduction of lifespan as compared to p38Kb mutant controls (compare blue line to red

and grey lines).

Fig 4. p38Kb is limiting for stv function and localization.

Protein aggregate number at **A)** 1 week and **B)** 5 weeks and protein aggregate size at

C) 1 week and **D)** 5 weeks measured in stv over-expression backgrounds. Over-

expression of stv leads to reduced aggregate number at 1 and 5 weeks and aggregate

size at 5 weeks. Co-over-expression of stv and p38Kb does not result in a further

decrease in protein aggregate number but trends towards decreased aggregate size at

both 1 and 5 weeks of age. Asterisks denote a p-value of ≤0.001 when compared to the

GAL4 control. **E)** stv localizes to the adult muscle Z-disc and M-line (white arrows) in

control animals. **F)** stv localization is disrupted in p38Kb mutants. **G)** Quantification of

average pixel density. **H)** Over-expression of stv alone has no significant effect on

lifespan (pink line as compared to yellow and black lines), however, co-over-expression

of stv and p38Kb results in a further increase in lifespan (compare red line to blue line).

Fig 5. p38Kb and stv regulate Lamin aggregation.

A) Immunoblot analysis of stv-RNAi and GAL4 controls flies muscle lysates probed with

anti-Lamin and **B)** quantified using densitometry. **C)** Immunoblot analysis of

p38Kb$^{Ex41/Ex41}$ control and p38Kb$^{\Delta45/\Delta45}$ mutant muscle lysates probed with anti-Lamin

and **D)** quantified using densitometry. **H)** Immunoblots of sucrose gradient fractions

from 3 week old p38Kb$^{Ex41/Ex41}$ control and p38Kb$^{\Delta45/\Delta45}$ mutant muscle probed with anti-

Lamin. **G)** Immunoblot analysis of p38Kb$^{Ex41/Ex41}$ control and p38Kb$^{\Delta45/\Delta45}$ mutant muscle

lysates probed with anti-phospho Lamin and **H)** quantified using densitometry. **I)**

Immunoblots of sucrose gradient fractions from 3 week old p38Kb$^{Ex41/Ex41}$ control and

p38Kb$^{\Delta45/\Delta45}$ mutant muscle probed with anti-phospho-Lamin.

Fig 6. p38Kb and stv physically interact with Lamin.

A) Endogenous HspB8-GFP, Hsc70-GFP and stv-GFP-FLAG fusion proteins were

immunoprecipitated from adult muscle lysate using anti-GFP coated beads.

Immunoblots were probed with anti-Lamin and anti-GFP. **B)** A nls-GFP or Lamin-GFP

was expressed in muscles using MHC-GAL4. GFP proteins were immunoprecipitated

from adult muscle lysates using anti-GFP coated beads. Immunoblots were probed with

anti-stv and anti-GFP. C) p38Kb^{KD} was expressed in muscles using Mef2-GAL4. p38Kb

was immunoprecipitated from adult muscle lysates using anti-FLAG coated beads.

Immunoblots were probed with anti-lamin and anti-FLAG.

Fig S1. CASA complex expression with age. Immunoblot analysis of **A)** endogenously

GFP tagged Hsc70-4, **B)** endogenously GFP tagged HspB8, **C)** stv, and **D)** the FLAG

tagged p38Kb kinase dead transgene with age. Asterisks denote a p value of ≤0.05 as

compared to the 3 day time point.

Fig S2. Localization of HspB8 in p38Kb mutants. Endogenously GFP-tagged HspB8

was analyzed in the muscle of three week old **A)** controls and **B)** p38Kb mutants. HspB8

(green) localizes to the Z-disc and M-line of the indirect flight muscle (actin in magenta)

in both control and p38Kb mutant animals.

References

1.   Taylor, R. C. & Dillin, A. Aging as an event of proteostasis collapse. *Cold Spring*

Harb. Perspect. Biol. **3**, (2011).

2.   Ben-Zvi, A., Miller, E. A. & Morimoto, R. I. Collapse of proteostasis represents an

early molecular event in Caenorhabditis elegans aging. *Proc. Natl. Acad. Sci. U. S.*

A. **106**, 14914–14919 (2009).

3.   Cohen, E., Bieschke, J., Perciavalle, R. M., Kelly, J. W. & Dillin, A. Opposing

activities protect against age-onset proteotoxicity. *Science* **313**, 1604–1610 (2006).

4.   Demontis, F. & Perrimon, N. FOXO/4E-BP signaling in Drosophila muscles regulates

organism-wide proteostasis during aging. *Cell* **143**, 813–825 (2010).

5.   Morley, J. F., Brignull, H. R., Weyers, J. J. & Morimoto, R. I. The threshold for

polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and

influenced by aging in Caenorhabditis elegans. *Proc. Natl. Acad. Sci. U. S. A.* **99**,

10417–10422 (2002).

6.   Palazzolo, I. *et al.* Overexpression of IGF-1 in muscle attenuates disease in a mouse

model of spinal and bulbar muscular atrophy. *Neuron* **63**, 316–328 (2009).

7.   Schnöder, L. *et al.* Deficiency of Neuronal p38α MAPK Attenuates Amyloid

Pathology in Alzheimer Disease Mouse and Cell Models through Facilitating

Lysosomal Degradation of BACE1. *J. Biol. Chem.* **291**, 2067–2079 (2016).

8.   Webber, J. L. & Tooze, S. A. New insights into the function of Atg9. *FEBS Lett.* **584**,

1319–1326 (2010).

9.   Liu, B., Cheng, Y., Zhang, B., Bian, H. & Bao, J. Polygonatum cyrtonema lectin

induces apoptosis and autophagy in human melanoma A375 cells through a

mitochondria-mediated ROS-p38-p53 pathway. *Cancer Lett.* **275**, 54–60 (2009).

10. Younce, C. W. & Kolattukudy, P. E. MCP-1 causes cardiomyoblast death via

autophagy resulting from ER stress caused by oxidative stress generated by

inducing a novel zinc-finger protein, MCPIP. *Biochem. J.* **426**, 43–53 (2010).

11. Duan, W.-J. *et al.* Silibinin activated p53 and induced autophagic death in human

fibrosarcoma HT1080 cells via reactive oxygen species-p38 and c-Jun N-terminal

kinase pathways. *Biol. Pharm. Bull.* **34**, 47–53 (2011).

12. Zhuang, Y., Li, Y., Li, X., Xie, Q. & Wu, M. Atg7 Knockdown Augments Concanavalin

A-Induced Acute Hepatitis through an ROS-Mediated p38/MAPK Pathway. *PloS One*

11, e0149754 (2016).

13. Henson, S. M. *et al.* p38 signaling inhibits mTORC1-independent autophagy in

senescent human CD8[+] T cells. *J. Clin. Invest.* **124**, 4004–4016 (2014).

14. Luo, Y. *et al.* Autophagy regulates ROS-induced cellular senescence via p21 in a

p38 MAPKα dependent manner. *Exp. Gerontol.* **46**, 860–867 (2011).

15. Rudolf, E., Rezáčová, K. & Cervinka, M. Activation of p38 and changes in

mitochondria accompany autophagy to premature senescence-like phenotype switch

upon chronic exposure to selenite in colon fibroblasts. *Toxicol. Lett.* **231**, 29–37

(2014).

16. Vrailas-Mortimer, A. *et al.* A muscle-specific p38 MAPK/Mef2/MnSOD pathway

regulates stress, motor function, and life span in Drosophila. *Dev. Cell* **21**, 783–795

(2011).

17. Belozerov, V. E., Ratkovic, S., McNeill, H., Hilliker, A. J. & McDermott, J. C. In vivo

interaction proteomics reveal a novel p38 mitogen-activated protein kinase/Rack1

pathway regulating proteostasis in Drosophila muscle. *Mol. Cell. Biol.* **34**, 474–484

(2014).

18. Guruharsha, K. G. *et al.* A protein complex network of Drosophila melanogaster. *Cell*

147, 690–703 (2011).

19. Arndt, V. *et al.* Chaperone-assisted selective autophagy is essential for muscle

maintenance. *Curr. Biol. CB* **20**, 143–148 (2010).

20. Stürner, E. & Behl, C. The Role of the Multifunctional BAG3 Protein in Cellular

Protein Quality Control and in Disease. *Front. Mol. Neurosci.* **10**, 177 (2017).

21. Behl, C. BAG3 and friends: co-chaperones in selective autophagy during aging and

disease. *Autophagy* **7**, 795–798 (2011).

22. Gamerdinger, M. *et al.* Protein quality control during aging involves recruitment of

the macroautophagy pathway by BAG3. *EMBO J.* **28**, 889–901 (2009).

23. Kettern, N., Dreiseidler, M., Tawo, R. & Höhfeld, J. Chaperone-assisted degradation:

multiple paths to destruction. *Biol. Chem.* **391**, 481–489 (2010).

24. Min, J.-N. *et al.* CHIP deficiency decreases longevity, with accelerated aging

phenotypes accompanied by altered protein quality control. *Mol. Cell. Biol.* **28**, 4018–

4025 (2008).

25. Terman, A., Gustafsson, B. & Brunk, U. T. Autophagy, organelles and ageing. *J.*

Pathol. **211**, 134–143 (2007).

26. Dietzl, G. *et al.* A genome-wide transgenic RNAi library for conditional gene

inactivation in Drosophila. *Nature* **448**, 151–156 (2007).

27. R. C. Team. R Foundation for Statistical Computing. (2015).

28. Lim, K. L. *et al.* Parkin mediates nonclassical, proteasomal-independent

ubiquitination of synphilin-1: implications for Lewy body formation. *J. Neurosci. Off.*

J. Soc. Neurosci. **25**, 2002–2009 (2005).

29. Olzmann, J. A. *et al.* Parkin-mediated K63-linked polyubiquitination targets misfolded

DJ-1 to aggresomes via binding to HDAC6. *J. Cell Biol.* **178**, 1025–1038 (2007).

30. Yung, C., Sha, D., Li, L. & Chin, L.-S. Parkin Protects Against Misfolded SOD1

Toxicity by Promoting Its Aggresome Formation and Autophagic Clearance. *Mol.*

Neurobiol. **53**, 6270–6287 (2016).

31. Hao, R. *et al.* Proteasomes activate aggresome disassembly and clearance by

producing unanchored ubiquitin chains. *Mol. Cell* **51**, 819–828 (2013).

32. Tan, J. M. M. *et al.* Lysine 63-linked ubiquitination promotes the formation and

autophagic clearance of protein inclusions associated with neurodegenerative

diseases. *Hum. Mol. Genet.* **17**, 431–439 (2008).

33. Katsuragi, Y., Ichimura, Y. & Komatsu, M. p62/SQSTM1 functions as a signaling hub

and an autophagy adaptor. *FEBS J.* **282**, 4672–4678 (2015).

34. Ulbricht, A. *et al.* Cellular mechanotransduction relies on tension-induced and

chaperone-assisted autophagy. *Curr. Biol. CB* **23**, 430–435 (2013).

35. Hattori, A., Mizuno, T., Akamatsu, M., Hisamoto, N. & Matsumoto, K. The

Caenorhabditis elegans JNK Signaling Pathway Activates Expression of Stress

Response Genes by Derepressing the Fos/HDAC Repressor Complex. *PLOS*

Genet. **9**, e1003315 (2013).

36. Coulson, M., Robert, S. & Saint, R. Drosophila starvin encodes a tissue-specific

BAG-domain protein required for larval food uptake. *Genetics* **171**, 1799–1812

(2005).

37. Sato, T. *et al.* DNAJB6 myopathy in an Asian cohort and cytoplasmic/nuclear

inclusions. *Neuromuscul. Disord. NMD* **23**, 269–276 (2013).

38. Selcen, D. *et al.* Mutation in BAG3 causes severe dominant childhood muscular

dystrophy. *Ann. Neurol.* **65**, 83–89 (2009).

39. Sarparanta, J. *et al.* Mutations affecting the cytoplasmic functions of the co-

chaperone DNAJB6 cause limb-girdle muscular dystrophy. *Nat. Genet.* **44**, 450–455,

S1-2 (2012).

40. Fanzani, A. *et al.* Hypertrophy and atrophy inversely regulate Caveolin-3 expression

in myoblasts. *Biochem. Biophys. Res. Commun.* **357**, 314–318 (2007).

41. Capanni, C. *et al.* Dysferlin in a hyperCKaemic patient with caveolin 3 mutation and

in C2C12 cells after p38 MAP kinase inhibition. *Exp. Mol. Med.* **35**, 538–544 (2003).

42. Suzuki, N. *et al.* Continuous administration of poloxamer 188 reduces overload-

induced muscular atrophy in dysferlin-deficient SJL mice. *Neurosci. Res.* **72**, 181–

186 (2012).

43. De Sandre-Giovannoli, A. *et al.* Lamin a truncation in Hutchinson-Gilford progeria.

Science **300**, 2055 (2003).

44. Eriksson, M. *et al.* Recurrent de novo point mutations in lamin A cause Hutchinson-

Gilford progeria syndrome. *Nature* **423**, 293–298 (2003).

45. Mounkes, L. C., Kozlov, S., Hernandez, L., Sullivan, T. & Stewart, C. L. A progeroid

syndrome in mice is caused by defects in A-type lamins. *Nature* **423**, 298–301

(2003).

46. Singla, A. *et al.* Lamin aggregation is an early sensor of porphyria-induced liver

injury. *J. Cell Sci.* **126**, 3105–3112 (2013).

47. Bank, E. M. *et al.* A laminopathic mutation disrupting lamin filament assembly

causes disease-like phenotypes in Caenorhabditis elegans. *Mol. Biol. Cell* **22**, 2716–

2728 (2011).

48. Casasola, A. *et al.* Prelamin A processing, accumulation and distribution in normal

cells and laminopathy disorders. *Nucleus* **7**, 84–102 (2016).

49. Dialynas, G. *et al.* Myopathic Lamin Mutations Cause Reductive Stress and Activate

the Nrf2/Keap-1 Pathway. *PLOS Genet.* **11**, e1005231 (2015).

50. Hübner, S., Eam, J. E., Hübner, A. & Jans, D. A. Laminopathy-inducing lamin A

mutants can induce redistribution of lamin binding proteins into nuclear aggregates.

Exp. Cell Res. **312**, 171–183 (2006).

51. Hübner, S., Eam, J. E., Wagstaff, K. M. & Jans, D. A. Quantitative analysis of

localization and nuclear aggregate formation induced by GFP-lamin A mutant

proteins in living HeLa cells. *J. Cell. Biochem.* **98**, 810–826 (2006).

52. Ahmed, M. S., Ikram, S., Bibi, N. & Mir, A. Hutchinson-Gilford Progeria Syndrome: A

Premature Aging Disease. *Mol. Neurobiol.* **55**, 4417–4427 (2018).

53. Barascu, A. *et al.* Oxidative stress induces an ATM-independent senescence

pathway through p38 MAPK-mediated lamin B1 accumulation. *EMBO J.* **31**, 1080–

1094 (2012).

54. Zaremba-Czogalla, M. *et al.* The Different Function of Single Phosphorylation Sites

of Drosophila melanogaster Lamin Dm and Lamin C. *PLoS ONE* **7**, (2012).

55. Li, Y. *et al.* Lamin Mutations Accelerate Aging via Defective Export of Mitochondrial

mRNAs through Nuclear Envelope Budding. *Curr. Biol. CB* **26**, 2052–2059 (2016).

56. Muñoz-Alarcón, A. *et al.* Characterization of lamin Mutation Phenotypes in

Drosophila and Comparison to Human Laminopathies. *PLOS ONE* **2**, e532 (2007).

57. Kitten, G. T. & Nigg, E. A. The CaaX motif is required for isoprenylation, carboxyl

methylation, and nuclear membrane association of lamin B2. *J. Cell Biol.* **113**, 13–23

(1991).

58. Holtz, D., Tanaka, R. A., Hartwig, J. & McKeon, F. The CaaX motif of lamin A

functions in conjunction with the nuclear localization signal to target assembly to the

nuclear envelope. *Cell* **59**, 969–977 (1989).

59. Meshorer, E. & Gruenbaum, Y. Gone with the Wnt/Notch: stem cells in

laminopathies, progeria, and aging. *J. Cell Biol.* **181**, 9–13 (2008).

60. Polychronidou, M., Hellwig, A. & Grosshans, J. Farnesylated nuclear proteins

Kugelkern and lamin Dm0 affect nuclear morphology by directly interacting with the

nuclear membrane. *Mol. Biol. Cell* **21**, 3409–3420 (2010).

61. Patterson, K. *et al.* The functions of Klarsicht and nuclear lamin in developmentally

regulated nuclear migrations of photoreceptor cells in the Drosophila eye. *Mol. Biol.*

Cell **15**, 600–610 (2004).

62. Cao, K., Capell, B. C., Erdos, M. R., Djabali, K. & Collins, F. S. A lamin A protein

isoform overexpressed in Hutchinson-Gilford progeria syndrome interferes with

mitosis in progeria and normal cells. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 4949–4954

(2007).

63. Machowska, M., Piekarowicz, K. & Rzepecki, R. Regulation of lamin properties and

functions: does phosphorylation do it all? *Open Biol.* **5**, (2015).

64. Saito, A., Kawai, K., Takayama, H., Sudo, T. & Osada, H. Improvement of

photoaffinity SPR imaging platform and determination of the binding site of

p62/SQSTM1 to p38 MAP kinase. *Chem. Asian J.* **3**, 1607–1612 (2008).

65. Rana, A., Rera, M. & Walker, D. W. Parkin overexpression during aging reduces

proteotoxicity, alters mitochondrial dynamics, and extends lifespan. *Proc. Natl. Acad.*

Sci. U. S. A. **110**, 8638–8643 (2013).

66. Burke, B. & Stewart, C. L. The nuclear lamins: flexibility in function. *Nat. Rev. Mol.*

Cell Biol. **14**, 13–24 (2013).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6