Cryptococcus neoformans **resist to drastic conditions by switching to viable but non-**

culturable cell phenotype

Benjamin Hommel[1,2,3*], Aude Sturny-Leclère[1*], Stevenn Volant[4], Nathanael Veluppillai[1],

Magalie Duchateau[5], Chen-Hsin Yu[6], Véronique Hourdel[5], Hugo Varet[4,7], Mariette Matondo[5],

John R Perfect[6], Arturo Casadevall[8], Françoise Dromer[1] & Alexandre Alanio[1,2,3,8]

[*] These authors participated equally to this work.

[1] Institut Pasteur, Molecular Mycology Unit, CNRS UMR2000, Paris, France

[2] Laboratoire de Parasitologie-Mycologie, Hôpital Saint-Louis, Groupe Hospitalier

Lariboisière, Saint-Louis, Fernand Widal, Assistance Publique-Hôpitaux de Paris (AP-HP),

Paris, France

[3] Université Paris Diderot, Sorbonne Paris Cité, Université de Paris, Paris, France

[4] Institut Pasteur - Bioinformatics and Biostatistics Hub - C3BI, USR 3756 IP CNRS, Paris,

France

[5] Institut Pasteur, Unité de spectrométrie de masse et Protéomique, Paris, France.

[6] Division of Infectious Diseases, Department of Medicine, Duke University Medical Center,

Durham, North Carolina, United States of America

[7] Institut Pasteur - Transcriptome and Epigenome Platform - Biomics Pole - C2RT, Paris,

France

[8] Department of Molecular Microbiology and Immunology, Johns Hopkins Bloomberg School

of Public Health, Baltimore, MD, USA

*Corresponding Author: Dr Alexandre Alanio, Molecular Mycology unit, Institut Pasteur, 25

rue du Dr Roux 75724 Paris Cedex 15; email: <u>alexandre.alanio@pasteur.fr</u>; Tel:

+33140613255; Fax: +33145688420

Keywords

Cryptococcus neoformans, viable but non culturable, reactivation, dormancy, growth,

latency, starvation, hypoxia, secretome, proteome, transcriptome, pantothenic acid, quorum

sensing

Abstract

31
Metabolically quiescent pathogens can persist in a viable non-replicating state for months or even

years. For certain infectious diseases, such as tuberculosis, cryptococcosis, histoplasmosis, latent

infection is a corollary of this dormant state, which has the risk for reactivation and clinical disease.

During murine cryptococcosis and macrophage uptake, stress and host immunity induce *C.*

neoformans heterogeneity with the generation of a sub-population of yeasts that manifests a

phenotype compatible with dormancy (low stress response, latency of growth). In this

subpopulation, mitochondrial transcriptional activity is regulated and this phenotype has been

considered as a hallmark of quiescence in stem cells. Based on these findings, we worked to

reproduce this phenotype in vitro and then standardize the experimental conditions to consistently

generate this dormancy in *Cryptococcus neoformans*.

We found that incubation of stationary phase yeasts (STAT) in nutriment limited conditions and

hypoxia for 8 days (8D-HYPOx) was able to produced cells that mimic the phenotype obtained in vivo.

In these conditions, mortality and/or apoptosis occurred in less than 5% of the yeasts compared to

30-40% of apoptotic or dead yeasts upon incubation in normoxia (8D-NORMOx). Yeasts in 8D-HYPOx

harbored a lower stress response, delayed growth and less that 1% of culturability on agar plates,

suggesting that these yeasts are viable but non culturable cells (VBNC). These VBNC were able to

reactivate in the presence of pantothenic acid, a vitamin that is known to be involved in quorum

sensing and a precursor of acetyl-CoA. Global metabolism of 8D-HYPOx cells showed some specific

requirements and was globally shut down compared to 8D-NORMOx and STAT conditions.

Mitochondrial analyses showed that the mitochondrial masse increased with mitochondria mostly

depolarized in 8D-HYPOx compared to 8D-NORMox, with increased expression of mitochondrial

genes. Proteomic and transcriptomic analyses of 8D-HYPOx revealed that the number of secreted

proteins and transcripts detected also decreased compared to 8D-NORMOx and STAT, and the

proteome, secretome and transcriptome harbored specific profiles that are engaged as soon as four

days of incubation. Importantly, acetyl-CoA and the fatty acid pathway involving mitochondria are

required for the generation and viability maintenance of VBNC.

All together, these data show that we were able to generate for the first time VBNC phenotype in

Cryptococcocus neoformans. This VBNC state is associated with a specific metabolism that should be

further studied to understand dormancy/quiescence in this yeast.

Introduction

All microorganisms are exposed to fluctuating environments and periodic external stresses, which inhibit their growth. Some microbes may overcome these stress periods by entering into a resistant non-replicative state, assessed by /associated with a decreased culturability (1). This decrease was first assumed to reflect microbial death as a consequence of a stochastic and induced nutrient deprivation decay (2). In quantum mechanics, the Schrödinger's experiment considered a cat simultaneously alive and dead until a measurement was made. A parallel experiment could be done with a dormant microorganism. Answering if it is alive or dead seems trivial but is unquestionably uneasy (3). In fact, there are two other alternative explanations. Non-culturable cells could be the resulting state of a genetically programmed death or could be viable but non culturable cells (VBNC) that are adapted to living / surviving but dormant cells like spores (4, 5). There is now clear evidence that VBNC are viable cells with an intact cell membrane and a low metabolic activity (6). Various environmental stressors can induce VBNC, such as starvation, hypoxia, stressful temperature, pH variations (7). In a finalistic approach, to enable perpetuation of the lineage, VBNC should be capable at some point to exit the dormant state to resume an active and replicating form when environmental conditions improve. This process could be described as reactivation or "resuscitation". At least 85 species of bacteria (8) and 10 species of fungi (9–11) have been reported capable of adapting a VBNC state. Hence, the VBNC state seems to be an intrinsic capability of many microorganisms including pathogenic ones (12–15). During infection, the "dormant" pathogens are able to tolerate both immune system attacks and prolonged exposure to antimicrobials (16). Pathogens are not known to cause disease when present in the VBNC state but virulence is retained and infection can be initiated again following reactivation (6).

Dormancy is an important pathogenic feature for some invasive fungal diseases, such as

cryptococcosis (17–19) and histoplasmosis (12), where dormant organisms are known to

reactivate years after primary infection. Cryptococcosis is caused by the ubiquitous

environmental basidiomycete yeast *Cryptococcus neoformans*. It occurs mainly in

immunocompromised individuals, and especially those with AIDS, with more than 200 000

new cases of meningoencephalitis and more than 180, 000 deaths per year worldwide (20).

Latency is an important phase of cryptococcosis pathogenesis. However, the "primary

complex" of infections has been rarely interrogated in contrast to tuberculosis (21).

Inhalation of infectious propagules leads to a primary pulmonary infection with a

granulomatous immune response (22) that either restrains infection or eradicates the yeast.

This immune response can fail when immunosuppression occurs leading to infection

reactivation and disease. Recently, evidence was reported for the existence of dormant

yeast cells of *C. neoformans* in models of *C. neoformans*/host interaction *in vivo* (mice) and

in vitro (macrophages) have been obtained (23). A population of dormant yeast cells

harboring characteristics that could be related to dormancy (low metabolic response,

growth latency, increased mitochondrial activity, increased autophagy, decreased

neoglucogenesis and reactivation by fetal calf serum) was described (23). To characterize

these cells, we needed reproducible *in vitro* conditions that would generate yeast cells with

a phenotype identical to that observed in the subpopulations of yeasts upon interaction with

hosts. Based on previous studies performed in skeletal muscle stem cells (24), we developed

an *in vitro* model using two consecutives stresses, i.e. nutrient starvation and hypoxia, and

used them to analyze the parameters influencing dormancy and reactivation of dormant

yeasts. Using phenotypic microarrays along with, proteome, secretome, and transcriptome

analyses, we further validated and characterized these yeast cells and their metabolic

pathways that are crucial for the generation, maintenance of and exit from this dormancy

state.

Results

Yeasts incubated in hypoxia were intact and had low metabolic activity and delayed

growth

Based on our previous identification of dormant cells in a mouse model of cryptococcosis

(23), we identified *in vitro* conditions that induced a phenotype in *C. neoformans* cells that

mimicked the phenotype observed *in vivo*. After defined pre-culture conditions, yeasts in

YPD liquid medium (stationary phase, STAT) were incubated in nutrient deprivation, hypoxia

(O_2<0.1%) and increased CO_2 concentration) at 30°C for 8 days (all subsequent will be

referred as 8D-HYPOx) (Supplementary Figure 1). Yeasts in hypoxia were then compared to

yeasts cultured in parallel in normoxia (referred as 8D-NORMOx), and to STAT.

Median body cell sizes were not different for yeasts in 8D-HYPOx, 8D-NORMOx compared to

STAT (4.5 µm; interquartile range (IQR) [3.8-5.1], 4.6 µm [4.1-5.0] and 4.6 µm [4.1-5.0],

respectively). Median capsule size was smaller in 8D-HYPOx (0.8 µm [IQR=0.7-1.0] and 8D-

NORMOx (0.9 µm [0.7-1.0]) compared to STAT, (1.0 µm [0.9-1.2]), (Supplementary Figure 2A

and B, p<0.0001). Specific staining revealed no differences in the shape of the nucleus

(Figure 1A), nor on the structure of the capsule based on E1 binding (Figure 1A,

Supplementary Figure 3A) but did show an increased quantity of vacuolar membrane (3.1 ±

0.4 increase in the geometric mean fluorescence 8D-HYPOx/8D-NORMOx cells compared to

STAT (Figure 1A, Supplementary Figure 3B). Plasma membrane was intact for the majority

(88%) of the cells in 8D-HYPOx, and for 60% of them in 8D-NORMOx compared to 98% in

STAT (Figure 1A, Supplementary Figure 3C). Transmission electron microscopy (TEM)

revealed a significantly thicker cell wall in 8D-HYPOx cells compared to STAT (median 206 nm

[170.5-245.0] and 163.5 nm [123.8-199.3], p<0.05, Supplementary Figure 2C) and the

presence of large intracytoplasmic vacuoles (Figure 1B).

Intracellular levels of glutathione as a marker of stress response in *C. neoformans* (23) was

assessed kinetically by CMFDA staining at day 0 (STAT) and 4, 6 and 8 d (D4, D6, D8). A

decrease in stress response was observed over time with an apparently homogenous

population harboring a low CMFDA intensity in 8D-HYPOx, as opposed to 8D-NORMOx

condition where a double population (low and high CMFDA intensity corresponding to low

and high stress response) was observed at D8 (Figure 1C). Growth curves in rich medium

(YPD) were determined at the same time points using mathematical modeling, with latency

(lag phase λ) as the major readout (Figure 1D). Median latency was significantly increased for

8D-NORMOx and 8D-HYPOx populations compared to STAT (1103 m [1083-1104] with a

continuous increase over time for yeast cells grown in 8D-HYPOx from 1387 m [1281-1445]

at D4, to 1878 m [1800-1898] at D8 whereas it plateaued in 8D-NORMOx (1194 m [1176-

1210] at D4, and 1288 m [960-1351] at 8D (Figure 1E).

Yeasts incubated in hypoxia were mostly viable but non-culturable cells (VBNC)

Cell viability and yeast apoptosis-like phenomenon measured over time evolved differently

in hypoxia and normoxia (Figure 2A). Viable non-apoptotic cells represented 98.7% [98.4-

99.0]) in the STAT condition, and almost the same proportion (98.9% [98.8-99.0]) at 8D-

HYPOx with a decrease at D2 D4 and D6, culminating at 74.6% [70.5-78.8] at D4. In

normoxia, that proportion decreased and plateaued at D8 at 63.6% [62.2-65.0]. The

proportion of apoptotic-like cells, negligible in STAT cells (0.8% [0.5-1.6]), peaked at D4 in

hypoxia (21.4% [17.5-25.4]) while it increased over time to reach a plateau at D6 in normoxia

(35.8% [34.9-36.7]). Dead cells represented less than 3% over time in hypoxia (2.7% [1.7-3.7]

at D4), while it reached 5% [4.2-5.9] at D6 in normoxia and was less than 1 % in STAT.

Culturability (proportion of culturable yeasts) assessed over time by colony forming unit

(CFU) on YPD agar was decreased in both conditions but significantly more in 8D-HYPOx than

in 8D-NORMOx (17.5% [15.4-19.7], vs. 0.8% [0.5-1.0] p<0.01) (Figure 2B)

The population of cells observed at 8D-HYPOx was heterogeneous in terms of CMFDA

staining, a small quantity harbored a high CMFDA staining (high stress response). We thus

hypothesized that they were the metabolically active cells responsible for the observed

growth. Consequently, yeast populations were sorted according to their CMFDA intensity

(low, medium and high). The latency of the three populations was comparable to that of the

unsorted (bulk) population and significantly higher than in the 8D-NORMOx and STAT

conditions (Figure 2C). Likewise, culturability assessed by plating the cells suspension on

Sabouraud agar plates was not significantly different in CMFDA^{low} (0.19±0.20), CMFDA^{medium}

(0.28±0.17), CMFDA^{high} (0.10±0.11) compared to the unsorted population (0.53±0.41)

(p>0.01), although significantly lower than in 8D-NORMOx (12.73% [7.62-18.62]) and STAT

(102.5% [93.25-111.8]) conditions (Figure 2C, p<0.01). Culturability assessed by counting the

proportion of culture-positive wells after 5 d of incubation for 3333 cells per well showed

also a decrease in culturability with 10 positive wells/18 seeded (55.5%) for the CMFDA^{low},

16/18 (88.8%) for the CMFDA^{medium}, 14/18 (77.7%) for the CMFDA^{high}, and 17/18 (94.4%) for

the unsorted populations whereas all wells (100%) grew in the 8D-NORMOx (6/6) and STAT

(3/3) conditions.

These results suggested that the combination of hypoxia and nutrient depletion led to the

production of a population mostly composed of what is called viable and non-culturable cells

(VBNC). A model for the evolution of the cell population phenotypes over time in hypoxia

and normoxia is proposed in Figure 2D.

Factors influencing latency

Latency of growth can be influenced by numerous factors. We first tested cell concentration

and culture medium. Serial 10-fold dilutions of the cells ($3x10^5$ to 3/well) resulted in an

increased latency for both 8D-HYPOx and STAT cells, in YPD and in minimal medium (MM)

(Supplementary Figure 4A-B). The latency was higher in MM than in YPD for 8D-HYPOx at all

the dilutions studied (Table 1) and for STAT at all dilutions except for $3x10^3$/well. Nonlinear

fitting permitted the extrapolation of the best fit curve and showed that a low nutrient

medium (MM) increased the latency by a differential of 2505 m for STAT and 2656 m for 8D-

HYPOx compared to a richer medium (YPD). There was almost no difference in latency for 1

cell between STAT and 8D-HYPOx: for 1 cell the latency (8D-HYPOx -STAT) was 1820 m for

YPD and 1971 m for MM.

8D-HYPOx can be reactivated by pantothenic acid (PA), the acetyl CoA precursor.

We hypothesized that the latency observed for the 8D-HYPOx population could be impacted

by the quantity of nutrients available in the medium, a parameter that could be simulated by

dilution of MM in water. MM dilutions resulted in a significant decrease in 8D-HYPOx latency

from 4463 m [3947-5488] in 100%MM to 3722 m [3612-4156] in 20%MM and 3358 m [3240-

3664] in 10%MM (p<0.01, Figure 3A). This decrease was also observed for STAT but to a

lesser extent (1593 m [1397-1720], 1326 m [1220-1510] and 1245 m [1045-1277] in

100%MM 20%MM and 10%MM, respectively, p <0.01).

One of the major characteristics of viable but not culturable cells (VBNC) is that they can be

reactivated under various conditions which should translate into decrease of latency. Thus,

we tested the effect on 8D-HYPOx of conditioned medium (freeze-dried and rehydrated MM

from STAT populations), murine macrophage cell lysates (J774 cell line), pantothenic acid

(PA, vitamin B5), thiamine (vitamin B1), sodium pyruvate, and L-DOPA. Latency decreased

only upon addition of J774 lysate, conditioned medium and PA. As conditioned medium and

J774 lysate are each complex media rich in PA, we decided to focus specifically on the effect

of PA, which was previously implicated in quorum sensing effects (25). Addition of PA to

100%MM showed a dramatic effect on the latency of 8D-HYPOx. For instance, latency was

decreased at concentrations of PA above 1.25×10^{-2} µM from 3994 m [3797-4332] (no PA)

and 4111 m [3866-4451] (1.25×10^{-4} µM) to 2213 m [1963-2426] for (1.25×10^{-2} µM), 2095 m

[2007-2170] (1.25 µM) and 2133 m [2035-2281] (125 µM), respectively (Figure 3B, p <0.01).

In comparison, less effect of PA was observed on STAT cells (from 1615 m [1532-1637] (no

PA) to 1433 m [1398-1456] (1.25×10^{-4} µM) and then 1036 m [1017-1045] (1.25×10^{-2} µM),

1048 m [1017-1045] (1.25 µM), 1018 m [982.5-1031] (125 µM)).

We compared the growth curves to investigate if a change in growth rate was induced by PA

or diluted MM. An increased slope would correspond to an increased growth rate, meaning

an increased speed of cell generation during exponential phase. MM dilutions (20%MM and

10%MM) significantly decreased growth rate of hypoxia cells compared to 100%MM but did

not alter STAT cells (Figure 3C). PA significantly increased growth rate of 8D-HYPOx at

concentrations above 1.25 µM and of STAT cells at concentrations above 1.25×10^{-2} µM

(Figure 3D, p < 0.01).

To understand whether the reactivation observed was due to an increased number of

growing cells or to an improved metabolic adaptation in 8D-HYPOx, limit dilutions

experiments were performed calculating the probability for one cell to grow (Supplementary

Figure 4C). Dilution of MM alone (10%MM) did not modify the median proportion of

culturable cells. Addition of PA increased significantly the median proportion of culturable

cells from 1.3 ‰ [0.8-1.9] to 2.5 ‰ [2.0-3.3] in 100%MM and from 1.1 ‰ [0.9-1, 4] to 2.5 ‰

[2.1-2.7] in 10%MM. (p <0.01, Figure 3E). Culturability of yeasts did not change upon

interaction with macrophages with a comparable median proportion of culturable yeast cells

in 8D-HYPOx before (1.04 ‰ [0.78-1.48]) and after 2 h macrophage uptake (1.21 ‰ [1.09-

1.33], p=0.35; Supplementary fFgure 4D).

Overall, when decreasing nutrient availability (diluting medium), latency decreased without

change in the number of culturable yeast cells, suggesting that culturable yeast cells were

better adapted to grow in poor than in richer medium. By contrast, addition of PA allowed

VBNC to switch to culturable cells (cell reactivation) with a decreased latency reflected an

increasing number of culturable cells. However, the 2-fold increase in culturability after

addition of PA was not sufficient to explain the extent of the decrease in latency suggesting

that PA influences also growth adaptation (Figure 3F). From now on, 8D-HYPOx yeast cells

are called VBNC for convenience.

VBNC showed specific metabolic requirements and general low metabolic activity

To explore the VBNC phenotype further, their metabolic profile was obtained using Biolog®

substrate utilization tests and compared with that of yeast cells in other conditions. Based

on the 761 metabolites tested and excluding 240 compounds positive in the dead cells

control, hierarchical clustering showed that VBNC possessed a unique profile (Figure 4A).

VBNC had significant lower global metabolic activity (median= 124.0 [94.0-173] and 87.0

[65.0-118.8]) compared yeast cells in 8D-NORMOx (246.0 [106.8-260.0] and 248.5 [123.0-

262.0] for the two biological replicates) and to STAT (262.0 [136.5-272.0]) ($p<0.01$, Figure

4B). Finally, we identified two compounds (trimetaphosphate; Adenosine-2', 3' cyclic

monophosphate) that were more metabolized by VBNC than by control yeast cells

(Supplementary Table 1).

VBNC have increased mitochondrial mass and depolarized mitochondria

Since the phenotypic microarrays based on mitochondrial activity measurement showed a

decrease in VBNC (8D-HYPOx), we tested further various mitochondrial parameters. ROS and

RNS production were slightly decreased in VBNC compared to STAT cells with median 152.3

[148.1-156.2] vs. 174.1 [168.9-177.8] using the ROS probe ($p<0.01$) and 87.1 [78.2-101.2] vs.

149.5 [98.7-206.5] using the RNS probe, respectively ($p<0.01$) (Supplementary Figure 5).

Using multispectral flow cytometry, we observed an increase in the mitochondrial mass in

VBNC and in 8D-NORMOx cells compared to STAT and dead yeast cells using Mitotracker

staining. JC-1 and TMRE staining revealed that the mitochondria in VBNC are mostly

depolarized (low TMRE and high JC-1 fluorescence) (Figure 5A-B). The expression of the

mitochondrial genes CYTb, NADH, mtLSU and COX1 significantly increased in 8D-HYPOx

compared to STAT ($p=0.029$) (Figure 5C), and DNA quantification remained similar ($p<0.05$).

VBNC have a reduced but a specific secretome and a specific proteome

Secreted protein in the supernatant (secretome) and cellular proteins (proteome) were

analyzed in STAT, VBNC (8D-HYPOx) and 8D-NORMOx using both qualitative and quantitative

analysis of the proteins recovered in triplicate.

In VBNC, a total of 1365 proteins were identified within the secretome and 3772 proteins

within the proteome. In the secretome, the median number of secreted proteins identified

(starting from 747 proteins [598-797] in STAT) decreased up to (411 [406-460]) in VBNC

compared to 8D-NORMOx (910 [909-928]) (Supplementary Figure 6A, p <0.001). The protein

concentrations in the secretome remained stable in VBNC and slightly increased in 8D-

NORMOx yeast cells (Supplementary Figure 6B, Supplementary Table 2). In the proteome,

the median number of proteins identified (starting from 2975 proteins [2951-3029] for STAT)

was stable over time in hypoxia and normoxia (3057 [3033-3171] in VBNC and 2968 [2960-

2976] in 8D-NORMOx Supplementary Figure 6C). The protein concentration in the proteome

(0.005 mg/mL [0.002-0.011] in STAT) tended to increase for both yeast cells reaching 0.017

mg/mL [0.008-0.018] for VBNC and 0.021 mg/mL [0.016—0.027] for control yeast cells)

(Supplementary Figure 6D).

We then analyzed the proteins that differed between STAT and VBNC/8D-NORMOx in the

secretome and the proteome. Overall, 17/ 1365 proteins present in the secretome, and

107/3772 present in the proteome were specifically found only in VBNC cells

(Supplementary Figure 7A-7B and Supplementary Table 3).

When secreted proteins and cellular proteins were pooled, the analysis found 226 proteins

present in all six conditions (Figure 6A). A total of 1654 proteins were found exclusively in

the proteome common to the 3 conditions, or exclusively in STAT (n=150), in control (n=75)

or in VBNC (n=103) (Figure 6A). A total of 17 proteins were detected exclusively in the

secretome common to the 3 conditions with 6 exclusively in STAT and hypoxia (but

different), and 16 exclusively in normoxia (Figure 6A, Supplementary Table 4).

For the subsequent analysis, the whole kinetics in hypoxia and normoxia was analyzed

(STAT, hypoxia and normoxia (D4, D6 and D8)). GO enrichment analysis of the proteins

present in in the secretome showed differences in the biological processes, molecular

functions and molecular components, whereas similar patterns and distributions were

observed for the proteome. Thus, in the secretome of hypoxia (D4, D6 and VBNC: D8) the

major enriched biological processes were translation and metabolic processing of

carbohydrates and in Normoxia (D4, D6 and D8), the major enriched biological processes

were related to oxidation-reduction mechanisms with no major change over time (Figure

6B). The enriched molecular functions and components were also different (supplementary

Figure 8A and B and Supplementary Table 5). For instance, in the proteome, in both hypoxia

and normoxia kinetic, the translation, oxidation-reduction biological processes (Figure 6C),

the nucleotide binding transferase activity and the catalytic activity for molecular functions

and the cytoplasm, ribosome, ribonucleoprotein and intracellular components were

significantly enriched (Supplementary Figure 8C and D and Supplementary Table 5).

Principal coordinate analysis (PCA) analysis of the global data revealed that the distance

between conditions for the secretome (Figure 7A) and the proteome (Figure 7C evolved

from STAT differently in hypoxia and normoxia conditions. An independent and specific

pattern was seen as soon as day 4 in hypoxia and normoxia. We then selected the proteins

present in all conditions and analyzed how the level of each protein evolved over time.

Overall, 85 secreted proteins were differentially produced, including 49 under normoxia and

36 in hypoxia (Supplementary Table 6) and 63 cellular proteins were differentially produced

including 47 significantly more produced in normoxia and 16 in hypoxia (Supplementary

Table 6). Based on these results, hierarchical clustering of the secretome (Figure 7B) and the

proteome (Figure 7D) was performed. It demonstrated that STAT specimens clustered

together with the normoxia specimens for the secretome, whereas they clustered with

hypoxia specimens for the proteome.

Among the 16 cellular proteins overproduced during hypoxia, one was a serine

carboxypeptidase (CNAG_06640); one is associated with intracellular transport

(CNAG_07846); 6 were associated with the processing of DNA / RNA and the cell-cycle

(CNAG_01455, CNAG_04570, CNAG_00819, CNAG_06079, CNAG_05311 and CNAG_04962),

and the 9th protein is linked to the mTOR pathway (CNAG_01148, also known as FK506-

binding protein) (Table 2). The other 7 proteins have no known roles (hypothetical proteins).

Cellular proteins overproduced in normoxia were subsequently, down produced in hypoxia.

These 47 proteins down produced in hypoxia were mostly proteins from the fatty acid

degradation pathway, the glyoxylate cycle: isocitrate lyase (CNAG_05303), malate synthase

(CNAG_05653) and a protein of the neoglucogenesis, the phophoenolpyruvate

carboxykinase (CNAG_04217).

Fatty acid pathway is required in VBNC

STRING analysis was used to visualize the predicted protein-protein interactions for the 63

cellular proteins differentially produced by VBNC (Figure 8A). Several metabolic pathways

were identified, but especially the fatty acids metabolic pathways: degradation and

metabolism of fatty acids, degradation of α-ketoacids (valine, leucine and iso-leucine),

metabolism of alpha-linolenic acid, beta-alanine and elongation of fatty acids as well as

peroxisomes. To examine whether VBNC stored fatty acids better than STAT or 8D-Normox,

Bodipy®. 505/515 that stains neutral lipids in lipids droplets was used but no difference was

observed (Figure 8B). The β oxidation pathway described in the KEGG database was used to

visualize the 63 cellular proteins differentially produced in VBNC and the 8 proteins for fatty

acid degradation metabolism (Figure 8C). The fatty acid degradation pathway (β oxidation) is

well-characterized in *Saccharomyces cerevisiae* with several genes found also in *C.*

neoformans genome (Figure 8D). Interestingly, the corresponding proteins in *C. neoformans*

were known to be under produced in VBNC. The 9 deletion mutants of proteins identified

from the KEGG database and by the orthology with *Saccharomyces* genome available in the

Madhani collection were tested in the model for their ability to generate VBNC. A significant

decreased in the proportion of living cell was measured for 04688Δ, 00490Δ, 07747Δ and

00524Δ, all directly involved in fatty acid degradation (Figure 8E, p<0.01), suggesting that the

fatty acid pathway is required in VBNC although the level of expression is lower than in STAT

cells.

VBNC cells have reduced but specific transcriptome

We then compared genes expression in the three conditions using transcriptome analysis. To

avoid bias related to decreased RNA content and not in gene expression, we performed a

specific experiment in which we spiked each sample with the same quantity of *S. cerevisiae*

prior to extraction allowing us to normalize the transcription of *C. neoformans* to that of *S.*

cerevisiae. The number of transcript was decreased in VBNC, and to a lesser extent in 8D-

NORMOx, compared to STAT and LOG conditions that harbored similar level of transcripts

(Figure 9A).

PCA analysis of the global transcriptomic data from non-spiked samples suggested an

evolution from LOG to STAT phases with a clear independent and specific route leading to

VBNC or control yeast cells (Figure 9B). GO enrichment analysis of upregulated transcripts

showed that the major biological processes involved were: signal transduction, ATP synthesis

coupled proton transport and regulation of ARF protein signal transduction in VBNC,

oxidation-reduction, protein folding and response to stress in control and oxidation-

reduction and translation in LOG (Figure 9C). Further analysis revealed the clustering into 7

clusters that clearly separate the transcriptional profile of the 4 experimental conditions

(LOG, STAT, VBNC, 8d-NORMOx, Figure 9D). Specifically, cluster 4 appeared to be composed

of genes upregulated in hypoxia compared to the other conditions which underlying

pathways are summarized in Table 3 and Supplementary Table 7.

Discussion

We report the existence of a *C. neoformans* phenotype whereby dormant cells *in vitro* are viable but not culturable cells (VNBC). Hence, *C. neoformans* joins the ranks of microbes where similar phenomena have been described (6, 8, 10, 12–16) with the proviso that for this organism this phenotype has obvious clinical importance given that latency is a common manifestation of human infection.

Latency of cryptococcosis that can be linked with dormancy of *C. neoformans* had been characterized clinically (26, 27) and epidemiologically (17, 18). This population of dormant yeasts is thought to be located in a pulmonary granuloma but other reservoirs could exist (23, 28). Recently, evidence for the existence of dormancy in a sub population of *C. neoformans* upon interaction with the host *in vivo* (murine infection) and with macrophages *in vitro* (23). To obtain homogenous population of dormant an *in vitro* protocol was set up here using stresses known to generate quiescence: nutrient starvation after a first culture in rich liquid culture medium and hypoxia (29–31). After eight days in hypoxia, the cells (8D-HYPOx) were alive with a preserved structure but with more vacuoles than stationary phase (STAT) cells. Then, when assessed for an apoptosis-like phenotype (32), viability and culturability, we concluded that 8D-HYPOx condition was able to generate a homogeneous population of viable but non-culturable cells (VBNC) with <1% apoptotic, > 99% viable and <0.01% cultural cells. This phenotype is associated with dormancy in other microbial populations (6–8, 10, 33). In contrast, 8D-NORMOx yeast population were very heterogeneous with an increased death rate and apoptosis and higher culturability. We observed that hypoxia turned almost the entire *C. neoformans* population into VBNC phenotype starting after 4 days of hypoxia suggesting that hypoxia was a protective factor against apoptosis. Hypoxia is an important factor to induce dormancy of mammalian stem

cells (34). Hypoxia is also present to some extend in the granuloma (35–37). In stem cells,

hypoxia is required for a strict metabolic regulation to maintain long-term quiescence and

self-renewal of populations. Hypoxia is also known to influence cell fate (differentiation, de-

differentiation) within the stem cell niche (38, 39). In addition, resistance to apoptosis is

known in mammalian cells by the induction of apoptosis inhibitors such as IAP-2 (40, 41) but

opposite results have been found with tumor cells (42). Nevertheless, by using these in vitro

conditions to generate what happened to be VBNC, we acknowledge that the phenotype

obtained could be different from the sub-population of dormant yeasts identified from the

lung of infected mice in our lab (23), although similarities are observed (low stress response

through CMFDA decreased fluorescence, latency of growth, increased mitochondrial

transcription). One important additional point is that there is technical limitation to work

with dormant yeasts from murine infection because of the low number of yeasts present in

the lung (10^4 cells recovered from the sacrifice of 42 animals) and the necessity to sort them

and by doing so impact their phenotype (23).

In VBNC-inducing conditions, the yeast population experienced an increased latency (lag

phase) over time. We investigated by different means if this delay was due to the decreased

number of cells able to grow or the decreased metabolic activity capacity. We validated in C.

neoformans that the difference in latency depended on nutrient availability (difference

between poor (MM) and rich (YPD) media), and on the number of cells able to grow

(dilutions of STAT yeasts). In addition, we showed that the metabolic activity of the cells was

also an important factor that participated in latency.

One of the major characteristics of cells manifesting the VBNC phenotype is their ability to

be reactivated - to divide again - in response to specific stimuli (7, 43, 44). Indeed, the final

demonstration of the existence of the VBNC phenotype came with the identification of

conditions that allowed reactivation. Among them, YPD-conditioned media, murine

macrophage lysate and PA had the greatest impact on latency under our conditions. YPD

medium and murine macrophage lysate are in fact complex media that also contained PA

(45), so we focused on PA alone. We hypothesized that PA was an important factor in

reactivating VBNC based on the fact that PA was known to decrease latency in *C.*

neoformans cells (46) and was also a known beneficial secreted compound by *C. neoformans*

(25). By adding PA to culture medium it decreased latency of VBNC but had less effect on

STAT.

We observed that MM dilutions increased growth capacities of VBNC (decreased latency). To

explain this counterintuitive observation, we propose that exposure to a high amount of

nutrients after long-term starvation was deleterious for yeast cells that had to resume

normal metabolism, possibly as a result of oxidative damage as metabolism increases, and

this hypothesis is further based on the fact that death was shown by others to be

accelerated in rich substrate (47–49).

Given that the critical issue in an reactivation experiments is to distinguish between an

increased number of cells from reactivation of the VBNC state from faster growth of already

present culturable cells, a method was developed to measure the probability of growth for

one yeast cell, inspired by the most probable number method (50). We found that the effect

of PA and medium dilution was different on VBNC cells compared to STAT cells. The effect of

PA on VBNC was a result from the doubling number of reactivated VBNC and from an

increase in growth rate. The weight of each factor remains to be determined. The proportion

of VBNC that was reactivated was in fact low (1.2‰). It is highly probable that resuscitation

of VBNC may require other factors that still need to be discovered. In our system, the

majority of the yeast population was probably not able to reactivate as hypoxia may have

pushed the yeasts too far into the process of dormancy for them to be reactivated or

without resuscitation (51). Indeed, the VBNC population may be injured while still alive and

unable to divide again (52). On the other hand, the MM dilution did not influence VBNC

reactivation culturability but rather decreased the growth rate by a faster rewire of the

metabolic state of the remaining culturable cells.

Despite the importance of *Cryptococcus*-protozoan interaction in the explanation of its

virulence and intracellular pathogenic strategy (53), we showed that macrophage uptake of

VBNC had no effect on culturability. This result is different from what was observed for other

microorganisms. For example, reactivation of VBNC was possible for the bacterium *L.*

monocytogenes and the protozoans *Acanthamoeaba polyphaga* (54) and *A. castellanii* (55).

We first screened our dormancy phenotype through the decrease in metabolic and stress

responses (23), as measured upon staining with the CMFDA fluorescent probe (56, 57).

Although it allowed us to detect that hypoxia was an important factor for the generation of

VBNC, we observed surprisingly that the culturability of the yeasts was independent of the

CMFDA level after sorting the different CMFDA populations. This suggested that CMFDA

staining is not a perfect marker of metabolic and stress response in *C. neoformans*.

Decreased metabolic activity in VBNC was demonstrated by several strategies. First, a

phenotypic array measurement (Biolog®) allowed us to show a reduction in the metabolism

of 761 compounds in VBNC. Second, we also noticed that the concentration of secreted or

intracellular proteins was lower in VBNC compared to 8D-NORMOx as reported during

quiescence in other organisms (58–61). We observed that the number of secreted proteins

present in the supernatant of VBNC decreased over time compared to the 8D-NORMOx

control. In parallel, the increase in the number of proteins present in the supernatant of 8D-

NORMOx could correspond to increased secretion but more likely to an increased release of

proteins due to a high cellular mortality observed in that condition. Third, using a unique

strategy of analysis for the transcriptome including the spike of a definite % of Log phase *S.*

cerevisiae cells, we demonstrated a global decreased transcriptional activity in 8D-HYPOx

compared to 8D-NORMOx, LOG and STAT conditions, identifying again the lower metabolic

activity of VBNC.

In 8D-HYPOx, we found that the mitochondrial phenotype was modified compared to STAT

condition, an included increased mitochondrial mass (increased mitotracker and JC-1

fluorescence) depolarization (TMRE low and JC-1 high), assessed by imagestream X and an

increased expression of mitochondrial genes without an increase in the quantity of

mitochondrial DNA between 8D-HYPOx and STAT. This observation could explain why ROS

and RNS levels are lower in VBNC compared to STAT. This depolarized state could also reflect

activation of autophagy (62). These results are reminiscent of those obtained in the dormant

subpopulation of yeasts observed in mice 7 days after infection and in macrophages, where

COX1 and autophagy genes (*ATG9* and *VPS13*) have been shown to be upregulated

compared to other subpopulations of yeasts (23).

We identified and quantified proteins from the secretome (supernatant) and proteome (cell

pellet) over time. A qualitative proteomic analysis of secreted and cellular proteins in the

three conditions: STAT, VBNC and 8D-NORMOx showed different protein profiles, which

allowed the identification of proteins specific for hypoxia including 7 secreted and 103

cellular proteins. Overall, 1365 secreted proteins were identified. This number is greater

than previously described (63–65) but the protein extraction methods differed (gel

extraction, TCA / acetone precipitation alone) and the identification procedures and the

culture media used (YPD or MM) were different. Nevertheless, of the 191 secreted proteins

previously identified in stationary phase in YPD   (63–65), 93.7% were found in our

corresponding sample (STAT with 565 proteins identified in all of the 3 replicates). In *C.*

neoformans, the secreted molecules can take several pathways to reach the periphery of the

cell and the extracellular space. There are conventional and unconventional secretory

pathways involving secretory machinery in protein export: Sec4, Sec6, Sec14, the Golgi

apparatus, and extracellular exosome vesicles (66–68). Our analysis of the secreted proteins

within the supernatants, since the TCA precipitation step used will precipitate soluble

proteins and also those in extracellular vesicles (69). Consequently, we are unable to

precisely identify their mechanism of secretion. These studies will be performed in the but

this opens future directions for study to understand the relationship between dormancy and

extracellular vesicles. Interestingly, a secreted protein, pqp1 peptidase (CNAG_00150) was

found overexpressed in hypoxia. This peptidase cleaved the pro-qsp1 quorum sensing

peptide into qsp1 (CNAG_03012), that acts on the whole cell population and regulates

virulence (70–72). However, we did not find qsp1 in the supernatants of the culture media

and yet it is present in micromolar amounts in the publication of Homer et al. (70). It is

possible that our extraction protocol was not optimized for small peptides (11 amino acids)

or that small peptides might have been eliminated through the various washings steps. The

analysis of the total proteome under hypoxia conditions allowed the identification of 3772

cellular proteins which is more than previously reported in *C. neoformans* (64, 73–75) and is

likely due to the optimized extraction method. PCoA analyzes of the secretome and

proteome kinetics showed a specific profile in VBNC and normoxia conditions compared to

STAT. The hierarchical clustering of the secretome reinforced the finding that VBNC was the

more distant condition compared to normoxia conditions and STAT. However, proteome

hierarchical clustering showed that STAT and day 4 hypoxia were close to each other and

that normoxia proteome was clearly different. The GO enrichment analysis revealed specific

functions of the secreted proteins in hypoxia compared to normoxia, whereas the pattern

was similar in the proteome: biological processes involving translation, carbohydrate

metabolism and glycolysis were enriched in the secretion of hypoxia cells.

The quantitative proteome analysis revealed that 16 proteins were more produced in

hypoxia, including CNAG_01148 also known as FK506-binding protein which is involved in

the mTOR pathway, and has a role in quiescence in *S. cerevisiae* (76). This protein is able to

bind to rapamycin (Sirolimus®) in both yeasts and mammalian cells promoting quiescence

(77, 78). In *C. neoformans*, rapamycin induces cell cycle arrest in the G1 phase (79).

Using STRING analysis of the protein-protein interactions provided evidence that the

regulation of fatty acids was the major metabolic pathway involved in VBNC. The regulation

of fatty acids is a phenomenon that occurs in *S. cerevisiae* quiescence by participating in

energy cellular homeostasis and membrane synthesis (29). In *Vibrio vulnificus*, changes in

fatty acid membrane composition contribute to maintaining the viability of the VBNC

state(33). A common feature among the metabolic fatty acids pathways is the participation

of acetyl-CoA (80). Acetyl-CoA is more than a cofactor in cell metabolism. It is central to the

physiology of mammalian cells (fatty acid metabolism, Krebs cycle, apoptosis, cell cycle,

damage- response DNA and epigenetics) (81). In *C. neoformans*, the genes of the three

principal production routes of acetyl-CoA in the cytosol are regulated (82): the β-oxidation of

fatty acids by the enzyme Mfe2, acetate by acetate synthase Acs1 and citrate by Acl1. Our

hypothesis isthat maintenance of fatty acid metabolism in hypoxia would allow the cells to

maintain membrane integrity and viability, with an increased mortality in deletion mutants

in key enzymes of the fatty acid pathway. If these results are in favor of the indispensability

of acetyl-CoA for the maintenance of *C. neoformans* infection, the relationship between

acetyl-CoA, pantothenic acid (PA) – a precursor of acetyl-CoA and dormant cells should be

further analyzed. The fact that PA was able to resuscitate VBNC cells, confirms the potential

relevance of the link made by the proteomic study between VBNC and metabolic pathways

of acetyl-CoA. Qualitative and quantitative changes in fatty composition occurs in *C.*

neoformans depending on the growth phase, with an increase during growth progression

(83). Bodipy staining showed no clear differences but demonstrated an important amount of

lipid droplets under each condition. KEGG pathway mapping of the 63 proteins regulated in

hypoxia showed an involvement of 8 proteins involved in fatty acid degradation. The 8

proteins involved were downregulated in hypoxia compared to normoxia. Another

reconstruction of the fatty acid degradation pathway using *Saccharomyces* genome

database involved 4 proteins also downregulated in hypoxia *FAA2* (CNAG_03019), *POX1*

(CNAG_07747), *FOX2* (CNAG_05721), *POT1* (CNAG_00490). The deletion mutants of these 9

proteins involved in both reconstructed pathways were tested in the *in vitro* model. The

mutants *pox1Δ*, *pot1Δ*, *CNAG_04688Δ* and *CNAG_00524Δ* had a decreased viability in

hypoxia compared to the parental strain KN99α strengthening the role of fatty acid

degradation in hypoxia and emphasizing the importance of downregulating fatty acid

degradation in VBNC phenotype. In *Candida albicans* macrophages internalization also

results in the induction of fatty acid degradation pathways (84) and similar patterns of

expression are observed in *C. neoformans* in the lungs of infected mice (85) and in response

to stress within macrophages and amoeba (86).

In summary, we describe conditions that promote *C. neoformans* to switch to a

homogenous population of viable but non culturable cells that can be considered dormant

and use those conditions to study the characteristic of these yeasts. VBNC cells manifest

what is potentially an extreme phenotype (dormancy) derived from stationary phase

(quiescence) but demonstrate the production of a very specific secretion pattern critical to

the pathobiology observed in the mammalian hosts. Our study identified genes involved in

the emergence and maintenance of the VBNC phenotype, which implicate a major role for

lipid metabolism in the emergence and maintenance of this state. The availability of

conditions that reliably induce VBNC provides a new research tool in the field of fungal and

C. neoformans biology to study the important features of dormancy and help understand

latency in cryptococcal pathogenesis.

Material and methods

Culture media and strains

Cryptococcus neoformans strains (Supplementary Table 8) were usually grown in liquid Yeast Peptone Dextrose (YPD: 1% yeast extract (BD Difco, Le Pont de Claix, France), 2% peptone (BD Difco), 2% D-glucose (Sigma, Saint Louis, Minnesota, USA)). Minimal medium (MM, 15mM D-glucose (Sigma), 10 mM $MgSO_4$ (Sigma), 29.4mM KH_2PO_4 (Sigma), 13mM Glycine (Sigma), 3.0 µM Thiamine (Sigma), pH5.5) was used for specific experiments (25).

In vitro generation of C. neoformans VBNC

Cryptococcus neoformans H99O strain stored in 20% glycerol at -80°C was cultured on Sabouraud agar plate at room temperature (step 1). After 2 to 5 days of culture, 10^7 cells were suspended in 10 mL of YPD in a T25cm^3 flask and cultured for 22 hours at 30 °C, 150 rpm with lateral shaking until stationary phase (final concentration≈2×10^8/mL) (step 2). One hundred µL of this first culture was added to 10 mL YPD (second culture) and incubated in the same conditions until stationary phase (Step 3). Then, the flasks were placed in a GENbag® (Biomérieux) anaerobic atmosphere generator to generate hypoxia and incubated for 8 days in the dark at 30°C. These yeast cells are called 8D-HYPOx and then VBNC for convenience. Control samples consisted of yeasts in stationary phase (step3, STAT) and yeasts incubated under the same conditions but in normoxia are called 8D-NORMOx (Step 4) (supplemental Figure 1). Of note, the atmosphere in the bag is enriched in CO_2 by the biochemical process of oxygen depletion.

Transmission electron microscopy

Freshly harvested cells from STAT and 8D-HYPOx were fixed overnight at 4°C with 2.5%

glutaraldehyde in 0.1 M sodium cacodylate buffer (CB) at pH 7.2. Then samples were washed

in 0.1 M CB (pH 7.2) and post-fixed in a 1% osmium tetroxide in CB at room temperature for

1 h. We pre-embedded the cells in 4% agar type 9 : the cells were mixed with agar, spun-

down by gentle centrifugation and the pellet stored at 4°C until the agar had solidified. We

excised 2 mm^3 sections from agar block, and processed them in the automatic MicroWave

tissue processor for electron microscopy (AMW, Leica Microsystems, Vienna, Austria) as

recommended by the manufacturer: samples were washed with water and dehydrated with

graded ethanol concentrations (25–100%), followed by a mixture of graded resin Spurr Agar

low viscosity Resin (Agar scientific, Gometz la ville, France) concentration in ethanol (35-

100%). The embedded agar blocks were polymerize at 60°C for 48h. Ultrathin sections (70

nm) were performed with an ultramicrotome 'Ultracut UC7' (Leica Microsystems, Vienna,

Austria), stained with 4% uranyl acetate and Reynold's lead citrate onto 100 mesh copper

grids coated with a thin glow carbon film (S162-1, Oxford Scientific, Eynsham, UK) and then

observed under Tecnai SPIRIT (FEI-Thermofisher Company) at 120 kV accelerating voltage on

a camera EAGLE 4Kx4K (FEI-Thermofisher Company).

Stainings and dyes

Unless otherwise stated, yeasts were washed in phosphate buffered saline (PBS) twice

before and 3 times after staining. Stress response was assessed using 5-

chloromethylfluorescein diacetate (Cell tracker™ green CMFDA, Life Technologies). Yeasts

(10^6) were labeled with 3.3μM CMFDA in 500μL of PBS for 30 m at 37°C in the dark. Capsule

with E1 monoclonal antibody as described previously (87). Nuclei were stained using 4,6-

diamidino-2-phenylindole (DAPI, Invitrogen, Carlsbad). Yeasts (10^7) were fixed with 3.7%

paraformaldehyde for 30 m before staining with 0.5µg/mL of DAPI for 5 m. MDY-64

(Thermofischer Scientific, Waltham) was used to stain the intracellular vacuolar membranes

by incubating 10^7 yeasts for 5 m with 5µM MDY-64. Lipid droplets were stained using

Bodipy® 505/515 (Invitrogen, Carlsbad) at 1µg/mL with 10^7 cells for 20 m at room

temperature. To assess viability, LIVE/DEAD® Fixable Violet Blue (416 nm/451 nm) diluted at

1:1000 and 250 µL was added to 10^7 yeasts pellets and incubated at 30°C in the dark.

Controls were dead yeasts generated upon exposure to 200 mM of oxygen peroxide (H_2O_2)

at 37°C, under 650 rpm agitation or heat killed 1h at 65°C.

Yeasts were analyzed by fluorescence microscopy and pictures taken using an AxioCam MRm

camera (Carl Zeiss, Oberkochen) on interferential contrast microscope (DMLB2 microscope;

Leica, Oberkochen) with Zeiss Axiovision software.

Mitochondrial mass and polarization were evaluated with Mitotracker Green (Invitrogen) at

40 nM, JC-1 (Invitrogen) at 10µg/ml and TMRE at 200nM in PBS on 10^7 yeasts for 30 m at

37°C and then washes 3 times in PBS.

Reactive oxygen (ROS) nitrogen species (RNS) measurements were also performed.

Microplate wells were filled with 10^4 yeasts in 100 µL PBS. Briefly, the probes for ROS (2',7'-

dichlorofluorescein diacetate, Sigma) and RNS (dihydrorhodamine 123, Invitrogen) were

diluted at 100 µM in PBS and methanol, respectively, and 20 µL were added to wells of a

microtiter plate containing 10^4 yeasts in 100 µL of PBS. Hydrogen peroxide (4M) was

included as a positive control. The plates were incubated at 37°C in the dark. After 1 hour,

fluorescence was measured with a fluorometer (Fluoroskan ascent™ fluorometer, Thermos

Fischer Scientific, Waltham, MA, USA) using excitation 485 nm and emission wavelengths of

530 nm (88). The data were expressed as arbitrary units of fluorescence ±SD.

Apoptosis assay

Yeasts (10^7) were fixed for 1 h at room temperature in 3.2% formaldehyde (Sigma) in PBS.

After washings in PBS, yeasts were incubated in 1 mL of 5% 2-mercaptoethanol in SPM

buffer (Sorbitol 1.2 M (Sigma), potassium phosphate monobasic 50 mM (Sigma), magnesium

chloride hexahydrate 1 mM (Sigma), pH 7.3) for 1 hr at 37°C. Yeasts were washed once in

SPM buffer and incubated for 1h at 30°C in 1 mL of a solution containing 2 mg of Zymolyase

20T (Euromedex®), 100 mg of lysing enzymes from *Trichoderma harzianum* (Sigma) and 1 mg

of BSA (Sigma) in spheroplasting buffer (sorbitol 1 M (Sigma), sodium citrate tribasic

dihydrate 100 mM (Fluka), EDTA 10 mM (Sigma), pH 5.8). All these incubations were

performed in the dark with agitation at 650 rpm in a thermomixer. The spheroplasts were

then washed 3 times in SPM buffer-permeabilized by incubation in 1 mL of 0.1% triton X-100

(Sigma) in PBS for 2 minutes on ice and washed twice in PBS. TUNEL-Fluorescein (In Situ Cell

Death Detection Kit, Roche Life Science) was used to measure apoptosis at the single cell

level following manufacturer's recommendations. Yeasts were resuspended in 50 μL of

solution of 10% of TdT and 90% dye-coupled dUTP in PBS, incubated 1 h at 37°C and then

washed twice in PBS. Apoptosis control was generated by incubation of $3x10^7$ yeasts in 2

mM H_2O_2 for 3 h at 37°C. TdT was omitted for the negative control. In each run, yeasts

treated with DNAse (3 U/mL, Ambion®) were used as a TUNEL positive control.

Flow cytometry measurements and cell sorting

Flow cytometry was used to quantify MDY64 staining, viability and apoptosis assays, yeasts -

using the Guava easyCyte 12HT Benchtop Flow Cytometer® (Guava, MERCK, Kenilworth,

New Jersey).

Multispectral flow cytometry (ImageStreamX, Amnis Corporation) was used to quantify

mitochondrial fluorescence as described before (23, 87). For each population of interest, the

geometric mean was calculated using the IDEAS software.

In specific experiments, yeasts were sorted on a BD FACS aria™ II flow cytometer. The gating

strategy included the exclusion of the aggregates and the doublets. Subsequently, the

sorting was carried out according to the intensity of CMFDA in 3 populations (CMFDA[high],

CMFDA[medium] and CMFDA[low]).

Comparison of growth curves and statistical modelling

Growth curves measurements were done using the Bioscreen® apparatus (Oy Growth Curves

Ab Ltd). Yeasts suspensions (300 μL at 10^4/mL) were added to wells of a plate and incubated

for 2 to 5 days at 30°C with continuous, high amplitude and fast speed agitation. Optical

densities (OD) were recorded at 600nm every 20 minutes. The R software with the Grofit

plug-in (89) was used and for each replicate and in each condition, we adapted the best

mathematical model according to the AIC (Akaike Information Criterion). The two models

that best modeled the growth curves were the logistic and Richards modified, where λ

represents the length of lag phase or latency of growth, μ the growth rate, A the maximum

OD (concentration), ν the shape parameter and t the time of incubation.

$$Logistic: y(t) = \frac{A}{1 + exp^{(\frac{4\mu}{A}(\lambda - t) + 2)}}$$

$$Richards\ modified: y(t) = A\left(1 + \nu + \frac{\mu}{A}(1+\nu)^{1+1/\nu}(\lambda - t)\right)^{\frac{-1}{\nu}}$$

These formulas were written in Graphpad Prism v6.02 to allow curve extrapolation and

determination of the latency parameter λ.

Culturability and reactivation experiments

To assess basic culturability, the suspensions recovered at the end of the assay were

enumerated and 3000 cells, counted with the Guava easyCyte 12HT Benchtop Flow

Cytometer®, were plated in duplicates on YPD agar. The number of colony forming units

(CFUs) was recorded after 5 days of incubation at 30°C. The experiment was performed

twice. Results were expressed as mean percent culturability. To assess the ability of 8D-

HYPOx cells to reactivate, we determined the probability for a cell to grow by seeding, for

each condition, 576 wells (six 96-well plates) with 100 yeasts/well and counting the number

of positive wells at the end of the incubation time, taking into account the Poisson's law.

Plates were incubated 5 days at 30°C in an Infors HT® multitron pro at 150 rpm. Two

independent experiments were performed for each condition. The probability (p) of a cell to

grow was calculated using the following formula, where X represents the number of positive

wells, Y the number of wells seeded.

$$X \sim B(100, p) \quad p = 1 - (1 - \frac{X}{Y})^{\frac{1}{100}}$$

Phagocytosis assay

Interaction with macrophages was performed as described previously (90) using the J774 cell

line, the monoclonal anti-capsular polysaccharide antibody E1 as an opsonin (91), calcofluor-

stained yeasts and a yeasts: macrophages ratio of 5:1. Phagocytosis was assessed by flow

cytometry using the Guava easyCyte 12HT Benchtop Flow Cytometer® (Merck).

Phenotypic microarrays

Using phenotypic microarray technology (Biolog®, Hayward,CA) we analyzed the metabolic

capacities of yeasts incubated VBNC/8D-NORMOx (both with biological duplicates), using as

controls either STAT or dead cells (8 days in hypoxia in a YPD pH = 5) (only 3 conditions, i.e.

VBNC, 8D-NORMOx and control could be tested in each series). The Biolog® apparatus allows

quantitative measurement of the metabolism over time based on the physiological state of

the yeast cells independently of cell division and based on mitochondrial activity (proton and

electron generation) (92). Reduction of tetrazolium dye that absorbed charge produced by

cell respiration results in a purple color change that is recorded over time. The optical

density measured in the well provide information on the intensity of the corresponding

metabolic activity. The effect of carbon, nitrogen, phosphorus, and sulfur sources on the

physiological state of the cell and cell respiration was examined in response to various

osmotic conditions and to different pHs (plates PM1, PM2A, PM3B, PM4A, PM6, PM7 and

PM8). A tetrazolium dye reacting to the redox status of the well turns purple when

respiration takes place which has been shown to be a good proxy for the ability of strains to

grow on a substrate (93). Phenotype microarray was done according to instructions from the

manufacturer (Biolog, Hayward, CA) with the following exceptions. Briefly, ≈7x10^8 cells were

washed 3 times in PBS and then resuspend in 27 mL of PBS. For each condition, the analysis

was performed based on the results obtained after 6 days of incubation. Overall, 761

compounds were tested and 240 compounds that were positive in the dead cells condition

were excluded from the analysis. The heatmap and clustering of the 521 remaining

conditions were performed using R studio v1.1.456 with ggplot package. For the endpoint

comparison, only positive wells in each condition were compared (393 in STAT, 279 in VBNC

duplicate 1, 200 in VBNC duplicate 2, 454 in 8D-NORMOx duplicate 1 and 380 in 8D-

NORMOx duplicate 2).

Mitochondrial nucleic acid quantification

Cells suspensions corresponding to STAT or VBNC (1 mL each in biological triplicates) were

centrifuged at 4 000rpm for 5 m at 4°C before a flash freeze step in -80°C ethanol solution

and storage at -80°C before extraction. The RNA extraction protocol was adapted from (94).

Briefly, samples were lyophilized during 18h. RNAs were extracted in 3 successive steps: i)

extraction performed in Trizol reagent (Invitrogen TriReagent (Sigma-Aldrich, St-Louis MO

63103 USA)), ii) "separation" by adding chloroform (Invitrogen) iii) overnight isopropanol

precipitation at -20°C.

The DNA protocol extraction was a manual method based on 3 steps. Briefly, a first stage

consisted in lysing the cell pellets by beat beating with MagNA Lyser Instrument (Roche

Diagnostics; 2 rounds at 7 000 rpm for 45 s each) in a solution of 2% Triton X-100 (Sigma)

and 1% of SDS (Biosolve BV Chimie, 5555 CE Valkenswaard, The Netherlands), NaCl 100mM;

Tris 10mM pH 8.0; EDTA 1mM. Then, an extraction step in phenol/chloroform solution

(Phenol:Chloroform:isoamyl alcohol pH 8.0, 1 mM EDA (Sigma-Aldrich, St-Louis MO 63103

USA) and a precipitation step in absolute ethanol and 5M of ammonium acetate solution

were done. RNAs and DNAs were quantified using the Nanodrop Spectophotometer

(ThermoFisher, Scientific, Inc.). All extracts were stored at -80°C before analysis.

qPCR and reverse transcriptase qPCR

Briefly, mtLSU primers were designed with Primers. 3 vs 4.1.0, while all other specific

primers were designed to be compatible with the Human Universal Probe Library set (90

probes, octamer, Roche Diagnostics) (Supplementary Table 9) (23). All reverse transcriptase

qPCR assays for a given target were performed simultaneously for all samples using the

LightCycler Multiplex RNA Virus Master Kit (Roche diagnostics). Each sample was normalized

with the geometric mean of the quantification cycle (Cq) of the corresponding GAPDH and

ACT housekeeping genes expression (95) and fold changes calculated according to Pfaffl

after determining the efficiency of each PCR assay (96).

Secretome and proteome analysis

Three biological replicates were performed kinetically upon incubation D4, D6 and D8 in

hypoxia and normoxia starting from step3 (Supplementary Figure 1) corresponding to

stationary phase (STAT). The suspensions were centrifuged at 4000rpm and the

supernatants (secretome) filtrated through a 0.22 µM membrane (Millex GV, Merck

Millipore, Burlington, USA). The corresponding cell pellets (proteome) were extracted in

parallel. For each biological replicate, three technical replicates were pooled to increase the

amount of recovered proteins for the supernatants.

All samples (supernatant or pellet) were lyophilized overnight after flashed freezing in liquid

nitrogen. We then adapted a protocol established to purify a sample rich in carbohydrates

(97). The lyophilized samples were resuspended in 0.5 mL of sterile milliQ water and the

following 3 steps-protocol was done: i) the samples were precipitated at 4°C by addition of 1

mL of 10% trichloroacetic acid (TCA) in acetone (kept at-20°C) followed by incubation for 5

m. After centrifugation at 20000g for 10 minutes at -20°C, the pellet was resuspended in

100% acetone with 10mM dithiothreitol (DTT), washed twice in this solution, and air dried;

ii) the pellet was then solubilized in sodium dodecyl sulfate (SDS) extraction buffer

containing 1% SDS, 0.15 M Tris HCl pH=8.8, 0.1M DTT, 1 mM ethylene diamine tetra acetic

acid (EDTA), 2 mM phenylmethylsulfonyl fluoride (PMSF), and incubated for 1 h at room

temperature and vortexed every 20 m. A 15000g centrifugation for 10 m at room

temperature allowed recovery of the supernatant; iii) Phenol was added at equal volume to

the supernatant and the tube was vortexed for 3 m. After centrifugation at 20000 g for 5 m

at room temperature, the denser phenolic phase was recovered in the lower part. An equal

volume of washing solution (Tris HCl pH 8.0, 10 mM, 1 mM EDTA, 0.7 M sucrose) was added

and the tube vortexed for 3 m. The phenolic phase was then recovered on the upper part

and transferred to a new tube. Then, 0.1M ammonium acetate in methanol (-20°C) was

added and the mixture stored at -20 ° C for 30 m. The precipitate was pelleted by 20000g

centrifugation for 10 m at -20°C, and washed first with 0.1M ammonium acetate in methanol

and then with 100% acetone. After a last centrifugation step (20000 g for 10 m at -20°C), the

pellet was air-dried and resuspended in 350 µl of denaturing buffer (8M urea, 100 mM

TrisHCl pH 8.0). Two µL of each sample were then used for dosage with the Biorad DC®

dosing kit according to the manufacturer's recommendations. The remaining solution was

stored at -80°C until analysis.

High pressure liquid chromatography coupled to tandem mass spectrometry (nano HPLC

MS/MS)

In solution protein digestion. Samples were re-suspended in 350 µL 8 M urea/ 100 mM Tris

HCl pH 8.5 after TCA/acetone precipitation and phenol extraction. Briefly, samples were

reduced with 5 mM tris(2-carboxyethyl)phosphine (TCEP) for 30 m at room temperature and

alkylated with 10 mM iodoacetamide for 30 m at room temperature in the dark. Then,

proteins were firstly digested for 5 hrs at 37°C with 250 ng rLys-C Mass Spec Grade

(Promega, Madison, USA). Samples were then diluted 4-fold with 100 mM Tris HCl pH 8.5 to

reach a concentration of 2M urea and then re-incubated overnight at 37°C with 1 µg

Sequencing Grade Modified Trypsin (Promega, Madison, WI, USA). A second incubation with

the same amount of trypsin (5 h at 37°C) was performed to ensure a complete digestion.

Digestion was stopped by adding formic acid to 5 % final concentration and peptides were

desalted and concentrated on Sep-Pak C_{18} SPE cartridge (Waters, Milford, MA, USA)

according to manufacturer's instructions.

Mass spectrometry data analysis. Tryptic peptides were analyzed on a Q Exactive Plus

instrument (Thermo Fisher Scientific, Bremen) coupled with an EASY nLC 1000

chromatography system (Thermo Fisher Scientific). Sample was loaded on an in-house

packed 30 cm nano-HPLC column (75 µm inner diameter) with C18 resin (1.9 µm particles,

100 Å pore size, Reprosil-Pur Basic C18-HD resin, Dr. Maisch GmbH, Ammerbuch-Entringen,

Germany) and equilibrated in 98 % solvent A (H2O, 0.1 % FA) and 2 % solvent B (ACN, 0.1 %

FA). Peptides were first eluted using a 2 to 18 % gradient of solvent B during 112 m, then a

18 to 30 % gradient of solvent B during 35 m, a 30 to 45 % gradient of solvent B during 15 m

and finally a 45 to 60 % gradient of solvent B during 5 m all at 250 nL.m^{-1} flow rate. The

instrument method for the Q Exactive Plus was set up in DDA mode (Data Dependent

Acquisition). After a survey scan in the Orbitrap (resolution 70 000), the 10 most intense

precursor ions were selected for HCD fragmentation with a normalized collision energy set

up to 27. Charge state screening was enabled, and precursors with unknown charge state or

a charge state of 1 and >7 were excluded. Dynamic exclusion was enabled for 45 s.

Data processing and analysis. All data were searched using Andromeda (98)with MaxQuant

software (99, 100) version 1.5.3.8 against the H99 strain proteins database (6980 entries)

and the H99 strain mitochondria proteins database (13 entries) concatenated with usual

known mass spectrometry contaminants and reversed sequences of all entries. Andromeda

searches were performed choosing trypsin as specific enzyme with a maximum number of

two missed cleavages. Possible modifications included carbamidomethylation (Cys, fixed),

oxidation (Met, variable) and Nter acetylation (variable). The mass tolerance in MS was set

to 20 ppm for the first search then 6 ppm for the main search and 10 ppm for the MS/MS.

Maximum peptide charge was set to seven and five amino acids were required as minimum

peptide length. The "match between runs" feature was applied for all samples with a

maximal retention time window of 2 m. One unique peptide to the protein group was

required for the protein identification. A false discovery rate (FDR) cutoff of 1 % was applied

at the peptide and protein levels.

MaxLFQ, Maxquant's label-free quantification (LFQ) algorithm was used to calculate protein

intensity profiles across samples (101). An LFQ value was assigned when the protein had at

least 2 peptides, of which 1 was unique. The median reproducibility by samples for

secretomic was 87.1% [82.5-90.6] and 86.5% [84.1-88.9] for cellular proteins.

Statistical analysis of differentially expressed proteins. Proteins for which we had no value

for a given condition and time were removed. Remaining missing values were imputed using

the structured least Squares algorithm (package imp4p) (102).

To get a global overview of the data, a principal coordinate analysis (PCoA) or a principal

component analysis (PCA) was used (103). The differential analysis was done with the linear

model approach proposed by the package R limma (v3.30.13) (104). We defined a model

that included condition (hypoxia or normoxia), time (D0=STAT, D4, D6, and D8) and replicate

(1 to 3) variables as main effects and interactions between conditions and times as well as

conditions and replicates. The latter interaction was useful to model the pairing between

measurements corresponding to the same condition. The proteins differentially expressed

over time are selected according to the following criteria: change with respect to the initial

value of log2 and a p <0.05. The p-values obtained were adjusted by the method of

Benjamini-Hochberg (105).

Analysis of enriched processes for secreted proteins "Gene Ontology (GO) enrichment

analysis"

The fungifun2 v2.2.8 BETA (106) site (https://elbe.hki-jena.de/fungifun/fungifun.php) was

used by selecting for each sample of secreted proteins the corresponding cellular protein

background and to study the cellular proteins the background was the whole database. For

this analysis we selected only the proteins identified in the 3 replicates per condition. For

each analysis, the following options were selected: p value <0.01, hypergeometric

distribution test, over-representation, p-read corrected by Benjamini-Hochberg, only directly

annotated association. Hypothetical proteins, which constitute 20-30% of the annotated

proteins, were not represented.

Venn Diagram

Venn diagram was generated using Jvenn (107) (http://jvenn.toulouse.inra.fr/app/index.html) with proteins only present in the 3 replicates for each condition (secretome and proteome).

Protein/protein interactions analysis

Analysis of protein-protein interactions of intracellular proteins regulated during hypoxia. STRING v10.5 software (https://string-db.org) was used (108). As this software used the strain JEC21 annotation, the list of orthologous proteins was transferred from H99 to JEC21 with a 62/63 protein match. The nodes directly related to the input were indicated with colors while the nodes of a higher iteration/depth were indicated in white.

KEGG pathway mapping and fatty acid degradation cycle reconstruction.

KEGG pathway database (https://www.genome.jp/kegg-bin) was used to map the pathways. It was rendered with the intracellular proteins significantly and differently produced in hypoxia: 63 proteins were transferred from H99 to JEC21. Nineteen proteins belonged to general metabolic pathway and more precisely 8 to fatty acid degradation ("map" pathways were not colored, *Cryptococcus*-specific pathways were colored green, entries were colored red). Another fatty acid degradation cycle was reconstructed from yeast genome database cycle (https://www.yeastgenome.org/) and the corresponding proteins were found in *Cryptococcus neoformans* H99 genome using BLAST (https://blast.ncbi.nlm.nih.gov/Blast.cgi).

 RNAseq analysis

For transcriptome analysis, yeasts incubated in 8D-HYPOx/8D-NORMOx, as well as STAT and

logarithmic (LOG) phase yeasts were studied in triplicates for two different sets, one

containing spiked *Saccharomyces cerevisiae* organisms (w_spike) and one with *C.*

neoformans cells only (w/o_spike). The former strategy was used to obtain an internal

standard of the level of expression of the whole transcripts and adjust the level of

expression of each gene to that of the constant expression of *S. cerevisiae*. In each spiked

sample, 15% of LOG phase *S. cerevisiae* cells were added to *C. neoformans* cells. Triplicates of

yeasts suspensions in the different conditions (total of 2.55×10^8 *C. neoformans* cells) were spiked

(w_spike) or not (w/o_spike) with 4.5×10^7 cells (representing then 15% of the cells) of *S. cerevisiae*

cells (culture in exponential phase). All samples were then processed as described above in the

"*Mitochondrial nucleic acid quantification*" section.

We used 1 µg of total RNA to purify polyadenylated mRNAs and to build an RNA library,

using TruSeq Stranded mRNA Sample Prep Kit (Illumina, #RS-122-9004DOC) as

recommended by the manufacturer. Directional library was checked for concentration and

quality on RNA chips with the Bioanalyzer Agilent. More precise and accurate quantification

was performed with sensitive fluorescent-based quantitation assays ("Quant-It" assays kit

and QuBit fluorometer, Invitrogen). Samples were normalized at 2 nM concentrations,

multiplexed 4 samples per lane and then denatured at a concentration of 1 nM using 0.1 M

NaOH at room temperature. Each sample was finally loaded on the flowcell at 9 pM.

Sequencing of the 24 samples was performed on the HiSeq 2500 sequencer (Illumina) in 65

bases single-end mode.

QC, mapping, counting, and global transcriptome level analysis

Reads were cleaned of adapter sequences and low-quality sequences using cutadapt version

1.11 (Marcel Martin. Cutadapt removes adapter sequences from high-throughput

sequencing reads (109)). Only sequences at least 25 nucleotides in length were considered

for further analysis. STAR version 2.5.0a (110), with default parameters, was used for

alignment on the reference genome (PRJNA411 from NCBI Bioproject). Read counts per gene

were measured using featureCounts version 1.4.6-p3 (111) from Subreads package

(parameters: -t gene -s 0). Results are summarized using multiqc version 0.7 (112). Count

data were analyzed using R version 3.3.1 and the Bioconductor package DESeq2 version

1.12.3 (113). Normalization and dispersion estimations were performed with DESeq2 using

the default parameters but statistical tests for differential expression were performed

without applying the independent filtering algorithm.

For w/o_spike samples, a generalized linear model was set in order to test for the

differential expression between STAT, 8D-HYPOx/8D-NORMOx and LOG conditions. For each

pairwise comparison, raw p-values were adjusted for multiple testing according to the

Benjamini and Hochberg (BH) procedure and genes with an adjusted p-value lower than 0.01

were considered differentially expressed.

For w_spike samples, mean normalized counts were calculated and the rank of each gene

determined for each gene in each condition. High and low rank values corresponded to

genes with the highest and lowest expression relative to spiked RNA, respectively.

Differential expression analysis

To identify the expression differences across conditions, the read counts were filtered and

normalized using the R package "DESeq2". Genes with read counts below a threshold of 10

in all samples were filtered out. After normalization of read counts, differentially expressed

genes for each condition were identified using the generalized linear model, which perform

pairwise comparisons among each of the conditions (STAT, 8D-HYPOx, 8D-NORMOx and

LOG). Genes passing the threshold, (FDR <5%), were considered significantly differentially

expressed and were then considered to be up or down-regulated according to the direction

of the fold change. The k-means clustering analyses were performed using R tools, "kmeans"

and "heatmap.2". To discover the underlying pathways of these differentially expressed

genes, we also carried out functional annotation analysis by summarizing the data, including

known H99 gene annotation, *S. cerevisiae* gene ortholog, Gene Ontology (GO), KEGG

pathway, and pfam domain, from FungiDB database (114). GO enrichment and KEGG

pathway enrichment for these genes were then assessed by using Fisher's Exact Test with

multiple testing correction of FDR threshold at 5%. Furthermore, conserved domains of each

C. neoformans gene were assessed by using the amino acid sequence to search the NCBI

Conserved Domain Database (CDD) (115). The presence of signal peptide sites was predicted

by SignalP 4.1 (116).

Statistical analysis

The statistical tests used for the comparisons between the different conditions were t-test,

Mann Whitney, Wilcoxon tests according to the distribution of the data.

Data access

All RNA sequence data from this study have been submitted to NCBI

(https://www.ncbi.nlm.nih.gov/geo) under accession number GSE118549.

The mass spectrometry proteomics data have been deposited to the ProteomeXchange

Consortium via the PRIDE partner repository with the dataset identifier PXD012570

(https://cran.r-project.org/).

Figure legends:

Figure 1: *Cryptococcus. neoformans* incubated for 8 days in hypoxia were intact with a low

stress response and delayed growth. A. Cells recovered after 8 days in hypoxia (8D-HYPOx)

were compared to those incubated in normoxia (8D-NORMOx), to stationary phase

(STAT=D0) cells. DAPI staining (DAPI) showed intact cells with a nucleus; labeling with the E1

(IgG1) anti-capsular antibody showed similar pattern in all conditions; vacuolar staining

MDY-64 showing more intense staining in 8D-HYPOx/8D-NORMOx than STAT yeasts;

LiveDead® violet staining showing high viability in 8D-HYPOx (HK = heat killed yeasts, BF =

bright field, bar = 10μm) **B.** Transmission electron microscopy of 8D-HYPOx cells and STAT

cells showing the presence of large intracytoplasmic vacuoles (bar = 0.5 μm) **C.** CMFDA

labeling showed that 8D-HYPOx yeasts had decreased stress response over time compared

to 8D-NORMOx. **D.** Latency parameter (λ) was calculated from growth curves in the different

conditions. **E.** Latency of growth increased over time for 8D-HYPOx yeasts compared to STAT

and 8D-NORMOx cells (*p <0.01). Experiments were done in biological and technical

triplicates. Each dot represents the value of λ for each replicate. Medians and interquartile

range (IQR) are represented.

Figure 2: *Cryptococcus neoformans* incubated for 8 days in hypoxia harbor a homogenous

phenotype of viable but non-culturable cell (VBNC). A. Kinetics analysis using multispectral

flow cytometry with viability (LIVE/DEAD®) and apoptosis (TUNEL) staining showing a

homogenous population of viable yeasts after D8 in hypoxia and a heterogeneous

population in normoxia with a mixture of viable, apoptotic like and dead yeasts. The

experiment was done in triplicate. Viable non-apoptotic cells represented a median

proportion of 87.7% [84.2-91.2] at D2, 74.6% [70.5-78.8] at D4, 90.9% [89.7-92.2] at D6, and

98.9% [98.8-99.0] at D8 in hypoxia, 89.4% [88.4-90.5] at D2, 83.5% [83.1-83.9] at D4, 59.6%

[59.2-60.0] at D6, 63.6% [62.2-65.0] at D8 in normoxia and 98.7% [98.4-99.0] in STAT.

Apoptotic-like cells represented a median proportion 9.9% [6.2-13.8] at D2, 21.4% [17.5-

25.4] at D4, 7.2% [6.5-7.9] at D6, 1.1% [0.9-1.2] at D8 in hypoxia, 8.8% [7.7-10.0] at D2,

13.8% [13.6-14] at D4, 35.8% [34.9-36.7] at D6, 32.9% [31.0-34.9] at D8 in normoxia and

0.8% [0.5-1.6] in STAT cells. The dead cells median proportion was 0.5% [0.4-0.6] in STAT,

1.9% [1.8-2.0] at D2, 2.7% [1.7-3.7] at D4, 1.8% [1.6-2.0] at D6, 0% at D8 in hypoxia1.7% [1.6-

1.8] at D2 , 2.2% [1.8-2.6] at D4, 5.0% [4.2-5.9] at D6, 3.2% [3.0-3.4] at D8 in normoxia.

B. Culturability was followed over 8 days by plating yeasts on Sabouraud agar (colony

forming unit method). The proportion of culturable cells decreased significantly over time

under hypoxia and significantly more than in 8D-NORMOx. Compared to STAT condition

(110.5% [108-113]), the median culturability was (33.2% [31.1-35.2] at D2, 4.3% [4.0 -4.6] at

D4, 0.8% [0.5-1.0] at D8) in hypoxia and (28.6% [17.2-40.0] at D2, 20.3% [17.9-22.8] at D4

and 17.5% [15.4-19.7] at D8 in normoxia cells) (* p<0.01).

C. Cell sorting according to the CMFDA intensity allowed showing a latency of growth after

8D-HYPOx and was increased regardless of the stress response and comparable in the three

populations (low, medium, high) to that of the unsorted population (bulk) compared to 8D-

NORMOx and STAT: CMFDA^{low} (2520 m [2257-2831]), CMFDA^{medium} (2401 m [2276-2553])

and CMFDA^{high} (2394 m [2129-2627]) was not significantly different from unsorted cells

(2269 m [2059-2344]) (p>0.01) although significantly different from 8D-NORMOx (1427 m

[1407-1462]) and STAT (1013 m [1107-1036]) conditions (* p<0.01). Two independent

experiments were pooled. Mean culturability (% mean CFU after plating) reported below the

plots was comparable in all hypoxia populations and lower than in the other conditions. The

ratio of positive wells/seeded wells are reported above the plots for each population

recovered at the end of the growth curve measurement. **D.** Proposed model for the

evolution of yeasts phenotypes upon incubation in hypoxia or normoxia leading to the

generation of a VBNC phenotype after 8D-HYPOx.

1015

Figure 3: *Cryptococcus neoformans* incubated for 8 days in hypoxia were reactivated by

pantothenic acid (PA) and culturable cells were rewired by minimal medium (MM)

dilution.

Growth curves were determined for STAT or cells incubated 8D-HYPOx. Latency and growth

rate (slope) were determined in different conditions. Experiments were always repeated

twice. Graph includes dot for the replicates, as well as median ± IQR (*p<0.01 compared to

control in the same condition). **A.** The dilution of minimal medium from 100% to 10%

(diluted MM) resulted in a progressive decrease of latency in 8D-HYPOx cells and to a lesser

extend in STAT cells. (*p <0.01 compared to 100%MM). Two independent experiments were

pooled. Each point represents the latency/well in two (STAT n=5, 8D-HYPOx n=15) **B.** 100-

fold dilution of pantothenic acid (starting at 125 µM) were added to the suspension prior to

growth curve determination. Pantothenic acid significantly decreased the latency at a

concentration of 1.25×10^{-2} µM or more in 8D-HYPOx and to a lesser extend in STAT cells (*

p<0.01). Two independent experiments were pooled. Each point represents the latency/well

in two (STAT n=9, 8D-HYPOx n=6). **C.** Growth rate was significantly decreased in 20%MM and

10%MM in 8D-HYPOx **D.** A significantly increased growth rate was observed after addition of

PA (1.25 µM or more) to 8D-HYPOx cells and at all concentrations for STAT. **E.** The addition

of pantothenic acid at 125µM but not the dilution of MM increased the culturability of 8D-

HYPOx based on the observation of the number of positive wells (100 cells per well) and the

calculation of the probability of one cell to grow. Each dot represents the calculated

culturability. Two independents experiments are pooled. **F.** Proposed model for the effect of

hypoxia and nutrient depletion. From now on, 8D-HYPOx are called VBNC (Viable but not

culturable cells). After 8 d of incubation in hypoxia and nutrient depletion, the decrease in

latency observed upon addition of pantothenic acid on VBNC is explained by the reactivation

of VBNC to culturable cells (increased culturability) and by the increased growth rate of

culturable cells (increased slope). The decrease in latency observed upon dilutions of MM

(decreased nutrient availability) is explained by a faster rewire of the culturable cells

(decreased latency corresponding to decreased time to resume growth) without increased

culturability or growth rate.

1045

Figure 4: Phenotypic array analysis of VBNC showed a lower metabolic activity. We used

the Biolog® technology to explore the metabolic requirements of VBNC compared to STAT

and 8D-NORMOx cells. We selected 8 plates and 761 different metabolites to scan the ability

of substrates to be used as an energetic source. The respiratory activity (measurement of

tetrazolium reduction in arbitrary units) was compared at the end of the incubation (6 days).

Two biological replicates (1 and 2) were done for the VBNC/8D-NORMOx conditions. **A.** The

heatmap of 521 selected compounds showed that VBNC exhibited a different metabolic

profile than STAT and 8D-NORMOx. **B.** The final endpoint analysis of the level of tetrazolium

reduction revealed a lower activity for VBNC cells compared to cells in other conditions.

VBNC and 8D-NORMOx conditions were done in duplicate (* p<0.01).

1056

Figure 5: Mitochondria of VBNC were depolarized with an increased mass and gene

expression. A. Mitochondria mass, and polarization were assessed in VBNC, 8D-NORMOx,

STAT and heat killed yeasts using Mitotracker green and TMRE and JC-1 stainings,

respectively and the ImageStream technology. In 8D-HYPOX (red histogram), the Mitotracker

staining is high, TMRE low and JC-1 high compared to STAT (blue histogram). **B.** Microscopic

visualization with Imagestream of the Mitotracker and TMRE stainings in VBNC and STAT

yeasts, showing the contrast in the intensity of the stainings between conditions. **C.**

Expression of five mitochondrial genes, cytochrome B (CYTb), ATP synthase 6 (ATP6), NADH

deshydrogenase (NADH), mitochondrial large sub unit (mtLSU), and cytochrome oxidase

(COX1) as expressed as the fold change to the geometric mean of the expression of GAPDH

and ACT1 housekeeping genes. The expression of CYTb, NADH, mtLSU and COX1 is

significantly increased in VBNC compared to STAT (* p=0.029).

1069

Figure 6: Secretome and proteome qualitative analysis showed specific features in VBNC.

A. Venn diagram of cellular and secreted proteins in stationary phase (STAT), VBNC and

control. Six secreted (clear dashed blue) and 103 cellular (deep blue) proteins were detected

specifically in VBNC. Six were secreted only in STAT (grey dashed line) and 150 were

detected in cellular protein only (dark line), 16 were secreted (pink dashed line) and 75

cellular (red line) only in 8D-NORMOx. Only proteins present in the 3 replicates for each

condition were selected and compared. **B.** GO enrichment analysis of secreted proteins

showed specific profile in VBNC. The major biological processes enriched in VBNC were

translation, metabolic process of carbohydrates and glycolytic biological processes whereas

oxidation-reduction was mainly enriched in 8D-NORMOx. **C.** GO enrichment analysis of

cellular proteins showed similar profiles in VBNC and 8D-NORMOx with an enrichment for

translation and oxidation-reduction.

1082

Figure 7: Secretome (A,B) and proteome (C,D) quantitative analysis showed an

independent and specific kinetic evolution associated with VBNC generation. Principal

coordinate analysis (PCoA) of secreted **A.** or cellular **C.** proteins and heatmap obtained by

Limma analysis of secretome **B.** and proteome **D.** data. For the secretome, 85 proteins were

differentially expressed overtime, including 49 significantly overexpressed in control and 36

in hypoxia. For the proteome, 63 cellular proteins were differentially expressed during

kinetics including 47 significantly overexpressed in normoxia and 16 in hypoxia. Inversely of

B, Data sets **D** showed that the STAT and hypoxia are closer than to normoxia.

1091

Figure 8: The fatty acid degradation pathway is required for VBNC viability. A. The Search

Tool for the Retrieval of Interacting Genes/Proteins (STRING) allowed the visualization of

predicted protein-protein interaction for the 63 proteins hypoxia-regulated identified. It

revealed an impact of hypoxia on diverse proteins as well as a prominent influence on fatty

acids metabolism. Different clusters were identified from the network mapping: pathways

linked to fatty acid metabolism (valine leucine and isoleucine degradation, propanoate

metabolism, fatty acid degradation and metabolism) and to carbon metabolism. **B.** Lipid

droplets stained with Bodipy® 505/515 showed the presence of lipid droplets in all

conditions. **C.** Eight entries among 63 proteins regulated in hypoxia mapped the fatty acid

degradation by using KEGG pathway mapping ("map" pathways are not colored,

Cryptococcus-specific pathways are colored green, the 8 entries are colored red) **D.** The fatty

acid degradation pathway reconstructed from *Saccharomyces cerevisiae* pathway. FA (Fatty

acid), FAA2 (Acyl Coa synthetase), POX1 (Fatty-acyl coenzyme A oxidase), FOX2 (3-

hydroxyacyl-CoA dehydrogenase, 3-hydroxyacyl-CoA dehydrogenase), POT1 (3-oxoacyl CoA

thiolase). **E.** Nine deletion mutants from fatty acid degradation pathway selected from KEGG

database and the orthology with *S. cerevisae* were assessed for their ability to survive

hypoxia in the model. The ratio of living cells for the mutants to that obtained with KN99α

was determined based on LIVE/DEAD staining after VBNC generation. A significant decreased

percentage of viable cells was measured for 04688Δ, 00490Δ, 07747Δ and 00524Δ (*p<0.01)

1111

Figure 9: The global transcription is decreased in VBNC but remains specific. A.

Transcriptional level of all conditions relative to that of *S. cerevisiae* spiked in each sample

showed a significant decrease in VBNC compared to other conditions **B.** PCA analysis of the

global transcriptomic data (wo_spikes) suggested that logarithmic cells (LOG) evolve to STAT

which evolve to VBNC or 8D-NORMOx **C.** GO enrichment analysis of upregulated transcript

(wo_spike) compared to STAT. The major enriched biological processes were signal

transduction, ATP synthesis coupled proton transport in VBNC and protein folding, oxidation

reduction process and response to stress in 8D-NORMOx. **D.** Heat map of the gene

expression revealed the presence of 7 clusters displaying different expression patterns for all

conditions. The color key represents normalized transformed counts**.**

Supplementary figures legends

1123

Supplementary figure 1: *In vitro* protocol for dormant cell generation. The protocol

followed several steps: (1) *Cryptocococcus neoformans* cells of a stock culture frozen at -80°C

were grown on Sabouraud agar plate for 2 to 5 days at room temperature; (2) 10^7 cells were

then suspended in 10 mL yeast extract peptone dextrose 2% (YPD) and incubated with

lateral shaking (150 rpm) at 30°C for 22 hours (stationary phase-STAT); (3) 100 µl, $2x10^7$ cells

were incubated again until the STAT and then placed in hypoxia or normoxia in the dark at

30°C (4) for up to 8 days when the resultant cells in hypoxia and normoxia were analyzed.

1131

1132   **Supplementary figure 2: Cell body size, capsule and cell wall thickness. A.** Median cell sizes

1133   were similar in 8D-HYPOx, 8D-NORMOx and STAT. **B.** Median capsule size was lower in 8D-

1134   HYPOx, 8D-NORMOx compared to STAT (*p<0.01, 100 cells measured). **C.** Cell wall was

1135   thicker in 8D-HYPOx compared to STAT (*p=0.0152, 50 cells measured).

1136

Supplementary figure 3: Flow cytometry diagrams assessing capsular, vacuolar structure

and viability in STAT, 8D-NORMOx and 8D-HYPOx. A. Histograms of fluorescence intensity

showing no difference in binding pattern after staining with anti-capsular polysaccharide E1

antibody **B.** More intense vacuolar staining with MDY-64 in 8D-HYPOx, 8D-NORMOx

conditions (one representative experiment out of the 3 independent experiments performed

is shown) **C.** Viability was assessed by membrane permeability staining (LIVE/DEAD, LVD)

showing almost 99% of live cells in STAT cells and 100% of dead cells in heat-killed cells (Left

panel). Plasma membrane was intact for more (87.2% [83.2-88.6]) cells in 8D-HYPOx, than in

8D-NORMOx (53.8% [50.9-60.5]). Experiments were done in triplicate and a representative

diagram is shown.

1147

Supplementary figure 4: Latency was influenced by medium and cell concentration

Growth of STAT and 8D-HYPOx cells was assessed using the BioScreen® apparatus. Serial

dilutions of 8D-HYPOx and STAT cells increased the latency in YPD (**A**) and in MM (**B**). Latency

curves extrapolations showed for both STAT and 8D-HYPOx cells, global latency was

decreased in YPD compared to MM. Each point represents the median ± IQR of the latency

of 3 independent experiments. **C.** Experimental set up used for the determination of the

probability of growth per cell (culturability). Hundred yeasts cells per well were plated in 96-

well plates in each condition (MM at 10 or 100% ± pantothenic acid (PA) at 125μM). The

number of positive wells per plate were determined and the probability of growth per plate

was calculated (see M&M section) **D.** 8D-HYPOx were exposed to macrophages during two

hours in the presence of opsonin. Culturability was similar in phagocytosed 8D-HYPOx cells

compared to controls. Each dot represents the calculated culturability. Two independents

experiments are pooled.

1161

1162　**Supplementary figure 5: ROS and RNS production were decreased in VBNC**

1163　Testing the ROS and RNS in VBNC vs STAT cells found a slight decreased in VBNC. ROS (left

1164　panel) and RNS (right panel) productions were measured using fluorescence probes and

1165　were significantly lower in VBNC compared to STAT cells respectively (*p<0.01). As expected,

1166　the addition of H_2O_2 as a positive control increased ROS and RNS levels.

1167

Supplementary figure 6: The number of identified proteins evolved differently between

secreted and cellular proteins in hypoxia and normoxia. The number of secreted protein

number significantly decreased over time in hypoxia (* p<0.01) (**A**) while the protein

concentration tended to increase in normoxia (**B**). The number of cellular proteins remained

stable in both conditions (**C**) while the protein concentration tended to increase (**D**). The

experiments were done in triplicates (median±IQR].

1174

1175　**Supplementary figure 7: Venn diagrams of STAT, VBNC and 8D-NORMOx for secreted and**

1176　**cellular proteins. A.** A hundred and seven cellular proteins were only present in hypoxia and

1177 2408 were common to the 3 experimental conditions. **B.** Seventeen secreted proteins were

1178 only present in hypoxia and 252 were common to the 3 experimental conditions.

1179

Supplementary figure 8: GO enrichment analysis for molecular function (A,B) and cellular

component (C,D) showed a particular pattern in hypoxia for secreted proteins (A,C) but

not for cellular (B,D) proteins. A. The major enriched molecular function process for

secreted proteins in hypoxia was structural constituents of ribosome and in normoxia

catalytic and oxidoreductase activity. **B.** For cellular component, the major enriched cellular

components were the ribosome, ribonucleoprotein complex and intracellular in hypoxia and

mainly ribosome in normoxia. **C.** The major enriched molecular function process for cellular

proteins in hypoxia were the same as normoxia: nucleotide binding, transferase activity and

catalytic activity. **D.** For cellular component, the major enriched cellular component for both

hypoxia and normoxia were cytoplasm, ribosome, ribonucleoprotein complex and

intracellular.

1191

Acknowledgements

We would like to acknowledge Christina A. Cuomo, Laurent Châtre, Guilhem Janbon, Jean-

Yves Coppée, Pierre Rocheteau, Pierre Henri Commere, Christine Schmitt, Olivier Gorgette,

Jacomine Krijnse-Locker, Caroline Proux for their comments on this work and help at

different steps of the study.

References

1.   Postgate JR. 1969. Chapter XVIII Viable counts and Viability, p. 611–628. *In* Methods in
Microbiology. Academic Press.
2.   Kaprelyants AS, Mukamolova GV, Kell DB. 1994. Estimation of dormant *Micrococcus luteus*
cells by penicillin lysis and by resuscitation in cell-free spent culture medium at high dilution.
FEMS Microbiol Lett 115:347.
3.   Emerson JB, Adams RI, Román CMB, Brooks B, Coil DA, Dahlhausen K, Ganz HH, Hartmann EM,
Hsu T, Justice NB, Paulino-Lima IG, Luongo JC, Lymperopoulou DS, Gomez-Silvan C, Rothschild-
Mancinelli B, Balk M, Huttenhower C, Nocker A, Vaishampayan P, Rothschild LJ. 2017.
Schrödinger's microbes: Tools for distinguishing the living from the dead in microbial
ecosystems. Microbiome 5:86.
4.   Roszak DB, Colwell RR. 1987. Survival strategies of bacteria in the natural environment.
Microbiol Rev 51:365–379.
5.   Nyström T. 2003. Nonculturable bacteria: programmed survival forms or cells at death's door?
BioEssays News Rev Mol Cell Dev Biol 25:204–211.
6.   Oliver JD. 2010. Recent findings on the viable but nonculturable state in pathogenic bacteria.
FEMS Microbiol Rev 34:415–425.
7.   Ayrapetyan M, Williams TC, Baxter R, Oliver JD. 2015. Viable but Nonculturable and Persister
Cells Coexist Stochastically and Are Induced by Human Serum. Infect Immun 83:4194–4203.
8.   Li L, Mendis N, Trigui H, Oliver JD, Faucher SP. 2014. The importance of the viable but non-
culturable state in human bacterial pathogens. Front Microbiol 5:258.
9.   Divol B, Lonvaud-Funel A. 2005. Evidence for viable but nonculturable yeasts in botrytis-
affected wine. J Appl Microbiol 99:85–93.
10.   Salma M, Rousseaux S, Grand AS-L, Divol B, Alexandre H. 2013. Characterization of the Viable
but Nonculturable (VBNC) State in *Saccharomyces cerevisiae*. PloS One 8:e77600.
11.   Serpaggi V, Remize F, Recorbet G, Gaudot-Dumas E, Sequeira-Le Grand A, Alexandre H. 2012.
Characterization of the "viable but nonculturable" (VBNC) state in the wine spoilage yeast
Brettanomyces. Food Microbiol 30:438–447.
12.   Brunet K, Alanio A, Lortholary O, Rammaert B. 2018. Reactivation of dormant/latent fungal
infection. J Infect 77:463–468.
13.   Lewenza S, Abboud J, Poon K, Kobryn M, Humplik I, Bell JR, Mardan L, Reckseidler-Zenteno S.
2018. Pseudomonas aeruginosa displays a dormancy phenotype during long-term survival in
water. PloS One 13:e0198384.
14.   Helaine S, Cheverton AM, Watson KG, Faure LM, Matthews SA, Holden DW. 2014.
Internalization of *Salmonella* by macrophages induces formation of nonreplicating persisters.
Science 343:204–208.
15.   Markus MB. 2018. Biological concepts in recurrent *Plasmodium vivax* malaria. Parasitology
145:1765–1771.
16.   Rittershaus ESC, Baek S-H, Sassetti CM. 2013. The normalcy of dormancy: common themes in
microbial quiescence. Cell Host Microbe 13:643–651.
17.   Garcia-Hermoso D, Janbon G, Dromer F. 1999. Epidemiological evidence for dormant
Cryptococcus neoformans infection. Cryptococcus Neoformans 37:3204–3209.
18.   Dromer F, Ronin O, Dupont B. 1992. Isolation of *Cryptococcus neoformans* var. *gattii* from an
Asian patient in France: evidence for dormant infection in healthy subjects. J Med Vet Mycol
Bi-Mon Publ Int Soc Hum Anim Mycol 30:395–397.
19.   Kwon-Chung KJ, Bennett JE. 1992. Medical mycology. Philadelphia : Lea & Febiger.
20.   Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning DW, Loyse A,
Boulware DR. 2017. Global burden of disease of HIV-associated cryptococcal meningitis: an
updated analysis. Lancet Infect Dis 17:873.

21.   Baker RD. 1976. The primary pulmonary lymph node complex of cryptococcosis. Am J Clin
Pathol 65:83–92.
22.   Goldman D, Lee SC, Casadevall A. 1994. Pathogenesis of pulmonary Cryptococcus neoformans
infection in the rat. Infect Immun 62:4755–4761.
23.   Alanio A, Vernel-Pauillac F, Sturny-Leclère A, Dromer F. 2015. Cryptococcus neoformans host
adaptation: toward biological evidence of dormancy. mBio 6:e02580-14.
24.   Rocheteau P, Vinet M, Chretien F. 2015. Dormancy and quiescence of skeletal muscle stem
cells. Results Probl Cell Differ 56:215–235.
25.   Albuquerque P, Nicola AM, Nieves E, Paes HC, Williamson PR, Silva-Pereira I, Casadevall A.
2013. Quorum sensing-mediated, cell density-dependent regulation of growth and virulence in
Cryptococcus neoformans. mBio 5:e00986-13.
26.   Baker RD, Haugen RK. 1955. Tissue Changes and Tissue Diagnosis in Cryptococcosis: A Study of
26 Cases. Am J Clin Pathol 25:14.
27.   Haugen RK, Baker RD. 1954. The Pulmonary Lesions in Cryptococcosis with Special References
to Subpleural Nodules. Am J Clin Pathol 24:1381.
28.   Idnurm A, Bahn Y-S, Nielsen K, Lin X, Fraser JA, Heitman J. 2005. Deciphering the model
pathogenic fungus *Cryptococcus neoformans*. Nat Rev Microbiol 3:753–764.
29.   De Virgilio C. 2012. The essence of yeast quiescence. FEMS Microbiol Rev 36:306–339.
30.   Grahl N, Shepardson KM, Chung D, Cramer RA. 2012. Hypoxia and fungal pathogenesis: to air
or not to air? Eukaryot Cell 11:560–570.
31.   Wayne LG, Hayes LG. 1996. An in vitro model for sequential study of shiftdown of
Mycobacterium tuberculosis through two stages of nonreplicating persistence. Infect Immun
64:2062–2069.
32.   Semighini CP, Averette AF, Perfect JR, Heitman J. 2011. Deletion of *Cryptococcus neoformans*
AIF ortholog promotes chromosome aneuploidy and fluconazole-resistance in a metacaspase-
independent manner. PLoS Pathog 7:e1002364.
33.   Day AP, Oliver JD. 2004. Changes in membrane fatty acid composition during entry of *Vibrio
vulnificus* into the viable but nonculturable state. J Microbiol Seoul Korea 42:69–73.
34.   Latil M, Rocheteau P, Châtre L, Sanulli S, Mémet S, Ricchetti M, Tajbakhsh S, Chrétien F. 2012.
Skeletal muscle stem cells adopt a dormant cell state post mortem and retain regenerative
capacity. Nat Commun 3:903.
35.   DuBois JC, Pasula R, Dade JE, Smulian AG. 2016. Yeast Transcriptome and In Vivo Hypoxia
Detection Reveals *Histoplasma capsulatum* Response to Low Oxygen Tension. Med Mycol
54:40–58.
36.   Qualls JE, Murray PJ. 2016. Immunometabolism within the tuberculosis granuloma: amino
acids, hypoxia, and cellular respiration. Semin Immunopathol 38:139–152.
37.   Sershen CL, Plimpton SJ, May EE. 2016. Oxygen Modulates the Effectiveness of Granuloma
Mediated Host Response to *Mycobacterium tuberculosis*: A Multiscale Computational Biology
Approach. Front Cell Infect Microbiol 6:6.
38.   Lane SW, Williams DA, Watt FM. 2014. Modulating the stem cell niche for tissue regeneration.
Nat Biotechnol 32:795–803.
39.   Mathieu J, Zhou W, Xing Y, Sperber H, Ferreccio A, Agoston Z, Kuppusamy KT, Moon RT,
Ruohola-Baker H. 2014. Hypoxia-inducible factors have distinct and stage-specific roles during
reprogramming of human cells to pluripotency. Cell Stem Cell 14:592–605.
40.   Dong Z, Wang JZ, Yu F, Venkatachalam MA. 2003. Apoptosis-Resistance of Hypoxic Cells:
Multiple Factors Involved and a Role for IAP-2. Am J Pathol 163:663–671.
41.   Wu J, Parungo C, Wu G, Kang PM, Laham RJ, Sellke FW, Simons M, Li J. 2004. PR39 inhibits
apoptosis in hypoxic endothelial cells: role of inhibitor apoptosis protein-2. Circulation
109:1660–1667.
42.   Graeber TG, Osmanian C, Jacks T, Housman DE, Koch CJ, Lowe SW, Giaccia AJ. 1996. Hypoxia-
mediated selection of cells with diminished apoptotic potential in solid tumours. Nature
379:88–91.

43.   Arana I, Orruño M, Pérez-Pascual D, Seco C, Muela A, Barcina I. 2007. Inability of *Escherichia*
coli to resuscitate from the viable but nonculturable state. FEMS Microbiol Ecol 62:1–11.

44.   Li L, Mendis N, Trigui H, Oliver JD, Faucher SP. 2014. The importance of the viable but non-
culturable state in human bacterial pathogens. Front Microbiol 5:258.

45.   Spry C, Kirk K, Saliba KJ. 2008. Coenzyme A biosynthesis: an antimicrobial drug target. FEMS
Microbiol Rev 32:56–106.

46.   Liew KL, Jee JM, Yap I, Yong PVC. 2016. In Vitro Analysis of Metabolites Secreted during
Infection of Lung Epithelial Cells by *Cryptococcus neoformans*. PloS One 11:e0153356.

47.   Calcott PH, Postgate JR. 1972. On substrate-accelerated death in *Klebsiella aerogenes*. J Gen
Microbiol 70:115–122.

48.   Mukamolova GV, Kaprelyants AS, Kell DB, Young M. 2003. Adoption of the transiently non-
culturable state — a bacterial survival strategy?

49.   Nofal M, Zhang K, Han S, Rabinowitz JD. 2017. mTOR Inhibition Restores Amino Acid Balance in
Cells Dependent on Catabolism of Extracellular Protein. Mol Cell 67:936-946.e5.

50.   Oblinger JL, Koburger JA. 1975. Understanding and Teaching the Most Probable Number
Technique. J Milk Food Technol 38:540–545.

51.   Pinto D, Santos MA, Chambel L. 2013. Thirty years of viable but nonculturable state research:
unsolved molecular mechanisms. Crit Rev Microbiol 41:61–76.

52.   Blackburn CW, McCarthy JD. 2000. Modifications to methods for the enumeration and
detection of injured Escherichia coli O157:H7 in foods. Int J Food Microbiol 55:285–290.

53.   Steenbergen JN, Shuman HA, Casadevall A. 2001. *Cryptococcus neoformans* interactions with
amoebae suggest an explanation for its virulence and intracellular pathogenic strategy in
macrophages. Proc Natl Acad Sci U S A 98:15245–15250.

54.   García MT, Jones S, Pelaz C, Millar RD, Abu Kwaik Y. 2007. *Acanthamoeba polyphaga*
resuscitates viable non-culturable *Legionella pneumophila* after disinfection. Environ Microbiol
9:1267–1277.

55.   Steinert M, Emödy L, Amann R, Hacker J. 1997. Resuscitation of viable but nonculturable
Legionella pneumophila Philadelphia JR32 by *Acanthamoeba castellanii*. Appl Environ
Microbiol 63:2047–2053.

56.   Schumann R, Rieling T, Görs S, Hammer A, Selig U, Schiewer U. 2003. Viability of bacteria from
different aquatic habitats. I. Environmental conditions and productivity. Aquat Microb Ecol
32:121–135.

57.   Sebastià J, Cristòfol R, Martín M, Rodríguez-Farré E, Sanfeliu C. 2003. Evaluation of fluorescent
dyes for measuring intracellular glutathione content in primary cultures of human neurons and
neuroblastoma SH-SY5Y. Cytometry A 51A:16–25.

58.   DeRisi JL, Iyer VR, Brown PO. 1997. Exploring the metabolic and genetic control of gene
expression on a genomic scale. Science 278:680–686.

59.   Boucherie H. 1985. Protein synthesis during transition and stationary phases under glucose
limitation in *Saccharomyces cerevisiae*. J Bacteriol 161:385–392.

60.   Fuge EK, Braun EL, Werner-Washburne M. 1994. Protein synthesis in long-term stationary-
phase cultures of *Saccharomyces cerevisiae*. J Bacteriol 176:5802–5813.

61.   Ju Q, Warner JR. 1994. Ribosome synthesis during the growth cycle of *Saccharomyces*
cerevisiae. Yeast Chichester Engl 10:151–157.

62.   Lee J, Giordano S, Zhang J. 2012. Autophagy, mitochondria and oxidative stress: cross-talk and
redox signalling. Biochem J 441:523–540.

63.   Rodrigues ML, Nakayasu ES, Oliveira DL, Nimrichter L, Nosanchuk JD, Almeida IC, Casadevall A.
2008. Extracellular vesicles produced by *Cryptococcus neoformans* contain protein
components associated with virulence. Eukaryot Cell 7:58–67.

64.   Geddes JMH, Croll D, Caza M, Stoynov N, Foster LJ, Kronstad JW. 2015. Secretome profiling of
Cryptococcus neoformans reveals regulation of a subset of virulence-associated proteins and
potential biomarkers by protein kinase A. BMC Microbiol 15:206.

65. Lev S, Crossett B, Cha SY, Desmarini D, Li C, Chayakulkeeree M, Wilson CF, Williamson PR,
Sorrell TC, Djordjevic JT. 2014. Identification of Aph1, a phosphate-regulated, secreted, and
vacuolar acid phosphatase in *Cryptococcus neoformans*. mBio 5:e01649-14.
66. Eisenman HC, Frases S, Nicola AM, Rodrigues ML, Casadevall A. 2009. Vesicle-associated
melanization in *Cryptococcus neoformans*. Microbiol Read Engl 155:3860–3867.
67. Rodrigues ML, Nakayasu ES, Oliveira DL, Nimrichter L, Nosanchuk JD, Almeida IC, Casadevall A.
2007. Extracellular vesicles produced by *Cryptococcus neoformans* contain protein
components associated with virulence. Eukaryot Cell 7:58–67.
68. Rodrigues ML, Djordjevic JT. 2012. Unravelling secretion in *Cryptococcus neoformans*: more
than one way to skin a cat. Mycopathologia 173:407–418.
69. de Gassart A, Geminard C, Fevrier B, Raposo G, Vidal M. 2003. Lipid raft-associated protein
sorting in exosomes. Blood 102:4336–4344.
70. Homer CM, Summers DK, Goranov AI, Clarke SC, Wiesner DL, Diedrich JK, Moresco JJ, Toffaletti
D, Upadhya R, Caradonna I, Petnic S, Pessino V, Cuomo CA, Lodge JK, Perfect J, Yates JR,
Nielsen K, Craik CS, Madhani HD. 2016. Intracellular Action of a Secreted Peptide Required for
Fungal Virulence. Cell Host Microbe 19:849–864.
71. Lee H, Chang YC, Nardone G, Kwon-Chung KJ. 2007. TUP1 disruption in Cryptococcus
neoformans uncovers a peptide-mediated density-dependent growth phenomenon that
mimics quorum sensing. Mol Microbiol 64:591.
72. Lee H, Chang YC, Varma A, Kwon-Chung KJ. 2009. Regulatory diversity of TUP1 in *Cryptococcus*
neoformans. Eukaryot Cell 8:1901–1908.
73. Selvan LDN, Kaviyil JE, Nirujogi RS, Muthusamy B, Puttamallesh VN, Subbannayya T, Syed N,
Radhakrishnan A, Kelkar DS, Ahmad S, Pinto SM, Kumar P, Madugundu AK, Nair B, Chatterjee
A, Pandey A, Ravikumar R, Gowda H, Prasad TSK. 2014. Proteogenomic analysis of pathogenic
yeast *Cryptococcus neoformans* using high resolution mass spectrometry. Clin Proteomics 11:5.
74. Selvan LDN, Sreenivasamurthy SK, Kumar S, Yelamanchi SD, Madugundu AK, Anil AK, Renuse S,
Nair BG, Gowda H, Mathur PP, Satishchandra P, Shankar SK, Mahadevan A, Keshava Prasad TS.
2015. Characterization of host response to *Cryptococcus neoformans* through quantitative
proteomic analysis of cryptococcal meningitis co-infected with HIV. Mol Biosyst 11:2529–2540.
75. Huang M, Hebert AS, Coon JJ, Hull CM. 2015. Protein Composition of Infectious Spores Reveals
Novel Sexual Development and Germination Factors in *Cryptococcus*. PLoS Genet
11:e1005490.
76. Jamieson DJ. 1998. Oxidative stress responses of the yeast *Saccharomyces cerevisiae*. Yeast
Chichester Engl 14:1511–1527.
77. De Virgilio C, Loewith R. 2006. The TOR signalling network from yeast to man. Int J Biochem
Cell Biol 38:1476–1481.
78. Rohde J, Heitman J, Cardenas ME. 2001. The TOR kinases link nutrient sensing to cell growth. J
Biol Chem 276:9583–9586.
79. García-Rodas R, Cordero RJB, Trevijano-Contador N, Janbon G, Moyrand F, Casadevall A,
Zaragoza O. 2014. Capsule growth in *Cryptococcus neoformans* is coordinated with cell cycle
progression. mBio 5:e00945-14.
80. Martinez DL, Tsuchiya Y, Gout I. 2014. Coenzyme A biosynthetic machinery in mammalian cells.
Biochem Soc Trans 42:1112–1117.
81. Srinivasan B, Sibon OCM. 2014. Coenzyme A, more than "just" a metabolic cofactor. Biochem
Soc Trans 42:1075–1079.
82. Kretschmer M, Wang J, Kronstad JW. 2012. Peroxisomal and mitochondrial β-oxidation
pathways influence the virulence of the pathogenic fungus *Cryptococcus neoformans*. Eukaryot
Cell 11:1042–1054.
83. Itoh T, Waki H, Kaneko H. 1975. Changes of Lipid Composition with Growth Phase of
Cryptococcus neoformans. Agric Biol Chem 39:2365–2371.
84. Lorenz MC, Bender JA, Fink GR. 2004. Transcriptional response of *Candida albicans* upon
internalization by macrophages. Eukaryot Cell 3:1076–1087.

85. Hu G, Cheng P-Y, Sham A, Perfect JR, Kronstad JW. 2008. Metabolic adaptation in *Cryptococcus*
neoformans during early murine pulmonary infection. Mol Microbiol 69:1456–1475.
86. da S Derengowski L, Paes HC, Albuquerque P, Tavares AHFP, Fernandes L, Silva-Pereira I,
Casadevall A. 2013. The transcriptional response of *Cryptococcus neoformans* to ingestion by
Acanthamoeba castellanii and macrophages provides insights into the evolutionary adaptation
to the mammalian host. Eukaryot Cell 12:761–774.
87. Hommel B, Mukaremera L, Cordero RJB, Coelho C, Desjardins CA, Sturny-Leclère A, Janbon G,
Perfect JR, Fraser JA, Casadevall A, Cuomo CA, Dromer F, Nielsen K, Alanio A. 2018. Titan cells
formation in *Cryptococcus neoformans* is finely tuned by environmental conditions and
modulated by positive and negative genetic regulators. PLoS Pathog 14:e1006982.
88. Ferreira GF, de Matos Baltazar L, Santos JRA, Monteiro AS, de Oliveira Fraga LA, Resende-
Stoianoff MA, Santos DA. 2013. The role of oxidative and nitrosative bursts caused by azoles
and amphotericin B against the fungal pathogen *Cryptococcus gattii*. J Antimicrob Chemother
68:1801–1811.
89. Kahm M, Hasenbrink G, Lichtenberg-Fraté H, Ludwig J, Kschischo M. 2010. grofit: Fitting
Biological Growth Curves withR. J Stat Softw 33.
90. Alanio A, Desnos-Ollivier M, Dromer F. 2011. Dynamics of *Cryptococcus neoformans*-
macrophage interactions reveal that fungal background influences outcome during
cryptococcal meningoencephalitis in humans. mBio 2:e00158-11.
91. Casadevall A, DeShaw M, Fan M, Dromer F, Kozel TR, Pirofski L a. 1994. Molecular and idiotypic
analysis of antibodies to *Cryptococcus neoformans* glucuronoxylomannan. Infect Immun
62:3864–3872.
92. Nielsen K, Cox GM, Litvintseva AP, Mylonakis E, Malliaris SD, Benjamin DK, Giles SS, Mitchell
TG, Casadevall A, Perfect JR, Heitman J. 2005. *Cryptococcus neoformans* {alpha} strains
preferentially disseminate to the central nervous system during coinfection. Infect Immun
73:4922–4933.
93. Bochner BR, Gadzinski P, Panomitros E. 2001. Phenotype microarrays for high-throughput
phenotypic testing and assay of gene function. Genome Res 11:1246–1255.
94. Moyrand F, Lafontaine I, Fontaine T, Janbon G. 2008. UGE1 and UGE2 regulate the UDP-
glucose/UDP-galactose equilibrium in *Cryptococcus neoformans*. Eukaryot Cell 7:2069–2077.
95. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F. 2002.
Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of
multiple internal control genes. Genome Biol 3:RESEARCH0034.
96. Pfaffl MW. 2001. A new mathematical model for relative quantification in real-time RT–PCR.
Nucleic Acids Res 29:e45.
97. Wu X, Xiong E, Wang W, Scali M, Cresti M. 2014. Universal sample preparation method
integrating trichloroacetic acid/acetone precipitation with phenol extraction for crop
proteomic analysis. Nat Protoc 9:362–374.
98. Cox J, Neuhauser N, Michalski A, Scheltema RA, Olsen JV, Mann M. 2011. Andromeda: a
peptide search engine integrated into the MaxQuant environment. J Proteome Res 10:1794–
1805.
99. Cox J, Mann M. 2008. MaxQuant enables high peptide identification rates, individualized
p.p.b.-range mass accuracies and proteome-wide protein quantification. Nat Biotechnol
26:1367–1372.
100. Tyanova S, Temu T, Cox J. 2016. The MaxQuant computational platform for mass
spectrometry-based shotgun proteomics. Nat Protoc 11:2301–2319.
101. Cox J, Hein MY, Luber CA, Paron I, Nagaraj N, Mann M. 2014. Accurate proteome-wide label-
free quantification by delayed normalization and maximal peptide ratio extraction, termed
MaxLFQ. Mol Cell Proteomics MCP 13:2513–2526.
102. Bø TH, Dysvik B, Jonassen I. 2004. LSimpute: accurate estimation of missing values in
microarray data with least squares methods. Nucleic Acids Res 32:e34.

103. Gower JC. 1966. Some distance properties of latent root and vector methods used in
multivariate analysis. Biometrika 325–38.
104. Ritchie ME, Phipson B, Wu, Hu Y, Law CW, Shi W, Smyth GK. 2015. limma powers differential
expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Res 43:e47.
105. Benjamini Y, Hochberg Y. 1995. Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. J R Stat Soc Ser B Methodol 57:289–300.
106. Priebe S, Kreisel C, Horn F, Guthke R, Linde J. 2014. FungiFun2: a comprehensive online
resource for systematic analysis of gene lists from fungal species. Bioinforma Oxf Engl 31:445–
446.
107. Bardou P, Mariette J, Escudié F, Djemiel C, Klopp C. 2014. jvenn: an interactive Venn diagram
viewer. BMC Bioinformatics 15.
108. Szklarczyk D, Franceschini A, Wyder S, Forslund K, Heller D, Huerta-Cepas J, Simonovic M, Roth
A, Santos A, Tsafou KP, Kuhn M, Bork P, Jensen LJ, von Mering C. 2014. STRING v10: protein-
protein interaction networks, integrated over the tree of life. Nucleic Acids Res 43:D447-52.
109. Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet.journal 17:10–12.
110. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.
2013. STAR: ultrafast universal RNA-seq aligner. Bioinformatics 29:15–21.
111. Liao Y, Smyth GK, Shi W. 2014. featureCounts: an efficient general purpose program for
assigning sequence reads to genomic features. Bioinforma Oxf Engl 30:923–930.
112. Ewels P, Magnusson M, Lundin S, Käller M. 2016. MultiQC: summarize analysis results for
multiple tools and samples in a single report. Bioinformatics 32:3047–3048.
113. Love MI, Huber W, Anders S. 2014. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. Genome Biol 15:550.
114. Basenko EY, Pulman JA, Shanmugasundram A, Harb OS, Crouch K, Starns D, Warrenfeltz S,
Aurrecoechea C, Stoeckert CJ, Kissinger JC, Roos DS, Hertz-Fowler C. 2018. FungiDB: An
Integrated Bioinformatic Resource for Fungi and Oomycetes. J Fungi Basel Switz 4.
115. Marchler-Bauer A, Bo Y, Han L, He J, Lanczycki CJ, Lu S, Chitsaz F, Derbyshire MK, Geer RC,
Gonzales NR, Gwadz M, Hurwitz DI, Lu F, Marchler GH, Song JS, Thanki N, Wang Z, Yamashita
RA, Zhang D, Zheng C, Geer LY, Bryant SH. 2017. CDD/SPARCLE: functional classification of
proteins via subfamily domain architectures. Nucleic Acids Res 45:D200–D203.
116. Nielsen H. 2017. Predicting Secretory Proteins with SignalP. Methods Mol Biol Clifton NJ
1611:59–73.

1485

A

STAT cellular **150**

6 STAT secreted

VBNC secreted **6**

VBNC cellular **103**

8D-NORMOx cellular **75**

Common cellular **1654**

Common cellular + secreted **226**

16 8D-NORMOx secreted

17 Common secreted

B **Secreted proteins**

D4
D6
VBNC
HYPOXIA

STAT

D8
D6
D4
NORMOXIA

- translation
- carbohydrate metabolic process
- proteolysis involved in cellular protein catabolic process
- pentose-phosphate shunt
- glycolytic process
- oxidation-reduction process
- tricarboxylic acid cycle

C **Cellular proteins**

D4
D6
VBNC
HYPOXIA

STAT

D8
D6
D4
NORMOXIA

- translation
- tRNA aminoacylation for protein translation
- cellular amino acid biosynthetic process
- protein folding
- oxidation-reduction process
- obsolete GTP catabolic process
- proteolysis involved in cellular protein catabolic process
- intracellular protein transport
- formation of translation preinitiation complex
- translational elongation

A

Edge Confidence

low (0.150)   high (0.700)

medium (0.400)   highest (0.900)

Node Color

colored nodes:
query proteins and first shell of interactors

white nodes:
second shell of interactors

- Valine, leucine and isoleucine degradation
- Propanoate metabolism
- Fatty acid metabolism
- Fatty acid degradation
- Carbon metabolism

B

BF | Bodipy

VBNC

8D-NORMOx

STAT

C

FATTY ACID DEGRADATION

D

Fatty acid degradation

trans-2,3-dehydroacyl-CoA → 3-oxoacyl-CoA

FOX2 (CNAG_05721)

POX1 (CNAG_07747)

POT1 (CNAG_00490)

FA → FA CoA
FAA1
FAA2 (CNAG_3019)
FAA3
FAA4
FAT1

acyl$_{n-2}$-CoA → 2,3,4-saturated fatty acyl-CoA

Acetyl coenzyme (Co) A

E

% of living cells ratio / KN99

KN99α, 02489Δ, 04688Δ, 06628Δ, 03019Δ, 00490Δ, 07747Δ, 04531Δ, 00524Δ, 05721Δ

Table 1 : Cell dilutions and latency related to the culture medium and their extrapolated equation

Cell dilutions	Latency (min)			
	YPD		MM	
	STAT	8D-HYPOx	STAT	8D-HYPOx
$3x10^5$	544 [409-732]	1764 [1605-1903]		
$3x10^4$	789 [641-958]	2030 [1902-2228]	585 [515-602]	2732 [2548-2903]
$3x10^3$	1037 [879-1223]	2356 [2264-2838]	1120 [1101-1154]	4124 [3741-4360]
$3x10^2$	1330 [1205-1501]	2680 [2381-3197]	2408 [2058-2502]	4668 [4353-5140]
$3x10^1$	1657 [1485-1801]	3557 [2924-3810]	3399 [3003-3618]	4483 [4354-5277]
3	2044 [1919-2241]		4127 [4093-4336]	
Equation (R^2)	Y=2127-296logX (0.90)	Y=3947-408logX (0.59)	Y=4632-941logX (0.97)	Y=6603-810logX (0.74)

Table 2: Metabolic pathways regulatd in hypoxia by STRING analysis

Metabolic pathways	Genes	Corrected *p* value (FDR 1%)
Fatty acid degradation	CNAG_00490, CNAG_00524, CNAG_03019, CNAG_02489, CNAG_06628, CNAG_07747, CNAG_04531, CNAG_04688	1,20E-09
Valine, leucine and isoleucine degradation	CNAG_00484, CNAG_00490, CNAG_00524, CNAG_03067, CNAG_06628, CNAG_04531, CNAG_04351, CNAG_04688	2,40E-08
Fatty acid metabolism	CNAG_00490, CNAG_00524, CNAG_03019, CNAG_07747, CNAG_04531, CNAG_04688	9,50E-07
Propanoate metabolism	CNAG_00797, CNAG_06628, CNAG_04531, CNAG_04351, CNAG_04688	4,00E-05
Carbon metabolism	CNAG_00797, CNAG_00826, CNAG_05653, CNAG_05303, CNAG_04531, CNAG_04351, CNAG_04217, CNAG_04688	4,00E-05
Biosynthesis of secondary metabolites	CNAG_00484, CNAG_00490, CNAG_00524, CNAG_00797, CNAG_03067, CNAG_02489, CNAG_06628, CNAG_04531, CNAG_04217, CNAG_04688, CNAG_06431	2,60E-04
Peroxisome	CNAG_00537, CNAG_03067, CNAG_03019, CNAG_06551, CNAG_07747	2,90E-04
Metabolic pathways	CNAG_00484, CNAG_00490, CNAG_00524, CNAG_00797, CNAG_00826, CNAG_03067, CNAG_03019, CNAG_01540, CNAG_02489, CNAG_06628, CNAG_07747, CNAG_05653, CNAG_05303, CNAG_04531, CNAG_04351, CNAG_04217, CNAG_04688	3,60E-04
beta-Alanine metabolism	CNAG_06628, CNAG_04531, CNAG_04351, CNAG_04688	4,50E-04
Microbial metabolism in diverse environments	CNAG_00490, CNAG_00797, CNAG_02489, CNAG_06628, CNAG_05653, CNAG_05303, CNAG_04531, CNAG_04217	4,70E-04
alpha-Linolenic acid metabolism	CNAG_00490, CNAG_07747	8,50E-04
Pyruvate metabolism	CNAG_00797, CNAG_06628, CNAG_05653, CNAG_04217	2,40E-03
Glycolysis / Gluconeogenesis	CNAG_00797, CNAG_02489, CNAG_06628, CNAG_04217	2,80E-03
Butanoate metabolism	CNAG_03067, CNAG_02045, CNAG_04531	7,20E-03
Fatty acid elongation	CNAG_00524, CNAG_04531	2,20E-02

Table 3: Transcriptomic pathways involved in the VBNC cluster 3 and genes the corresponding upregulated genes.

Source	ID	Name	Result gene list	Fold enrichment	Odds ratio	P-value	Benjamini Hochberg correction
GO	GO:0006468	protein phosphorylation	CNAG_00769,CNAG_01905,CNAG_02389,CNAG_02820,CNAG_03216,CNAG_03670,CNAG_04040,CNAG_04755,CNAG_05439,CNAG_06193,CNAG_06310,CNAG_06490,CNAG_06845,	3.59	3.84	1.3E-04	1.8E-03
GO	GO:0016310	phosphorylation	CNAG_00769,CNAG_01905,CNAG_02389,CNAG_02820,CNAG_03216,CNAG_03670,CNAG_04040,CNAG_04755,CNAG_05439,CNAG_06193,CNAG_06310,CNAG_06490,CNAG_06845,	2.9	3.08	8.7E-04	6.1E-03
GO	GO:0050789	regulation of biological process	CNAG_00332,CNAG_00547,CNAG_00818,CNAG_01258,CNAG_01533,CNAG_01611,CNAG_01841,CNAG_03212,CNAG_03763,CNAG_03902,CNAG_04036,CNAG_04243,CNAG_04496,CNAG_05535,CNAG_05695,CNAG_05861,CNAG_06188,CNAG_06531,CNAG_06606,CNAG_06691,CNAG_06814,CNAG_07015,CNAG_07320,CNAG_07470,CNAG_07506,CNAG_07921,	1.79	1.95	3.2E-03	6.4E-03
GO	GO:0023051	regulation of signaling	CNAG_00547,CNAG_01258,CNAG_01611,CNAG_03763,CNAG_04243,CNAG_07015,	4.54	4.7	3.3E-03	6.4E-03
GO	GO:0048583	regulation of response to stimulus	CNAG_00547,CNAG_01258,CNAG_01611,CNAG_03763,CNAG_04243,CNAG_07015,	4.54	4.7	3.3E-03	6.4E-03
GO	GO:0010646	regulation of cell communication	CNAG_00547,CNAG_01258,CNAG_01611,CNAG_03763,CNAG_04243,CNAG_07015,	4.54	4.7	3.3E-03	6.4E-03
GO	GO:0009966	regulation of signal transduction	CNAG_00547,CNAG_01258,CNAG_01611,CNAG_03763,CNAG_04243,CNAG_07015,	4.54	4.7	3.3E-03	6.4E-03
GO	GO:0065007	biological regulation	CNAG_00332,CNAG_00547,CNAG_00818,CNAG_01258,CNAG_01533,CNAG_01611,CNAG_01841,CNAG_03212,CNAG_03763,CNAG_03902,CNAG_04036,CNAG_04243,CNAG_04496,CNAG_05535,CNAG_05695,CNAG_05861,CNAG_06188,CNAG_06531,CNAG_06606,CNAG_06691,CNAG_06814,CNAG_07015,CNAG_07320,CNAG_07470,CNAG_07506,CNAG_07921,	1.74	1.91	4.3E-03	6.4E-03

GO	GO:0036211	protein modification process	CNAG_00498,CNAG_00769,CNAG_01385,CNAG_01905,CNAG_02389,CNAG_02820,CNAG_03216,CNAG_03670,CNAG_04040,CNAG_04755,CNAG_05439,CNAG_06193,CNAG_06310,CNAG_06490,CNAG_06845,	2.2	2.33	4.6E-03	6.4E-03
GO	GO:0006464	cellular protein modification process	CNAG_00498,CNAG_00769,CNAG_01385,CNAG_01905,CNAG_02389,CNAG_02820,CNAG_03216,CNAG_03670,CNAG_04040,CNAG_04755,CNAG_05439,CNAG_06193,CNAG_06310,CNAG_06490,CNAG_06845,	2.2	2.33	4.6E-03	6.4E-03
GO	GO:0035023	regulation of Rho protein signal transduction	CNAG_00547,CNAG_01258,CNAG_04243,CNAG_07015,	6.63	6.78	5.2E-03	6.6E-03
GO	GO:0050794	regulation of cellular process	CNAG_00332,CNAG_00547,CNAG_01258,CNAG_01533,CNAG_01611,CNAG_01841,CNAG_03212,CNAG_03763,CNAG_03902,CNAG_04036,CNAG_04243,CNAG_05535,CNAG_05695,CNAG_05861,CNAG_06188,CNAG_06531,CNAG_06606,CNAG_06691,CNAG_06814,CNAG_07015,CNAG_07320,CNAG_07470,CNAG_07506,CNAG_07921,	1.75	1.89	6.0E-03	7.0E-03
GO	GO:0051056	regulation of small GTPase mediated signal transduction	CNAG_00547,CNAG_01258,CNAG_03763,CNAG_04243,CNAG_07015,	4.57	4.7	7.0E-03	7.0E-03
GO	GO:1902531	regulation of intracellular signal transduction	CNAG_00547,CNAG_01258,CNAG_03763,CNAG_04243,CNAG_07015,	4.57	4.7	7.0E-03	7.0E-03
KEGG	ec00900__PK__KEGG	Terpenoid backbone biosynthesis	CNAG_02662,CNAG_03216,CNAG_05601,CNAG_06534,CNAG_07377,	4.74	5.26	4.7E-03	