Transcriptomic changes resulting from *STK32B* overexpression identifies pathways potentially relevant to essential tremor

Calwing Liao[1,2], Faezeh Sarayloo[1,2], Veikko Vuokila[2], Daniel Rochefort[2], Fulya Akçimen[1,2], Simone Diamond[2], Alexandre D. Laporte[2], Dan Spiegelman[2], Qin He[3], Hélène Catoire[2], Patrick A. Dion[2,4], Guy A. Rouleau[1,2,4]

[1]Department of Human Genetics, McGill University, Montréal, Quebec, Canada

[2]Montreal Neurological Institute, McGill University, Montréal, Quebec, Canada.

[3]Department of Biomedical Sciences, Université de Montréal, Montréal, Quebec, Canada

[4]Department of Neurology and Neurosurgery, McGill University, Montréal, Quebec, Canada

Corresponding Author:

Dr. Guy A. Rouleau, MD, PhD, FRCPC, OQ

Department of Neurology and Neurosurgery

McGill University

Montréal, Québec, Canada

H3A 2B4

E-mail: guy.rouleau@mcgill.ca

Keywords: *STK32B*, essential tremor, transcriptome, *FUS*

Conflict of Interests: All authors report no conflict of interests.

Funding Sources: Canadian Institutes of Health Research

Abstract

Essential tremor (ET) is a common movement disorder that has a high heritability. A number of genetic studies have associated different genes and loci with ET, but few have investigated the biology of any of these genes. *STK32B* was significantly associated with ET in a large GWAS study and was found to be overexpressed in ET cerebellar tissue. Here, we overexpressed *STK32B* in human cerebellar DAOY cells and used an RNA-Seq approach to identify differentially expressed genes by comparing the transcriptome profile of these cells to the one of control DAOY cells. Pathway and gene ontology enrichment identified axon guidance, olfactory signalling and calcium-voltage channels as significant. Additionally, we show that overexpressing *STK32B* affects transcript levels of previously implicated ET genes such as *FUS*. Our results investigate the effects of overexpressed *STK32B* and suggest that it may be involved in relevant ET pathways and genes.

Introduction

Essential tremor (ET) is one of the most common movement disorders and is typically characterized by a kinetic tremor in the hand or arms[1]. The severity tends to increase with age and may involve different regions such as the head, voice or jaw[2]. Previous studies investigating the histology of post-mortem tissue have pinpointed the cerebellum as a region of interest for ET[3]. Specifically, abnormalities with Purkinje cell axons and dendrites were found in ET brains[4]. Furthermore, the olivocerebellar circuitry has been implicated in ET pathology and many calcium voltage channels are highly expressed in this circuitry[5].

The genetic etiology of ET has remained largely elusive and most studies have focused on common or rare variants and on looking for genetic overlap with other disorders[6,7]. Twin studies have shown that ET has a concordance of 69–93% in monozygotic twins and 27–29% in dizygotic twins, which suggests that both genetic and environmental factors drive the onset and development of this complex trait[8]. A recent genome-wide association study (GWAS) identified a significant locus in *STK32B* and found that ET patients overexpressed *STK32B* in cerebellar tissue by comparison to healthy controls[9]. *STK32B* is transcribed and translated into YANK2, a serine/threonine kinase, that has not been well characterized. There have been several exome-wide studies that implicated different genetic variants as causes of ET. The first ET-implicated gene found through exome

sequencing was the Fused in Sarcoma gene (*FUS*)[10]. However, it is unclear whether these "ET"

genes interact or have indirect effects on the expression of each other.

To understand the effects of overexpressed *STK32B* and identify pathways potentially relevant to

ET, we overexpressed the gene in human cerebellar DAOY cells and compared transcriptomic

changes in overexpressed cells and empty-vector controls using RNA sequencing. Several

interesting pathways such as axon guidance, calcium ion transmembrane transport, and olfactory

transduction were significantly enriched after overexpression of *STK32B*. We also identified

previously implicated ET genes whose expression is dysregulated through the overexpression of

STK32B, suggesting that overexpressed *STK32B* may have relevant downstream effects.

Results

Confirming overexpression of *STK32B*

The *STK32B* stable cell lines had higher RNA expression of *STK32B*, which was detected by

reverse-transcriptase qPCR (Supplementary Figure 1). RNA sequencing (RNA-Seq) also

confirmed that the stable cell lines have higher *STK32B* RNA levels compared to controls; it was

the most significantly differentially expressed gene (Figure 1–2).

Quality control of RNA-seq analyses

The QQ-plot for differentially expressed genes did not show stratification or potential biases

(Supplementary Figure 2). The M-A plot showed slight enrichment of downregulated genes

compared to upregulated genes (Supplementary Figure 3). A principal component (PC) plot

showed separation of controls and *STK32B* cell lines (Figure 3). Additionally, the dendrogram with

heatmap shows that controls and *STK32B* cell lines are distinct (Supplementary Figure 4). The

mean-variance plot shows the shrinkage under the sleuth model (Supplementary Figure 5).

Overexpression of *STK32B* affects expression of previously identified ET-associated genes

A total of 3,794 genes were found to be differentially expressed with a q-value <0.05. There were

425 genes with a $\beta > |0.5|$ and q-value <0.05. Amongst the 425 genes, three potentially relevant ET

genes were dysregulated: *FUS* (q-value = 0.007, β=0.812) and two calcium voltage channel genes

that are enriched in the olivocerebellar circuitry, *CACNA1C* (q-value = 0.011, β= 0.503) and

CACNA1A (q-value = 0.030, β= 0.702). Several pathways and gene ontologies were enriched for
differentially expressed genes (DEGs) due to the overexpression of *STK32B*, including olfactory
transduction (P=2.68E-39), axon guidance (P=9.50E-36), and calcium ion transmembrane
transport (P=3.02E-31) (Table 1).
89
Gene network analyses
To identify which cluster of DEGs drove the significant pathways, gene network analysis based
on co-expression was done and identified 14 different clusters within the differentially expressed
genes (Figure 4). Analysis of the top 3 largest clusters for pathway enrichments found several
similar pathways in different databases including axon guidance (P<1.09E-12) in cluster 1 and
calcium signalling pathway in cluster 3 (2.06E-05).

Discussion
The genetic etiology of ET is complex and likely explained by a combination of copy number
variants, rare variants, gene-gene interactions and common variant drivers. One of the most
significantly enriched pathways we identified was olfactory signalling and transduction. Previous
reports have shown conflicting evidence about olfactory loss in ET[11,12]. It may be that ET patients
with dysregulated olfactory signalling have overexpressed *STK32B*, which may contribute to the
subset of ET patients with olfactory loss.
103
Interestingly, *FUS* was amongst the genes dysregulated when *STK32B* was overexpressed. In the
exome familial ET study that identified *FUS*, we also found reduced mRNA levels for *FUS*[10].
Similarly, the expression of *FUS* is lower when *STK32B* is overexpressed, suggesting an indirect
relationship between the two genes. Additionally, two calcium voltage-gated channel genes,
CACNA1C AND *CACNA1A* were overexpressed. Enrichment analyses found calcium ion
transmembrane transport to be highly significant in GO Processes, suggesting that *STK32B* may
play a role upstream of these genes. In the olivocerebellar circuitry, a system implicated in ET, is
enriched for both of these genes[5,13]. *CACNA1A* has been shown to be predominantly expressed in
Purkinje cells, a cell type relevant to ET[13]. Another enriched pathway was axon guidance. A
previous study identified the *TENM4* missense variant segregating within ET families[14]. *TENM4*
is a regulator of axon guidance and myelination and Tenm4 knockout mice showed an ET

phenotype, suggesting that axon guidance is important in ET[14]. Similarly, overexpressed *STK32B*

could be dysregulating important pathways relevant to axon guidance.

By sub-stratifying the overexpressed genes into different clusters that are co-expressed, several

interesting pathways involving the cardiovascular system were identified. Pathways such as right

ventricular cardiomyopathy in KEGG and angiogenesis in GO Processes were found among the

top five significant pathways. In certain ET patients, beta-blockers can reduce tremor magnitude

and frequency. Beta-blockers lower blood pressure and are used to treat irregular heart rhythm and

other cardiovascular phenotypes. A common treatment for ET is propranolol, a beta-blocker. This

was a drug developed for cardiovascular health that affects adrenergic activity, but still reduces

tremor in ET individuals. This could suggest that overexpressed *STK32B* affects cardiovascular

health, which may in turn affect or lead to the ET phenotype or that *STK32B* may be pleiotropic

and affect the nervous and cardiovascular system differently.

Due to the heterogeneity of ET, it is likely that not all ET-affected individuals have overexpression

of *STK32B*. One limitation of this study is that DAOY cells are a cancerous cell line and may not

be the ideal model for understanding the transcriptome of Purkinje cells. Future studies on the

effects of *STK32B* overexpression could use induced pluripotent stem cells differentiated into cells

with Purkinje cells properties; such cells would likely have less biological noise. However, our

findings support the idea that *STK32B* is a gene of interest for a subset of ET cases, and further

investigations into how *STK32B* may interact with other genes are warranted.

Methods and Materials

Cell Culture and plasmid construction

The DAOY cell line was cultured in Eagle's Minimum Essential Medium (EMEM) with 10% fetal

bovine serum (FBS) at 37 degrees Celsius and 5% CO_2 with the Glico Pen-Strep-Glutamine

cocktail at 1X. Cells were passaged every 2 days at 80-90% confluence and at the same time. The

cDNA of *STK32B* (NM_001306082.1) was inserted into a pcDNA3.1(+) vector containing a

neomycin resistance gene, used as a selectable marker. This transcript was picked because it had

the highest expression in the cerebellum based on GTEx 53 v7. The plasmid was transformed and

expanded in XL10-Gold *Escherichia coli* strain (Stratagene). The cDNA of the plasmid was sent

for Sanger sequencing to validate the gene sequence.

Transfection and deriving stable cell lines

The cells were transfected for 48 hours with 8 ug of DNA of using the jetPRIME transfection

reagent (Polyplus). A green fluorescent protein (GFP) control vector was used to assess and

estimate transfection efficiency and optimize transfection parameters. Stable cell lines were

established by subjecting transfected cells to G418 antibiotic at 1 μg/mL for 10 days and maintained

at a concentration of 400 μg/mL for an additional 2 weeks. A kill curve was done to determine the

ideal antibiotic selection. In parallel, empty pcDNA3.1(+) vector control lines were also grown and

underwent antibiotic selection.

RNA extraction and sequencing

RNA was extracted using the RNeasy Mini Kit (Qiagen). The RNA concentration was measured

using the Synergy H4 microplate reader. RNA for the top three overexpressed *STK32B* cell lines

and three empty-vector controls was sent to Macrogen Inc. for sequencing. RT-qPCR was used to

confirm overexpression in the cell liens compared to controls. Library preparation was done with

the TruSeq Stranded Total RNA Kit (Ilumina) with Ribo-Zero depletion. Sequencing was done on

the NovaSeq 6000 at 150bp paired-end reads with a total of 200M reads. Samples were randomized

for cell lysis, RNA extraction, Ribo-Zero depletion, library preparation and sequencing to account

for potential batch effects.

Data processing and quality control

The FASTQ files were pseudo-aligned with Salmon using the Ensembl v94 annotation of the

human genome[16]. The parameters of Salmon included the following: 200 bootstraps with mapping

validation, GC bias correction, sequencing bias correction, 4 range factorization bins, a minimum

score fraction of 0.95 and VB optimization with a VB prior of 1e-5. The estimated counts were

analyzed using sleuth to identify DEGs[17]. A likelihood ratio test was used to identify differentially

expressed transcripts and a Wald test was used to get β values ($\log_2(x + 0.5)$). Additionally,

transcript-level aggregation was done against the Ensembl v94 annotation to determine gene-level

differential expression. P-values were corrected using the Benjamini-Hochberg procedure to

account for false discovery rate (FDR). Q-value (p-values corrected for FDR) significance was set

for an FDR of <0.05. A principal component (PC) plot and heatmap was done to identify clusters.

Pathway and gene set enrichment analysis

Pathway and gene set enrichment for relevant databases was done using Gene Network 2.0. To

investigate genes with larger fold changes, only differentially expressed genes with a $\beta > |0.5|$ were

analyzed for pathway enrichments. Network clustering was done by grouping genes based on co-

expression. Pathway analyses were also conducted with Gene Network 2.0 for the clusters using a

Wilcoxin test.

Acknowledgments
187 This work was supported by a Canadian Institutes of Health Research Foundation Scheme grant

188 (#332971). G.A.R. holds a Canada Research Chair in Genetics of the Nervous System and the

189 Wilder Penfield Chair in Neurosciences. C.L. is a recipient of the Frederick Banting and Charles

190 Best Canada Graduate Scholarship from the Canadian Institutes of Health Research (CIHR).

Data availability statement
Any data is available upon request of the corresponding author.

References

1.　Elble, R. J. What is Essential Tremor? *Curr. Neurol. Neurosci. Rep.* **13,** 353 (2013).
2.　Haubenberger, D. & Hallett, M. Essential Tremor. *N. Engl. J. Med.* **378,** 1802–1810 (2018).
3.　Gironell, A. The GABA Hypothesis in Essential Tremor: Lights and Shadows. *Tremor*
Other Hyperkinet. Mov. (N. Y). **4,** 254 (2014).
4.　Axelrad, J. E. *et al.* Reduced Purkinje cell number in essential tremor: a postmortem
study. *Arch. Neurol.* **65,** 101–7 (2008).
5.　Schmouth, J.-F., Dion, P. A. & Rouleau, G. A. Genetics of essential tremor: From
phenotype to genes, insights from both human and mouse studies. *Prog. Neurobiol.* **119–**
120, 1–19 (2014).
6.　Liao, C. *et al.* Investigating the association and causal relationship between restless legs
syndrome and essential tremor. *Parkinsonism Relat. Disord.* (2018).
doi:10.1016/j.parkreldis.2018.10.022
7.　Houle, G. *et al.* Teneurin transmembrane protein 4 is not a cause for essential tremor in a
Canadian population. *Mov. Disord.* **32,** 292–295 (2017).
8.　Tanner, C. M. *et al.* Essential tremor in twins: an assessment of genetic vs environmental
determinants of etiology. *Neurology* **57,** 1389–91 (2001).

9.   Müller, S. H. *et al.* Genome-wide association study in essential tremor identifies three new
loci. *Brain* **139,** 3163–3169 (2016).
10.   Merner, N. D. *et al.* Exome Sequencing Identifies FUS Mutations as a Cause of Essential
Tremor. *Am. J. Hum. Genet.* **91,** 313–319 (2012).
11.   Applegate, L. M. & Louis, E. D. Essential tremor: Mild olfactory dysfunction in a
cerebellar disorder. *Parkinsonism Relat. Disord.* **11,** 399–402 (2005).
12.   Shah, M., Muhammed, N., Findley, L. J. & Hawkes, C. H. Olfactory tests in the diagnosis
of essential tremor. *Parkinsonism Relat. Disord.* **14,** 563–568 (2008).
13.   Mori, Y. *et al.* Primary structure and functional expression from complementary DNA of a
brain calcium channel. *Nature* **350,** 398–402 (1991).
14.   Hor, H. *et al.* Missense mutations in TENM4 , a regulator of axon guidance and central
myelination, cause essential tremor. *Hum. Mol. Genet.* **24,** 5677–5686 (2015).
15.   Kim, J., Basak, J. M. & Holtzman, D. M. The role of apolipoprotein E in Alzheimer's
disease. *Neuron* **63,** 287–303 (2009).
16.   Patro, R., Duggal, G., Love, M. I., Irizarry, R. A. & Kingsford, C. Salmon provides fast
and bias-aware quantification of transcript expression. *Nat. Methods* **14,** 417–419 (2017).
17.   Pimentel, H., Bray, N. L., Puente, S., Melsted, P. & Pachter, L. Differential analysis of
RNA-seq incorporating quantification uncertainty. *Nat. Methods* **14,** 687–690 (2017).
230

Data availability statement

232 C.L. conducted the experiments, analyses and drafted the manuscript. F.S., V.V., D.R., F.A., S.D.,

233 Q.H., H.C. helped with experiments. A.D.L. and D.S. helped with bioinformatic analyses.

234 P.A.D., G.A.R oversaw the study and drafting the manuscript.

Figures and Tables

Figure 1. The RNA expression of *STK32B* overexpressed cells compared to empty-vector controls based on RNA sequencing data. Error bars show the technical variance determined with 200 bootstraps. Abbreviations: TPM, transcript per million.

Figure 2. Volcano plot of comparing overexpressed *STK32B* to empty-vector controls.
Differentially expressed genes (q < 0.05) are shown in red. Beta value was determined with a
Wald's test.

**Figure 3. Principle component analysis plot of RNA sequencing data of *STK32B*
overexpressed cells and empty-vector controls.** Overexpressed cells shown in blue and controls
shown in red.

Figure 4. Gene network of differentially expressed genes based on co-expression of RNA sequencing data for 31,499 samples. Each unique colour represents a cluster.

261 **Table 1. Gene ontology and pathway analyses of differentially expressed genes.**

Gene Set ID	Gene Set Name	P	Database
KEGG_OLFACTORY_TRANSDUCTION	Olfactory transduction	2.68E-39	KEGG
KEGG_PATHWAYS_IN_CANCER	Pathways in cancer	2.96E-28	
KEGG_FOCAL_ADHESION	Focal adhesion	1.19E-26	
KEGG_BLADDER_CANCER	Bladder cancer	6.53E-25	
KEGG_ECM_RECEPTOR_INTERACTION	Ecm-receptor interaction	8.38E-25	
GO:0001650	fibrillar center	1.13E-29	GO Cellular
GO:0031012	extracellular matrix	8.57E-27	
GO:0001725	stress fiber	1.24E-26	
GO:0005925	focal adhesion	2.82E-23	
GO:0015629	actin cytoskeleton	4.02E-22	
REACTOME:R-HSA-381753	Olfactory Signaling Pathway	3.22E-46	Reactome
REACTOME:R-HSA-418555	G alpha (s) signalling events	1.19E-29	
REACTOME:R-HSA-3000170	Syndecan interactions	1.53E-27	
REACTOME:R-HSA-2022090	Assembly of collagen fibrils and other multimeric structures	4.31E-25	
REACTOME:R-HSA-1474290	Collagen formation	2.35E-24	
GO:0050911	detection of chemical stimulus involved in sensory perception of smell	1.00E-44	Go Processes
GO:0007411	axon guidance	9.50E-36	
GO:0048146	positive regulation of fibroblast proliferation	1.31E-33	
GO:0070588	calcium ion transmembrane transport	3.02E-31	
GO:0050907	detection of chemical stimulus involved in sensory perception	3.14E-31	

265 **Table 2. Gene ontology and pathway analyses of differentially expressed gene for the three largest clusters based on co-**
266 **expression.**

Gene Set Name	P	Database	Cluster
Regulation of actin cytoskeleton	1.03E-13	KEGG	1
Pathways in cancer	1.23E-13		
Axon guidance	1.09E-12		
Focal adhesion	4.96E-12		
Arrhythmogenic right ventricular cardiomyopathy (ARVC)	4.63E-09		
fibrillar center	1.13E-29	GO Cellular	
extracellular matrix	8.57E-27		
stress fiber	1.24E-26		
focal adhesion	2.82E-23		
actin cytoskeleton	4.02E-22		
actin binding	3.66E-15	GO Function	
laminin binding	2.37E-13		
insulin-like growth factor binding	9.72E-13		
collagen binding	3.73E-12		
protein heterodimerization activity	5.54E-12		
cell migration involved in sprouting angiogenesis	7.07E-18	GO Processes	
cell migration	2.94E-16		
inactivation of MAPK activity	3.31E-15		
cytoskeleton organization	6.34E-15		
negative regulation of cell migration	9.12E-15		
Cell-extracellular matrix interactions	7.16E-15	Reactome	
Signal Transduction	1.98E-13		
Non-integrin membrane-ECM interactions	2.58E-12		
Signaling by Rho GTPases	1.84E-11		

EPHA-mediated growth cone collapse	2.30E-11		
Rna polymerase	4.57E-13	KEGG	2
Pyrimidine metabolism	1.37E-12		
Alzheimer's disease	3.55E-11		
Purine metabolism	4.11E-11		
Gnrh signaling pathway	1.84E-09		
nucleolus	5.96E-19	GO Cellular	
preribosome, large subunit precursor	2.21E-18		
small-subunit processome	5.13E-18		
DNA-directed RNA polymerase III complex	5.10E-15		
DNA-directed RNA polymerase I complex	1.85E-14		
pseudouridine synthase activity	7.75E-16	GO Function	
RNA binding	1.25E-15		
snoRNA binding	4.90E-14		
rRNA binding	3.55E-13		
ATP-dependent RNA helicase activity	5.23E-13		
RNA secondary structure unwinding	1.57E-16	GO Processes	
tRNA modification	7.73E-16		
positive regulation of gene expression, epigenetic	3.09E-15		
ribosomal large subunit biogenesis	3.17E-15		
maturation of SSU-rRNA from tricistronic rRNA transcript (SSU-rRNA, 5.8S rRNA, LSU-rRNA)	3.32E-15		
rRNA modification in the nucleus and cytosol	3.38E-18	Reactome	
tRNA processing	1.65E-17		
tRNA modification in the nucleus and cytosol	1.66E-16		
rRNA processing	9.91E-15		
tRNA processing in the nucleus	1.69E-14		
RIG-I-like receptor signaling pathway	1.24E-10	KEGG	

Cytosolic DNA-sensing pathway	1.92E-07	
O-glycan biosynthesis	1.163E-06	
Focal adhesion	1.36893E-06	
Calcium signaling pathway	2.06986E-05	
caveola	2.00E-08	
proteinaceous extracellular matrix	7.41E-08	
postsynaptic membrane	2.45E-06	GO Cellular
microfibril	4.99E-06	
extracellular matrix	1.12E-05	
type I interferon receptor binding	3.64E-07	
neuropeptide binding	5.89E-07	
scavenger receptor activity	6.23E-07	GO Function
calcium-dependent phospholipid binding	9.44E-07	
RNA binding	2.46E-06	
response to mechanical stimulus	8.42E-09	
lung development	1.55E-08	
negative regulation of neuron apoptotic process	5.29E-08	GO Processes
cellular response to hypoxia	7.29E-08	
regulation of type I interferon-mediated signaling pathway	1.11E-07	
O-glycosylation of TSR domain-containing proteins	3.73E-10	
Defective B3GALTL causes Peters-plus syndrome (PpS)	5.15E-10	
DDX58/IFIH1-mediated induction of interferon-alpha/beta	1.29E-09	Reactome
O-linked glycosylation	4.91E-08	
Diseases associated with O-glycosylation of proteins	1.04E-07	

268