1

Title: Experimental evidence that belowground competition can influence the evolution of root traits in two closely related morning glory species

Sara M. Colom[1,2], Regina S. Baucom[1]

[1]4034 Biological Sciences Building
Ecology and Evolutionary Biology Department
University of Michigan
Ann Arbor, MI 48109
scolom@umich.edu
1 (734) 647-8490 (phone)
[2]corresponding author

Abstract

Although root traits play a critical role in mediating plant-plant interactions and resource acquisition from the soil environment, research examining if and how belowground competition can influence the evolution of root traits remains largely unexplored. Here we examine the potential that root traits may evolve as a target of selection from interspecific competition using *Ipomoea purpurea* and *I. hederacea,* two closely related morning glory species that commonly co-occur in the United States. We show that belowground competitive interactions between the two species can alter the pattern of selection on root traits in each species. Specifically, competition with *I. purpurea* changes the pattern of selection on root angle in *I. hederacea*, and competitive interactions with *I. hederacea* changes the pattern of selection on root size in *I. purpurea*. However, we did not uncover evidence that intraspecific competition altered the pattern of selection on any root traits within *I. hederacea*. Overall, our results suggest that belowground competition between closely related species can influence the phenotypic evolution of root traits in natural populations. Our findings provide a microevolutionary perspective of how competitive belowground interactions may impact plant fitness, potentially leading to patterns of plant community structure.

Keywords: character displacement, root trait evolution, natural selection, *Ipomoea*

89
90   **Introduction**
91
One of the key reasons plant species are thought to coexist in a given habitat is through a niche
partitioning (Aarssen 1997; Huston 1997; Tilman et al. 1997b, 2001; Loreau 2000). Such niche
partitioning is hypothesized to occur following competitive exclusion (competitive exclusion under
limiting similarity; Gause 1936; Hutchinson 1957; Hardin 1960; Macarthur and Levins 1967), or from
trait divergence stemming from competitive interactions between species (*i.e.* character displacement;
Brown and Wilson 1965; Pfennig and Pfennig 2009). Because of the relevance of these ideas to the
formation of plant community structure, there is a large literature examining competitive interactions
among plants (Faget et al. 2013). Most of this work, however, focuses on above-ground interactions, and
as a result, little is known about root-root interactions belowground.
101
Roots, which provide a vital resource acquisition function for the plant (Fitter, 2002), also serve
as a structure through which plants experience competitive interactions with neighboring plants, whether
indirectly through alterations of the soil environment—*i.e.* reduction of water and nutrients—or directly
by the excretion of signaling and/or allelopathic molecules (Shenk, 2006 & Callaway, 2002). Thus,
specific features of the root system likely represent an important target of selection from belowground
competition. However, current evidence determining if belowground competition serves as an agent of
selection on root traits is strikingly absent from the literature.
109
The plant root system is a complex multicellular organ that can be roughly characterized into both
size and architectural traits. Root surface area, width, and root length are size proxies whereas traits
describing the spatial organization the root system, such as root angle, lateral root branching pattern, and
internode branching distance are root architecture traits (Fitter et al. 1991; Lynch, 1997). These
morphological and architectural root phenotypes strongly influence how a plant accesses nutrients and
water (Fitter et al. 1991; BassiriRad, 2005; Manschadi et al. 2006; *reviewed in* Lynch 2007; Kellermeier
and Amtmann, 2013). For example, shallow root architectures are linked to increases in the uptake of
immobile resources such as phosphorus (Lynch and Brown, 2001; Beebe et al. 2006; Lambers et al.
2006), whereas deep root architectures are linked to an increase in water uptake (Rojas Pierce et al. 2005;
Ho et al. 2005). Thus, shallow root systems may be more advantageous and lead to higher fitness in
nutrient-limited soils, and deeper root systems may provide a fitness advantage in water limited
environments. Further, traits such as root surface area and root system size, both increase resource uptake
and drought tolerance since they allow the plant to access more soil; as such, larger root systems would be
expected to provide a fitness advantage. How these different root traits may influence fitness in the field
is most often studied in crop plants (Lynch 2007), leading to a significant gap in our understanding of the
factors that influence root trait evolution in natural plant populations. In light of this, how belowground
interactions between competing species can mediate plant resource acquisition patterns and potentially
alter selection on root architecture and size traits—especially in wild plant species—is largely unknown.
128
129   There are plausible reasons to expect competitive interactions to influence the adaptive evolution
130 of root traits. The intensity of competition between plants is greater when rooting zones overlap (reviewed
131 in Casper and Jackson, 1997; Casper et al. 2003; Rubio et al. 2001), indicating that occupying the same
132 belowground niche has a deleterious effect on plant fitness. Although studies hint that differences in root

systems between competitors leads to higher fitness (*reviewed in* Silvertown, 2014), most of the research characterizes the root system at a course level (*e.g.* belowground biomass, root length density; Poorter and Ryser, 2015), and has yet to include specific root size and architectural traits. Competition between co-occurring, closely related species can be especially intense due to greater overlap in physical space or niche use (Macarthur and Levins, 1967; Pfennig and Pfennig 2009; Burns and Strauss, 2011). To reduce the effects of this competition, selection would be expected to favor the divergence of root traits (*e.g.*, root angle, root length, and overall root system size) that play important roles in water and nutrient acquisition.

       Despite this expectation, there are other explanations for particular root trait shapes or sizes in a species. As above, roots may evolve shallower, deeper, or larger root systems (among other changes) to optimize resource uptake in particular soil environments (Manschadi et al. 2006; Ferguson et al. 2016). Thus, specific root traits may reflect responses to factors in the environment that are distinct from competition. The only way to differentiate competition from other environmental factors that influence root trait evolution is to manipulate the presence of the competitor under otherwise identical conditions and determine if the pattern of selection on the trait is altered as a result (Wade and Kalisz, 1990; Dudley, 1996; Mauricio and Rausher, 1997). Markedly, while there are many studies assessing how competition influences plant fitness (reviewed in Casper and Jackson, 1997 and Faget et al. 2013), there are no studies, to our knowledge, that have examined the potential that competition from a closely related species acts as an agent of selection on root morphology.

       The purpose of this work is to determine if belowground competition between two morning glory species—*Ipomoea purpurea* and *I. hederacea*—can influence the phenotypic evolution of root traits in either species. *I. purpurea* and *I. hederacea* are two closely related vines and are common weeds of agriculture in the southeastern and Midwest US. They are most commonly found growing naturally in agricultural fields or in areas of high disturbance (Baucom et al. 2011). In some fields, both species are found to co-occur and intensely compete by vining together above ground; in other fields only one of the species may be present (personal observation, RS Baucom). Previous work has established that competition from one species can alter the pattern of selection on the other. Smith and Rausher (2008) manipulated the presence of *I. purpurea* and experimentally showed that competition between the two species for pollinators can lead to divergence in the floral morphology of *I. hederacea*. Because these species interact in other ways, and share similar morphology as well as resource needs, it is likely that other competitive interactions between the two can lead to trait divergence—namely, root trait divergence following belowground competitive interactions.

       Here, we examine the potential that competitive interactions between these two closely related species can drive the evolution of root traits. We first characterize the extent of phenotypic overlap in early growth root traits between *I. purpurea* and *I. hederacea* and then examine the potential for genetic variation underlying these traits. We likewise investigate the potential that natural selection can drive the evolution of root traits in field conditions. We specifically asked the following questions: How do root traits vary within and between species, and to what extent do the species exhibit phenotypic overlap? Is there evidence for genetic variation underlying root traits of either species, indicating that traits can respond to selection? Does belowground interspecific competition between *I. purpurea* and *I. hederacea* impose selection on root traits, and is there evidence that within-species competition (specifically, *I.*

hederacea-I. hederacea competition) similarly acts as an agent of selection? Because the adaptive
potential of traits can be obscured by plasticity when in competition, we also examine the potential that
the presence of a competitor can directly impact root phenotypes. We found early growth root traits to
differ significantly between the species, but still exhibit considerable overlap; we likewise found evidence
for genetic variation in root traits. Importantly, we show that patterns of selection on root traits change
according to the presence of an interspecific competitor, and that this varies according to species:
competition with *I. purpurea* changes the pattern of selection on root angle in *I. hederacea*, and
competitive interactions with *I. hederacea* alters the pattern of selection on root size in *I. purpurea*.
Interestingly, we found no evidence that within-species competition in *I. hederacea* influenced the pattern
of selection on root traits. These differential patterns of selection on root traits supports the idea that
competitive interactions between two closely related species can influence root trait evolution.

Study system

The common morning glory, *Ipomoea purpurea* (L.) Roth (Convolvulaceae) and ivy leaf morning
glory, *I. hederacea* (L.) Jacquin are self-compatible annual climbing vines that commonly co-occur
throughout the eastern United States. The two closely related sister species occur in similar habitat types
(*e.g.* side of train tracks, agricultural fields, road sides and waste areas). Both species germinate between
the months of May and August and begin to flower about six weeks after germination and continue to
flower until they die at first frost. The species have similar above-ground growth patterns and produce
long stems that branch occasionally. *I. purpurea* is larger (up to 10' long) compared to *I. hederacea* (up to
6' long). Belowground, *I. purpurea* and *I. hederacea* have fibrous root systems consisting of a primary
root with branched lateral roots, and both species vary greatly in the degree of lateral root branching
(*personal observation*).

The history of coexistence between *I. purpurea* and *I. hederacea* is only partially known.
Evidence suggests that *I. purpurea* is native to Central America (Gray 1886; Barkley 1986; Hickman
1993, Fang, 2013), and it has been present in the eastern United States since at least the early 1700s
(Pursh 1814). In contrast, *I. hederacea* has been in the United States for at least 150 years (Bright 1998)
but whether *I. hederacea* is native to the United States (Mohr 1901; Stevens 1948) or was introduced
from tropical America is uncertain (Shreve et al. 1910; Strausbaugh and Core 1964; Wunderline 1982;
Mahler 1984).

Materials and methods

*Plant material and growth conditions—*We performed complementary greenhouse and field studies to
investigate the potential that root traits of these two sister *Ipomoea* species could respond to natural
selection. To generate experimental seeds for our common garden and field experiments we selfed 25 and
35 maternal lines of *I. purpurea* and *I. hederacea*, respectively, which were previously sampled as seed
from five populations located in Pennsylvania and Ohio. Seeds were scarified and planted in a
randomized design in the Matthaei Botanical Gardens (Ann Arbor, MI) greenhouse in November of 2015
and plants were allowed to set seed from selfing for all subsequent experiments.

Greenhouse experiment

We performed a greenhouse experiment to characterize early growth root traits between and
within *I. purpurea* and *I. hederacea* in the summer of 2016. We planted replicate once-selfed seeds in
custom built rhizotrons containing generic potting soil (Figure 1) in greenhouse conditions at the Matthaei
Botanical Gardens (Ann Arbor, MI). Rhizotrons consisted of 8" x 10" frames made out of cut pieces of
corrugated plastic and a transparent polystyrene sheet held to the frame by duct tape. Each rhizotron was
filled with approximately 20.45 grams of soil and a single seed was placed in the center of the rhizotron
approximately one inch below the soil surface against the transparent polystyrene sheet.

We planted three replicates per maternal line per species in the rhizotrons and positioned the
rhizotrons in custom-built wooden frames at 30° in a completely randomized design (see Figure A1 in
appendix A for root image from rhizotron, and instructions on building rhizotron frames in app. Figure
B1; both apps. A and B are available online). We replicated this experiment in its entirety, twice. Thus,
we planted 150 individuals of *I. purpurea* (3 biological replicates x 25 maternal lines x 2 experimental
replicates) and 210 individuals of *I. hederacea* (3 biological replicates x 35 maternal lines x 2
experimental replicates) for a total of 360 individuals. We watered each individual daily by hand to
standardize water availability across all individuals for three weeks.

*Greenhouse root phenotyping—*One week after germination, we scanned each rhizotron to measure
below-ground root traits using a CanoScan
LIDE 110® scanner bed. For each image, we
traced the roots in Photoshop version CS6 to
facilitate automated quantification of root size
based on their pixels in ImageJ version 3.0
(Abràmoff and Magalhães 2004).
246

We focused on root size and root
architecture by measuring root system pixels
(root size), root system width, primary root
length and root angle (Figure 1). We elected to
focus on these specific traits because they are
relatively straightforward to measure across
different growing conditions and they also play
a vital role in plant resource use and uptake
(Wasson et al. 2012; Paez-Garcia, et al. 2015).
Prior to data collection, ImageJ was first
globally calibrated with the set scale tool in
order to obtain measurement in metric units for
all of the following procedures. To obtain
primary root length, root system width and root
angle, first we used the multi-pointer tool and
placed a total of four points along the root tips
of the root system image in the following order:
1) primary root at the root stem surface, 2) root

Figure 1. Example of an *Ipomoea* individual growing in rhizotron containing soil (left) and root traced in photoshop (right). Landmarks are placed to estimate root system width (distance between landmark B and D), primary root length (vertical distance between landmark A and C) and left and right root angle (inner angle formed between A, C with respect to B and D, separately for the left and and right root angles, respectively). The grey dashed line indicates the distance calculated for root system width and primary root length, and the red shaded region and solid red arcs indicate the root angle estimated for left and right angle, respectively.

tip of the left outermost root, 3) primary root tip and 4) root tip of the right outermost root.

We used R to calculate primary root length, root system width and root angle. For primary root length, we calculated the vertical distance between the primary root at the soil surface and the tip of the primary root. We estimated root system width as the euclidean distance between the outermost lateral root tips. To estimate root angle we used the cosine formula,

$$\cos^{-1}\left(\frac{b2+c2-a2}{2bc}\right),$$

on a right triangle formed by the primary root (the longest root perpendicular to the soil surface), and each of the outer lateral roots. Here, b is the distance between the primary root and the outermost lateral root tip, c is a measure of the length of the primary root calculated above, and a is the length between the outermost lateral root tip and the primary root tip (Figure 1). Root angle was calculated for both the right and left lateral roots, separately (R script in supplementary), averaged, and reported in degrees. Finally, to obtain root size, we converted the traced root images into binary images, selected 'area' as a measurement output in the 'measurement' option, and performed the 'Analyze Particles' function in ImageJ. This function quantified the total number of black pixels (all pixels from the root system) and reported the values in centimeters squared.

Field experiment

*Field design and planting—*We conducted a field experiment in the summer of 2017 to characterize root trait variation in *I. purpurea* and *I. hederacea* grown in field conditions and to determine if natural selection acts on root traits in the context of interspecific competition. We planted replicate selfed seed from maternal lines of *I. purpurea* and *I. hederacea* in two treatments: 'alone', competition (hereafter, 'competition' treatment) in a field plot at Matthaei Botanical Gardens (Ann Arbor, MI) on June 2, 2017. We likewise planted replicate selfed seed from *I. hederacea* maternal lines in the presence of intraspecific competition to determine if, at least for this species, within-species competition could influence the evolution of root traits. We used eight maternal lines of *I. purpurea* and *I. hederacea* from a single population from Pennsylvania (PA4). We decided to use maternal lines from this population since preliminary greenhouse data demonstrated high phenotypic overlap between both species for this population (Figure B2). We planted eight replicates of each maternal line per species across four blocks for our alone treatment (8 maternal lines x 2 biological replicates x 4 blocks x 2 species = 128 plants). For our interspecific competition treatment we paired each maternal line of *I. purpurea* with each maternal line of *I. hederacea* for each possible pairwise combination, and planted 2 replicates of each pairing across 4 blocks (64 unique interspecific competition combinations x 2 biological replicates x 4 blocks = 512 individuals x 2 species = 1024 plants). For our intraspecific competition treatment we planted 2 replicates of each possible combination of the 8 *I. hederacea* maternal lines within each block (28 unique interspecific competition combinations x 2 biological replicates x 4 blocks = 224 experimental units x 2 plants = 448 plants). Thus, each pairwise competition pairing was replicated 8 times across the experiment. It is important to note that although we were likewise interested in the potential that *intraspecific* competition in *I. purpurea* could influence root trait evolution in this species, we elected to examine this only in *I. hederacea* due to both field space limitations and the experimental difficulty of phenotyping large numbers of root systems in the field.

Seeds were planted into experimental units (*i.e.* cell with either a single plant for the absence of
competition, or two plants for the competition environment) which were arrayed across the four spatial
blocks in a completely random block design. Experimental units were spaced uniformly 1m from each
other; individuals in competition treatments were planted four inches apart from each other within their
experimental unit. One week after planting we scored germination. Due to intense drought, < 48% of
seeds germinated overall. We thus planted a second experimental cohort (cohort 2) on June 19, 2017 to
increase sample size. We ended with a total 1177 plants of which 670 plants were in interspecific
competition, 341 plants were in intraspecific competition, and 166 plants in the alone treatment; of our
final sample, 56% came from cohort 1 and 44% came from cohort 2. Throughout the timespan of the field
experiment we kept the immediate surroundings of each experimental unit (~6 inches from base of plants)
clear of weeds. Three weeks after the first planting date we placed 1m tall bamboo stakes at the base of
every experimental plant at a 45° angle, which allowed us to train vines of competing plants away from
another, thus removing the potential for above-ground competition.
321

*Field root excavation—*To characterize the phenotypic variation of root traits of *I. purpurea* and *I.*
hederacea grown in field conditions with and without competition, we adapted the 'shovelomics'
excavation method described by Colombi et al. (2015). We harvested roots after three weeks of growth on
a subset of plants in the field. For root phenotyping, we sampled individuals only from cohort 1 because
individuals from cohort 2 were small and not reproductively mature, whereas most individuals of cohort 1
had developed flower buds. We sampled between two to four replicates per maternal line from both
competition treatments—specifically, we sampled a total of 165 *I. purpurea* individuals, (N = 23 and N =
142 from the alone and interspecific competition treatments, respectively), and a total of 304 *I. hederacea*
individuals (N = 31, N = 132, N = 141 from alone, interspecific and intraspecific competition treatments,
respectively). To excavate roots, we cut the stem two inches from the soil surface, marked the side of the
stem facing the competitor with a permanent marker and then dug the root system with a shovel by

placing the shovel head at 45 degree
angle, six inches from the plant stem.
This method unearthed the first six
inches of the root system. The
excavated root core was then shaken
gently to remove adhering soil and
placed in plastic bags for root imaging.
340

Field root imaging and phenotyping—
Root imaging was carried out indoors
with the use of a cubic photo shooting
tent (MVPOWER, 16"x16"x16") in order
to standardize imaging between samples
and facilitate the use of REST (Colombi
et al. 2015), an automated root
phenotyping program developed to
characterize the root systems of plants
grown in the field. To image the root
system, roots were hung in the center of

Figure 2. REST output showing the original image of an *Ipomoea* species excavated from the field (A) and its binary form with an arc superimposed by REST to obtain the outer right and left root angles (here 30 degrees on the left and 48.3 degrees on the right) from the horizontal place in red. (B) The blue box shows the region of interest detected automatically by REST program of which total root based pixels were quantified from within to obtain root size ('area convex hull' in REST) and measure root system width ('max root width' in REST) based on the distance between the right and left outermost roots in the box.

the photo shooting box and photographed with a Canon EOS Rebel XSi 12.2 MP (18-55mm IS Lens).
Images were imported into REST (Colombi et al. 2015), and we manually specified where the stem at the
soil surface was for each image (shown in Figure 2). After user specification of the stem/soil surface,
REST draws a rectangular region of interest for pixel analysis to standardize measurements across
images. All root measurements were quantified from the pixels lying within this region of interest. REST
returned the root angle (right and left root angle), root system width ('Max width') and a root system size
proxy ('Area convex hull'), among other morphological and architectural traits. We focused on these
three root traits because they are similar to the traits captured in our greenhouse study. Root angle (left
and right) are determined by calculating the outermost angle between the top most lateral root and the soil
surface plane at the plant stem, and then subtracting this value from a perpendicular line (i.e. 90°) drawn
at the plant stem. The root system width and root system size estimates captured in REST are also the
same as the measures taken in the greenhouse rhizotron study as they were estimated from the total
number of root derived pixels, respectively. We do not have data for primary root length from plants
grown in the field because this trait was destroyed in the process of sampling the roots.
366
Field plant fitness data—We began to collect mature fruit in the month of September and continued to do
so until late October when all plants have senesced. The entire aboveground of all remaining plants were
collected by first frost and seeds were manually removed, cleaned, and counted with a seed counter to
obtain an estimate of total fitness. We sampled 165 individuals of *I. purpurea* (N = 23 and N = 142, from
alone and competition treatments, respectively), and 304 individuals of *I. hederacea* (N = 31, N = 132,
and N = 141, from alone, interspecific and intraspecific competition treatments, respectively). Ultimately,
we sampled seeds from a total of 508 plants; 27% of these individuals came from cohort 1 and 72% from
cohort 2.
375
376 **Data analysis**
377
Greenhouse experiment—All statistical analyses were carried out in R (version 3.3.1). We fit separate
linear mixed models using the lmer function of the lme4 package (Bates et al. 2015) for each of the root
traits measured to test for the presence of variation in root traits between species, populations, and
maternal lines. Each respective linear mixed model consisted of the root trait as the response variable,
species, population and experimental replicate ('Experiment') as fixed effects and maternal line as a
random effect. To ascertain the significance of the predictor variables we used F-statistics for the fixed
effects and a log likelihood ratio test (using the anova and ranova functions from the lmerTest package;
Kuznetsova et al. 2016) for the random effect. We ran additional linear mixed models for each species
separately to test for evidence of maternal line variation within *I. purpurea* and *I. hederacea*, where root
trait was the response variable, population and experiment were fixed effects and maternal line was a
random effect. We further examined how roots varied between species in trait space by performing
principal component analysis (PCA) using a correlation matrix of all root traits measured in the
greenhouse with the 'PCA' function from the FactoMineRPackage (Lê et al 2008).
391
Field experiment—To examine how root traits vary between *I. purpurea* and *I. hederacea* grown in field
conditions, and to determine if root phenotypes were influenced by competitive interactions, we ran
mixed linear models as above. We fit a separate model for each root trait where the trait was the response
variable and block, treatment, and block *x* treatment interaction were fixed effects and maternal line and

maternal line *x* treatment interaction were random effects. Preliminary analyses indicated there were no
significant maternal line *x* treatment interactions for any trait. We thus elected to exclude these effects
from our final models. As above, we visualized phenotypic variation in root traits between species when
grown in field conditions by performing principal component analysis (PCA) with a correlation matrix on
all traits including root system size, root system width and average root angle. In addition, we generated a
correlation matrix using the family mean values for all the root traits measured for each species separately
to examine relationships between the three traits.
403
Selection analyses—We used genotypic selection analyses (Lande and Arnold, 1983; Rausher 1992) to
estimate selection gradients on each root trait in each competition environment, and ANCOVA to
determine if competition and experimental block altered selection on root traits of the two species. We
elected to perform a joint selection analysis using maternal line averages of the root traits because it
allowed us to examine direct selection acting on each trait while controlling for environmentally induced
biases (Rausher 1992). We estimated selection gradients on root system width, root system size, and root
angle of both species in each competitive treatment environment (alone and interspecific competition) by
performing multiple regression with the focal root traits included as predictor variables and relative
fitness (total seed number) as the dependent variable. For all selection analysis, we used mean
standardized root trait values and untransformed relative fitness. Preliminary analysis indicated that
individuals of both species from the second cohort produced significantly fewer total seeds than
individuals from the first cohort (*I. purpurea*: F = 100.3, p-value < 0.001; *I. hederacea*: F = 213.9, p-
value < 0.001), but preliminary analyses also provided no evidence that selection gradients differed
between cohorts within either species for any root trait. Thus we elected to combine cohorts in the
genotypic selection analyses (cohort 1 N = 141 and cohort 2 N = 367). Further, while we examined the
potential for non-linear selection influencing root traits in preliminary analyses, we did not find evidence
of either stabilizing or disruptive selection acting and thus present only the results of linear selection
analyses.
422
We used analysis of covariance (ANCOVA) to determine if the direction and/or intensity of
selection varied between the presence and absence of competition (Wade and Kalisz, 1990) separately for
each species. For *I. purpurea*, we compared selection gradients between plants grown in interspecific
competition or grown alone, and for *I. hederacea*, we compared selection gradients from inter- or
intraspecific competition with that of plants grown alone. In each analysis, models included competition
treatment, block, the standardized root trait values, and all interactions as predictors of relative fitness.
Significant interactions between the competition treatment and standardized root traits indicate that
selection gradients differed between treatments. Block and block *x* treatment interactions were likewise
included within the ANCOVAs.
432
433 **Results**
434
Greenhouse experiment—In our greenhouse rhizotron study assessing early root traits, we found
significant variation between species in root system width and average root angle (Table 1). The root
system of *I. hederacea* was wider (8.49 cm, Table 1) than that of *I. purpurea* (6.92 cm, Table 1) and the
overall root size of *I. hederacea* was on average greater (3.95 cm^2, Table 1) than *I. purpurea* (2.37 cm^2,
Table 1). *I. purpurea* exhibited lateral roots that were closer to the soil surface (root angle: 37.33 degrees
on average, Table 1) compared to *I. hederacea* (30.44 degrees on average, Table 1). Although species

varied in the above traits (Table 1), a visualization of the four root traits in a PCA identified considerable
overlap of root phenotypes between species (Figure B1-A). The root system width, root angle and
primary root length loaded most strongly on the first principal component, which captured 37.3% of the
total variation (Figure B1-B), and root size loaded most strongly on the second principal component,
which explained 29.1% of the total variation (Figure B1-C). These first two PCA's can thus serve as
descriptors of root system architecture (i.e. spatial arrangement of root system) and root size, respectively.
447
448 **Table 1**. Species differences in *I. purpurea* and *I. hederacea* root traits measured from rhizotrons in the greenhouse.
449 Least-squares means ±1 SE for each trait by species and F-statistics and likelihood ratio test statistics (χ^2) values
450 showing the effects of experimental replicate, species, population, and maternal line variation on plant phenotypes.
451 Maternal lines were nested within populations.

	Species		F-statistics			
Trait	*I. purpurea*	*I. hederacea*	Experiment	Species	Population	Mat. Line
Root system width (cm)	6.92 ± 0.28	8.49 ± 0.21	**317.78**	**16.70**	**6.78**	**$\chi^2 = 4.96$**
Primary root length (cm)	11.35 ± 0.61	11.39 ± 0.47	**22.74**	0.01	0.77	**$\chi^2 = 5.28$**
Root angle (degrees)	37.45 ± 1.33	30.41 ± 1.02	**62.15**	**15.26**	0.21	**$\chi^2 = 12.51$**
Root system size (cm^2)	2.37 ± 0.21	3.95 ± 0.16	**35.32**	**31.47**	**7.30**	$\chi^2 = 2.22$

452 The denominator degrees of freedom (DenDF) was estimated using Satterthwaite's method for F-tests to obtain
453 significance values of fixed-effects. χ^2 represent the log-likelihood chi statistic.
454
Additionally, we found evidence for both population and maternal line variation in root traits. We
found significant population variation for root system width and root system size, and variation among
maternal lines for root system width, root angle, and primary root length (Table 1). Separate mixed
models, performed per species, identified significant maternal line variation within *I. purpurea* for root
system width ($\chi^2 = 7.30$, p-value = 0.01), root size ($\chi^2 = 3.70$, p-value = 0.05) and root angle ($\chi^2 =$
4.05, p-value = 0.04). We identified maternal line variation within *I. hederacea* for root angle ($\chi^2 = 8.63$,
p-value < 0.01), and marginally significant maternal line variation for primary root length ($\chi^2 = 3.10$, p-
value = 0.07).
463
Field experiment—A visualization of root system width, size, and root angle in a principal component
analysis showed a high overlap between species in root phenotypes (Figure B2) in plants grown in the
field. We identified maternal line variation in root system width (Table 2); a within species examination
revealed this result to be driven by *I. purpurea* ($\chi^2 = 4.69$, p-value = 0.03). We further found a significant
and strong correlation between root size and root width (r = 0.85, p-value < 0.001; Table B1) in *I.*
purpurea, whereas there was evidence for strong and significant positive correlations between all root

traits within *I. hederacea* (root width and root angle r = 0.60; root size and root width r = 0.58; root size
and root width r = 0.85; p-value < 0.001 for all pairwise traits; Table B1).
472
473 **Table 2**. Influence of competitive treatment on *I. purpurea* and *I. hederacea* root traits when grown in the field. F-
474 statistics and chi-square values showing the effects of block, species, competitive environment and maternal line
475 variation on plant phenotypes. Significant fixed and random effects (p-value < 0.05) are indicated in bold.

	Block		Species		Competition		Mat. Line	
Trait	F	P	F	P	F	P	X^2	P
Root system width (cm)	1.50	0.21	0.87	0.36	0.76	0.47	**5.81**	**0.02**
Root system size (cm^2)	**29.92**	**<0.001**	0.04	0.84	2.33	0.10	0.02	0.89
Root angle (degrees)	2.54	0.06	0.06	0.80	2.04	0.13	0.07	0.79
Seed number	**9.41**	**<0.001**	0.30	0.59	**9.81**	**<0.001**	**5.43**	**0.02**

476
With the exception of a marginally significant treatment effect on root size (F = 2.33, p-value =
0.09; Table 2), we found that interspecific competition in the field did not strongly influence root
phenotypes of either species. A closer examination of the linear mixed models within species suggested
that this treatment effect likely impacts *I. hederacea* (F = 2.10, p-value = 0.12) but not *I. purpurea* (F =
0.04, p-value = 0.84). However, there was a strong effect of competition on fitness, with *I. purpurea*
experiencing a fitness reduction of 65.5%, and *I. hederacea* a reduction of 63.5% when in interspecific
competition. *I. hederacea* planted in intraspecific competition likewise experienced a significant fitness
reduction (60.3% lower than plants grown alone). Intraspecific competition between *I. hederacea* plants
led to slightly lower fitness than when grown in interspecific competition (*i.e.* 6% reduction in intra-
versus interspecific competition), but this difference was not significant (Table 3).
487
488 **Table 3.** Influence of competitive treatment on *I. purpurea* and *I. hederacea* root traits when grown in the field.
489 Least-squares means ±1 SE for each trait in each treatment.

	I. purpurea		*I. hederacea*		
Trait	Alone	Competition	Alone	Competition (inter)	Competition (intra)
Root system width (cm)	7.12 ±0.37	7.00 ±.0.25	7.43 ±0.36	7.31 ±0.25	7.03 ±0.25
Root system size (cm^2)	77.49 ±3.93	76.69 ±2.16	78.08±3.77	77.28 ±2.22	71.07 ±2.23
Root angle (degrees)	26.62 ±2.05	24.43 ±1.14	27.00 ±1.96	24.81 ±1.16	22.63 ±1.17
Seed number	750.01 ±68.42	491.15 ±53.88	709.84 ±68.42	450.98 ±53.48	428.26 ±55.10

490
Selection on root traits in field conditions—From our selection gradient analyses, we identified positive
linear selection on root angle in *I. hederacea* in the presence of interspecific competition (β = 0.22, p-
value = 0.03; Table 4) but no evidence of selection when when grown alone (β = -0.01, p-value = 0.88;
Table 4). ANCOVA revealed a significant treatment interaction for root angle in *I. hederacea* (F = 4.47,
p-value = 0.04; Table B2), supporting the idea that the pattern of selection for root angle differs according
to competitive context for this species. Further, we found a marginally significant block by treatment
interaction (F = 2.37, p-value = 0.08; Table B2) for root angle in *I. hederacea* when in interspecific
competition, suggesting that environmental differences can influence the strength and/or direction of
selection on this trait. In comparison, we found no evidence of selection when *I. hederacea* was grown in
intraspecific competition (β = -0.28, p-value = 0.29; Table 4; Table B2).
501
502 **Table 4.** Influence of interspecific and intraspecific competitive environment on selection gradients (multivariate
503 analyses) in *I. purpurea* and *I. hederacea*. Selection gradients that are significantly different than zero are shown in
504 bold (p-value < 0.05) whereas gradients that are marginally significant (p-value < 0.09) shown with an (^)).
505 Standard errors of the estimate presented in parentheses.Values in bold indicate significant treatment by trait, block
506 by trait, and treatment by block by trait interactions.

	I. purpurea		*I. hederacea* (inter)		*I. hederacea* (intra)
Trait	Alone	Competition	Alone	Competition	Competition
Root system width (cm)	-0.27 (0.23)	0.36 (0.21)	0.17 (0.12)	-0.21 (0.15)	-0.02 (0.14)
Root system size (cm^2)	0.59^ (0.32)	-0.14 (0.22)	-0.09 (0.13)	0.07 (0.10)	0.02 (0.18)
Root angle (degrees)	0.02 (0.10)	-0.09 (0.08)	0.01 (0.08)	**0.22 (0.09)**	-0.28 (0.25)

507
508
For *I. purpurea*, we found marginally significant positive selection for root system size when
grown in the absence of interspecific competition (β = 0.59, p-value = 0.09; Table 4), but no evidence of
selection on root size when in the presence of competition (β = -0.14, p-value = 0.52; Table 4). ANCOVA
revealed a significant treatment interaction with root size (F = 6.01, p-value = 0.02; Table B2) indicating
that selection on root size differs according to competitive context in this species. We likewise found a
significant treatment *x* block interaction for root system width within *I. purpurea* (F = 7.30, p-value
<0.01; Table B2) and a marginally significant treatment *x* block effect for root angle (F = 3.02, p-value =
0.08; Table B2), indicating that the pattern of selection on these two traits are impacted by both
competitive context and other unmeasured environmental variables (*i.e.* block effect).

Discussion

Interspecific competition imposes selection on root traits in I. purpurea *and* I. hederacea

      Given the functional importance of root systems, we hypothesized that competition between two closely related species could impose selection on root traits, and that selection could promote divergence in such traits. However, there are few, if any, examinations of the potential for selection on root traits in field conditions. Thus, we characterized the phenotypic variation in root traits of two closely related species, determined if genetic variation in these traits was present both within the greenhouse and in the field, and examined the potential that competition changed the pattern of selection on roots. From our greenhouse experiment, we found early growth root traits to differ significantly between the species, and found evidence for genetic variation underlying traits—both between population variation and maternal line variation. Results from our manipulative field experiment showed that in the absence of competition there was a trend for positive linear selection acting on root size in *I. purpurea,* but no evidence for the same pattern of selection in the presence of competition. Interestingly, we found selection acting on different traits in *I. hederacea*: in this species, we found positive selection acting on root angle when in the presence of competition with *I. purpurea*, but no evidence of selection on this trait in the absence of competition. Somewhat surprisingly, we found no evidence of selection on root angle (or any root trait) in *I. hederacea* when in the presence of *intraspecific* competition (i.e., *I. hederacea-I. hederacea* competition). Thus, competition below-ground from *I. purpurea* promotes the evolution of broader root angles (*i.e.* a more shallow root system) in *I. hederacea*, but the same effect is not seen in *I. hederacea* when in within-species competition.

      Because water, ion and minerals are heterogeneously spread in the soil according to their chemical and physical properties, differences in root architecture between plants determines what specific resources are readily available for uptake and in turn how plants compete for such resources (Lynch, 2005). This provides a likely explanation for the pattern of selection we identified for *I. hederacea* in the presence of interspecific competition; because shallow lateral roots enable the exploitation of nutrients near the soil surface, individuals with shallow roots may be at a fitness advantage when in the presence of a competitor compared to individuals with deeper root systems. In support of this idea, shallow rooting systems have been shown to be advantageous in common bean, maize and rice when grown in environments that are limited by phosphorus and other resources that accumulate in the topsoil (Rubio et al. 2003; Lynch and Brown, 2001; York et al. 2015; Sandhu et al. 2016).

      From an ecological standpoint, it is somewhat puzzling that selection favors a larger root system in *I. purpurea* when competition is absent, but not when competition is present. Larger root systems allow for greater exploitation of soil nutrients and water, and have been shown to be correlated with increased fitness in other species (Svačina and Chloupek, 2014; Ehdaie and Waines, 2008). As such, we expected to identify selection for larger root systems regardless of competitive environment. A potential explanation for our findings is that root traits that were not measured here—primary root length, lateral root placement, and/or hair root density—may play an important role in resource uptake in the presence of competition. An investment in greater root foraging precision, as well as selection on traits that optimize resource uptake efficiency could potentially reduce the deleterious effects of competition. Thus, it is possible that root size is not under selection when these two species compete because selection is instead

acting on traits that increase resource uptake efficiency (Fitter et al. 1991; Hodge et al. 2002; York et al.
2015).

Although we identified selection on only two traits—root size and angle—the strong correlations
we uncovered between root width, size, and lateral root angle suggests traits not under direct selection
will likely evolve due to indirect selection. We identified strong positive correlations between root width,
size, and angle in *I. hederacea*, indicating that width and size may evolve indirectly given selection on
root angle. In *I. purpurea*, the strong positive correlation between root width and size, and pattern of
positive selection on root size, suggests that root width should likewise experience indirect positive
selection. That we found no evidence of correlations between root angle and root width and size in *I.*
purpurea suggests root angle may evolve with fewer constraints in this species. It is likewise notable that
we uncovered genetic variation underlying only root width in *I. purpurea* in the field experiment;
however, this result is not particularly surprising given that genetic variation in field conditions is often
obscured by high environmental variation (Conner, 2003). Notably, in our greenhouse experiment, we
found evidence for both population and maternal line variation on root traits in both species, suggesting
these traits have the capacity to evolve either through selective pressures or as a result of genetic drift.

Further, while we identified different patterns of selection across the competitive environments
between the two species, we found suggestive, but limited, evidence for plasticity in the root traits of
either species as a result of competition. Plant root systems can impact the root growth of other closely
neighboring plants either indirectly *via* altering the physical and chemical soil environment and/or
directly through the excretion of signaling and/or allelopathic molecules (Schenk 2006; reviewed in
Cahill and McNickle 2013 and Depuydt, 2014). As such, we expected to find a significant treatment
effect on root trait phenotypes, and thus evidence of phenotypic plasticity in root architecture and size
traits. Other experiments characterizing root phenotypes in a range of species have found mixed results
when plants are grown in competition, ranging from genotypic- and species-specific responses in root
growth to no response whatsoever (Mahall and Callaway 1991; Falik et al. 2003; Bartelheimer et al. 2006;
Fang et al. 2013; Belter and Cahill 2015; Litav and Harper 1967; Semchenko et al. 2007). Therefore, the
results we report here suggest that these two *Ipomoea* species may lack a mechanism to modify their root
growth in competition, that plasticity may be occurring in other, unmeasured traits, or simply that the
effect sizes on root trait changes due to competition were small, and high variance due to other
environmental factors reduced our ability to identify significant plasticity in root traits given competition.

Importantly, the belowground plant-plant competition imposed by our experimental design led to
reduced fitness of both species—around 60% fewer seeds produced by each species in the presence of
competition (whether interspecific and intraspecific)—indicating that although we did not uncover root
trait plasticity, there was clearly a cost imposed by the presence of belowground competition between and
within species. We note, however, that the strongest trend in reduced root size occurred when *I.*
hederacea was planted in *intraspecific* competition relative to interspecific competition. This suggests *I.*
hederacea may potentially be decreasing overall plant growth as an adaptive response to reduce
intraspecific competition. Such a potential plastic response within *I. hederacea* when in competition with
a congener may explain why we did not detect evidence for selection from intraspecific competition on
root traits in this species.

Overall, our finding of different patterns of selection acting on root traits in the different
competitive treatments indicates that plant-plant competition can act as a selective agent on root traits.
That we identified selection on different root traits between species is consistent with the idea of niche
partitioning, which predicts greater divergence in resource associated traits between species to reduce
competition for limiting resources (Macarthur and Levins 1967). Multiple field studies examining the
relationship between different rooting depths of various co-occurring plant species have shown that a
decrease in overlap between rooting zones of neighboring plants positively impacts plant yield and
biomass (*i.e.* plant fitness; Fargione and Tilman 2005; Mueller et al. 2015). Our results extend this finding
to show that interactions between two closely related, co-occurring species elicits selection for different
patterns of root traits. Hence, it is possible that competition between the two *Ipomoea* sister species
promotes the divergence in resource-related root traits.
623
Finally, although our research provides the first experimental evidence that belowground
competition can influence the evolution of root traits in these two related species, we are not showing the
outcome of such competitive interactions across many natural populations. More specifically, while our
study supports the idea that the adaptive process can occur in root traits as a response to belowground
competition, we do not explicitly test for broad-scale patterns that would suggest such interactions have
led to trait divergence (*i.e.*, divergence in root traits where the species co-occur versus similarity in areas
where they do not co-occur). Future work testing for patterns of phenotypic evolution in root traits
between multiple naturally occurring populations of these two species is thus needed to draw conclusions
for, if and how competition belowground has influenced the evolution of root traits in natural populations
across the landscape.
634
To our knowledge, this is the first study to explicitly test the potential that root traits may exhibit
evidence of selection as a result of competitive interactions. This is surprising, since there are an
abundance of studies showing belowground competition between plants can play a major role in plant
fitness and overall community structure. One major reason research on the evolution of root traits may be
lacking is due to the highly laborious nature of sampling and phenotyping root traits. The development of
root sampling and imaging technologies is now facilitating the study of root traits across many
agricultural plants, and as such we expect more examples of similar work in wild species. The results we
present here can serve as a first step for addressing the evolution of root phenotypes in light of important
ecological processes such as plant-plant competition.

References

Aarssen, L. W. 1997. High productivity in grassland ecosystems: affected by species diversity or productive species? Oikos 80: 182–183.

Arthur, W. 1982. The evolutionary consequences of interspecific competition. Adv. Ecol. Res. 12:127–187.

BassiriRad, H. 2005. Nutrient acquisition by plants: an ecological perspective. Springer science & business media.

Bates, D., M. Mächler, B. Bolker, and S.Walker. 2015. Fitting linear mixed-effects models using lme4. Journal of Statistical Software, Articles 67 (1): 1–48.

Baucom, R. S., S-M Chang, J. M. Kniskern, M. D. Rausher, and J. R. Stinchcombe. 2011. Morning glory as a powerful model in ecological genomics: tracing adaptation through both natural and artificial selection. Heredity 107 (5): 377–85.

Baylis, A.D. 2000. Why glyphosate is a global herbicide: strengths, weaknesses and prospects. Pest Management Science 56 (4): 299–308.

Beebe, S.E., M. Rojas-Pierce, X. Yan, M.W. Blair, F. Pedraza, F. Muñoz, J. Tohme, and J.P. Lynch. 2006. Quantitative trait loci for root architecture traits correlated with phosphorus acquisition in common bean. Crop Science 46: 413–23.

Briones, O., C. Montaña, and E. Ezcurra. 1996. Competition between three chihuahuan desert species: evidence from plant size-distance relations and root distribution. Journal of Vegetation Science: Official Organ of the International Association for Vegetation Science 7 (3): 453–60.

Brown, W. L., and E. O. Wilson. 1956. Character displacement. Systematic Zoology 5 (2): 49–64.

Cahill, Jr, J.F., and G.G. McNickle, 2011. The behavioral ecology of nutrient foraging by plants. Annual Review of Ecology, Evolution, and Systematics 42 (1): 289–311.

Callaway, R. M., S. C. Pennings, and C. L. Richards. 2003. Phenotypic plasticity and interactions among plants. Ecology.

Callaway, R.M. 2002. The detection of neighbors by plants. Trends in Ecology & Evolution 17 (3): 104–5.

Casper, B.B., and R.B. Jackson. 1997. Plant competition underground. Annual Review of Ecology and Systematics 28 (1): 545–70.

Conner, J.K., R. Franks, and C. Stewart. 2003. Expression of additive genetic variances and covariances
for wild radish floral traits: comparison between field and greenhouse environments. Evolution;
International Journal of Organic Evolution 57 (3): 487–95.

Day, T., and K.A. Young. 2004. Competitive and facilitative evolutionary diversification. BioScience
54:101–109.

Depuydt, S. 2014. Arguments for and against self and non-self root recognition in plants. Frontiers in
Plant Science 5: 614.

Dudley, S. A. and A. L. File. 2007. Kin recognition in an annual plant. *Biol. Lett.* 3: 435–438.

Dudley, S. A. 1996. Differing selection on plant physiological traits in response to environmental water
availability: a test of adaptive hypotheses. Evolution 50: 92–102.

Ehdaie, B., and J.G. Waines. 2008. larger root system increases water–nitrogen uptake and grain yield in
bread wheat. In: Appels R et al (eds) 11th international wheat genetics symposium 2008. Sydney
University Press, Brisbane, p 659.

Faget, M., N.A. Kerstin, A.Walter, J.M. Herrera, S. Jahnke, U. Schurr, and V.M. Temperton. 2013. Root-
root interactions: extending our perspective to be more inclusive of the range of theories in ecology and
agriculture using in-vivo analyses. Annals of Botany 112 (2): 253–66.

Fargione, J., and D. Tilman, 2005. Niche differences in phenology and rooting depth promote coexistence
with a dominant c4 bunchgrass. Oecologia 143 (4): 598–606.

Ferguson, L, G., Sancho, M.T. Rutter, and C.T. Murren. 2016. Root architecture, plant size and soil
nutrient variation in natural populations of arabidopsis thaliana. Evolutionary Ecology 30 (1): 155–71.

Fitter, A. H., T. R. Stickland, M. L. Harvey, and G. W. Wilson. 1991. Architectural analysis of plant root
systems 1. Architectural correlates of exploitation efficiency. The New Phytologist 118 (3): 375–82.

Fitter, A.H. 2002. Characteristics and functions of root systems. In Plant Roots, 49–78. CRC Press.

Fitter, A.H., L. Williamson, B. Linkohr, and O. Leyser. 2002. Root system architecture determines fitness
in an arabidopsis mutant in competition for immobile phosphate ions but not for nitrate ions. Proceedings.
Biological Sciences / The Royal Society 269 (1504): 2017–22.

Gause, G. F. 1936. The struggle for existence. Soil Science 41 (2): 159.

Hardin, G. 1960. The competitive exclusion principle. Science 131 (3409): 1292–97.

Ho, M.D., J.C. Rosas, K.M. Brown, and J.P. Lynch. 2005. Root architectural tradeoffs for water and
phosphorus acquisition. Functional Plant Biology: FPB 32 (8): 737–48.

743
744 Hodge, A., D. Robinson, B.S. Griffiths and A. H. Fitter. 1999. Why plants bother: root proliferation
745 results in increased nitrogen capture from an organic patch when two grasses compete. Plant, Cell &
746 Environment 22 (7): 811–20.
747
748 Hoss, N.E., A-K. Kassim, D.E. Peterson, and T.M. Loughin. 2003. Efficacy of glyphosate, glufosinate,
749 and imazethapyr on selected weed species. Weed Science 51 (1): 110–17.
750
751 Huston, M.A. 1997. Hidden treatments in ecological experiments: re-evaluating the ecosystem function of
752 biodiversity. Oecologia 110 (4): 449–60.
753
754 Kellermeier, F., F. Chardon, and A. Amtmann. 2013. Natural variation of arabidopsis root architecture
755 reveals complementing adaptive strategies to potassium starvation. Plant Physiology 161 (3): 1421–32.
756
757 Kuznetsova, Alexandra, Per Brockhoff, and R. Christensen. 2017. lmerTest package: tests in linear mixed
758 effects models. Journal of Statistical Software, Articles 82 (13): 1–26.
759
760 Lambers, H., M.W. Shane, M.D. Cramer, S.J. Pearse, and E.J. Veneklaas. 2006. Root structure and
761 functioning for efficient acquisition of phosphorus: matching morphological and physiological traits.
762 Annals of Botany 98 (4): 693–713.
763
764 Lande, R., and S.J. Arnold. 1983. The measurement of selection on correlated characters. Evolution 37
765 (6): 1210–26
766
767 Lynch, J. 1995. Root architecture and plant productivity. Plant Physiol. 109: 7–13
768
769 Lynch, J.P. 2007. Roots of the second green revolution. Australian Journal of Botany 55 (5): 493–512
770
771 Lynch, J.P., and K.M. Brown. 2001. Topsoil foraging – an architectural adaptation of plants to low
772 phosphorus availability. Plant and Soil 237 (2): 225–37
773
774 Mamolos, A. P., G. K. Elisseou, and D. S. Veresoglou. 1995. Depth of root activity of coexisting
775 grassland species in relation to n and p additions, measured using nonradioactive tracers. The Journal of
776 Ecology 83 (4): 643–52
777
778 Manschadi, A.M., J. Christopher, P. deVoil, and G.L. Hammer. 2006. The role of root architectural traits
779 in adaptation of wheat to water-limited environments. Functional Plant Biology: FPB 33 (9): 823–37
780
781 Mauricio, R., and M.D. Rausher. 1997. Experimental manipulation of putative selective agents provides
782 evidence for the role of natural enemies in the evolution of plant defense. Evolution 51 (5): 1435–44
783
784 Macarthur, R., and R. Levins. 1967. The limiting similarity, convergence, and divergence of coexisting
785 species. The American Naturalist 101 (921): 377–85
786

787   Mueller, K.E., D. Tilman, D.A. Fornara, and S.E. Hobbie. 2013a. Root depth distribution and the
788   diversity—productivity relationship in a long-term grassland experiment. Ecology 94 (4): 787–93
789
790   Nobel, P. 1997. Root distribution and seasonal production in the northwestern sonoran desert for a c3
791   subshrub, a c4 bunchgrass, and a cam leaf succulent. American Journal of Botany 84 (7): 949
792
793   Peterson, B.G., and C. Peter 2018. PerformanceAnalytics: Econometric Tools for Performance and Risk
794   Analysis. R package version 1.5.2
795
796   Pfennig, K. S., and D. W. Pfennig. 2009. Character displacement: ecological and reproductive responses
797   to a common evolutionary problem. The Quarterly Review of Biology.
798
799   Poorter, H., and P. Ryser. 2015. The limits to leaf and root plasticity: what is so special about specific
800   root length? The New Phytologist 206 (4): 1188–90
801
802   Rausher, M.D., 1992. The measurement of selection on quantitative traits: biases due to environmental
803   covariances between traits and fitness. Evolution 46 (3): 616–26
804
805   Rubio, G., H. Liao, X. Yan, and J.P. Lynch. 2003. Topsoil foraging and its role in plant competitiveness
806   for phosphorus in common bean. Crop Science 43: 598–607
807
808   Sandhu, N., K. A. Raman, R.O. Torres, A. Audebert, A. Dardou, A. Kumar, and A. Henry. 2016. Rice
809   root architectural plasticity traits and genetic regions for adaptability to variable cultivation and stress
810   conditions. Plant Physiology 171 (4): 2562–76
811
812   Schenk, H. J. 2006. Root competition: beyond resource depletion: root competition: beyond resource
813   depletion. The Journal of Ecology 94 (4): 725–39
814
815   Schenk, H. J., and R.B. Jackson. 2002. Rooting depths, lateral root spreads and below-ground/above-
816   ground allometries of plants in water-limited ecosystems. The Journal of Ecology 90 (3): 480–94
817
818   Sebastien Le, J., Josse, F. Husson. 2008 FactoMineR: An R package for multivariate analysis. Journal of
819   Statistical Software, 25(1), 1-18
820
821   Silvertown, J. 2004. Plant coexistence and the niche. Trends in Ecology & Evolution 19 (11): 605–11
822
823   Slatkin, M. 1980. Ecological character displacement. Ecology 61:163–177
824
825   Svačina, P., T. Středa, and O. Chloupek. 2014. Uncommon selection by root system size increases barley
826   yield. Agronomy for Sustainable Development 34 (2): 545–51
827
Tilman, D., C. L. Lehman, and K. T. Thomson. 1997. Plant diversity and ecosystem productivity:
theoretical considerations. Proceedings of the National Academy of Sciences of the United States of
America 94 (5): 1857–61

Wade, M.J., and S. Kalisz. 1990. The causes of natural selection. Evolution 44 (8): 1947–55

Wasson, A. P., R. A. Richards, R. Chatrath, S. C. Misra, S. V. Sai Prasad, G. J. Rebetzke, J. A.
Kirkegaard, J. Christopher, and M. Watt. 2012. Traits and selection strategies to improve root systems
and water uptake in water-limited wheat crops. Journal of Experimental Botany 63 (9): 3485–98

York, L.M., T. Galindo-Castañeda, J.R. Schussler, and J.P. Lynch. 2015. Evolution of us maize (zea mays
l.) Root architectural and anatomical phenes over the past 100 years corresponds to increased tolerance of
nitrogen stress. Journal of Experimental Botany 66 (8): 2347–58

875 **Appendix A**
876 **Custom made rhizotron frame instructions**
877
Figure A1. Procedure and materials required for construction of rhizotron frames. Four wood boards of 8" x 1" x
10', 80 ½" x 4' wooden dowels, wood glue, power drill, measuring tape, and a pencil will be required to build a total
of four rhizotron frames. First saw each 10' wooden board in half to obtain a total of eight 5' boards and then saw
each 4' wooden dowel in half to obtain a total of 160 2' dowels. Measure and draw out 40 evenly spaced dots on the
10' board approximately two inches from the top edge. For every of the 40 evenly spaced dots, draw a dot 4"
beneath it at 30 degrees relative to the dots above. Drill a ½" hole where each dot is located. Make sure that each
board looks identical with their holes matching up. Finally, line up two boards 2' apart and insert the ends of the
dowels on the corresponding holes of the boards using wood glue at the tips to make sure they stay in place. Repeat
this process until dowels have been placed in all the holes in order to create the frame.
887

901 **Figure A2.** The rhizotron greenhouse experiment. Each rhizotron was maintained at a thirty degree angle
902 with the use of custom built wooden frames (right).

917
918
919 **Appendix B**
920
921 **Supporting Tables and Figures**
922
923 **Figure B1:** PCA summarizing root trait space between *I. hederacea* and *I. purpurea* from population PA4 grown in
924 field conditions where 62.50% and 28.40% of the phenotypic variation is explained by PCA1 and PCA2, respectively.
925 Bar graphs (B & C) show the loading scores of each root trait for PCA1 and PCA2. B) root system size and root
926 system width load heaviest in PCA1, C) and average root angle loads heaviest in PCA2.
927

951
Figure B2: PCA results summarizing the variation of all the root traits measured in the greenhouse experiment
(primary root length, root angle, root system size and root system width) between *I. hederacea* and *I. purpurea*. A)
The PCA biplot shows the first two PCAs' and how individuals of *I.purpurea* (black solid circles) and *I.hederacea*
(grey solid triangles) vary in this trait space. A) 37.3% and 29.1% of the phenotypic variation is explained by PCA1
and PCA2, respectively, and the overlapping ellipses representing the 95 % confidence intervals of each species
indicates high phenotypic similarity between *I.purpurea* and *I.hederacea*. Bar graphs (B & C) show the loading scores
that each root trait contributes to PCA1 and PCA2, respectively. B) Root system width,root angle and primary root
length load strongest in PCA1,C) whereas root angle loads strongest in PCA 2. PCA 1 can be used as an indicator of
root system architecture and PCA 2 can serve as an indicator of root size.

974
975

976
977
978 **Table B1** Genetic correlation matrix between root traits for *I. purpurea* (above diagonal) and for *I. hederacea*
979 (below diagonal). Pearson correlations were calculated on the family means of non-transformed root traits for each
980 species separately, and across treatments. Significant correlations (p-value < 0.05) are indicated in bold.
981

	Root angle	Root width	Root size
Root angle	—	-0.01	0.1
Root system width	**0.59**	—	**0.85**
Root system size	**0.60**	**0.80**	—

989 **Table B2:** Influence of interspecific competition within *I. purpurea* and interspecific and intraspecific competition
990 in *I. hederacea*, on the pattern of selection acting on specific root traits in these species. F-statistics from the
991 ANCOVA testing the effects of competition, block, and root trait X competition on selection gradients are shown.
992 Values in bold indicate significant treatment by trait, block by trait, and treatment by block by trait interactions
993 whereas effects that are marginally significant (p-value < 0.09) shown with an (^)).

	I. purpurea		*I. hederacea* (alone-inter)		*I. hederacea* (alone-intra)	
Root system width (cm)	F	P	F	P	F	P
Competition	2.76	0.10	0.82	0.37	1.35	0.25
Block	0.57	0.25	0.35	0.79	0.52	0.67
Competition X Block	**7.30**	**<0.01**	0.16	0.92	0.38	0.76
Root system size (cm^2)	F	P	F	P	F	P
Competition	**6.01**	**0.02**	0.58	0.45	0.87	0.35
Block	1.76	0.72	0.29	0.84	0.12	0.95
Competition X Block	1.24	0.40	0.12	0.95	0.23	0.87
Root angle	F	P	F	P	F	P
Competition	1.06	0.31	**4.47**	**0.04**	0.01	0.90
Block	0.74	0.62	1.79	0.16	**3.78**	0.01
Competition X Block	3.02^	0.08	2.37^	0.08	0.02	1.00