# Sex-biased gene expression is repeatedly masculinized in asexual females

Darren J. Parker[1,2,*], Jens Bast[1], Kirsten Jalvingh[1], Zoé Dumas[1], Marc Robinson-Rechavi[1,2], and Tanja Schwander[1]

1. Department of Ecology and Evolution, University of Lausanne, Lausanne, Switzerland,

2. Swiss Institute of Bioinformatics, Lausanne, Switzerland

*  Corresponding author: DarrenJames.Parker@unil.ch

## Abstract

Males and females feature strikingly different phenotypes, despite sharing most of their genome. The resolution of this apparent paradox is through differential gene expression, whereby genes are expressed at different levels in each sex. This resolution, however, is likely to be incomplete, leading to conflict between males and females over the optimal expression of genes. We test the hypothesis that gene expression in females is constrained from evolving to its optimum level due to sexually antagonistic selection on males, by examining changes sex-biased gene expression in five obligate asexual species of stick insect, which do not produce males. We predicted that the transcriptome of asexual females would be feminized as asexual females do not experience any sexual conflict. Contrary to our prediction we find that asexual females feature masculinized gene expression, and hypothesise that this is due to shifts in female optimal gene expression levels following the suppression of sex.

## Introduction

Genetic constraints between developmental stages, sexes, and castes arise as a result of a shared genome[1]. Species are able to mitigate these constraints by differentially expressing suites of genes in specific contexts to produce and maintain different phenotypes. This resolution, however, may be incomplete when regulatory control of gene expression is not sufficiently labile as to allow for optimal expression in each phenotypic context, leading to intralocus conflict[2].

This phenomenon has been most widely studied between the sexes, where strong sexual dimorphism is generally underlain by sex-biased gene expression[3]. Sex-biased gene expression is thought to have been largely driven by selection to resolve intralocus sexual conflict[4]. As such, contemporary sex-biased gene expression is expected to represent a combination of both resolved and partially un-resolved sexual conflict[5,6]. In the latter case, suboptimal gene expression levels are maintained by opposing selection in males and females, with the relative strengths of selection acting on each sex determining the difference between optimal and observed (suboptimal) expression levels.

Sexually antagonistic selection has the potential to constrain the optimal expression of large portions of a species' transcriptome and to thereby generate sub-optimal phenotypes in each sex. However, whether un-resolved conflict is pervasive can be difficult to investigate in natural populations, due to the relatively small (but numerous) effects of individual loci[5,7]. An ideal situation to address this question would be to examine how the transcriptome evolves following the cessation of sexual conflict. This is the case in asexually reproducing species when derived from a sexual ancestor. Because asexual species consist only of females, there is no sexual conflict and selection can optimize the female phenotype independently of any correlated effects in males. Despite the potential of this approach, previous studies have only used sexual species, examining how the transcriptome changes under experimentally altered levels of sexual selection[8–11].

The premise of these studies is that because sexual selection is typically stronger on males than females[12], a reduction in sexual selection (e.g. by enforcing monogamy) will disproportionately affect males, resulting in a shift in gene expression towards the female optimum. While this optimum is unknown, it is assumed that female-biased genes are generally beneficial for females, and male-biased genes for males[4,13], such that shifts towards female optima would generate a feminization of gene expression (increased female-biased and decreased male-biased expression). The empirical support for this hypothesis, however, remains mixed[8–11], and the most recent study[9] further showed that shifts in sex-biased gene expression under altered sexual selection varied among tissues and conditions. However, a potential constraint in these studies is that even under reduced sexual selection, selection still acts on male phenotypes as fertile males still need to be produced in each generation. Thus, many genes potentially subject to sexually antagonistic selection therefore remain unaffected by reduced sexual selection, with genes negatively affecting male viability or fertility being obvious

69     examples. This constraint does not apply to recently derived asexual species as *all* aspects of
70     sexual conflict present in the sexual species are absent in the asexual species.
71
Here we use *Timema* stick insects to examine how sex-biased genes change in expression
following a transition to asexuality. *Timema* comprise multiple independent transitions to
asexuality[14], allowing us to examine how idiosyncratic any shifts in sex-biased gene expession
are. Furthermore, male *Timema* have a single X and no Y chromosome (XX / X0 sex
determination)[15], avoiding any potential difficulties arising from sex-limited regions of the
genome. We first identify genes with sex-biased expression in five sexual *Timema* species by
sequencing the transcriptomes of three different tissue types in each sex. We then study the
fate of these sex-biased genes in close asexual relatives of each sexual species to test whether
their expression is consistently feminized. This allows us to test for the first time the importance
of intralocus sexual conflict on gene expression changes following a loss of sex.
82
83

## Results

### Gene expression is repeatedly masculinized in asexual females

To examine changes in sex-biased gene expression in asexual females, we first identified orthologous genes in each of the five sexual-asexual sister species pairs using reciprocal best Blast hits. We then classified genes as being sex-biased by comparing male and female expression in each sexual species (FDR < 0.05, absolute fold-change > 2). Sex-biased genes were identified separately for each sexual species and each of the three tissue types (whole bodies, reproductive tract and leg tissue; see Methods). As expected, given their different roles and morphology in males and females, reproductive tracts featured large numbers of sex-biased genes in all species (2843-3845, corresponding to approximately 30% of each transcriptome; Supplemental Table 1). Legs and whole bodies had fewer genes with sex-biased gene expression overall, but with considerable variation among species (0.5-12%; Supplemental Table 1). Variation among species could be due to variation in sexually dimorphic physiology between species but is also likely driven (at least partially) by differences in between sample variance (Supplemental Table 2). Note that because sex-biased genes were identified separately for each sexual species this approach cannot be used to determine if sex-biased genes are the same across species. We therefore investigated if sex-biased genes are the same between species as a second step (see below). Although the genes may be different, we can examine if sex-biased genes in different species are involved in similar functions by comparing the GO terms of sex-biased genes across species. Sex-biased genes in each species and tissue were significantly enriched for many functions (136-445 significant GO terms for male-biased genes, 138-726 for female-biased genes; Supplemental Table 3). However, only a few of them overlapped between species (Supplemental Fig. 1) (though the overlap was greater than expected by chance (FDR < 0.05, Supplemental Table 4)).

We then examined whether sex-biased genes change in expression between sexual and asexual females. Surprisingly, we found that the transcriptomes of asexual females were strongly masculinized. The expression of female-biased genes was significantly reduced in all five independently evolved asexual species and in each tissue (14 out of 15 instances, the exception being the whole-body comparison between *T. podura* and *T. genevievae*, which shows reduced female-biased gene expression, but not significantly (FDR = 0.076), Fig. 1, Supplemental Table 5). By contrast, male-biased genes significantly increased in expression in most tissues of asexual females (10 out of 15 instances), although they also significantly decreased in two instances (in *T. shepardi* reproductive tracts and *T. tahoe* legs) (Fig. 1, Supplemental Table 5).

In addition to sex-biased genes, one class of interesting genes is sex-limited genes (genes expressed in only one of the two sexes). The expression of sex-limited genes depends on different genetic architectures in males and females. Sex-limited genes are therefore expected to be free from sexual conflict over expression levels and thus may be expected to show different shifts in expression in asexual females than sex-biased genes. In particular, as female-limited genes are free from sexual conflict we expect that there will be no overall change in

expression between sexual and asexual females, if relaxation of sexual conflict is the main
driver of changes in asexual females. Note that sex-limited genes were identified separately
from sex-biased genes to avoid inflating the dispersion of the model used to identify sex-biased
genes (see Methods). Overall, we find only a few sex-limited genes (0-50), with most of these in
the reproductive tracts (Supplemental Tables 6 and 7). Like female-biased genes, female-
limited genes also show a significant reduction in expression in asexual females in most cases
(8 out of the 9 instances with more than one female-limited gene) (Fig. 2, Supplemental Fig. 2,
Supplemental Table 6). Almost all male-limited genes show very little to no expression in
asexual females, and are expressed at much lower levels than found in males (Fig. 2,
Supplemental Fig. 2, Supplemental Table 6), suggesting that few, if any, male-limited genes
have been co-opted for new functions in asexual females.
139
Our analyses show that gene expression in asexual females is generally masculinized. This
effect is particularly clear for female-biased genes which decrease in expression across five
different species and three different tissue types, showing the masculinization of sex-biased
gene expression in asexuals is very repeatable. Given this unexpected finding, we verified that
our results were not biased by the gene sets we chose to use, which excluded genes with very
low expression in asexual females, and genes without an ortholog between sexual and asexual
sister species (see Methods). Exclusion of these genes could bias our results if shifts in gene
expression disproportionately occur in these genes. To examine the impact of these factors we
firstly repeated our analyses without excluding genes with low expression in asexual females.
Generally excluded genes were were few in number (1-6%) and more likely to be male-biased
(Supplemental Tables 8-9). Repeating our analyses with these genes included found that shifts
in sex-biased gene expression in asexuals remained qualitatively the same as when they were
excluded (Supplementary Fig. 3). Secondly, we mapped reads from all samples of a sexual-
asexual species pair to a single reference (the full transcriptome of either the sexual or the
asexual species). With this strategy there is no need to identify orthologs between sexual and
asexual species pairs. Repeating our analyses using the full sexual or asexual transcriptome,
we found very few sex-biased genes had no expression in asexual females (Supplemental
Tables 10-13), and we obtained qualitatively similar results as in the main analysis
(Supplementary Figs 4-5). Taken together these analyses show that the masculinization of gene
expression we observe is not biased by our gene set selection.
160
161  ***Masculinization of sex-biased gene expression does not depend on gene identity***
162
In the above analyses, each species-pair was treated separately, which allowed us to maximize
the number of genes used in comparing changes in sex-biased gene expression between
sexual and asexual females. In doing so we use five different reference gene sets (pairwise
orthologs between sexual and asexual sister species, see Methods), which prevents us from
examining whether repeated changes to the same sex-biased genes are responsible for the
expression shifts we observe in asexual females.
169
To answer this question we firstly repeated the above analyses using only genes with 1-to-1
orthology between all ten species (between 2886 and 3003 expressed genes depending on

tissue, see Methods). Results based on this reduced gene set are qualitatively the same as
using the full gene set, i.e. an overall masculinization of sex-biased gene expression in asexual
females (Supplemental Fig. 6). As in the previous analyses the reproductive tract featured more
sex-biased genes (784-1071) than whole bodies and legs (43-375) (Supplemental Table 14).
This pattern is further illustrated by the fact that reproductive tract samples cluster first by sex
and then phylogeny, whereas it is the opposite for legs (Fig. 3). Whole-body samples show a
more mixed pattern with most samples clustering firstly by sex but with one species (*T. podura*,
which has the fewest sex-biased genes in this tissue) clustering firstly by phylogeny. Despite the
lower power of this smaller gene set (compared to the full gene set), expression of female-
biased genes was significantly reduced in asexual females in 11 out of 15 instances. Male-
biased gene expression significantly increased in asexual females in 9 out of 15 instances
(Supplemental Fig. 6, Supplemental Table 15).
184
The overlap between sex-biased genes from different species is significantly greater than
expected by chance but rather small in size (Figs. 4A and 4B, Supplemental Table 16).
Importantly for our analyses, the small overlap between species means that the consistently
masculinized gene expression we observe in asexual females is largely independent of gene
identity. This finding is strengthened by an examination of the shifts in expression for genes sex-
biased in 1, 2, 3, 4 or 5 sexual species, which show that the masculinization seen in asexual
females is stronger for genes that are sex-biased in fewer sexual species (Fig. 4C). These
findings suggest that the shifts in sex-biased gene expression we see are likely due to the
property of them being sex-biased, rather than them being involved in the same specific
biological process.
195
196  ***Functional analysis of sex-biased genes which change in expression in asexual females***
197
A plausible explanation for decreased expression of female-biased genes in asexual females is
selection against traits used for sexual reproduction. In asexual *Timema*, several sexual traits
are known to be reduced, including the production of volatile and contact pheromones[16]. Here
we observe that female-biased genes are indeed enriched for terms linked to the production of
sexual phenotypes (e.g. pheromone biosynthetic process, reproductive behavior, etc,
Supplemental Table 3), however, to more specifically identify functions that may be affected by
shifts in gene expression, we examined the GO terms specifically enriched in female-biased
genes with decreased expression in asexual females.
206
Depending on species and tissue, between 0 and 160 GO terms were significantly enriched,
with far fewer terms enriched in legs than in whole-bodies or reproductive tracts (Supplemental
Table 17), as expected given the smaller number of sex-biased genes in legs. There are no
consistently enriched GO terms between all species (Supplemental Fig. 7A), and although some
terms can be easily associated with reduction of sexual traits (e.g. olfactory behavior,
chemosensory behavior, detection of stimulus involved in sensory perception), the majority of
terms have no clear link to sexual trait reduction. Most enriched terms instead are related to
metabolic and developmental processes. This could potentially be a signature of a shift in
energy budget in asexual females which no longer have to produce costly sexual traits

(Supplemental Table 17). Male-biased genes that increased in expression were enriched for
between 0 and 81 terms, and again, no terms were shared between all species, and very few
between any pair of species (Supplemental Table 18, Supplemental Fig. 7B).
219
The removal of sexual conflict is expected to cause the feminization of gene expression in
asexual females. Although overall the pattern of expression change we observe is opposite to
this prediction, it is possible that a feminization of gene expression still occurs for a small subset
of genes, but its effect is masked by the larger effect of masculinization. We specifically
examine the subset of sex-biased genes that follow the expected pattern of feminization, by
looking at processes enriched for female-biased genes that increase in expression and male-
biased genes that decrease in expression in asexual females. We would expect that genes
showing feminization would be enriched for processes associated with sexual conflict. Both
male- and female-biased genes showed an enrichment of many terms (between 0 and 360, and
between 1 and 195, respectively), including some that could be associated with sexual conflict
(e.g. sexual reproduction, female mating behavior, etc). However, the majority of terms have no
clear link to sexual conflict, and again no terms were shared between all species (Supplemental
Table 19, Supplemental Table 20, Supplemental Fig. 8).
233
Taken together, the functional enrichment analyses suggest that the changes in gene
expression we observe are involved in a diverse set of processes in each of the species. This is
in line with what we observe from the gene expression analyses, which show that sex-biased
genes have little overlap between species, and that the largest shifts in gene expression are in
genes that are sex-biased in the fewest species. In addition to being different between species,
the enriched GO terms were not particularly informative for determining if they are involved in
sexual trait decay or sexual antagonism. This reflects the relative difficulty in obtaining functional
annotations in *Timema*, due to their evolutionary distance from a well characterised insect
model system, meaning that most functions are broad and difficult to attribute to specific roles in
sexual traits or sexual antagonism.
244
245  ***No disproportionate decay of sex-biased genes in asexuals***
246
Sex-biased genes in sexual species often evolve rapidly, due to strong sexual selection and/or
sexual antagonism which drives positive selection for amino-acid changes[17]. In asexual species,
sex-biased genes are also expected to evolve rapidly, but due to reduced purifying selection on
redundant sexual traits underlain by sex-biased genes. Although interesting, identifying
differences in evolutionary rates between gene classes in asexual species is difficult due to the
overall elevated rates in asexual species (including in *Timema*[18]), and because genes are
inherited as a single linkage group which reduces the power to detect differences in evolutionary
rate between genes. Here we found evidence for an elevated rate of dN/dS in asexual species
and in sex-biased genes (Supplemental Fig. 9, Supplemental Tables 21-23). We do not see any
evidence for an interaction between sex-bias and reproductive mode (Supplemental Fig. 9,
Supplemental Tables 21-23), indicating that the increase in dN/dS for sex-biased genes is
similar in sexual and asexual species.
259

## Discussion

Conflict over gene expression levels between males and females is thought to drive the evolution of sex-biased gene expression[4]. While sex-biased expression is expected to reduce the amount of intralocus sexual conflict, it is unlikely to be complete for many genes, meaning that some proportion of sex-biased genes are likely subject to sexually antagonistic selection[5,6]. Here we chose to investigate how sex-biased gene expression changes in asexual species which experience no sexual conflict. We predicted that transcriptomes of asexual females would be feminized as sex-biased genes in asexual females would no longer be constrained by countervailing selection pressures in males. Contrary to our prediction we found evidence for an overall masculinization of sex-biased gene expression in asexual females. This pattern of masculinization was very consistent across each of the five independently derived asexual species, and three tissue types we examined. In addition, masculinization was not driven by changes in expression of the same genes in each species, showing that it is the property of being sex-biased *per se* that is most likely to be responsible for the shifts in expression we observe.

Taken together, our results provide strong evidence for a masculinization of gene expression in asexual species. The strength of this finding does not mean there is no sexual conflict over optimal levels in sexual species, but rather that changes in asexual females driven by a release of conflict are negligible relative to changes driven by other mechanisms. The presence of such alternate mechanisms can best be illustrated by the fact that female-limited genes (that should experience no sexual conflict over gene expression level in sexual species), show a consistent masculinization similar to sex-biased genes. We suggest that this is because although reproducing asexually does remove the pressure of sexual conflict, it also removes the need for many of the sexual traits sexually dimorphic gene expression underlies. Consequently, while we expected gene expression in asexual females to be free to move to a female optimum, it is also likely that the optimal female phenotype is different for sexual and asexual females.

Female asexual *Timema* show reductions in several sexual traits including a reduced sperm storage organ, and reduced volatile and contact pheromone production[16]. Since sexually dimorphic traits are largely a product of sex-biased gene expression[3], a link between reduced female sexual traits and reduced female-biased gene expression is a plausible explanation for the decreased expression of female-biased genes we observe. It is less clear why we also see an accompanying increase of expression in male-biased genes in asexual females. We suggest three, non-mutually exclusive, speculative explanations for this. Firstly, increased expression of male-biased genes may arise as a result of sexual trait reduction in cases where high expression of a gene in males acts to suppress the development of a trait, or when low expression in females acts to enhance a female sexual trait. In such genes selection for sexual trait reduction in asexual females would be expected to produce an increase in expression. A second potential explanation is that in sexual species there are a number of products produced by males and then transferred to females that are important for female fertility. For instance, in many insects, ovulation and oviposition are stimulated by substances in the male ejaculate such as juvenile hormone, prostaglandins, and myotropins[19,20]. Since these products are not provided

by males in asexual species, females may need to increase expression of the genes that
produce these products to compensate. While this explanation could explain some of the
increased expression of male-biased genes we observe in the reproductive tract, it is unlikely to
be a general explanation for the increased expression of male-biased genes across all species
and tissues. Finally, a third potential explanation is that if males and females in sexual species
have separate niches, a transition to asexuality would allow asexual females to expand into the
male niche. Differential niche use is likely to, at least in part, be mediated by sex-biased gene
expression, meaning that asexual females would need to masculinize their gene expression in
order to occupy the vacant niche left by males. While differences in male and female niche use
have not been extensively studied in *Timema*, sexual dimorphism in mandible shape has been
reported, implying that there may be some differential use of niche-space in sexual *Timema*
species[21]. Future work examining sexual niche usage and gene expression is needed to
evaluate this hypothesis.
317
For the reasons detailed above, we believe that female trait reduction is the most likely
explanation for the majority of changes in sex-biased gene expression we observe, rather than
the cessation of sexual conflict. Similar to our findings, a recent study[9] found that experimentally
reduced sexual selection also produced an overall masculinization of gene expression in *D.*
*pseudoobscura*. However, Veltsos et al.[9] interpret their findings as a consequence of reduced
conflict, and attribute masculinization to the dynamic nature of sexually antagonistic selection
causing unpredictable changes in sex-biased gene expression. An alternative explanation,
however, is that the masculinization of gene expression in females seen in Veltsos et al.
corresponds to a reduction of sexual traits under reduced sexual selection, similar to our
findings in *Timema*. Previous studies have reported reduced sexual traits in females evolving
under reduced sexual selection in the *D. pseudoobscura* lines studied by Veltsos et al.[11,22]. As
such, both Veltsos et al. and our results highlight that the shifts in sex-biased gene expression
we observe in the absence, or under reduced levels, of sexual conflict may be in part due to a
shift in the optimal trait levels in females. Such shifts in female optima under different sexual
selection scenarios are important to consider as an explanation even for studies that observe
the expected feminization of sex-biased gene expression[8,10,11]. This is because reducing sexual
selection can also favour the increased expression of female sexual traits under some
conditions. In these situations, the feminization of sex-biased gene expression can be due to
changes in sexual trait optima rather than due to a reduction in the amount of intralocus sexual
conflict. Changes to optimal trait values under different selective scenarios are difficult to predict
*a priori*[23], meaning future studies will require careful examination of optimal phenotypes under
different selective scenarios in order to correctly interpret any changes in sex-biased gene
expression.
341
In conclusion, we find that sex-biased gene expression is repeatedly masculinized following a
transition to asexuality, and suggest that this result is driven primarily by a reduction of female
sexual traits. While we observe similar patterns of masculinization across all five asexual
species, the genes involved were mostly different, reflecting the dynamic nature of sex-biased
gene expression. In line with this, the functional processes associated with expression change
in each species were also diverse. Finally, our study highlights the importance of considering

explanations other than intralocus sexual conflict for explaining shifts in sex-biased gene
expression, since differences in sexual conflict are also likely to be accompanied by changes in
sexual trait optima which may enhance or mask changes caused by a reduction or cessation of
intralocus sexual conflict.

## Methods

Individuals for whole-body and tissue-specific samples were collected from the field as last instar juveniles in spring 2013 and 2014, respectively (collection locations for all samples are given in Supplemental Table 24). All individuals were raised in common garden conditions (23°C, 12h:12h, 60% humidity, fed with Ceanothus cuttings) until eight days following their final moult. Prior to RNA extraction, individuals were fed with an artificial medium for two days to avoid RNA contamination with gut content and then frozen at -80°C. For leg samples, three legs were used from each individual (one foreleg, one midleg, and one hindleg). Reproductive tracts were dissected to consist of ovaries, oviducts and spermatheca in females and testes and accessory glands in males. Note the same individuals were used for leg and reproductive tract samples. To ensure individuals were reproductively active at the time of sampling, all sexual individuals were allowed to mate, and asexual and sexual females were observed to lay eggs. When analyses were repeated using virgin sexual females, we obtained qualitatively similar results (see Supplemental Material and Supplemental Fig. 10).

### *RNA extraction and sequencing*

We generated three biological replicates per species and tissue from pooled individuals (1-9 individuals per replicate, a total of 516 individuals, in 150 replicates in total (including the virgin sexual females); see Supplemental Table 24). To extract RNA, samples were flash-frozen in liquid nitrogen followed by addition of Trizol (Life Technologies) before being homogenized using mechanical beads (Sigmund Lindner). Chloroform and ethanol were then added to the samples and the aqueous layer transferred to RNeasy MinElute Columns (Qiagen). RNA extraction was then completed using an RNeasy Mini Kit following the manufacturer's instructions. RNA quantity and quality was measured using NanoDrop (Thermo Scientific) and Bioanalyzer (Agilent). Strand-specific library preparation and single-end sequencing (100 bp, HiSeq2000) were performed at the Lausanne Genomic Technologies Facility.

The 150 libraries produced a total of just under 5 billion single-end reads. 6 whole-body and 6 tissue-specific libraries produced significantly more reads than the average for the other samples. To reduce any influence of this on downstream analyses, these libraries were sampled down to approximately the average number of reads for whole-body or tissue-specific libraries respectively using seqtk (https://github.com/lh3/seqtk Version: 1.2-r94).

### *Transcriptome references*

*De novo* reference transcriptome assemblies for each species were generated previously[18]. Our expression analyses were conducted using two sets of orthologs. Firstly, we identified orthologs between sexual and asexual sister species using reciprocal Blast as described in Parker et al.[24]. Secondly, we used the 3010 one-to-one orthologs present in all 10 *Timema* species as identified by Bast et al.[18]. The identified ortholog sequences varied in length among different species. Since length variation might influence estimates of gene expression, we aligned orthologous sequences using PRANK (v.100802, default options)[25] and trimmed them using

alignment_trimmer.py[26] to remove overhanging gaps at the ends of the alignments. If an
alignment contained a gap of greater than 3 bases then sequence preceding or following the
alignment gap (whichever was shortest) was discarded. Any orthologous sequences that had a
trimmed length of less than 300 bp were also discarded. Finally, before mapping, genes with
significant Blast hits to rRNA sequences were removed from the trimmed transcriptome
references.
402
403 ***Read trimming and mapping***
404
Before mapping, adapter sequences were trimmed from raw reads with CutAdapt[27]. Reads were
then quality trimmed using Trimmomatic v 0.36[28], clipping leading or trailing bases with a phred
score of <10 from the read, before using a sliding window from the 5' end to clip the read if 4
consecutive bases had an average phred score of <20. Any reads with a sequence length of
<80 after trimming were discarded. Reads from each libret were then mapped to the
transcriptome references using Kallisto (v. 0.43.1)[29] with the following options -l 210 -s 25 --bias
--rf-stranded for whole-body samples and -l 370 -s 25 --bias --rf-stranded for tissue-specific
samples (the -l option was different for whole-body and tissue-specific samples as the fragment
length for these libraries was different).
414
415 ***Differential expression analysis***
416
Expression analyses were performed using the Bioconductor package EdgeR (v. 3.18.1)[30] in R
(v. 3.4.1)[31]. Firstly, to identify sex-biased genes we compared male and female expression
separately for each tissue in each sexual species. Genes with counts per million less than 0.5 in
2 or more libraries per sex were excluded from expression analyses. Normalization factors for
each library were computed using the TMM method. To estimate dispersion we then fit a
generalized linear model (GLM) with a negative binomial distribution with sex as an explanatory
variable and used a GLM likelihood ratio test to determine the significance of sex on gene
expression for each gene. P-values were then corrected for multiple tests using Benjamini and
Hochberg's algorithm[32]. Sex-biased genes were then defined as genes that showed a greater
than 2 fold difference in expression between males and females with an FDR < 0.05. We chose
this threshold in order to select a robust set of sex-biased genes, and to reduce the effect of
sex-biased allometry[33]. Note that analyses using just an FDR threshold to define sex-biased
genes produced qualitatively similar results (Supplemental Tables 25-26).
430
431 Clustering of expression values ($\log_2$ CPM) was performed using Ward's hierarchical clustering
432 of Euclidean distances with the R package pvclust (v. 2.0.0)[34], with bootstrap resampling
433 (method.hclust="ward.D2", method.dist="euclidean", nboot=10000), and visualized using R
434 package pheatmap (v. 1.0.8)[35].
435
436 To quantify how sex-biased genes change in expression in asexual females we then compared
437 gene expression in sexual and asexual females separately for each species pair and each
438 tissue. We also compared the change in expression in asexual females for male- and female-

439    biased genes to unbiased genes using a Wilcoxon test, corrected for multiple tests using
440    Benjamini and Hochberg's algorithm[32].
441
442    ***Analysis of sex-limited genes***
443
Sex-limited genes were classified as genes that had at least 2 FKPM (Fragments Per Kilobase
Million) in each replicate of one sex and 0 FKPM in each replicate of the other sex. FKPM
values were calculated using EdgeR. The expression levels of female-limited genes in sexual
and asexual females, and male-limited genes in sexual males and asexual females was
compared using a Wilcoxon test, corrected for multiple tests using Benjamini and Hochberg's
algorithm[32].
450
451    ***Sequence evolution of sex-biased genes***
452
To test if sex-biased genes experience reduced purifying selection in asexuals, we examined if
sex-biased genes have elevated dN/dS ratios in asexuals. To do this we firstly fit a binomial
glmm (dN/dS values were transformed to fall into two categories: zero or non-zero), with
reproductive mode, sex-bias and their interaction as fixed effects and gene identity as a random
effect. Secondly, we firstly fit a glmm with a gamma distribution to the dN/dS values that were
greater than zero, with the same fixed and random effects as the binomial model. All glmms
were fit using the lme4 package (v. 1.1.14)[36] in R, and significance of terms was determined
using a log-likelihood ratio test. dN/dS values were calculated for each of the one-to-one
orthologs using codeml implemented in the PAML package[37] to generate maximum likelihood
estimates of dN/dS for each terminal branch in the phylogeny (using the "free model") as
described in Bast et al.[18].
464
465    ***GO term analysis***
466
Genes were functionally annotated using Blast2GO (version 4.1.9)[38] as described in Parker et
al.[24]. Briefly, sequences from each sexual species were compared with BlastX to either NCBI's
nr-arthropod or *Drosophila melanogaster* (drosoph) databases, to produce two sets of functional
annotations, one derived from all arthropods and one specifically from *Drosophila melanogaster*.
The *D. melanogaster* GO term annotation generated around four times more annotations per
sequence than NCBI's nr-arthropod database. We therefore conducted all subsequent analyses
using the GO terms derived from *D. melanogaster*, but note that results using the annotations
from all arthropods were qualitatively the same (see Supplemental Fig. 11).
475
476    To identify overrepresented GO terms we conducted gene set enrichment analyses (GSEA)
477    using the R package TopGO (v. 2.28.0)[39], using the elim algorithm to account for the GO
478    topology. GO terms were considered to be significantly enriched when $p < 0.05$. Repeating our
479    GO term analyses using the more liberal 'weight01' algorithm produced qualitatively the same
480    results (results not shown).
481

Since we defined sex-biased genes with both FDR and FC thresholds, we ranked sex-biased
genes for the GSEA to take both FDR and FC into account. To identify overrepresented GO
terms for female-biased genes, genes were ranked by FDR in four subsets: female-biased with
FC > 2, female-biased with FC < 2, male-biased with FC < 2, and male-biased with FC > 2.
Female-biased gene subsets were ranked so that small FDR values were ranked highly, male-
biased gene subsets were ranked so that small FDR values were ranked low in the list. The four
lists were then joined together in the order given above, and assigned a unique rank. This
ranked list produces a list where strongly female-biased genes with are at the top, followed by
weakly female-biased genes, then weakly male-biased genes, and finally strongly male-biased
genes at the bottom. To identify overrepresented GO terms for male-biased genes the ranked
list for female-biased genes was simply inverted. Finally, to examine the GO terms
overrepresented in sex-biased genes which changed expression in asexuals, female- and male-
biased genes were ranked by fold-change between sexual and asexual females.
495
496 To determine if the overlap of sets of sex-biased genes or GO terms was greater than expected
497 by chance we used the SuperExactTest package (v. 0.99.4; [40]) in R which calculates the
498 probability of multi-set intersections. P-values for were multiple test corrected using Benjamini
499 and Hochberg's algorithm implemented in R.
500

## Data
502 Raw reads have been deposited in SRA under accession codes SRR5748941-SRR5749000,
503 for whole-body samples and SRR5786827-SRR5786961 for tissue-specific samples. Scripts for
504 the analyses in this paper are available at
505 https://github.com/DarrenJParker/Timema_Sex_Biased_Gene_Exp, and will be archived at
506 Zenodo after acceptance.
507

## Acknowledgments
509 This study was supported by Swiss FNS grants PP00P3_170627, PP00P3_139013, and
510 CRSII3_160723. We would like to thank Chloe Larose and Bart Zijlstra for their
511 assistance in the field.

**Figures**

**Figure 1** | **Expression shifts in sex-biased genes in asexual females.** Positive values on the
y-axis indicate increased expression in asexual females. Asterisks indicate the significance level
(FDR) of Wilcoxon tests comparing the change in expression in female-biased (red) and male-
biased (blue) genes to unbiased genes (*** < 0.001, ** < 0.01, * < 0.05). Species names are
abbreviated as follows: Tte = *T. tahoe*, Tms = *T. monikensis*, Tdi = *T. douglasi*, Tsi = *T.
shepardi*, and Tge = *T. genevievae*.

**Figure 2 | Expression of sex-limited genes in the reproductive tract.** A) Expression of
female-limited genes in sexual females (SF, red) and asexual females (AF, purple), B)

Expression of male-limited genes in sexual females (SF, red) and asexual females (AF, purple),

C) Expression of male-limited genes in sexual males (SM, blue) and asexual females (AF,

purple). Asterisks indicate the significance level (FDR) of Wilcoxon tests (*** < 0.001, ** < 0.01,

* < 0.05). Species names are given as abbreviations in the form sexual-species - asexual

species at the left-hand side (Tbi = *T. bartmani*, Tce = *T. cristinae*, Tps = *T. poppensis*, Tcm = *T.*

*californicum*, Tpa = *T. podura*, Tte = *T. tahoe*, Tms = *T. monikensis*, Tdi = *T. douglasi*, Tsi = *T.*

*shepardi*, and Tge = *T. genevievae*). Note this figure depicts only the results from the

reproductive tract. For whole-bodies see Supplemental Figure 2. Legs were not plotted due to

the small number od sex-limited genes in this tissue (Supplemental Tables 6-7).

**Figure 3** | Heatmaps and hierarchical clustering of gene expression (log$_2$ CPM) for whole-body, reproductive tract, and leg samples. Values on each node show the bootstrap support from 10,000 replicates. Species names are abbreviated as follows: Tbi = *T. bartmani*, Tce = *T. cristinae*, Tps = *T. poppensis*, Tcm = *T. californicum*, Tpa = *T. podura*

539 **Figure 4** | A) Venn-diagrams showing the overlap of female-biased genes B) Venn-diagrams
540 showing the overlap of male-biased genes C) Boxplots showing the change in expression of
541 female-biased (reds) and male-biased genes (blues) in asexual females when a genes is female
542 or male-biased in 1, 2, 3, 4 or 5 sexual species.

**References**

1.   Lande, R. Sexual dimorphism, sexual selection, and adaptation in polygenic characters.

*Evolution* **34**, 292–305 (1980).

2.   Rice, W. R. & Chippindale, A. K. Intersexual ontogenetic conflict. *J. Evol. Biol.* **14**, 685–693

(2001).

3.   Mank, J. E. The transcriptional architecture of phenotypic dimorphism. *Nature Ecology &*

*Evolution* **1**, 0006 (2017).

4.   Connallon, T. & Knowles, L. L. Intergenomic conflict revealed by patterns of sex-biased

gene expression. *Trends Genet.* **21**, 495–499 (2005).

5.   Innocenti, P. & Morrow, E. H. The sexually antagonistic genes of *Drosophila melanogaster*.

*PLoS Biol.* **8**, e1000335 (2010).

6.   Griffin, R. M., Dean, R., Grace, J. L., Rydén, P. & Friberg, U. The shared genome is a

pervasive constraint on the evolution of sex-biased gene expression. *Mol. Biol. Evol.* **30**,

2168–2176 (2013).

7.   Cheng, C. & Kirkpatrick, M. Sex-specific selection and sex-biased gene expression in

humans and flies. *PLoS Genet.* **12**, e1006170 (2016).

8.   Hollis, B., Houle, D., Yan, Z., Kawecki, T. J. & Keller, L. Evolution under monogamy

feminizes gene expression in *Drosophila melanogaster*. *Nat. Commun.* **5**, 3482 (2014).

9.   Veltsos, P., Fang, Y., Cossins, A. R., Snook, R. R. & Ritchie, M. G. Mating system

manipulation and the evolution of sex-biased gene expression in *Drosophila*. *Nat. Commun.*

**8**, 2072 (2017).

10.  Moghadam, H. K., Pointer, M. A., Wright, A. E., Berlin, S. & Mank, J. E. W chromosome

expression responds to female-specific selection. *Proc. Natl. Acad. Sci. U. S. A.* **109**, 8207–

8211 (2012).

11.  Immonen, E., Snook, R. R. & Ritchie, M. G. Mating system variation drives rapid evolution

of the female transcriptome in *Drosophila pseudoobscura*. *Ecol. Evol.* **4**, 2186–2201 (2014).

12. Janicke, T., Häderer, I. K., Lajeunesse, M. J. & Anthes, N. Darwinian sex roles confirmed

across the animal kingdom. *Sci Adv* **2**, e1500983 (2016).

13. Dean, R., Hammer, C., Higham, V. & Dowling, D. K. Masculinization of gene expression is

associated with male quality in *Drosophila melanogaster*. *Evolution* (2018).

doi:10.1111/evo.13618

14. Schwander, T., Henry, L. & Crespi, B. J. Molecular evidence for ancient asexuality in

*Timema* stick insects. *Curr. Biol.* **21**, 1129–1134 (2011).

15. Schwander, T. & Crespi, B. J. Multiple direct transitions from sexual reproduction to

apomictic parthenogenesis in *Timema* stick insects. *Evolution* **63**, 84–103 (2009).

16. Schwander, T., Crespi, B. J., Gries, R. & Gries, G. Neutral and selection-driven decay of

sexual traits in asexual stick insects. *Proc. Biol. Sci.* **280**, 20130823 (2013).

17. Ellegren, H. & Parsch, J. The evolution of sex-biased genes and sex-biased gene

expression. *Nat. Rev. Genet.* **8**, 689–698 (2007).

18. Bast, J. *et al.* Consequences of asexuality in natural populations: insights from stick insects.

*Mol. Biol. Evol.* **35**, 1668–1677 (2018).

19. Arnqvist, G. & Rowe, L. *Sexual conflict*. (Princeton University Press, 2005).

20. Gillott, C. Male accessory gland secretions: modulators of female reproductive physiology

and behavior. *Annu. Rev. Entomol.* **48**, 163–184 (2003).

21. Roy, D., Seehausen, O. & Nosil, P. Sexual dimorphism dominates divergent host plant use

in stick insect trophic morphology. *BMC Evol. Biol.* **13**, 135 (2013).

22. Hunt, J., Snook, R. R., Mitchell, C., Crudgington, H. S. & Moore, A. J. Sexual selection and

experimental evolution of chemical signals in *Drosophila pseudoobscura*. *J. Evol. Biol.* **25**,

2232–2241 (2012).

23. Dale, J., Dey, C. J., Delhey, K., Kempenaers, B. & Valcu, M. The effects of life history and

sexual selection on male and female plumage colouration. *Nature* **527**, 367–370 (2015).

24. Parker, D. J. *et al.* Repeated evolution of asexuality involves convergent gene expression

changes. *Mol. Biol. Evol.* **36** , 350–364 (2019).

25. Löytynoja, A. & Goldman, N. An algorithm for progressive multiple alignment of sequences

with insertions. *Proc. Natl. Acad. Sci. U. S. A.* **102**, 10557–10562 (2005).

26. Parker, D. J. fasta_tools v1.2. *Zenodo. https://zenodo.org/record/59775* (2016).

doi:10.5281/zenodo.162913

27. Martin, M. Cutadapt removes adapter sequences from high-throughput sequencing reads.

*EMBnet.journal* **17**, 10–12 (2011).

28. Bolger, A. M., Lohse, M. & Usadel, B. Trimmomatic: a flexible trimmer for Illumina

sequence data. *Bioinformatics* **30**, 2114–2120 (2014).

29. Bray, N. L., Pimentel, H., Melsted, P. & Pachter, L. Near-optimal probabilistic RNA-seq

quantification. *Nat. Biotechnol.* **34**, 525–527 (2016).

30. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140

(2010).

31. R Core Team. R: A Language and Environment for Statistical Computing. (2017).

32. Benjamini, Y. & Hochberg, Y. Controlling the false discovery rate: A practical and powerful

approach to multiple testing. *J. R. Stat. Soc. Series B Stat. Methodol.* **57**, 289–300 (1995).

33. Montgomery, S. H. & Mank, J. E. Inferring regulatory change from gene expression: the

confounding effects of tissue scaling. *Mol. Ecol.* **25**, 5114–5128 (2016).

34. Suzuki, R. & Shimodaira, H. Pvclust: an R package for assessing the uncertainty in

hierarchical clustering. *Bioinformatics* **22**, 1540–1542 (2006).

35. Kolde, R. Pheatmap: pretty heatmaps. *R package version* **61**, (2012).

36. Bates, D., Mächler, M., Bolker, B. & Walker, S. Fitting linear mixed-effects models using

lme4. *Journal of Statistical Software, Articles* **67**, 1–48 (2015).

37. Yang, Z. PAML 4: phylogenetic analysis by maximum likelihood. *Mol. Biol. Evol.* **24**, 1586–

1591 (2007).

38. Götz, S. *et al.* High-throughput functional annotation and data mining with the Blast2GO

suite. *Nucleic Acids Res.* **36**, 3420–3435 (2008).

39. Alexa, A., Rahnenführer, J. & Lengauer, T. Improved scoring of functional groups from

gene expression data by decorrelating GO graph structure. *Bioinformatics* **22**, 1600–1607

(2006).

40. Wang, M., Zhao, Y. & Zhang, B. Efficient test and visualization of multi-set intersections.

*Sci. Rep.* **5**, 16923 (2015).