1

High transcriptional error rates

vary as a function of gene expression level

Authors: K.M. Meer[1,2], P.G. Nelson[1], K. Xiong[3,4], J. Masel[1]*

Affiliations:

[1]Department of Ecology & Evolutionary Biology, University of Arizona, 1041 E Lowell St Tucson

AZ 85721 USA.

[2]Present address: Computational Bioscience Program, University of Colorado Anschutz Medical

Campus, 12800 E 19th Ave, Mail Stop 8303, Aurora, CO 80045 USA.

[3] Department of Molecular & Cellular Biology, University of Arizona, 1007 E Lowell St Tucson AZ

85721 USA.

[4]Present address: Computational Biology & Bioinformatics Program, Yale University, 266

Whitney Avenue, New Haven, CT 06520 USA.

*Correspondence to: masel@email.arizona.edu

18   **Abstract**

Errors in gene transcription can be costly, and organisms have evolved to prevent their

occurrence or mitigate their costs. The simplest interpretation of the drift barrier hypothesis

suggests that species with larger population sizes would have lower transcriptional error rates.

However, *Escherichia coli* seems to have a higher transcriptional error rate than species with

lower effective population sizes, e.g. *Saccharomyces cerevisiae*. This could be explained if

selection in *E. coli* were strong enough to maintain adaptations that mitigate the consequences

of transcriptional errors through robustness, on a gene by gene basis, obviating the need for

low transcriptional error rates and associated costs of global proofreading. Here we note that if

selection is powerful enough to evolve local robustness, selection should also be powerful

enough to locally reduce error rates. We therefore predict that transcriptional error rates will

be lower in highly abundant proteins on which selection is strongest. However, we only expect

this result when error rates are high enough to significantly impact fitness. As expected, we find

such a relationship between expression and transcriptional error rate for non C→U errors in *E.*

coli but not in *S. cerevisiae*. We do not find this pattern for C→U changes in *E. coli*, presumably

because most deamination events occurred during sample preparation, but do for C→U

changes in *S. cerevisiae*, supporting the interpretation that C→U error rates estimated with an

improved protocol, and which occur at rates comparable to *E. coli* non C→U errors, are

biological.

Keywords: nearly neutral theory; weak selection; Cir-Seq; RNA editing; phenotypic mutation

 Main Text

Errors are costly, and we therefore expect natural selection to reduce their rate. However,

selection cannot achieve everything. In particular, it is only able to purge deleterious mutations

when their selection coefficient s is significantly greater than one divided by the "effective

population size". This numerical limit to selection may reflect not just the number of individuals

in a population, but also competing selection at linked sites (Good and Desai 2014; Lynch 2007).

The "nearly neutral theory" holds that deleterious mutations close to this limit are abundant

(Ohta 1973), and the "drift barrier hypothesis" holds that differences in the precise location of

this limit explain important differences among species (Lynch 2007). For example, codon usage

bias is stronger in species believed to have higher effective population sizes (Vicario et al.

2007), indicating stronger selection to purge slightly deleterious synonymous mutations.

Rajon and Masel (2011) highlighted the distinction between a "global" solution that ameliorates

a problem at many loci at once, and a set of "local" solutions that solve them one at a time.

Because mutations affecting single loci are likely to have smaller fitness consequences than

mutations with genome-wide effects, the drift barrier forms a more formidable barrier to local

solutions than it does to global solutions. When local solutions evolve (in populations with large

effective population sizes), they can obviate the need for global solutions. This yields the

counterintuitive prediction that when global solutions are examined, it may be species with low

effective population sizes that show the most extreme adaptations. Specifically, rates of error in

transcription and translation could be higher in species with high effective population sizes,

since reducing error rates by kinetic proofreading is a costly global solution (Rajon and Masel

2011).

Here we focus on mistranscription errors, where during transcription, the wrong nucleic acid is

incorporated at a single site. This can lead to non-functional proteins, incurring three types of

costs. First is the energetic cost of futile transcription and translation (Wagner 2007); which can

be significant in bacteria with large population sizes (Lynch and Marinov 2015; Petrov and Hartl

2000). Second, there is the opportunity cost of not using ribosomes to make other gene

products (Dekel and Alon 2005; Kafri et al. 2016; Scott et al. 2014). Third, there is the cost of

disposing of a misfolded and potentially toxic protein (Drummond and Wilke 2009; Geiler-

Samerotte et al. 2011; Tomala and Korona 2013). Rajon and Masel (2011) predicted that in

populations with smaller effective population sizes and more loci, costly proofreading might

evolve to reduce the rate of mistranscription and hence the frequency with which these three

costs are born, while in populations with very large effective population sizes and fewer loci,

local solutions might evolve to reduce the cost of each mistranscription event, allowing their

rate to stay high.

This prediction seems to have been confirmed for mistranscription (Xiong et al. 2017), whose

rate of 8.2×10^{-5} in *Escherichia coli* (Traverse and Ochman 2016b) is far higher than that in

Saccharomyces cerevisiae (3.9×10^{-6}) (Gout et al. 2017) or *Caenorhabditis elegans* (4.1×10^{-6})

(Gout et al. 2013), which have lower effective population sizes. Indeed, the rate is higher even

than that of *Buchnera aphidicola* (4.7×10^{-5}) (Traverse and Ochman 2016b). *Buchnera* is a highly

mutationally degraded species in which the drift barrier is an obstacle to the maintenance of

fidelity in many other important cellular functions (McCutcheon and Moran 2012); this high

rate in *Buchnera* may thus indicate that the drift barrier forms an obstacle even to global

solutions (Xiong et al. 2017). All these error rates except for that of *C. elegans* (Gout et al.

2013), were estimated using Cir-Seq (Acevedo and Andino 2014), and should therefore be

comparable, although sample preparation techniques differ in vulnerability to deamination.

If the drift barrier theory of Rajon and Masel (2011) explains the high rate of mistranscription in

E. coli, this implies that selection in *E. coli* must be potent enough to be sensitive to the

consequences of transcription errors in a local (i.e. site-specific) way, not just to its global rate.

Local solutions to mistranscription fall into two categories: local robustness to the

consequences of mistranscription when it occurs (this evolved robustness is hypothesized to be

responsible for permitting globally high mistranscription rates), and locally reduced

mistranscription rates at the sites most sensitive to it.

Here we test whether selection is able to maintain locally lower transcriptional error rates in

highly expressed genes. Selection to purge deleterious mutations is generally more effective in

highly expressed genes, as evidenced, for example, by stronger codon bias (Cutter and

Charlesworth 2006; Duret and Mouchiroud 1999; Ran et al. 2014; Sharp et al. 2010), which

lowers translational error rates (Zhang et al. 2016). Somatic mutations (Frigola et al. 2017),

alternative transcriptional start sites (Xu et al. 2019), post-transcriptional modifications (Liu and

Zhang 2018a, b), alternative mRNA polyadenylation (Xu and Zhang 2018), and translation errors

(Mordret et al. 2018) also occur at lower rates at sites where they are likely to have larger

effects. We similarly predict that because high mistranscription rates matter more for highly

expressed genes, highly expressed genes should evolve a lower rate of mistranscription. We

make this prediction for *E. coli*, where mistranscription rates are globally high and thus so is

local selection pressure. In contrast, we do not expect a relationship between expression level

and mistranscription rate in *S. cerevisiae*, where mistranscription rates are globally much lower.

Mistranscription rate data in *E. coli* were taken from Traverse and Ochman (2016a), who used

Cir-Seq (Acevedo and Andino 2014) to distinguish mistranscription events from sequencing

errors. Within the largest and highest-quality batch of their data (see Methods), data from four

experimental conditions (minimal vs. rich media, and midlog vs. stationary phase) were

sometimes analyzed separately and sometimes pooled. Mistranscription rates are much higher

for C$\rightarrow$U substitutions: ~10^{-4} rather than ~10^{-5} for other mistranscription types. Since C$\rightarrow$U

changes likely represent post-transcriptional deamination events, they are probably not

mistranscription errors per se, and they are also potentially more sensitive to preparation

artifacts (Chen et al. 2014), so we excluded them from most of our analysis.

To further ensure the quality of the data, we exclude "hotspot" nucleotide sites experiencing

significantly ($p<10^{-9}$) more errors of one type than expected from our model fitted as described

below. This eliminates recent mutations, inaccurate mapping of reads to the genome, or other

artifacts of the experiment or pipeline, as well as eliminating any sites subject to programmed

post-transcriptional RNA editing. We identified 5 protein-coding and 2,390 non-coding sites

that met this "hotspot" criteria for at least one experimental condition, and excluded these

sites from our analysis. The high rate of apparent mistranscription hotspots in non-coding genes

has been interpreted (Traverse and Ochman 2016a) as a consequence of the fact that *E. coli* has

multiple polymorphic rRNA operons, making mapping of reads inaccurate. We therefore restrict

our analysis to protein-coding genes only.

We modeled the number of errors observed per nucleotide site as count data, using a

generalized linear model. The number of errors expected is the product of the number of

observations of that nucleotide site, and the modeled mistranscription rate, the latter a linear

function of log protein abundance, experimental condition, and substitution type (see Methods,

model fitting culminated in the "final model" given in Equation 1, which includes protein

abundance, and experimental condition and substitution type each having effects on intercept

but not slope). The dependence on protein abundance (Figure 1; slope of 0 rejected with p

$=2\times10^{-12}$ in final model) supports our prediction from drift barrier theory, a result that gets

slightly stronger if we omit our hotspot removal procedure. The 11 non-C→U substitution types

have substantially different mistranscription rates; fitting different intercepts for each type

(while leaving their slopes the same) is strongly supported for inclusion in our final model (p =

2×10^{-16}). Similarly, different intercepts for different experimental conditions are supported ($p =$

1.5×10^{-3}).

Figure 1. Highly expressed *E. coli* genes are subject to lower mistranscription rates. Dashed "simple model" shows a linear model fitted to a pooled dataset of 11 substitution types (i.e. excluding C→U) for all four conditions in batch #1. Dotted line shows the linear model with both condition and substitution type affecting the intercept but not the slope, plotted as a weighted average over conditions and substitutions, with weights proportional to the frequencies of opportunity to occur (i.e. by the numbers of reads of sites with A/C/G/U). Solid line shows the pooled data, binned by protein abundance as described in the Methods, and plotted according to mean protein abundance and the mean and 95% CI of the mistranscription rate within each bin. Data were divided into 10 bins; because of the limited availability of reads for low-expression genes, data within the first three bins were pooled. Note that mistranscription rate is per possible error, so the total mistranscription rate per nucleotide is around three times larger.

Figure 2. While different substitution types occur at different rates, all but C→U occur less often for highly abundant *E. coli* proteins. Dotted lines show our final linear model, in which the slope is the same across all error types. To display this model, we averaged the intercept over the four conditions, weighted according to the numbers of reads in each condition. Dashed lines show linear models with both the slope and intercept fitted separately for each error type using data pooled across all four conditions. Solid lines show the mean mistranscription rates, binned by protein abundance as described in the Methods, plotted according to mean protein abundance within each bin; error bars show 95% CI. Data were divided into 8 bins; because of the limited availability of reads for low-expression genes, data within the first three bins were pooled. Note that mistranscription rate is per possible error, so total mistranscription rate per nucleotide is around three times larger. The C→U error plot is shown inset due to very different y-axis values.

Figure 3. Different experimental conditions for *E. coli* yield similar mistranscription rates.
Dotted lines show our final model, in which the slope is constant across all conditions. To
display this model, we averaged the intercept over all error types, weighted by the frequencies
of opportunity for the error type to occur (i.e. by the numbers of reads of sites with A/C/G/U).
Dashed lines show linear models with both the slope and intercept fitted separately for each
condition using data pooled across all error types. Solid lines show the mean mistranscription
rates, binned by protein abundance as described in the Methods, plotted according to mean
protein abundance within each bin; error bars show 95% CI. Data were divided into 8 bins;
because of the limited availability of reads for low-expression genes, data within the first three
bins were pooled. Note that mistranscription rate is per possible error, so total mistranscription
rate per nucleotide is around three times larger.

Also fitting different slopes for different substitution types yields only a marginally significant

additional improvement over our final model, mostly due to the steeper slope for G→A

substitutions (Figure 2; p=0.051). Similarly, also fitting different slopes for each experimental

condition only marginally improves the fit relative to our final model (p = 0.026) with statistical

significance driven by a steeper slope in the minimal-static condition, which had far fewer data

points than the other conditions (Figure 3). The negligible differences in slope obtained from

pooling vs. stratifying (Figure 1) indicates that the slope is not spuriously generated by

systematic differences among the combined substitution types and experimental conditions.

Traverse and Ochman (2016a) reported that mistranscription errors were more commonly

synonymous (32%) than would be predicted if errors occurred at random across the genome

(24%). When controlling for the effects of substitution type, condition, and protein abundance

in our final model of mistranscription rates, the synonymous vs. non-synonymous status of the

potential mistranscription error did not predict the error rate (p = 0.08). Indeed, following our

data processing and quality filters, the overall frequency with which a mistranscription error

was synonymous was 23.4%, suggesting that the previously reported excess of synonymous

mistranscription events was due to data quality issues. In any case, whatever molecular

mechanism is responsible for variation in mistranscription rates, it seems to act at the level of

the gene rather than at the level of the nucleotide site.

Molecular chaperones play a critical role in mitigating the harm from mistranscription by

reducing misfolding. Genes that are chaperone clients might tolerate higher mistranscription

rates. Alternatively, sensitivity to mistranscription might select both for a lower

mistranscription rate and chaperone use. We found no support for either hypothesis; adding an

intercept term for *GroEL* chaperonin use was not a significant improvement on top of our final

model (p = 0.1).

As discussed in the Introduction, Cir-Seq data on the yeast *S. cerevisiae* indicates a much lower

mistranscription rate than *E. coli* (Gout et al. 2017), suggesting that it uses a global solution,

reducing site-specific selection pressures on mistranscription rates. We therefore do not predict

a relationship between gene expression and local mistranscription rate in this species, and do

not find one for the 11 non C→U substitution types (Figures 4 and 5; p=0.2 in our linear model

controlling for substitution type as a fixed effect).

Figure 4. No relationship between protein abundance and mistranscription rate in *S. cerevisiae* non-C$\rightarrow$U errors. Dashed line shows a simple linear model fitted to a pooled dataset of the 11 non-C$\rightarrow$U substitution types. Dotted line shows the linear model that corresponds to the *E. coli* "Final Model", with substitution type affecting the intercept but not the slope. To display this model, we took a weighted average of the intercept over substitution types as a function of the frequencies of opportunity to occur. Solid line shows the pooled data, binned by protein abundance as described in the Methods, and plotted according to mean protein abundance and the mean and 95% CI of the mistranscription rate within each bin. Data were divided into 10 bins; because of the limited availability of reads for low-expression genes, data within the first two bins were pooled. The y-axis spans one order of magnitude, as in Figure 1, making the two figures visually comparable in terms of the magnitude of a slope or lack thereof. Note that mistranscription rate is per possible error, so total mistranscription rate per nucleotide is around three times larger.

Figure 5. *S. cerevisiae* mistranscription rates broken down by base-substitution error types. Dotted lines show our final linear model, in which the intercept varies by error type, but the slope is the same across all error types. Dashed lines show linear models with both the slope and intercept fitted separately for each error type. Solid lines show the mean mistranscription rates, binned by protein abundance as described in the Methods, plotted according to mean protein abundance within each bin; error bars show 95% CI. Data were divided into 8 bins; because of the limited availability of reads for low-expression genes, data within the first three bins were pooled. Note that mistranscription rate is per possible error, so total mistranscription rate per nucleotide is around three times larger. The C→U error plot is shown inset due to radically different y-axis values than for the other error types.

However, C→U substitutions, which occur at much higher rates than other substitution types

and hence are subject to more selection even in *S. cerevisiae*, are less frequent for highly

abundant proteins (Figure 5; p= 0.006 for model restricted to C→U data). This confirms that the

protocol of Gout et al. (2017) succeeded in avoiding deamination events during sample

preparation (which should not depend on protein abundance), where that of Traverse &

Ochman (2016a) did not.

The high rate of mistranscription errors in *E. coli* came as a surprise to many (Traverse and

Ochman 2016a, b). This naturally raises the hypothesis that it is the data that are in error. While

the Cir-Seq technique is effective in preventing sequencing errors from inflating estimated

mistranscription rates (Acevedo and Andino 2014), it does not eliminate artifacts of the sample

preparation and analysis such as mutations occurring during the Cir-Seq experiment, nor

inaccurate mapping of reads to the genome. While these could artificially inflate estimated

mistranscription rates, we are not aware of any plausible mechanism by which the degree of

such inflation would be a function of protein abundance. Our results thus confirm the credibility

of the data, and hence of the statement that *E. coli* has a strikingly high non-C→U

mistranscription rate. After applying our quality filters, we calculate the total rate of all non-

C→U errors as 4.1×10^{-5} per site, or 8.6×10^{-5} if C→U errors are also included. In contrast, in *S.*

cerevisiae, we calculate from the data of Gout et al. (2017) a non-C→U mistranscription rate of

2.3×10^{-6}, or 3.5×10^{-6} with the C→U error type included.

The dependence of the *E. coli* mistranscription rate on the strength of selection (as reflected by

protein abundance), but not the *S. cerevisiae* mistranscription rate, is consistent with proposed

drift barrier explanations (McCandlish and Plotkin 2016; Rajon and Masel 2011; Xiong et al.

2017). In particular, *E. coli* is smaller and is generally accepted to have a larger effective

population size than *S. cerevisiae*. *E. coli* also has fewer loci, occurring within 4453 genes in K-12

(Riley et al. 2006) compared to 5178 genes in *S. cerevisiae* (Engel et al. 2014), which makes it

easier to evolve robustness at each one. A typical *E. coli* mRNA encodes 0.02% of the total

protein molecules in a cell, while a yeast mRNA encodes only 0.01% (Lu et al. 2007), meaning

that each transcription error in *E. coli* has a proportionately greater impact at one moment in

time. While a typical yeast mRNA has a longer half-life and so makes proteins over a longer time

(Siwiak and Zielenkiewicz 2013), this is not enough to counteract all other factors making local

solutions easier to evolve in *E. coli*.

We have shown that local mistranscription rates vary in a systematic way on a per-gene basis,

but have not determined the mechanisms by which expression error rates vary.

Mistranscription rates are affected at the gene level by the presence or absence of specific RNA

polymerase subunits (Thomas et al. 1998; Walmacq et al. 2009) or transcription factors

(Bubunenko et al. 2017; Irvin et al. 2014; Roghanian et al. 2015), and at a level higher than

nucleotide site but lower than gene by sequence characteristics such as long mononucleotide

repeats (Ackermann and Chao 2006; Gu et al. 2010). The degree to which such factors vary

systematically is unknown.

We have also shown that the local mistranscription rates even of highly expressed *E. coli* genes

are higher than the global mistranscription rate in *S. cerevisiae*, suggesting that *E. coli* genes are

somehow more robust to the consequences of mistranscription than are *S. cerevisiae* genes.

One sequence-based measure of robustness to transcription errors is the use or avoidance of

"fragile" codons that are one substitution away from a stop codon (Cusack et al. 2011). Serine,

arginine, glycine, and leucine can be encoded by either fragile or non-fragile codons. Of these

four amino acids, preference for fragile vs. robust codons is independent of %GC content only

in serine (TCT, TCC, AGT, and AGC are robust, while TCA and TCG are fragile), making serine

ideal for comparing codon fragility between species. We find that 25.7% of serine residues are

encoded by fragile codons in the *E. coli* proteome and 31.6% in *S. cerevisiae*. Dividing by the

expected frequency of robust serine residues (1/3) shows that *E. coli* uses fragile serine codons

77% as often as expected, compared to 95% in *S. cerevisiae*. These numbers suggest that

avoidance of fragile codons might explain at least some of the robustness of the *E. coli* genome.

However, the robustness associated with *E. coli*'s global solution is not so complete as to

eliminate selection for locally lower mistranscription rates in the genes subject to the strongest

selection, leading to the trend detected here.

Methods

Scripts used in these analyses are available at https://github.com/MaselLab/Meer-et-al-

Transcriptional-Error-Rates.

E. coli mistranscription data

Pre-processed data were obtained from Traverse and Ochman (2016a), that included how many

times each of the 4,641,652 nucleotide loci in the K-12 MG155 reference genome (GenBank

accession: NC.000913.3) was observed, and how often each nucleotide was seen there. We

assigned these loci to 4,140 protein coding genes and 178 non-coding genes using the

annotation of GenBank accession NC.000913.3. We analyzed the 3,935,551 nucleotide loci

within annotated non-overlapping protein-coding ORFs, and 47,344 nucleotide loci from non-

coding genes based on annotated 'start' and 'stop' positions. We excluded any sites that were

present in overlapping genes, as we could not assign a single error rate or protein abundance in

such cases.

Traverse and Ochman (2016a) data were obtained in multiple batches (referred to as

"replicates" in their data tables), with results reported only on two of the batches. Batch #2 had

approximately half as much data and twice the error rate of batch #1, so we restrict our

analysis to batch #1 only. Combining the data from each of the four experimental conditions

(minimal vs. rich media, and midlog vs. stationary phase) within batch #1 effectively yielded

15,742,204 protein-coding sites and 189,376 non-coding sites, where "site" is used here as

shorthand for condition×nucleotide locus, i.e. to describe the set of reads of a nucleotide locus

within just one experimental condition.

We excluded any site that had no reads and any protein-coding transcript site with no protein

abundance measure, leaving 5,994,463 coding and 182,233 non-coding sites. Each site can

experience three different substitution error types (e.g. C$\rightarrow$U, C$\rightarrow$A, and C$\rightarrow$G), which we

treated separately, yielding 17,983,389 coding and 546,699 non-coding "possible errors" for

analysis. Note that data for the three alternative errors at the same site are not, strictly

speaking, independent, because the occurrence of one error reduces the denominator for the

other two. However, at low error rates, this effect is negligible.

Mutations occurring during the Cir-Seq experiment, inaccurate mapping of reads to the

genome, or other artifacts of the experiment or pipeline can result in the appearance of

mistranscription "hot spots" that are best removed. We calculated the likelihoods of seeing that

many or more errors for each of the 18,530,088 possible errors being analyzed, using a

significance cutoff of 10^{-9} to ensure that only $10^{-9}\times18,530,088=0.02$ possible errors are falsely

excluded, or slightly more if there is genuine biological variation in mistranscription rates

beyond that captured by our linear model. We calculated likelihoods from a cumulative

binomial distribution based on the number of reads at that site and the rate of error expected

at that site from our model. When a possible error was excluded with likelihood $< 10^{-9}$, we

excluded the entire nucleotide locus (i.e. all three possible substitutions in all four conditions).

We performed an iterative procedure, first fitting a model of constant error rate for all non

C$\rightarrow$U errors and a separate error rate for C$\rightarrow$U errors, using expectations from this model to

exclude outliers, then using the cleaned-up data to develop a more sophisticated error rate

model of all conditions/substitution types, and using the revised expectations from this model

to update which loci should be excluded etc. until convergence. In the final iteration, one or

more possible errors was determined to be an outlier at 5 protein-coding and 2,390 non-coding

loci. For protein-coding outliers, we excluded all possible errors at each of the 5 outlier loci, i.e.

up to 60 possible errors (3 possible errors at 5 loci in 4 conditions). Some sites had no transcript

reads in some conditions, resulting in only 48 rather than 60 possible errors being excluded by

this procedure, leaving 17,983,341 possible errors in protein-coding transcript regions for

analysis. Excluding C→U substitutions, due to their significantly higher error rate and likelihood

of occurring post-transcriptionally, further reduced this to 16,466,559 non-C→U possible errors

for analysis.

S. cerevisiae mistranscription data

Similarly pre-processed transcript data were obtained from Gout et al. (2017), who recorded

how many times each nucleotide locus was observed in the S288C reference genome (GenBank

accession: GCA_000146045.2), to which the wild-type BY4741 strain used in their experiment is

very closely related. Only one experimental condition was used in this study. Using the same

methodology as for the *E. coli* data, we used the accession to assign nucleotide sites to the

5,983 protein-coding nuclear gene regions based on the annotated 'start' and 'stop' positions.

This process identified 8,853,931 nucleotide loci within annotated protein-coding ORFs,

resulting in 26,561,793 possible errors for analysis.

Excluding any transcript site without reads or with unreported or zero protein abundance left

us with 18,649,818 possible errors. Using our outlier detection protocol, we identified 44 loci

containing possible errors as outliers and excluded all possible errors at the associated loci (132

possible errors in total), leaving 18,649,686 possible errors for analysis.

C$\rightarrow$U errors were also identified as having a substantially higher error rate in the yeast data

(1.8×10^{-5} versus 2.3×10^{-6} for other mistranscription types), and were excluded from our analysis,

resulting in 17,394,875 non-C$\rightarrow$U possible errors.

Protein abundance data

Integrated protein abundance data were taken from PaxDB (Wang et al. 2015).

GroEL client status

We labelled the 1,929,741 possible errors associated with 252 *E. coli* proteins as having GroEL

client status, based on the identification of those proteins by Kerner et al. (2005) as specific

interactors with the GroEL chaperonin.

Statistical model

We modeled the error rate at site *i* within gene *j* as a linear function of the log-abundance of

protein *j*, i.e.

$$\frac{E_i}{R_i} = \rho + \beta \ln(Abundance_j)$$

390   where E_i is the number of reads containing a particular error and R_i is the total number of

391   reads at that nucleotide site.

393   To better model the error function in the linear model, we multiply both sides by R_i:

$$E_i \sim R_i + R_i \log_{10}(Abundance_j) + \varepsilon_{Poisson}$$

395   The observed number of errors E_i has the properties of count data, and so can be modeled as a

396   sample from a Poisson distribution. We fitted the statistical model above using a generalized

397   linear model function in R (glm, stats package), specifying the family of the model as

398   "poisson(link = identity)". For *E. coli*, experimental condition and type of error (excluding C→U)

399   were added as fixed effects to yield:

$$E_i \sim type : R_i + cond : R_i + R_i \log_{10}(Expression_j) + \varepsilon_{Poisson} \tag{1}$$

Slope as a function of expression level could also be made dependent on type and/or condition,

but adding these terms resulted in either insignificant or marginally significant improvements to

the *E. coli* model fit. For *S. cerevisiae*, the condition term does not apply, and expression was

not supported a predictive in the model. P-values associated with adding or removing terms to

a linear model were obtained using the anova command with the Chisq option to compare

nested models in R. P-values associated with comparisons to the final models for *E. coli* and *S.*

cerevisiae are given in the text.

Data Binning

We binned data by protein abundance for visualization and comparison to the fitted models. All

possible errors were sorted by the abundance value of the corresponding protein. Bin

boundaries were evenly spaced along our log-abundance axis between the 5% quantile and the

95% quantile, with data beyond these quantiles included in the edge bins. For each bin, one

point was plotted with y-value equal to the mean and 95% confidence interval of the

mistranscription rate and an x-value equal to the geometric mean of protein abundance. The

number of mistranscription errors observed is expected to follow a binomial distribution with r

trials, each with probability p of an error. We thus estimated a standard error of $\sqrt{(1-\hat{p})\hat{p}/r}$,

where r is the total number of reads within the bin and $\hat{p}$ is the observed error frequency

within the bin. To generate the 95% confidence interval we multiplied this standard error by

1.96. To keep standard errors for low-abundance bins reasonably low, data from several low-

abundance bins were combined.

Acknowledgements: This work was supported by the Undergraduate Biology Research Program

at the University of Arizona, the John Templeton Foundation [39667, 60814], and the National

Institutes of Health (GM104040). We thank Christophe Herman for helpful discussions and

430   **References**

Acevedo A, Andino R. 2014. Library preparation for highly accurate population sequencing of
RNA viruses. Nat Protoc. 9:1760–1769. doi: 10.1038/nprot.2014.118

Ackermann M, Chao L. 2006. DNA sequences shaped by selection for stability. PLoS Genetics.
2:e22.

Bubunenko MG, et al. 2017. A Cre Transcription Fidelity Reporter Identifies GreA as a Major
RNA Proofreading Factor in *Escherichia coli*. Genetics. 206:179-187. doi:
10.1534/genetics.116.198960

Chen G, et al. 2014. Cytosine Deamination Is a Major Cause of Baseline Noise in Next-
Generation Sequencing. Molecular Diagnosis & Therapy. 18:587-593. doi: 10.1007/s40291-014-
0115-2

Cusack BP, Arndt PF, Duret L, Crollius HR. 2011. Preventing dangerous nonsense: selection for
robustness to transcriptional error in human genes. PLoS Genetics. 7:e1002276.

Cutter AD, Charlesworth B. 2006. Selection intensity on preferred codons correlates with
overall codon usage bias in *Caenorhabditis remanei*. Curr Biol. 16:2053-7. doi:
10.1016/j.cub.2006.08.067

Dekel E, Alon U. 2005. Optimality and evolutionary tuning of the expression level of a protein.
Nature. 436:588-592.

Drummond DA, Wilke CO. 2009. The evolutionary consequences of erroneous protein
synthesis. Nat Rev Genet. 10:715-24. doi: 10.1038/nrg2662

Duret L, Mouchiroud D. 1999. Expression pattern and, surprisingly, gene length shape codon
usage in Caenorhabditis, Drosophila, and Arabidopsis. Proc Natl Acad Sci U S A. 96:4482-4487.
doi: 10.1073/pnas.96.8.4482

Engel SR, et al. 2014. The reference genome sequence of *Saccharomyces cerevisiae*: then and
now. G3: Genes, Genomes, Genetics. 4:389-398.

Frigola J, et al. 2017. Reduced mutation rate in exons due to differential mismatch repair. Nat
Genet. 49:1684-1692. doi: 10.1038/ng.3991

Geiler-Samerotte KA, et al. 2011. Misfolded proteins impose a dosage-dependent fitness cost
and trigger a cytosolic unfolded protein response in yeast. Proc Natl Acad Sci U S A. 108:680-
685. doi: 10.1073/pnas.1017570108

Good BH, Desai MM. 2014. Deleterious Passengers in Adapting Populations. Genetics.
198:1183-1208. doi: 10.1534/genetics.114.170233

Gout J-F, et al. 2017. The landscape of transcription errors in eukaryotic cells. Sci Adv.
3:e1701484. doi: 10.1126/sciadv.1701484

Gout J-F, et al. 2013. Large-scale detection of in vivo transcription errors. Proc Natl Acad Sci U S
A. 110:18584-18589. doi: 10.1073/pnas.1309843110

Gu T, et al. 2010. Avoidance of long mononucleotide repeats in codon pair usage. Genetics.
186:1077-1084.

Irvin JD, et al. 2014. A genetic assay for transcription errors reveals multilayer control of RNA
polymerase II fidelity. PLoS Genet. 10:e1004532. doi: 10.1371/journal.pgen.1004532

Kafri M, Metzl-Raz E, Jona G, Barkai N. 2016. The Cost of Protein Production. Cell Rep. 14:22-31.
doi: 10.1016/j.celrep.2015.12.015

Kerner MJ, et al. 2005. Proteome-wide Analysis of Chaperonin-Dependent Protein Folding in
Escherichia coli. Cell. 122:209-220. doi: 10.1016/j.cell.2005.05.028

Liu Z, Zhang J. 2018a. Human C-to-U Coding RNA Editing Is Largely Nonadaptive. Mol Biol Evol.
35:963-969. doi: 10.1093/molbev/msy011

Liu Z, Zhang J. 2018b. Most m^{6}A RNA Modifications in Protein-Coding Regions Are Evolutionarily
Unconserved and Likely Nonfunctional. Mol Biol Evol. 35:666-675. doi:
10.1093/molbev/msx320

Lu P, et al. 2007. Absolute protein expression profiling estimates the relative contributions of
transcriptional and translational regulation. Nature Biotechnology. 25:117-124. doi:
10.1038/nbt1270

Lynch M. 2007. *The origins of genome architecture*. Sunderland: Sinauer Associates.

Lynch M, Marinov GK. 2015. The bioenergetic costs of a gene. Proc Natl Acad Sci U S A.
112:15690-5. doi: 10.1073/pnas.1514974112

McCandlish DM, Plotkin JB. 2016. Transcriptional errors and the drift barrier. Proc Natl Acad Sci
U S A. 113 3136-3138. doi: 10.1073/pnas.1601785113

McCutcheon JP, Moran NA. 2012. Extreme genome reduction in symbiotic bacteria. Nat Rev
Microbiol. 10:13-26. doi: 10.1038/nrmicro2670

Mordret E, et al. 2018. Systematic detection of amino acid substitutions in proteome reveals a
mechanistic basis of ribosome errors. bioRxiv.255943. doi: 10.1101/255943

Ohta T. 1973. Slightly Deleterious Mutant Substitutions in Evolution. Nature. 246:96-98. doi:
10.1038/246096a0

Petrov DA, Hartl DL. 2000. Pseudogene evolution and natural selection for a compact genome. J
Hered. 91:221-227.

Rajon E, Masel J. 2011. The evolution of molecular error rates and the consequences for
evolvability. Proc Natl Acad Sci U S A. 108:1082-1087. doi: 10.1073/pnas.1012918108

Ran W, Kristensen DM, Koonin EV. 2014. Coupling between protein level selection and codon
usage optimization in the evolution of bacteria and archaea. mBio. 5:e00956-14. doi:
10.1128/mBio.00956-14

Riley M, et al. 2006. Escherichia coli K-12: a cooperatively developed annotation snapshot—
2005. Nucleic Acids Res. 34:1-9.

Roghanian M, Zenkin N, Yuzenkova Y. 2015. Bacterial global regulators DksA/ppGpp increase
fidelity of transcription. Nucleic Acids Res. 43:1529-36. doi: 10.1093/nar/gkv003

Scott M, Klumpp S, Mateescu EM, Hwa T. 2014. Emergence of robust growth laws from optimal
regulation of ribosome synthesis. Mol Syst Biol. 10:747. doi: 10.15252/msb.20145379

Sharp PM, Emery LR, Zeng K. 2010. Forces that influence the evolution of codon bias. Philos
Trans R Soc Lond B Biol Sci. 365:1203-12. doi: 10.1098/rstb.2009.0305

Siwiak M, Zielenkiewicz P. 2013. Transimulation - Protein Biosynthesis Web Service. PLoS ONE.
8:e73943. doi: 10.1371/journal.pone.0073943

Thomas MJ, Platas AA, Hawley DK. 1998. Transcriptional fidelity and proofreading by RNA
polymerase II. Cell. 93:627-637.

Tomala K, Korona R. 2013. Evaluating the fitness cost of protein expression in *Saccharomyces*
cerevisiae. Genome Biol Evol. 5:2051-60. doi: 10.1093/gbe/evt154

Traverse CC, Ochman H. 2016a. Conserved rates and patterns of transcription errors across
bacterial growth states and lifestyles. Proc Natl Acad Sci U S A. 113:3311-3316. doi:
10.1073/pnas.1525329113

Traverse CC, Ochman H. 2016b. Correction for Traverse and Ochman, Conserved rates and
patterns of transcription errors across bacterial growth states and lifestyles. Proc Natl Acad Sci
U S A. 113:E4257-E4258. doi: 10.1073/pnas.1609677113

Vicario S, Moriyama EN, Powell JR. 2007. Codon usage in twelve species of Drosophila. BMC
Evol Biol. 7:226. doi: 10.1186/1471-2148-7-226

Wagner A. 2007. Energy costs constrain the evolution of gene expression. J Exp Zool Part B.
308B:322-324. doi: 10.1002/jez.b.21152

Walmacq C, et al. 2009. Rpb9 subunit controls transcription fidelity by delaying NTP
sequestration in RNA polymerase II. J Biol Chem. 284:19601-12. doi: 10.1074/jbc.M109.006908

Wang M, et al. 2015. Version 4.0 of PaxDb: Protein abundance data, integrated across model
organisms, tissues, and cell-lines. Proteomics. 15:3163-3168. doi: 10.1002/pmic.201400441

Xiong K, McEntee JP, Porfirio DJ, Masel J. 2017. Drift Barriers to Quality Control When Genes
Are Expressed at Different Levels. Genetics. 205:397-407. doi: 10.1534/genetics.116.192567

Xu C, Park J-K, Zhang J. 2019. Evidence that alternative transcriptional initiation is largely
nonadaptive. PLoS Biology. 17:e3000197. doi: 10.1371/journal.pbio.3000197

Xu C, Zhang J. 2018. Alternative Polyadenylation of Mammalian Transcripts Is Generally
Deleterious, Not Adaptive. Cell Syst. 6:734-742. doi: 10.1016/j.cels.2018.05.007

Zhang D, et al. 2016. The Effect of Codon Mismatch on the Protein Translation System. PLoS
ONE. 11:e0148302. doi: 10.1371/journal.pone.0148302