Title:

Intra- and Inter-Specific Investigations of Skeletal DNA Methylation and Femur Morphology in Primates

Running Title:

Skeletal DNA Methylation Variation in Primates

Authors and Affiliations:

Genevieve Housman[1,2*], Ellen E. Quillen[3], and Anne C. Stone[1,2]

[1]School of Human Evolution and Social Change, Arizona State University, Tempe, AZ, USA.

[2]Center for Evolution and Medicine, Arizona State University, Tempe, AZ, USA.

[3]Department of Genetics, Texas Biomedical Research Institute, San Antonio, TX, USA.

*Corresponding author:

Genevieve Housman, Section of Genetic Medicine, University of Chicago, 920 East 58th Street, CLSC

317, Chicago, IL 60637, USA. Email: ghousman@uchicago.edu

Author Notes:

Genevieve Housman is currently affiliated with the University of Chicago, and Ellen E. Quillen is

currently affiliated with Wake Forest School of Medicine.

Abstract:

Objectives

Epigenetic mechanisms influence the development and maintenance of complex phenotypes and may also

contribute to the evolution of species-specific phenotypes. With respect to skeletal traits, little is known

about the gene regulation underlying these hard tissues or how tissue-specific patterns are associated with

bone morphology or vary among species. To begin exploring these topics, this study evaluates one

epigenetic mechanism, DNA methylation, in skeletal tissues from five nonhuman primate species which

display anatomical and locomotor differences representative of their phylogenetic groups.

Materials and Methods

First, we test whether intra-specific variation in skeletal DNA methylation is associated with intra-specific

variation in femur morphology. Second, we identify inter-specific differences in DNA methylation and

assess whether these lineage-specific patterns may have contributed to species-specific morphologies.

Specifically, we use the Illumina Infinium MethylationEPIC BeadChip to identify DNA methylation

patterns in femur trabecular bone from baboons (n=28), macaques (n=10), vervets (n=10), chimpanzees

(n=4), and marmosets (n=6).

Results

Significant differentially methylated positions (DMPs) were associated with a subset of morphological

variants, but these likely have small biological effects and may be confounded by other variables

associated with morphological variation. Conversely, several species-specific DMPs were identified, and
these are found in genes enriched for functions associated with complex skeletal traits.
45
46 *Discussion*
47 Overall, these findings reveal that while intra-specific epigenetic variation is not readily associated with
48 skeletal morphology differences, some inter-specific epigenetic differences in skeletal tissues exist and
49 may contribute to evolutionarily distinct phenotypes. This work forms a foundation for future
50 explorations of gene regulation and skeletal trait evolution in primates.
51
52 **Key Words:** DNA methylation, nonhuman primates, evolution, epigenome, bone
53
Introduction:
Primates distinguish themselves from other mammals with their unique suite of anatomical
features that initially enabled arboreal niche occupation and subsequently evolved to fit a myriad of
habitats and forms of locomotion, the most unique being hominin bipedalism. The resulting
morphological variation that has evolved across taxa has its foundation in underlying skeletal anatomies.
Such skeletal morphologies are used to both characterize extant primate diversity and reconstruct the
anatomy and locomotor capabilities of extinct primate species (Ankel-Simons, 2007; Fleagle, 1999). The
range of mechanisms that enable the development of skeletal features are not entirely understood, though.
Skeletal features related to varied body forms are often described as the result of environmental
adaptations. However, skeletal morphology is more accurately defined as the result of complex processes,
including environmental, genetic, and epigenetic mechanisms. While environmental (Lewton, 2017;
Lewton, Ritzman, Copes, Garland, & Capellini, 2018) and genetic (Joganic et al., 2017; Ritzman et al.,
2017) factors have been readily explored, the important roles of epigenetic factors, such as DNA
methylation, in skeletal tissue development and maintenance have only recently been identified (Delgado-
Calle et al., 2013; Rushton et al., 2014; Simon & Jeffries, 2017).
For instance, epigenetic processes are influential in regulating skeletal muscle development
(Palacios & Puri, 2006) which can impact the adjacent skeletal scaffolding system. Several genes
involved in human skeletal development appear to be differentially methylated across fetal and adult
developmental stages (de Andrés et al., 2013). Lastly, methylation variation in humans and model
organisms has been implicated in several skeletal pathologies and disorders, such as osteoporosis and
osteoarthritis (Delgado-Calle et al., 2013; den Hollander et al., 2014; Jeffries et al., 2016; Moazedi-Fuerst
et al., 2014; Morris et al., 2017; Ostanek, Kranjc, Lovšin, Zupan, & Marc, 2018; Rushton et al., 2014;
Simon & Jeffries, 2017). Some of these studies are the first to assess methylation patterns in human
skeletal tissues, and such steps are crucial for identifying the relationship between epigenetic variation
and skeletal phenotypic variation.
The contributions of epigenetics to primate phenotypic variation were first considered by King
and Wilson (1975), who proposed that anatomical and behavioral differences between humans and
chimpanzees were more likely "based on changes in the mechanisms controlling the expression of genes
than on sequence changes in proteins" (King & Wilson, 1975). Studies to understand methylation
variation across species began soon afterwards (Gama-Sosa et al., 1983). General changes to mammalian
epigenomes have been examined (Sharif, Endo, Toyoda, & Koseki, 2010), but most epigenetics work in

primates has focused on humans – how it varies across distinct tissues within individuals (Slieker et al.,
2013), across different individuals (Petronis et al., 2003), across populations (Heyn et al., 2013), in
relation to aging processes (Fraga et al., 2005), and in relation to diet (Shelnutt et al., 2004), as well as
how it is inherited across generations (Flanagan et al., 2006). These studies have identified interesting
intra-specific methylation variation present in humans.
Similarly, epigenetic variation has been identified among primate species. Inter-specific variation
in epigenetic signatures was initially inferred from underlying genomic sequences (Bell et al., 2012). For
instance, several promoter CpG densities vary across primates. These likely relate to regulatory
methylation differences across species as primate CpG densities correlate with methylation levels (Weber
et al., 2007). Additionally, gene expression studies, which primarily focus on brain tissues (Babbitt et al.,
2010; Cáceres et al., 2003) and a small set of other soft tissues (Blake et al., 2017; Blekhman, Oshlack,
Chabot, Smyth, & Gilad, 2008; Karere et al., 2013; Pavlovic, Blake, Roux, Chavarria, & Gilad, 2018;
Tung, Zhou, Alberts, Stephens, & Gilad, 2015), have also noted regulatory differences across species.
Methylation differences in brain tissues have evolved across primates and contributed to resultant brain
phenotypes and disease vulnerabilities (Enard et al., 2004; Farcas et al., 2009; Madrid, Chopra, & Alisch,
2018; Mendizabal et al., 2016; Provencal et al., 2012; Zeng et al., 2012). Thus, methylation-phenotype
relationships can be identified in primates. Primate methylation patterns in blood cells and other soft
tissues have also been studied, but not to the same degree (Fukuda et al., 2013; Gao et al., 2017;
Hernando-Herraez et al., 2013; Lea, Altmann, Alberts, & Tung, 2016; Lindskog et al., 2014; Martin et al.,
2011; Molaro et al., 2011; Pai, Bell, Marioni, Pritchard, & Gilad, 2011; Vilgalys, Rogers, Jolly,
Mukherjee, & Tung, 2018).
Interestingly, two studies using soft tissues and blood identified differential methylation and
expression of genes essential for skeletal development (*RUNX1*, *RUNX3*, and *COL2A1*) between some
primates (Hernando-Herraez et al., 2013; Lindskog et al., 2014). Additionally, the emerging field of
ancient epigenetics, which reconstructs methylation patterns from ancient DNA degradation patterns in
hominin remains (Smith, Monroe, & Bolnick, 2015; Gokhman et al., 2014; Gokhman, Meshorer, &
Carmel, 2016; Gokhman et al., 2017), have found that other skeletal developmental genes (*HOXD*
complex) are differentially methylated among modern humans and ancient hominins (Gokhman et al.,
2014). These findings suggest that primates do exhibit distinct epigenetic patterns and that primate
skeletal epigenetics is an important area of research for uncovering developmental and evolutionary
patterns relevant for complex skeletal traits, such as anatomy, locomotion, and disease. Despite these
potential avenues of insight, studies of nonhuman primate (NHP) skeletal epigenetics are limited
(Housman, Havill, Quillen, Comuzzie, & Stone, 2018).
Overall, there are clear knowledge gaps in our understanding of NHP skeletal complexity in
relation to epigenetic variation and epigenetic differences between phylogenetically diverse NHP species.
The present study begins to remedy this by assessing how genome-wide and gene-specific DNA
methylation in primate skeletal tissues varies intra- and inter-specifically and in relation to femur form.
Specifically, for this study, we explored the evolution of the epigenome and its relation to
nonpathological skeletal traits by identifying DNA methylation patterns in femur trabecular bone from
baboons, macaques, vervets, chimpanzees, and marmosets and assessing intra- and inter-specific
methylation variation and its relation to morphology. With this study design, we aimed to assess whether
skeletal methylation variation within species is associated with skeletal morphology variation, to what

degree skeletal methylation varies among species, and whether species-specific changes in methylation
further inform our understanding of skeletal morphology evolution. Additionally, this dataset expands the
number of characterized skeletal methylation patterns in NHPs for future primate skeletal epigenetics
investigations.
131
132 **Materials and Methods:**
133
134 *Ethics Statement*
135 NHP tissue samples included were opportunistically collected at routine necropsy of these animals. No
136 animals were sacrificed for this study, and no living animals were used in this study. Chimpanzee tissues
137 were collected opportunistically during routine necropsy prior to the September 2015 implementation of
138 Fish and Wildlife Service rule 80 FR 34499.
139
NHP Samples
NHP samples come from captive colonies of chimpanzees (*Pan troglodytes*), baboons (*Papio*
spp.), rhesus macaques (*Macaca mulatta*), and marmosets (*Callithrix jacchus*) from the Southwest
National Primate Research Center in Texas, as well as vervets (*Chlorocebus aethiops*) from the Wake
Forest/UCLA Vervet Research Colony in North Carolina. Femora were opportunistically collected at
routine necropsy of these animals and stored in -20°C freezers at the Texas Biomedical Research Institute
after dissection. These preparation and storage conditions ensured the preservation of skeletal DNA
methylation patterns. Samples include baboons (n=28), macaques (n=10), vervets (n=10), chimpanzees
(n=4), and marmosets (n=6). Age ranges span adulthood for each species and are comparable between
each group (Figure 1, Table S1). Scaled ages were also calculated for each sample (Table S1) using
published life expectancies for each species (Rowe, 1996). Both sexes are represented (female: n=33,
male: n=24, unknown: n=1).
152
Assessment of Femur Morphologies
On the right femora of NHP samples, 29 linear morphology traits (Figure 2, Table S6) were
measured using calipers. These measurements characterize overall femur shape (McHenry & Corruccini,
1978; Terzidis et al., 2012). All measurements were collected by one researcher, and intra-observer error
for each measurement was determined by performing triplicate measurements on approximately 10% of
the samples in each species. These measurements were spaced throughout the entire data collection
period. Error was calculated as the mean absolute difference divided by the mean (Corner, Lele, &
Richtsmeier, 1992; White & Folkens, 2000). All measurements that were retained for downstream
analyses had errors of less than 5%, and the only measurement excluded was macaque intercondylar
notch depth (error = 6.62%) (File S2).
163
Genome-Wide DNA Methylation Profiling
Trabecular bone cores were obtained from the medial condyles on the right distal femora of each
NHP sample using a drill press that cored transversely through the condyle leaving the articular surface
preserved (Supplemental Text). Cortical bone was removed from cores using a Dremel, and the remaining
trabecular bone cores were pulverized into bone dust using a SPEX SamplePrep Freezer/Mill. DNA was

extracted from femoral trabecular bone using a phenol-chloroform protocol optimized for skeletal tissues
(Barnett and Larson 2012). Genome-wide DNA methylation was assessed using Illumina Infinium
MethylationEPIC microarrays (EPIC array) (Supplemental Text). The array data discussed in this
publication have been deposited in NCBI's Gene Expression Omnibus and are accessible through GEO
SuperSeries accession number GSE103332, which includes the SubSeries accession numbers
GSE103279, GSE103271, GSE103280, GSE94677, GSE103328, and GSE103287.
175

Methylation Data Processing
Using previously described methods (Housman et al., 2018), raw EPIC array data were
normalized and converted to β values which represent the average methylation levels at each site (0 =
completely unmethylated sites, 1 = fully methylated sites), and M values which are the log transformed
ratio of methylated signal to unmethylated signal. Probes with failed detection levels (p-value > 0.05) in
greater than 10% of samples and samples with greater than 30% of probes with detection level failures
were removed from downstream analyses. Using previously described methods (Housman et al., 2018),
probes with sequence mismatches to NHP genomes, which could produce biased methylation
measurements, were computationally filtered out and excluded from downstream analyses (Supplemental
Text, Figures S1-S3, Tables S2-S5, File S1). Additionally, cross-reactive probes (McCartney et al., 2016),
probes containing SNPs at the CpG site, probes detecting SNP information, probes detecting methylation
at non-CpG sites, and probes targeting sites within the sex chromosomes were removed (Aryee et al.,
2014; Fortin, Triche, & Hansen, 2016). Lastly, we assessed whether NHP SNPs overlapped with targeted
probes (Supplemental Text, Table S2, File S1). The resulting finalized datasets (Figure S4) were then
used in further statistical analyses.
191

Statistical Analysis of Differential Methylation
In order to identify sites that were significantly differentially methylated across comparative
groups, we designed and tested general linear models (GLMs) which related the variables of interest
(morphological measures and species membership) to the DNA methylation patterns for each site, while
accounting for the effects of additional variables, batch effects, and latent variables (Maksimovic,
Phipson, & Oshlack, 2016). Sites found to have significant associations were classified as significant
differentially methylated positions (DMPs). Overall, GLMs were used to estimate differences in
methylation levels associated with femur morphologies within each taxonomic group (intra-specific) and
between each taxonomic group (inter-specific). Specific details about each set of GLMs are provided in
the following sections with additional details in the Supplemental Text.
202

Intra-Specific Analyses
For the intra-specific analyses, variables included in each GLM were the femur morphologies
within each taxonomic group, sex, age, and steady state weight when known, as well as unknown latent
variables calculated using the iteratively re-weighted least squares approach in the sva package in R (Jaffe
& Irizarry, 2014; Leek, Johnson, Parker, Jaffe, & Storey, 2012; Leek & Storey, 2007, 2008). Latent
variables estimated for each morphology were included to help mitigate any unknown batch and cell
heterogeneity effects on methylation variation at each site (Table S7). Each GLM design matrix was fit to
corresponding M value array data by generalized least squares using the limma package in R (Huber et

al., 2015; Phipson, Lee, Majewski, Alexander, & Smyth, 2016; Ritchie et al., 2015), and the estimated
coefficients and standard errors for each morphology were computed. Lastly, for each coefficient, an
empirical Bayes approach was applied using the limma package in R (Lönnstedt & Speed, 2002;
McCarthy & Smyth, 2009; Phipson, Lee, et al., 2016; Smyth, 2004) to compute moderated t-statistics,
log-odds ratios of differential methylation, and associated p-values adjusted for multiple testing
(Benjamini & Hochberg, 1995). Significant DMPs for the effect of each morphology were defined as
those having log fold changes in M values corresponding to an adjusted p-value of less than 0.05. Lastly,
the gene ontology (GO) and KEGG pathway enrichment for significant CpGs was determined using the
missMethyl package in R (Benjamini & Hochberg, 1995; Geeleher et al., 2013; Phipson, Maksimovic, &
Oshlack, 2016; Ritchie et al., 2015; Young, Wakefield, Smyth, & Oshlack, 2010), which takes into
account the differing number of probes per gene present on the array.
222
Inter-Specific Analyses
For the inter-specific analyses, variables included in the GLM were taxonomic grouping, sex,
age, known batch effects (e.g., array number and position), and unknown latent variables calculated using
the method described above. An additional GLM which replaced the age variable with ages scaled to
species-specific life expectancies (Rowe, 1996) was also tested. The six latent variables estimated were
included to help mitigate any unknown batch and cell heterogeneity effects on methylation variation at
each site. The GLM design matrix was fit to the M value array data using the method described above,
and the estimated coefficients and standard errors for taxonomic group effects were computed. As
described above, moderated t-statistics, log-odds ratios of differential methylation, and associated p-
values adjusted for multiple testing were computed, and significant DMPs for the effect of taxonomy
were defined as those having log fold changes in M values corresponding to an adjusted p-value of less
than 0.05.
To determine only those methylation differences that represent fixed changes between genera, we
used methods similar to those described in (Hernando-Herraez et al., 2013). Briefly, significant DMPs
were identified between all possible pairwise comparisons of taxa (n=10: baboon-macaque, baboon-
vervet, baboon-chimpanzee, baboon-marmoset, macaque-vervet, macaque-chimpanzee, macaque-
marmoset, vervet-chimpanzee, vervet-marmoset, chimpanzee-marmoset). A significant DMP was then
defined as taxon-specific if it was found to be significant in all four pairwise comparisons containing the
taxon of interest but not found in any of the remaining pairwise comparisons. The GO and KEGG
pathway enrichment of these DMPs was then determined as described above.
Additionally, global changes in methylation were calculated using distance matrices (Hernando-
Herraez et al., 2013) of the methylation levels for all finalized 39,802 filtered probes. These changes were
assessed at a species-level by averaging the β values per probe within each species. We then used
Euclidean distances to calculate the difference between every two species. Neighbor joining trees were
estimated from these distances using the ape package in R (Paradis, Claude, & Strimmer, 2004). For each
resulting tree, 1000 bootstraps were performed to determine confidence values for each branch. Global
changes in methylation were also assessed at the individual-level using Euclidean distances to calculate
the difference between every two individuals.
251
DNA Methylation Profiling and Analyses of HOXD10

Based on the inter-specific DNA methylation patterns identified in this study and those identified
in other evolutionary anthropological studies (Gokhman et al. 2014), the *HOXD10* gene was selected for
subsequent DNA methylation profiling and analysis at a higher resolution using gene-specific sequencing
techniques. Specifically, primers were designed and optimized to PCR amplify regions spanning across
the entire *HOXD10* gene, as well as upstream and downstream several hundred bases (hg19
chr2:176980532-176985117), in each NHP species for regular and bisulfite treated DNA (Tables S14-
S17). All gene-specific assays were performed in a subset of the samples tested using the EPIC array and
included chimpanzees (n=3), baboons (n=3), macaques (n=3), vervets (n=3), and marmosets (n=3) (Table
S1). Additionally, a subset of these assays that targeted the region around one test locus (cg02193236)
were performed in all of the samples (Table S1). As described above, DNA was extracted from femoral
trabecular bone using a phenol-chloroform protocol optimized for skeletal tissues (Barnett and Larson
2012). DNA was bisulfite converted using the EZ DNA MethylationTM Gold Kit according to the
manufacturer's instructions (Zymo Research). Successful PCR amplification was confirmed using gel
electrophoresis. Gene-specific PCR products were then purified using an exonuclease I and shrimp
alkaline phosphatase protocol and sequenced on the Applied Biosystems 3730 capillary sequencer at the
DNA Laboratory at Arizona State University.
Regular and bisulfite sequences were aligned to the appropriate NHP references within the
Enredo-Pecan-Orthus (EPO) whole-genome multiple alignments of several primate genomes [Ensembl
Compara.8_primates_EPO] (Paten, Herrero, Beal, et al. 2008; Paten, Herrero, Fitzgerald, et al. 2008)
using MEGA7 (Kumar et al. 2016). Manual annotation of these sequences within each sample confirmed
that the gene sequences belong to the appropriate primate species and that the regular and bisulfite treated
sequences only differ in cytosine composition. The number and distribution of methylated loci throughout
the *HOXD10* gene were then identified and compared within and among species to provide a higher
resolution of methylation variation within this targeted gene. Lastly, with respect to the test locus
(cg02193236) evaluated in all samples, methylation levels across this region were compared with
corresponding methylation levels of this region as determined using the EPIC array.
279
Results:
The aim of this study was to identify DNA methylation patterns in skeletal tissues from several
NHP species in order to determine how skeletal methylation varies at different taxonomic scales and in
relation to complex skeletal traits. We evaluated DNA methylation patterns in femur trabecular bone from
five NHP species – baboons, macaques, vervets, chimpanzees, and marmosets (Figure 1, Table S1). We
used the Illumina Infinium MethylationEPIC BeadChip (EPIC array), which was determined to be
effective at identifying DNA methylation patterns in NHP DNA (Supplemental Text, Figures S1-S4,
Tables S2-S5, File S1). Further, we selected probes appropriate for intra- and inter-specific comparison,
which removed the effect of sequence differences among individuals and species as a reason for
methylation differences. With these data, we first tested whether intra-specific variation in skeletal DNA
methylation is associated with intra-specific variation in femur morphology. Second, we identified inter-
specific differences in DNA methylation and assessed whether these lineage-specific patterns may have
contributed to species-specific morphologies.
293
Genome-Wide Intra-Specific Differential Methylation and Morphological Variation

Measurements of 29 linear morphology traits (Figure 2, Table S6) were collected from each NHP
right femur. All measurements had less than 5% error, except those for intercondylar notch depth in
macaques (Figure S5, File S2). Significant DMPs associated with each intra-specific linear morphology
were interrogated from 189,858 sites in in baboons, 190,898 sites in macaques, 191,639 sites in vervets,
576,804 sites in chimpanzees, and 68,709 sites in marmosets (Tables S7-S8). In baboons (n=28), 1 DMP
was hypomethylated with increasing bicondylar femur length and increasing maximum femur length. In
macaques (n=10), 1 DMP was hypermethylated with increasing proximal femur width, 1 DMP was
hypermethylated with increasing medial condyle width, and 6 DMPs were hypomethylated with
increasing medial condyle width. In vervets (n=10), 1 DMP was hypomethylated with increasing superior
shaft width, 2 DMPs were hypomethylated with increasing inferior shaft width, and 1 DMP was
hypermethylated with increasing anatomical neck height. In chimpanzees (n=4), 216 DMPs were
hypomethylated and 57 DMPs were hypermethylated with increasing anatomical neck length. Lastly, in
marmosets (n=6), no DMPs were associated with morphological variation (Table S9, File S3).
While the maximum absolute change in mean methylation ($\Delta\beta$) for most of these DMPs is greater
than 10% ($\Delta\beta = 0.1$), the change in methylation between the individual with the largest morphology
measurement and the individual with the smallest morphology measurement is less than 0.1 for several
DMPs (File S3). Tests for enrichment of gene ontology (GO) and KEGG pathway functions were done
for the 3 intra-specific morphologies that had more than 2 DMPs associated with them. However, no GO
biological processes were found to be enriched in DMPs associated with either macaque medial condyle
width or chimpanzee anatomical neck length. Additionally, KEGG pathway functions were only found to
be enriched in DMPs associated with chimpanzee anatomical neck length, and these pathways were
predominantly involved in immune system cell signaling and differentiation (Table S10).
317

Genome-Wide Inter-Specific Differential Methylation
To determine how methylation varies inter-specifically, differential methylation was interrogated
from the 39,802 probes that were shared among NHP species (Table S11, File S4). Species-specific
DMPs were determined by identifying DMPs that were significant in all 4 pairwise comparisons
containing the taxon of interest but not in any of the remaining pairwise comparisons. These methods
identified 650 species-specific DMPs in baboons, 257 in macaques, 639 in vervets, 2,796 in chimpanzees,
and 13,778 in marmosets (Table S11). Similar results were obtained using a model that accounted for
scaled age based on life expectancy instead of exact age (660 baboon-specific DMPs, 260 macaque-
specific DMPs, 651 vervet-specific DMPs, 2,720 chimpanzee-specific DMPs, and 13,689 marmoset-
specific DMPs) (File S4). Thus, only the species-specific DMPs identified using the model that accounted
for exact age were used in downstream analyses. Additionally, given the disproportionate number of
marmoset-specific DMPs identified, caution should be used when interpreting the biological relevance of
these data. Comparably, methylation patterns also distinguish broader taxonomic groups, such as
cercopithecoids (baboons, macaques, and vervets), apes (chimpanzees), and platyrrhines (marmosets).
Specifically, 2,655 DMPs were found to be specific to cercopithecoids, 1,869 were found to be ape-
specific, and 14,985 were found to be specific to platyrrhines (File S5). It should be noted that the ape and
platyrrhine categories only include one taxon each – chimpanzee and marmoset, respectively. Similar
results were obtained using a model that accounted for scaled age based on life expectancy instead of

exact age (2,660 cercopithecoid-specific DMPs, 1,797 ape-specific DMPs, and 14,963 platyrrhine-
specific DMPs) (File S5).
Species-specific DMPs spanned 7,320 genes with an average of 2.2±2.0 (s.d.) significant probes
per gene (Table S12). Over half of these genes contain just 1 significant species-specific DMP (n=3,816),
and a portion of these are due to only 1 EPIC array probe targeting each gene (n=1,788). The remaining
genes that contain at least 2 significant species-specific DMPs (n=3,503) have an average of 3.4±2.3 (s.d.)
significant probes per gene (range: 2-34). Additionally, these species-specific DMPs covered a range of
locations with respect to genes and CpG islands (Table S12), indicating that these species-specific
changes in methylation are distributed throughout the genome. Using various $\Delta\beta$ cutoff thresholds
decreases the final number of species-specific DMPs to varying degrees (Table S11, File S4). Counter to
these decreases in numbers, though, species-specific DMPs cover a range of locations with respect to
genes and CpG islands regardless of the $\Delta\beta$ cutoff threshold (Table S12).
Overall, across $\Delta\beta$ cutoff thresholds, more species-specific DMPs were found in the platyrrhine
marmosets, followed by the great ape chimpanzees, and lastly the cercopithecoid baboons, macaques, and
vervets. Additionally, the proportions of hypermethylated and hypomethylated species-specific DMPs
within each taxon remain fairly constant across $\Delta\beta$ cutoff thresholds (Table S11). In baboons, macaques,
vervets, and chimpanzees, more than half of all species-specific DMPs show patterns of
hypermethylation, and in marmosets, more than half of all species-specific DMPs show patterns of
hypomethylation. The only disruption in these trends is in chimpanzees when no $\Delta\beta$ threshold is applied.
Nevertheless, species-specific DMPs with different $\Delta\beta$ cutoff thresholds do differ in their abilities to
cluster animals into taxonomic groups (Figure S6, File S4). For all thresholds, apes, cercopithecoids, and
platyrrhines cluster into distinct groups. However, within cercopithecoids, vervets only cluster into a
distinct species group with a $\Delta\beta \geq 0.2$ threshold, and the baboon-macaque clade require a $\Delta\beta \geq 0.3$
threshold. Lastly, species-specific clustering of baboons, macaques, and vervets only occurs with a $\Delta\beta \geq$
0.4 threshold (Figure 3).
Additionally, global changes in methylation across all 39,802 probes were evaluated. Average
global changes within each species reveal that apes, cercopithecoids, and platyrrhines are
phylogenetically distinct from one another, and these divergences are well supported (Figure 4).
Similarly, when phylogenetic relationships are evaluated using the global changes in methylation of
individual animals, distinct lineages are again formed between apes, cercopithecoids, and platyrrhines
(Figure S7). However, within the cercopithecoid clade, several poorly supported branches result in
baboons, macaques, and vervets not forming distinct lineages. Phylogenetic separation of these
cercopithecoid species into distinct lineages is only possible when the methylation changes considered are
reduced to only include species-specific DMPs with a $\Delta\beta \geq 0.4$ threshold (Figure S8).
Species-specific DMPs also show associations with genes that have a wide array of GO biological
processes (File S6) and KEGG pathway functions (File S7). Cellular adhesion is a primary GO function
found to be highly enriched in species-specific DMPs from baboons, macaques, vervets, chimpanzees,
and marmosets. Species-specific DMPs for chimpanzees and marmosets are also enriched for genes
involved in the regulation of transcription and gene expression. Additionally, enrichment of genes
involved in anatomical developmental processes is found in baboons, vervets, chimpanzees, and
marmosets. Chimpanzees and marmosets show further enrichment of genes contributing to pattern
specification processes, limb development, and skeletal system development. Moreover, marmoset

species-specific DMPs are enriched for genes with functions very closely related to skeletal development,
such as osteoblast differentiation and ossification, as well as genes involved in metabolism and the
development of other organ systems including skeletal muscles, nerves, the brain, the heart, blood vessels,
kidneys, eyes, and ears (File S6). Several enriched pathways reinforce these molecular functions, and
additional pathways related to cancers and other disease were also identified (File S7).
Lastly, out of the species-specific DMPs identified, some were found to overlap with those
previously identified as being differentially methylated among primates. These included genes within the
HOXD cluster which have been found to be differentially methylated among modern and ancient
hominins (Gokhman et al., 2014). Specifically, 1 baboon-specific DMP is hypermethylated in *HOXD9*,
and 2 chimpanzee-specific DMPs are hypomethylated in *HOXD9* (File S4). Additional genes in the
HOXD cluster (*HOXD8* and *HOXD10*) contain marmoset-specific DMPs, and genes such as *ARTN*,
COL2A1, and *GABBR1* which have been found to be differentially methylated among modern humans
and great apes (Hernando-Herraez et al., 2013), also contain marmoset-specific DMPs (File S4).
However, given the unusual nature of the marmoset data, these results will not be discussed further.
392
Inter-Specific DNA Methylation Profiling of HOXD10
Based on the inter-specific DNA methylation patterns identified in this study (Figure 5, Table
S13) and those identified in other evolutionary anthropological studies (Gokhman et al. 2014), the
HOXD10 gene was selected for subsequent DNA methylation profiling and analysis at a higher resolution
using gene-specific sequencing techniques. Regular and bisulfite sequences of several regions in the
HOXD10 gene were generated (Tables S14-S17). First, loci across the entire *HOXD10* gene were
examined from a subset of EPIC array samples – 3 baboons, 3 macaques, 3 vervets, 3 chimpanzees, and 3
marmosets (Figure 6, Tables S1 and S18, File S8-S9). Additionally, one test locus (cg02193236) and its
surrounding region in *HOXD10* were surveyed in all NHP samples in order to confirm the reliability of
the EPIC array in assessing DNA methylation levels (Figure S9, Tables S1 and S19, File S10-S11). This
test, which showed similar methylation patterns to those identified with the EPIC array, supplemented
previous methylation array replication studies that have found resulting data to be highly correlated in
replicate chimpanzee tissue samples (>98% correlated) (Pai et al., 2011) and in replicate macaque tissue
samples (97% correlated) (Ong et al., 2014). Following the alignment of these sequences to the
appropriate NHP references, the presence and absence of methylation across the *HOXD10* gene in each
animal was determined (Tables S18-S19, Files S8-S11). These data reveal that across the *HOXD10* gene,
NHPs display generally low methylation with some clustered increased amounts of methylation upstream
of the gene and at the start of the gene body (Figure 6). Additionally, in marmosets, the bisulfite sequence
findings are consistent with methylation patterns identified using the EPIC array, so these data are unable
to further clarify the unusual distribution of methylation levels discovered in these samples (Figure S4).
413
Discussion:
This study explored skeletal gene regulation and its relation to complex traits and species
differences by evaluating genome-wide and gene-specific DNA methylation patterns in trabecular bone
from several NHP species. These species are phylogenetically representative of the order and display
anatomical and locomotor differences. Specifically, while the sampled cercopithecoids (baboons,
macaques, vervets) display varying levels of terrestrial and arboreal quadrupedal locomotion, the sampled

ape species (chimpanzees) is a knuckle-walking quadruped that also vertical climbs and occasionally
walks bipedally (Cawthon Lang, 2006). Even more distinct are the sampled platyrrhines (marmosets)
which locomote via vertical clinging and leaping (Cawthon Lang, 2005). Lemurs and lorises are the only
major taxonomic primate groups not examined in this study. Regardless, the current sample set provided a
unique opportunity to examine skeletal epigenetic differences in relation to bone morphology at an
evolutionary scale. In this research we assume that adult bone methylation patterns reflect gene regulation
that enables the maintenance of post-maturation skeletal morphologies. Some of these patterns may be
impacted by environmental forces, like diet and biomechanical loading, but we attempted to control for
these variables by only including NHPs from captive colonies with similar environmental exposures.
Nevertheless, we are unable to ascertain how substantially the adult skeletal methylation in our sample set
deviates from the regulatory patterns that initially arose during early skeletal element development.
Despite these limitations, the novel data collected in this study can help to inform the relationship
between epigenetic variation and skeletal traits. First, we evaluated the association between intra-specific
variation in skeletal DNA methylation and intra-specific variation in femur morphology. Second, we
assessed inter-specific DNA methylation differences and the potential contribution of these lineage-
specific patterns to species-specific morphologies.
436
437   ***Intra-specific DMPs are not readily associated with morphological variants.***
438   With respect to intra-specific morphology, very few sites were found to be differentially
methylated. DMPs were only identified in association with baboon bicondylar femur length, baboon
maximum femur length, macaque proximal femur width, macaque medial condyle width, vervet superior
shaft width, vervet inferior shaft width, vervet anatomical neck height, and chimpanzee anatomical neck
length (Table S8). Additionally, most of these associations only identified one DMP. This limited number
of associations may be due to the small amount of variation identified in each morphology. This point is
supported in that almost all the morphologies with methylation associations also have the highest intra-
specific variation in size (Figure S5, Files S2-S3). Further, because the animals included in this study
were born and raised in captivity, their limited exposure to environmental variation may limit both the
range of phenotypic variation and methylation variation present within species. Alternatively, the limited
number of associations may simply be due to the small sample sizes within each species. Increasing the
number of individuals included in this study would improve the power to detect morphology-related
DMPs. Indeed, for our largest NHP sample set (baboons: n=28, 189,858 CpG sites evaluated), there is
only 11% power to detect a 10% change in methylation ($\Delta\beta = 0.1$) between two groups given an FDR
cutoff of 0.05 (Graw, Henn, Thompson, & Koestler, 2019). Further, larger sample sizes might also reveal
larger amounts of variation within each morphology. Thus, investigations employing larger sample sets
will provide more insight on these fronts. Nevertheless, it may be the case that DNA methylation
variation does not have a large influence on nonpathological femur morphology within NHPs.
456   Of the few intra-specific methylation patterns associated with morphology, they likely have weak
functional effects. Research has shown that individual site-specific methylation changes are not readily
associated with differential gene expression (Bork et al., 2010; Chen et al., 2011; Koch et al., 2011).
Rather, differential gene expression is made possible through the accumulation of several methylation
changes within promotor regions (Suzuki & Bird, 2008) or across the gene body (Singer, Kosti, Pachter,
& Mandel-Gutfreund, 2015). Detecting accumulations of methylation changes across genes should have

been possible in the current study as 4-19 probes targeted each gene on average (Table S4). Thus, the
individual sites identified in this study are not due to experimental limitations and likely have limited
impacts on gene regulation. This is further supported by the degree of methylation variation observed
among these DMPs. Sites that have an average change in mean methylation less than 10% ($\Delta\beta < 0.1$) are
thought to have little biological relevance (Hernando-Herraez et al., 2013). Thus, those DMPs with small
changes in methylation likely have little to no biological function. Further, some of the DMPs with $\Delta\beta \geq$
0.1 appear to be highly influenced by small subsets of the sample sets. For example, in the case of
macaque proximal width, the majority of methylation change is due to two individuals (1 female and 1
male) that have extremely low methylation at cg19349877 as compared to other macaques (File S3).
Additionally, in the case of chimpanzee anatomical neck length, the 273 DMPs identified are largely
affected by one individual that has a longer anatomical neck than other chimpanzees. Lastly, the limited
enrichment of GO biological processes and KEGG pathway functions among detected DMPs indicates a
lack of common function among these associated regions. Overall, these finding suggest that while some
DMPs appear to be associated with intra-specific morphological variation in NHPs, not enough evidence
is present to support them having a functional role in the development and maintenance of this
morphological variation.
478
Differential methylation is observed among NHP species.
With respect to inter-specific variation, several sites were found to have significant species-
specific methylation differences. Specifically, out of the 39,802 sites examined, 650 species-specific
DMPs were identified in baboons, 257 in macaques, 639 in vervets, 2,796 in chimpanzees, and 13,778 in
marmosets, and these span 7,320 genes (Tables S11-S12). These numbers of species-specific DMPs did
not substantially change when accounting for age scaled to life expectancy instead of exact age (File S4).
However, many of these DMPs had biologically insignificant changes in mean methylation. Thus, several
$\Delta\beta$ cutoff thresholds were considered – from a 10% change in mean methylation ($\Delta\beta \geq 0.1$) up to a 40%
change in mean methylation ($\Delta\beta \geq 0.4$) – which reduced the overall numbers of species-specific DMPs.
Regardless of $\Delta\beta$ cutoff threshold, more species-specific DMPs were found in the platyrrhine marmosets,
followed by the great ape chimpanzees, and lastly the cercopithecoid baboons, macaques, and vervets.
This trend in numbers of species-specific DMPs is expected given the known phylogenetic relationships
between primates, with cercopithecoids more closely related to apes and platyrrhines more distantly
related to both groups (Perelman et al., 2011; Rogers & Gibbs, 2014). However, the number of marmoset-
specific DMPs is substantially larger than those for other taxa. While this discrepancy may reflect
marmosets having more species-specific changes, aspects of the experimental design may also contribute
to it. Marmosets are the only platyrrhine included in this study, so although some marmoset-specific
DMPs are truly specific to marmosets, others may be specific to all platyrrhines. Comparably,
cercopithecoid-specific DMPs and catarrhine-specific DMPs may be cancelled out from this study, as
such changes would be shared between all cercopithecoids and catarrhines, respectively. Second, since
marmosets are the most phylogenetically distant from humans as compared to the other NHPs included in
this study (Perelman et al., 2011; Rogers & Gibbs, 2014), the probe filtering steps may have biased
downstream data in favor of finding significant results primarily in marmosets. Third, the marmoset data
itself has a slightly different normalized distribution, with more mean methylation levels of 50%, than
that for other NHPs (Figure S4), which may be due to the chimerism often observed in marmosets. Levels

of chimerism in marmoset skeletal tissues have not been examined. However, blood-derived tissues are

known to have high levels of chimerism (>50% chimeric), as compared to epithelial tissues which have

the lowest levels of all tissues examined thus far (12% chimeric) (Malukiewicz et al., 2015; Ross, French,

& Orti, 2007). Even if trabecular bone has low chimerism, its close proximity to blood cells in bone

marrow increases the chance of highly chimeric cell contamination. Additionally, although when assessed

computationally, the filtered array probes examined in this study appear to hybridize sufficiently for

marmosets (Figure S3), there may be other unknown biological or technical issues that impede proper

DNA methylation analyses from the EPIC array in marmosets and may have inflated the overall number

of marmoset-specific DMPs identified. Given the disproportionate number of marmoset-specific DMPs

identified and the unusual nature of these data, interpreting the biological relevance of these data requires

caution.

The number of species-specific DMPs identified in this study, is comparable to those identified in

a previous study that assessed methylation patterns in blood from chimpanzees, bonobos, gorillas, and

orangutans using the 450K array and similar alignment criteria filtering methods with a focus on sites that

had a $\Delta\beta \geq 0.1$ (Hernando-Herraez et al., 2013). This research used a final set of 99,919 probes that were

shared across all great ape species and covered 12,593 genes with at least 2 probes per gene. Out of these,

2,284 species-specific DMPs were found in humans, 1,245 in chimpanzees and bonobos, 1,374 in

gorillas, and 5,501 in orangutans (Hernando-Herraez et al., 2013). In the current study, when a $\Delta\beta \geq 0.1$

threshold was used, only 1,572 chimpanzee-specific DMPs were identified (Table S11). This number is

lower than the number found in previous research comparing chimpanzees to other great apes (Hernando-

Herraez et al., 2013), species that are evolutionarily closer to chimpanzees than are cercopithecoids and

platyrrhines (Perelman et al., 2011; Rogers & Gibbs, 2014). However, the total number of sites examined

in the present study is approximately one-third of that examined in the prior great ape study. Thus, a 3-

fold increase in the number of sites examined might identify a 3-fold increase in chimpanzee-specific

DMPs as compared to cercopithecoids and platyrrhines, which would be closer to expectations

(Hernando-Herraez et al., 2013).

Additionally, the number of DMPs that distinguish species from one another in the current study

is substantially smaller than the numbers of DMPs identified between different skeletal tissue types,

between different age cohorts, and between individuals with different skeletal disease states within a NHP

species (Housman et al., 2018). This finding is to be expected since differences in DNA methylation

which regulate gene expression (Singer et al., 2015; Suzuki & Bird, 2008) should increase or decrease

with the degree of differences between cellular functions among comparative groups (Zhang et al., 2013).

In the case of different tissues, substantial DNA methylation differences likely promote distinct gene

regulation that is necessary for cells in different tissues to promote different functions. In the case of

different age cohorts, slightly fewer DNA methylation differences within the same skeletal tissue may

allow cells to emphasize efforts on growth and development in juveniles, as compared to maintenance in

adults, without altering the general skeletal-related functions of this tissue. In the case of osteoarthritic

disease states, even fewer regulatory changes may be needed to initiate the dysregulation of tissue

function. Finally, the comparatively small number of DNA methylation differences between adult,

skeletally healthy, NHP species seems reasonable given that other studies have also noted the presence of

more regulatory variation within species than between species (Uebbing et al., 2016). In summary,

epigenetic and regulatory differences, which control and enable age-dependent organ functions, should be

greater within a species when comparing between tissue types, age cohorts, and disease states, than when
comparing between species.
548    When evaluating how well the methylation patterns at species-specific DMPs cluster samples, a
$\Delta\beta \geq 0.4$ threshold is necessary to achieve clustering into distinct species (Figure 3). Less stringent $\Delta\beta$
thresholds are able to separate apes, platyrrhines, and cercopithecoids into distinct groups, but the
cercopithecoids do not form monophyletic species groups at these thresholds (Figure S6). Global changes
in methylation show similar phylogenetic patterns. While average species methylation patterns reveal a
well-supported tree topology that reflects known phylogenetic relationships between taxa (Perelman et al.,
2011; Rogers & Gibbs, 2014) (Figure 4), global changes in methylation among individual animals do not
distinguish cercopithecoids into distinct monophyletic groups (Figure S7). As before, parsing down these
global changes to only species-specific DMPs with $\Delta\beta \geq 0.4$ fixes the phylogeny (Figure S8). In previous
studies, global methylation changes were able to separate great apes into species-specific phylogenetic
groups (Hernando-Herraez et al., 2013). Conversely, the need for a higher $\Delta\beta$ cutoff threshold to
distinguish species in the current study may be due to evolutionary reasons.
560    The divergence times between cercopithecoids (baboons, macaques, and vervets) are comparable
to those between great apes (humans, chimpanzees, gorillas, and orangutans) (Perelman et al., 2011).
Thus, divergence times do not explain why global changes in methylation are unable to resolve species-
specific phylogenetic clades in the current study as compared to previous research. On the other hand,
while nonhuman great apes have experienced higher rates of molecular evolution as compared to humans
(Elango, Thomas, NISC Comparative Sequencing Program, & Yi, 2006), baboons and macaques have
slow rates of molecular evolution as compared to other cercopithecoids (Elango, Lee, Peng, Loh, & Yi,
2009), which may correspond to slower rates of epigenetic evolution. This might make baboons and
macaques appear more similar to vervets than expected, and further, this may make resolving the
phylogenetic divergences between cercopithecoids more difficult than that between the great apes.
Additionally, the number of sites included in the present study (39,802) as compared to previous studies
(99,919) (Hernando-Herraez et al., 2013) may limit the ability of the present study to fully resolve
species-specific lineages. This is reinforced by the low support of several branches in the phylogeny
based on global changes in methylation across individual animals (Figure S7). Conversely, sample size
was likely not a contributing factor to the discrepancies between the current study (n=58) and previous
studies (n=32) (Hernando-Herraez et al., 2013), as the current study has a slightly larger sample size.
However, the number of individuals per species was more uniform in previous studies (Hernando-Herraez
et al., 2013) than in the current study.
578    Alternatively, the fact that global changes in methylation are unable to fully resolve
cercopithecoid species-specific phylogenetic clades in the current study, may instead indicate that not
enough time has passed for cercopithecoid species to evolve fixed epigenetic changes between taxa in this
tissue. Additionally, it is possible that epigenetic variation at many of the sites examined in this study are
under balancing selection in cercopithecoids which prevents these markers from accurately resolving the
evolutionary divergences between these species. Previous research of gene regulation differences between
species has found that some deviations in gene expression may be under directional or balancing selection
(Romero, Ruvinsky, & Gilad, 2012; Whitehead & Crawford, 2006). Given the relatively low variance in
methylation across species in combination with lineage-specific changes in mean methylation at the
species-specific DMPs identified in this study, it is possible that these loci are experiencing similar

selective pressures (Romero et al., 2012). Nevertheless, most inter-specific regulatory differences appear
to be under stabilizing selection or neutral evolution (Brawand et al., 2011; Gilad, 2012; Romero et al.,
2012), and this cannot be ruled out in the present study.
591
Inter-specific DMPs are found in genes enriched for functions associated with skeletal traits.
The evolution of methylation changes along specific NHP lineages is associated with several
functions that may contribute to species-specific phenotypic differences (Files S6-S7). First, several
skeletal tissue functions are enriched in species-specific DMPs. Among almost all NHPs, cellular
adhesion functions are highly enriched. Cellular adhesion is necessary for cells to attach to other cells or
extracellular matrix, which is a necessity for bone cells (Mbalaviele, Shin, & Civitelli, 2006).
Additionally, in baboons, vervets, chimpanzees, and marmosets, anatomical developmental processes are
enriched, and in chimpanzees and marmosets, pattern specification processes, limb development, and
skeletal system development are enriched. Lastly, in marmosets, specific skeletal functions, such as
osteoblast differentiation and ossification are also enriched. Overall, these functions validate that most
patterns of differential methylation relate to skeletal tissue function, regulation, development, and
maintenance, as well as to larger anatomical developmental processes. Additionally, functions not
specific to the skeletal system were identified. In chimpanzees and marmosets, transcription and gene
expression regulatory functions were enriched. Further, in marmosets, functions related to the
development of skeletal muscles, nerves, the brain, the heart, blood vessels, kidneys, eyes, and ears were
also enriched. All together, these findings suggest that many species-specific changes in methylation may
contribute to the regulation of complex phenotypic changes. While this relationship was not observed for
intra-specific skeletal morphology variation, other skeletal traits not examined in this study may be
related. Nevertheless, many of the genes associated with the described functions only contain an average
of 1-2 differentially methylated sites. As described above, individual site-specific methylation changes are
not readily associated with differential gene expression (Bork et al., 2010; Chen et al., 2011; Koch et al.,
2011). Therefore, the enriched functions identified are likely not true biological effects due to methylation
differences on their own. Rather they hint at biological effects that may be the result of the combined
effects of several genetic, epigenetic, and other regulatory processes.
Finally, some of the genes containing species-specific DMPs overlap with those previously
identified as being differentially methylated in other tissues among primates. Specifically, the *HOXD*
cluster which is involved in limb development shows differential methylation among species. In the
current study, *HOXD9* shows species-specific hypermethylation at 1 site in baboons and species-specific
hypomethylation at 2 sites in chimpanzees, while in previous work, *HOXD9* shows hypermethylation in
archaic hominins as compared to modern humans (Gokhman et al., 2014). In the *HOXD* cluster, *HOXD10*
also contains interesting inter-specific DNA methylation patterns (Figure 5, Table S13), has an active role
in anatomical development, and has been found to be differentially methylated among hominins
(Gokhman et al. 2014). Thus, it was selected for subsequent DNA methylation profiling and analysis at a
higher resolution using gene-specific sequencing techniques.
HOXD10 specifically codes for a protein that functions as a sequence-specific transcription factor
which is expressed in the developing limb buds and is involved in differentiation and limb development.
In the current study, each NHP shows low to intermediate methylation levels across the gene body
(Figure 5, Table S13). A similar pattern is observed in the gene-specific methylation data, which further

reveals that *HOXD10* is not highly methylated in NHPs. However, across all taxa, some clusters of
hypermethylation are found upstream of the gene and at the start of the gene body, with marmosets on
average displaying more methylation in the gene body than other taxa (Figure 6, Table S18). The
consistency between bisulfite sequence and EPIC array data in marmosets does not clarify the unusual
nature of the marmoset EPIC array data (Figure S4), so more data are needed in these samples to better
characterize marmoset *HOXD10* methylation patterns. In the *HOXD10* gene body of hominins, humans
display hypomethylation, Neandertals displayed intermediate methylation levels, and Denisovans
displayed high levels of methylation (Gokhman et al., 2014). The variation of methylation patterns in this
gene body suggest that intermediate methylation levels may be a more ancestral epigenetic state for this
region in the primate lineage, while the extreme hypermethylation of this region in Denisovans and the
extreme hypomethylation of this region in humans may be derived epigenetic states. Previous work has
proposed that methylation differences in *HOXD10* may be associated with phenotypic distinctions
between modern human and archaic hominin limbs (Gokhman et al., 2014). While the current study did
not find substantial associations between methylation variation and aspects of femur morphology within
NHP species, further work to understand the role of differential methylation of *HOXD10* in promoting
morphological changes of the limb should be explored.
646
Conclusions and future directions.
In conclusion, while only a few significant associations were identified between methylation and
femur morphologies, several significant differences in methylation were observed inter-specifically.
Moreover, these species-specific DMPs were found in genes enriched for functions associated with
complex skeletal traits. This is the first study to characterize DNA methylation patterns in skeletal tissues
from a taxonomically diverse set of NHPs, and it is the first study to directly compare these patterns to the
nonpathological morphologies of the skeletal elements from which the tissues were derived. This design
enabled an initial exploration of skeletal gene regulation and its relation to complex traits and species
differences for which little else is currently known.
Ultimately, this work forms a foundation for future explorations of gene regulation and skeletal
trait evolution in primates. The dataset collected and investigated here provides an initial sampling of
skeletal methylation patterns in NHPs, and future primate skeletal epigenetics investigations can add to
these data at population- and evolutionary-scales. At a population-level, increasing the number of
individuals from one species can provide further insight into how genetic variation interacts with gene
regulation to contribute to a wider variety of healthy and pathological skeletal traits. Additionally,
sampling individuals of different ages would provide further insight into how gene regulation across
development contributes to skeletal anatomy. Adult bone methylation patterns are limited in their ability
to fully inform how gene regulation contributes to the differential development of skeletal morphologies.
In the present study, we assume that regulatory patterns present in adult skeletal tissues are involved in (or
at least not inhibiting) the maintenance of adult skeletal morphology, and that the regulatory patterns that
enable the maintenance of adult skeletal traits reflect those that enabled the development of those traits.
Future work in the field of primate skeletal epigenetics could explicitly test the degree to which adult
bone methylation patterns reflect patterns that arise during early skeletal element development. Lastly, at
an evolutionary-level, increasing the number of represented species, both extant and extinct, will further

inform our understanding of gene regulation evolution in primates and how patterns of variation
contribute to skeletal phenotypes.
673

Acknowledgements:
This work was supported by the National Institutes of Health (P01HL028972 to Anthony G.
Comuzzie, P40OD010965 to Matthew J. Jorgensen, P40RR019963 to Jay Kaplan); the Leakey
Foundation (Research Grant for Doctoral Students to G.H.); the Wenner-Gren Foundation (Gr. 9310 to
G.H.); the Nacey Maggioncalda Foundation (James F. Nacey Fellowship to G.H.); the International
Primatological Society (to G.H.); Sigma Xi (Grant-in-Aid of Research to G.H.); the ASU Center for
Evolution and Medicine (Venture Fund to G.H.); the ASU Graduate Research and Support Program (to
G.H.). Additionally, this investigation used resources that were supported by the Southwest National
Primate Research Center grant P51OD011133 from the Office of Research Infrastructure Programs,
National Institutes of Health.
We thank Eric D. Johnson and members of the Department of Genetics at the Texas Biomedical
Research Institute, including Anthony G. Comuzzie, Anne Sheldrake, Jaydee Foster, Kara Peterson, Mel
Carless, and Laura Cox, for helpful discussions. We also thank Megann Phillips for assistance with PCR
primer design.
Newly reported data have been made available on NCBI's Gene Expression Omnibus and are
accessible through the GEO SuperSeries accession number GSE103332, which includes the following
SubSeries accession numbers: GSE103279 (intra-specific baboon DNA methylation data), GSE103271
(intra-specific macaque DNA methylation data), GSE103280 (intra-specific vervet DNA methylation
data), GSE94677 (intra-specific chimp DNA methylation data), GSE103328 (intra-specific marmoset
DNA methylation data), and GSE103287 (inter-specific DNA methylation data).

References:
Ankel-Simons, F. (2007). *Primate Anatomy* (3rd ed.). New York: Academic Press.
Aryee, M. J., Jaffe, A. E., Corrada-Bravo, H., Ladd-Acosta, C., Feinberg, A. P., Hansen, K. D., &
Irizarry, R. A. (2014). Minfi: A flexible and comprehensive Bioconductor package for the
analysis of Infinium DNA methylation microarrays. *Bioinformatics*, *30*(10), 1363–1369. doi:
10.1093/bioinformatics/btu049
Babbitt, C. C., Fedrigo, O., Pfefferle, A. D., Boyle, A. P., Horvath, J. E., Furey, T. S., & Wray, G. A.
(2010). Both Noncoding and Protein-Coding RNAs Contribute to Gene Expression Evolution in
the Primate Brain. *Genome Biology and Evolution*, *2*(0), 67–79. doi: 10.1093/gbe/evq002
Bell, C. G., Wilson, G. A., Butcher, L. M., Roos, C., Walter, L., & Beck, S. (2012). Human-specific CpG
"beacons" identify loci associated with human-specific traits and disease. *Epigenetics*, *7*(10),
1188–1199. doi: 10.4161/epi.22127
Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B
(Methodological)*, *57*(1), 289–300.
Blake, L. E., Thomas, S. M., Blischak, J. D., Hsiao, C. J., Chavarria, C., Myrthil, M., … Pavlovic, B. J.
(2017). A Comparative Study Of Endoderm Differentiation In Humans And Chimpanzees.
BioRxiv, 135442. doi: 10.1101/135442
Blekhman, R., Oshlack, A., Chabot, A. E., Smyth, G. K., & Gilad, Y. (2008). Gene Regulation in
Primates Evolves under Tissue-Specific Selection Pressures. *PLoS Genetics*, *4*(11), e1000271.
doi: 10.1371/journal.pgen.1000271

Bork, S., Pfister, S., Witt, H., Horn, P., Korn, B., Ho, A. D., & Wagner, W. (2010). DNA methylation
pattern changes upon long-term culture and aging of human mesenchymal stromal cells. *Aging*
Cell, *9*(1), 54–63. doi: 10.1111/j.1474-9726.2009.00535.x
Brawand, D., Soumillon, M., Necsulea, A., Julien, P., Csárdi, G., Harrigan, P., … Kaessmann, H. (2011).
The evolution of gene expression levels in mammalian organs. *Nature*, *478*(7369), 343–348. doi:
10.1038/nature10532
Cáceres, M., Lachuer, J., Zapala, M. A., Redmond, J. C., Kudo, L., Geschwind, D. H., … Barlow, C.
(2003). Elevated gene expression levels distinguish human from non-human primate brains.
Proceedings of the National Academy of Sciences, *100*(22), 13030–13035.
Cawthon Lang, K. A. (2005). Primate Factsheets: Common marmoset (Callithrix jacchus) Taxonomy,
Morphology, & Ecology. Retrieved January 13, 2019, from
http://pin.primate.wisc.edu/factsheets/entry/common_marmoset
Cawthon Lang, K. A. (2006). Primate Factsheets: Chimpanzee (Pan troglodytes) Taxonomy,
Morphology, & Ecology. Retrieved January 13, 2019, from
http://pin.primate.wisc.edu/factsheets/entry/chimpanzee
Chen, Y., Choufani, S., Ferreira, J. C., Grafodatskaya, D., Butcher, D. T., & Weksberg, R. (2011).
Sequence overlap between autosomal and sex-linked probes on the Illumina
HumanMethylation27 microarray. *Genomics*, *97*(4), 214–222. doi: 10.1016/j.ygeno.2010.12.004
Corner, B., Lele, S., & Richtsmeier, J. (1992). Measuring precision of three-dimensional landmark data.
Journal of Quantitative Anthropology, *3*, 347–359.
de Andrés, M. C., Kingham, E., Imagawa, K., Gonzalez, A., Roach, H. I., Wilson, D. I., & Oreffo, R. O.
C. (2013). Epigenetic Regulation during Fetal Femur Development: DNA Methylation Matters.
PLoS ONE, *8*(1), e54957. doi: 10.1371/journal.pone.0054957
Delgado-Calle, J., Fernández, A. F., Sainz, J., Zarrabeitia, M. T., Sañudo, C., García-Renedo, R., …
Riancho, J. A. (2013). Genome-wide profiling of bone reveals differentially methylated regions in
osteoporosis and osteoarthritis. *Arthritis & Rheumatism*, *65*(1), 197–205. doi: 10.1002/art.37753
den Hollander, W., Ramos, Y. F. M., Bos, S. D., Bomer, N., van der Breggen, R., Lakenberg, N., …
Meulenbelt, I. (2014). Knee and hip articular cartilage have distinct epigenomic landscapes:
Implications for future cartilage regeneration approaches. *Annals of the Rheumatic Diseases*,
73(12), 2208–2212. doi: 10.1136/annrheumdis-2014-205980
Elango, N., Lee, J., Peng, Z., Loh, Y.-H. E., & Yi, S. V. (2009). Evolutionary rate variation in Old World
monkeys. *Biology Letters*, *5*(3), 405–408. doi: 10.1098/rsbl.2008.0712
Elango, N., Thomas, J. W., NISC Comparative Sequencing Program, & Yi, S. V. (2006). Variable
molecular clocks in hominoids. *Proceedings of the National Academy of Sciences*, *103*(5), 1370–
1375. doi: 10.1073/pnas.0510716103
Enard, W., Fassbender, A., Model, F., Adorján, P., Pääbo, S., & Olek, A. (2004). Differences in DNA
methylation patterns between humans and chimpanzees. *Current Biology*, *14*(4), R148–R149.
Farcas, R., Schneider, E., Frauenknecht, K., Kondova, I., Bontrop, R., Bohl, J., … Haaf, T. (2009).
Differences in DNA Methylation Patterns and Expression of the CCRK Gene in Human and
Nonhuman Primate Cortices. *Molecular Biology and Evolution*, *26*(6), 1379–1389. doi:
10.1093/molbev/msp046
Flanagan, J. M., Popendikyte, V., Pozdniakovaite, N., Sobolev, M., Assadzadeh, A., Schumacher, A., …
Petronis, A. (2006). Intra- and Interindividual Epigenetic Variation in Human Germ Cells.
American Journal of Human Genetics, *79*(1), 67–84.
Fleagle, J. G. (1999). *Primate Adaptation and Evolution*. New York: Academic Press.
Fortin, J.-P., Triche, T., & Hansen, K. (2016). Preprocessing, normalization and integration of the
Illumina HumanMethylationEPIC array. *BioRxiv*, 065490. doi: 10.1101/065490
Fraga, M. F., Ballestar, E., Paz, M. F., Ropero, S., Setien, F., Ballestar, M. L., … Esteller, M. (2005).
Epigenetic differences arise during the lifetime of monozygotic twins. *Proceedings of the*
National Academy of Sciences of the United States of America, *102*(30), 10604–10609. doi:
10.1073/pnas.0500398102

Fukuda, K., Ichiyanagi, K., Yamada, Y., Go, Y., Udono, T., Wada, S., … Sasaki, H. (2013). Regional
DNA methylation differences between humans and chimpanzees are associated with genetic
changes, transcriptional divergence and disease genes. *Journal of Human Genetics*, *58*(7), 446–
454. doi: 10.1038/jhg.2013.55

Gama-Sosa, M. A., Midgett, R. M., Slagel, V. A., Githens, S., Kuo, K. C., Gehrke, C. W., & Ehrlich, M.
(1983). Tissue-specific differences in DNA methylation in various mammals. *Biochimica et*
Biophysica Acta (BBA)-Gene Structure and Expression, *740*(2), 212–219.

Gao, F., Niu, Y., Sun, Y. E., Lu, H., Chen, Y., Li, S., … Tan, T. (2017). De novo DNA methylation
during monkey pre-implantation embryogenesis. *Cell Research*. doi: 10.1038/cr.2017.25

Geeleher, P., Hartnett, L., Egan, L. J., Golden, A., Raja Ali, R. A., & Seoighe, C. (2013). Gene-set
analysis is severely biased when applied to genome-wide methylation data. *Bioinformatics*
(Oxford, England), *29*(15), 1851–1857. doi: 10.1093/bioinformatics/btt311

Gilad, Y. (2012). Using Genomic Tools to Study Regulatory Evolution. *Methods in Molecular Biology*
(Clifton, N.J.), *856*, 335–361. doi: 10.1007/978-1-61779-585-5_14

Gokhman, D., Agranat-Tamir, L., Housman, G., Garcia-Perez, R., Nissim-Rafinia, M., Mallick, S., …
Carmel, L. (2017). Extensive Regulatory Changes in Genes Affecting Vocal and Facial Anatomy
Separate Modern from Archaic Humans. *BioRxiv*, 106955. doi: 10.1101/106955

Gokhman, D., Lavi, E., Prüfer, K., Fraga, M. F., Riancho, J. A., Kelso, J., … Carmel, L. (2014).
Reconstructing the DNA methylation maps of the Neandertal and the Denisovan. *Science (New*
York, N.Y.), *344*(6183), 523–527. doi: 10.1126/science.1250368

Gokhman, D., Meshorer, E., & Carmel, L. (2016). Epigenetics: It's Getting Old. Past Meets Future in
Paleoepigenetics. *Trends in Ecology & Evolution*, *31*(4), 290–300. doi:
10.1016/j.tree.2016.01.010

Graw, S., Henn, R., Thompson, J. A., & Koestler, D. C. (2019). pwrEWAS: A user-friendly tool for
comprehensive power estimation for epigenome wide association studies (EWAS). *BMC*
Bioinformatics, *20*(1), 218. doi: 10.1186/s12859-019-2804-7

Hernando-Herraez, I., Prado-Martinez, J., Garg, P., Fernandez-Callejo, M., Heyn, H., Hvilsom, C., …
Marques-Bonet, T. (2013). Dynamics of DNA Methylation in Recent Human and Great Ape
Evolution. *PLOS Genet*, *9*(9), e1003763. doi: 10.1371/journal.pgen.1003763

Heyn, H., Moran, S., Hernando-Herraez, I., Sayols, S., Gomez, A., Sandoval, J., … Esteller, M. (2013).
DNA methylation contributes to natural human variation. *Genome Research*, *23*(9), 1363–1372.
doi: 10.1101/gr.154187.112

Housman, G., Havill, L. M., Quillen, E. E., Comuzzie, A. G., & Stone, A. C. (2018). Assessment of DNA
Methylation Patterns in the Bone and Cartilage of a Nonhuman Primate Model of Osteoarthritis.
CARTILAGE. doi: 10.1177/1947603518759173

Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., … Morgan, M. (2015).
Orchestrating high-throughput genomic analysis with Bioconductor. *Nature Methods*, *12*(2), 115–
121. doi: 10.1038/nmeth.3252

Jaffe, A. E., & Irizarry, R. A. (2014). Accounting for cellular heterogeneity is critical in epigenome-wide
association studies. *Genome Biology*, *15*(2), R31. doi: 10.1186/gb-2014-15-2-r31

Jeffries, M. A., Donica, M., Baker, L. W., Stevenson, M. E., Annan, A. C., Humphrey, M. B., …
Sawalha, A. H. (2016). Genome-Wide DNA Methylation Study Identifies Significant Epigenomic
Changes in Osteoarthritic Subchondral Bone and Similarity to Overlying Cartilage. *Arthritis &*
Rheumatology, *68*(6), 1403–1414. doi: 10.1002/art.39555

Joganic, J. L., Willmore, K. E., Richtsmeier, J. T., Weiss, K. M., Mahaney, M. C., Rogers, J., &
Cheverud, J. M. (2017). Additive genetic variation in the craniofacial skeleton of baboons (genus
Papio) and its relationship to body and cranial size. *American Journal of Physical Anthropology*,
1–17. doi: 10.1002/ajpa.23349

Karere, G. M., Glenn, J. P., Birnbaum, S., Rainwater, D. L., Mahaney, M. C., VandeBerg, J. L., & Cox,
L. A. (2013). Identification of candidate genes encoding an LDL-C QTL in baboons. *Journal of*
Lipid Research, *54*(7), 1776–1785.

King, M. C., & Wilson, A. C. (1975). Evolution at two levels in humans and chimpanzees. *Science*,
188(4184), 107–116. doi: 10.1126/science.1090005
Koch, C. M., Suschek, C. V., Lin, Q., Bork, S., Goergens, M., Joussen, S., … Wagner, W. (2011).
Specific Age-Associated DNA Methylation Changes in Human Dermal Fibroblasts. *PLOS ONE*,
6(2), e16679. doi: 10.1371/journal.pone.0016679
Lea, A. J., Altmann, J., Alberts, S. C., & Tung, J. (2016). Resource base influences genome-wide DNA
methylation levels in wild baboons (Papio cynocephalus). *Molecular Ecology*, *25*(8), 1681–1696.
doi: 10.1111/mec.13436
Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E., & Storey, J. D. (2012). The sva package for
removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics, *28*(6), 882–883. doi: 10.1093/bioinformatics/bts034
Leek, J. T., & Storey, J. D. (2007). Capturing Heterogeneity in Gene Expression Studies by Surrogate
Variable Analysis. *PLoS Genetics*, *3*(9), e161. doi: 10.1371/journal.pgen.0030161
Leek, J. T., & Storey, J. D. (2008). A general framework for multiple testing dependence. *Proceedings of
the National Academy of Sciences*, *105*(48), 18718–18723. doi: 10.1073/pnas.0808709105
Lewton, K. L. (2017). The effects of captive versus wild rearing environments on long bone articular
surfaces in common chimpanzees (Pan troglodytes). *PeerJ*, *5*, e3668. doi: 10.7717/peerj.3668
Lewton, K. L., Ritzman, T., Copes, L. E., Garland, T., & Capellini, T. D. (2018). Exercise-induced
loading increases ilium cortical area in a selectively bred mouse model. *American Journal of
Physical Anthropology*, 1–9. doi: 10.1002/ajpa.23770
Lindskog, C., Kuhlwilm, M., Davierwala, A., Fu, N., Hegde, G., Uhlén, M., … Pontén, F. (2014).
Analysis of candidate genes for lineage-specific expression changes in humans and primates.
Journal of Proteome Research, *13*(8), 3596–3606.
Lönnstedt, I., & Speed, T. (2002). Replicated microarray data. *Statistica Sinica*, *12*(1), 31–46.
Madrid, A., Chopra, P., & Alisch, R. S. (2018). Species-Specific 5 mC and 5 hmC Genomic Landscapes
Indicate Epigenetic Contribution to Human Brain Evolution. *Frontiers in Molecular
Neuroscience*, *11*. doi: 10.3389/fnmol.2018.00039
Maksimovic, J., Phipson, B., & Oshlack, A. (2016). A cross-package Bioconductor workflow for
analysing methylation array data. *F1000Research*, *5*, 1281. doi: 10.12688/f1000research.8839.1
Malukiewicz, J., Boere, V., Fuzessy, L. F., Grativol, A. D., Silva, I. de O. e, Pereira, L. C. M., … Stone,
A. C. (2015). Natural and Anthropogenic Hybridization in Two Species of Eastern Brazilian
Marmosets (Callithrix jacchus and C. penicillata). *PLOS ONE*, *10*(6), e0127268. doi:
10.1371/journal.pone.0127268
Martin, D. I. K., Singer, M., Dhahbi, J., Mao, G., Zhang, L., Schroth, G. P., … Boffelli, D. (2011).
Phyloepigenomic comparison of great apes reveals a correlation between somatic and germline
methylation states. *Genome Research*, *21*(12), 2049–2057. doi: 10.1101/gr.122721.111
Mbalaviele, G., Shin, C. S., & Civitelli, R. (2006). Perspective: Cell–Cell Adhesion and Signaling
Through Cadherins: Connecting Bone Cells in Their Microenvironment. *Journal of Bone and
Mineral Research*, *21*(12), 1821–1827. doi: 10.1359/jbmr.060811
McCarthy, D. J., & Smyth, G. K. (2009). Testing significance relative to a fold-change threshold is a
TREAT. *Bioinformatics*, *25*(6), 765–771. doi: 10.1093/bioinformatics/btp053
McCartney, D. L., Walker, R. M., Morris, S. W., McIntosh, A. M., Porteous, D. J., & Evans, K. L.
(2016). Identification of polymorphic and off-target probe binding sites on the Illumina Infinium
MethylationEPIC BeadChip. *Genomics Data*, *9*, 22–24. doi: 10.1016/j.gdata.2016.05.012
McHenry, H. M., & Corruccini, R. S. (1978). The femur in early human evolution. *American Journal of
Physical Anthropology*, *49*(4), 473–487. doi: 10.1002/ajpa.1330490407
Mendizabal, I., Shi, L., Keller, T. E., Konopka, G., Preuss, T. M., Hsieh, T.-F., … Yi, S. V. (2016).
Comparative Methylome Analyses Identify Epigenetic Regulatory Loci of Human Brain
Evolution. *Molecular Biology and Evolution*, *33*(11), 2947–2959. doi: 10.1093/molbev/msw176

Moazedi-Fuerst, F. C., Hofner, M., Gruber, G., Weinhaeusel, A., Stradner, M. H., Angerer, H., …
Graninger, W. B. (2014). Epigenetic differences in human cartilage between mild and severe OA.
Journal of Orthopaedic Research, *32*(12), 1636–1645. doi: 10.1002/jor.22722
Molaro, A., Hodges, E., Fang, F., Song, Q., McCombie, W. R., Hannon, G. J., & Smith, A. D. (2011).
Sperm Methylation Profiles Reveal Features of Epigenetic Inheritance and Evolution in Primates.
Cell, *146*(6), 1029–1041. doi: 10.1016/j.cell.2011.08.016
Morris, J. A., Tsai, P.-C., Joehanes, R., Zheng, J., Trajanoska, K., Soerensen, M., … Bell, J. T. (2017).
Epigenome-wide association of DNA methylation in whole blood with bone mineral density.
Journal of Bone and Mineral Research, n/a-n/a. doi: 10.1002/jbmr.3148
Ong, M.-L., Tan, P. Y., MacIsaac, J. L., Mah, S. M., Buschdorf, J. P., Cheong, C. Y., … Holbrook, J. D.
(2014). Infinium Monkeys: Infinium 450K Array for the Cynomolgus macaque (*Macaca*
fascicularis). *G3: Genes|Genomes|Genetics*, *4*(7), 1227–1234. doi:
10.1534/g3.114.010967
Ostanek, B., Kranjc, T., Lovšin, N., Zupan, J., & Marc, J. (2018). Chapter 18—Epigenetic Mechanisms in
Osteoporosis. In A. Moskalev & A. M. Vaiserman (Eds.), *Epigenetics of Aging and Longevity*
(pp. 365–388). doi: 10.1016/B978-0-12-811060-7.00018-8
Pai, A. A., Bell, J. T., Marioni, J. C., Pritchard, J. K., & Gilad, Y. (2011). A Genome-Wide Study of DNA
Methylation Patterns and Gene Expression Levels in Multiple Human and Chimpanzee Tissues.
PLOS Genetics, *7*(2), e1001316. doi: 10.1371/journal.pgen.1001316
Palacios, D., & Puri, P. L. (2006). The epigenetic network regulating muscle development and
regeneration. *Journal of Cellular Physiology*, *207*(1), 1–11. doi: 10.1002/jcp.20489
Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics (Oxford, England)*, *20*(2), 289–290.
Pavlovic, B. J., Blake, L. E., Roux, J., Chavarria, C., & Gilad, Y. (2018). A Comparative Assessment of
Human and Chimpanzee iPSC-derived Cardiomyocytes with Primary Heart Tissues. *Scientific*
Reports, *8*(1), 1–14. doi: 10.1038/s41598-018-33478-9
Perelman, P., Johnson, W. E., Roos, C., Seuánez, H. N., Horvath, J. E., Moreira, M. A. M., … Pecon-
Slattery, J. (2011). A Molecular Phylogeny of Living Primates. *PLoS Genetics*, *7*(3), e1001342.
doi: 10.1371/journal.pgen.1001342
Petronis, A., Gottesman, I. I., Kan, P., Kennedy, J. L., Basile, V. S., Paterson, A. D., & Popendikyte, V.
(2003). Monozygotic twins exhibit numerous epigenetic differences: Clues to twin discordance?
Schizophrenia Bulletin, *29*(1), 169–178.
Phipson, B., Lee, S., Majewski, I. J., Alexander, W. S., & Smyth, G. K. (2016). Robust hyperparameter
estimation protects against hypervariable genes and improves power to detect differential
expression. *The Annals of Applied Statistics*, *10*(2), 946–963. doi: 10.1214/16-AOAS920
Phipson, B., Maksimovic, J., & Oshlack, A. (2016). missMethyl: An R package for analyzing data from
Illumina's HumanMethylation450 platform. *Bioinformatics*, *32*(2), 286–288. doi:
10.1093/bioinformatics/btv560
Provencal, N., Suderman, M. J., Guillemin, C., Massart, R., Ruggiero, A., Wang, D., … Szyf, M. (2012).
The Signature of Maternal Rearing in the Methylome in Rhesus Macaque Prefrontal Cortex and T
Cells. *Journal of Neuroscience*, *32*(44), 15626–15642. doi: 10.1523/JNEUROSCI.1470-12.2012
Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). Limma powers
differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
Research, gkv007. doi: 10.1093/nar/gkv007
Ritzman, T. B., Banovich, N., Buss, K. P., Guida, J., Rubel, M. A., Pinney, J., … Stone, A. C. (2017).
Facing the facts: The Runx2 gene is associated with variation in facial morphology in primates.
Journal of Human Evolution, *111*, 139–151. doi: 10.1016/j.jhevol.2017.06.014
Rogers, J., & Gibbs, R. A. (2014). Comparative primate genomics: Emerging patterns of genome content
and dynamics. *Nature Reviews Genetics*, *15*(5), 347–359. doi: 10.1038/nrg3707
Romero, I. G., Ruvinsky, I., & Gilad, Y. (2012). Comparative studies of gene expression and the
evolution of gene regulation. *Nature Reviews Genetics*, *13*(7), 505–516. doi: 10.1038/nrg3229

Ross, C. N., French, J. A., & Orti, G. (2007). Germ-line chimerism and paternal care in marmosets
(Callithrix kuhlii). *Proceedings of the National Academy of Sciences*, *104*(15), 6278–6282. doi:
10.1073/pnas.0607426104
Rowe, N. (1996). *The Pictorial Guide to Living Primates*. East Hampton, New York: Pogonias Press.
Rushton, M. D., Reynard, L. N., Barter, M. J., Refaie, R., Rankin, K. S., Young, D. A., & Loughlin, J.
(2014). Characterization of the Cartilage DNA Methylome in Knee and Hip Osteoarthritis:
Methylation Profile of OA Cartilage. *Arthritis & Rheumatology*, *66*(9), 2450–2460. doi:
10.1002/art.38713
Sharif, J., Endo, T. A., Toyoda, T., & Koseki, H. (2010). Divergence of CpG island promoters: A
consequence or cause of evolution? *Development, Growth & Differentiation*, *52*(6), 545–554. doi:
10.1111/j.1440-169X.2010.01193.x
Shelnutt, K. P., Kauwell, G. P. A., Gregory III, J. F., Maneval, D. R., Quinlivan, E. P., Theriaque, D. W.,
… Bailey, L. B. (2004). Methylenetetrahydrofolate reductase 677C→T polymorphism affects
DNA methylation in response to controlled folate intake in young women. *The Journal of
Nutritional Biochemistry*, *15*(9), 554–560. doi: 10.1016/j.jnutbio.2004.04.003
Simon, T. C., & Jeffries, M. A. (2017). The Epigenomic Landscape in Osteoarthritis. *Current
Rheumatology Reports*, *19*(6), 30. doi: 10.1007/s11926-017-0661-9
Singer, M., Kosti, I., Pachter, L., & Mandel-Gutfreund, Y. (2015). A diverse epigenetic landscape at
human exons with implication for expression. *Nucleic Acids Research*, gkv153. doi:
10.1093/nar/gkv153
Slieker, R. C., Bos, S. D., Goeman, J. J., Bovée, J. V., Talens, R. P., van der Breggen, R., … Heijmans, B.
T. (2013). Identification and systematic annotation of tissue-specific differentially methylated
regions using the Illumina 450k array. *Epigenetics & Chromatin*, *6*, 26. doi: 10.1186/1756-8935-
6-26
Smith, R. W. A., Monroe, C., & Bolnick, D. A. (2015). Detection of Cytosine Methylation in Ancient
DNA from Five Native American Populations Using Bisulfite Sequencing. *PLoS ONE*, *10*(5).
doi: 10.1371/journal.pone.0125344
Smyth, G. K. (2004). Linear models and empirical bayes methods for assessing differential expression in
microarray experiments. *Statistical Applications in Genetics and Molecular Biology*, *3*, Article3.
doi: 10.2202/1544-6115.1027
Suzuki, M. M., & Bird, A. (2008). DNA methylation landscapes: Provocative insights from epigenomics.
Nature Reviews Genetics, *9*(6), 465–476. doi: 10.1038/nrg2341
Terzidis, I., Totlis, T., Papathanasiou, E., Sideridis, A., Vlasis, K., & Natsis, K. (2012). Gender and Side-
to-Side Differences of Femoral Condyles Morphology: Osteometric Data from 360 Caucasian
Dried Femori. *Anatomy Research International*, *2012*, e679658. doi: 10.1155/2012/679658
Tung, J., Zhou, X., Alberts, S. C., Stephens, M., & Gilad, Y. (2015). The genetic architecture of gene
expression levels in wild baboons. *ELife*, *4*, e04729. doi: 10.7554/eLife.04729
Uebbing, S., Künstner, A., Mäkinen, H., Backström, N., Bolivar, P., Burri, R., … Ellegren, H. (2016).
Divergence in gene expression within and between two closely related flycatcher species.
Molecular Ecology, *25*(9), 2015–2028. doi: 10.1111/mec.13596
Vilgalys, T. P., Rogers, J., Jolly, C., Mukherjee, S., & Tung, J. (2018). *Evolution of DNA methylation in
baboons*. doi: 10.1101/400093
Weber, M., Hellmann, I., Stadler, M. B., Ramos, L., Pääbo, S., Rebhan, M., & Schübeler, D. (2007).
Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the
human genome. *Nature Genetics*, *39*(4), 457–466. doi: 10.1038/ng1990
White, T., & Folkens, P. (2000). *Human osteology* (2nd ed.). San Diego: Academic Press.
Whitehead, A., & Crawford, D. L. (2006). Neutral and adaptive variation in gene expression. *Proceedings
of the National Academy of Sciences of the United States of America*, *103*(14), 5425–5430. doi:
10.1073/pnas.0507648103
Young, M. D., Wakefield, M. J., Smyth, G. K., & Oshlack, A. (2010). Gene ontology analysis for RNA-
seq: Accounting for selection bias. *Genome Biology*, *11*, R14. doi: 10.1186/gb-2010-11-2-r14

Zeng, J., Konopka, G., Hunt, B. G., Preuss, T. M., Geschwind, D., & Yi, S. V. (2012). Divergent Whole-
Genome Methylation Maps of Human and Chimpanzee Brains Reveal Epigenetic Basis of
Human Regulatory Evolution. *The American Journal of Human Genetics*, *91*(3), 455–465. doi:
10.1016/j.ajhg.2012.07.024
Zhang, B., Zhou, Y., Lin, N., Lowdon, R. F., Hong, C., Nagarajan, R. P., … Wang, T. (2013). Functional
DNA methylation differences between tissues, cell types, and across individuals discovered using
the M&M algorithm. *Genome Research*, *23*(9), 1522–1540. doi: 10.1101/gr.156539.113
976

977   **Figure Legends:**

979
980   **Figure 1. Nonhuman Primate Sample Set Ages.** Boxes represent one standard deviation from the
981   average age, and whiskers depict the full range of ages for each species. Baboons (n=28) are 16.90±5.02
982   years, chimpanzees (n=4) are 11.31±1.87 years, macaques (n=10) are 14.75±2.65 years, marmosets (n=6)
983   are 3.34±1.41 years, and vervets (n=10) are 9.31±10.30 years. Additional details can be found in Table
984   S1.

986
987   **Figure 2. Nonhuman Primate Morphological Measurements.** See Table S6 for a detailed description
988   of these measurements.

Figure 3. Methylation Levels at Species-Specific DMPs with Δβ≥0.4 Identified in the Inter-Specific Study. Heatmap depicting the DNA methylation levels (β values) of all species-specific DMPs with average absolute Δβ values greater than 0.4 between each taxonomic group (x-axis) in all nonhuman primate samples (n=58). The sex and age of each nonhuman primate are also provided (y-axis). Red indicates higher methylation at a DMP, while blue indicates lower methylation at a DMP. The dendrogram of all samples (y-axis) clusters individuals based on the similarity of their methylation patterns. Samples cluster based on species-level taxonomic groupings and as predicted based on known species phylogenetic histories.

Figure 4. Phylogeny Based on Average Species-Level Global Changes in Methylation. Observed phylogenetic relationship among nonhuman primates when considering average species-level global changes in methylation. This tree was constructed using the methylation levels for all finalized 39,802 filtered probes. We averaged the β values per probe within a species, used Euclidean distances to calculate the difference between every two species, and estimated a neighbor joining tree using this

1005 distance matrix. For the resulting tree, 1000 bootstraps were performed to determine confidence values
1006 for each branch. The number provide at each node indicates the number of bootstrap replicates that
1007 support it out of the 1000 performed.

1008

1009
Figure 5. Genome-Wide Methylation Levels Across *HOXD10* in Nonhuman Primates. Plot of the
methylation levels of significant DMPs across the *HOXD10* gene (hg19 chr2:176981492-176984670).
Plot shows the average β values for each DMP with error bars indicating 1 standard deviation in each
direction for each comparative group (teal = baboon, orange = chimpanzee, purple = macaque, pink =
marmoset, and light green = vervet). DMP chromosomal position in relation to the *HOXD10* gene is also
depicted. This gene is of interest because it has been found to be differentially methylated in ancient and
modern hominin species (Gokhman et al., 2014). Of the sites depicted here, 5 DMPs were found to show
significant species-specific methylation in marmosets. Of the 5 species-specific DMPs in the *HOXD10*
gene of marmosets, 4 have Δβ between 0.2 and 0.3 (**) and 1 has a Δβ < 0.1 (*). See Table S13 for
additional information. In the *HOXD10* gene, the two exons are denoted with the thickest bars (exon 1:
hg19 chr2:176981561-176982306; exon 2: hg19 chr2:176983681-176983959), the UTRs are denoted
with bars of intermediate thickness (5'UTR: hg19 chr2:176981491-176981561; 3'UTR: hg19
chr2:176983959-176984670), and the one intron is denoted with the thinnest bar (intron 1: hg19
chr2:176982306-17698368).

Figure 6. Gene-Specific Methylation Levels Across *HOXD10* in Nonhuman Primates. Bar plot of DNA methylation across the *HOXD10* gene (hg19 chr2:176981492-176984670), as well as upstream and downstream several hundred bases (hg19 chr2:176980532-176985117). Bars depict the presence (tall bar), partial presence (medium bar), or absence (low bar) of methylation at human derived CpG sites in 15 nonhuman primate samples – 3 baboons, 3 macaques, 3 vervets, 3 chimpanzees, and 3 marmosets. While regular sequencing was very successful, bisulfite sequencing was less successful, with several sequence reads uninterpretable. As such, nonhuman primate methylation data is only available for a subset of the CpGs known in humans. Partial presence of methylation was called when sequencing fluorescence peaks for cytosine and thymine were both present at a particular site and one was at least half the size of the other. Overall, these data provide additional information regarding gene-specific methylation levels across *HOXD10*. CpG sites that were also targeted by the EPIC array are highlighted in yellow and include cg18115040 (chr2, position 176981328), cg25371634 (chr2, position 176981422), cg13217260 (chr2, position 176981469), cg03918304 (chr2, position 176981654), cg17489939 (chr2, position 176981919), cg26708100 (chr2, position 176983815), cg10393811 (chr2, position 176983927), cg08992581 (chr2, position 176983949), and cg06005169 (chr2, position 176984634). See Table S18 and Files S8-S9 for additional information.