Title:

Intra- and Inter-Specific Investigations of Skeletal DNA Methylation and Femur Morphology in Primates

Running Title:

Skeletal DNA Methylation Variation in Primates

Authors and Affiliations:

Genevieve Housman[1,2*], Ellen E. Quillen[3], and Anne C. Stone[1,2]

[1]School of Human Evolution and Social Change, Arizona State University, Tempe, AZ, USA.

[2]Center for Evolution and Medicine, Arizona State University, Tempe, AZ, USA.

[3]Department of Genetics, Texas Biomedical Research Institute, San Antonio, TX, USA.

*Corresponding author:

Genevieve Housman, Section of Genetic Medicine, University of Chicago, 920 East 58th Street, CLSC

317, Chicago, IL 60637, USA. Email: ghousman@uchicago.edu

Author Notes:

Genevieve Housman is currently affiliated with the University of Chicago, and Ellen E. Quillen is

currently affiliated with Wake Forest School of Medicine.

Abstract:

Objectives

Epigenetic mechanisms influence the development and maintenance of complex phenotypes and may also

contribute to the evolution of species-specific phenotypes. With respect to skeletal traits, little is known

about the gene regulation underlying these hard tissues or how tissue-specific patterns are associated with

bone morphology or vary among species. To begin exploring these topics, this study evaluates one

epigenetic mechanism, DNA methylation, in skeletal tissues from five nonhuman primate species which

display anatomical and locomotor differences representative of their phylogenetic groups.

Materials and Methods

First, we test whether intra-specific variation in skeletal DNA methylation is associated with intra-specific

variation in femur morphology. Second, we identify inter-specific differences in DNA methylation and

assess whether these lineage-specific patterns may have contributed to species-specific morphologies.

Specifically, we use the Illumina Infinium MethylationEPIC BeadChip to identify DNA methylation

patterns in femur trabecular bone from baboons (n=28), macaques (n=10), vervets (n=10), chimpanzees

(n=4), and marmosets (n=6).

Results

Significant differentially methylated positions (DMPs) were associated with a subset of morphological

variants, but these likely have small biological effects and may be confounded by other variables

associated with morphological variation. Conversely, several species-specific DMPs were identified, and
these are found in genes enriched for functions associated with complex skeletal traits.
45
46 *Discussion*
47 Overall, these findings reveal that while intra-specific epigenetic variation is not readily associated with
48 skeletal morphology differences, some inter-specific epigenetic differences in skeletal tissues exist and
49 may contribute to evolutionarily distinct phenotypes. This work forms a foundation for future
50 explorations of gene regulation and skeletal trait evolution in primates.
51
52 **Key Words:** DNA methylation, nonhuman primates, evolution, epigenome, bone
53
Introduction:
Primates distinguish themselves from other mammals with their unique suite of anatomical
features that initially enabled arboreal niche occupation and subsequently evolved to fit a myriad of
habitats and forms of locomotion, the most unique being hominin bipedalism. The resulting
morphological variation that has evolved across taxa has its foundation in underlying skeletal anatomies.
Such skeletal morphologies are used to both characterize extant primate diversity and reconstruct the
anatomy and locomotor capabilities of extinct primate species (Ankel-Simons, 2007; Fleagle, 1999; Leigh
& Shea, 1995; Schultz, 1930, 1937). The range of mechanisms that enable the development of skeletal
features are not entirely understood, though. Skeletal features related to varied body forms are often
described as the result of environmental adaptations. However, skeletal morphology is more accurately
defined as the result of complex processes, including environmental, genetic, and epigenetic mechanisms.
While environmental (Henriksen, Creaby, Lund, Juhl, & Christensen, 2014; Lewton, 2017; Lewton,
Ritzman, Copes, Garland, & Capellini, 2018; Macrini et al., 2013) and genetic (Joganic et al., 2017;
Ritzman et al., 2017; S. M. Smith, Garic, Berres, & Flentke, 2014) factors have been readily explored, the
important roles of epigenetic factors, such as DNA methylation, in skeletal tissue development and
maintenance have only recently been identified (Delgado-Calle et al., 2013; Goldring & Marcu, 2012;
García-Ibarbia et al., 2013; Liu et al., 2013; Loughlin & Reynard, 2015; Ostanek, Kranjc, Lovšin, Zupan,
& Marc, 2018; Ramos et al., 2014; Reynard, Bui, Syddall, & Loughlin, 2014; Reynard, 2017; Simon &
Jeffries, 2017).
For instance, epigenetic processes are influential in regulating skeletal muscle development
(Brand-Saberi, 2005; Ling et al., 2012; Palacios & Puri, 2006; Pandorf, Haddad, Wright, Bodell, &
Baldwin, 2009; Zwetsloot, Laye, & Booth, 2009) which can impact the adjacent skeletal scaffolding
system. Several genes involved in human skeletal development appear to be differentially methylated
across fetal and adult developmental stages (de Andrés et al., 2013). Lastly, methylation variation in
humans and model organisms has been implicated in several skeletal pathologies and disorders, such as
rheumatoid arthritis, osteoporosis, and osteoarthritis (Bovée, Hogendoorn, Wunder, & Alman, 2010;
Delgado-Calle et al., 2013; den Hollander et al., 2014; Dimitriou, Jones, McGonagle, & Giannoudis,
2011; Fernández-Tajes et al., 2014; García-Ibarbia et al., 2013; Goldring & Marcu, 2012; Iliopoulos,
Malizos, Oikonomou, & Tsezou, 2008; Jeffries et al., 2016, 2014; Kasaai, Gaumond, & Moffatt, 2013;
Liu et al., 2013; Loughlin & Reynard, 2015; Moazedi-Fuerst et al., 2014; Morris et al., 2017; Ralston &
Uitterlinden, 2010; Ramos et al., 2014; Reynard, 2017; Reynard et al., 2014; Rivadeneira et al., 2009;

Rushton et al., 2014). Some of these studies are the first to assess methylation patterns in human skeletal
tissues (Delgado-Calle et al., 2013; den Hollander et al., 2014; Fernández-Tajes et al., 2014; García-
Ibarbia et al., 2013; Jeffries et al., 2014; Moazedi-Fuerst et al., 2014; Rushton et al., 2014), and such steps
are crucial for identifying the relationship between epigenetic variation and skeletal phenotypic variation.

The contributions of epigenetics to primate phenotypic variation were first considered by King
and Wilson (1975), who proposed that anatomical and behavioral differences between humans and
chimpanzees were more likely "based on changes in the mechanisms controlling the expression of genes
than on sequence changes in proteins" (King & Wilson, 1975). Studies to understand methylation
variation across species began soon afterwards (Gama-Sosa et al., 1983). General changes to mammalian
epigenomes have been examined (Sharif, Endo, Toyoda, & Koseki, 2010), but most epigenetics work in
primates has focused on humans – how it varies across distinct tissues within individuals (Lister et al.,
2009; Slieker et al., 2013), across different individuals (Oates et al., 2006; Petronis et al., 2003; Weksberg
et al., 2002), across populations (Heyn et al., 2013; Rakyan et al., 2004), in relation to aging processes
(Fraga et al., 2005), and in relation to diet (Jacob et al., 1998; Rampersaud, Kauwell, Hutson, Cerda, &
Bailey, 2000; Shelnutt et al., 2004), as well as how it is inherited across generations (Flanagan et al.,
2006; Gibbs et al., 2010; Suter, Martin, & Ward, 2004; van Dongen et al., 2014). These studies have
identified interesting intra-specific methylation variation present in humans.

Similarly, epigenetic variation has been identified among primate species. Inter-specific variation
in epigenetic signatures was initially inferred from underlying genomic sequences (Bell et al., 2012;
Haygood, Fedrigo, Hanson, Yokoyama, & Wray, 2007; Prendergast et al., 2007). For instance, several
promoter CpG densities vary across primates. These likely relate to regulatory methylation differences
across species as primate CpG densities correlate with methylation levels (Weber et al., 2007).
Additionally, gene expression studies, which primarily focus on brain tissues (Babbitt et al., 2010;
Cáceres et al., 2003; Warner et al., 2009) and a small set of other soft tissues (Blekhman et al. 2008;
Karere et al. 2010; Karere et al. 2012; Karere et al. 2013; Tung et al. 2015), have also noted regulatory
differences across species. Methylation differences in brain tissues have evolved across primates and
contributed to resultant brain phenotypes and disease vulnerabilities (Enard et al., 2004; Farcas et al.,
2009; Kothapalli et al., 2007; Madrid, Chopra, & Alisch, 2018; Mendizabal et al., 2016; Provencal et al.,
2012; Zeng et al., 2012). Thus, methylation-phenotype relationships can be identified in primates. Primate
methylation patterns in blood cells and other soft tissues have also been studied, but not to the same
degree (Fukuda et al., 2013; Gao et al., 2017; Hernando-Herraez et al., 2013, p.; Lea, Altmann, Alberts, &
Tung, 2016; Lindskog et al., 2014; Martin et al., 2011; Molaro et al., 2011; Pai, Bell, Marioni, Pritchard,
& Gilad, 2011; Vilgalys, Rogers, Jolly, Mukherjee, & Tung, 2018).

Interestingly, two studies using soft tissues and blood identified differential methylation and
expression of genes essential for skeletal development (*RUNX1*, *RUNX3*, and *COL2A1*) between some
primates (Hernando-Herraez et al., 2013; Lindskog et al., 2014). Additionally, the emerging field of
ancient epigenetics, which reconstructs methylation patterns from ancient DNA degradation patterns in
hominin remains (R. W. A. Smith, Monroe, & Bolnick, 2015; Gokhman et al., 2014; Gokhman,
Meshorer, & Carmel, 2016; Gokhman et al., 2017), have found that other skeletal developmental genes
(*HOXD* complex) are differentially methylated among modern humans and ancient hominins (Gokhman
et al., 2014). These findings suggest that primates do exhibit distinct epigenetic patterns and that primate
skeletal epigenetics is an important area of research for uncovering developmental and evolutionary

patterns relevant for complex skeletal traits, such as anatomy, locomotion, and disease. Despite these
potential avenues of insight, studies of nonhuman primate (NHP) skeletal epigenetics are limited
(Housman, Havill, Quillen, Comuzzie, & Stone, 2018).
Overall, there are clear knowledge gaps in our understanding of NHP skeletal complexity in
relation to epigenetic variation and epigenetic differences between phylogenetically diverse NHP species.
The present study begins to remedy this by assessing how genome-wide and gene-specific DNA
methylation in primate skeletal tissues varies intra- and inter-specifically and in relation to femur form.
Specifically, for this study, we explored the evolution of the epigenome and its relation to
nonpathological skeletal traits by identifying DNA methylation patterns in femur trabecular bone from
baboons, macaques, vervets, chimpanzees, and marmosets and assessing intra- and inter-specific
methylation variation and its relation to morphology. With this study design, we aimed to assess whether
skeletal methylation variation within species is associated with skeletal morphology variation, to what
degree skeletal methylation varies among species, and whether species-specific changes in methylation
further inform our understanding of skeletal morphology evolution. Additionally, this dataset expands the
number of characterized skeletal methylation patterns in NHPs for future primate skeletal epigenetics
investigations.
143
144 **Materials and Methods:**
145
146 *Ethics Statement*
147 NHP tissue samples included were opportunistically collected at routine necropsy of these animals. No
148 animals were sacrificed for this study, and no living animals were used in this study. Chimpanzee tissues
149 were collected opportunistically during routine necropsy prior to the September 2015 implementation of
150 Fish and Wildlife Service rule 80 FR 34499.
151
NHP Samples
NHP samples come from captive colonies of chimpanzees (*Pan troglodytes*), baboons (*Papio*
spp.), rhesus macaques (*Macaca mulatta*), and marmosets (*Callithrix jacchus*) from the Southwest
National Primate Research Center in Texas, as well as vervets (*Chlorocebus aethiops*) from the Wake
Forest/UCLA Vervet Research Colony in North Carolina. Femora were opportunistically collected at
routine necropsy of these animals and stored in -20°C freezers at the Texas Biomedical Research Institute
after dissection. These preparation and storage conditions ensured the preservation of skeletal DNA
methylation patterns. Samples include baboons (n=28), macaques (n=10), vervets (n=10), chimpanzees
(n=4), and marmosets (n=6). Age ranges span adulthood for each species and are comparable between
each group (Figure 1, Table S1). Scaled ages were also calculated for each sample (Table S1) using
published life expectancies for each species (Rowe, 1996). Both sexes are represented (female: n=33,
male: n=24, unknown: n=1).
164
Assessment of Femur Morphologies
On the right femora of NHP samples, 29 linear morphology traits (Figure 2, Table S6) were
measured using calipers. These measurements characterize overall femur shape (McHenry & Corruccini,
1978; Terzidis et al., 2012). All measurements were collected by one researcher, and intra-observer error

for each measurement was determined by performing triplicate measurements on approximately 10% of
the samples in each species. These measurements were spaced throughout the entire data collection
period. Error was calculated as the mean absolute difference divided by the mean (Corner, Lele, &
Richtsmeier, 1992; White & Folkens, 2000). All measurements that were retained for downstream
analyses had errors of less than 5%, and the only measurement excluded was macaque intercondylar
notch depth (error = 6.62%) (File S2).
175
Genome-Wide DNA Methylation Profiling
Trabecular bone cores were obtained from the medial condyles on the right distal femora of each
NHP sample using a drill press that cored transversely through the condyle leaving the articular surface
preserved (Supplemental Text). Cortical bone was removed from cores using a Dremel, and the remaining
trabecular bone cores were pulverized into bone dust using a SPEX SamplePrep Freezer/Mill. DNA was
extracted from femoral trabecular bone using a phenol-chloroform protocol optimized for skeletal tissues
(Barnett and Larson 2012). Genome-wide DNA methylation was assessed using Illumina Infinium
MethylationEPIC microarrays (EPIC array) (Supplemental Text). The array data discussed in this
publication have been deposited in NCBI's Gene Expression Omnibus and are accessible through GEO
SuperSeries accession number GSE103332, which includes the SubSeries accession numbers
GSE103279, GSE103271, GSE103280, GSE94677, GSE103328, and GSE103287.
187
Methylation Data Processing
Using previously described methods (Housman et al., 2018), raw EPIC array data were
normalized and converted to β values which represent the average methylation levels at each site (0 =
completely unmethylated sites, 1 = fully methylated sites), and M values which are the log transformed
ratio of methylated signal to unmethylated signal. Probes with failed detection levels (p-value > 0.05) in
greater than 10% of samples and samples with greater than 30% of probes with detection level failures
were removed from downstream analyses. Using previously described methods (Housman et al., 2018),
probes with sequence mismatches to NHP genomes, which could produce biased methylation
measurements, were computationally filtered out and excluded from downstream analyses (Supplemental
Text, Figures S1-S3, Tables S2-S5, File S1). Additionally, cross-reactive probes (McCartney et al., 2016),
probes containing SNPs at the CpG site, probes detecting SNP information, probes detecting methylation
at non-CpG sites, and probes targeting sites within the sex chromosomes were removed (Aryee et al.,
2014; Fortin, Triche, & Hansen, 2016). Lastly, we assessed whether NHP SNPs overlapped with targeted
probes (Supplemental Text, Table S2, File S1). The resulting finalized datasets (Figure S4) were then
used in further statistical analyses.
203
Statistical Analysis of Differential Methylation
In order to identify sites that were significantly differentially methylated across comparative
groups, we designed and tested general linear models (GLMs) which related the variables of interest
(morphological measures and species membership) to the DNA methylation patterns for each site, while
accounting for the effects of additional variables, batch effects, and latent variables (Maksimovic,
Phipson, & Oshlack, 2016). Sites found to have significant associations were classified as significant
differentially methylated positions (DMPs). Overall, GLMs were used to estimate differences in

methylation levels associated with femur morphologies within each taxonomic group (intra-specific) and
between each taxonomic group (inter-specific). Specific details about each set of GLMs are provided in
the following sections with additional details in the Supplemental Text.
214
Intra-Specific Analyses
For the intra-specific analyses, variables included in each GLM were the femur morphologies
within each taxonomic group, sex, age, and steady state weight when known, as well as unknown latent
variables calculated using the iteratively re-weighted least squares approach in the sva package in R (Jaffe
& Irizarry, 2014; Leek, Johnson, Parker, Jaffe, & Storey, 2012; Leek & Storey, 2007, 2008). Latent
variables estimated for each morphology were included to help mitigate any unknown batch and cell
heterogeneity effects on methylation variation at each site (Table S7). Each GLM design matrix was fit to
corresponding M value array data by generalized least squares using the limma package in R (Huber et
al., 2015; Phipson, Lee, Majewski, Alexander, & Smyth, 2016; Ritchie et al., 2015), and the estimated
coefficients and standard errors for each morphology were computed. Lastly, for each coefficient, an
empirical Bayes approach was applied using the limma package in R (Lönnstedt & Speed, 2002;
McCarthy & Smyth, 2009; Phipson, Lee, et al., 2016; Smyth, 2004) to compute moderated t-statistics,
log-odds ratios of differential methylation, and associated p-values adjusted for multiple testing
(Benjamini & Hochberg, 1995). Significant DMPs for the effect of each morphology were defined as
those having log fold changes in M values corresponding to an adjusted p-value of less than 0.05. Lastly,
the gene ontology (GO) and KEGG pathway enrichment for significant CpGs was determined using the
missMethyl package in R (Benjamini & Hochberg, 1995; Geeleher et al., 2013; Phipson, Maksimovic, &
Oshlack, 2016; Ritchie et al., 2015; Young, Wakefield, Smyth, & Oshlack, 2010), which takes into
account the differing number of probes per gene present on the array.
234
Inter-Specific Analyses
For the inter-specific analyses, variables included in the GLM were taxonomic grouping, sex,
age, known batch effects (e.g., array number and position), and unknown latent variables calculated using
the method described above. An additional GLM which replaced the age variable with ages scaled to
species-specific life expectancies (Rowe, 1996) was also tested. The six latent variables estimated were
included to help mitigate any unknown batch and cell heterogeneity effects on methylation variation at
each site. The GLM design matrix was fit to the M value array data using the method described above,
and the estimated coefficients and standard errors for taxonomic group effects were computed. As
described above, moderated t-statistics, log-odds ratios of differential methylation, and associated p-
values adjusted for multiple testing were computed, and significant DMPs for the effect of taxonomy
were defined as those having log fold changes in M values corresponding to an adjusted p-value of less
than 0.05.
To determine only those methylation differences that represent fixed changes between genera, we
used methods similar to those described in (Hernando-Herraez et al., 2013). Briefly, significant DMPs
were identified between all possible pairwise comparisons of taxa (n=10: baboon-macaque, baboon-
vervet, baboon-chimpanzee, baboon-marmoset, macaque-vervet, macaque-chimpanzee, macaque-
marmoset, vervet-chimpanzee, vervet-marmoset, chimpanzee-marmoset). A significant DMP was then
defined as taxon-specific if it was found to be significant in all four pairwise comparisons containing the

taxon of interest but not found in any of the remaining pairwise comparisons. The GO and KEGG
pathway enrichment of these DMPs was then determined as described above.
255   Additionally, global changes in methylation were calculated using distance matrices (Hernando-
Herraez et al., 2013) of the methylation levels for all finalized 39,802 filtered probes. These changes were
assessed at a species-level by averaging the β values per probe within each species. We then used
Euclidean distances to calculate the difference between every two species. Neighbor joining trees were
estimated from these distances using the ape package in R (Paradis, Claude, & Strimmer, 2004). For each
resulting tree, 1000 bootstraps were performed to determine confidence values for each branch. Global
changes in methylation were also assessed at the individual-level using Euclidean distances to calculate
the difference between every two individuals.
263
264   *DNA Methylation Profiling and Analyses of HOXD10*
265   Based on the inter-specific DNA methylation patterns identified in this study and those identified
in other evolutionary anthropological studies (Gokhman et al. 2014), the *HOXD10* gene was selected for
subsequent DNA methylation profiling and analysis at a higher resolution using gene-specific sequencing
techniques. Specifically, primers were designed and optimized to PCR amplify regions spanning across
the entire *HOXD10* gene, as well as upstream and downstream several hundred bases (hg19
chr2:176980532-176985117), in each NHP species for regular and bisulfite treated DNA (Tables S14-
S17). All gene-specific assays were performed in a subset of the samples tested using the EPIC array and
included chimpanzees (n=3), baboons (n=3), macaques (n=3), vervets (n=3), and marmosets (n=3) (Table
S1). Additionally, a subset of these assays that targeted the region around one test locus (cg02193236)
were performed in all of the samples (Table S1). As described above, DNA was extracted from femoral
trabecular bone using a phenol-chloroform protocol optimized for skeletal tissues (Barnett and Larson
2012). DNA was bisulfite converted using the EZ DNA MethylationTM Gold Kit according to the
manufacturer's instructions (Zymo Research). Successful PCR amplification was confirmed using gel
electrophoresis. Gene-specific PCR products were then purified using an exonuclease I and shrimp
alkaline phosphatase protocol and sequenced on the Applied Biosystems 3730 capillary sequencer at the
DNA Laboratory at Arizona State University.
281   Regular and bisulfite sequences were aligned to the appropriate NHP references within the
Enredo-Pecan-Orthus (EPO) whole-genome multiple alignments of several primate genomes [Ensembl
Compara.8_primates_EPO] (Paten, Herrero, Beal, et al. 2008; Paten, Herrero, Fitzgerald, et al. 2008)
using MEGA7 (Kumar et al. 2016). Manual annotation of these sequences within each sample confirmed
that the gene sequences belong to the appropriate primate species and that the regular and bisulfite treated
sequences only differ in cytosine composition. The number and distribution of methylated loci throughout
the *HOXD10* gene were then identified and compared within and among species to provide a higher
resolution of methylation variation within this targeted gene. Lastly, with respect to the test locus
(cg02193236) evaluated in all samples, methylation levels across this region were compared with
corresponding methylation levels of this region as determined using the EPIC array.
291
292   **Results:**
293   The aim of this study was to identify DNA methylation patterns in skeletal tissues from several
NHP species in order to determine how skeletal methylation varies at different taxonomic scales and in

relation to complex skeletal traits. We evaluated DNA methylation patterns in femur trabecular bone from
five NHP species – baboons (n=28), macaques (n=10), vervets (n=10), chimpanzees (n=4), and
marmosets (n=6) (Figure 1, Table S1). We used the Illumina Infinium MethylationEPIC BeadChip (EPIC
array), which was determined to be effective at identifying DNA methylation patterns in NHP DNA
(Supplemental Text, Figures S1-S4, Tables S2-S5, File S1). Further, we selected probes appropriate for
intra- and inter-specific comparison, which removed the effect of sequence differences among individuals
and species as a reason for methylation differences. With these data, we first tested whether intra-specific
variation in skeletal DNA methylation is associated with intra-specific variation in femur morphology.
Second, we identified inter-specific differences in DNA methylation and assessed whether these lineage-
specific patterns may have contributed to species-specific morphologies.
305
Genome-Wide Intra-Specific Differential Methylation and Morphological Variation
Measurements of 29 linear morphology traits (Figure 2, Table S6) were collected from each NHP
right femur. All measurements had less than 5% error, except those for intercondylar notch depth in
macaques (Figure S5, File S2). Significant DMPs associated with each intra-specific linear morphology
were interrogated from 189,858 sites in in baboons, 190,898 sites in macaques, 191,639 sites in vervets,
576,804 sites in chimpanzees, and 68,709 sites in marmosets (Tables S7-S8). In baboons, 1 DMP was
hypomethylated with increasing bicondylar femur length and increasing maximum femur length. In
macaques, 1 DMP was hypermethylated with increasing proximal femur width, 1 DMP was
hypermethylated with increasing medial condyle width, and 6 DMPs were hypomethylated with
increasing medial condyle width. In vervets, 1 DMP was hypomethylated with increasing superior shaft
width, 2 DMPs were hypomethylated with increasing inferior shaft width, and 1 DMP was
hypermethylated with increasing anatomical neck height. In chimpanzees, 216 DMPs were
hypomethylated and 57 DMPs were hypermethylated with increasing anatomical neck length. Lastly, in
marmosets, no DMPs were associated with morphological variation (Table S9, File S3).
While the maximum absolute change in mean methylation ($\Delta\beta$) for most of these DMPs is greater
than 10% ($\Delta\beta = 0.1$), the change in methylation between the individual with the largest morphology
measurement and the individual with the smallest morphology measurement is less than 0.1 for several
DMPs (File S3). Tests for enrichment of gene ontology (GO) and KEGG pathway functions were done
for the 3 intra-specific morphologies that had more than 2 DMPs associated with them. However, no GO
biological processes were found to be enriched in DMPs associated with either macaque medial condyle
width or chimpanzee anatomical neck length. Additionally, KEGG pathway functions were only found to
be enriched in DMPs associated with chimpanzee anatomical neck length, and these pathways were
predominantly involved in immune system cell signaling and differentiation (Table S10).
329
Genome-Wide Inter-Specific Differential Methylation
To determine how methylation varies inter-specifically, differential methylation was interrogated
from the 39,802 probes that were shared among NHP species (Table S11, File S4). Species-specific
DMPs were determined by identifying DMPs that were significant in all 4 pairwise comparisons
containing the taxon of interest but not in any of the remaining pairwise comparisons. These methods
identified 650 species-specific DMPs in baboons, 257 in macaques, 639 in vervets, 2,796 in chimpanzees,
and 13,778 in marmosets (Table S11). Similar results were obtained using a model that accounted for

scaled age based on life expectancy instead of exact age (660 baboon-specific DMPs, 260 macaque-
specific DMPs, 651 vervet-specific DMPs, 2,720 chimpanzee-specific DMPs, and 13,689 marmoset-
specific DMPs) (File S4). Thus, only the species-specific DMPs identified using the model that accounted
for exact age were used in downstream analyses. Comparably, when evaluating methylation patterns that
distinguish more broad taxonomic groups, 2,655 DMPs were found to be specific to Old World monkeys
(OWMs), 1,869 were found to be ape-specific, and 14,985 were found to be specific to New World
monkeys (NWMs) (File S5). It should be noted that the ape and NWM categories only include one taxon
each – chimpanzee and marmoset, respectively. Similar results were obtained using a model that
accounted for scaled age based on life expectancy instead of exact age (2,660 OWM-specific DMPs,
1,797 ape-specific DMPs, and 14,963 NWM-specific DMPs) (File S5).
Species-specific DMPs spanned 7,320 genes with an average of 2.2±2.0 (s.d.) significant probes
per gene (Table S12). Over half of these genes contain just 1 significant species-specific DMP (n=3,816),
and a portion of these are due to only 1 EPIC array probe targeting each gene (n=1,788). The remaining
genes that contain at least 2 significant species-specific DMPs (n=3,503) have an average of 3.4±2.3 (s.d.)
significant probes per gene (range: 2-34). Additionally, these species-specific DMPs covered a range of
locations with respect to genes and CpG islands (Table S12), indicating that these species-specific
changes in methylation are distributed throughout the genome. Using various $\Delta\beta$ cutoff thresholds
decreases the final number of species-specific DMPs to varying degrees (Table S11, File S4). Counter to
these decreases in numbers, though, species-specific DMPs cover a range of locations with respect to
genes and CpG islands regardless of the $\Delta\beta$ cutoff threshold (Table S12).
Overall, across $\Delta\beta$ cutoff thresholds, more species-specific DMPs were found in the NWM
marmosets, followed by the great ape chimpanzees, and lastly the OWM baboons, macaques, and vervets.
Additionally, the proportions of hypermethylated and hypomethylated species-specific DMPs within each
taxon remain fairly constant across $\Delta\beta$ cutoff thresholds (Table S11). In baboons, macaques, vervets, and
chimpanzees, more than half of all species-specific DMPs show patterns of hypermethylation, and in
marmosets, more than half of all species-specific DMPs show patterns of hypomethylation. The only
disruption in these trends is in chimpanzees when no $\Delta\beta$ threshold is applied. Nevertheless, species-
specific DMPs with different $\Delta\beta$ cutoff thresholds do differ in their abilities to cluster animals into
taxonomic groups (Figure S6, File S4). For all thresholds, apes, OWMs, and NWMs cluster into distinct
groups. However, within OWMs, vervets only cluster into a distinct species group with a $\Delta\beta \geq 0.2$
threshold, and the baboon-macaque clade require a $\Delta\beta \geq 0.3$ threshold. Lastly, species-specific clustering
of baboons, macaques, and vervets only occurs with a $\Delta\beta \geq 0.4$ threshold (Figure 3).
Additionally, global changes in methylation across all 39,802 probes were evaluated. Average
global changes within each species reveal that apes, OWMs, and NWMs are phylogenetically distinct
from one another, and these divergences are well supported (Figure 4). Similarly, when phylogenetic
relationships are evaluated using the global changes in methylation of individual animals, distinct lineages
are again formed between apes, OWMs, and NWMs (Figure S7). However, within the OWM clade,
several poorly supported branches result in baboons, macaques, and vervets not forming distinct lineages.
Phylogenetic separation of these OWM species into distinct lineages is only possible when the
methylation changes considered are reduced to only include species-specific DMPs with a $\Delta\beta \geq 0.4$
threshold (Figure S8).

Species-specific DMPs also show associations with genes that have a wide array of GO biological
processes (File S6) and KEGG pathway functions (File S7). Cellular adhesion is a primary GO function
found to be highly enriched in species-specific DMPs from baboons, macaques, vervets, chimpanzees,
and marmosets. Species-specific DMPs for chimpanzees and marmosets are also enriched for genes
involved in the regulation of transcription and gene expression. Additionally, enrichment of genes
involved in anatomical developmental processes is found in baboons, vervets, chimpanzees, and
marmosets. Chimpanzees and marmosets show further enrichment of genes contributing to pattern
specification processes, limb development, and skeletal system development. Moreover, marmoset
species-specific DMPs are enriched for genes with functions very closely related to skeletal development,
such as osteoblast differentiation and ossification, as well as genes involved in metabolism and the
development of other organ systems including skeletal muscles, nerves, the brain, the heart, blood vessels,
kidneys, eyes, and ears (File S6). Several enriched pathways reinforce these molecular functions, and
additional pathways related to cancers and other disease were also identified (File S7).
Out of the species-specific DMPs identified, some were found to overlap with those previously
identified as being differentially methylated among primates. These include *ARTN*, *COL2A1*, and
GABBR1 which have been found to be differentially methylated among modern humans and great apes
(Hernando-Herraez et al., 2013), as well as *HOXD8*, *HOXD9*, and *HOXD10* which have been found to be
differentially methylated among modern and ancient hominins (Gokhman et al., 2014). Specifically, 1
marmoset-specific DMP is hypomethylated in *ARTN*, 3 marmoset-specific DMPs are hypermethylated in
COL2A1, 2 baboon-specific DMPs and 3 marmoset-specific DMPs are hypermethylated in *GABBR1*, and
2 marmoset-specific DMPs are hypomethylated in *GABBR1* (File S4). Additionally, 3 marmoset-specific
DMPs are hypermethylated in *HOXD8*, 1 baboon-specific DMP and 2 marmoset-specific DMPs are
hypermethylated in *HOXD9*, 2 chimpanzee-specific DMPs are hypomethylated in *HOXD9*, and 5
marmoset-specific DMPs are hypermethylated in *HOXD10*. Of these *HOXD10* species-specific DMPs, 4
have $\Delta\beta$ between 0.2 and 0.3 and 1 has a $\Delta\beta < 0.1$ (Figure 5, Table S13).
403
Inter-Specific DNA Methylation Profiling of HOXD10
Because of the high number of species-specific DMPs within *HOXD10*, methylation patterns
across this gene were assessed at a higher resolution using gene-specific sequencing techniques. Regular
and bisulfite sequences of several regions in the *HOXD10* gene were generated (Tables S14-S17). First,
loci across the entire *HOXD10* gene were examined from a subset of EPIC array samples – 3 baboons, 3
macaques, 3 vervets, 3 chimpanzees, and 3 marmosets (Figure 6, Tables S1 and S18, File S8-S9).
Additionally, one test locus (cg02193236) and its surrounding region in *HOXD10* were surveyed in all
NHP samples in order to confirm the reliability of the EPIC array in assessing DNA methylation levels
(Figure S9, Tables S1 and S19, File S10-S11). This test, which showed similar methylation patterns to
those identified with the EPIC array, supplemented previous methylation array replication studies that
have found resulting data to be highly correlated in replicate chimpanzee tissue samples (>98%
correlated) (Pai et al., 2011) and in replicate macaque tissue samples (97% correlated) (Ong et al., 2014).
Following the alignment of these sequences to the appropriate NHP references, the presence and absence
of methylation across the *HOXD10* gene in each animal was determined (Tables S18-S19, Files S8-S11).
These data reveal that across the *HOXD10* gene, NHPs display generally low methylation with some

clustered increased amounts of methylation upstream of the gene and at the start of the gene body (Figure
6).
421
422   **Discussion:**
423   This study explored skeletal gene regulation and its relation to complex traits and species
differences by evaluating genome-wide and gene-specific DNA methylation patterns in trabecular bone
from several NHP species. These species are phylogenetically representative of the order and display
anatomical and locomotor differences. Specifically, while the sampled OWMs (baboons, macaques,
vervets) display varying levels of terrestrial and arboreal quadrupedal locomotion, the sampled ape
species (chimpanzees) is a knuckle-walking quadruped that also vertical climbs and occasionally walks
bipedally (Cawthon Lang, 2006). Even more distinct are the sampled NWMs (marmosets) which
locomote via vertical clinging and leaping (Cawthon Lang, 2005). Lemurs and lorises are the only major
taxonomic primate groups not examined in this study. Regardless, the current sample set provided a
unique opportunity to examine skeletal epigenetic differences in relation to bone morphology at an
evolutionary scale. In this research we assume that adult bone methylation patterns reflect gene regulation
that enables the maintenance of post-maturation skeletal morphologies. Some of these patterns may be
impacted by environmental forces, like diet and biomechanical loading, but we attempted to control for
these variables by only including NHPs from captive colonies with similar environmental exposures.
Nevertheless, we are unable to ascertain how substantially the adult skeletal methylation in our sample set
deviates from the regulatory patterns that initially arose during early skeletal element development.
Despite these limitations, the novel data collected in this study can help to inform the relationship
between epigenetic variation and skeletal traits. First, we evaluated the association between intra-specific
variation in skeletal DNA methylation and intra-specific variation in femur morphology. Second, we
assessed inter-specific DNA methylation differences and the potential contribution of these lineage-
specific patterns to species-specific morphologies.
444
445   *Intra-specific DMPs are not readily associated with morphological variants.*
446   With respect to intra-specific morphology, very few sites were found to be differentially
methylated. DMPs were only identified in association with baboon bicondylar femur length, baboon
maximum femur length, macaque proximal femur width, macaque medial condyle width, vervet superior
shaft width, vervet inferior shaft width, vervet anatomical neck height, and chimpanzee anatomical neck
length (Table S8). Additionally, most of these associations only identified one DMP. This limited number
of associations may be due to the small sample sizes within each species or to the small amount of
variation identified in each morphology. This latter point is supported in that almost all the morphologies
with methylation associations also have the highest intra-specific variation in size (Figure S5, Files S2-
S3). Further, because the animals included in this study were born and raised in captivity, their limited
exposure to environmental variation may limit both the range of phenotypic variation and methylation
variation present within species. Alternatively, it may be the case that DNA methylation variation does
not have a large influence on nonpathological femur morphology within NHPs.
458   Of the few intra-specific methylation patterns associated with morphology, they likely have weak
functional effects. Research has shown that individual site-specific methylation changes are not readily
associated with differential gene expression (Bork et al., 2010; Chen et al., 2011; Koch et al., 2011).

Rather, differential gene expression is made possible through the accumulation of several methylation

changes within promotor regions (Suzuki & Bird, 2008) or across the gene body (Singer, Kosti, Pachter,

& Mandel-Gutfreund, 2015). Detecting accumulations of methylation changes across genes should have

been possible in the current study as 4-19 probes targeted each gene on average (Table S4). Thus, the

individual sites identified in this study are not due to experimental limitations and likely have limited

impacts on gene regulation. This is further supported by the degree of methylation variation observed

among these DMPs. Sites that have an average change in mean methylation less than 10% ($\Delta\beta < 0.1$) are

thought to have little biological relevance (Hernando-Herraez et al., 2013). Thus, those DMPs with small

changes in methylation likely have little to no biological function. Further, some of the DMPs with $\Delta\beta \geq$

0.1 appear to be highly influenced by small subsets of the sample sets. For example, in the case of

macaque proximal width, the majority of methylation change is due to two individuals (1 female and 1

male) that have extremely low methylation at cg19349877 as compared to other macaques (File S3).

Additionally, in the case of chimpanzee anatomical neck length, the 273 DMPs identified are largely

affected by one individual that has a longer anatomical neck than other chimpanzees. Lastly, the limited

enrichment of GO biological processes and KEGG pathway functions among detected DMPs indicates a

lack of common function among these associated regions. Overall, these finding suggest that while some

DMPs appear to be associated with intra-specific morphological variation in NHPs, not enough evidence

is present to support them having a functional role in the development and maintenance of this

morphological variation.

Differential methylation is observed among NHP species.

With respect to inter-specific variation, several sites were found to have significant species-

specific methylation differences. Specifically, out of the 39,802 sites examined, 650 species-specific

DMPs were identified in baboons, 257 in macaques, 639 in vervets, 2,796 in chimpanzees, and 13,778 in

marmosets, and these span 7,320 genes (Tables S11-S12). These numbers of species-specific DMPs did

not substantially change when accounting for age scaled to life expectancy instead of exact age (File S4).

However, many of these DMPs had biologically insignificant changes in mean methylation. Thus, several

$\Delta\beta$ cutoff thresholds were considered – from a 10% change in mean methylation ($\Delta\beta \geq 0.1$) up to a 40%

change in mean methylation ($\Delta\beta \geq 0.4$) – which reduced the overall numbers of species-specific DMPs.

Regardless of $\Delta\beta$ cutoff threshold, more species-specific DMPs were found in the NWM marmosets,

followed by the great ape chimpanzees, and lastly the OWM baboons, macaques, and vervets. This trend

in numbers of species-specific DMPs is expected given the known phylogenetic relationships between

primates, with OWMs more closely related to apes and NWMs more distantly related to both groups

(Perelman et al., 2011; Rogers & Gibbs, 2014). However, the number of marmoset-specific DMPs is

substantially larger than those for other taxa. While this discrepancy may reflect marmosets having more

species-specific changes, aspects of the experimental design may also contribute to it. Marmosets are the

only NWM included in this study, so although some marmoset-specific DMPs are truly specific to

marmosets, others may be specific to all NWMs. Comparably, OWM-specific DMPs and catarrhine-

specific DMPs may be cancelled out from this study, as such changes would be shared between all

OWMs and catarrhines, respectively. Second, since marmosets are the most phylogenetically distant from

humans as compared to the other NHPs included in this study (Perelman et al., 2011; Rogers & Gibbs,

2014), the probe filtering steps may have biased downstream data in favor of finding significant results

primarily in marmosets. Third, the marmoset data itself has a slightly different normalized distribution,
with more mean methylation levels of 50%, than that for other NHPs (Figure S4), which may be due to
the chimerism often observed in marmosets. Levels of chimerism in marmoset skeletal tissues have not
been examined. However, blood-derived tissues are known to have high levels of chimerism (>50%
chimeric), as compared to epithelial tissues which have the lowest levels of all tissues examined thus far
(12% chimeric) (Malukiewicz et al., 2015; Ross, French, & Orti, 2007). Even if trabecular bone has low
chimerism, its close proximity to blood cells in bone marrow increases the chance of highly chimeric cell
contamination. Additionally, although when assessed computationally, the filtered array probes examined
in this study appear to hybridize sufficiently for marmosets (Figure S3), there may be other unknown
biological or technical issues that impede proper DNA methylation analyses from the EPIC array in
marmosets and may have inflated the overall number of marmoset-specific DMPs identified.
The number of species-specific DMPs identified in this study, is comparable to those identified in
a previous study that assessed methylation patterns in blood from chimpanzees, bonobos, gorillas, and
orangutans using the 450K array and similar alignment criteria filtering methods with a focus on sites that
had a $\Delta\beta \geq 0.1$ (Hernando-Herraez et al., 2013). This research used a final set of 99,919 probes that were
shared across all great ape species and covered 12,593 genes with at least 2 probes per gene. Out of these,
2,284 species-specific DMPs were found in humans, 1,245 in chimpanzees and bonobos, 1,374 in
gorillas, and 5,501 in orangutans (Hernando-Herraez et al., 2013). In the current study, when a $\Delta\beta \geq 0.1$
threshold was used, only 1,572 chimpanzee-specific DMPs were identified (Table S11). This number is
lower than the number found in previous research comparing chimpanzees to other great apes (Hernando-
Herraez et al., 2013), species that are evolutionarily closer to chimpanzees than are OWMs and NWMs
(Perelman et al., 2011; Rogers & Gibbs, 2014). However, the total number of sites examined in the
present study is approximately one-third of that examined in the prior great ape study. Thus, a 3-fold
increase in the number of sites examined might identify a 3-fold increase in chimpanzee-specific DMPs
as compared to OWMs and NWMs, which would be closer to expectations (Hernando-Herraez et al.,
2013).
Additionally, the number of DMPs that distinguish species from one another in the current study
is substantially smaller than the numbers of DMPs identified between different skeletal tissue types,
between different age cohorts, and between individuals with different skeletal disease states within a NHP
species (Housman et al., 2018). This finding is to be expected since differences in DNA methylation
which regulate gene expression (Singer et al., 2015; Suzuki & Bird, 2008) should increase or decrease
with the degree of differences between cellular functions among comparative groups (Zhang et al., 2013).
In the case of different tissues, substantial DNA methylation differences likely promote distinct gene
regulation that is necessary for cells in different tissues to promote different functions. In the case of
different age cohorts, slightly fewer DNA methylation differences within the same skeletal tissue may
allow cells to emphasize efforts on growth and development in juveniles, as compared to maintenance in
adults, without altering the general skeletal-related functions of this tissue. In the case of osteoarthritic
disease states, even fewer regulatory changes may be needed to initiate the dysregulation of tissue
function. Finally, the comparatively small number of DNA methylation differences between adult,
skeletally healthy, NHP species seems reasonable given that other studies have also noted the presence of
more regulatory variation within species than between species (Uebbing et al., 2016). In summary,
epigenetic and regulatory differences, which control and enable age-dependent organ functions, should be

greater within a species when comparing between tissue types, age cohorts, and disease states, than when
comparing between species.
When evaluating how well the methylation patterns at species-specific DMPs cluster samples, a
$\Delta\beta \geq 0.4$ threshold is necessary to achieve clustering into distinct species (Figure 3). Less stringent $\Delta\beta$
thresholds are able to separate apes, NWMs, and OWMs into distinct groups, but the OWMs do not form
monophyletic species groups at these thresholds (Figure S6). Global changes in methylation show similar
phylogenetic patterns. While average species methylation patterns reveal a well-supported tree topology
that reflects known phylogenetic relationships between taxa (Perelman et al., 2011; Rogers & Gibbs,
2014) (Figure 4), global changes in methylation among individual animals do not distinguish OWMs into
distinct monophyletic groups (Figure S7). As before, parsing down these global changes to only species-
specific DMPs with $\Delta\beta \geq 0.4$ fixes the phylogeny (Figure S8). In previous studies, global methylation
changes were able to separate great apes into species-specific phylogenetic groups (Hernando-Herraez et
al., 2013). Conversely, the need for a higher $\Delta\beta$ cutoff threshold to distinguish species in the current study
may be due to evolutionary reasons.
The divergence times between OWMs (baboons, macaques, and vervets) are comparable to those
between great apes (humans, chimpanzees, gorillas, and orangutans) (Perelman et al., 2011). Thus,
divergence times do not explain why global changes in methylation are unable to resolve species-specific
phylogenetic clades in the current study as compared to previous research. On the other hand, while
nonhuman great apes have experienced higher rates of molecular evolution as compared to humans
(Elango, Thomas, NISC Comparative Sequencing Program, & Yi, 2006), baboons and macaques have
slow rates of molecular evolution as compared to other OWMs (Elango, Lee, Peng, Loh, & Yi, 2009),
which may correspond to slower rates of epigenetic evolution. This might make baboons and macaques
appear more similar to vervets than expected, and further, this may make resolving the phylogenetic
divergences between OWMs more difficult than that between the great apes. Additionally, the number of
sites included in the present study (39,802) as compared to previous studies (99,919) (Hernando-Herraez
et al., 2013) may limit the ability of the present study to fully resolve species-specific lineages. This is
reinforced by the low support of several branches in the phylogeny based on global changes in
methylation across individual animals (Figure S7). Conversely, sample size was likely not a contributing
factor to the discrepancies between the current study (n=58) and previous studies (n=32) (Hernando-
Herraez et al., 2013), as the current study has a slightly larger sample size. However, the number of
individuals per species was more uniform in previous studies (Hernando-Herraez et al., 2013) than in the
current study.
Alternatively, the fact that global changes in methylation are unable to fully resolve OWM
species-specific phylogenetic clades in the current study, may instead indicate that not enough time has
passed for OWM species to evolve fixed epigenetic changes between taxa in this tissue. Additionally, it is
possible that epigenetic variation at many of the sites examined in this study are under balancing selection
in OWMs which prevents these markers from accurately resolving the evolutionary divergences between
these species. Previous research of gene regulation differences between species has found that some
deviations in gene expression may be under directional or balancing selection (Romero, Ruvinsky, &
Gilad, 2012; Whitehead & Crawford, 2006). Given the relatively low variance in methylation across
species in combination with lineage-specific changes in mean methylation at the species-specific DMPs
identified in this study, it is possible that these loci are experiencing similar selective pressures (Romero

et al., 2012). Nevertheless, most inter-specific regulatory differences appear to be under stabilizing
selection or neutral evolution (Brawand et al., 2011; Gilad, 2012; Romero et al., 2012), and this cannot be
ruled out in the present study.
590
591 ***Inter-specific DMPs are found in genes enriched for functions associated with skeletal traits.***
592    The evolution of methylation changes along specific NHP lineages is associated with several
functions that may contribute to species-specific phenotypic differences (Files S6-S7). First, several
skeletal tissue functions are enriched in species-specific DMPs. Among almost all NHPs, cellular
adhesion functions are highly enriched. Cellular adhesion is necessary for cells to attach to other cells or
extracellular matrix, which is a necessity for bone cells (Mbalaviele, Shin, & Civitelli, 2006).
Additionally, in baboons, vervets, chimpanzees, and marmosets, anatomical developmental processes are
enriched, and in chimpanzees and marmosets, pattern specification processes, limb development, and
skeletal system development are enriched. Lastly, in marmosets, specific skeletal functions, such as
osteoblast differentiation and ossification are also enriched. Overall, these functions validate that most
patterns of differential methylation relate to skeletal tissue function, regulation, development, and
maintenance, as well to larger anatomical developmental processes. Additionally, functions not specific to
the skeletal system were identified. In chimpanzees and marmosets, transcription and gene expression
regulatory functions were enriched. Further, in marmosets, functions related to the development of
skeletal muscles, nerves, the brain, the heart, blood vessels, kidneys, eyes, and ears were also enriched.
All together, these findings suggest that many species-specific changes in methylation may contribute to
the regulation of complex phenotypic changes. While this relationship was not observed for intra-specific
skeletal morphology variation, other skeletal traits not examined in this study may be related.
Nevertheless, many of the genes associated with the described functions only contain an average of 1-2
differentially methylated sites. As described above, individual site-specific methylation changes are not
readily associated with differential gene expression (Bork et al., 2010; Chen et al., 2011; Koch et al.,
2011). Therefore, the enriched functions identified are likely not true biological effects due to methylation
differences on their own. Rather they hint at biological effects that may be the result of the combined
effects of several genetic, epigenetic, and other regulatory processes.
615    Finally, some of the genes containing species-specific DMPs overlap with those previously
identified as being differentially methylated in other tissues among primates. Specifically, the
neurotrophic factor *ARTN* shows species-specific hypomethylation at 1 site in marmosets, and in previous
work it shows species-specific hypermethylation at 3 sites in humans as compared to other great apes
(Hernando-Herraez et al., 2013). Additionally, *COL2A1* which codes for the predominant type of collagen
in cartilage, shows species-specific hypermethylation at 3 sites in marmosets, and in previous work it
shows species-specific hypermethylation of 4 sites in humans as compared to other great apes (Hernando-
Herraez et al., 2013). The neuronal receptor *GABBR1* shows more complicated methylation patterns. In
the current study, *GABBR1* shows species-specific hypermethylation at 2 sites in baboons,
hypermethylation at 3 sites in marmosets, and hypomethylation at 2 sites in marmosets. In previous work,
GABBR1 shows hypomethylation in orangutans and hypermethylation in chimpanzees and bonobos
(Hernando-Herraez et al., 2013). Lastly, 3 genes within the *HOXD* cluster, which are involved in limb
development, also show complicated methylation patterns. In the current study, *HOXD8* shows species-
specific hypermethylation at 3 sites in marmoset, *HOXD9* shows species-specific hypermethylation at 1

site in baboons and 2 sites in marmosets and species-specific hypomethylation at 2 sites in chimpanzees,

and *HOXD10* shows species-specific hypermethylation at 5 sites in marmosets. In previous work,

HOXD8 shows hypomethylation in modern and archaic hominins while *HOXD9* and *HOXD10* show

hypermethylation in archaic hominins as compared to modern humans (Gokhman et al., 2014).

In the *HOXD* cluster, *HOXD10* contains the largest number of NHP species-specific methylation

differences observed in this study, has an active role in anatomical development, and has been found to be

differentially methylated among hominins. Thus, it was selected for subsequent DNA methylation

profiling and analysis at a higher resolution using gene-specific sequencing techniques. *HOXD10*

specifically codes for a protein that functions as a sequence-specific transcription factor which is

expressed in the developing limb buds and is involved in differentiation and limb development. In the

current study, each NHP shows low to intermediate methylation levels across the gene body (Figure 5,

Table S13). A similar pattern is observed in the gene-specific methylation data, which further reveals that

HOXD10 is not highly methylated in NHPs. However, across all taxa, some clusters of hypermethylation

are found upstream of the gene and at the start of the gene body, with marmosets on average displaying

more methylation in the gene body than other taxa (Figure 6, Table S18). In the *HOXD10* gene body of

hominins, humans display hypomethylation, Neandertals displayed intermediate methylation levels, and

Denisovans displayed high levels of methylation (Gokhman et al., 2014). The variation of methylation

patterns in this gene body suggest that intermediate methylation levels may be a more ancestral epigenetic

state for this region in the primate lineage, while the extreme hypermethylation of this region in

Denisovans and the extreme hypomethylation of this region in humans may be derived epigenetic states.

Previous work has proposed that methylation differences in *HOXD10* may be associated with phenotypic

distinctions between modern human and archaic hominin limbs (Gokhman et al., 2014). While the current

study did not find substantial associations between methylation variation and aspects of femur

morphology within NHP species, further work to understand the role of differential methylation of

HOXD10 in promoting morphological changes of the limb should be explored.

Conclusions and future directions.

In conclusion, while only a few significant associations were identified between methylation and

femur morphologies, several significant differences in methylation were observed inter-specifically.

Moreover, these species-specific DMPs were found in genes enriched for functions associated with

complex skeletal traits. This is the first study to characterize DNA methylation patterns in skeletal tissues

from a taxonomically diverse set of NHPs, and it is the first study to directly compare these patterns to the

nonpathological morphologies of the skeletal elements from which the tissues were derived. This design

enabled an initial exploration of skeletal gene regulation and its relation to complex traits and species

differences for which little else is currently known.

Ultimately, this work forms a foundation for future explorations of gene regulation and skeletal

trait evolution in primates. The dataset collected and investigated here provides an initial sampling of

skeletal methylation patterns in NHPs, and future primate skeletal epigenetics investigations can add to

these data at population- and evolutionary-scales. At a population-level, increasing the number of

individuals from one species can provide further insight into how genetic variation interacts with gene

regulation to contribute to a wider variety of healthy and pathological skeletal traits. Additionally,

sampling individuals of different ages would provide further insight into how gene regulation across

development contributes to skeletal anatomy. Adult bone methylation patterns are limited in their ability
to fully inform how gene regulation contributes to the differential development of skeletal morphologies.
In the present study, we assume that regulatory patterns present in adult skeletal tissues are involved in (or
at least not inhibiting) the maintenance of adult skeletal morphology, and that the regulatory patterns that
enable the maintenance of adult skeletal traits reflect those that enabled the development of those traits.
Future work in the field of primate skeletal epigenetics could explicitly test the degree to which adult
bone methylation patterns reflect patterns that arise during early skeletal element development. Lastly, at
an evolutionary-level, increasing the number of represented species, both extant and extinct, will further
inform our understanding of gene regulation evolution in primates and how patterns of variation
contribute to skeletal phenotypes.
681
Acknowledgements:

This work was supported by the National Institutes of Health (P01 HL028972 to Anthony G.
Comuzzie); the Leakey Foundation (Research Grant for Doctoral Students to G.H.); the Wenner-Gren
Foundation (Gr. 9310 to G.H.); the Nacey Maggioncalda Foundation (James F. Nacey Fellowship to
G.H.); the International Primatological Society (to G.H.); Sigma Xi (Grant-in-Aid of Research to G.H.);
the ASU Center for Evolution and Medicine (Venture Fund to G.H.); the ASU Graduate Research and
Support Program (to G.H.). Additionally, this investigation used resources that were supported by the
Southwest National Primate Research Center grant P51 OD011133 from the Office of Research
Infrastructure Programs, National Institutes of Health.
We thank Eric D. Johnson and members of the Department of Genetics at the Texas Biomedical
Research Institute, including Anthony G. Comuzzie, Anne Sheldrake, Jaydee Foster, Kara Peterson, Mel
Carless, and Laura Cox, for helpful discussions. We also thank Megann Phillips for assistance with PCR
primer design.

Newly reported data have been made available on NCBI's Gene Expression Omnibus and are
accessible through the GEO SuperSeries accession number GSE103332, which includes the following
SubSeries accession numbers: GSE103279 (intra-specific baboon DNA methylation data), GSE103271
(intra-specific macaque DNA methylation data), GSE103280 (intra-specific vervet DNA methylation
data), GSE94677 (intra-specific chimp DNA methylation data), GSE103328 (intra-specific marmoset
DNA methylation data), and GSE103287 (inter-specific DNA methylation data).
701

References:

Ankel-Simons, F. (2007). *Primate Anatomy* (3rd ed.). New York: Academic Press.
Aryee, M. J., Jaffe, A. E., Corrada-Bravo, H., Ladd-Acosta, C., Feinberg, A. P., Hansen, K. D., &
Irizarry, R. A. (2014). Minfi: A flexible and comprehensive Bioconductor package for the
analysis of Infinium DNA methylation microarrays. *Bioinformatics*, *30*(10), 1363–1369. doi:
10.1093/bioinformatics/btu049
Babbitt, C. C., Fedrigo, O., Pfefferle, A. D., Boyle, A. P., Horvath, J. E., Furey, T. S., & Wray, G. A.
(2010). Both Noncoding and Protein-Coding RNAs Contribute to Gene Expression Evolution in
the Primate Brain. *Genome Biology and Evolution*, *2*(0), 67–79. doi: 10.1093/gbe/evq002

Bell, C. G., Wilson, G. A., Butcher, L. M., Roos, C., Walter, L., & Beck, S. (2012). Human-specific CpG
"beacons" identify loci associated with human-specific traits and disease. *Epigenetics*, *7*(10),
1188–1199. doi: 10.4161/epi.22127
Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful
Approach to Multiple Testing. *Journal of the Royal Statistical Society. Series B*
(Methodological), *57*(1), 289–300.
Blekhman, R., Oshlack, A., Chabot, A. E., Smyth, G. K., & Gilad, Y. (2008). Gene Regulation in
Primates Evolves under Tissue-Specific Selection Pressures. *PLoS Genetics*, *4*(11), e1000271.
doi: 10.1371/journal.pgen.1000271
Bork, S., Pfister, S., Witt, H., Horn, P., Korn, B., Ho, A. D., & Wagner, W. (2010). DNA methylation
pattern changes upon long-term culture and aging of human mesenchymal stromal cells. *Aging*
Cell, *9*(1), 54–63. doi: 10.1111/j.1474-9726.2009.00535.x
Bovée, J. V. M. G., Hogendoorn, P. C. W., Wunder, J. S., & Alman, B. A. (2010). Cartilage tumours and
bone development: Molecular pathology and possible therapeutic targets. *Nature Reviews*
Cancer, *10*(7), 481–488. doi: 10.1038/nrc2869
Brand-Saberi, B. (2005). Genetic and epigenetic control of skeletal muscle development. *Annals of*
Anatomy - Anatomischer Anzeiger, *187*(3), 199–207. doi: 10.1016/j.aanat.2004.12.018
Brawand, D., Soumillon, M., Necsulea, A., Julien, P., Csárdi, G., Harrigan, P., … Kaessmann, H. (2011).
The evolution of gene expression levels in mammalian organs. *Nature*, *478*(7369), 343–348. doi:
10.1038/nature10532
Cáceres, M., Lachuer, J., Zapala, M. A., Redmond, J. C., Kudo, L., Geschwind, D. H., … Barlow, C.
(2003). Elevated gene expression levels distinguish human from non-human primate brains.
Proceedings of the National Academy of Sciences, *100*(22), 13030–13035.
Cawthon Lang, K. A. (2005). Primate Factsheets: Common marmoset (Callithrix jacchus) Taxonomy,
Morphology, & Ecology. Retrieved January 13, 2019, from
http://pin.primate.wisc.edu/factsheets/entry/common_marmoset
Cawthon Lang, K. A. (2006). Primate Factsheets: Chimpanzee (Pan troglodytes) Taxonomy,
Morphology, & Ecology. Retrieved January 13, 2019, from
http://pin.primate.wisc.edu/factsheets/entry/chimpanzee
Chen, Y., Choufani, S., Ferreira, J. C., Grafodatskaya, D., Butcher, D. T., & Weksberg, R. (2011).
Sequence overlap between autosomal and sex-linked probes on the Illumina
HumanMethylation27 microarray. *Genomics*, *97*(4), 214–222. doi: 10.1016/j.ygeno.2010.12.004
Corner, B., Lele, S., & Richtsmeier, J. (1992). Measuring precision of three-dimensional landmark data.
Journal of Quantitative Anthropology, *3*, 347–359.
de Andrés, M. C., Kingham, E., Imagawa, K., Gonzalez, A., Roach, H. I., Wilson, D. I., & Oreffo, R. O.
C. (2013). Epigenetic Regulation during Fetal Femur Development: DNA Methylation Matters.
PLoS ONE, *8*(1), e54957. doi: 10.1371/journal.pone.0054957
Delgado-Calle, J., Fernández, A. F., Sainz, J., Zarrabeitia, M. T., Sañudo, C., García-Renedo, R., …
Riancho, J. A. (2013). Genome-wide profiling of bone reveals differentially methylated regions in
osteoporosis and osteoarthritis. *Arthritis & Rheumatism*, *65*(1), 197–205. doi: 10.1002/art.37753
den Hollander, W., Ramos, Y. F. M., Bos, S. D., Bomer, N., van der Breggen, R., Lakenberg, N., …
Meulenbelt, I. (2014). Knee and hip articular cartilage have distinct epigenomic landscapes:

Implications for future cartilage regeneration approaches. *Annals of the Rheumatic Diseases*,
73(12), 2208–2212. doi: 10.1136/annrheumdis-2014-205980

Dimitriou, R., Jones, E., McGonagle, D., & Giannoudis, P. V. (2011). Bone regeneration: Current
concepts and future directions. *BMC Medicine*, *9*(1), 1.

Elango, N., Lee, J., Peng, Z., Loh, Y.-H. E., & Yi, S. V. (2009). Evolutionary rate variation in Old World
monkeys. *Biology Letters*, *5*(3), 405–408. doi: 10.1098/rsbl.2008.0712

Elango, N., Thomas, J. W., NISC Comparative Sequencing Program, & Yi, S. V. (2006). Variable
molecular clocks in hominoids. *Proceedings of the National Academy of Sciences*, *103*(5), 1370–
1375. doi: 10.1073/pnas.0510716103

Enard, W., Fassbender, A., Model, F., Adorján, P., Pääbo, S., & Olek, A. (2004). Differences in DNA
methylation patterns between humans and chimpanzees. *Current Biology*, *14*(4), R148–R149.

Farcas, R., Schneider, E., Frauenknecht, K., Kondova, I., Bontrop, R., Bohl, J., … Haaf, T. (2009).
Differences in DNA Methylation Patterns and Expression of the CCRK Gene in Human and
Nonhuman Primate Cortices. *Molecular Biology and Evolution*, *26*(6), 1379–1389. doi:
10.1093/molbev/msp046

Fernández-Tajes, J., Soto-Hermida, A., Vázquez-Mosquera, M. E., Cortés-Pereira, E., Mosquera, A.,
Fernández-Moreno, M., … Blanco, F. J. (2014). Genome-wide DNA methylation analysis of
articular chondrocytes reveals a cluster of osteoarthritic patients. *Annals of the Rheumatic
Diseases*, *73*(4), 668–677. doi: 10.1136/annrheumdis-2012-202783

Flanagan, J. M., Popendikyte, V., Pozdniakovaite, N., Sobolev, M., Assadzadeh, A., Schumacher, A., …
Petronis, A. (2006). Intra- and Interindividual Epigenetic Variation in Human Germ Cells.
American Journal of Human Genetics, *79*(1), 67–84.

Fleagle, J. G. (1999). *Primate Adaptation and Evolution*. New York: Academic Press.

Fortin, J.-P., Triche, T., & Hansen, K. (2016). Preprocessing, normalization and integration of the
Illumina HumanMethylationEPIC array. *BioRxiv*, 065490. doi: 10.1101/065490

Fraga, M. F., Ballestar, E., Paz, M. F., Ropero, S., Setien, F., Ballestar, M. L., … Esteller, M. (2005).
Epigenetic differences arise during the lifetime of monozygotic twins. *Proceedings of the
National Academy of Sciences of the United States of America*, *102*(30), 10604–10609. doi:
10.1073/pnas.0500398102

Fukuda, K., Ichiyanagi, K., Yamada, Y., Go, Y., Udono, T., Wada, S., … Sasaki, H. (2013). Regional
DNA methylation differences between humans and chimpanzees are associated with genetic
changes, transcriptional divergence and disease genes. *Journal of Human Genetics*, *58*(7), 446–
454. doi: 10.1038/jhg.2013.55

Gama-Sosa, M. A., Midgett, R. M., Slagel, V. A., Githens, S., Kuo, K. C., Gehrke, C. W., & Ehrlich, M.
(1983). Tissue-specific differences in DNA methylation in various mammals. *Biochimica et
Biophysica Acta (BBA)-Gene Structure and Expression*, *740*(2), 212–219.

Gao, F., Niu, Y., Sun, Y. E., Lu, H., Chen, Y., Li, S., … Tan, T. (2017). De novo DNA methylation
during monkey pre-implantation embryogenesis. *Cell Research*. doi: 10.1038/cr.2017.25

García-Ibarbia, C., Delgado-Calle, J., Casafont, I., Velasco, J., Arozamena, J., Pérez-Núñez, M. I., …
Riancho, J. A. (2013). Contribution of genetic and epigenetic mechanisms to Wnt pathway
activity in prevalent skeletal disorders. *Gene*, *532*(2), 165–172. doi: 10.1016/j.gene.2013.09.080

Geeleher, P., Hartnett, L., Egan, L. J., Golden, A., Raja Ali, R. A., & Seoighe, C. (2013). Gene-set
analysis is severely biased when applied to genome-wide methylation data. *Bioinformatics*
(Oxford, England), *29*(15), 1851–1857. doi: 10.1093/bioinformatics/btt311
Gibbs, J. R., van der Brug, M. P., Hernandez, D. G., Traynor, B. J., Nalls, M. A., Lai, S.-L., … Singleton,
A. B. (2010). Abundant Quantitative Trait Loci Exist for DNA Methylation and Gene Expression
in Human Brain. *PLoS Genetics*, *6*(5), e1000952. doi: 10.1371/journal.pgen.1000952
Gilad, Y. (2012). Using Genomic Tools to Study Regulatory Evolution. *Methods in Molecular Biology*
(Clifton, N.J.), *856*, 335–361. doi: 10.1007/978-1-61779-585-5_14
Gokhman, D., Agranat-Tamir, L., Housman, G., Garcia-Perez, R., Nissim-Rafinia, M., Mallick, S., …
Carmel, L. (2017). Extensive Regulatory Changes in Genes Affecting Vocal and Facial Anatomy
Separate Modern from Archaic Humans. *BioRxiv*, 106955. doi: 10.1101/106955
Gokhman, D., Lavi, E., Prüfer, K., Fraga, M. F., Riancho, J. A., Kelso, J., … Carmel, L. (2014).
Reconstructing the DNA methylation maps of the Neandertal and the Denisovan. *Science (New*
York, N.Y.), *344*(6183), 523–527. doi: 10.1126/science.1250368
Gokhman, D., Meshorer, E., & Carmel, L. (2016). Epigenetics: It's Getting Old. Past Meets Future in
Paleoepigenetics. *Trends in Ecology & Evolution*, *31*(4), 290–300. doi:
10.1016/j.tree.2016.01.010
Goldring, M. B., & Marcu, K. B. (2012). Epigenomic and microRNA-mediated regulation in cartilage
development, homeostasis, and osteoarthritis. *Trends in Molecular Medicine*, *18*(2), 109–118.
doi: 10.1016/j.molmed.2011.11.005
Haygood, R., Fedrigo, O., Hanson, B., Yokoyama, K.-D., & Wray, G. A. (2007). Promoter regions of
many neural- and nutrition-related genes have experienced positive selection during human
evolution. *Nature Genetics*, *39*(9), 1140–1144. doi: 10.1038/ng2104
Henriksen, M., Creaby, M. W., Lund, H., Juhl, C., & Christensen, R. (2014). Is there a causal link
between knee loading and knee osteoarthritis progression? A systematic review and meta-analysis
of cohort studies and randomised trials. *BMJ Open*, *4*(7), e005368.
Hernando-Herraez, I., Prado-Martinez, J., Garg, P., Fernandez-Callejo, M., Heyn, H., Hvilsom, C., …
Marques-Bonet, T. (2013). Dynamics of DNA Methylation in Recent Human and Great Ape
Evolution. *PLOS Genet*, *9*(9), e1003763. doi: 10.1371/journal.pgen.1003763
Heyn, H., Moran, S., Hernando-Herraez, I., Sayols, S., Gomez, A., Sandoval, J., … Esteller, M. (2013).
DNA methylation contributes to natural human variation. *Genome Research*, *23*(9), 1363–1372.
doi: 10.1101/gr.154187.112
Housman, G., Havill, L. M., Quillen, E. E., Comuzzie, A. G., & Stone, A. C. (2018). Assessment of DNA
Methylation Patterns in the Bone and Cartilage of a Nonhuman Primate Model of Osteoarthritis.
CARTILAGE. doi: 10.1177/1947603518759173
Huber, W., Carey, V. J., Gentleman, R., Anders, S., Carlson, M., Carvalho, B. S., … Morgan, M. (2015).
Orchestrating high-throughput genomic analysis with Bioconductor. *Nature Methods*, *12*(2), 115–
121. doi: 10.1038/nmeth.3252
Iliopoulos, D., Malizos, K. N., Oikonomou, P., & Tsezou, A. (2008). Integrative MicroRNA and
Proteomic Approaches Identify Novel Osteoarthritis Genes and Their Collaborative Metabolic
and Inflammatory Networks. *PLoS ONE*, *3*(11), e3740. doi: 10.1371/journal.pone.0003740

Jacob, R. A., Gretz, D. M., Taylor, P. C., James, S. J., Pogribny, I. P., Miller, B. J., … Swendseid, M. E.
(1998). Moderate folate depletion increases plasma homocysteine and decreases lymphocyte
DNA methylation in postmenopausal women. *The Journal of Nutrition*, *128*(7), 1204–1212.

Jaffe, A. E., & Irizarry, R. A. (2014). Accounting for cellular heterogeneity is critical in epigenome-wide
association studies. *Genome Biology*, *15*(2), R31. doi: 10.1186/gb-2014-15-2-r31

Jeffries, M. A., Donica, M., Baker, L. W., Stevenson, M. E., Annan, A. C., Beth Humphrey, M., …
Sawalha, A. H. (2016). Genome-Wide DNA Methylation Study Identifies Significant Epigenomic
Changes in Osteoarthritic Subchondral Bone and Similarity to Overlying Cartilage. *Arthritis &*
Rheumatology, *68*(6), 1403–1414. doi: 10.1002/art.39555

Jeffries, M. A., Donica, M., Baker, L. W., Stevenson, M. E., Annan, A. C., Humphrey, M. B., …
Sawalha, A. H. (2014). Genome-Wide DNA Methylation Study Identifies Significant Epigenomic
Changes in Osteoarthritic Cartilage. *Arthritis & Rheumatology*, *66*(10), 2804–2815. doi:
10.1002/art.38762

Joganic, J. L., Willmore, K. E., Richtsmeier, J. T., Weiss, K. M., Mahaney, M. C., Rogers, J., &
Cheverud, J. M. (2017). Additive genetic variation in the craniofacial skeleton of baboons (genus
Papio) and its relationship to body and cranial size. *American Journal of Physical Anthropology*,
1–17. doi: 10.1002/ajpa.23349

Karere, G. M., Glenn, J. P., Birnbaum, S., Rainwater, D. L., Mahaney, M. C., VandeBerg, J. L., & Cox,
L. A. (2013). Identification of candidate genes encoding an LDL-C QTL in baboons. *Journal of*
Lipid Research, *54*(7), 1776–1785.

Karere, G. M., Glenn, J. P., VandeBerg, J. L., & Cox, L. A. (2010). Identification of baboon microRNAs
expressed in liver and lymphocytes. *Journal of Biomedical Science*, *17*(1), 1.

Karere, G. M., Glenn, J. P., VandeBerg, J. L., & Cox, L. A. (2012). Differential microRNA response to a
high-cholesterol, high-fat diet in livers of low and high LDL-C baboons. *BMC Genomics*, *13*(1),
1.

Kasaai, B., Gaumond, M.-H., & Moffatt, P. (2013). Regulation of the Bone-restricted IFITM-like (Bril)
Gene Transcription by Sp and Gli Family Members and CpG Methylation. *Journal of Biological*
Chemistry, *288*(19), 13278–13294. doi: 10.1074/jbc.M113.457010

King, M. C., & Wilson, A. C. (1975). Evolution at two levels in humans and chimpanzees. *Science*,
188(4184), 107–116. doi: 10.1126/science.1090005

Koch, C. M., Suschek, C. V., Lin, Q., Bork, S., Goergens, M., Joussen, S., … Wagner, W. (2011).
Specific Age-Associated DNA Methylation Changes in Human Dermal Fibroblasts. *PLOS ONE*,
6(2), e16679. doi: 10.1371/journal.pone.0016679

Kothapalli, K. S. D., Anthony, J. C., Pan, B. S., Hsieh, A. T., Nathanielsz, P. W., & Brenna, J. T. (2007).
Differential Cerebral Cortex Transcriptomes of Baboon Neonates Consuming Moderate and High
Docosahexaenoic Acid Formulas. *PLoS ONE*, *2*(4), e370. doi: 10.1371/journal.pone.0000370

Lea, A. J., Altmann, J., Alberts, S. C., & Tung, J. (2016). Resource base influences genome-wide DNA
methylation levels in wild baboons (Papio cynocephalus). *Molecular Ecology*, *25*(8), 1681–1696.
doi: 10.1111/mec.13436

Leek, J. T., Johnson, W. E., Parker, H. S., Jaffe, A. E., & Storey, J. D. (2012). The sva package for
removing batch effects and other unwanted variation in high-throughput experiments.
Bioinformatics, *28*(6), 882–883. doi: 10.1093/bioinformatics/bts034

Leek, J. T., & Storey, J. D. (2007). Capturing Heterogeneity in Gene Expression Studies by Surrogate
878   Variable Analysis. *PLoS Genetics*, *3*(9), e161. doi: 10.1371/journal.pgen.0030161
879   Leek, J. T., & Storey, J. D. (2008). A general framework for multiple testing dependence. *Proceedings of*
880   *the National Academy of Sciences*, *105*(48), 18718–18723. doi: 10.1073/pnas.0808709105
881   Leigh, S. R., & Shea, B. T. (1995). Ontogeny and the evolution of adult body size dimorphism in apes.
882   *American Journal of Primatology*, *36*(1), 37–60. doi: 10.1002/ajp.1350360104
883   Lewton, K. L. (2017). The effects of captive versus wild rearing environments on long bone articular
884   surfaces in common chimpanzees (Pan troglodytes). *PeerJ*, *5*, e3668. doi: 10.7717/peerj.3668
885   Lewton, K. L., Ritzman, T., Copes, L. E., Garland, T., & Capellini, T. D. (2018). Exercise-induced
886   loading increases ilium cortical area in a selectively bred mouse model. *American Journal of*
887   *Physical Anthropology*, 1–9. doi: 10.1002/ajpa.23770
888   Lindskog, C., Kuhlwilm, M., Davierwala, A., Fu, N., Hegde, G., Uhlén, M., … Pontén, F. (2014).
889   Analysis of candidate genes for lineage-specific expression changes in humans and primates.
890   *Journal of Proteome Research*, *13*(8), 3596–3606.
891   Ling, B. M. T., Bharathy, N., Chung, T.-K., Kok, W. K., Li, S., Tan, Y. H., … Taneja, R. (2012). Lysine
892   methyltransferase G9a methylates the transcription factor MyoD and regulates skeletal muscle
893   differentiation. *Proceedings of the National Academy of Sciences*, *109*(3), 841–846. doi:
894   10.1073/pnas.1111628109
895   Lister, R., Pelizzola, M., Dowen, R. H., Hawkins, R. D., Hon, G., Tonti-Filippini, J., … Ecker, J. R.
896   (2009). Human DNA methylomes at base resolution show widespread epigenomic differences.
897   *Nature*, *462*(7271), 315–322. doi: 10.1038/nature08514
898   Liu, Y., Aryee, M. J., Padyukov, L., Fallin, M. D., Hesselberg, E., Runarsson, A., … Feinberg, A. P.
899   (2013). Epigenome-wide association data implicate DNA methylation as an intermediary of
900   genetic risk in rheumatoid arthritis. *Nature Biotechnology*, *31*(2), 142–147. doi: 10.1038/nbt.2487
901   Lönnstedt, I., & Speed, T. (2002). Replicated microarray data. *Statistica Sinica*, *12*(1), 31–46.
902   Loughlin, J., & Reynard, L. N. (2015). Osteoarthritis: Epigenetics of articular cartilage in knee and hip
903   OA. *Nature Reviews Rheumatology*, *11*(1), 6–7. doi: 10.1038/nrrheum.2014.189
904   Macrini, T. E., Coan, H. B., Levine, S. M., Lerma, T., Saks, C. D., Araujo, D. J., … Havill, L. M. (2013).
905   Reproductive status and sex show strong effects on knee OA in a baboon model. *Osteoarthritis*
906   *and Cartilage / OARS, Osteoarthritis Research Society*, *21*(6), 839–848. doi:
907   10.1016/j.joca.2013.03.003
908   Madrid, A., Chopra, P., & Alisch, R. S. (2018). Species-Specific 5 mC and 5 hmC Genomic Landscapes
909   Indicate Epigenetic Contribution to Human Brain Evolution. *Frontiers in Molecular*
910   *Neuroscience*, *11*. doi: 10.3389/fnmol.2018.00039
911   Maksimovic, J., Phipson, B., & Oshlack, A. (2016). A cross-package Bioconductor workflow for
912   analysing methylation array data. *F1000Research*, *5*, 1281. doi: 10.12688/f1000research.8839.1
913   Malukiewicz, J., Boere, V., Fuzessy, L. F., Grativol, A. D., Silva, I. de O. e, Pereira, L. C. M., … Stone,
914   A. C. (2015). Natural and Anthropogenic Hybridization in Two Species of Eastern Brazilian
915   Marmosets (Callithrix jacchus and C. penicillata). *PLOS ONE*, *10*(6), e0127268. doi:
916   10.1371/journal.pone.0127268

Martin, D. I. K., Singer, M., Dhahbi, J., Mao, G., Zhang, L., Schroth, G. P., … Boffelli, D. (2011).
Phyloepigenomic comparison of great apes reveals a correlation between somatic and germline
methylation states. *Genome Research*, *21*(12), 2049–2057. doi: 10.1101/gr.122721.111
Mbalaviele, G., Shin, C. S., & Civitelli, R. (2006). Perspective: Cell–Cell Adhesion and Signaling
Through Cadherins: Connecting Bone Cells in Their Microenvironment. *Journal of Bone and*
Mineral Research, *21*(12), 1821–1827. doi: 10.1359/jbmr.060811
McCarthy, D. J., & Smyth, G. K. (2009). Testing significance relative to a fold-change threshold is a
TREAT. *Bioinformatics*, *25*(6), 765–771. doi: 10.1093/bioinformatics/btp053
McCartney, D. L., Walker, R. M., Morris, S. W., McIntosh, A. M., Porteous, D. J., & Evans, K. L.
(2016). Identification of polymorphic and off-target probe binding sites on the Illumina Infinium
MethylationEPIC BeadChip. *Genomics Data*, *9*, 22–24. doi: 10.1016/j.gdata.2016.05.012
McHenry, H. M., & Corruccini, R. S. (1978). The femur in early human evolution. *American Journal of*
Physical Anthropology, *49*(4), 473–487. doi: 10.1002/ajpa.1330490407
Mendizabal, I., Shi, L., Keller, T. E., Konopka, G., Preuss, T. M., Hsieh, T.-F., … Yi, S. V. (2016).
Comparative Methylome Analyses Identify Epigenetic Regulatory Loci of Human Brain
Evolution. *Molecular Biology and Evolution*, *33*(11), 2947–2959. doi: 10.1093/molbev/msw176
Moazedi-Fuerst, F. C., Hofner, M., Gruber, G., Weinhaeusel, A., Stradner, M. H., Angerer, H., …
Graninger, W. B. (2014). Epigenetic differences in human cartilage between mild and severe OA.
Journal of Orthopaedic Research, *32*(12), 1636–1645. doi: 10.1002/jor.22722
Molaro, A., Hodges, E., Fang, F., Song, Q., McCombie, W. R., Hannon, G. J., & Smith, A. D. (2011).
Sperm Methylation Profiles Reveal Features of Epigenetic Inheritance and Evolution in Primates.
Cell, *146*(6), 1029–1041. doi: 10.1016/j.cell.2011.08.016
Morris, J. A., Tsai, P.-C., Joehanes, R., Zheng, J., Trajanoska, K., Soerensen, M., … Bell, J. T. (2017).
Epigenome-wide association of DNA methylation in whole blood with bone mineral density.
Journal of Bone and Mineral Research, n/a-n/a. doi: 10.1002/jbmr.3148
Oates, N. A., van Vliet, J., Duffy, D. L., Kroes, H. Y., Martin, N. G., Boomsma, D. I., … Chong, S.
(2006). Increased DNA Methylation at the AXIN1 Gene in a Monozygotic Twin from a Pair
Discordant for a Caudal Duplication Anomaly. *American Journal of Human Genetics*, *79*(1),
155–162.
Ong, M.-L., Tan, P. Y., MacIsaac, J. L., Mah, S. M., Buschdorf, J. P., Cheong, C. Y., … Holbrook, J. D.
(2014). Infinium Monkeys: Infinium 450K Array for the Cynomolgus macaque (*Macaca*
fascicularis). *G3 : Genes|Genomes|Genetics*, *4*(7), 1227–1234. doi:
10.1534/g3.114.010967
Ostanek, B., Kranjc, T., Lovšin, N., Zupan, J., & Marc, J. (2018). Chapter 18—Epigenetic Mechanisms in
Osteoporosis. In A. Moskalev & A. M. Vaiserman (Eds.), *Epigenetics of Aging and Longevity*
(pp. 365–388). doi: 10.1016/B978-0-12-811060-7.00018-8
Pai, A. A., Bell, J. T., Marioni, J. C., Pritchard, J. K., & Gilad, Y. (2011). A Genome-Wide Study of DNA
Methylation Patterns and Gene Expression Levels in Multiple Human and Chimpanzee Tissues.
PLOS Genetics, *7*(2), e1001316. doi: 10.1371/journal.pgen.1001316
Palacios, D., & Puri, P. L. (2006). The epigenetic network regulating muscle development and
regeneration. *Journal of Cellular Physiology*, *207*(1), 1–11. doi: 10.1002/jcp.20489

Pandorf, C. E., Haddad, F., Wright, C., Bodell, P. W., & Baldwin, K. M. (2009). Differential epigenetic
modifications of histones at the myosin heavy chain genes in fast and slow skeletal muscle fibers
and in response to muscle unloading. *AJP: Cell Physiology*, *297*(1), C6–C16. doi:
10.1152/ajpcell.00075.2009

Paradis, E., Claude, J., & Strimmer, K. (2004). APE: Analyses of Phylogenetics and Evolution in R
language. *Bioinformatics (Oxford, England)*, *20*(2), 289–290.

Perelman, P., Johnson, W. E., Roos, C., Seuánez, H. N., Horvath, J. E., Moreira, M. A. M., … Pecon-
Slattery, J. (2011). A Molecular Phylogeny of Living Primates. *PLoS Genetics*, *7*(3), e1001342.
doi: 10.1371/journal.pgen.1001342

Petronis, A., Gottesman, I. I., Kan, P., Kennedy, J. L., Basile, V. S., Paterson, A. D., & Popendikyte, V.
(2003). Monozygotic twins exhibit numerous epigenetic differences: Clues to twin discordance?
Schizophrenia Bulletin, *29*(1), 169–178.

Phipson, B., Lee, S., Majewski, I. J., Alexander, W. S., & Smyth, G. K. (2016). Robust hyperparameter
estimation protects against hypervariable genes and improves power to detect differential
expression. *The Annals of Applied Statistics*, *10*(2), 946–963. doi: 10.1214/16-AOAS920

Phipson, B., Maksimovic, J., & Oshlack, A. (2016). missMethyl: An R package for analyzing data from
Illumina's HumanMethylation450 platform. *Bioinformatics*, *32*(2), 286–288. doi:
10.1093/bioinformatics/btv560

Prendergast, J. G., Campbell, H., Gilbert, N., Dunlop, M. G., Bickmore, W. A., & Semple, C. A. (2007).
Chromatin structure and evolution in the human genome. *BMC Evolutionary Biology*, *7*(1), 72.
doi: 10.1186/1471-2148-7-72

Provencal, N., Suderman, M. J., Guillemin, C., Massart, R., Ruggiero, A., Wang, D., … Szyf, M. (2012).
The Signature of Maternal Rearing in the Methylome in Rhesus Macaque Prefrontal Cortex and T
Cells. *Journal of Neuroscience*, *32*(44), 15626–15642. doi: 10.1523/JNEUROSCI.1470-12.2012

Rakyan, V. K., Hildmann, T., Novik, K. L., Lewin, J., Tost, J., Cox, A. V., … Beck, S. (2004). DNA
Methylation Profiling of the Human Major Histocompatibility Complex: A Pilot Study for the
Human Epigenome Project. *PLOS Biology*, *2*(12), e405. doi: 10.1371/journal.pbio.0020405

Ralston, S. H., & Uitterlinden, A. G. (2010). Genetics of Osteoporosis. *Endocrine Reviews*, *31*(5), 629–
662. doi: 10.1210/er.2009-0044

Ramos, Y. F. M., den Hollander, W., Bovée, J. V. M. G., Bomer, N., van der Breggen, R., Lakenberg, N.,
… Meulenbelt, I. (2014). Genes Involved in the Osteoarthritis Process Identified through Genome
Wide Expression Analysis in Articular Cartilage; the RAAK Study. *PLoS ONE*, *9*(7), 1–12. doi:
10.1371/journal.pone.0103056

Rampersaud, G. C., Kauwell, G. P., Hutson, A. D., Cerda, J. J., & Bailey, L. B. (2000). Genomic DNA
methylation decreases in response to moderate folate depletion in elderly women. *The American*
Journal of Clinical Nutrition, *72*(4), 998–1003.

Reynard, L. N. (2017). Analysis of genetics and DNA methylation in osteoarthritis: What have we learnt
about the disease? *Seminars in Cell & Developmental Biology*, *62*, 57–66. doi:
10.1016/j.semcdb.2016.04.017

Reynard, L. N., Bui, C., Syddall, C. M., & Loughlin, J. (2014). CpG methylation regulates allelic
expression of GDF5 by modulating binding of SP1 and SP3 repressor proteins to the

osteoarthritis susceptibility SNP rs143383. *Human Genetics*, *133*(8), 1059–1073. doi:
10.1007/s00439-014-1447-z

Ritchie, M. E., Phipson, B., Wu, D., Hu, Y., Law, C. W., Shi, W., & Smyth, G. K. (2015). Limma powers
differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids*
Research, gkv007. doi: 10.1093/nar/gkv007

Ritzman, T. B., Banovich, N., Buss, K. P., Guida, J., Rubel, M. A., Pinney, J., … Stone, A. C. (2017).
Facing the facts: The Runx2 gene is associated with variation in facial morphology in primates.
Journal of Human Evolution, *111*, 139–151. doi: 10.1016/j.jhevol.2017.06.014

Rivadeneira, F., Styrkársdottir, U., Estrada, K., Halldórsson, B. V., Hsu, Y.-H., Richards, J. B., …
Uitterlinden, A. G. (2009). Twenty bone-mineral-density loci identified by large-scale meta-
analysis of genome-wide association studies. *Nature Genetics*, *41*(11), 1199–1206. doi:
10.1038/ng.446

Rogers, J., & Gibbs, R. A. (2014). Comparative primate genomics: Emerging patterns of genome content
and dynamics. *Nature Reviews Genetics*, *15*(5), 347–359. doi: 10.1038/nrg3707

Romero, I. G., Ruvinsky, I., & Gilad, Y. (2012). Comparative studies of gene expression and the
evolution of gene regulation. *Nature Reviews Genetics*, *13*(7), 505–516. doi: 10.1038/nrg3229

Ross, C. N., French, J. A., & Orti, G. (2007). Germ-line chimerism and paternal care in marmosets
(Callithrix kuhlii). *Proceedings of the National Academy of Sciences*, *104*(15), 6278–6282. doi:
10.1073/pnas.0607426104

Rowe, N. (1996). *The Pictorial Guide to Living Primates*. East Hampton, New York: Pogonias Press.

Rushton, M. D., Reynard, L. N., Barter, M. J., Refaie, R., Rankin, K. S., Young, D. A., & Loughlin, J.
(2014). Characterization of the Cartilage DNA Methylome in Knee and Hip Osteoarthritis:
Methylation Profile of OA Cartilage. *Arthritis & Rheumatology*, *66*(9), 2450–2460. doi:
10.1002/art.38713

Schultz, A. H. (1930). The Skeleton of the Trunk and Limbs of Higher Primates. *Human Biology*, *2*(3),
303–438.

Schultz, A. H. (1937). Proportions, Variability and Asymmetries of the Long Bones of the Limbs and the
Clavicles in Man and Apes. *Human Biology*, *9*(3), 281–328.

Sharif, J., Endo, T. A., Toyoda, T., & Koseki, H. (2010). Divergence of CpG island promoters: A
consequence or cause of evolution? *Development, Growth & Differentiation*, *52*(6), 545–554. doi:
10.1111/j.1440-169X.2010.01193.x

Shelnutt, K. P., Kauwell, G. P. A., Gregory III, J. F., Maneval, D. R., Quinlivan, E. P., Theriaque, D. W.,
… Bailey, L. B. (2004). Methylenetetrahydrofolate reductase 677C→T polymorphism affects
DNA methylation in response to controlled folate intake in young women. *The Journal of*
Nutritional Biochemistry, *15*(9), 554–560. doi: 10.1016/j.jnutbio.2004.04.003

Simon, T. C., & Jeffries, M. A. (2017). The Epigenomic Landscape in Osteoarthritis. *Current*
Rheumatology Reports, *19*(6), 30. doi: 10.1007/s11926-017-0661-9

Singer, M., Kosti, I., Pachter, L., & Mandel-Gutfreund, Y. (2015). A diverse epigenetic landscape at
human exons with implication for expression. *Nucleic Acids Research*, gkv153. doi:
10.1093/nar/gkv153

Slieker, R. C., Bos, S. D., Goeman, J. J., Bovée, J. V., Talens, R. P., van der Breggen, R., … Heijmans, B.
T. (2013). Identification and systematic annotation of tissue-specific differentially methylated

regions using the Illumina 450k array. *Epigenetics & Chromatin*, *6*, 26. doi: 10.1186/1756-8935-6-26

Smith, R. W. A., Monroe, C., & Bolnick, D. A. (2015). Detection of Cytosine Methylation in Ancient DNA from Five Native American Populations Using Bisulfite Sequencing. *PLoS ONE*, *10*(5). doi: 10.1371/journal.pone.0125344

Smith, S. M., Garic, A., Berres, M. E., & Flentke, G. R. (2014). Genomic factors that shape craniofacial outcome and neural crest vulnerability in FASD. *Frontiers in Genetics*, *5*. doi: 10.3389/fgene.2014.00224

Smyth, G. K. (2004). Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Statistical Applications in Genetics and Molecular Biology*, *3*, Article3. doi: 10.2202/1544-6115.1027

Suter, C. M., Martin, D. I. K., & Ward, R. L. (2004). Germline epimutation of MLH1 in individuals with multiple cancers. *Nature Genetics*, *36*(5), 497–501. doi: 10.1038/ng1342

Suzuki, M. M., & Bird, A. (2008). DNA methylation landscapes: Provocative insights from epigenomics. *Nature Reviews Genetics*, *9*(6), 465–476. doi: 10.1038/nrg2341

Terzidis, I., Totlis, T., Papathanasiou, E., Sideridis, A., Vlasis, K., & Natsis, K. (2012). Gender and Side-to-Side Differences of Femoral Condyles Morphology: Osteometric Data from 360 Caucasian Dried Femori. *Anatomy Research International*, *2012*, e679658. doi: 10.1155/2012/679658

Tung, J., Zhou, X., Alberts, S. C., Stephens, M., & Gilad, Y. (2015). The genetic architecture of gene expression levels in wild baboons. *ELife*, *4*, e04729. doi: 10.7554/eLife.04729

Uebbing, S., Künstner, A., Mäkinen, H., Backström, N., Bolivar, P., Burri, R., … Ellegren, H. (2016). Divergence in gene expression within and between two closely related flycatcher species. *Molecular Ecology*, *25*(9), 2015–2028. doi: 10.1111/mec.13596

van Dongen, J., Ehli, E., Slieker, R., Bartels, M., Weber, Z., Davies, G., … Boomsma, D. (2014). Epigenetic Variation in Monozygotic Twins: A Genome-Wide Analysis of DNA Methylation in Buccal Cells. *Genes*, *5*(2), 347–365. doi: 10.3390/genes5020347

Vilgalys, T. P., Rogers, J., Jolly, C., Mukherjee, S., & Tung, J. (2018). *Evolution of DNA methylation in baboons*. doi: 10.1101/400093

Warner, L. R., Babbitt, C. C., Primus, A. E., Severson, T. F., Haygood, R., & Wray, G. A. (2009). Functional consequences of genetic variation in primates on tyrosine hydroxylase (TH) expression in vitro. *Brain Research*, *1288*, 1–8. doi: 10.1016/j.brainres.2009.06.086

Weber, M., Hellmann, I., Stadler, M. B., Ramos, L., Pääbo, S., Rebhan, M., & Schübeler, D. (2007). Distribution, silencing potential and evolutionary impact of promoter DNA methylation in the human genome. *Nature Genetics*, *39*(4), 457–466. doi: 10.1038/ng1990

Weksberg, R., Shuman, C., Caluseriu, O., Smith, A. C., Fei, Y.-L., Nishikawa, J., … Squire, J. (2002). Discordant KCNQ1OT1 imprinting in sets of monozygotic twins discordant for Beckwith-Wiedemann syndrome. *Human Molecular Genetics*, *11*(11), 1317–1325.

White, T., & Folkens, P. (2000). *Human osteology* (2nd ed.). San Diego: Academic Press.

Whitehead, A., & Crawford, D. L. (2006). Neutral and adaptive variation in gene expression. *Proceedings of the National Academy of Sciences of the United States of America*, *103*(14), 5425–5430. doi: 10.1073/pnas.0507648103

Young, M. D., Wakefield, M. J., Smyth, G. K., & Oshlack, A. (2010). Gene ontology analysis for RNA-
seq: Accounting for selection bias. *Genome Biology*, *11*, R14. doi: 10.1186/gb-2010-11-2-r14
Zeng, J., Konopka, G., Hunt, B. G., Preuss, T. M., Geschwind, D., & Yi, S. V. (2012). Divergent Whole-
Genome Methylation Maps of Human and Chimpanzee Brains Reveal Epigenetic Basis of
Human Regulatory Evolution. *The American Journal of Human Genetics*, *91*(3), 455–465. doi:
10.1016/j.ajhg.2012.07.024
Zhang, B., Zhou, Y., Lin, N., Lowdon, R. F., Hong, C., Nagarajan, R. P., … Wang, T. (2013). Functional
DNA methylation differences between tissues, cell types, and across individuals discovered using
the M&M algorithm. *Genome Research*, *23*(9), 1522–1540. doi: 10.1101/gr.156539.113
Zwetsloot, K. A., Laye, M. J., & Booth, F. W. (2009). Novel epigenetic regulation of skeletal muscle
myosin heavy chain genes. Focus on "Differential epigenetic modifications of histones at the
myosin heavy chain genes in fast and slow skeletal muscle fibers and in response to muscle
unloading." *AJP: Cell Physiology*, *297*(1), C1–C3. doi: 10.1152/ajpcell.00176.2009

1095

1096 **Figure Legends:**

1097

1098
1099 **Figure 1. Nonhuman Primate Sample Set Ages.** Boxes represent one standard deviation from the
1100 average age, and whiskers depict the full range of ages for each species. Baboons (n=28) are 16.90±5.02
1101 years, chimpanzees (n=4) are 11.31±1.87 years, macaques (n=10) are 14.75±2.65 years, marmosets (n=6)
1102 are 3.34±1.41 years, and vervets (n=10) are 9.31±10.30 years. Additional details can be found in Table
1103 S1.

1104

1105
1106 **Figure 2. Nonhuman Primate Morphological Measurements.** See Table S6 for a detailed description
1107 of these measurements.

1108

1109

Figure 3. Methylation Levels at Species-Specific DMPs with Δβ≥0.4 Identified in the Inter-Specific

Study. Heatmap depicting the DNA methylation levels (β values) of all species-specific DMPs with

average absolute Δβ values greater than 0.4 between each taxonomic group (x-axis) in all nonhuman

primate samples (n=58). The sex and age of each nonhuman primate are also provided (y-axis). Red

indicates higher methylation at a DMP, while blue indicates lower methylation at a DMP. The

dendrogram of all samples (y-axis) clusters individuals based on the similarity of their methylation

patterns. Samples cluster based on species-level taxonomic groupings and as predicted based on known

species phylogenetic histories.

1118

1119

Figure 4. Phylogeny Based on Average Species-Level Global Changes in Methylation. Observed

phylogenetic relationship among nonhuman primates when considering average species-level global

changes in methylation. This tree was constructed using the methylation levels for all finalized 39,802

filtered probes. We averaged the β values per probe within a species, used Euclidean distances to

calculate the difference between every two species, and estimated a neighbor joining tree using this

distance matrix. For the resulting tree, 1000 bootstraps were performed to determine confidence values

for each branch. The number provide at each node indicates the number of bootstrap replicates that

support it out of the 1000 performed.

1128

1129

Figure 5. Genome-Wide Methylation Levels Across *HOXD10* in Nonhuman Primates. Plot of the
methylation levels of significant DMPs across the *HOXD10* gene (hg19 chr2:176981492-176984670).
Plot shows the average β values for each DMP with error bars indicating 1 standard deviation in each
direction for each comparative group (teal = baboon, orange = chimpanzee, purple = macaque, pink =
marmoset, and light green = vervet). DMP chromosomal position in relation to the *HOXD10* gene is also
depicted. This gene is of interest because it has been found to be differential methylation in ancient and
modern hominin species (Gokhman et al., 2014). Of the sites depicted here, 5 DMPs were found to show
significant species-specific methylation in marmosets. Of the 5 species-specific DMPs in the *HOXD10*
gene of marmosets, 4 have Δβ between 0.2 and 0.3 (**) and 1 has a Δβ < 0.1 (*). See Table S13 for
additional information. In the *HOXD10* gene, the two exons are denoted with the thickest bars (exon 1:
hg19 chr2:176981561-176982306; exon 2: hg19 chr2:176983681-176983959), the UTRs are denoted
with bars of intermediate thickness (5'UTR: hg19 chr2:176981491-176981561; 3'UTR: hg19
chr2:176983959-176984670), and the one intron is denoted with the thinnest bar (intron 1: hg19
chr2:176982306-17698368).

1144

1145

Figure 6. Gene-Specific Methylation Levels Across *HOXD10* in Nonhuman Primates. Bar plot of
DNA methylation across the *HOXD10* gene (hg19 chr2:176981492-176984670), as well as upstream and
downstream several hundred bases (hg19 chr2:176980532-176985117). Bars depict the presence (tall

bar), partial presence (medium bar), or absence (low bar) of methylation at human derived CpG sites in
15 nonhuman primate samples – 3 baboons, 3 macaques, 3 vervets, 3 chimpanzees, and 3 marmosets.
While regular sequencing was very successful, bisulfite sequencing was less successful, with several
sequence reads uninterpretable. As such, nonhuman primate methylation data is only available for a
subset of the CpGs known in humans. Partial presence of methylation was called when sequencing
fluorescence peaks for cytosine and thymine were both present at a particular site and one was at least
half the size of the other. Overall, these data provide additional information regarding gene-specific
methylation levels across *HOXD10*. CpG sites that were also targeted by the EPIC array are highlighted
in yellow and include cg18115040 (chr2, position 176981328), cg25371634 (chr2, position 176981422),
cg13217260 (chr2, position 176981469), cg03918304 (chr2, position 176981654), cg17489939 (chr2,
position 176981919), cg26708100 (chr2, position 176983815), cg10393811 (chr2, position 176983927),
cg08992581 (chr2, position 176983949), and cg06005169 (chr2, position 176984634). See Table S18 and
Files S8-S9 for additional information.