1 **Title**
2 <u>Discovery of several thousand highly diverse circular DNA viruses</u>
3

Authors
Michael J. Tisza[1], Diana V. Pastrana[1], Nicole L. Welch[1], Brittany Stewart[1], Alberto Peretti[1], Gabriel J.
Starrett[1], Yuk-Ying S. Pang[1], Arvind Varsani[2,3], Siddharth R. Krishnamurthy[4], Patricia A. Pesavento[5], David
H. McDermott[6], Philip M. Murphy[6], Jessica L. Whited [7,8,9], Bess Miller [7,8], Jason M. Brenchley[10], Stephan P.
Rosshart[11], Barbara Rehermann[11], John Doorbar[12], Blake A. Ta'ala[13], Olga Pletnikova[14], Juan Troncoso[14],
Susan M. Resnick[15], Anca M. Segall[16], Christopher B. Buck[1]*
10
Affiliations
[1]Lab of Cellular Oncology, NCI, NIH, Bethesda, MD 20892-4263
[2]The Biodesign Center of Fundamental and Applied Microbiomics, School of Life Sciences, Center for
Evolution and Medicine, Arizona State University, 1001 S. McAllister Ave, Tempe, AZ 85287-5001, USA
[3]Structural Biology Research Unit, Department of Clinical Laboratory Sciences, University of Cape Town,
Rondebosch, 7701, Cape Town, South Africa.
[4]Metaorganism Immunity Section, Laboratory of Immune System Biology, NIAID, NIH, Bethesda, MD
20892-4263
[5]Department of Pathology, Microbiology, and Immunology, University of California, Davis, CA 95616,
USA.
[6]Molecular Signaling Section, Laboratory of Molecular Immunology, NIAID, Bethesda, Maryland, USA.
[7]Department of Orthopedic Surgery, Harvard Medical School, The Harvard Stem Cell Institute, Brigham
and Women's Hospital, 60 Fenwood Road, Boston, MA, 02115, USA.
[8]Broad Institute of MIT and Harvard, 7 Cambridge Center, Cambridge, MA, 02142, USA.
[9]Department of Stem Cell and Regenerative Biology, Harvard University, 7 Divinity Avenue, Cambridge,
MA, 02138, USA.
[10]Barrier Immunity Section, Lab of Viral Diseases, NIAID, NIH, MD 20892
[11]Immunology Section, Liver Diseases Branch, NIDDK, NIH, Bethesda, MD 20892-1800
[12]Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge, CB2 1QP, UK.
[13]Mililani Mauka Elementary, Mililani, HI, 96789, USA.
[14]Department of Pathology (Neuropathology), Johns Hopkins University School of Medicine, Baltimore,
MD, USA.
[15]Laboratory of Behavioral Neuroscience, National Institute on Aging, Baltimore, MD 21224, USA.
[16]Viral Information Institute & Department of Biology, San Diego State University, San Diego, CA, 92182-
4614, USA.
*Corresponding author: buckc@mail.nih.gov

Abstract
Virologists have posited the existence of millions of distinct viral species, but fewer than 9000 viral
species are catalogued in GenBank's RefSeq database. We selectively enriched for and amplified the
genomes of circular DNA viruses in over 70 animal samples, ranging from cultured soil nematodes to
human tissue specimens. Over 2500 complete circular genomes, each representing a new viral taxon,
were assembled, thoroughly annotated, and deposited in GenBank. The new genomes belong to dozens
of established and emerging viral families. Some of the genomes have unexpected gene compositions
that appear to be the result of recombination between ssDNA viruses and ssRNA viruses. In addition,
hundreds of circular DNA elements that do not encode any discernable similarities to previously
characterized sequences were identified. To characterize these "dark matter" sequences, we used an
artificial neural network to identify candidate viral capsid proteins and cell culture assays to validate that

some predicted capsid proteins formed virus-like particles. These data further the understanding of viral
sequence diversity and allow for more comprehensive analysis of the virosphere.
51
Main
There has been a rush to utilize massive parallel sequencing approaches to better understand the
complex microbial communities associated with humans and other animals. Although the bacterial
populations in these surveys have become increasingly recognizable[1], a substantial fraction of the reads
and *de novo* assembled contigs in many metagenomics efforts are binned as genetic "dark matter," with
no recognizable similarity to characterized sequences[2,3]. Some of this dark matter undoubtedly consists
of viral sequences, which have remained poorly characterized due to their enormous diversity[4]. Recent
efforts have shown that our understanding of viral diversity, even of viruses known to directly infect
humans, has been woefully incomplete[5]. To increase the power of future studies seeking to find
additional associations between microbes and host animal phenotypes, representative sequences for all
clades of the virosphere need be identified, annotated, and made publicly accessible.
Virus discovery has typically proven to be more difficult than discovery of cellular organisms.
Whereas all known cellular organisms encode conserved sequences (such as ribosomal RNAs) that can
readily be identified through sequence analysis, viruses, as a whole, do not have any universally
conserved sequence components[6,7]. Nevertheless, some success has been achieved in RNA virus
discovery by probing for the conserved sequences of their distinctive RNA-dependent RNA polymerase
or reverse transcriptase genes in metatranscriptomic data[8]. Also, many bacteriophages of the order
Caudovirales, such as the families Siphoviridae, Podoviridae, and Myoviridae, are extensively
characterized due to their and their hosts' culturability and their detectability using viral plaque assays[9-
11]. In contrast, many small DNA viruses are not easily cultured[12], use diverse genome replication
strategies, and typically lack polymerase genes[13]. An additional challenge is that small DNA viruses with
segmented genomes may have segments that do not encode recognizable homologs of known viral
genes. Therefore, small DNA viruses are particularly sparse in reference databases.
Despite the apparent challenges in detecting small DNA viruses, many have physical properties
that can be leveraged to facilitate their discovery. In contrast to the nuclear genomes of animals, many
DNA virus genomes have circular topology, which allows selective enrichment through rolling circle
amplification (RCA) methods. Further, the unique ability of viral capsids to protect nucleic acids from
nuclease digestion and to mediate the migration of the viral genome through ultracentrifugation
gradients or size exclusion columns allows physical isolation of viral genomes. In this study, we use
physical and topological features common to many DNA viruses to uncover a diverse array of complete
circular viral genomes in various human and non-human animal samples. A high-throughput
bioinformatics pipeline was developed to classify and annotate over 2500 candidate viral genomes.
84
Results
Virion enrichment and sequencing
We have previously developed methods for discovery of viruses (primarily polyomaviruses and
papillomaviruses) from skin swabs and complex tissue specimens[14]. The approach begins with
detergent-mediated cell disruption and nuclease digestion of non-encapsidated nucleic acid. This is
followed by ultracentrifugation of virions through Optiprep gradients. The purified virions are then
subjected to protein digestion, and the encapsidated DNA is amplified by random-primed RCA using
bacteriophage phi29 DNA polymerase, which selectively amplifies circular DNA templates and conducts
second strand synthesis on ssDNA. The amplified DNA is then deep-sequenced, and reads are *de novo*
assembled into contigs with SPAdes[15]. This approach was applied to a large set of human and animal
samples (Table S1). The samples sources were varied, but emphasis was placed on using a variety of
tissues from vertebrates and model organisms.

For the current study, we developed a bioinformatics pipeline, Cenote-Taker (a portmanteau of
cenote, a naturally occurring circular water pool, and *note-taker*), to identify and annotate *de novo*
assembled contigs with terminal direct repeats consistent with circular DNA molecules. In this pipeline,
putative circular sequences were queried against GenBank's nucleotide database using BLASTN to
remove circles with extensive nucleotide identity (>90% across any 500 bp window) to known
sequences. Sequences with >90% identity to previously reported viral sequences represented less than
1.5% of circular contigs. Open reading frames (ORFs) from remaining unidentified circular DNA
sequences >240 nucleotides (nt) in length were translated and used for RPS-BLAST queries of GenBank's
Conserved Domain Database. ORFs that did not yield E values better than 1×10^{-4} in RPS-BLAST were
subjected to BLASTP searches of viral sequences in GenBank's nr database[16-18]. ORFs that were not
confidently identified in BLAST searches were subjected to searches using HHBlits[19]. The results were
used to annotate and name each sequence in a human-readable genome map as well as a format
suitable for submission to GenBank. After checking the Cenote-Taker output of each genome, minor
revisions were made and files were submitted to GenBank (BioProjects PRJNA393166 and
PRJNA396064).
112
<u>Characterization of DNA viruses in animal metagenomes</u>
Of the novel circular sequences detected in the survey, 1844 encode genes with similarity to proteins of
ssDNA viruses and 55 encode genes with similarity to dsDNA viral proteins (Fig. 1A). The large majority
of genomes from this study are highly divergent from previously reported sequences (Fig. S1). The most
common taxon was the viral family *Microviridae*, a class of small bacteriophages, with 670 complete
genomes. This represents a dramatic expansion beyond the 459 non-redundant microvirus genomes
previously listed in GenBank (of which 44 were listed in the RefSeq database) (Fig. 1B). We discovered
562 genomes that had similarity to Circular Rep-Encoding Single-Stranded (CRESS) viruses that have yet
to be assigned into specific families, i.e. unclassified CRESS viruses (compared to the 85 listed in RefSeq).
Three hundred genomes displayed recognizable similarity to members of the *Circoviridae*, a class of
animal-tropic CRESS viruses, adding to the 427 non-redundant circovirus genomes previously listed in
GenBank (167 genomes were listed in the RefSeq database). Other genomes that were uncovered
represent *Anelloviridae* (n=170), *Inoviridae* (n=70), *Genomoviridae* (n=58), *Siphoviridae* (n=18),
unclassified phage (n=14), unclassified virus (n=6), *Geminiviridae* (n=8), *Podoviridae* (n=10), *Myoviridae*
(n=7), *Papillomaviridae* (n=4), and *CrAssphage* (n=2) (Fig. 1B).
It is difficult to assign a host to most of the viruses from this study due to their divergence from
known viral sequences. However, three viruses had exact matches to CRISPR spacers in bacterial
genomes (Siphoviridae sp. isolate ctcj11:Shewanella sp. W3-18-1, Inoviridae sp. isolate ctce6:
Shewanella baltica OS195, Microviridae sp. isolate ctbe523:Paludibacter propionicigenes WB4) and one
virus had an exact match to the CRISPR spacer of an archaeon (Caudovirales sp. isolate
cthg227:Methanobrevibacter sp. AbM4), implying that these organisms are infected by these viruses.
In addition to circular genomes with recognizable similarity to known viruses, 609 circular
contigs appeared to represent elements that lacked discernable similarity to known viruses (Fig. 1A).
136
The vast majority of the *de novo* assembled circular genomes were <10 kb in length (Fig S2). This
reflects the tendency of ssDNA genomes to be small (the largest reported is ~10 kb)[7]. However, this
observation may also be due to the greater difficulty in *de novo* assembling complete gapless contigs for
larger genomes, as large genomes require more reads for uninterrupted coverage and are more likely to
contain difficult-to-assemble regions. Further, RCA, which we used to amplify viral genomes, may
preferentially amplify smaller circular genomes. Despite these technical obstacles, our detection of a
new tailed bacteriophage with a 419kb genome (Myoviridae sp. isolate ctbc4, GenBank Accession:

MH622943), along with 45 other >10 kb circular sequences (Fig S2), indicates that the methods used for
the current work can detect large viral genomes.
There has been a recent renewal of interest in the hypothesis that viruses may be etiologically
associated with degenerative brain diseases, such as Alzheimer's disease[20,21]. Samples of brain tissue
from individuals who died of Alzheimer's disease (n=6) and other forms of dementia (n=6) were
subjected to virion enrichment and deep sequencing. Although complete or partial genomes of known
papillomaviruses, Merkel cell polyomavirus, and/or anelloviruses were observed in some samples
(Supplementary File S1), no novel complete viral genomes were recovered (Table S1). No recognizably
virus-like sequences were observed in a follow-up RNAseq analysis of the samples. It is likely that the
observed viral DNA sequences represent viral particles from blood vessels in these tissues or from
environmental sources. It is important to note that the methods used in the present study focused on
virion enrichment and thus would not detect latent or past viral infections.
156
<u>Assignment of recognizable sequences to clusters</u>
Sequence similarity network analyses were used to display related groups of genes from viral genomes
(Fig. 2, Fig S3). Protein sequences from RefSeq (ssDNA viruses) and from viral genomes discovered here
were aligned pairwise (each sequence was compared to all other sequences) using EFI-EST[22]. The output
was visualized with Cytoscape[23], with each node representing an amino acid sequence. Nodes
(sequences) with significant amino acid identity (E value $<1\times10^{-5}$) were connected with lines. This
method generated dozens of clusters of related sequences, one of which is displayed in Fig. 2. Note that
not all sequences in a given cluster necessarily have a significant similarity to all other sequences in that
cluster. Rather, many sequences have a second- or third-degree relatedness to one another. The nodes
added by this study populate and provide greater connectivity in the networks for many viral protein
families.
168
<u>Large CRESS viruses</u>
Although no single family of viruses accounts for the majority of genomes in this study, these results
expand the knowledge of the vast diversity of CRESS viruses, which appear to be ubiquitous among
eukaryotes[24-27]. The CRESS virus supergroup, which is primarily defined by a characteristic superfamily 3
DNA helicase (Rep) protein, encompasses the recognized families *Circoviridae*, *Geminiviridae*, and many
others. Characterized CRESS viruses have small icosahedral virions (<25 nm in diameter) with a simple
T=1 geometry[28]. This capsid architecture likely limits genome size, as nearly all previously reported
CRESS genomes/genome segments are under 3.5 kb. Exceptions to this size rule are bacilladnaviruses,
which have 4.5 - 6 kb genomes[29] and cruciviruses, which have 3.5 - 5.5 kb genomes[30]. Interestingly, the
genomes of these larger CRESS viruses encode capsid genes that appear to have been acquired
horizontally from RNA viruses[31]. In our dataset, eight CRESS-like circular genomes exceed 6 kb in length
(Fig. 3C-J), making them the largest CRESS genomes ever identified. Further, this study's large CRESS
genomes are apparently attributable to several independent acquisitions of capsid genes from other
taxa and/or capsid gene duplication events.
Notably, a large CRESS genome (CRESS virus isolate ctdh33, associated with rhabditid
nematodes that were serially cultured from a soil sample) encoded three separate genes with predicted
homology to the capsid gene of satellite tobacco necrosis virus (STNV, an RNA virus satellite) (Fig 3C).
The STNV capsid protein, in turn, exhibits similarity to geminivirus capsid proteins[13]. The three predicted
STNV capsid homologs in the nematode virus are highly divergent from one another, with only 28-30%
amino acid similarity. A possible explanation for this observation is that the capsid gene array is the
result of gene duplication events. Similar STNV capsid-like genes have previously been observed as a
single copy in unclassified <3 kb CRESS viruses[32] and several additional examples of such genes were

detected in the present study. Network analysis of new sequences suggests a possible model in which
the STNV capsid gene originated from an ancestral CRESS DNA virus (Fig S4).
To further characterize the soil nematode virus, the three candidate capsid proteins were
overexpressed in combination in human-derived 293TT cells[33]. Particles apparently composed of all
three capsid proteins were able to sediment in Optiprep gradients. Further, analysis by negative stain
electron microscopy (EM) revealed ~50 nm particles (Fig S5A). Interestingly these particles were found
to contain nonspecific nuclease-resistant RNA (Fig S5B). This supports the concept that CRESS virus
capsid proteins might theoretically be re-targeted to packing RNA virus genomes or vice versa.
CRESS genomes ctba10, ctcc19, ctbj26, ctcd34, and ctbd1037 (ranging from 3.5 - 6.2 kb in
length) also each encode two divergent capsid gene homologs (Fig. 3D, Fig S6). Single genomes encoding
multiple capsid genes with related but distinct amino acid sequences have been observed in RNA
viruses[34], but we believe that this is the first time it has been reported in ssDNA viruses.
Two related large CRESS viruses (ctdb796 and ctce741) encode capsid proteins similar to those
of bacilladnaviruses (Fig 3G, I). Interestingly, the Rep genes of the two viruses do not show detectable
similarity to known bacilladnavirus Reps and are instead similar to the Reps of certain unclassified CRESS
viruses, suggesting that CRESS ctdb796 and CRESS ctce741 are representatives of a new hybrid CRESS
virus family. Two other CRESS virus genomes (ctca5 and ctgh4) encode capsid genes that show amino
acid similarity to distinct groups of icosahedral T=3 ssRNA virus capsids[35] (tombus- and tombus-like
viruses) (Fig 3A,F, Fig. S7). Further, a 6.6 kb CRESS virus (ctbd466) (Fig 3H) was found to encode a gene
with similarity to the capsid region of the polyprotein of a newly described ssRNA virus, ciliovirus[35,36].
Protein fold predictor Phyre2[37] showed a top hit (58% confidence) for the capsid protein of a norovirus
(ssRNA virus with T=3 icosahedral capsid). Taken together, these findings suggest multiple instances of
chimerization between ssDNA and ssRNA viruses.
Two CRESS genomes (ctbb30 and ctbc27) from separate Rhesus macaque stool samples
combine CRESS Rep genes with several genes specific to *Inoviridae* (a family of circular ssDNA
bacteriophages), including inovirus-like capsid genes, which encode proteins that form a filamentous
virion (Fig 3B, J). Other inovirus-like genes encoded in these genomes include homologs of zonular
occludens toxin (ZOT, a packaging ATPase) and RstB (a DNA-binding protein required for host genome
integration)[38] (Fig. 3). Curiously, TBLASTX searches using CRESS ctbb30 and ctbc27 sequences yielded
large segments of similarity to various bacterial chromosomes (e.g., GenBank accession numbers
AP012044 and AP018536), presumably representing integrated prophages. This suggests that CRESS
ctbb30 and ctbc27 represent a previously undescribed prokaryote-infecting branch of the CRESS virus
supergroup.
224
Network analysis of genetic "dark matter"
We defined potential viral "dark matter" in the survey as circular contigs with no hits with E values
$<1\times10^{-5}$ in BLASTX searches of a database of viral and plasmid proteins. To categorize the predicted
proteins of the 609 dark matter elements, we used pairwise comparison with EFI-EST. A great majority
of translated gene sequences could be categorized into dark matter protein clusters (DMPCs) containing
four or more members (Fig. 4A). Further, groups of related dark matter elements (i.e. dark matter
genome groups (DMGGs)), much like viral families, could be delineated by the presence of a conserved,
group-specific marker gene. For example, DMPC1 can be thought of as the marker gene for DMGG1.
Certain DMPCs tend to co-occur on the same DMGG. For instance, DMPC7 and DMPC17 ORFs are always
observed in genomes with a DMPC1 ORF (i.e., DMGG1) (Fig. 4B). This *pro tempore* categorization
method is useful for visualizing the data, but we stress that is not necessarily taxonomically definitive.
Powerful new bioinformatic methods for detecting distant homology rely on the observation
that protein structures are typically more conserved than primary amino acid sequences. HHpred, for
example, compares the predicted fold of a query sequence to a library of known protein structures[39].

Alignments were prepared for each major DMPC to identify conserved residues and each alignment was
used for HHPRED queries. This approach identified that ten DMPCs are likely viral capsid proteins and 11
are rolling circle replicases (Fig. 4A).
While most of the circular dark matter in the survey could be characterized using these
methods, dark matter contigs represent a small remaining fraction in some samples (Fig. S9).
244
<u>Functional analysis of candidate capsid proteins by expression in cell culture</u>
In contrast to viral genes with conserved enzymatic functions, sequences of the capsid genes are
often poorly conserved, even within a given viral family[40]. Moreover, it appears that capsid proteins
have arisen repeatedly through capture and modification of different host cell proteins[31]. This makes it
challenging to detect highly divergent capsid proteins using alignment-based approaches. We therefore
turned to an alignment-independent approach known as iVireons, an artificial neural network trained by
comparing alignment-independent variables between a large set of known viral structural proteins and
known non-structural proteins[41] (https://vdm.sdsu.edu/ivireons/). As an example of the approach,
iVireons scores for DMPCs associated with DMGG1 are shown in Fig. 4C. Other sets of iVireons scores
can be seen in Fig. S8.
Of the 17 DMGGs for which HHPRED did not identify capsid genes, iVireons predicted that ten
contain at least one DMPC predicted to encode some type of virion structural protein (median score of
cluster >0.70). This allowed us to generate the testable hypothesis that some of these predicted
structural proteins would form virus-like particles (VLPs) if expressed in cell culture.
A subset of predicted capsid genes were expressed in *E. coli* and subjected to size exclusion
chromatography. Electron microscopic analysis of the proteins showed distinct, albeit irregular, VLPs for
three of these predicted capsid genes, whereas a negative control protein did not form particles (Fig. 5).
The results suggest that, in multiple cases, iVireons correctly predicted the identity of viral capsid
proteins.
264
Discussion
Massive parallel DNA sequencing surveys characterizing microbial communities typically yield a
significant fraction of reads that cannot be mapped to known genes. The present study sought to
provide the research community with an expanded catalog of viruses with circular DNA genomes
associated with humans and animals, as well as a means to characterize future datasets. We hope that
the availability of this expanded viral sequence catalog will facilitate future investigation into
associations between viral communities and disease states.
At present, GenBank's RefSeq database includes complete sequences for 8112 viral genomes,
most of which fit into 131 families recognized by the International Committee on Taxonomy of Viruses
(ICTV)[42]. The current study, which focused on circular DNA viruses with detergent-resistant capsids,
found 2508 new complete circular genomes which contribute new information and understanding to a
broad range of established and emerging taxa.
The predominance of small (<10kb) genomes with remote similarity to ssDNA viruses in our
dataset probably reflects the underappreciated ubiquity of small ssDNA viruses in nature. These viruses
are often overlooked in studies that only characterize sequences that are closely related to reference
genomes. In addition, ssDNA is not detected by some current DNA sequencing technologies unless
second-strand synthesis (such as the RCA approach used in the current study) is conducted.
While many of the viruses discovered in this study appear to be derived from prokaryotic
commensals, it is important to note that bacteriophages can contribute to human and animal diseases
by transducing toxins, antimicrobial resistance proteins, or genes that alter the physiology of their
bacterial hosts[43]. Furthermore, interaction between animal immune systems and bacteriophages
appears to be extensive[44].

Over 100 distinct human anellovirus sequences were found in human blood. Anelloviruses have yet to be causally associated with any human disease, but this study indicates that we are likely still just scratching the surface of the sequence diversity of human anelloviruses. It will be important to fully catalog this family of viruses to address the field's general assumption that they are harmless.

Several of the CRESS viruses detected in this study are larger than any other CRESS DNA virus genomes that have been described previously. In some cases, the larger size of these genomes may have been enabled by a process involving capsid gene duplication events. Further, CRESS virus acquisition of T=3 capsids from ssRNA *Nodaviridae* and *Tombusviridae* families has been previously suggested as the origin of bacilladnaviruses and cruciviruses, respectively. We present evidence of additional recombination events between CRESS viruses and ssRNA viruses, ssRNA satellites, and ssDNA bacteriophages. These data suggest that CRESS viruses are at the center of a tangled evolutionary history of viruses in which genomes change not just via gradual point mutations but also through larger scale recombination and chimerization events.

It is likely that some dark matter sequences detected in this study share a common ancestor with known viruses but are too divergent to retain discernable sequence similarity. In some cases, the dark matter circles may represent a more divergent segment of a virus with a multipartite genome. Alternatively, some of these sequences likely represent entirely new viral lineages that have not previously been recognized.

This study suggests that, whether they are in our communities, in our homes, on our dinner tables, or in us, there are likely hordes of undiscovered viruses still waiting to be seen.

Methods

Sample collection and sequencing

De-identified human swabs and tissue specimens were collected under the approval of various Institutional Review Boards (Supplemental Table 1). Animal tissue samples were collected under the guidance of various Animal Care and Use Committees.

Nematodes were cultured out of soil samples collected in Bethesda, Maryland, USA on OP50-Seeded NGM-lite plates (C. elegans kit, Carolina Biological Supply).

Viral particles were concentrated by subjecting nuclease-digested detergent-treated lysate to ultracentrifugation over an Optiprep step gradient, as previously described https://ccrod.cancer.gov/confluence/display/LCOTF/Virome[14]. DNA from individually collected fractions of the gradient was amplified by RCA using phi29 polymerase (TempliPhi, GE Health Sciences). While we expected most viral particles to travel to the middle of the gradient based on previous experiments, RCA was conducted on individual fractions spanning the gradient, in an attempt to detect viruses with different biophysical properties. Pooled fractions were sequenced with Illumina technology on either MiSeq or NextSeq500 sequencers. Contigs were assembled using SPAdes with the 'plasmid' setting. Circularity was confirmed by assessing assembly graphs using Bandage[45].

Analysis of brain samples

Brain samples were initially analyzed by Optiprep gradient purification, RCA amplification, and deep sequencing, as described above. JC polyomavirus, which has previously been reported in brain samples[46], can display high buoyancy in Optiprep gradients[47]. Fractions from near the top of the Optiprep gradient were subjected to an alternative method of virion enrichment using microcentrifuge columns (Pierce) packed with 2 ml of Sepharose 4B Bead suspension exchanged into PBS. Fractions were

335   clarified at 5000 x g for 1 minute, and 200 µl of clarified extract was loaded onto the gel bed. The
336   column was spun at 735 x g and the eluate was digested with proteinase K, ethanol-precipitated, and
337   subjected to RCA. No additional viral sequences were detected by this method.
338
339   The brain samples were also subjected to confirmatory analysis by RNA sequencing. RNA was extracted
340   with Qiagen Lipid Tissue RNeasy Mini Kit and subjected to human ribosomal RNA depletion with Thermo
341   RiboMinus. The library was prepared with NEBNext Ultra™ II Directional RNA Library Prep Kit for Illumina
342   and subjected to massive parallel sequencing on the Illumina HiSeq platform.
343
<u>Cenote-Taker, Virus Discovery and Annotation Pipeline</u>
Cenote-Taker, a bioinformatics pipeline written for this project, was used for collection and detailed
annotation of each circular sequence. The flow of the program can be described as follows:
(1) Removes contigs (assembled with SPAdes) smaller than 1000 nts
(2) Predicts which contigs are circular based on overlapping ends
(3) Determines whether circular contig has any ORFs of 80 AA or larger or else discards sequence
(4) Uses BLASTN against GenBank 'nt' to disregard any circular sequences that are >90% identical to
known sequences across a >500 bp window
(5) Uses Circlator [48] to rotate circular contigs so that a non-intragenic start codon of one of the ORFs will
be the wrap point
(6) Uses BLASTX against a custom virus + plasmid database (derived from GenBank 'nr' and RefSeq) to
attempt to assign the circular sequence to a known taxon
(7) Translates each ORF of 80 AA or larger
(8) Uses RPS-BLAST to predict function of each ORF by aligning to known NCBI Conserved Domains
(9) Generates a tbl file of RPS-BLAST results
(10) Takes ORFs without RPS-BLAST hits and queries the GenBank 'nr viral' database with BLASTP
(11) Generates a tbl file of BLASTP results
(12) Takes ORFs without any BLASTP hits and queries HHblits (databases: uniprot20, pdb70, scop70,
pfam_31, NCBI_CD)
(13) Generates a tbl file of HHblits results
(14) Complies with a GenBank request to remove annotations for ORFs-within-ORFs that do not contain
conserved sequences
(15) Combines all tbl files into a master tbl file
(16) Generates a name for each virus based on taxonomic results and nature of sample
(17) Generates properly formatted fsa and tbl files in a separate directory
(18) Uses tbl2asn to make gbf, val, and sqn files
370
371   A 'beta version' of Cenote-Taker is available at the GitHub website:
372   (https://github.com/mtisza1/Cenote-Taker). This GitHub project is intended to demonstrate the steps
373   used to analyze the described sequences. Work is underway to make the program portable for general
374   use in other high-performance computing environments. Interested parties may contact the authors
375   about Cenote-Taker.
376
377   This work utilized the computational resources of the NIH HPC Biowulf cluster. (http://hpc.nih.gov).
378   Genome maps were drawn using MacVector 16.

Anelloviruses

Analysis of linear contigs in the survey found many instances of recognizable viral sequences. One noteworthy example were anelloviruses, where many contigs terminated near the GC-rich stem-loop structure that is thought to serve as the origin of replication. This segment of the anellovirus genome is presumably incompatible with the short read deep sequencing technologies used in this study. Nearly complete anellovirus genomes, defined as having a complete ORF1 gene and at least 10-fold coverage, were deposited in GenBank (Table S1).

Expressing Potential Viral Structural Proteins in human 293TT cells

Codon-modified expression constructs encoding the three predicted STNV-like capsid proteins from the nematode virus (CRESS virus isolate ctdh33) were designed according to a previously reported "as different as possible" algorithm https://ccrod.cancer.gov/confluence/display/LCOTF/Pseudovirus+Production. 293TT cells were transfected with combinations of STNV capsid-like protein expression constructs for roughly 48 hours. Cells were lysed in a small volume of PBS with 0.5% Triton X-100 or Brij-58 and Benzonase DNase/RNase (Sigma). After several hours of maturation at neutral pH, the lysate was clarified at 5000 x g for 10 min. The clarified lysate was loaded onto a 27-33-39% Optiprep gradient in PBS with 0.8 M NaCl. Gradient fractions were collected by bottom puncture of the tube and screened by PicoGreen nucleic acid stain (Invitrogen), BCA, and SDS-PAGE analysis. Electron microscopic analysis was then performed.

Expressing Potential Viral Structural Proteins in *E. coli*

Several genes that were identified by iVireons as being potential viral structural proteins were cloned into plasmids with a T7 polymerase-responsive promoter. Plasmids were transfected into T7 Express lysY/I^q *E. coli*, which express T7 polymerase under the induction of IPTG. Bacteria were grown at 37ºC in LB broth until OD600 = 0.5. Flasks were cooled to room temperature, IPTG was added to 1 mM, and cultures were shaken at room temperature for approximately 16 hours. Cells were then pelleted for immediate processing.
Total protein was extracted with a BPER (Pierce) and nuclease solution. Then, virion-sized particles were enriched from the clarified lysate using size exclusion chromatography with 2% agarose beads https://ccrod.cancer.gov/confluence/display/LCOTF/GelFiltration. Fractions were analyzed using Coomassie-stained SDS-PAGE gels for presence of a unique band corresponding to the expressed protein. Fractions of interest were analyzed using negative stain electron microscopy.

Electron Microscopy

Five µl samples were adsorbed onto a carbon-deposited copper grid for one minute. Sample was then washed 5 times on water droplets then stained with 0.5% uranyl acetate for 1 second. The negatively stained samples were examined on a FEI Tecnai T12 transmission electron microscope.

Sequencing

Illumina sequencing was conducted at the CCR Genomics Core at the National Cancer Institute, NIH, Bethesda, MD 20892.

Acknowledgements

426
427 **References**
428
429 1   Lloyd-Price, J. *et al.* Strains, functions and dynamics in the expanded Human
430   Microbiome Project. *Nature* **550**, 61-66, doi:10.1038/nature23889 (2017).
431 2   Krishnamurthy, S. R. & Wang, D. Origins and challenges of viral dark matter. *Virus Res*
432   **239**, 136-142, doi:10.1016/j.virusres.2017.02.002 (2017).
433 3   Oh, J. *et al.* Biogeography and individuality shape function in the human skin
434   metagenome. *Nature* **514**, 59-64, doi:10.1038/nature13786 (2014).
435 4   Simmonds, P. *et al.* Consensus statement: Virus taxonomy in the age of metagenomics.
436   *Nat Rev Microbiol* **15**, 161-168, doi:10.1038/nrmicro.2016.177 (2017).
437 5   Pastrana, D. V. *et al.* Metagenomic Discovery of 83 New Human Papillomavirus Types
438   in Patients with Immunodeficiency. *mSphere* **3**, doi:10.1128/mSphereDirect.00645-18
439   (2018).
440 6   O'Leary, N. A. *et al.* Reference sequence (RefSeq) database at NCBI: current status,
441   taxonomic expansion, and functional annotation. *Nucleic Acids Res* **44**, D733-745,
442   doi:10.1093/nar/gkv1189 (2016).
443 7   Brister, J. R., Ako-Adjei, D., Bao, Y. & Blinkova, O. NCBI viral genomes resource.
444   *Nucleic Acids Res* **43**, D571-577, doi:10.1093/nar/gku1207 (2015).
445 8   Shi, M. *et al.* Redefining the invertebrate RNA virosphere. *Nature*,
446   doi:10.1038/nature20167 (2016).
447 9   Pope, W. H. *et al.* Whole genome comparison of a large collection of
448   mycobacteriophages reveals a continuum of phage genetic diversity. *Elife* **4**, e06416,
449   doi:10.7554/eLife.06416 (2015).
450 10   Grose, J. H. & Casjens, S. R. Understanding the enormous diversity of bacteriophages:
451   the tailed phages that infect the bacterial family Enterobacteriaceae. *Virology* **468-470**,
452   421-443, doi:10.1016/j.virol.2014.08.024 (2014).
453 11   Grose, J. H., Jensen, G. L., Burnett, S. H. & Breakwell, D. P. Genomic comparison of 93
454   Bacillus phages reveals 12 clusters, 14 singletons and remarkable diversity. *BMC
455   Genomics* **15**, 855, doi:10.1186/1471-2164-15-855 (2014).
456 12   Bedell, M. A. *et al.* Amplification of human papillomavirus genomes in vitro is dependent
457   on epithelial differentiation. *J Virol* **65**, 2254-2260 (1991).
458 13   Koonin, E. V., Dolja, V. V. & Krupovic, M. Origins and evolution of viruses of eukaryotes:
459   The ultimate modularity. *Virology* **479-480**, 2-25, doi:10.1016/j.virol.2015.02.039 (2015).
460 14   Peretti, A., FitzGerald, P. C., Bliskovsky, V., Buck, C. B. & Pastrana, D. V. Hamburger
461   polyomaviruses. *The Journal of general virology* **96**, 833-839, doi:10.1099/vir.0.000033
462   (2015).
463 15   Bankevich, A. *et al.* SPAdes: a new genome assembly algorithm and its applications to
464   single-cell sequencing. *J Comput Biol* **19**, 455-477, doi:10.1089/cmb.2012.0021 (2012).
465 16   Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. Basic local alignment
466   search tool. *J Mol Biol* **215**, 403-410, doi:10.1016/S0022-2836(05)80360-2 (1990).
467 17   Marchler-Bauer, A. & Bryant, S. H. CD-Search: protein domain annotations on the fly.
468   *Nucleic Acids Res* **32**, W327-331, doi:10.1093/nar/gkh454 (2004).
469 18   Marchler-Bauer, A. *et al.* CDD: NCBI's conserved domain database. *Nucleic Acids Res*
470   **43**, D222-226, doi:10.1093/nar/gku1221 (2015).
471 19   Remmert, M., Biegert, A., Hauser, A. & Söding, J. HHblits: lightning-fast iterative protein
472   sequence searching by HMM-HMM alignment. *Nat Methods* **9**, 173-175,
473   doi:10.1038/nmeth.1818 (2011).

20 Itzhaki, R. F. *et al.* Microbes and Alzheimer's Disease. *J Alzheimers Dis* **51**, 979-984,
doi:10.3233/JAD-160152 (2016).
21 Eimer, W. A. *et al.* Alzheimer's Disease-Associated beta-Amyloid Is Rapidly Seeded by
Herpesviridae to Protect against Brain Infection. *Neuron* **99**, 56-63 e53,
doi:10.1016/j.neuron.2018.06.030 (2018).
22 Gerlt, J. A. *et al.* Enzyme Function Initiative-Enzyme Similarity Tool (EFI-EST): A web
tool for generating protein sequence similarity networks. *Biochim Biophys Acta* **1854**,
1019-1037, doi:10.1016/j.bbapap.2015.04.015 (2015).
23 Su, G., Morris, J. H., Demchak, B. & Bader, G. D. Biological network exploration with
Cytoscape 3. *Curr Protoc Bioinformatics* **47**, 8.13.11-24,
doi:10.1002/0471250953.bi0813s47 (2014).
24 Krupovic, M., Ghabrial, S. A., Jiang, D. & Varsani, A. Genomoviridae: a new family of
widespread single-stranded DNA viruses. *Arch Virol* **161**, 2633-2643,
doi:10.1007/s00705-016-2943-3 (2016).
25 Zerbini, F. M. *et al.* ICTV Virus Taxonomy Profile: Geminiviridae. *The Journal of general*
virology **98**, 131-133, doi:10.1099/jgv.0.000738 (2017).
26 Rosario, K. *et al.* Revisiting the taxonomy of the family Circoviridae: establishment of the
genus Cyclovirus and removal of the genus Gyrovirus. *Arch Virol* **162**, 1447-1463,
doi:10.1007/s00705-017-3247-y (2017).
27 Varsani, A. & Krupovic, M. Smacoviridae: a new family of animal-associated single-
stranded DNA viruses. *Arch Virol* **163**, 2005-2015, doi:10.1007/s00705-018-3820-z
(2018).
28 Khayat, R. *et al.* The 2.3-angstrom structure of porcine circovirus 2. *J Virol* **85**, 7856-
7862, doi:10.1128/JVI.00737-11 (2011).
29 Tomaru, Y. *et al.* Isolation and characterization of a single-stranded DNA virus infecting
Chaetoceros lorenzianus Grunow. *Appl Environ Microbiol* **77**, 5285-5293,
doi:10.1128/aem.00202-11 (2011).
30 Quaiser, A., Krupovic, M., Dufresne, A., Francez, A. J. & Roux, S. Diversity and
comparative genomics of chimeric viruses in Sphagnum-dominated peatlands. *Virus*
Evol **2**, vew025, doi:10.1093/ve/vew025 (2016).
31 Kazlauskas, D. *et al.* Evolutionary history of ssDNA bacilladnaviruses features horizontal
acquisition of the capsid gene from ssRNA nodaviruses. *Virology* **504**, 114-121,
doi:10.1016/j.virol.2017.02.001 (2017).
32 Kraberger, S. *et al.* Characterisation of a diverse range of circular replication-associated
protein encoding DNA viruses recovered from a sewage treatment oxidation pond. *Infect*
Genet Evol **31**, 73-86, doi:10.1016/j.meegid.2015.01.001 (2015).
33 Buck, C. B., Pastrana, D. V., Lowy, D. R. & Schiller, J. T. Efficient intracellular assembly
of papillomaviral vectors. *J Virol* **78**, 751-757 (2004).
34 Agranovsky, A. A., Lesemann, D. E., Maiss, E., Hull, R. & Atabekov, J. G. "Rattlesnake"
structure of a filamentous plant RNA virus built of two capsid proteins. *Proc Natl Acad*
Sci U S A **92**, 2470-2473 (1995).
35 Makino, D. L., Larson, S. B. & McPherson, A. The crystallographic structure of Panicum
Mosaic Virus (PMV). *J Struct Biol* **181**, 37-52, doi:10.1016/j.jsb.2012.10.012 (2013).
36 Greninger, A. L. & DeRisi, J. L. Draft Genome Sequences of Ciliovirus and Brinovirus
from San Francisco Wastewater. *Genome Announc* **3**, doi:10.1128/genomeA.00651-15
(2015).
37 Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. The Phyre2 web
portal for protein modeling, prediction and analysis. *Nature protocols* **10**, 845-858,
doi:10.1038/nprot.2015.053 (2015).
38 Falero, A. *et al.* DNA binding proteins of the filamentous phages CTXphi and VGJphi of
Vibrio cholerae. *J Bacteriol* **191**, 5873-5876, doi:10.1128/JB.01206-08 (2009).

39   Zimmermann, L. *et al.* A Completely Reimplemented MPI Bioinformatics Toolkit with a
New HHpred Server at its Core. *J Mol Biol* **430**, 2237-2243,
doi:10.1016/j.jmb.2017.12.007 (2018).
40   Buck, C. B. *et al.* The Ancient Evolutionary History of Polyomaviruses. *PLoS Pathog* **12**,
e1005574, doi:10.1371/journal.ppat.1005574 (2016).
41   Seguritan, V. *et al.* Artificial neural networks trained to detect viral and phage structural
proteins. *PLoS Comput Biol* **8**, e1002657, doi:10.1371/journal.pcbi.1002657 (2012).
42   King, A. M. Q. *et al.* Changes to taxonomy and the International Code of Virus
Classification and Nomenclature ratified by the International Committee on Taxonomy of
Viruses (2018). *Arch Virol*, doi:10.1007/s00705-018-3847-1 (2018).
43   Waldor, M. K. & Mekalanos, J. J. Lysogenic conversion by a filamentous phage
encoding cholera toxin. *Science* **272**, 1910-1914 (1996).
44   Hodyra-Stefaniak, K. *et al.* Mammalian Host-Versus-Phage immune response
determines phage fate in vivo. *Sci Rep* **5**, 14802, doi:10.1038/srep14802 (2015).
45   Wick, R. R., Schultz, M. B., Zobel, J. & Holt, K. E. Bandage: interactive visualization of
de novo genome assemblies. *Bioinformatics* **31**, 3350-3352,
doi:10.1093/bioinformatics/btv383 (2015).
46   Chalkias, S. *et al.* ViroFind: A novel target-enrichment deep-sequencing platform reveals
a complex JC virus population in the brain of PML patients. *PLoS One* **13**, e0186945,
doi:10.1371/journal.pone.0186945 (2018).
47   Geoghegan, E. M. *et al.* Infectious Entry and Neutralization of Pathogenic JC
Polyomaviruses. *Cell Rep* **21**, 1169-1179, doi:10.1016/j.celrep.2017.10.027 (2017).
48   Hunt, M. *et al.* Circlator: automated circularization of genome assemblies using long
sequencing reads. *Genome Biol* **16**, 294, doi:10.1186/s13059-015-0849-0 (2015).
549

Figure 1: Novel viruses associated with animal samples

Gross characterization of viruses discovered in this project compared to NCBI RefSeq virus database entries. (A) Pie chart representing the number of viral genomes in broad categories. (B) Bar graph showing the number of new representatives of known viral families. Note that genomes in this study were assigned taxonomy based on at least one region with a BLASTX hit with an E value $<1\times10^{-5}$, suggesting commonality with a known viral family. Some may ultimately be characterized as representatives of novel families

Figure 2: Network analysis of a putative category of CRESS capsid proteins

EFI-EST softwar[22] was used to perform "all against all" alignments of all RefSeq ssDNA virus proteins and all predicted proteins identified in this study. The figure shows one example of a cluster that encompasses a class of known CRESS virus capsid proteins. Additional clusters encompassing other known viral proteins are shown in Fig. S3.

Geminiviridae coat protein

Unclassified CRESS virus capsid protein

● RefSeq
● This study

Figure 3: Genome maps of large CRESS DNA genomes
CRESS-like 'Rep' genes are displayed in yellow-orange, virion structural genes shown in green, other common viral genes shown in pink, other genes in grey.

Figure 4: Dark matter analysis

(A) Sequence similarity network analysis for genes from dark matter circular sequences (minimum cluster size = 4). Clusters are colored based on assigned dark matter genome group (DMGG). Structural predictions from HHPRED are indicated (>85% probability). *Rep* = rolling circle replicases typical of CRESS viruses or ssDNA plasmids. *Capsid* = single-jellyroll capsid protein. *Attachment* = cell attachment proteins typified by Inoviridae. *DNA-Binding* = DNA-binding domain. *PLA2* = phospholipase A2. *FtsL* = FtsL Cell division protein. Clusters that contain a representative protein that was successfully expressed as a virus-like particle are outlined by a dashed rectangle (See Fig. 5). (B) Maps of three examples of DMGG1 with DMPCs labeled (linearized for display). (C) DMGG1 iVireons 'structure' score summary by protein cluster. Scores range from -1 (unlikely to be a virion structural protein) to 1 (likely to be a virion structural protein). Additional iVireons score summaries can be found in Fig. S8.

A

DMGG1 · DMGG2 · DMGG3 · DMGG4 · DMGG5 · DMGG6 · DMGG7 · DMGG8 · DMGG9 · DMGG10 · DMGG11 · DMGG12 · DMGG13 · DMGG14 · DMGG15 · DMGG16 · DMGG17 · DMGG18 · DMGG19 · DMGG20 · DMGG21 · DMGG22 · DMGG23 · DMGG24 · DMGG25 · DMGG26 · DMGG27

DMPC1, DMPC2, DMPC7, DMPC9, DMPC10, *Rep*, DMPC11, *DNA-Binding*, *Capsid*, DMPC15, DMPC17, DMPC18, *DNA-Binding*, *Rep*, DMPC21, DMPC24, *Attachment*, DMPC27, DMPC31, DMPC32, DMPC33, *Capsid*, DMPC35, DMPC37, *Rep*, *FtsL*, DMPC40, DMPC42, DMPC43, DMPC44, DMPC45, *Capsid*, DMPC48, DMPC52, *PLA2*, DMPC55, *Rep*, DMPC61, *Attachment*, DMPC64, DMPC65, *Rep*, DMPC67, *Capsid*, DMPC70, DMPC71, DMPC79, DMPC81

B

DMGG1 isolate ctbj340
DMPC1 — DMPC79 — DMPC17, Rep
500 1000 1500 2000 2500

DMGG1 isolate ctcg74
DMPC1 — DMPC7 — DMPC24, Attach
500 1000 1500 2000

DMGG1 isolate ctcf180
DMPC1 — DMPC7
500 1000 1500 2000

C

Genome Group #1 Protein Clusters' Structural Prediction

iVireons ANN 'Structural' score

DMPC1, DMPC7, DMPC13, DMPC17, DMPC24, DMPC26, DMPC28, DMPC30, DMPC61, DMPC63, DMPC71, DMPC74, DMPC76, DMPC79, DMPC80

Figure 5: Expression of putative capsid proteins

Images taken by negative stain electron microscopy. Genome maps are linearized for display purposes. Expressed genes are colored green. Other genes are colored blue. iVireons scores are listed in parentheses. (A-C) Images represent virus-like particles from iVireons-predicted viral structural genes. (D) Merkel Cell Polyomavirus Small-T (a viral non-structural protein) is shown as a negative control.

A — DMGC11 isolate ctgh70. DMPP27 (0.96); hypothetical protein.

B — DMGC2 Isolate ctbf349. DMPP77; DMPP6; DMPP22; DMPP2 (0.97).

C — DMGC3 isolate ctjb153. DMPP75; DMPP21 (0.99); DMPP3; DMPP4.

D — Merkel Cell Polyomavirus (non-structural). VP2; VP1; ALTO; Small-T (0.03).

Supplemental Figure 1: Divergence of proteins encoded by circular contigs

BLASTX summary of each circular DNA molecule recovered from virus enriched samples. Sequences were queried against a database of viral and plasmid sequences. Only hits with E values $< 1\times10^{-5}$ were plotted. Here, BLASTX only reports the most significant stretch of amino acid sequence from each circular contig, and, therefore, other regions of each contig can be assumed to be equally or less conserved.

Length, in nucleotides, of circular DNA sequences representing putative viral genomes from this study.

Supplementary Figure 3: Network analysis of other categories of viral proteins
The figure depicts the degree to which selected groups of novel proteins reported in the current study are related to known viral protein categories in the RefSeq database. See Figure 2 for additional information.

RefSeq
This study

**Microviridae
major capsid**

**CRESS
virus
replication
associated
protein**

**Microviridae
replication
associated
protein**

**CRESS virus
capsid #2**

**Microviridae
minor capsid (pilot)**

**Anelloviridae
ORF1/2/3**

**Parvoviridae
capsid/PLA2**

**Genomoviridae
capsid**

**CRESS virus
capsid #3**

**Inoviridae
ZOT (Packaging ATPase)**

**CRESS virus
capsid #4**

<u>Supplemental Figure 4: STNV capsid sequence is within the CRESS DNA capsid "realm"</u>
A sequence similarity network made using EFI-EST (E value<1×10⁻⁵) shows capsid protein sequences of related CRESS DNA viruses in RefSeq, STNV and homologous RNA satellites from GenBank, as well as protein sequences from viral genomes discovered in the present study thought to contain the STNV fold (RPS-BLAST top hit = CDD Accession:cl27634; E value<1×10⁻⁵).

<u>Supplemental Figure 5: Nematode-associated virus capsid expression</u>
Putative capsid proteins from CRESS virus isolate ctdh33 associated with rhabditid nematodes (Fig. 3) were expressed in a human cell line. (A) Negative Stain EM images of virus-like particles resulting from the expression of all three genes predicted to encode for STNV-like capsid proteins. (B) Agarose gel electrophoretic analysis of nucleic acids extracted from ctdh33 VLPs shows that purified VLPs from fraction 4 of the Optiprep gradient contain RNA. Note that Benzonase digests both DNA and RNA, DnaseI only digests DNA and Rnase only digests RNA.

A

B

= DNA ladder
= Fraction 3 + Benzonase
= Fraction 3 + DNaseI
= Fraction 3 + RNase
= Fraction 4 + Benzonase
= Fraction 4 + DNaseI
= Fraction 4 + RNase

Supplemental Figure 6: Genome maps of additional CRESS genomes encoding multiple capsid genes
CRESS 'Rep' genes are shown in yellow-orange, predicted virion structural genes in green, other candidate genes in grey.

Supplemental Figure 7: Novel acquisitions of RNA virus capsids

Sequence similarity network made in EFI-EST (E value <1×10^{-5}) showing capsid protein sequences of selected ssRNA viruses (Nodaviridae, Tombusviridae, tombus-like viruses) and ssDNA viruses (bacilladnavirus and crucivirus) together with protein sequences from viral DNA genomes observed in the present study with similarity to RNA virus coat protein (NCBI CDD: cl20209). CRESS virus isolate ctca5 capsid gene and CRESS virus isolate ctgh4 capsid gene had no detectible similarity to any known DNA virus sequences.

Supplementary Figure 8 (related to Figure 4C): iVireons scores of DMGGs with candidate viral structural gene(s)
Box-and-whisker plots of iVireons 'Structural' scores for individual DMPCs (numbers on x-axes) grouped by DMGG. Scores (y-axes) range from -1 (unlikely to be a virion structural protein) to 1 (likely to be a virion structural protein). DMGG2 and DMGG3 have been combined due to inferred chimerism.

Supplemental Figure 9: Sample Characterizaition by iterative BLAST Searches

Contigs of over 1000 nts from each sample were subject to an iterative BLAST search chararcterization. First, BLASTN was performed against the RefSeq database. Contigs without hits were then queried by BLASTX against all of GenBank 'nr' database. Contigs without hits were then queried by BLASTX against a database of proteins from genomes reported in this study. The proportion of total reads mapping to each contig was calculated and used for this plot. Individual inspection of contigs shows that most hits in the 'Translated AA alignment to GenBank 'nr' "Bacteria"' were likely plasmid or prophage proteins. The proportions of hits in each category are sensitive to stringency settings and to which databases are chosen for the analysis. The key aims of the figure are to display the proportion of reads the current survey rendered classifiable and the fraction of remaining dark matter reads in various samples.