Modeling epistasis in mice and yeast using the proportion of two or more distinct genetic backgrounds: evidence for "polygenic epistasis"

Christoph D. Rau, Natalia M. Gonzales, Joshua S. Bloom, Danny Park, Julien Ayroles, Abraham A. Palmer, Aldons J. Lusis, Noah Zaitlen[*]

Christoph D. Rau: Department of Anesthesiology, UCLA, chrau@ucla.edu

Natalia M. Gonzales: Department of Human Genetics, University of Chicago, ngon@uchicago.edu

Joshua S. Bloom: Department of Human Genetics, UCLA, Jbloom@mednet.ucla.edu

Julien F. Ayroles: Department of Ecology and Evolutionary Biology, Princeton University. jayroles@princeton.edu

Danny Park: Department of Medicine, UCSF, sihyung@gmail.com

Abraham A. Palmer: Department of Psychiatry, and Institute for Genomic Medicine, UCSD, aap@ucsd.edu

Aldons J. Lusis: Department of Human Genetics, UCLA, jlusis@mednet.ucla.edu

Noah Zaitlen: Department of Neurology, UCLA, nzaitlen@mednet.ucla.edu, Corresponding Author

Abstract

Background: The majority of quantitative genetic models used to map complex traits assume that alleles have similar effects across all individuals. Significant evidence suggests, however, that epistatic interactions modulate the impact of many alleles. Nevertheless, identifying epistatic interactions remains computationally and statistically challenging. In this work, we address some of these challenges by developing a statistical test for *polygenic epistasis* that determines whether the effect of an allele is altered by the global genetic ancestry proportion from distinct progenitors.

Results: We applied our method to data from mice and yeast. For the mice, we observed 49 significant genotype-by-ancestry interaction associations across 14 phenotypes as well as over 1,400 Bonferroni-corrected genotype-by-ancestry interaction associations for mouse gene expression data. For the yeast, we observed 92 significant genotype-by-ancestry interactions across 38 phenotypes. Given this evidence of epistasis, we test for and observe evidence of rapid selection pressure on ancestry specific polymorphisms within one of the cohorts, consistent with epistatic selection.

Conclusions: Unlike our prior work in human populations, we observe widespread evidence of ancestry-modified SNP effects, perhaps reflecting the greater divergence present in crosses using mice and yeast.

Keywords: Epistasis, Genetic Ancestry, GWAS, Mouse Genetics

Author Summary: Many statistical tests which link genetic markers in the genome to differences in traits rely on the assumption that the same polymorphism will have identical effects in different individuals. However, there is substantial evidence indicating that this is not the case. Epistasis is the phenomenon in which multiple polymorphisms interact with one another to amplify or negate each other's effects on a trait. We hypothesized that individual SNP effects could be changed in a polygenic manner, such that the proportion of as genetic ancestry, rather than specific markers, might be used to capture epistatic interactions. Motivated by this possibility, we develop a new statistical test that allowed us to examine the genome to identify polymorphisms which have different effects depending on the ancestral makeup of each individual. We use our test in two different populations of inbred mice and a yeast panel and demonstrate that these sorts of variable effect polymorphisms exist in 14 different physical traits in mice and 38 phenotypes in yeast as well as in murine gene expression. We use the term "polygenic epistasis" to distinguish these interactions from the more conventional two- or multi-locus interactions.

Introduction

Genetic association studies in humans and model organisms have identified a number of significant links between individual polymorphisms and phenotypic variability. A fundamental assumption of many of these studies is that an allele will have a similar effect in each member of the population, that is, that epistatic and other higher-order interactions across the genome can largely be ignored[1–4]. Prior studies to specifically detect epistasis in flies[5], yeast[6], mice[7,8], and humans[9] have considered pairwise interactions between loci to begin to identify interacting loci, yet this approach is hampered by a substantial increase in the threshold of significance due to the increased numbers of tests (from n to at least $\binom{n}{2}$, where n are the number of SNPs tested). In this study, we consider detection of epistasis by testing the effects of interactions between individual polymorphisms and global ancestry (θ), which we defined as the percentage of the genome inherited from a given ancestor or ancestral population.

Recently, we developed a SNP-ancestry interaction model and applied it to a human dataset, detecting only modest evidence of ancestry-specific genetic effects[10]. There is substantial evidence of interactions with ancestry in model organisms. For example, we observed radically different phenotypic consequences of null alleles of *Tcf7l2* and *Cacna1c* when expressed on different inbred strain backgrounds[11]. Recently, a number of powerful genetic resources have been developed for model systems to map variation in complex traits, such as the Hybrid Mouse Diversity Panel[12], the Collaborative Cross[13,14], the BXD recombinant inbred panel[15], various outbred mouse and rat populations[16], or the Drosophila Synthetic Population Resource[17] . The variation in ancestry present in these populations provide a unique opportunity to test for ancestry-driven epistasis, but no existing method is suitable for this task.

To address this problem, we incorporate a linear mixed model capable of accounting for the highly structured relatedness of constructed model organism[18] into our previous approach. We then

examine two distinct mouse cohorts as well as a panel of yeast crosses and test for interactions between

the effects of individual polymorphisms and the percentage of the genome for each individual that

originated with a specific founder. Although our initial model was designed to identify both genetic

background and environmental effects, the controlled environment in which model organisms are raised is

intended to minimize sources of environmental variance. This test, which we have called $Gx\theta$, acts as a

surrogate measure of the concurrent action of many other SNPs, and allows us to ask whether or not the

effect size of a given SNP changes as a function of overall genetic ancestry. In contrast to a pairwise

epistatic test, where significance means a detected interaction between two specific loci, a natural

interpretation of a significant $Gx\theta$ interaction is widespread epistasis of the tested genotype G with many

loci across the genome, as we previously showed[10].

We first evaluated the statistical properties of our method on simulated phenotypes based on real

recombinant inbred line data sets[12]. The $Gx\theta$ test successfully distinguishes between the marginal

effects driven only by a given SNP or ancestry and effects that arise from their interaction. We then

applied the test to two different populations of mice: recombinant inbred (RI) lines that are a subset of the

HMDP[12] (143 phenotypes), and a 50^{th} generation intercross between the inbred LG/J and SM/J mouse

strains[19], whose ancestry proportions have been clearly and precisely determined[19] (75 phenotypes).

To show that our approach is broadly applicable, we also apply this test to a yeast panel consisting of

thousands of progeny derived from 15 distinct yeast crosses[20] (40 phenotypes). Utilizing gene

expression data from the HMDP cohort in two different conditions, we further demonstrate that our

approach replicates findings across similar populations.

Given this evidence of epistasis, we examined the ancestry distribution at specific sites in the

mouse HMDP RI lines. Under neutrality, on average, we would expect equal representation of founder

ancestry in the panel, however, we observed that for many positions in the genome there is a statistically

significant depletion or overrepresentation of specific founder ancestries. We interpret this as evidence of

selection during the process of RI strain derivation and we identify regions of the genome with strong

selection up to 4 Mb in length. These regions are evidence of strong Gxθ interactions which inhibit or

promote the transmission of specific alleles to subsequent generations. These regions are enriched for

genes involved in cancer and organogenesis and are enriched (P=1.8E-4) for metallopeptidases, which

play key roles in fertility and neo/perinatal lethality(21–23).

Our observation of genomic locations where the effect size of a given polymorphism is driven by

overall genomic context and ancestry highlights the importance of studying epistasis and other effects in

model organisms. Studying these context-dependent effects in human cohorts has been notoriously

difficult, but our results indicate that given the right design and analytical framework these effects can be

highly significant, and has implications for our ability to build predictive models from genotype to

phenotype and account for ancestry effects in population studies.

Results

Method Overview:

Our objective is to determine whether a given SNP i has a different effect size as a function of

changes in overall genetic background. As described in our previous work(10), this is effectively

achieved through an interaction test between each SNP and global genetic ancestry θ. In a standard test

of the association between a polymorphism X_i and a phenotype **y**, we can model this relationship as

$y = \mu + \beta_i X_i + u + e$, where μ is the mean phenotypic value, β is the effect size of the SNP, u is a

random variable accounting for relatedness and e is a random variable accounting for all other sources of

error in the study. We extend this analysis to account for interactions between a polymorphism and

ancestry by adding terms to our model such that the relationship now becomes

$$y = \mu + \beta_i X_i + \delta\theta + \beta_{Gx\theta}{}_i \theta * X_i + u + e$$

where δ is the global weight of the ancestry effect, θ are the ancestries for all N individuals and $\beta_{Gx\theta}$ are

the weights of the Gxθ effect. Our Gxθ test is then an LRT test with a null of $\beta_{Gx\theta i} = 0$ and an alternate

of $\beta_{Gx\theta i} \neq 0$.

To account for multiple testing, we follow the best practices of the respective model organism

communities, specifically we use the previously reported genome-wide significance threshold for the

HMDP of 4.2E-6 (24) or the AIL of 8.06E-5 (19). For Yeast, we follow the example previously reported

of Bloom et al (25) and use an FDR of 5%.

Simulated Data:

To examine the properties of our approach in a model organism cross, we applied the method to

phenotypes simulated using real genotypes that reflect the underlying relatedness present within the

recombinant inbred (RI) lines of the Hybrid Mouse Diversity Panel (HMDP). Random SNPs with a

minor allele frequency (MAF) between 25% and 75% were selected from our panel and phenotypes

constructed at 200 values of either a main SNP effect β_G or a SNP-ancestry interaction effect $\beta_{G\times\theta}$ where θ

is the proportion of all SNPs arising from C57BL/6J (see Methods) mice for interaction with each

individual SNP with 1,000 simulations at each β (200,000 total simulations) (see Methods).

We observe that changes to either β_G or $\beta_{G\times\theta}$ independently did not affect the power of identifying

an association for $\beta_{G\times\theta}$ or β_G, respectively, indicating that these two terms are correctly being estimated

independently of one another. We briefly explored incorporating a second genetic relationship matrix

(GRM) accounting for population structure arising from descent from a single ancestor (the B6

mouse)(26), however we observed no significant improvement in power when incorporating this second

GRM (Figure 1a, 1b).

To evaluate what forms of interactions the test is powered to identify we examined a range of

epistatic architectures, focusing specifically on our method's ability to recover higher-order interactions, a

type of epistasis which traditional models struggle to accommodate. For 100,000 simulations, a SNP was

selected at random from SNPs with MAF between 25 and 75% and combined with 1-10 additional SNPs

(10,000 each) and phenotypes generated for these SNP groupings followed by analysis by the $G\times\theta$

algorithm (see Methods). As the number of interacting SNPs increases, the power to detect the individual

SNP decreases as expected (Figure 1c), however the power to detect a Gxθ locus rises (Figure 1d),

suggesting that our model is powerfully situated to identify SNPs involved in complex higher-order

interactions, especially those consisting of more than 5 interacting partners (Figure S1).

Application to *in vivo* Data:

Mouse Populations

We applied the Gxθ test to two large panels of mice to identify Gxθ effects (i.e. instances where a

given polymorphism interacts epistatically with one or many other loci as captured in the model by θ, the

global ancestry). Our first cohort is the HMDP, a set of 150+ commercially available inbred strains(27).

Numerous GWAS have been performed in the HMDP, including several using PYLMM, which forms the

core of our algorithm(24,28,29). The largest component of the HMDP is comprised of 122 RI strains (28

AxB, 71 BxD, 12 BxH, 11 CxB). Each RI was constructed from re-derivation of novel inbred lines via

brother-sister mating following an F_2 cross between the sub-panel's parental lines. In the case of the

HMDP, one of the parental strains for each RI strain was the commonly studied strain C57BL/6J (**B6**).

Using B6 as the common ancestral line to the RI panels, we calculated θs for each RI strain (Fig 2a). As

expected, the average ancestry attributable to B6 was roughly 50% (50.63%) and roughly normally

distributed. We removed a single outlier, BXD32/TyJ, whose B6 Ancestry of 25.41% reflects a

previously known additional backcross to DBA/2J, resulting in a strain that is 75% DBA and 25% B6.

Each study using the HMDP uses a different subset of the entire panel, and we selected a study on heart

failure induced by the chronic beta adrenergic agonist isoproterenol(30) for analysis as it used the most RI

strains compared to other published HMDP data. We used 123 clinical phenotypes in conjunction with

microarray-derived gene expressions measured in the left ventricle and on average tested 67 RI strains per

phenotype.

The second cohort consists of 1,063 animals from the $F_{50} - F_{56}$ generation of an advanced

intercross line (**AIL**) created by crossing the LG/J and SM/J inbred mouse strains (19). Unlike the RI

strains from the HMDP, AILs are maintained in a manner that minimizes inbreeding and promotes

genetic diversity at each generation. We arbitrarily set LG/J as the ancestral strain of interest and

calculated θs for each of the 1,063 mice in the panel (Fig 2b). For this study, we focused on a diverse set

of 133 phenotypes that had been measured in these mice. We describe the results of the Gxθ associations

in each panel before demonstrating replication of interactions in a phenotypic trait as well as in expression

data.

Evidence of SNP x Ancestry Interactions in Phenotypic Data

128   We applied the Gxθ method to the 123 observed heart-failure related phenotypes from the HMDP

RI strains. We observed well-calibrated statistics, with λ_{GC} equal to 0.978 and $\lambda_{Gx\theta C}$ equal to 1.045 after

pooling the p values across all phenotypes. We observed 44 significant Gxθ loci across 9 phenotypes:

E/A ratio, free fatty acid content in the blood, cardiac fibrosis, fractional shortening of the heart during

contraction, heart rate, internal diameter of the left ventricle, left ventricular mass and left and right atrial

weights (Table 1 and S1). These Gxθ loci were largely distinct from previously reported GWAS loci in

the same phenotypes in this panel of mice[28,31], yet contained a number of highly relevant genes, as

discussed below.

136   By way of example, we focus on two important phenotypes from the HMDP panel. Cardiac

fibrosis is a marker of cardiac dysfunction. Genes identified through the Gxθ screen (Figure 3B) as

potential candidates for cardiac fibrosis include: *Crisp2* (rs6295287, p=6.74E-7), a secreted biomarker of

cardiovascular disease[32], *Top2b* (rs31538570, p=2.14E-06), which plays a cardioprotective role in

response to stress[33], *Rarb* (rs31538570, p=2.14E-06), a known regulator of inflammation with

unknown function in the heart[34], and *Fibrosin* (rs33146511, p=2.54E-6), a major component of the

fibrosis pathway[35]. An increase in left ventricular mass in response to catecholamine challenge is the

primary marker of cardiac hypertrophy in the HMDP. A single Gxθ locus(rs31313229, p=2.91E-06) on

chromosome 3 (Figure 3C) contains the gene *Lphn2*, which has a role in the promotion of cellular

adhesion in response to external stimuli.[36] Crucially, none of these loci, many of which contain

candidates of particular interest or relevance to heart failure, were reported in the original study,

suggesting the possibility of new targets for further therapeutic research.

We next applied the Gxθ method to 133 phenotypes measured in the LG/J x SM/J AIL, including

the sex of the animal as a covariate as, unlike the HMDP data, the AIL consists of both male and female

animals. As with the HMDP data, we observed well-calibrated statistics in this cross, with λ_{GC} equal to

0.995 and $\lambda_{Gx\theta C}$ equal to 1.033. Despite the larger sample size, we observed only 4 significant loci across

4 phenotypes: activity levels in a saline-injected animal on day 3 of a conditioned place preference test;

average weight of animals across 5 different time points roughly a week apart; glucose (mg/dL) in blood

after a 4 hour fast [Figure 3D]; weight at ~68 days of age (Table 1 and S1).

Yeast Populations:

We also used the Gxθ algorithm to examine 15 previously reported yeast crosses(20) ranging

from 650 to 950 progeny per cross for evidence of significant Gxθ loci and an average of 71,000

segregating genetic variants. We examined the responses of these progeny to forty different chemical

stimuli(20) and recovered 92 loci across 38 phenotypes which met a 5% FDR threshold for that

cross/phenotype combination (Table S2). At a more stringent level of significance (5% FDR across an

entire cross), we identify 9 loci across 7 phenotypes (LiCl, EGTA, Maltose, MnSO4, Paraquat, YNB,

YPD) which show strong evidence of Gxθ interactions (Table 1, Figure 3A).

Gxθ Associations in HMDP Cohort Gene Expression

We next examined gene expression in the hearts of the HMDP cohort using previously obtained

transcriptome microarrays from both control and treated conditions(37). Each cohort consists of

approximately 70 RI lines, with 66 lines overlapping between the two cohorts (full lists of strains in Table

S2).   We examined 13,155 expressed and varying (Coefficient of Variation > 5%) genes from the left

ventricles of the HMDP cohorts using the ~170k SNPs with MAF >= 0.05. We observed 1,486

significant associations with 18 genes at a Bonferroni-corrected P value of 3.2E-10 (135,130 associations

with 1,350 genes at GW-significant threshold of 4.2E-6) in the control cohort and 597 significant

associations with 39 genes at the same threshold in the treated cohort (32,043 associations with 1,042

genes at the genome-wide significant threshold of 4.2E-6(24)) (Full results available in Table S10).

Phenotypic variability was larger in the catecholamine treated cohort of the heart failure HMDP

study (average Coefficient of Variation in control was 22.6% vs 28.6% for the treated cohort)(30), which

could partially explain the lower number of significant associations observed in that group of mice.

Notably, we saw no enrichment for SNPs in the *cis*-regions of the examined genes in contrast to what is

observed in standard mapping. Genes in the control cohort with significant Gx□ associations are

enriched for mitochondria-associated genes (P=1.8E-10 (P=.032 in treated)), suggesting a role of Gx□

interactions in the regulation of mitochondrial dynamics. Genes in the treated cohort with significant

associations were enriched for genes involved in post-transcriptional modifications to RNA including

RNA splicing (P=7.4E-3), highlighting the importance of alternative splicing to the response to

catecholamine challenge(38).

Replication of Gx□θ Associations across eQTLs from HMDP Cohort

To demonstrate that the method is able to replicate Gxθ results across cohorts, we examined the

reproducibility of expression Gxθ QTLs in the treated and untreated RI lines of the HMDP. Of the 1,486

associations observed in the control data, we observe 305 (21%) with Gxθ interaction (P<.05) in the

treated cohort (36 at FDR< 5%), indicating a strong replication (p = 2.2E-16 from a binomial test)

between the two cohorts despite differences in genetic background and the potentially reduced power in

the treated cohort caused by the effects of the catecholamine drug.

Distortion from Expected Allelic Frequencies

After population admixture, allelic frequencies will shift in the resulting population. This can be

due to the effects of selection against deleterious allelic combinations (i.e. fitness epistasis), meiotic drive

(i.e. intragenomic conflict resulting in the transmission of one or more alleles over another during

meiosis). Alternatively, allele frequencies may shift through random genetic drift. We hypothesize that

Gxθ loci can be driven by fitness epistasis, or instances where allelic variation in one locus affects fitness

at other loci across the genome. This hypothesis is in line with prior research(39–41) which has explored

allelic imbalance or gametic ratio distortion as a function of epistasis and fitness. Examining the genomes

of admixed population for signs of significant allelic shift may reveal interesting loci that can inform long

standing questions in evolutionary and population genetics about the forces that maintains variation in

fitness in populations, as well as the presence of non-equal chromosomal inheritance.

We therefore conducted a test to search for individual loci with enriched or depleted B6 ancestry

in the RI strains. As expected, the average B6 ancestry across all SNPs was 50.68% +/- 8.1%, which is

indistinguishable from the ancestry by strain average of 50.63% +/- 6.7% (Fig 4a). At the level of

individual loci, however, we observed significant variation in B6 ancestry across the genome (Fig 4b),

with some SNPs displaying very low or very high frequency (Fig 4c). We calculated the statistical

likelihood of detecting the observed ancestries at all loci across the genome (λ=1.07, Fig 4d).

We subsequently performed a Wald test for ancestry frequency (θ) at each SNP i where $W =$

$\frac{(\bar{\theta}-\theta_i)^2}{var(\theta)}$. We used the global variance in the denominator, which assumes that most SNPs lie under the

null and which is conservative if this assumption is violated and could be replaced with a more powerful

test. For each SNP, we examined strains which were not invariant for that locus and identified 614 SNPs

from the 170k original SNPs with significantly altered ancestries at an FDR of 5% (Table S3). We

suspect that these SNPs do not represent passive effects such as genetic drift due to the nature of how RI

panels are created, with a single F2 generation of mice followed by brother-sister mating and eventual

fixation as similar effects on allelic balance have been attributed to genetic incompatibilities in other

mouse populations(42). Allelic balance of paternal alleles across the RI panel, therefore, may be modeled

using a Bernoulli distribution. In the HMDP, with 122 RI lines, the odds of a 30% imbalance (80% B6 vs

20% other) would be 8.3E-12, which, even across 200,000 SNPs results in 2E-6 expected SNPs with this

level of imbalance across the entire genome.

Nine loci were identified in which 5 or more SNPs were located together (Table 2). These 9

regions contain the majority (525, 86%) of all significant affected loci and indicate regions where

significant allelic imbalances are occurring. One region of particular interest lies on chromosome 18 and

contains 246 SNPs (40% of all significantly altered SNPs) with an average B6 ancestry of 82.5%. SNPs

in this region are found between B6 and C3H/HeJ and B6 and BALB/cJ. This locus contains a number of

interesting candidate genes potentially relating to organismal fitness, including *Epc1*, a transcription

factor linked to DNA repair, muscle differentiation and cancer suppression(43,44). It is also the only

gene within the locus whose expression has a suggestive Gxθ association with the locus itself (P=0.014).

When all genes within 2 MB (the average Linkage Disequilibrium (LD) block size in the

HMDP(27)) of boundaries of these allelic imbalance loci are examined as a whole, we observe significant

enrichments for genes involved in cancer (corrected P-values range from 1.3E-8 to 1.6E-7). These genes,

whose functions range from cell cycle control to organismal growth and DNA damage mediation and

repair are not unexpected given their crucial role in organismal survival. If the B6 or alternative allele

deleteriously affect the function of one or more of these genes when combined with new alleles from

other strains located elsewhere on the genome, then it is reasonable to assume that the ability for an

individual to pass on its genetic information to the next generation may be significantly reduced. Other

enriched gene ontology categories include the crucial-for-life categories of respiratory tube development

(2.7E-4) and heart development (7.6E-4), likely mediated through changes in Matrix Metalloproteinase

activity (3.9E-8). These enzymes are canonically responsible for the regulation of the extracellular

matrix, but have been linked to the modulation of responses to many bioactive molecules and have

important roles in fertility, embryonic development and neo/perinatal mortality in addition to many other

diseases(23). Taken together, these enrichments suggest that we may be observing depletion due to

epistatic interactions with other loci which were neutral or beneficial in the original inbred lines, but

which become deleterious for a given allele when placed into a new genetic environment.

Discussion

We present a test which we call Gxθ, that leverages admixed populations such as inbred mouse

strains to identify sites of "polygenic epistasis" where a SNP interacts with an unknown number of other

loci summarized into a single genomic ancestry score, θ. One major advantage of this approach is that,

unlike other epistasis-focused association approaches, it does not increase the number of tests when

compared to a typical GWAS, so similar significance thresholds may be used.

The existence of epistatic interactions and their role in human diseases and phenotypes have been

known for many years, with notable examples of gene-gene interactions in Alzheimer's disease(45),

Bardet-Biedl syndrome(46) as well as classic interactions governing hair color and skin pigmentation.

Despite these examples and many more like them, epistatic interactions have proven difficult to find using

association techniques(47). One notable exception has been the success of using consomic strains of

mice, where one or more chromosome in strain A is replaced with a chromosome from another strain

B(48). This approach, which increases or decreases the contribution of a given strain to phenotypic and

genetic variance in a clear and controlled manner has resulted in the identification of numerous epistatic

interactions(49,50).

Our method sought to circumvent this traditional pitfall in Epistasis GWAS by examining each

SNP only once for an interaction with a global genomic ancestry to identify interactions or "polygenic

epistasis". Our model is able to identify much higher order interactions than can be observed using

traditional SNP-SNP interaction models. As demonstrated through simulation, our model in fact

increases in power as the complexity of the epistatic interactions increases, likely reaching peak power in

the hypothetical scenario where every SNP interacts with the tested SNP to affect phenotypic expression.

When applied to human populations, a different form of our test was able to find several significant

associations(10)

In this study, we examined two populations of mice: the recombinant inbred lines of the Hybrid

Mouse Diversity Panel(27) and an Advanced Intercross Lines (AIL) based on LG/J and SM/J(19) as well

as a yeast RIL mapping panel founded by fifteen yeast strains(20). We observed significant Gxθ

interactions in 14 of the 15 yeast crosses that affected over half of the examined phenotypes, reinforcing

the widespread role of genotype-by-genotype interactions in regulating quantitative traits across diverse

genetic backgrounds and phenotypes. In the mouse cohorts, however, despite nearly ten times the number

of genetically distinct mice as well as a larger number of phenotypes and genotypes in the AIL, we

observed approximately eleven-fold more significant Gxθ peaks in the HMDP phenotypes than in the AIL

phenotypes (44 vs 4).

Several reasons may account for this difference. First, outbred mice have significantly lower

power due to increased phenotypic variance caused by high rates of heterozygosity at alleles while inbred

populations have increased power due to a lack of this heterozygosity. This can also be seen in our

regular GWAS results(19,30,31). Second, the different phenotypes studied in each cohort will have

different genetic architecture and experimental noise (e.g. the AIL study includes many behavioral traits

while the HMDP does not). Third, significantly more recombination events have occurred in the AIL

compared to the HMDP, leading to a tighter distribution of background and a larger number of

independent tests. Fourth, the genetic backgrounds of the two cohorts are different, which might also

contribute to differences in observed numbers of Gxθ interactions. Although individually, LG and SM

are only slightly less genetically diverged from one another as any pair of strains that make up the RI

panels of the HMDP (Table S4), the presence of five ancestral lines in the HMDP compared to only two

in the F50 cross results in much more genetic diversity in the HMDP compared to the F50 mice. Finally,

we observe a higher variance in ancestral background in the mice HMDP when compared to the AIL. As

our method relies on differences in ancestral background to identify sites with different effect sizes in

different genetic contexts, the reduced variance in the AIL lines necessarily corresponds to a decrease in

the power to detect Gxθ interactions. Taken together, our method is best suited to datasets with relatively

low heterozygosity, clear and numerous differences in genetic background, and with higher variance in

the percentage of SNPs attributable to a given ancestry.

Furthermore, we identify nine regions across the genome of the RI lines of the HMDP which

show a strong selection for or against a particular parental allele. We propose that this is a result of an

epistatic fitness interaction, where individual loci in one ancestral strain are genetically incompatible with

alleles from another strain. We considered other alternatives explanations, some of which were explored

in prior work(41). One alternative would be population structure, where recombination may not have had

sufficient time to randomly assort alleles after a bottleneck event, though we think that this is an unlikely

explanation for our results. One of the key benefits of working with synthetic mapping populations such

as RI panels is that the crossing scheme (e.g. fennel cross, round-robin)is designed to minimize such

effects(40). Another possibility would be assortative mating, where mate selection drives the selection

for or depletion of a particular allele. While possible, and not explicitly arising from epistasis per se, it

would still produce a fitness differential across allelic combinations and be picked up by our analysis.

Finally, epistatic interactions, the focus of this manuscript, can lead to changes in fitness as demonstrated

by prior work(41) which examined Drosophila RI panels in which all expected genotypic combinations

for a diallelic cross were generated. They demonstrated a strong correlation between the degree of allelic

imbalance and the effect size of the fitness effect of each allelic combination. It follows that similar

scenarios should exist in mouse and yeast panels as well. One possible extension of this approach would

be to study fitness in the Collaborative Cross strains, another animal population in which a large

proportion of the strains went extinct, presumably due to genetic incompatibilities.

In the future, our approach could be leveraged to study additional extant populations or newly

created cohorts. For example, our tool could be used to examine whether genetic modifications such as

knockouts have differential effects across the mice of a cross, and we are actively exploring its utility in

this regard.

Conclusion

The results of our study suggest that heterogenous SNP effects due to differing ancestries is

pervasive in model organism populations. This observation is consistent with prior observations of

epistatic interactions in inbred strains of mice as well as examples of ancestry interaction in human

studies(10). However, the number and magnitude of ancestry interactions we found was much larger than

those found in human studies; we hypothesize this is driven by the larger genetic distance of the

ancestors. Further analyses of the sites of polygenic epistasis may reveal novel epistatic interactions

which drive phenotypic expression, and suggests that careful attention to genetic ancestry should be

considered when studying the role of an individual polymorphism on a phenotype.

Methods

Model Organism Populations and Ancestry Calling

Model organism data were drawn from previously reported studies(19,20,30). The LGxSM AIL

consists of 1,063 G50-56 mice derived from an original F1 intercross between the LG and SM inbred

lines. The Hybrid Mouse Diversity Panel consists of over 150 strains of commercially available inbred

mice(27), of which 122 strains were recombinant inbred lines and suitable for our study. The yeast

crosses originated from 15 original strains which were bred in pairs to create 15 new crosses. SNPwise

ancestries were determined by identifying all SNPs which differed between parental lines (AIL: LG and

SM or HMDP: C57BL/6 and A, C3H, DBA/2 or BALB/c or Yeast: See Table S5). Genotypes from the

G50-56.RI lines or yeast crosses were filtered for these SNPs and ancestries calculated using either LG or

C57BL/6 or one of the yeast strains as the strain background of interest.

Simulation Framework

 Power Calculations

We created sets of simulated phenotypes based on the genotypes of the HDMP RI panel, which is

an admixed population in which the B6 strain, on average, contributes 50% of each strain's DNA.   For

each simulated phenotypes, we drew a SNP (75% > MAF > 25%) at random from the HMDP genotypes

and created a phenotype based on β, the genetic effect size, ☐ the effect size of the interaction between

global ancestry (θ) and the chosen SNP and a multivariate normal (mvn) derived from three variance

terms: σ_g^2, the proportion of variance attributable to genetic effects σ_θ^2 , the proportion of variance

attributable to Gxθ effects and σ_ε^2, the residual proportional variance attributable to all combined sources

of error and variation not considered in this study. Phenotypes were generated both with and without the

Gxθ variance term to ascertain the necessity of incorporating a second GRM ($\mathbf{K}^A$), which would correct

for relatedness in ancestry, into the algorithm.

We simulated four distinct phenotypes for our analysis

1) Phenotypes generated by including a SNP Effect

349
$$y = \beta X + mvn\left(0, \sigma_g^2 \mathbf{K} + \sigma_e^2 \mathbf{I}\right) \ or$$

$$y = \beta X + mvn(0, \sigma_g^2 \mathbf{K} + \sigma_\theta^2 \mathbf{K}^A + \sigma_e^2 \mathbf{I})$$

350   2) Phenotypes generated by including a Gxθ Effect

351
$$y = \beta_{Gx\theta}\theta * X + mvn(0, \sigma_g^2 \mathbf{K} + \sigma_e^2 \mathbf{I}) \ or$$

$$y = \beta_{Gx\theta}\theta * X + mvn(0, \sigma_g^2 \mathbf{K} + \sigma_\theta^2 \mathbf{K}^A + \sigma_e^2 \mathbf{I})$$

In each phenotype, σ_g^2 was set to 0.4. When incorporated, σ_θ^2 was set to 0.2 and σ_ε^2 was set to the

remainder of the variance (0.6 or 0.4). The power of our model and independence of our β and $\beta_{Gx\theta}$

terms were queried by varying either β or $\beta_{Gx\theta}$ from 0 to 1 (200 values set 0.005 apart) with 1,000

simulated phenotypes at each step (200,000 total simulations per phenotype).

Epistasis Simulation

Using the same panel and set of SNPs described above, we drew one SNP at random to be our

test SNP and one to ten additional SNPs to be simulated epistatically interacting SNPs (10,000

simulations per set of interacting SNPs, 100,000 simulations in total).   For each simulation we created a

composite SNP in which only strains with the minor (non-B6) allele in every one of the tests and

interacting SNPs had that allele in the composite SNP. We used this composite SNP to generate a

phenotype as described above and selecting an effect size to generate a χ^2 test statistic of 20.

Gxθ Model

The equation to determine the effects of a SNP and a SNP x Ancestry term on a phenotype can be

written as:

$$y_k = \mu + \sum_{i=1}^{M} \beta_i X_i + \delta\theta_K + \sum_{i=1}^{M} \beta_{Gx\theta_i} \theta_i X_{ik} + \varepsilon_k$$

Where y_k is the phenotype of individual k, μ is the mean phenotypic value, M is the number of markers, β

are the weights on the SNPs, X is the *m by n* array of SNP genotypes, δ is the global weight of the

ancestry effect, θ is the ancestries for all N individuals. $\beta_{Gx\theta}$ are the weights of the Gxθ effect and ε is

the combined error term. We want to identify SNPs where $\beta_{Gx\theta_i} \neq 0$ as these are sites where Ancestry is

interacting with our genotypes.

Motivated by our model above, we can write a new model for the effect of a single SNP i on a

phenotypic trait as:

$$y = \mu + \beta_i X_i + \delta\theta + \beta_{Gx\theta_i}\theta * X_i + u + e$$

375   For an individual SNP i. Here, θ is the column vector of ancestries, and θ * X is the element-wise

376   product. The random effect u accounts for relatedness of individuals based on SNPs. Our Gxθ test is then

377   an LRT test with a null of $\beta_{Gx\theta_i} = 0$ and an alternate of $\beta_{Gx\theta_i} \neq 0$.

Reporting Gxθ Results and Identifying Genes of Interest

GWAS thresholds were previously determined to be 4.1E-6 in the case of the HMDP(24) or

8.06E-6 in the case of the AIL(19). An FDR of 5% was used in the yeast crosses as previously

reported(25) Genes of Interest were identified by examining nearby genes (within 2 MB) for cis-eQTLs

as previously reported in the HMDP(30) and/or by literature and database analyses using bioGPS

(www.biogps.org) which draws data from PubMed and other publicly available databases to help

researchers understand the role of their genes of interest.

Determining Significance of Allelic Imbalance Sites

Allelic imbalance significance was determined by performing a Wald test for ancestry frequency

(θ) at each SNP i where $W = \frac{(\bar{\theta} - \theta_i)^2}{var(\theta)}$ and only strains where the SNP was not invariant were included.

Allelic imbalances were considered to be significant after Benjamini-Hochberg correction at a cutoff of

5%.

Gene Set Enrichment Analysis:

All gene enrichments were performed using the GeneAnalytics platform(51), which uses a

binomial distribution to assess the enrichment for a user-supplied list of genes within GO terms,

Superpaths and other biologically relevant categories. Reported P-values have been corrected for

multiple comparisons using the Benjamini-Hochberg correction.

Declarations:

Ethics Approval and Consent to Participate: Not Applicable

Consent for Publication: Not Applicable

Availability of Data and Materials:

Data from the HMDP may be accessed at https://systems.genetics.ucla.edu/

Data from the AIL cross may be accessed at http://palmerlab.org/protocols-data/

Data from the Yeast crosses may be accessed at https://github.com/joshsbloom/yeast-16-parents

The Gxθ algorithm may be found at https://github.com/ChristophRau/GxTheta

Competing Interests: The authors declare they have no competing interests

Funding: This work was supported by NIDA (AAP: R01DA021336; NMG: F31DA036358), NIGMS

(NMG: T32GM007197, JFA: R35GM124881), and the NIH (CDR: K99HL138301, NZ:

K25HL121295, U01HG009080, and R01HG006399, AJL: 1U54 DK120342, 1R01 HL147883 and

1R01 DK117850). The funding bodies played no role in the study other than funding.

Author Contributions:

CDR designed the project, created the Gxθ software, acquired the HMDP data and analyzed and

interpreted the data used in this manuscript as well as drafted the manuscript. NMG assisted in the design

of the AIL cohort, acquired the AIL data and substantially edited the manuscript. JB designed and

acquired the yeast cross data and substantially edited the manuscript. DP assisted in conceptualizing the

Gxθ algorithm and substantially revised the manuscript. JA provided conceptual support and

substantially revised the manuscript. AAP assisted in the design of the AIL cohort and substantially

revised the manuscript. AJL assisted in the design of the HMDP cohort and substantially revised the

manuscript. NZ conceptualized and designed the Gxθ algorithm, interpreted the data and helped to draft

and substantially revise the manuscript. All authors read and approved the final manuscript.

Acknowledgements:

The authors would like to thank Yibin Wang for his constructive critiques of the algorithm and the

manuscript.

References

1.	Polderman TJC, Benyamin B, de Leeuw C a, Sullivan PF, van Bochoven A, Visscher PM, et al. Meta-analysis of the heritability of human traits based on fifty years of twin studies. Nat Genet [Internet]. 2015;47(7):702–9. Available from: http://www.nature.com/doifinder/10.1038/ng.3285

2.	Hill WG, Goddard ME, Visscher PM. Data and theory point to mainly additive genetic variance for complex traits. PLoS Genet [Internet]. 2008 Feb 29 [cited 2018 Oct 2];4(2):e1000008. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18454194

3.	Fish AE, Capra JA, Bush WS. Are Interactions between cis-Regulatory Variants Evidence for Biological Epistasis or Statistical Artifacts? Am J Hum Genet [Internet]. 2016 Oct 6 [cited 2018 Oct 2];99(4):817–30. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27640306

4.	Mäki-Tanila A, Hill WG. Influence of gene interaction on complex trait variation with multilocus models. Genetics [Internet]. 2014 Sep [cited 2018 Oct 2];198(1):355–67. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24990992

5.	Huang W, Richards S, Carbone MA, Zhu D, Anholt RRH, Ayroles JF, et al. Epistasis dominates the genetic architecture of Drosophila quantitative traits. Proc Natl Acad Sci U S A. 2012;

6.	Costanzo M, VanderSluis B, Koch EN, Baryshnikova A, Pons C, Tan G, et al. A global genetic interaction network maps a wiring diagram of cellular function. Science (80-). 2016;

7.	Tyler AL, Donahue LR, Churchill GA, Carter GW. Weak Epistasis Generally Stabilizes Phenotypes in a Mouse Intercross. PLoS Genet [Internet]. 2016 Feb [cited 2018 Oct 2];12(2):e1005805. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26828925

8.	Varón-González C, Navarro N. Epistasis regulates the developmental stability of the mouse craniofacial shape. Heredity (Edinb) [Internet]. 2018 Sep 12 [cited 2018 Dec 4];1. Available from: http://www.nature.com/articles/s41437-018-0140-8

9.	Hemani G, Shakhbazov K, Westra H-J, Esko T, Henders AK, McRae AF, et al. Detection and replication of epistasis influencing transcription in humans. Nature [Internet]. 2014 Apr 10 [cited 2014 Oct 20];508(7495):249–53. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3984375&tool=pmcentrez&rendertype=abstract

10.	Park DS, Eskin I, Kang EY, Gamazon ER, Eng C, Gignoux CR, et al. An ancestry-based approach for detecting interactions. Genet Epidemiol [Internet]. 2018 Feb 1 [cited 2018 Oct 2];42(1):49–63. Available from: http://doi.wiley.com/10.1002/gepi.22087

11.	Sittig LJ, Carbonetto P, Engel KA, Krauss KS, Barrios-Camacho CM, Palmer AA. Genetic Background Limits Generalizability of Genotype-Phenotype Relationships. Neuron [Internet]. 2016 Sep 21 [cited 2018 Jul 25];91(6):1253–9. Available from: https://www.sciencedirect.com/science/article/pii/S0896627316305050?via%3Dihub

12.	Lusis AJ, Seldin MM, Allayee H, Bennett BJ, Civelek M, Davis RC, et al. The Hybrid Mouse Diversity Panel: A Resource for Systems Genetics Analyses of Metabolic and Cardiovascular Traits. J Lipid Res [Internet]. 2016 Jun [cited 2016 Oct 13];58(12):7250–7. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27099397

13.	Philip VM, Sokoloff G, Ackert-Bicknell CL, Striz M, Branstetter L, Beckmann MA, et al. Genetic analysis in the Collaborative Cross breeding population. Genome Res. 2011;

14.	Churchill GA, Airey DC, Allayee H, Angel JM, Attie AD, Beatty J, et al. The Collaborative Cross, a community resource for the genetic analysis of complex traits. Nature Genetics. 2004.

15.	Ashbrook DG, Arends D, Prins P, Mulligan MK, Roy S, Williams EG, et al. The expanded BXD family of mice: A cohort for experimental systems genetics and precision medicine. bioRxiv. 2019 Jul 8;672097.

16.	Solberg Woods LC, Palmer AA. Using Heterogeneous Stocks for Fine-Mapping Genetically Complex Traits. In: Methods in Molecular Biology. Humana Press Inc.; 2019. p. 233–47.

17.	King EG, Merkes CM, McNeil CL, Hoofer SR, Sen S, Broman KW, et al. Genetic dissection of a

model complex trait using the Drosophila Synthetic Population Resource. Genome Res. 2012;

18. Kang HM, Zaitlen NA, Wade CM, Kirby A, Heckerman D, Daly MJ, et al. Efficient control of population structure in model organism association mapping. Genetics [Internet]. 2008 Mar [cited 2012 Mar 13];178(3):1709–23. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2278096&tool=pmcentrez&rendertype =abstract

19. Gonzales NM, Seo J, Hernandez-Cordero AI, Pierre CL St., Gregory JS, Distler MG, et al. Genome wide association analysis in a mouse advanced intercross line. Nat Commun [Internet]. 2018 Sep 10 [cited 2018 Oct 2];in press. Available from: https://www.biorxiv.org/content/early/2018/09/10/230920

20. Bloom JS, Boocock J, Treusch S, Sadhu MJ, Day L, Oates-Barker H, et al. Rare variants contribute disproportionately to quantitative trait variation in yeast. bioRxiv [Internet]. 2019 Apr 15 [cited 2019 Aug 16];607291. Available from: https://www.biorxiv.org/content/10.1101/607291v1

21. Jha SK, Rauniyar K, Jeltsch M. Key molecules in lymphatic development, function, and identification. Ann Anat - Anat Anzeiger [Internet]. 2018 Sep 26 [cited 2018 Jul 25];219:25–34. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29842991

22. Afratis NA, Selman M, Pardo A, Sagi I. Emerging insights into the role of matrix metalloproteases as therapeutic targets in fibrosis. Matrix Biol [Internet]. 2018 Aug [cited 2018 Jul 25];68–69:167–79. Available from: http://www.ncbi.nlm.nih.gov/pubmed/29428229

23. Cerdà-Costa N, Gomis-Rüth FX. Architecture and function of metallopeptidase catalytic domains. Protein Sci [Internet]. 2014 Feb [cited 2018 Jul 25];23(2):123–44. Available from: http://www.ncbi.nlm.nih.gov/pubmed/24596965

24. Bennett BJ, Farber CR, Orozco L, Kang HM, Ghazalpour A, Siemers N, et al. A high-resolution association mapping panel for the dissection of complex traits in mice. Genome Res [Internet]. 2010 Feb [cited 2011 Jul 2];20(2):281–90. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2813484&tool=pmcentrez&rendertype =abstract

25. Bloom JS, Boocock J, Treusch S, Sadhu MJ, Day L, Oates-Barker H, et al. Rare variants contribute disproportionately to quantitative trait variation in yeast. Elife. 2019;

26. Sul JH, Bilow M, Yang W-Y, Kostem E, Furlotte N, He D, et al. Accounting for Population Structure in Gene-by-Environment Interactions in Genome-Wide Association Studies Using Mixed Models. Schork NJ, editor. PLOS Genet [Internet]. 2016 Mar 4 [cited 2019 Feb 5];12(3):e1005849. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26943367

27. Ghazalpour A, Rau CDCD, Farber CRCR, Bennett BJBJ, Orozco LDLD, Van Nas A, et al. Hybrid mouse diversity panel: a panel of inbred mouse strains suitable for analysis of complex genetic traits. Mamm Genome [Internet]. 2012 Oct [cited 2012 Nov 30];23(9–10):680–92. Available from: http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3586763&tool=pmcentrez&rendertype =abstract

28. Rau CD, Wang J, Avetisyan R, Romay MC, Martin L, Ren S, et al. Mapping genetic contributions to cardiac pathology induced by beta-adrenergic stimulation in mice. Circ Cardiovasc Genet. 2015;8(1).

29. Parks BWW, Sallam T, Mehrabian M, Psychogios N, Hui STT, Norheim F, et al. Genetic Architecture of Insulin Resistance in the Mouse. Cell Metab [Internet]. 2015 Feb 3 [cited 2015 Feb 4];21(2):334–46. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25651185

30. Rau CD, Wang J, Avetisyan R, Romay MC, Martin L, Ren S, et al. Mapping genetic contributions to cardiac pathology induced by beta-adrenergic stimulation in mice. Circ Cardiovasc Genet [Internet]. 2015 Dec 5 [cited 2014 Dec 8];8(1):40–9. Available from: http://circgenetics.ahajournals.org/cgi/doi/10.1161/CIRCGENETICS.113.000732

31. Wang JJ-C, Rau C, Avetisyan R, Ren S, Romay MCMC, Stolin G, et al. Genetic Dissection of

Cardiac Remodeling in an Isoproterenol-Induced Heart Failure Mouse Model. PLoS Genet [Internet]. 2016 Jul [cited 2016 Oct 13];12(7):1–30. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27385019

32. Gibbs GM, Scanlon MJ, Swarbrick J, Curtis S, Gallant E, Dulhunty AF, et al. The cysteine-rich secretory protein domain of Tpx-1 is related to ion channel toxins and regulates ryanodine receptor Ca2+ signaling. J Biol Chem [Internet]. 2006 Feb 17 [cited 2018 Oct 2];281(7):4156–63. Available from: http://www.ncbi.nlm.nih.gov/pubmed/16339766

33. Zhang S, Liu X, Bawa-Khalfe T, Lu L-S, Lyu YL, Liu LF, et al. Identification of the molecular basis of doxorubicin-induced cardiotoxicity. Nat Med [Internet]. 2012 Nov 28 [cited 2018 Oct 2];18(11):1639–42. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23104132

34. Albino-Sanchez M, Vazquez-Hernandez J, Ocadiz-Delgado R, Serafin-Higuera N, León-Galicia I, Garcia-Villa E, et al. Decreased RARβ expression induces abundant inflammation and cervical precancerous lesions. Exp Cell Res [Internet]. 2016 Aug 1 [cited 2018 Oct 2];346(1):40–52. Available from: http://www.ncbi.nlm.nih.gov/pubmed/27207583

35. Wyler DJ. Fibrosin, a Novel Fibrogenic Protein: Discovery, Cloning and Implications for Fibrotic Disorders. Int Arch Allergy Immunol [Internet]. 1996 Dec [cited 2018 Oct 2];111(4):326–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/8957104

36. White GR, Varley JM, Heighway J. Genomic structure and expression profile of LPHH1, a 7TM gene variably expressed in breast cancer cell lines. Biochim Biophys Acta [Internet]. 2000 Apr 25 [cited 2018 Oct 2];1491(1–3):75–92. Available from: http://www.ncbi.nlm.nih.gov/pubmed/10760572

37. Rau CD, Romay MC, Tuteryan M, Wang JJ-C, Santolini M, Ren S, et al. Systems Genetics Approach Identifies Gene Pathways and Adamts2 as Drivers of Isoproterenol-Induced Cardiac Hypertrophy and Cardiomyopathy in Mice. Cell Syst [Internet]. 2017 Nov [cited 2016 Nov 29];4(1):121-128.e4. Available from: http://linkinghub.elsevier.com/retrieve/pii/S2405471216303593

38. Lara-Pezzi E, Gómez-Salinero J, Gatto A, García-Pavía P. The Alternative Heart: Impact of Alternative Splicing in Heart Disease. J Cardiovasc Transl Res [Internet]. 2013 Dec 18 [cited 2018 Oct 2];6(6):945–55. Available from: http://www.ncbi.nlm.nih.gov/pubmed/23775418

39. Schumer M, Brandvain Y. Determining epistatic selection in admixed populations. Mol Ecol. 2016;

40. Ehrenreich IM. Epistasis: Searching for interacting genetic variants using crosses. Genetics. 2017.

41. Corbett-Detig RB, Zhou J, Clark AG, Hartl DL, Ayroles JF. Genetic incompatibilities are widespread within species. Nature. 2013;

42. Srivastava A, Morgan AP, Najarian ML, Sarsani VK, Sigmon JS, Shorter JR, et al. Genomes of the mouse collaborative cross. Genetics. 2017;

43. Wang Y, Alla V, Goody D, Gupta SK, Spitschak A, Wolkenhauer O, et al. Epigenetic factor EPC1 is a master regulator of DNA damage response by interacting with E2F1 to silence death and activate metastasis-related gene signatures. Nucleic Acids Res [Internet]. 2016 Jan 8 [cited 2018 Oct 2];44(1):117–33. Available from: http://www.ncbi.nlm.nih.gov/pubmed/26350215

44. Kee HJ, Kim J-R, Nam K-I, Park HY, Shin S, Kim JC, et al. Enhancer of Polycomb1, a Novel Homeodomain Only Protein-binding Partner, Induces Skeletal Muscle Differentiation. J Biol Chem [Internet]. 2007 Mar 9 [cited 2018 Oct 2];282(10):7700–9. Available from: http://www.ncbi.nlm.nih.gov/pubmed/17192267

45. Combarros O, Cortina-Borja M, Smith AD, Lehmann DJ. Epistasis in sporadic Alzheimer's disease. Neurobiol Aging [Internet]. 2009 Sep [cited 2018 Oct 2];30(9):1333–49. Available from: http://www.ncbi.nlm.nih.gov/pubmed/18206267

46. Badano JL, Leitch CC, Ansley SJ, May-Simera H, Lawson S, Lewis RA, et al. Dissection of epistasis in oligogenic Bardet–Biedl syndrome. Nature [Internet]. 2006 Jan 4 [cited 2018 Oct 2];439(7074):326–30. Available from: http://www.nature.com/articles/nature04370

47. Mackay TF, Moore JH. Why epistasis is important for tackling complex human disease genetics.

Genome Med [Internet]. 2014 Jun 9 [cited 2018 Oct 2];6(6):125. Available from: http://genomemedicine.biomedcentral.com/articles/10.1186/gm561

48.   Buchner DA, Nadeau JH. Contrasting genetic architectures in different mouse reference populations used for studying complex traits. Genome Res [Internet]. 2015 Jun 1 [cited 2018 Oct 2];25(6):775–91. Available from: http://www.ncbi.nlm.nih.gov/pubmed/25953951

49.   Ackerman KG, Huang H, Grasemann H, Puma C, Singer JB, Hill AE, et al. Interacting genetic loci cause airway hyperresponsiveness. Physiol Genomics [Internet]. 2005 Mar 21 [cited 2018 Oct 2];21(1):105–11. Available from: http://www.ncbi.nlm.nih.gov/pubmed/15657107

50.   Street VA, Kujawa SG, Manichaikul A, Broman KW, Kallman JC, Shilling DJ, et al. Resistance to Noise-Induced Hearing Loss in 129S6 and MOLF Mice: Identification of Independent, Overlapping, and Interacting Chromosomal Regions. J Assoc Res Otolaryngol [Internet]. 2014 Oct 21 [cited 2018 Oct 2];15(5):721–38. Available from: http://link.springer.com/10.1007/s10162-014-0472-x

51.   Fuchs SBA, Lieder I, Stelzer G, Mazor Y, Buzhor E, Kaplan S, et al. GeneAnalytics: An Integrative Gene Set Analysis Tool for Next Generation Sequencing, RNAseq and Microarray Data. Omi A J Integr Biol. 2016;

Figure 1. Simulation Results A+B) Power calculations based on simulated data with variable main SNP effects β_G (A) or variable Ancestry-SNP effects $\beta_{Gx\theta}$ (B). Blue and Purple power curves are the power curves for detecting a significant SNP effect, while the orange and green curves are the power curves for detecting a significant $Gx\theta$ effect. Two phenotypic models, one incorporating 1 GRM (1K) correcting for relatedness in the SNPs (green, purple) and one incorporating 2 GRMs (2K) correcting for relatedness in both SNPs and Ancestry (red, blue) were used. (C+D) Effect of increasing numbers of epistatically interacting SNPs on the algorithms ability to detect a significant main SNP effect (C) or $Gx\theta$ effect (D). The straight line in C and D represents the line where expected and observed P values are the same, while the grey band indicates the region in which the P-value distribution cannot be distinguished from the expected P-value distribution.

Figure 2. Ancestral Strain Contributions by Strain. A) The 122 strains of the RI panel of the HMDP, **B)** The 1063 animals in the F_{50} – F_{56} generation of the AIL Cross

Figure 3. Gxθ Results in the Yeast, HMDP and LGxSM AIL Cohorts. A) Lithium Chloride Exposure in Yeast B) Cardiac Fibrosis and C) Left Ventricular Weight in the HMDP D) Glucose in the AIL. Significance Threshold in Yeast (A)= 1.2E-5 (FDR 5%) HMDP (B,C)= 4.2E-6(24), AIL (D)=9.1E-6 (FDR 5%)

Figure 4. Allelic Frequency Imbalances of individual sites across the genome. A) A histogram of B6 ancestry across the geneome. **B)** Genome-wide plot of ancestry **C)** Zoomed in region on chr 12 with significant depletion of B6 allele. **D)** QQ plot of all Pvalues. Genomic Inflation Factor λ=1.07

Supporting Information Legends

Figure S1. Genomic Inflation Factors of SNP and Gxθ associations with increasingly complex SNP-SNP interactions.

Table S1. All observed Gxθ associations between clinical traits and SNPs in the AIL, HMDP and Yeast cohorts

Table S2. Strains contained in the recombinant inbred panels of the HMDP and their presence in the control or isoproterenol-treated cohorts of the data.

Table S3 Alleles with significantly altered ancestries at FDR 5% in the HMDP

Table S4. Number of SNPs between strains in the AIL and HMDP founder strains as determined by the mouse diversity array. Homozygous differences are 1, heterozygous differences are 0.5

Table S5. Information on the yeast crosses used in this manuscript

Table S6-S9. Summary statistics for Gxθ results from figure 3

Table S10. All significant Gxθ associations for HMDP transcriptome data

Figure 1

Figure 2

Figure 3

Figure 4

Table 1. Most significant Gxθ associations for each phenotype observed in the AIL or HMDP cohorts as well as all Yeast Gxθ loci passing a cross-wide 5% FDR threshold See table S1 for complete list of loci. Act3.2 is 'activity levels in control animal on day 3 of a conditioned place preference test', LVIDs is 'left ventricular internal dimension at systole' and PT is post isoproterenol treatment.

Cohort	Phenotype	Chromosome	Basepair	rsID	P
AIL	Act3.2	11	93607101	rs27076868	6.22E-07
AIL	Average Body Weight	6	133867165	rs30057768	1.62E-06
AIL	Glucose	9	70318603	rs50320206	3.03E-07
AIL	Weight at 68 days	6	133899245	rs47230944	2.09E-06
HMDP	E wave 3 weeks PT	19	21360156	rs36743940	5.71E-10
HMDP	E/A ratio 3 weeks PT	6	8462274	rs47261338	7.10E-08
HMDP	Fibrosis control	17	41256115	rs6295287	6.74E-07
HMDP	Fractional shortening control	X	48286693	rs30272504	3.69E-06
HMDP	Free fatty acids control	11	113035483	rs27020574	4.09E-06
HMDP	Heart rate 1 weeks PT	18	9408717	rs29769121	2.37E-06
HMDP	Heart rate control	X	48286693	rs30272504	2.68E-06
HMDP	Left atrial weight 3 weeks PT	16	54550810	rs51319671	2.18E-06
HMDP	LVIDs control	X	48286693	rs30272504	2.92E-06
HMDP	Left ventricular mass 1 weeks PT	3	148140470	rs31313229	2.91E-06
HMDP	Left ventricular mass 2 weeks PT	17	72564961	rs50549031	8.65E-08
HMDP	Right atrial weight 3 weeks PT	16	49837267	rs50479702	4.13E-07
Yeast	Cross 2999: EGTA	XIV	622639	NA	2.91E-07
Yeast	Cross 2999: Lithium Chloride	IV	519675	NA	1.04E-07
Yeast	Cross 2999: Lithium Chloride	XII	341678	NA	1.16E-05
Yeast	Cross 2999: Manganese Sulfate	XIV	622639	NA	2.30E-06
Yeast	Cross 3043: Manganese Sulfate	XVI	206053	NA	4.17E-10
Yeast	Cross 3043: Paraquat	XV	410166	NA	3.14E-08
Yeast	Cross 3043: YNB	XV	561457	NA	3.67E-07
Yeast	Cross 3043: YPD	XV	561457	NA	2.50E-06
Yeast	Cross B: Maltose	VII	1067754	NA	1.20E-07

Table 2. **Regions with Significant and Sustained B6 Depletion or Selection**

Chromosome	Start	End	Number of SNPs	Average B6 Ancestry	Average Significance	Genes of Interest
1	10763382	10945149	21	10.66	7.96E-05	*Cpa6*
1	59458339	60046978	54	16.22	5.50E-04	*Bmpr2*
1	134463208	134712156	10	87.52	0.0048	*Nfasc*
11	64495476	64538963	5	18.18	0.0011	*Myocd*
11	110824338	110984075	18	85.75	0.0077	*Kcnj2*
12	79002481	80362465	72	18.43	0.0011	*Gphn*
12	106889886	107633641	49	19.15	0.0014	*Pigh*
15	4743169	6121864	50	18.14	0.0011	*Prkaa1*
18	5465201	8965173	246	82.50	0.017	*Epc1*