1

4 Male evolution under relaxed selection: Evidence for degeneration in sperm produced by male snails

5 from asexual lineages

7 *Joseph Jalinsky, John M. Logsdon, Jr. and Maurine Neiman*

8 *Department of Biology, University of Iowa, Iowa City, IA, 52242, USA*

9 *Corresponding author: joseph-jalinsky@uiowa.edu*

Abstract

How drastic changes in selective regimes affect trait evolution is an important open biological question.

We take advantage of naturally occurring and repeated transitions from sexual to asexual reproduction

in *Potamopyrgus antipodarum*, a New Zealand freshwater snail to address how relaxed selection on

male-specific traits influences sperm morphology. The occasional production of male offspring by the

otherwise all-female asexual lineages allows a unique and powerful opportunity to assess the fate of

sperm traits in a context where males are unnecessary. These comparisons revealed that the sperm

produced by "asexual" males are markedly distinct from sexual counterparts. In particular, the asexual

male sperm harbored markedly higher phenotypic variation and was much more likely to be grossly

abnormal. Together, these data suggest that transitions to asexual reproduction might be irreversible at

least in part because male function is likely to be compromised. More broadly, our results are consistent

with a scenario where relaxed selection translates into rapid trait degeneration.

Evolution by natural selection generates adaptation and often underlies the initial appearance and

subsequent elaboration of complex traits. While Charles Darwin was one of the very first to link

selection and adaptation, he also recognized the importance of variation away from presumed optimal

phenotypes (Darwin 1859). One of the most striking examples of this phenomenon is the eye of

Astyanax mexicanus, a freshwater fish characterized by both surface and cave ecotypes. One of the

main differences between the *A. mexicanus* surface and cave forms is that the cave dwellers have lost

eye structures, likely linked to the perpetual absence of light in their habitat (Jeffery 2005, Cartwright *et*

al. 2017). The process that yields the dismantling of once useful traits, so-called regressive (or

degenerative) evolution, is generally thought to be a consequence of a shift in selection regime and can

result in the degeneration of ancestrally important traits. Examples of trait loss *via* regressive evolution

are common in nature and across diverse taxonomic groups, including humans (Prout 1964, Wilson

1982, Remko *et al.* 2005, Ansaloni & Catena 2009; Stanley 2014). Transitions to relaxed selection are

often attributed to environmental or life-history alterations that render historically important

phenotypes irrelevant (Lahti *et al.* 2009; Ellers *et al.* 2012; Helliwell 2013).

We investigate how shifts in reproductive mode (here, sexual to asexual reproduction) alter the

selective regime experienced by traits that are used only for sexual reproduction. Our particular focus is

on the nature and extent of the loss of "maleness" in asexual lineages where females reproduce without

any need for males. In this scenario, males, and the genes underlying male-specific traits, are

presumably extraneous (Carson *et al.* 1982). This is the basis for the expectation that in these asexual

lineages, traits needed only for sexual reproduction ("sex-specific traits"; *e.g.*, genes used only for

meiosis or male-specific traits like sperm (Schurko & Logsdon 2008; Tvedte *et al.* 2017) will experience a

shift from purifying to relaxed selection at the onset of asexuality. If maintenance of these now-neutral

sex-specific traits is not otherwise costly, and if the genes underlying the sex-specific traits are not

pleiotropic with respect to traits important outside the context of sex, the sex-specific traits will

subsequently evolve primarily *via* genetic drift and mutation instead of selection (van der Kool &

Schwander 2014). These expectations have been evaluated with respect to sexual signaling in bush

crickets (Lehmann *et al.* 2007) and wasps (Pannebakker *et al.* 2005), sperm-storage organs and

fertilization success of males produced by otherwise all-female asexual lineages in stick insects

(Schwander *et al.* 2013), and sperm formation in oribatid mites (Taberly 1988; Maraun 2003; Heethoff

2007). While these studies indicate that transitions to asexuality often but not always (*e.g.* Lehmann *et*

al. 2007) underlie sex-specific trait degeneration, the conclusions that can be drawn are limited by the

inability to directly compare the sexual and asexual forms (*e.g.*, sexuals and asexuals are allopatric, are

not closely related, or the asexuals are hybrids).

How sperm and seminal fluid proteins evolve under relaxed selection is of particularly broad interest

because these traits are among the most important mediators of fertilization, sperm competition, sexual

conflict, and speciation (Swanson & Vacquier 2002). Accordingly, characterizing sperm evolution in the

context of asexuality will provide a rare glimpse into the consequences of instantaneous shifts in

selection for traits that are usually critically important determinants of evolutionary trajectories.

There are two major barriers facing the characterization of the evolution of sperm and related traits in

an asexual background. First, obligately asexual lineages derived from dioecious taxa are typically all or

nearly all females (*e.g.*, Gottlieb & Zchori-Fein 2001; Schon *et al.* 2009; Neiman *et al.* 2012). Second,

even if male asexuals exist, powerful and rigorous characterization of how relaxed selection influences

sperm evolution requires the direct comparisons that can only come from sperm produced by

ecologically and phenotypically similar males from closely related sexual lineages. Perhaps because

many asexual taxa are either of hybrid origin or do not have close sexual relatives (reviewed in Neiman

& Schwander 2011), this type of study has yet to be conducted.

We are able to overcome both of these barriers by taking advantage of the occasional production of

male offspring by obligately asexual *Potamopyrgus antipodarum* (Neiman *et al.* 2012). This New Zealand

freshwater snail provides a powerful means of directly evaluating the impact of shifts in reproductive

mode because otherwise similar obligately sexual and obligately asexual snails regularly coexist in

natural populations, enabling direct comparisons between sexual and asexual individuals, genomes, and

populations. Asexual *P. antipodarum* are the product of multiple separate transitions from sexual

ancestors, allowing us to treat distinct asexual lineages as replicated experiments into asexuality

(Neiman *et al.* 2011; Paczesniak *et al.* 2013). What makes *P. antipodarum* particularly well suited for this

study is that asexual female lineages occasionally (~1.5% of all offspring, depending on the asexual

lineage involved; Neiman *et al.* 2012) produce males, perhaps *via* loss of sex chromosomes during

oogenesis (Soper *et al.* 2013). We leverage this unusual phenomenon to characterize morphological

variation of sperm in sexually *vs.* asexually produced males, with the expectation that the latter have

experienced a relatively recent history of relaxed selection on male-specific traits and genes. Based on

this logic, we predict that the sperm produced by asexual males will harbor more variation and be more

likely to be abnormal than sperm from their sexual counterparts.

Materials and Methods

Snail collection and ploidy/reproductive mode determination

Potamopyrgus antipodarum used in this study were collected either from New Zealand lake populations

(11 lakes; 33 sexual males) or from individual and separately derived asexual lineages (8 lineages; 24

asexual males). The latter are recently descended (<5 generations) from asexual snails collected from

New Zealand lake populations that were subsequently cultured in our laboratory at the University of

Iowa. All snails were housed in plastic 10L tanks filled with oxygenated carbon-filtered tap water in a

16C room with a 16:8-hr light:dark cycle and fed dried *Spirulina* algae *ad libitum* (following Zachar &

Neiman 2013). We focused our search for asexual males in lineages that had been previously

characterized as relatively likely to harbor asexual males (Neiman *et al.* 2012). We also emphasized

asexual lineages descended from lake populations that population genetic studies (Paczesniak *et al.*

2013) demonstrated harbor asexual subpopulations that represent separate *in situ* transitions from

sexual to asexual reproduction. For the sexual males, we used field collections from lakes that were

known to contain relatively high frequencies (>10-20%) of sexual males (Neiman *et al.* 2011).

We identified males by sexing each individual from each field collection (N = 1370 snails) or lineage (N =

4505 snails) that was large enough (>/=3mm shell length) to have reached sexual maturity (*e.g.*, Larkin

et al. 2016). Sexual *P. antipodarum* are diploid and asexuals are either triploid or tetraploid (Neiman *et*

al. 2011), allowing us to use comparisons between triploid and tetraploid-produced sperm to at least in

part control for effects of asexuality *vs.* ploidy elevation on sperm phenotype. We used a Becton

Dickinson ARIA II flow cytometer on dissected head tissue (following Bankers *et al.* 2017) to assign ploidy

(diploid, triploid, tetraploid) and, thus, reproductive mode (diploid = sexual; triploid, tetraploid =

asexual) to each male. The overall incidence of males in asexual lineages was ~1% (59/5875), and

asexual lineages varied in the relative frequency of male production (Supp Table 1). Both observations

are consistent with a similar survey in Neiman *et al*. (2012). Because our goal was to replicate males

three times within each asexual lineage that we used, we excluded 35 males that either represented a

lineage in which more than three males were found (13 asexual lineages; these "extra" males were

excluded arbitrarily) or were from a lineage in which fewer than three males were found (six asexual

lineages). This left us with a total of 24 asexually produced male *P. antipodarum* for our study. For sexual

males, we used the first three arbitrarily selected sexual males that we identified per each of seven field

collections, for a total of 21 sexual male *P. antipodarum*. Hereafter, we use "set" to indicate one three-

male replicate from each unique lineage or lake of origin and ploidy combination: our diploid *P.*

antipodarum sample represents seven male sets (each from a different field collection), triploid *P.*

antipodarum are represented by six male sets (each from a different asexual lineage and lake of origin),

and tetraploid *P. antipodarum* are represented by two male sets (from two different asexual lineages,

each from a different lake), for a total of 15 *P. antipodarum* sets and 45 males (Figure 1).

We provided a separate control for nuclear DNA content, which can have major influences on sperm

phenotype (Alavi *et al.* 2013), by including males and sperm from the closely related (Haase 2008) and

diploid obligate sexual species *P. estuarinus* (three sets, for a total of nine males) and *Halopyrgus*

pupoides (one set, three males)*. Because *P. estuarinus* and *H. pupoides* have only about 60% of the

nuclear genome DNA content of diploid *P. antipodarum* (Logsdon *et al.* 2017), we used comparisons

across these three diploid taxa to account for the effect of nuclear genome DNA content on sperm

phenotype. These outgroup snails were collected from their native New Zealand estuary habitat

(Waikuku Beach, New Zealand) in January 2018.

Sperm extraction and imaging

We used forceps to extract the bodies of male snails from their shells and then dissected away the

sperm duct, which transports sperm and seminal fluid from the testis to the penis. We centered the

sperm duct tissue on a glass microscope slide and added ~5µl of DAPI staining solution (following Soper

et al. 2013). We used repeated pipetting of this solution to release sperm from the ruptured duct and

distribute the sperm across the slide. We then mounted the slide on an Olympus IX71 fluorescence

microscope and used the DAPI filter to locate the fluorescent sperm heads and the polarized filter to

photograph sperm for subsequent analysis. We arbitrarily selected and then photographed a minimum

of 12 whole sperm cells for every male. We excluded sperm cells that were entangled with other sperm

or were obscured by non-sperm cellular detritus. We used sperm from at least three males (one male

set) for each unique lake and ploidy combination, for a total of at least 36 sperm per set.

Phenotype measurements

We imported photographs into ImageJ (Rueden *et al.* 2017) and measured head length (HL; head tip to

head base), head width (HW; width at midpoint of head), head circumference (HC; perimeter of head),

and tail length (TL; length from head base to tail end) for each sperm cell. We also used Mahalanobis

distance (MD; distance between a distribution and a focal point) to provide a multi-dimensional

generalization that represents a composite of all four of the sperm traits that we measured (De

Maesschalck *et al.* 2000). Finally, we used our consistent observations of a number of qualitatively

distinct phenotypic classes of sperm abnormalities to sort sperm into "normal" (most common

phenotype of smooth cylindrical head and smooth tail of consistent width; 79% of all sperm) *vs.* one of

six "abnormal" ultrastructural categories: >1 tail, distorted neck, oblong head, tail bump, two heads,

wrinkled head (*e.g.*, Menkveld *et al.* 1990). This approach allows us to distinguish large-scale differences

in complex sperm morphology that might not be evident from our measurements of individual sperm

traits.

Statistical analyses

We used a RAxML tree generated from Illumina whole-genome sequencing data from 26 *P. antipodarum*

(10 sexual individuals, 16 asexual individuals (11 triploid and 5 tetraploid)) to provide a phylogenetic

framework needed to control for potential effects of phylogenetic relationship on sperm phenotype

(Supp Figure 1). This tree also demonstrates that we have likely captured a broad sample of sexual and

asexual *P. antipodarum* genetic diversity, including a number of separately derived asexual lineages. We

used this tree in Phylosignal (Keck *et al.* 2016), an R package that tests for phylogenetic signal, to assess

whether sperm morphology was determined at least in part by shared ancestry. We attached mean trait

values for each of the sperm traits measured below to a subsampled phylogenetic tree, which

represented directly (same asexual lineage) or indirectly (shared lake of origin, justified by multiple lines

of evidence for a predominant effect of source population on *P. antipodarum* genotype, *e.g.*, Dybdahl &

Lively 1995; Paczesniak *et al.* 2013; Bankers *et al*. 2017) nine of the 15 *P. antipodarum* sets from which

we sampled males (*i.e.*, sets with both genetic data and sperm measurements). We did not find any

significant phylogenetic correlations underlying these sperm traits (Supp Figure 2). Together, the

outcome of these analyses indicate that phylogenetic relatedness is not a main driver of variation in

sperm morphology. We also used hierarchical clustering of sperm morphology data to produce a

cladogram that we used to assess whether phylogenetic relatedness is the primary driver of across-

lineage/lake sperm variation (Figure 2, Supp Figure 3).

We applied ANOVA and MANOVA models to evaluate how lake/lineage of origin affected each of the

four individual sperm traits that we measured (HL, HW, HC, TL) along with the composite MD trait. We

first used M/ANOVA for the trait values estimated for each set to assess whether the three different

males within each set varied in sperm morphology. We then applied M/ANOVA to groupings of diploid,

triploid, and tetraploid *P. antipodarum* to evaluate whether ploidy affects sperm morphology. Finally,

we used an unpaired *t*-test to compare means for sexual and asexual *P. antipodarum* sperm traits to

assess the effect of reproductive mode on sperm morphology. Prior to each of the M/ANOVA analyses,

we used a Shapiro-Wilk test applying a Bonferroni-corrected threshold *p*-value of 0.0026 to account for

multiple comparisons (19 within-set comparisons per each of the 95 Shapiro-Wilk tests) for each of the

five traits to evaluate whether the data met the assumption of a normal distribution. For the 35 tests

that violated this assumption, we applied a Box-Cox transformation to attempt to achieve normality,

which was successful in 14 cases. For the 21 cases that still violated normality assumptions, we used

Kruskal-Wallis tests to compare population medians. We then used principal component analysis (PCA)

to visualize among-species (*P. antipodarum*, *P. estuarinus*, *H. pupoides*) differences that maximize

variance for the four multidimensional sperm trait values in order to detect species-specific differences.

Finally, we used Linear Discriminate Analysis (LDA) by grouping diploid, triploid, and tetraploid *P.*

antipodarum into predefined classes. This analysis minimizes within-ploidy variance and maximizes

across-ploidy variance to find linear combinations of variables that best explain the data. We used this

analysis to assess how differing ploidy level translates to differences in sperm morphology. All statistical

analyses were implemented in R 3.4.1 (R Core Team 2013) or SPSS 24.0 (IBM Corp. 2016)

Results

We evaluated a total of 750 sperm from each of 57 male snails (21 diploid sexuals, 18 triploid asexuals,

six tetraploid asexuals, nine *P. estuarinus*, three *H. pupoides* (Figure 1). representing 12 lakes and three

estuaries of origin as well as three species. Because we did not find any asexual male in field collections

despite evaluating >380 field-collected male *P. antipodarum* from eight lake populations, we had to

compare field-collected sexual individuals to laboratory-reared asexual males. While we cannot formerly

exclude the possibility that these differences in rearing environment between sexually and asexually

produced males could influence sperm production, we felt that it was important to only use field-

collected sexual males because of the distinct possibility that the inbreeding that is a necessary

component of lab culture of sexual lineages of these snails would influence snail phenotype in a manner

that does not reflect natural variation. Because asexual *P. antipodarum* reproduce via apomictic

parthenogenesis (Phillips & Lambert 1998), inbreeding, and inbreeding depression, is not a concern for

laboratory-reared asexuals. Because male-specific traits that are encoded in the genomes of asexual

lineages presumably experience weak or even no selection, we also believe that it is unlikely - though

not impossible - that laboratory-imposed selection would affect these male traits in a manner that is

qualitatively different than male traits harbored in asexual lineages in the field.

Our analyses revealed extensive qualitative (Supp Figure 4) and quantitative variation (Supp Table 2) in

sperm morphology. Multi-dimensional hierarchical clustering of primary sperm traits (HL, HW, HC, and

TL) suggested that reproductive mode (*vs.* ploidy or shared phylogenetic history) is the primary driver of

sperm morphology in *P. antipodarum* (Figure 2, Supp Figure 3). This conclusion is bolstered by the fact

that diploid sexual *P. antipodarum* sperm clustered with the two diploid sexual outgroup species instead

of with the asexual polyploid conspecifics.

Sperm morphology is primarily driven by reproductive mode

We used comparisons across the three ploidy levels represented by *P. antipodarum* and sperm from the

diploid outgroup species *P. estuarinus* and *H. pupoides* to account for effects of nuclear genome DNA

content on sperm morphology. That nuclear genome DNA content *per se* is not the sole driver of sperm

morphology in these snails is demonstrated by our findings that 1) sperm produced by *H. pupoides* and

P. estuarinus, which seem to share the same relatively small nuclear DNA content, were nevertheless

morphologically distinct (Figure 2, Supp Figure 5), 2) head width and tail length were statistically

indistinguishable between triploid and tetraploid *P. antipodarum* (Supp Figure 6), 3) the CV for triploid

and tetraploid *P. antipodarum* is statistically indistinguishable for all traits except tail length (Supp Figure

7) and 4) different ploidies overlap in morphometric space (Figure 3). We did detect some evidence for

ploidy effects on sperm trait means and variances: the LDA revealed ploidy-based clustering (Figure 3),

and the ANOVA demonstrated a main effect of ploidy on all sperm traits (Supp Table 2 B). Our CV-based

analyses indicated that sperm produced by diploid male *P. antipodarum* harbors significantly lower

variance (*e.g.*, HL mean = 4.48μm, SD = 0.17) than sperm produced by triploid (HL mean = 5.29μm, SD =

0.44) and tetraploid (HL mean = 5.99μm, SD = 0.38) males (Supp Figure 7).

Asexual *P. antipodarum* often produce abnormal sperm

Visual inspection of *P. antipodarum* sperm revealed substantial variation. Most (N=429; 76%) of these

sperm had an overtly normal phenotype, represented by a smooth and cylindrical head and a flagellum

of consistent width from the neck to tip (Supp Figure 4 A). We sorted the most common deviations we

observed from this normal phenotype into six discrete categories that emerged during our imaging and

analysis stages: >1 tail (3.5% of all sperm), distorted neck (2.5%), oblong head (1.5%), tail bump (10%),

two heads (1%), and wrinkled head (6.5%) (see Supp Figure 4 B-D). There was markedly higher incidence

of abnormal sperm produced by triploid (N=95; 43% of all triploid sperm) and tetraploid (N=24; 27%) *P.*

antipodarum relative to the sperm produced by diploid *male P. antipodarum* (N=16; 6%) (Fisher's exact

test p = < 0.001) (Figure 4). Misshapen heads were the only abnormality shared across all three ploidies,

while the most common abnormality, tail bump, is restricted to triploid asexual *P. antipodarum*.

Among- and within-species variation in sperm morphology

We used a principal component analysis to visualize and compare multivariate sperm phenotypes across

sexual *P. antipodarum* and the two closely related sexual outgroup species. This analysis revealed a

distinct species-level signal, with no overlap between the clusters of sperm data corresponding to each

species (Supp Figure 5). Our ANOVA and MANOVA analyses demonstrated statistically significant

differences in each of the five sperm traits amongst these three species (HL: p = 0.0005, HW: p = 0.001,

HC: p = 0.01, TL: p = 7.6e-08, MD: p = 4.5e-13).

We used a MANOVA to address whether there was intraspecific (individual level) variation

within each of the 45 *P. antipodarum* male sets. This analysis revealed significant across-male variation

within four diploid and four triploid sets (Supp Table 2). Sperm tail length was the trait most often

variable across sets (5/19 populations), and sperm head circumference was the least likely to feature

across-set variation (1/19). The MD analyses revealed only two cases of significant within-population

variation (asexual Haupiri and asexual Brunner). The CV analyses indicated that sperm head length and

head circumference are the least variable traits and head width and tail length are the most variable

traits across all *P. antipodarum* sperm (Supp Figure 8).

Discussion

We used the powerful *Potamopyrgus antipodarum* system to perform a direct comparison of male-

specific traits across otherwise similar males that differ in ploidy level and reproductive mode. We found

evidence for a major effect of sexual *vs*. asexual reproduction on sperm morphology, with sperm

produced by asexual males harboring substantially more variation and much more likely to feature

large-scale structural abnormalities. That a substantial fraction of asexual-produced sperm is abnormal

is consistent with a scenario where male-specific traits are no longer useful in an asexual lineage. We

also used comparisons between the three different *P. antipodarum* ploidies and three different diploid

sexual taxa included in our study to assess the extent to which increased nuclear genome DNA content

in the asexual *P. antipodarum*, *versus* asexuality *per se*, might affect sperm morphology. That asexual-

produced sperm are generally larger, and that these size differences to some extent scale with ploidy

(Supp Figure 6) is consistent with this possibility. Even so, the significant differences in sperm trait values

(*e.g.*, head length) between *P. estuarinus* and *H. pupoides*, which have virtually identical genome sizes

(Logsdon *et al.* 2017), indicates that other factors besides nuclear genome content are also important in

determining sperm morphology.

Our study is especially novel in comparing sperm phenotypes across reproductive modes as well as

ploidy level, allowing us to at least in part decouple effects of polyploidy from those of asexuality. We

can draw parallels and contrasts with other ploidy-focused sperm studies: Alavi *et al.* (2013) showed

that tetraploid European weatherfish (*Misgurnus fossilis*) produce sperm that are 4.8% larger than

sperm produced by their triploid counterparts, and Flajshans *et al.* (2008) found that triploid male

Prussian carp (*Carassius gibelio*) make sperm with larger volume than sperm generated by diploid and

tetraploid males. Considered together along with these two earlier studies, we can use our results to

conclude that nuclear genome DNA content is not the sole driver of the patterns we observed. Our

study is especially novel in allowing us to compare both across different ploidy levels as well as among

lineages with different nuclear genome content but the same ploidy level. We were also able to exclude

a major role of phylogeny in driving any of the patterns that our analyses revealed. Together,

comparisons provide a powerful means of disentangling effects of reproductive mode from nuclear

genome DNA content from ploidy elevation.

The abnormal phenotypes that often characterized sperm from asexual male *P. antipodarum* do appear

to be affected in part by ploidy: sperm from triploid asexual males spanned a wider range of abnormal

phenotypes and were more than twice as likely to be abnormal than sperm from tetraploid

counterparts. Cytogenetic characterization of some components of spermatogenesis in diploid sexual *vs.*

triploid asexual *P. antipodarum* have demonstrated that sperm production in the latter is associated

with aberrant meiotic pairing (Soper *et al.* 2013). Meiosis is disrupted when odd number of

chromosomes pair (Gupta 1985; Comai 2005; Henry *et al.* 2005), hinting that meiotic disruption is one

possible explanation for the differences between triploid and tetraploid-produced sperm in *P.*

antipodarum.

The morphologically distinct and often abnormal sperm produced by asexual *P. antipodarum* could also

be at least in part a consequence of mutation accumulation in the genes underlying sperm production,

which are expected to be under relaxed selection in all (or nearly all)-female asexual populations

(Schurko & Logsdon 2008; Lahti *et al.* 2009). Under this scenario, genes that influence sperm traits in

asexuals should be accumulating mutations at a higher rate than sexuals. This mutation accumulation

process would be evidenced by the degeneration and eventual pseudogenization of genes and traits

related to sperm function in asexual populations.

Our discovery that asexual-produced sperm harbor more morphological variation and are more likely to

be abnormal than sexually produced sperm is consistent with this expectation. The fact that the male

offspring occasionally produced by female asexual *P. antipodarum* are still investing in sperm suggests

that the time period over which the genes underlying sperm production have been subject to relaxed

selection is relatively short and/or that selection favoring the loss of this trait is weak. The latter would

not be surprising in light of the fact that male production is quite rare (~<1.5% of all offspring) in asexual

P. antipodarum. A powerful test of the former could come from comparisons of males produced by

different asexual *P. antipodarum* lineages, which are often recently derived (Paczesniak *et al.* 2013) but

also vary in the time since derivation from sexual *P. antipodarum* (Neiman *et al.* 2005). If time since

derivation to asexuality is a key determinant of the extent of male trait degeneration, we predict that

males from older asexual lineages should tend to produce sperm that harbors relatively high variance

and is abnormal at a higher frequency than sperm produced by males from younger asexual lineages.

While we cannot formally exclude the possibility that ploidy elevation *per se* is a major driver of sperm

abnormality, the fact that diploid male sexual *P. antipodarum* produce normal sperm at a similar

frequency to the two outgroup species suggests that nuclear genome DNA content alone does not

account for the high frequency of sperm abnormalities observed in asexual male *P. antipodarum*. The

wide sperm phenotype variation that we observed across asexual males also hints at a process (*i.e.*

mutation accumulation) that is stochastic compared to a direct consequence of ploidy elevation.

Together, our results indicate that asexual male *P. antipodarum* are likely experiencing decay of male-

specific traits, highlighting the likelihood that transitions to asexual reproduction results in striking

differences in selective regimes between sexual and asexual populations. This finding is of direct

relevance to evaluating how relaxed selection influences phenotypes in natural populations and

suggests that transitions back to sexual reproduction in asexual populations are likely to face challenges

associated with degeneration of sex-specific traits. Critical next steps include the characterization of

molecular evolution of sperm-related genes in sexual *vs.* asexual populations and assessment of

whether older asexual populations have experienced a relatively high level of trait decay.

Acknowledgments

We gratefully acknowledge help in snail collections from Mike Winterbourn, Mary Morgan-Richards,

Katelyn Larkin, Kyle McElroy, Jeremy Richardson, Laura Bankers, and Kaitlin Hatcher. We thank Sarit

Smolikove for microscope use, Julie Meachen for statistical advice, and Deanna Soper for assistance in

sperm extraction protocols. The ploidy identification data presented herein were obtained at the Flow

Cytometry Facility, which is a Carver College of Medicine/Holden Comprehensive Cancer Center core

research facility at the University of Iowa. The Facility is funded through user fees and the generous

financial support of the Carver College of Medicine, Holden Comprehensive Cancer Center, and Iowa

City Veteran's Administration Medical Center. This project was funded by NSF-MCB 1122176 and NSF

DEB-1753851

References Cited

Alavi, Sayyed Mohammad Hadi, *et al.* "Sperm morphology, motility, and velocity in naturally occurring

polyploid European weatherfish (*Misgurnus fossilis* L.)." *Theriogenology* 80.2 (2013): 153-160.

Altmann, Thomas, *et al.* "Easy determination of ploidy level in *Arabidopsis thaliana* plants by means of

pollen size measurement." *Plant Cell Reports* 13.11 (1994): 652-656.

Ansaloni, Luca, F. Catena, and A. D. Pinna. "What is the function of the human vermiform

appendix?." *European Surgical Research* 43.2 (2009): 67-71.

Bankers, Laura, *et al.* "Genomic evidence for population-specific responses to co-evolving parasites in a

New Zealand freshwater snail." *Molecular Ecology* 26.14 (2017): 3663-3675.

Bast, Jens, *et al.* "Consequences of asexuality in natural populations: insights from stick

insects." *Molecular Biology and Evolution* 35.7 (2018): 1668-1677.

Berg, Johannes, Stana Willmann, and Michael Lässig. "Adaptive evolution of transcription factor binding

sites." *BMC Evolutionary Biology* 4.1 (2004): 42.

Carson, Hampton L., L. S. Chang, and T. W. Lyttle. "Decay of female sexual behavior under

parthenogenesis." *Science* 218.4567 (1982): 68-70.

Cartwright, Reed A., *et al.* "The importance of selection in the evolution of blindness in cavefish." *BMC*

Evolutionary Biology 17.1 (2017): 45.

Comai, Luca. "The advantages and disadvantages of being polyploid." *Nature Reviews Genetics* 6.11

(2005): 836.

Darwin, Charles. "On the origin of the species by natural selection." (1859).

De Maesschalck, Roy, D. Jouan-Rimbaud, and D. L. Massart. "The mahalanobis distance." *Chemometrics*

and Intelligent Laboratory Systems 50.1 (2000): 1-18.

Doniger, Scott W., J. Huh, and J. C. Fay. "Identification of functional transcription factor binding sites

using closely related *Saccharomyces* species." *Genome Research* 15.5 (2005): 701-709.

Dybdahl, Mark F., and C. M. Lively. "Diverse, endemic and polyphyletic clones in mixed populations of a

freshwater snail (*Potamopyrgus antipodarum*)." *Journal of Evolutionary Biology*8.3 (1995): 385-398.

Ellers, Jacintha, *et al.* "Ecological interactions drive evolutionary loss of traits." *Ecology Letters* 15.10

(2012): 1071-1082.

Fong, Daniel W., T. C. Kane, and D. C. Culver. "Vestigialization and loss of nonfunctional

characters." *Annual Review of Ecology and Systematics* 26.1 (1995): 249-268.

Flajšhans, Martin, *et al.* "Characteristics of sperm of polyploid Prussian carp *Carassius gibelio." Journal of*

Fish Biology 73.1 (2008): 323-328.

Foulkes, M. Raine. "Pollen size as an indicator of ploidy level in rhexia, a small genus of flowering plants

endemic to the North American coastal plain, in north Florida and coastal Georgia." *Georgia Journal of*

Science 76.1 (2018): 93.

Gottlieb, Yuval, and E. Zchori-Fein. "Irreversible thelytokous reproduction in *Muscidifurax*

uniraptor." *Entomologia Experimentalis et Applicata* 100.3 (2001): 271-278.

Gould, Frank W. "Pollen size as related to polyploidy and speciation in the *andropogon saccharoides—A.*

barbinodis complex." *Brittonia* 9.2 (1957): 71-75.

Gupta, Prabhat, and G. Fedak. "Genetic control of meiotic chromosome pairing in polyploids in the

genus *Hordeum*." *Canadian Journal of Genetics and Cytology* 27.5 (1985): 515-530.

Heethoff, Michael, *et al.* "High genetic divergences indicate ancient separation of parthenogenetic

lineages of the oribatid mite *Platynothrus peltifer* (Acari, Oribatida)." *Journal of Evolutionary*

Biology 20.1 (2007): 392-402.

Helliwell, Katherine E., G. L. Wheeler, and A. G. Smith. "Widespread decay of vitamin-related pathways:

coincidence or consequence?." *Trends in Genetics* 29.8 (2013): 469-478.

Henry, Isabelle M., *et al.* "Aneuploidy and genetic variation in the *Arabidopsis thaliana* triploid

response." *Genetics* 170.4 (2005): 1979-1988.

IBM Corp. IBM SPSS Statistics for Windows, Version 24.0. Armonk, NY: IBM Corp.

Jeffery, William R. "Regressive evolution in *Astyanax* cavefish." *Annual Review of Genetics* 43 (2009): 25-

47.

Lahti, David C., *et al.* "Relaxed selection in the wild." *Trends in Ecology & Evolution* 24.9 (2009): 487-496.

Larkin, Katelyn, C. Tucci, and M. Neiman. "Effects of polyploidy and reproductive mode on life history

trait expression." *Ecology and Evolution* 6.3 (2016): 765-778.

Law, Jennifer H., and B. J. Crespi. "The evolution of geographic parthenogenesis in *Timema* walking-

sticks." *Molecular Ecology* 11.8 (2002): 1471-1489.

Lehmann, Gerlind, U.C. Johannes Strauß, and R. Lakes-Harlan. "Listening when there is no sexual

signalling? Maintenance of hearing in the asexual bushcricket *Poecilimon intermedius*." *Journal of*

Comparative Physiology A 193.5 (2007): 537-545.

Maraun, Mark, *et al.* "Radiation in sexual and parthenogenetic oribatid mites (Oribatida, Acari) as

indicated by genetic divergence of closely related species." *Experimental & Applied Acarology* 29.3-4

(2003): 265-277.

Marinho, Ricardo, *et al.* "Pollen and stomata morphometrics and polyploidy in Eriotheca (Malvaceae-

Bombacoideae)." *Plant Biology* 16.2 (2014): 508-511.

Muller, Hermann Joseph. "Some genetic aspects of sex." *The American Naturalist* 66.703 (1932): 118-

138.

Neiman, Maurine, J. Jokela, and C. M. Lively. "Variation in asexual lineage age in *Potamopyrgus*

antipodarum, a New Zealand snail." *Evolution* 59.9 (2005): 1945-1952.

Neiman, Maurine, S. Meirmans, and P. G. Meirmans. "What can asexual lineage age tell us about the

maintenance of sex?." *Annals of the New York Academy of Sciences* 1168.1 (2009): 185-200.

Neiman, Maurine, and T. Schwander. "Using parthenogenetic lineages to identify advantages of

sex." *Evolutionary Biology* 38.2 (2011): 115-123.

Neiman, Maurine, *et al.* "Wide variation in ploidy level and genome size in a New Zealand freshwater

snail with coexisting sexual and asexual lineages." *Evolution* 65.11 (2011): 3202-3216.

Neiman, Maurine, *et al.* "Male offspring production by asexual *Potamopyrgus antipodarum*, a New

Zealand snail." *Heredity* 109.1 (2012): 57.

Neiman, Maurine, and T. Schwander. "Using parthenogenetic lineages to identify advantages of sex."

Evolutionary Biology 38 (2011):115-123.

Ohta, Tomoko. "Near-neutrality in evolution of genes and gene regulation." *Proceedings of the National*

Academy of Sciences 99.25 (2002): 16134-16137.

Paczesniak, Dorota, *et al.* "Discordance between nuclear and mitochondrial genomes in sexual and

asexual lineages of the freshwater snail *Potamopyrgus antipodarum*." *Molecular Ecology* 22.18 (2013):

4695-4710.

Pannebakker, Bart, *et al.* "Sexual functionality of *Leptopilina clavipes* (Hymenoptera: Figitidae) after

reversing *Wolbachia*-induced parthenogenesis." *Journal of Evolutionary Biology*1 8.4 (2005): 1019-1028.

Phillips, Ngaire R., and D. M. Lambert. "Genetics of *Potamopyrgus antipodarum* (Gastropoda:

Prosobranchia): evidence for reproductive modes." *New Zealand Journal of Zoology* 16.3 (1989): 435-

445.

Prout, Timothy. "Observations on structural reduction in evolution." *The American Naturalist* 98.901

(1964): 239-249.

Leys, Remko, *et al.* "Regressive evolution of an eye pigment gene in independently evolved eyeless

subterranean diving beetles." *Biology Letters* 1.4 (2005): 496-499.

Rueden, Curtis T., *et al.* "ImageJ2: ImageJ for the next generation of scientific image data." *BMC*

Bioinformatics 18.1 (2017): 529.

Schwander, Tanja, and B. J. Crespi. "Multiple direct transitions from sexual reproduction to apomictic

parthenogenesis in *Timema* stick insects." *Evolution* 63.1 (2009): 84-103.

Schurko, Andrew M., and J. M. Logsdon. "Using a meiosis detection toolkit to investigate ancient asexual

"scandals" and the evolution of sex." *Bioessays* 30.6 (2008): 579-589.

Schön, Isa, G. Rossetti, and K. Martens. "Darwinulid ostracods: ancient asexual scandals or scandalous

gossip?." *Lost Sex*. Springer, Dordrecht (2009): 217-240.

Schwander, Tanja, *et al.* "Neutral and selection-driven decay of sexual traits in asexual stick

insects." *Proceedings of the Royal Society of London B: Biological Sciences* 280.1764 (2013): 20130823.

Soper, Deanna M., *et al.* "Spermatozoa production by triploid males in the New Zealand freshwater snail

Potamopyrgus antipodarum." *Biological Journal of the Linnean Society* 110.1 (2013): 227-234.

Stanley, Shane Gareth. *Vestigial hind limbs in the appendicular skeleton of three genera and four species*

of African skink (Squamata, Scincidae). Diss. Fayetteville State University, 2014.

Taberly, Georges. "Researchs on the thelytokous parthenogenesis of two species of Oribatid mites:

Trhypochthonius tectorum(Berlese) and *Platynothrus peltifer*(Koch). IV. Observations on atavic

males." *Acarologia* 29.1 (1988): 95-112.

Tvedte, Eric S., A. A. Forbes, and J. M. Logsdon Jr. "Retention of core meiotic genes across diverse

Hymenoptera." *Journal of Heredity* 108.7 (2017): 791-806.

van der Kooi, Casper J., and Tanja Schwander. "On the fate of sexual traits under asexuality." *Biological*

Reviews 89.4 (2014): 805-819.

Villeneuve, Anne M., and Kenneth J. Hillers. "Whence meiosis?." *Cell* 106.6 (2001): 647-650.

Wilson, Carl L. "Vestigial structures and the flower." *American Journal of Botany* (1982): 1356-1365.

Wright, Sewall. "Pleiotropy in the evolution of structural reduction and of dominance." *The American*

Naturalist 98.899 (1964): 65-69.

Zachar, Nicholas, and M. Neiman. "Profound effects of population density on fitness-related traits in an

invasive freshwater snail." *PloS One* 8.11 (2013): e80067.

Figure Legends

Figure 1. Schematic representing our experimental design, which included sperm from 57 individual

males (each male represented by small colored boxes) from three different species, two reproductive

modes, three ploidy levels, 12 lakes, and three estuaries. We characterized at least 12 sperm per male

and three males per set, for a total of at least 36 sperm per set. Each set is represented by the boxed

lake of origin and three small colored boxes representing the three males in the set. *P. est = P.*

estuarinus from three different New Zealand estuaries (CC = Waikuku Beach; Hutt = Hutt River, NZ; Fox =

Foxton). *H. pupoides* was collected from Waikuku Beach.

Figure 2. Hierarchical clustering dendrogram of sperm head morphology. Each square represents the

mean trait values of the >/= 36 sperm taken from each of the three males used for each unique

ploidy/reproductive mode/population set. Labeling follows Figure 1.

Figure 3. Linear discriminant function analysis of transformed sperm traits (head length, head width,

head circumference, tail length) for diploid, triploid, and tetraploid *P. antipodarum*. Histogram

represents the cumulative numbers of sperm in each bin for LD1. Arrows represent each variable (HL,

HW, HC, TL), length of arrows represents degree of discrimination, and arrow angle is the direction of

the variables in 2-D space.

Figure 4. Proportion and characterization of sperm phenotypes that we categorized as "abnormal".

Figure 1. Schematic representing our experimental design, which included sperm from 57 individual males (each male represented by small colored boxes) from three different species, two reproductive modes, three ploidy levels, 12 lakes, and three estuaries. We characterized at least 12 sperm per male and three males per set, for a total of at least 36 sperm per set. Each set is represented by the boxed lake of origin and three small colored boxes representing the three males in the set. *P. est* = *P. estuarinus* from three different New Zealand estuaries (CC = Waikuku Beach; Hutt = Hutt River, NZ; Fox = Foxton). *H. pupoides* was collected from Waikuku Beach.

Figure 2. Hierarchical clustering dendrogram of sperm head morphology. Each square represents the mean trait values of the >/= 36 sperm taken from each of the three males used for each unique ploidy/reproductive mode/population set. Labeling follows Fig. 1.

Figure 3. Linear discriminant function analysis of transformed sperm traits (head length, head width, head circumference, tail length) for diploid, triploid, and tetraploid *P. antipodarum*. Histogram represents the cumulative numbers of sperm in each bin for LD1. Arrows represent each variable (HL, HW, HC, TL), length of arrows represents degree of discrimination, and arrow angle is the direction of the variables in 2-D space.

Figure 4. Proportion and characterization of sperm phenotypes that we categorized as "abnormal".