Rift Valley fever virus minigenome system for investigating the role of L protein residues in viral transcription and replication

Hanna Jérôme[1], Martin Rudolf[1], Michaela Lelke[1], Meike Pahlmann[1], Carola Busch[1], Sabrina Bockholt[1], Stephanie Wurr[1], Stephan Günther[1,2]*, Maria Rosenthal[1&], Romy Kerber[1,2&]

[1] Department of Virology, Bernhard Nocht Institute for Tropical Medicine, Hamburg, Germany

[2] German Center for Infection Research (DZIF), Partner site Hamburg – Lübeck – Borstel – Riems, Hamburg, Germany

* Corresponding author

E-mail: guenther@bni.uni-hamburg.de (SG)

& These senior authors contributed equally to this work.

Short title: An ambisense minigenome system for Rift Valley fever virus

Abstract

Replicon systems are important molecular tools for investigating the function of virus proteins and regulatory elements involved in viral RNA synthesis. We developed an ambisense minigenome system for Rift Valley fever virus (RVFV) to dissect the effects of L gene mutations on viral transcription versus replication. The S RNA segment with regulatory elements for ambisense gene expression served as backbone for the minigenome. Expression of the luciferase reporter gene allowed the overall activity of the RVFV replication complex to be assessed, while northern blot analysis enabled differentiation between synthesis of viral mRNA and replication intermediates. The functionality of the system was demonstrated by probing residues predictably involved in the active site of the cap-snatching endonuclease in the N-terminus of the L protein. Corresponding mutations led to a selective defect in the viral mRNA synthesis as described for other viruses of the *Bunyavirales* order. The analysis of further L gene mutants revealed an essential and specific role of a C-terminal region in the RVFV L protein in viral transcription. In summary, the established minigenome system is suitable for functional testing of the relevance of residues for viral transcription and replication. It may be used to validate hypotheses arising from structural or biochemical investigations of the RVFV replication complex.

Introduction

Rift Valley fever virus (RVFV) is an important human and animal pathogen in Sub-Saharan Africa and on the Arabian Peninsula. RVFV infections cause death and abortion in ruminants and pseudoruminants and outbreaks may be associated with a high economic burden. In humans, the virus may cause febrile illness including hemorrhagic fever with fatal outcome [1, 2]. As medical countermeasures to prevent or treat the disease in humans are lacking, it is listed on the WHO blueprint for urgent research and development [3].

RVFV belongs to the family of *Phenuiviridae* within the *Bunyavirales* order and contains a tri-segmented single stranded RNA genome with negative polarity. The small (S) RNA segment utilizes an ambisense coding strategy; the nucleocapsid protein (N) is encoded in antisense orientation and the non-structural protein (NSs) in sense orientation. Both genes are separated by an intergenic region (IGR). The middle and large (M and L) RNA segments contain genes for the glycoprotein precursor, the nonstructural protein NSm, and the large L protein (~250 kDa), respectively. N and L proteins together with the viral RNA constitute the viral replication complexe, the structural unit for genome replication and transcription [4].

Several minigenome systems have been established for RVFV and were used to study the function of proteins and regulatory elements involved in RVFV genome replication and transcription [5-9]. However, these systems have not been optimized for comprehensive mutagenesis studies aiming at measuring simultaneously genome replication and transcription: (i) the measurement range (i.e. the ratio between positive and negative control) of the reporter gene assay is rather small [7, 9], although performance could be improved upon depletion of cellular protein kinase R [9], (ii) the read-out requires complex experimental procedures (i.e. virus-like particle production and transfer on indicator cells, chloramphenicol acetyltransferase [CAT] assay, and/or infection with Modified Vaccinia virus Ankara expressing T7 RNA polymerase [MVA-T7]) [5, 6, 8, 9], and/or (iii) the systems require transfection of plasmids hampering generation of larger numbers of L gene mutants [6-9].

In this article, we report the establishment of an ambisense minigenome system for RVFV based on

the viral genomic S segment. It has been designed in analogy to the T7 RNA polymerase-driven

minigenome system published for Lassa virus [10]. It provides a measurement range of two to three

log units even after transfection of PCR products for expression of L protein, which is important for

rapid and large-scale mutagenesis of the L gene. Luciferase reporter gene expression allows for

assessment of the overall activity of the RVFV replication complex, while northern blot analysis of

viral RNA facilitates easy discrimination between products of viral transcription and replication. A

mutagenesis study for the L protein was conducted to demonstrate that the established RVFV

system is suitable for studies with the aim of dissecting the molecular mechanisms of replication

and transcription.

Materials and methods

Direct sequencing of RVFV

Overlapping fragments of RVFV-ZH501-BNI S, M, and L RNA, respectively, were reverse

transcribed, amplified, and sequenced directly. In order to generate a S RNA based minigenome

with an authentic promoter of strain ZH501-BNI, the conserved 5' and 3' termini of S RNA were

sequenced as described previously [11]. In brief, purified virus RNA was treated with 5 units of

tobacco acid pyrophosphatase (Epicentre) to generate 5' monophosphorylated termini.

Subsequently 5' and 3' termini were ligated using 10 units of T4 RNA ligase (New England

Biolabs) at 37°C for 1 h. The resulting intramolecular ligation site was reverse transcribed and

amplified, and the PCR product was sequenced. Sequences of primers used can be obtained upon

request. Compared to the sequences of RVFV strain ZH501 deposited in GenBank (accession

numbers DQ375406, DQ380200, DQ380149), the S, M and L RNA sequences for RVFV strain

ZH501-BNI show the following nucleotide (amino acid) differences: S RNA, C616A (NSs protein:

C194Stop); M RNA, A715T (envelope polyprotein: Q232L), T895C (envelope polyprotein:

L292P), G3614A (envelope polyprotein: Stop1198Stop).

Construction of plasmids for RVFV minigenome system

Vero E6 cells in 75-cm^2 tissue culture flasks were inoculated with RVFV-ZH501-BNI. After 4
days, the supernatant was cleared by low-speed centrifugation and virus was pelleted by overnight
ultracentrifugation. The pellet was resuspended in water, and virus RNA was purified by using the
QIAamp viral RNA kit (Qiagen) according to the manufacturer's instructions. Purified RNA was
reverse transcribed and the resulting cDNA amplified using the Superscript III One-Step RT-PCR
System with Platinum Taq (Invitrogen). Amplified RVFV genes for N and L proteins were cloned
into expression vector pCITE-2a containing a T7 RNA polymerase promoter, an internal ribosomal
entry site (IRES), and a favorable Kozak consensus sequence, resulting in pCITE-RVFV-N and
pCITE-RVFV-L, respectively. The final sequences of N and L genes in pCITE 2a vector matched
the consensus sequence of strain ZH501-BNI. The RVFV minigenome plasmid (pRVFV-MG) is
based on the genomic RVFV S RNA integrated into the vector pX12ΔT [12]. The pRVFV-MG
contains the T7 RNA polymerase promoter followed by a single G residue, 5' untranslated region
(5'-UTR) including the conserved 5' terminus, CAT gene, 3' end of the NSs gene (55 nucleotides),
the IGR, 3' end of the N gene (49 nucleotides in reverse orientation), Renilla luciferase (Ren-Luc)
gene in reverse orientation, 3'-UTR including the conserved 3' terminus, hepatitis delta ribozyme
(HDR), and T7 RNA polymerase transcription termination sequence (T7t) (Fig 1). The sequence of
RVFV IGR and flanking nucleotides of the N and NSs genes were synthesized by GeneArt
(Thermo Fisher) for stepwise assembly of the complete minigenome in the pX12ΔT vector,
resulting in pRVFV-MG. Correctness of all sequences was ascertained by sequencing. Sequences of
primers used for cloning are available upon request.

Fig 1. Components of the RVFV ambisense minigenome system. Schematic representation of
minigenome system constructs as used for transfection into BSR-T7/5 cells. The system comprises

the RVFV minigenome plasmid pRVFV-MG based on RVFV S RNA, the plasmid for expression

of the N protein (pCITE-RVFV-N), and the PCR products for expression of the L gene (RVFV-L-

PCR). Plasmid pCITE-FF expressing firefly luciferase serves as a transfection control (not

depicted). Functional elements are abbreviated as follows: T7p, T7 RNA polymerase promoter;

IRES, internal ribosmal entry site; UTR, untranslated region with conserved termini; CAT,

chloramphenicol acetyltransferase gene; IGR, intergenic region, Ren-Luc, Renilla luciferase gene;

HDR, hepatitis delta ribozyme; T7t, T7 RNA polymerase transcription termination sequence.

Minigenome assay

Mutant L genes were generated by mutagenic PCR using pCITE-RVFV-L as a template. The PCR

products containing the functional cassette for expression of mutant L protein (T7 RNA polymerase

promoter, IRES, and L gene) were purified, quantified spectrophotometrically, and used for

transfection without prior cloning as described previously [13]. The presence of the artificial

mutation was ascertained by sequencing. BSR-T7/5 cells stably expressing T7 RNA polymerase

[12] were transfected per well of a 24-well plate with 250 ng of L gene PCR product, 750 ng of

pRVFV-MG expressing Renilla luciferase (Ren-Luc), 500 ng of pCITE-RVFV-N expressing N

protein, and 10 ng of pCITE-FF-Luc expressing firefly luciferase as an internal transfection control.

All transfection were performed with Lipofectamine 2000 (Thermo Fisher) according to the

manufacturer's instructions. At 24 h after transfection, cells were either used to purify total RNA

using an RNeasy Mini Kit (Qiagen) for northern blotting, or lysed in 100 µl of passive lysis buffer

(Promega) per well, and assayed for firefly luciferase and Ren-Luc activity using the dual-luciferase

reporter assay system (Promega). To compensate for differences in transfection efficiency or cell

density, Ren-Luc levels were corrected with the firefly luciferase levels (resulting in standardized

relative light units [sRLU]). RNA (1–2 µg) was separated in a 1.5%-agarose–formaldehyde gel and

transferred onto a Roti-Nylon Plus membrane (Roth) for northern blot analysis. After pre-

hybridization, blots were hybridized with a [32]P-labeled riboprobe targeting the Ren-Luc gene. RNA

bands were visualized by autoradiography using a Typhoon scanner (GE Healthcare).

Expression of L protein

To verify expression of L protein mutants, BSR-T7/5 cells in a well of a 24-well were transfected

with 500 ng of PCR product encoding L protein mutants tagged at the C-terminus with a 3xFLAG

sequence. Cells were additionally inoculated with MVA-T7 [14] prior to the transfection to enhance

L protein expression levels. At 24 h after transfection, cytoplasmic lysate was separated in a 3–8%

Tris-acetate polyacrylamide gel, transferred to a nitrocellulose membrane (Whatman), and detected

by immunoblotting using peroxidase-conjugated anti-FLAG M2 antibody (1:10,000) (A8592;

Sigma-Aldrich). For visualization of the L protein bands by chemiluminescence, the SuperSignal

West Femto substrate (Pierce) and a FUSION SL image acquisition system (Vilber Lourmat) were

used.

Results and discussion

Essential components for an ambisense minireplicon system for RVFV, namely L gene, N gene,

and minigenome, were integrated into appropriate vectors for T7 RNA polymerase-driven

expression in BSR-T7/5 cells (Fig 1). The S RNA segment containing regulatory elements for

ambisense gene expression was used as backbone for the minigenome. The latter contained CAT

and Ren-Luc genes in sense and antisense orientation to measure transcriptional activity of the

system, although only Ren-Luc expression, which is solely dependent on the activity of the RVFV

polymerase, was measured in this study. Expression of firefly luciferase from co-transfected

plasmid served as internal control for transfection efficacy. To demonstrate functionality of the

system, experiments were conducted with wild-type L protein. An L protein mutant containing a

mutation in the catalytic site of the RNA-dependent RNA polymerase (D1133N) served as a

negative control. Wild-type L protein mediated high levels of Ren-Luc expression (up to 1,500,000

light units), while the inactive mutant showed 100–1,000 fold less Ren-Luc expression. RNA

products of genome replication (antigenome) and transcription (Ren-Luc mRNA) generated by

wild-type L protein were clearly visible as distinct signals in northern blot (Fig 2 and S1 Table),

similar to the ambisense Lassa virus minigenome system [10, 15, 16]. Transcription signals for the

D1133N mutant were absent but an unspecific background signal at the antigenome position was

sometimes observed on the northern blot, which was taken into account in the quantification of

antigenome signal intensity. The precise nature of the unspecific material is not known.

To validate the RVFV minigenome system, we tested L protein mutants with exchanges of residues

presumably involved in the endonuclease active site. The endonuclease domain has been located in

the N-terminal ~250 residues of RVFV L protein [9] and is predictably required for viral

transcription, as demonstrated for the corresponding domain of Lassa virus L protein [15, 17].

Putative catalytic residues D111, E125 and K143 were selected based on amino acid alignments of

RVFV with La Crosse virus and hantavirus endonuclease domains as well as prior functional and

structural information on the catalytic site [9, 18, 19]. Mutation of these residues to alanine resulted

in a strong decrease in Ren-Luc activity (Fig 2 and S1 Table). On northern blot, replication products

were detected, whereas Ren-Luc mRNA was absent consistent with the low level of Ren-Luc

activity. In summary, alanine substitution of RVFV L protein residues predictably involved in

endonuclease activity led to a selective defect in viral transcription. The transcription-defective

phenotype could be demonstrated using the ambisense minigenome system.

Fig 2. Determination of transcription and replication activity of L protein mutants using the

RVFV ambisense minigenome system. (A) The activity of L protein mutants in viral replication

and transcription was measured via Ren-Luc reporter gene expression. The Ren-Luc activity is

shown in the bar graph (mean and standard deviation of standardized relative light units [sRLU] as

a percentage of the wild-type (WT) in 3 to 14 independent transfection experiments). Signals for

antigenomic RNA (position AG, representing viral replication) and Ren-Luc mRNA (position

mRNA, representing viral transcription) were detected by northern blotting using a radiolabeled

riboprobe hybridizing to the Ren-Luc gene. A defective L protein with a mutation in the catalytic

site of the RNA-dependent RNA polymerase (D1133N) served as a negative control (neg.).

Northern blots were performed two to three times per mutant and signals on northern blots were

quantified using ImageJ2 software [20]. The quantitative data are presented in S1 Table. The

methylene blue-stained 28S rRNA (28S) served as a marker for gel loading and RNA transfer.

Additionally, immunoblot analysis of 3xFLAG-tagged L protein mutants is shown (L). Mutants

with an mRNA defective phenotype are marked with an asterisk. For experimental details see

methods section. Dotted lines indicate removal of irrelevant lanes for presentation purposes.

Original blots are included in the supporting information (S1 File). **(B)** Further L protein mutants

were tested in the RVFV ambisense minigenome system essentially as described in (A). The bar

graph represents mean and standard deviation of 6 independent measurements for the Ren-Luc

activity. Northern blot analysis was performed twice; the figure depicts one representative

experiment. The quantitative data are presented in S1 Table. Dotted lines indicate removal of

irrelevant lanes for presentation purposes. Original blots are included in the supporting information

(S1 File).

Besides the endonuclease, a cap-binding function is important for viral transcription. It has been

hypothesized that the C terminus of bunyavirus L protein is involved in this function [21, 22].

Furthermore, a role for the C terminus of Lassa virus L protein in viral transcription was proposed

based on a mutagenesis study using the Lassa virus minigenome system [16]. A typical structural

motif of the cap-binding site comprises two aromatic amino acid side chains forming a sandwich

with the guanine moiety of a cap structure. Additionally, the triphosphate moiety of the cap

structure is often interacting with positively charged amino acids. However, a cap-binding site does

not feature a specific sequence motif; therefore it is not possible to predict residues potentially

involved in cap-binding just based on sequence.

We used the RVFV minigenome system to investigate whether the C-terminal region of RVFV L

protein might play a role in viral cap-snatching. Based on an alignment of phlebovirus L protein

sequences, 34 partially or completely conserved aromatic and positively charged amino acids were

selected for an alanine mutation screen (S1 Fig). Ten of these residues were found to be important

for transcription but not replication of the viral genome (Fig 2 and S1 Table). Four residues were

aromatic or heteroaromatic (Y1728, Y1800, H1858 and F2007) and six were positively charged

(K1680, K1682, K1839, R1841, R1942 and R2068). A subset of these residues was additionally

modified to serine and aspartic acid. All these modifications resulted in a selective defect in mRNA

synthesis, which confirmed the involvement of the C-terminal region between positions 1680 and

2068 of the RVFV L protein in viral transcription (Fig 2 and S1 Table).

Conclusions

We developed an ambisense minigenome system for RVFV, which is suitable for screening of L

gene mutations for their impact on viral transcription and replication. As a proof of principle, the

system was used to probe residues potentially involved in the endonuclease active site.

Corresponding mutations resulted in a selective defect in viral transcription, as has been reported

for other viruses of the *Bunyavirales* order [15, 23]. Additionally, the system facilitated identifying

residues in the C-terminal region of RVFV L protein being important for viral transcription but not

replication. However, these data is no proof for the existence of a cap-binding site. They merely

demonstrate a specific role of the identified amino acids during viral transcription. Further

conclusions require biochemical and structural data.

In summary, the established RVFV ambisense minigenome system (i) is suitable to screen L

mutants without cloning, (ii) yields sufficient signal strength without depletion of the cellular

protein kinase R [9], and (iii) allows for technically simple discrimination between viral transcription and replication. Therefore this system is well suited to validate hypotheses arising from structural or biochemical investigation of the RVFV replication complex.

Acknowledgement

We thank Martin Meyer and Beate Becker-Ziaja for excellent technical assistance. We thank Sophia Reindl for fruitful discussions and also acknowledge support by Stephanie Jansen and Stefanie Becker.

References

1.　Ikegami T, Makino S. The pathogenesis of Rift Valley fever. Viruses. 2011;3(5):493-519.

2.　Gaudreault NN, Indran SV, Balaraman V, Wilson WC, Richt JA. Molecular aspects of Rift Valley fever virus and the emergence of reassortants. Virus Genes. 2018.

3.　Mehand MS, Al-Shorbaji F, Millett P, Murgue B. The WHO R&D Blueprint: 2018 review of emerging infectious diseases requiring urgent research and development efforts. Antiviral Res. 2018;159:63-7.

4.　Ikegami T. Molecular biology and genetic diversity of Rift Valley fever virus. Antiviral Res. 2012;95(3):293-310.

5.　Lopez N, Muller R, Prehaud C, Bouloy M. The L protein of Rift Valley fever virus can rescue viral ribonucleoproteins and transcribe synthetic genome-like RNA molecules. J Virol. 1995;69(7):3972-9.

6.　Accardi L, Prehaud C, Di Bonito P, Mochi S, Bouloy M, Giorgi C. Activity of Toscana and Rift Valley fever virus transcription complexes on heterologous templates. J Gen Virol. 2001;82(Pt 4):781-5.

7.   Ikegami T, Peters CJ, Makino S. Rift valley fever virus nonstructural protein NSs promotes

viral RNA replication and transcription in a minigenome system. J Virol. 2005;79(9):5606-15.

8.   Gauliard N, Billecocq A, Flick R, Bouloy M. Rift Valley fever virus noncoding regions of

L, M and S segments regulate RNA synthesis. Virology. 2006;351(1):170-9.

9.   Klemm C, Reguera J, Cusack S, Zielecki F, Kochs G, Weber F. Systems to establish

bunyavirus genome replication in the absence of transcription. J Virol. 2013;87(14):8205-12.

10.   Hass M, Golnitz U, Muller S, Becker-Ziaja B, Gunther S. Replicon system for Lassa virus.

Journal of Virology. 2004;78(24):13793-803.

11.   Meyer BJ, Southern PJ. Concurrent sequence analysis of 5' and 3' RNA termini by

intramolecular circularization reveals 5' nontemplated bases and 3' terminal heterogeneity for

lymphocytic choriomeningitis virus mRNAs. J Virol. 1993;67(5):2621-7.

12.   Buchholz UJ, Finke S, Conzelmann KK. Generation of bovine respiratory syncytial virus

(BRSV) from cDNA: BRSV NS2 is not essential for virus replication in tissue culture, and the

human RSV leader region acts as a functional BRSV genome promoter. J Virol. 1999;73(1):251-9.

13.   Hass M, Lelke M, Busch C, Becker-Ziaja B, Gunther S. Mutational evidence for a structural

model of the Lassa virus RNA polymerase domain and identification of two residues, Gly1394 and

Asp1395, that are critical for transcription but not replication of the genome. Journal of Virology.

2008;82(20):10207-17.

14.   Sutter G, Ohlmann M, Erfle V. Non-replicating vaccinia vector efficiently expresses

bacteriophage T7 RNA polymerase. FEBS Lett. 1995;371(1):9-12.

15.   Lelke M, Brunotte L, Busch C, Gunther S. An N-terminal region of Lassa virus L protein

plays a critical role in transcription but not replication of the virus genome. Journal of Virology.

2010;84(4):1934-44.

16.   Lehmann M, Pahlmann M, Jerome H, Busch C, Lelke M, Gunther S. Role of the C terminus

of Lassa virus L protein in viral mRNA synthesis. J Virol. 2014;88(15):8713-7.

17.   Holm T, Kopicki JD, Busch C, Olschewski S, Rosenthal M, Uetrecht C, et al. Biochemical

and structural studies reveal differences and commonalities among cap-snatching endonucleases

from segmented negative-strand RNA viruses. J Biol Chem. 2018;293(51):19686-98.

18.   Fernandez-Garcia Y, Reguera J, Busch C, Witte G, Sanchez-Ramos O, Betzel C, et al.

Atomic Structure and Biochemical Characterization of an RNA Endonuclease in the N Terminus of

Andes Virus L Protein. PLoS Pathog. 2016;12(6):e1005635.

19.   Reguera J, Gerlach P, Rosenthal M, Gaudon S, Coscia F, Gunther S, et al. Comparative

Structural and Functional Analysis of Bunyavirus and Arenavirus Cap-Snatching Endonucleases.

PLoS Pathog. 2016;12(6):e1005636.

20.   Rueden CT, Schindelin J, Hiner MC, DeZonia BE, Walter AE, Arena ET, et al. ImageJ2:

ImageJ for the next generation of scientific image data. BMC Bioinformatics. 2017;18(1):529.

21.   Reguera J, Weber F, Cusack S. Bunyaviridae RNA polymerases (L-protein) have an N-

terminal, influenza-like endonuclease domain, essential for viral cap-dependent transcription. PLoS

Pathog. 2010;6(9):e1001101.

22.   Ferron F, Weber F, de la Torre JC, Reguera J. Transcription and replication mechanisms of

Bunyaviridae and Arenaviridae L proteins. Virus Res. 2017;234:118-34.

23.   Morin B, Coutard B, Lelke M, Ferron F, Kerber R, Jamal S, et al. The N-terminal domain of

the arenavirus L protein is an RNA endonuclease essential in mRNA transcription. PLoS Pathog.

2010;6(9):e1001038.

24.   Simossis VA, Heringa J. PRALINE: a multiple sequence alignment toolbox that integrates

homology-extended and secondary structure information. Nucleic Acids Research. 2005;33(Web

Server issue):W289-94.

25.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, et al. Fast, scalable generation

of high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol.

2011;7:539.

26. McWilliam H, Li W, Uludag M, Squizzato S, Park YM, Buso N, et al. Analysis Tool Web

Services from the EMBL-EBI. Nucleic Acids Res. 2013;41(Web Server issue):W597-600.

27. Bawono P, Heringa J. PRALINE: a versatile multiple sequence alignment toolkit. Methods

Mol Biol. 2014;1079:245-62.

28. Robert X, Gouet P. Deciphering key features in protein structures with the new ENDscript

server. Nucleic Acids Res. 2014;42(Web Server issue):W320-4.

29. Drozdetskiy A, Cole C, Procter J, Barton GJ. JPred4: a protein secondary structure

prediction server. Nucleic Acids Res. 2015;43(W1):W389-94.

Supporting Information

S1 Table. Functional analysis of L protein mutants in the RVFV ambisense minigenome system.

330 [1] For each mutant, 3 to 14 independent transfection experiments were performed. Ren-Luc values represent mean with standard deviation (n = 3-14). Northern blots were performed at least twice for every mutant. A selective defect in mRNA synthesis was defined as reduction in Ren-Luc level (≤1–35%) despite wild-type like antigenome synthesis (37–280%) and reduction of the mRNA-to-antigenome ratio (≤1–35%). Mutants with a selective defect in mRNA synthesis are shown in boldface on grey background.

[2] Standardized relative light unit (sRLU) value (wild-type L protein = 100%).

[3] sRLU value of mutant divided by sRLU value of the negative control mutant (D1133N) containing a mutation in the catalytic site of the RNA-dependent RNA polymerase.

[4] Antigenome signals in northern blots were quantified via intensity profiles using ImageJ2 software (wild-type L protein = 100%). Background signals at the position of the antigenome of the respective northern blot were either subtracted from all other replication signals or data were evaluated without background subtraction (numbers in parentheses).

[5] RNA signals on northern blots were quantified using ImageJ2 software and the mRNA-to-

antigenome signal ratio was calculated. The wild-type ratio was set at 100% for each experiment

(i.e., the signal ratio of a mutant was normalized with the wild-type ratio) to render independent

experiments comparable. Background signals at the position of the antigenome were subtracted

from all other replication signals. Data without background correction are also shown (numbers in

parentheses).

S1 Figure. Alignment of C terminal sequences of phlebovirus L proteins. The figure presents a

secondary structure-guided alignment of the C-terminal sequences of L proteins from 22

phleboviruses (Uniprot accession numbers are given). The alignment was created by manually

combining results of ClustalOmega and PRALINE programs [24-27] and data are presented by

ESPript (http://espript.ibcp.fr) [28]. The corresponding secondary structure predictions were

calculated by Jpred4 [29] and are depicted below the sequences (β-sheets as arrows, α-helices as

barrels, loops as lines). All positions refer to RVFV strain ZH-501 full-length L protein. Impact of

residue-to-alanine substitution on RVFV L protein activity is indicated by triangles above the

sequences (white, mild or no effect; grey, general defect in L protein activity; red, selective defect

in viral transcription).

S1 File. Original northern and western blots. Lanes of northern blots that have been used for

quantification and statistical analysis (included in S1 Table) are labeled with the respective L

protein mutation, "WT" for L protein wild-type, or "neg." for negative control mutant D1133N.

Lanes of northern blots and methylene blue stained northern blot membranes presented in Fig 2 are

framed by dotted rectangles. Western blot bands used in Fig 2 are also framed by dotted rectangles

and labeled.