Developmental plasticity in metabolism but not in energy reserve accumulation in a

seasonally polyphenic butterfly

Running title: Metabolic pre-diapause polyphenism

Sami M. Kivelä[a, b], Karl Gotthard[c] and Philipp Lehmann[c]

[a] Department of Zoology, Institute of Ecology and Earth Sciences, University of Tartu,

Vanemuise 46, EE-51014 Tartu, Estonia

[b] Present address: Department of Ecology and Genetics, University of Oulu, PO Box 3000,

FI-90014 University of Oulu, Finland

[c] Department of Zoology, Stockholm University, SE-10691 Stockholm, Sweden

Corresponding author: Sami M. Kivelä

e-mail: sami.kivela@oulu.fi

Key words: allometry, diapause, direct development, fat reserve, life history, metabolic rate,

polyphenism

Author contributions: All authors conceived the study and designed the experiment. SMK

and PL carried out the experiment. PL ran the body composition analyses and analysed the

respirometric measurements. SMK ran statistical analyses. All authors participated in writing

the manuscript.

Summary statement: The switch between diapause and direct development results in developmental pre-diapause plasticity in life history and metabolism but not in reserve accumulation in a seasonally polyphenic butterfly.

Abstract

The evolution of seasonal polyphenisms (discrete phenotypes in different annual generations)

associated with alternative developmental pathways of diapause (overwintering) and direct

development is favoured in temperate insects. Seasonal life history polyphenisms are

common and include faster growth and development under direct development than diapause.

However, the physiological underpinnings of this difference remain poorly known despite its

significance for understanding the evolution of polyphenisms. We measured respiration and

metabolic rates through the penultimate and final larval instars in the butterfly *Pieris napi* and

show that directly developing larvae grew and developed faster and had a higher metabolic

rate than larvae entering pupal diapause. The metabolic divergence appeared only in the final

instar, that is, after the induction of developmental pathway that takes place in the

penultimate instar in *P. napi*. The accumulation of fat reserves during the final larval instar

was similar under diapause and direct development, which was unexpected as diapause is

predicted to select for exaggerated reserve accumulation. This suggests that overwinter

survival may be more dependent on metabolic suppression during diapause than the size of

energy reserves. The results, nevertheless, demonstrate that physiological changes coincide

with the divergence of life histories between the alternative developmental pathways, thus

elucidating the proximate basis of seasonal life history polyphenisms.

Introduction

Temperate and high latitudes are characterized by the seasonal cycle between favourable and unfavourable conditions. Organisms living in these environments year-round must be able to tolerate adverse conditions. In short-lived organisms, survival over the adverse season is usually possible only in a specific life stage that is able to enter a state of diapause, dormancy or hibernation that can tolerate adverse conditions and survive long periods on stored energy reserves (Tauber et al., 1986; Danks, 1987; Hahn and Denlinger, 2007; Varpe, 2017). In organisms that can complete several generations during the favourable season, not all generations experience the adverse winter conditions. Hence, a developmental switch (Nijhout, 2003) resulting in alternative developmental pathways – one leading to overwintering and delayed maturation and one leading to direct maturation – evolves. Because individuals completing their life cycles during the favourable season face different selection pressures compared to overwintering individuals, the evolution of alternative developmental pathways should result in phenotypic divergence (polyphenism) between directly developing and overwintering individuals (Moran, 1992; Kivelä et al., 2013, 2016a).

Seasonal polyphenism, the expression of different phenotypes in different generations through the season, in morphology and life history is widespread in arthropods, particularly in insects (e.g. West-Eberhard, 2003; Arbačiauskas, 2004; Simpson et al., 2011). Insect seasonal life history polyphenisms are associated with alternative developmental pathways of diapause (overwintering) and direct development (e.g. Wiklund et al., 1991; Blanckenhorn and Fairbairn, 1995; Aalberg Haugen et al., 2012; Friberg et al., 2012; Välimäki et al., 2013; reviewed by Kivelä et al., 2013). Alternative life histories in the diapause and direct development pathways have a physiological basis, like life history phenotypes in general (see e.g. contributions in Flatt and Heyland, 2011). These physiological differences culminate in the life stage that is capable of entering diapause, because surviving a long adverse period in the dormant state of diapause necessitates a very different physiology compared to direct development into reproductive adult (e.g. Han and Bauce, 1998; Hahn and Denlinger, 2007, 2011; Liu et al., 2007, 2016; Rozsypal et al., 2013; Lehmann et al., 2016, 2018).

Life history divergence between the diapause and direct development pathways may, however, start accumulating already before the attainment of the developmental stage that is capable of diapause. For instance, individuals entering direct development often have shorter

development times and higher growth rates compared to individuals that will enter diapause later in development (Wiklund et al., 1991; Blanckenhorn and Fairbairn, 1995; Aalberg Haugen et al., 2012; Friberg et al., 2012; Välimäki et al., 2013; reviewed by Kivelä et al., 2013). Such pre-diapause trait differentiation is expected because the developmental switch between diapause and direct development is typically triggered by environmental cues of seasonal change already before the attainment of the diapause-capable life stage (Tauber et al., 1986; Friberg et al., 2011). Hence, we would expect divergence in life history and physiology to start accumulating immediately after the induction of the developmental pathway, but the developmental timing of trait divergence between diapause and direct development remains insufficiently understood. Depending on species, life history traits start diverging between the alternative developmental pathways either early (Nylin et al., 1989; see also Lindestad et al., 2019) or late in juvenile development (Friberg et al., 2012; Esperk et al., 2013). Development and growth rates are expected to be associated with other biological rates, like metabolic rate (see e.g. Brown et al., 2004; but see Glazier, 2015 for a potential lack of association between metabolic rate and other biological rates). Nevertheless, it remains unknown whether the relatively short development time and high growth rate under direct development are associated with a high metabolic rate that also diverges after the induction of alternative developmental pathways. Alternatively, directly developing individuals may just follow the same metabolic and growth trajectories as diapausing ones but metamorphose earlier.

Besides metabolic rate, the diapause and direct development pathways are expected to show also other pre-diapause physiological differences. Overwinter survival necessitates energy reserves that can fuel metabolism through the whole dormant period during which feeding is either not possible at all or insufficient to fully fuel metabolism (Tauber et al., 1986; Danks, 1987; Hahn and Denlinger, 2007, 2011; Lehmann et al., 2016, 2018; see Liu et al., 2016 for energy reserves for summer diapause). These energy reserves must be accumulated before the beginning of diapause, which may result in differences between pathways in the allocation of resources to building reserves (Tauber et al., 1986; Danks, 1987; Hahn and Denlinger, 2007, 2011; Liu et al., 2007).

Here, we address potential pre-diapause physiological divergence between the diapause and direct development pathways by using the green-veined white butterfly *Pieris napi* (Linnaeus) as study system. *Pieris napi* produces at least two generations per year in a large

part of its Palearctic distribution (Wiklund et al., 1991; Tolman and Lewington, 2002; Kivelä et al., 2015). It overwinters exclusively as a pupa, but the induction of the developmental pathway takes place during the penultimate larval instar in response to photoperiod and temperature (Friberg et al., 2011). Individuals following the diapause and direct development pathways have different life histories, including a shorter larval development time under direct development than diapause (Wiklund et al., 1991; Friberg et al., 2011; Kivelä et al., 2015). However, we do not know whether the relatively high rates of development and growth under direct development are associated with a high metabolic rate, and whether metabolic rate diverges between the diapause and direct development pathways at the same time as the divergence in development and growth rate appears. This information is vital to fully understand the physiology of alternative developmental pathways. In *P. napi*, diapause pupae contain more storage lipids than pupae that develop directly into adult six days after pupation, lipidomes and metabolomes differing between the developmental pathways in the same phase of pupal development as well (Lehmann et al., 2016, 2018). It remains unknown if these differences appear already in the larval stage during which lipid reserves are accumulated in *P. napi* (Kivelä et al., 2016b) and many other insects (Tauber et al., 1986; Hahn and Denlinger, 2007, 2011). Interestingly, however, metabolic rate and metabolome are similar in diapausing and directly developing *P. napi* pupae right after pupation (Lehmann et al., 2016, 2018). This is surprising given that the induction of the developmental pathway takes place already in the larval stage (Friberg et al., 2011) and some physiological differences appear in the final larval instar as physiology of pupation induction differs between diapause and direct development in this species (Kivelä et al., 2017). Hence, *P. napi* clearly has potential for inter-pathway physiological differentiation already in the larval stage, but whether there are differences in metabolic rate and reserve accumulation already in the larval stage needs to be elucidated.

We measure metabolic rates of growing *P. napi* larvae around the induction of either pupal diapause or direct development in the penultimate larval instar and continue the measurements through the final instar. This design enables us to assess whether metabolic rate is higher under direct development than diapause and whether potential metabolic divergence between diapause and direct development begins directly after the induction of alternative developmental pathways has taken place in the penultimate instar or later. We also measure body lipid content through the final larval instar to compare energy reserve accumulation in the alternative developmental pathways. The fat body is the major energy

reserve in insects (Hahn and Denlinger, 2007, 2011; Arrese and Soulages, 2010), and since

lipid is the main fuel during diapause in *P. napi* (Lehmann et al., 2016), body lipid content

should be a good indicator of energy reserve size. Together, these two traits provide insight

into the metabolic strategy of diapause in *P. napi*, and whether the energetic challenge is met

through (i) strong metabolic suppression, (ii) extensive lipid accumulation, or (iii) a

combination of both.

Materials and methods

Experimental design

Six diapause-generation adult *P. napi* females were captured in Stockholm, Sweden

(59°22'N, 18°4'E) and individually placed in 1 L containers provided with the natural larval

host plant *Alliaria petiolata* as an oviposition substrate in a laboratory. Eggs were frequently

monitored and neonate larvae were individually moved to 0.5 L containers with ad libitum *A.*

petiolata. Each family (offspring of a single female) was split between two climate cabinets

(Termaks Series KB8400L; Termaks, Bergen, Norway): one with diapause-inducing short-

day conditions (12 h day length at 20°C) and the other with long-day conditions that induce

direct development (23 h day length at 20°C). The larvae were monitored daily, and fresh *A.*

petiolata was provided ad libitum. As soon as the larvae started to moult to the penultimate

(IV) instar, they were monitored with at most six hour intervals during the photophase and,

immediately after finishing the moult, taken to the experiment. At this stage, siblings from

both diapause- and direct development-inducing conditions were further split between control

and respirometry groups: the first larva to finish the moult to instar IV within each family was

randomized to one group, thereafter every second sibling to finish the moult being assigned

to alternative groups.

Larvae in the control group were weighed (Precisa XB 120A; Precisa, Dietikon, Switzerland)

twice per day with ca. 8 h interval during photophase in the short-day climate cabinet. This

procedure continued through instars IV and V until the larvae became prepupae. In the

respirometry group, the procedure of measuring the larvae was otherwise similar as in the

control group but weighing of the larvae was always preceded by a measurement in a

respirometer (larvae were weighed immediately after the respirometric measurement; see

below for details of respirometric measurements). Larvae were kept in the climate cabinets

under the assigned diapause- or direct development-inducing conditions all the time, except

during the measurements.

Prepupae were monitored twice a day for pupation and, approximately 48 h after pupation,

the pupae were weighed and sex was determined. Pupae were housed under the same

conditions in the same climate cabinets where they were during the larval period. Pupae were

monitored daily for adult eclosion. At 20°C, directly developing individuals typically eclose

within two weeks after pupation. Therefore, we considered all individuals that did not eclose

and showed no signs of development within three weeks after pupation to be in diapause. All

such individuals were still in diapause after six months.

The experiment was repeated with two temporally separate cohorts so that the start of the

experiment in the second cohort was 14 days later than that in the first cohort. Different

families were included in the two cohorts. The first and the second cohorts included 2 and 4

families, respectively. In the first cohort, 10 control group larvae and 10 respirometry group

larvae were exposed to both diapause- and direct development-inducing conditions (5

larvae/family in each combination of developmental pathway and group). In the second

cohort, the group-specific number of larvae was 12 (2–5 larvae/family in each combination of

developmental pathway and group). Due to mortality in the first cohort (two direct-

development control larvae), we got data from 42 control and 44 respirometry larvae through

the instars IV and V in total.

Respirometry

Differential mode flow-through respirometry was used to measure carbon dioxide (CO_2)

production and oxygen (O_2) consumption of individual larvae. Outdoor air was sucked into

the system by SS-4 pump (Sable Systems, Las Vegas, USA) set to 500 ml min^{-1}, and

scrubbed of H_2O using drierite (WA Hammond Drierite, Xenia, USA). Then, the air inflow

was split into two lines. Both lines were pushed at a steady rate of 150 ml/min using two

separate mass flow controllers (840 Series; Sierra Instruments Inc, California, USA). The

first line went into a MUX multiplexer (Sable Systems) and through the measurement

chamber (volume 10 ml), located in a climatic cabinet (Panasonic MIR-154-PE, Panasonic,

Osaka, Japan) set to 20°C. Thereafter, it again was scrubbed of H_2O with magnesium

perchlorate (Sigma-Aldrich) before entering the sample line of a Li-7000 CO_2 analyser

(LiCor, Lincoln, USA). The second line proceeded in the same way, mimicking the exact

length of the sample line (including an empty measurement chamber), before entering the

reference line of the Li-7000 CO_2 analyser. Both lines then proceeded through ascarite CO_2

scrubbers (Acros Organics, New Jersey, USA), and entered an Oxzilla FC-2 O_2 analyser

(Sable Systems) after which air was ejected to the room. Both CO_2 and O_2 analysers were

calibrated before the start of the experiment. Differential CO_2 and O_2 were calculated by

subtracting the output of the reference line from the output of the sample line. Sampling rate

was 1 Hz for all measurements. The measurement protocol entailed a 2 min baseline

measurement of an empty chamber (preliminary experiments showed that this was long

enough), followed by an 8 min measurement of the chamber containing the animal, and

ending with a 3 min measurement of the empty chamber (see Fig. S1). The raw data was first

lag corrected to synchronize CO_2 and O_2 traces and then drift corrected (there was minimal

drift during the 15 minute measurements). The trace was then baseline corrected against the

empty chamber value, fractioned and multiplied with flow rate to convert the raw output to

ml CO_2 or O_2 h^{-1} (Lighton 2008).

We aimed to minimise the time larvae had to be without access to food, and the consequent

negative effects on growth. Therefore, larvae were not starved before respirometric

measurements, yet they were without food during the measurement (lasting approximately 15

minutes including handling). The number of faecal pellets produced during the measurement

was recorded to take into account potential variation in the digestive cycle and CO_2 released

from the faeces. We did not measure movement of larvae during measurements as a previous

study using a similar setup showed that larvae move little in the tubes and, in that study, the

movement effect was either statistically insignificant or very small (Kivelä et al., 2016b).

We calculated RQ only from measurements where signal-to-noise ratio was >5 as these

measurements were considered as reliable. The signal-to-noise ratio was calculated as the

difference in measurement mean and baseline mean, divided by the baseline standard

deviation. All CO_2 measurements exceeded this threshold but, for O_2, the signal was

generally too weak until the end of the penultimate instar (the smallest larva with a successful

O_2 measurement was 18.4 mg).

Analysis of body lipid and water content

In addition to larvae used in control and respirometric measurements, we reared extra larvae under the above-mentioned diapause and direct development conditions. Once these larvae entered the final (V) instar, a time point (day during the instar V or a two-day old pupa) for freezing was randomized for each individual. The larvae were weighed twice a day until they were frozen in liquid nitrogen at the pre-determined time points. The numbers of frozen final-instar larvae and pupae were 99 and 14 in the diapause pathway, respectively, the corresponding direct development pathway numbers being 94 and 17.

Samples were kept at -80°C until analysis. Water and total lipid content of larvae was measured after chloroform:methanol (Chl:Meth) extraction (Folch et al., 1957) according to the following protocol. First, larvae were weighed (fresh mass; Precisa XB 120A balance), and then dried for 72 h at 55°C. Larvae were then reweighed (dry mass), placed in small glass vials (20 ml) containing 4 ml Chl:Meth (2:1 [vol:vol]), crushed thoroughly with a glass rod, and left for 72 h in a fume hood at 20°C. Chl:Meth effectively dissolves lipids in the sample into the liquid phase. This includes neutral lipids and phospholipids, but also glycerol, carbohydrates and amino acids (Newman et al., 1972), so this method overestimates total lipids (Williams et al., 2011). In *P. napi*, there appears to be no significant accumulation of these compounds right after pupation regardless of pathway (Lehmann et al., 2018) and, therefore, the method should be suitable for between-pathway comparisons even though the absolute lipid values need to be treated with caution. A previous study, using mass-spectrometric methods, verifies that neutral lipids constitute the absolute major component of total lipids in young pupae of *P. napi* (Lehmann et al., 2016). After the Chl:Meth submersion, 250 µl of the suspension was retrieved with a syringe (Hamilton, Model 1825 RN SYR, 22s ga) and applied in pre-weighed (Cahn 28 Automatic Electrobalance), aluminium cups (350 µl VWR Aluminium round micro weigh). Between each sample the syringe was rinsed with Chl:Meth twice. The aluminium cups were left to evaporate in a fume hood for 4 hours and reweighed.

Total lipid mass in the sample was calculated by multiplying the mass of lipids in the cup with the dilution factor (4000/250 = 16). Relative lipid content was calculated as the ratio of

total lipid mass to dry mass. Relative water content was calculated as 1 – ([dry mass] / [fresh

mass]).

Statistical analyses

We used linear mixed-effects models fitted with the maximum likelihood method by using

the function 'lme' (package 'nlme'; Pinheiro et al., 2016) in R version 3.3.2 (R Core Team,

2016) in all data analyses. The multimodel inference approach (e.g. Burnham and Anderson,

2002) was used throughout.

We compared life history traits associated with growth and development (development time

through instars IV and V, growth rate in instars IV and V, time from the beginning of instar

IV until the attainment of peak larval mass in instar V, peak larval mass, and pupal mass)

between the diapause and direct development pathways, and between the control and

respirometry groups. Individual-specific growth rate was calculated as the regression slope of

ln(mass) on time (days) in the focal instar by including all observations in the regression in

instar IV, and the first half of them in instar V to get a growth rate estimate for the free

growth period before growth deceleration preceding pupation (see Fig. S2). In analyses of life

history traits, the "global model" (i.e. the most extensive model considered) included group

(control/respirometry), developmental pathway (diapause/direct), sex (female/male) and all

interactions among them, as well as cohort (first/second; not included in any interactions), as

fixed effects. Family was set as a random effect to control for non-independence of siblings.

The set of all meaningful models (i.e. following the principle of hierarchy) simpler than the

global model was derived with the function 'dredge' (package 'MuMIn'; Barton, 2016; see

Oline Resource, Table S1, for the sets of models with Akaike weights >0.05). The models,

including the global model, were averaged with the function 'model.avg' (Barton, 2016).

When needed, we took heteroscedasticity into account by adding weights determined by a

variance function (either 'varIdent' or 'varExp' [Pinheiro et al., 2016]) into the models. The

sets of models with Akaike weight >0.05 are presented in Table S1.

For CO_2 production rate, we analysed instars IV and V separately and included only

observations during the growth period in the analysis. Data were ln-transformed to facilitate

allometric inferences (see Kilmer and Rodríguez, 2016 for arguments favouring this approach

over reduced major axis regression). Fixed effects in the global model included centered ln-

transformed mass, developmental pathway, sex and all interactions among them, as well as

the square of centered ln-transformed mass included in interactions with developmental

pathway and sex. Cohort (not included in any interactions) and the number of frass produced

during the respirometric measurement (a covariate) were also included in the fixed effects. To

control for repeated within-individual measures, random effects included random individual-

specific intercepts and slopes in relation to centered ln-transformed mass. The inclusion of

random slopes significantly improved model goodness-of-fit over random-intercepts-only

model both in instar IV and V data when measured by Akaike information criterion

(ΔAIC>2). The inclusion of random individual-specific slopes is also important to avoid

overconfident estimates of fixed effects (Schielzeth and Forstmeier, 2009). To reduce

correlations among random intercepts and slopes, ln-transformed mass was centered

(subtracting the mean from each observation). Centering a variable only changes the mean

but not the variance. Therefore, centering of ln(mass) does not affect the regression slopes of

other traits on it, and allometric inferences remain, consequently, unaffected by centering.

Model averaging and modelling of heteroscedasticity was performed as explained above, and

the sets of models with Akaike weight >0.05 are presented in Table S2.

Variation in RQ was analysed similarly as explained above for CO_2, except that the models

included only random individual-specific intercepts because the inclusion of random

individual-specific slopes did not improve model goodness-of-fit (AIC values were lower for

the random intercept models). Because there was no need to include random individual-

specific slopes in the models for RQ, mass was not centered to make interpretation of the

results easier. In instar IV, we added weights by the 'varIdent' variance function (Pinheiro et

al., 2016) to the models, because modelling the higher residual variance in females than in

males improved the estimation accuracy of random effects, yet AIC did not suggest a

difference between the weighted and unweighted models (ΔAIC= 0.919). The sets of models

with Akaike weight >0.05 are presented in Table S3.

In the analyses of body composition data (body water and lipid content) from instar V larvae,

fixed effects of the global model included developmental pathway, ln-transformed fresh

mass, the square of ln-transformed fresh mass and interactions between developmental

pathway and each of ln-transformed fresh mass and its square. Family was set as a random

effect. Weights by the 'varExp' variance function (Pinheiro et al., 2016) significantly

improved model goodness-of-fit (ΔAIC=43.0 for water content and ΔAIC=11.5 for lipid

content). We also compared relative water and lipid contents between similarly-sized instar V

larvae and pupae. Here, only larvae whose mass was between the minimum and maximum of

pupal masses in these data were included in the analysis. The global model included

developmental pathway, developmental stage (instar V larva/pupa) and their interaction as

fixed effects, and family as a random effect. Model averaging and modelling of

heteroscedasticity was performed as explained above, and the sets of models with Akaike

weight >0.05 are presented in Tables S4 and S5.

Results

Life history divergence between alternative developmental pathways

In both the control and respirometry group, all individuals exposed to the short day length

manipulation entered diapause, while the long day manipulation invariably induced direct

development. Individuals developing directly into adults had higher larval growth rates than

individuals entering pupal diapause both in instars IV and V (Table 1; Fig 1). Development

time through instars IV and V was shorter under direct development than diapause, and

directly developing individuals reached the peak larval mass in instar V sooner after entering

instar IV than the ones entering diapause (Table 1; Fig. 2). Instar duration was shorter under

direct development than diapause both in instar IV (direct development mean=2.11 [95% CI

2.03, 2.19] days; diapause mean=2.62 [2.52, 2.72] days) and V (direct development mean=

3.38 [3.19, 3.57] days; diapause mean= 4.64 [4.49, 4.79] days), the relative difference in

instar durations becoming more pronounced in instar V compared to instar IV (instar duration

under direct development is 73% and 81% of that under diapause development in instars V

and IV, respectively). Despite having higher growth rate, the fast development of directly

maturing individuals led to a low peak larval mass compared to that attained by individuals

entering diapause (Table 1; Fig. 2). Males had higher pupal masses than females, and there

was also a weak tendency for the pupal mass to be higher under diapause than direct

development (Table 1; Fig. 2).

Metabolic divergence between alternative developmental pathways

We use whole-organism CO_2 production rate as a proxy for metabolic rate because there is a

strong and positive correlation between these two traits (estimated correlation coefficient

based on resampling to avoid pseudoreplication = 0.961 [95% percentile CI 0.934, 0.980];

Fig. S3). Furthermore, O_2 consumption rate, that is needed in the calculation of metabolic

rate, could not be reliably measured early in instar IV with the set-up used here. Therefore, to

facilitate inter-pathway comparisons through the two investigated instars, we base our

inferences on CO_2 production rate.

CO_2 production rate increased non-linearly through instar IV; a hyperallometric increase

early in the instar changed to a hypoallometric increase later in the instar, and the ontogenetic

increase in CO_2 production ceased altogether at the end of the instar (Table 2; Fig. 3a, c; see

also Fig. S4). There was no difference between diapause and direct development in CO_2

production in instar IV (Table 2). In instar V, CO_2 production had a qualitatively similar

relationship to body mass as in instar IV, yet the non-linearity of the relationship was weaker,

and there was a difference between the alternative developmental pathways, CO_2 production

being higher under direct development than diapause (Table 2; Fig. 3b, d).

The respiratory quotient (RQ) in instar IV was independent of developmental pathway and

body mass, the latter being apparently due to the data concentrating at the latter part of the

instar (Table 2; Fig. 4a). The larvae became large enough for reliable measurement of O_2

consumption (and RQ) only in the latter part of the instar IV, which is why we do not have

RQ data from the beginning of the instar IV. In instar V, RQ showed a peaked relationship to

body mass, RQ values being higher in the second than in the first cohort (Table 2; Fig. 4b).

There was no difference in RQ values between diapause and direct development.

Body water and lipid contents in alternative developmental pathways

The absolute amount of water in the body did not differ between diapause and direct

development and increased nearly isometrically, yet non-linearly, over the instar V, the

scaling becoming increasingly hypoallometric at the end of the instar (Table 3; Fig. 5a).

Absolute mass of lipids in the body did not differ between diapause and direct development

either and increased linearly and hyperallometrically through instar V (allometric

exponent=1.37; Table 3; Fig. 5b).

When comparing two days old pupae with similarly-sized instar V larvae, relative water

content was lower in diapause pupae than in larvae or directly developing pupae (Table 3;

Fig. 5c). A similar comparison in the proportion of lipids in dry body matter revealed that the

proportion of lipids was higher in instar V larvae than in pupae (Table 3; Fig. 5d), but there

was no statistical support for a difference between the developmental pathways (Table 3).

Discussion

We found the expected life history divergence between the diapause and direct development

pathways through the last two larval instars in *P. napi*; larval growth and development were

faster and body size smaller in individuals destined for direct development rather than pupal

diapause. This life history differentiation between the alternative developmental pathways

was associated with physiological differentiation, metabolic rate (defined as CO_2 production)

being higher under direct development than diapause. As expected, metabolic rate of larvae

entering direct development and pupal diapause differed only in the final larval instar, and

not in the penultimate instar. This is because the penultimate instar is the typical time window

during which the developmental pathway is induced in *P. napi* (Friberg et al., 2011), resulting

in a limited post-induction period for phenotypic divergence to become apparent in the

penultimate instar. Putting these results together with the observation that the inter-pathway

difference in development time became more pronounced in the final larval instar and the fact

that the induction of a developmental pathway is largely independent of rates of growth and

development in *P. napi* (Friberg et al., 2012; Kivelä et al., 2015), a high metabolic rate seems

associated with fast growth and rapid development in individuals set for direct maturation

after the induction of the developmental pathway. A high metabolic rate may be the cause for

a high growth rate, or vice versa (Glazier, 2015). Either way, the relatively low metabolic rate

in the diapause pathway and the lack of exaggerated lipid accumulation suggest that

metabolic suppression is an important component of the diapause phenotype.

Ontogenetic allometric scaling of metabolism to body size was complex. The scaling of

metabolic rate to body size was non-linear, early-instar hyperallometry turning into

hypoallometry at the end of an instar (see also Glazier, 2005). This non-linear pattern was

consistent both in the penultimate and final instars, yet the non-linearity was exaggerated in

the penultimate instar. In an earlier study concentrating only on the diapause pathway, we

found a similar non-linear scaling of metabolic rate to body mass in the penultimate instar in

the diapause pathway and a linear scaling in the final instar (Kivelä et al., 2016b). A larger

sample size in the present study is a likely explanation why we could detect a clearer

levelling-off of metabolic rate in the penultimate instar (Figs 3a, c and S3) and a clear but

weak non-linearity also in the final instar (Fig. 3b, d). Non-linearity of scaling may arise due

to an increasing relative amount of metabolically inactive matter in the body towards the end

of instars (see Glazier, 2005). Accordingly, the proportion of lipids in body matter largely

follow this pattern. Moreover, the present data from the penultimate instar (see also Fig. S4)

are consistent with the oxygen-dependent induction of moulting hypothesis, predicting that

the tracheal respiratory system, that primarily grows during moults, reaches its maximum

capacity at the end of the instar and so sets an upper limit for respiration rate (Greenlee and

Harrison, 2004, 2005; Callier and Nijhout, 2011; Kivelä et al., 2016c).

Even though metabolic rate differed between the developmental pathways, there was no

difference in the accumulation of lipids between diapause and direct development, although

lipids constitute the main energy reserves used during diapause in many insects (Hahn and

Denlinger, 2007, 2011; Arrese and Soulages, 2010), including *P. napi* (Lehmann et al.,

2016). This is not exceptional (Hahn and Denlinger, 2007), yet unexpected given the ecology

of *P. napi* where diapausing pupae must survive 5 to 10 months on the energy reserves

acquired during the larval stage. In addition, many diapause-generation individuals pupate

early in the summer and therefore also experience a long pre-winter warm period in pupal

diapause. This may be particularly stressful as higher temperatures should lead to increased

fat reserve depletion in ectotherms (Han and Bauce, 1998; Irwin and Lee, 2003; Bosch et al.,

2010; Williams et al., 2014; see also Liu et al., 2016). Although there was no inter-pathway

difference in the accumulation of energy reserves, final-instar *P. napi* larvae accumulate lots

of lipids in their bodies during this instar, as indicated by the hyperallometric increase in total

lipid mass in the body (Fig. 5b). Respiratory quotient (RQ) values exceeding one suggest that

either metabolism is partially anaerobic (see e.g. Nielsen & Christian, 2007) or larvae convert

carbohydrates to lipids (Zebe, 1953; Wigglesworth, 1972). Conversion of carbohydrates to

lipids seems more likely as the proportion of lipids in dry body matter peaks similarly in

halfway through the instar V as RQ (compare Fig. 4b to Fig. S5; see also Kivelä et al.,

2016b). In the pupal stage, diapause pupae had lower water (and possibly higher lipid; see Fig. 4d) contents than directly developing ones soon after pupation and this difference emerged during pupation as larval water (and lipid) content did not differ between the developmental pathways. Reduced water content of diapause pupae may be a strategy to improve cold tolerance (see Danks, 2000; Bennett et al., 2005; Rozsypal et al., 2013).

Seasonal polyphenisms in life history and morphology associated with the diapause and direct development pathways have been demonstrated repeatedly in insects (Wiklund et al., 1991; Blanckenhorn and Fairbairn, 1995; West-Eberhard, 2003; Simpson et al., 2011; Aalberg Haugen et al., 2012; Friberg et al., 2012; Välimäki et al., 2013; Kivelä et al., 2013). Here, we show that the phenotypic divergence between the developmental pathways extends to metabolism, and the metabolic divergence appears after the induction of a developmental pathway has taken place. Pre-diapause metabolic divergence between larvae entering diapause and direct development that we demonstrate here persists after pupation (Lehmann et al., 2016). This is necessary because diapause pupae must survive up to 10 months with reserves similar to those of directly developing pupae that instead eclose as adults within a few weeks.

The generally extensive phenotypic divergence between alternative developmental pathways suggest that many components of the physiological and developmental machinery are independent enough for pathway-specific evolution. It is possible that pathway-specific selection changes metabolic rate, which would imply that metabolic rate is a key component of physiology affecting the evolution of life history polyphenisms (see also Arbačiauskas, 2004; Furness et al., 2015; but see Saastamoinen et al., 2013), and possibly life history evolution across species (Brown et al., 2004; Maino et al., 2014). Yet, it remains also possible that metabolic rate responds to changes in growth rate (Glazier, 2015), and that growth rate is the key to phenotypic differentiation between the alternative developmental pathways. Further studies are necessary for a rigorous assessment of these alternatives.

Acknowledgements

We thank Loke von Schmalensee for help in running the experiment.

Competing interests

No competing interests declared

Funding

583 This study was financed by the international fellowship program at Stockholm University
(SMK), the Estonian Research Council (grant PUT1474, SMK), Academy of Finland (grants
No. 314833 and 319898, SMK), the Knut and Alice Wallenberg foundation (grant
2012.0058, KG) and the Bolin Centre for Climate Research at Stockholm University (KG).

References

Aalberg Haugen, I. M., Berger, D. and Gotthard, K. (2012). The evolution of alternative
developmental pathways: footprints of selection on lifehistory traits in a butterfly. *J. Evol. Biol.* **25,** 1377–1388.

Arrese, E. L. and Soulages, J. L. (2010). Insect fat body: energy, metabolism, and regulation. *Annu. Rev. Entomol.* **55,** 207–225.

Arbačiauskas, K. (2004). Seasonal phenotypes of *Daphnia*: post-diapause and directly developing offspring. *J. Limnol.* **63**, 7–15.

Barton, K. (2016). MuMIn: Multi-Model Inference. R package version 1.15.6. https://CRAN.R-project.org/package=MuMIn

Bennett, V. A., Sformo, T., Walters, K., Toien, O., Jeannet, K., Hochstrasser, R., Pan, Q., Serianni, A. S., Barnes, B. M. and Duman, J. G. (2005). Comparative overwintering physiology of Alaska and Indiana populations of the beetle *Cucujus clavipes* (Fabricius): roles of antifreeze proteins, polyols, dehydration and diapause. *J. Exp. Biol.* **208**, 4467–4477.

Blanckenhorn, W. U. and Fairbairn, D. J. (1995). Life history adaptation along a

latitudinal cline in the water strider *Aquarius remigis* (Heteroptera: Gerridae). *J. Evol.*

Biol. **8**, 21–41.

Bosch, J., Sgolastra, F. and Kemp, W. P. (2010). Timing of eclosion affects diapause

development, fat body consumption and longevity in *Osmia lignaria*, a univoltine,

adult-wintering solitary bee. *J. Insect. Physiol.* **56**, 1949–1957.

Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M. and West, G. B. (2004). Toward

a metabolic theory of ecology. *Ecology* **85**, 1771–1789.

Burnham, K. P., Anderson, D. R. (2002). *Model Selection and Multimodel Inference, A*

Practical Information-Theoretic Approach, 2nd edn. New York, US: Springer.

Callier, V. and Nijhout, H. F. (2011). Control of body size by oxygen supply reveals size-

dependent and size-independent mechanisms of molting and metamorphosis. *Proc. Nat.*

Acad. Sci. USA **108**, 14664–14669.

Danks, H. V. (1987). *Insect dormancy: an ecological perspective*. Ottawa, Canada:

Biological Survey of Canada.

Danks, H. V. (2000). Dehydration in dormant insects. *J. Insect Physiol.* **46**, 837–852.

Esperk, T., Stefanescu, C., Teder, T., Wiklund, C., Kaasik, A. and Tammaru, T. (2013).

Distinguishing between anticipatory and responsive plasticity in a seasonally

polyphonic butterfly. *Evol. Ecol.* **27**, 315–332.

Flatt, T. and Heyland, A. (eds.) (2011). *Mechanisms of life history evolution: the genetics*

and physiology of life history traits and trade-offs. Oxford, UK: Oxford University

Press.

Folch, J., Lees, M. and Sloane Stanley, G. H. (1957). A simple method for the isolation and

purification of total lipids from animal tissue. *J. Biol. Chem.* **226**, 497–509.

Friberg, M,, Aalberg Haugen, I. M., Dahlerus, J., Gotthard, K. and Wiklund, C. (2011).

Asymmetric diapause decision-making in butterfly larvae. *Oecologia* **165**, 301–310.

Friberg, M., Dahlerus, J. and Wiklund, C. (2012). Strategic larval decision-making in a

bivoltine butterfly. *Oecologia* **169**, 623–635.

Furness, A. I., Reznick, D. N., Springer, M. S. and Meredith, R. W. (2015). Convergent

evolution of alternative developmental trajectories associated with diapause in African

and South American killifish. *Proc. R. Soc. B* **282**, 20142189.

Glazier, D. S. (2005). Beyond the '3/4-power law': variation in the intra- and interspecific

scaling of metabolic rate in animals. *Biol. Rev.* **80**, 611–662.

Glazier, D. S. (2015). Is metabolic rate a universal 'pacemaker' for biological processes?
Biol. Rev. **90**, 377-407.

Greenlee, K. J. and Harrison, J. F. (2004). Development of respiratory function in the
American locust *Schistocerca americana* II. Within-instar effects. *J. Exp. Biol.* **207**,
509–517.

Greenlee, K. J. and Harrison, J. F. (2005). Respiratory changes throughout ontogeny in the
tobacco hornworm caterpillar, *Manduca sexta. J. Exp. Biol.* **208**, 1385–1392.

Hahn, D. A. and Denlinger, D. L. (2007). Meeting the energetic demands of insect
diapause: nutrient storage and utilization. *J. Insect Physiol.* **53**, 760–773.

Hahn, D. A. and Denlinger, D. L. (2011). Energetics of insect diapause. *Annu. Rev.*
Entomol. **56**, 103–121.

Han, E.-N. and Bauce, E. (1998). Timing of diapause initiation, metabolic changes and
overwintering survival of the spruce budworm, *Choristoneura fumiferana. Ecol.*
Entomol. **23**, 160-167.

Irwin, J. T. and Lee, R. E. Jr. (2003). Cold winter microenvironments conserve energy and
improve overwintering survival and potential fecundity of the goldenrod gall fly,
Eurosta solidaginis. Oikos **100**, 71–78.

Kilmer, J. T. and Rodríguez, R. L. (2016). Ordinary least squares regression is indicated
for studies of allometry. *J. Evol. Biol.* **30**, 4–12.

Kivelä, S. M., Välimäki, P. and Gotthard, K. (2013). Seasonality maintains alternative life-
history phenotypes. *Evolution* **67**, 3145–3160.

Kivelä, S. M., Välimäki, P. and Gotthard, K. (2016a). Evolution of alternative insect life
histories in stochastic seasonal environments. *Ecol. Evol.* **6**, 5596–5613.

Kivelä, S. M., Lehmann, P. and Gotthard, K. (2016b). Do respiratory limitations affect
metabolism of insect larvae before moulting? An empirical test at the individual level.
J. Exp. Biol. **219**, 3061–3071.

Kivelä, S. M., Friberg, M., Wiklund, C., Leimar, O. and Gotthard, K. (2016c). Towards
a mechanistic understanding of insect life history evolution: oxygen-dependent
induction of moulting explains moulting sizes. *Biol. J. Linn. Soc.* **117**, 586–600.

Kivelä, S. M., Svensson, B., Tiwe, A. and Gotthard, K. (2015). Thermal plasticity of
growth and development varies adaptively among alternative developmental pathways.
Evolution **69**, 2399–2413.

Kivelä, S. M., Friberg, M., Wiklund, C. and Gotthard, K. (2017). Adaptive

developmental plasticity in a butterfly: mechanisms for size and time at pupation differ

between diapause and direct development. *Biol. J. Linn. Soc.* **122**, 46–57.

Lehmann, P., Pruisscher, P., Posledovich, D., Carlsson, M., Käkelä, R., Tang, P., Nylin,

S., Wheat, C. W., Wiklund, C. and Gotthard, K. (2016). Energy and lipid

metabolism during direct and diapause development in a pierid butterfly. *J. Exp. Biol.*

219, 3049–3060.

Lehmann, P., Pruisscher, P., Koštal, V., Moos, M., Šimek, P., Nylin, S., Agren, R.,

Väremo, L., Wiklund, C., Wheat, C. W. and Gotthard, K. (2018). Metabolome

dynamics in the butterfly *Pieris napi*: distinguishing maintenance, termination and post-

diapause phases. *J. Exp. Biol.* **221**, doi:10.1242/jeb.169508

Lighton, J. R. B. (2008). *Measuring metabolic rates: A manual for scientists*. New York,

US: Oxford University Press.

Lindestad, O., Wheat, C. W., Nylin, S. and Gotthard, K. (2019). Local adaptation of

photoperiodic plasticity maintains life cycle variation within latitudes in a butterfly.

Ecology **100**, e02550.

Liu, Z., Gong, P., Wu, K., Wei, W., Sun, J. and Li, D. (2007). Effects of larval host plants

on over-wintering preparedness and survival of the cotton bollworm, *Helcoverpa*

armigera (Hübner) (Lepidoptera: Noctuidae). *J. Insect. Physiol.* **53**, 1016-1026.

Liu, Z., Xin, Y., Zhang, Y., Fan, J. and Sun, J. (2016). Summer diapause induced by high

temperatures in the oriental tobacco budworm: ecological adaptation to hot summers.

Sci. Rep. **6**, 27443.

Maino, J. L., Kearney, M. R., Nisbet, R. M. and Kooijman, S. A. L. M. (2014).

Reconciling theories for metabolic scaling. *J. Anim. Ecol.* **83**, 20–29.

Moran, N. A. (1992). The evolutionary maintenance of alternative phenotypes. *Am. Nat.* **139**,

971–989.

Newman, H. A. I., Gordon, E. A., Heggen, D. W., and Keller, M. D. (1972). Rapid

extraction of triglycerides from human adipose tissue with petroleum ether. *Clin. Chem.*

18, 290–292.

Nielsen,, M. G. and Christian, K. A. (2007). The mangrove ant, *Camponotus anderseni*,

switches to anaerobic respiration in response to elevated CO2 levels. *J. Insect Physiol.*

53, 505-508.

Nijhout, H. F. (2003). Development and evolution of adaptive polyphenisms. *Evol. Dev.* **5**,

9–18.

Nylin, S., Wickman P.-O. and Wiklund C. (1989). Seasonal plasticity in growth and
development of the speckled wood butterfly, *Pararge aegeria* (Satyrinae). *Biol. J. Linn.*
Soc. **38**, 155-171.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D. and R Core Team (2016). nlme: Linear and
Nonlinear Mixed Effects Models. R package version 3.1-128, URL: http://CRAN.R-
project.org/package=nlme

R Core Team (2016). *R: A language and environment for statistical computing*. R
Foundation for Statistical Computing, Vienna, Austria, URL https://www.R-
project.org/

Rozsypal, J., Koštál, V., Zahradníčková, H. and Šimek, P. (2013). Overwintering strategy
and mechanisms of cold tolerance in the codling moth (*Cydia pomonella*). *PLoS ONE*
8, e61745.

Schielzeth, H. and Forstmeier, W. (2009). Conclusions beyond support: overconfident
estimates in mixed models. *Behav. Ecol.* **20**, 416–420.

Saastamoinen, M., Brommer, J. E., Brakefield, P. M. and Zwaan, B. J. (2013).
Quantitative genetic analysis of responses to larval food limitation in a polyphenic
butterfly indicates environment- and trait-specific effects. *Ecol. Evol.* **3**, 3576–3589.

Simpson, S. J., Sword, G. A. and Lo, N. (2011). Polyphenism in insects. *Curr. Biol.* **21**,
R738–R749.

Tauber, M. J., Tauber, C. A. and Masaki, S. (1986). *Seasonal adaptations of insects*. New
York, US: Oxford Univ. Press.

Tolman, T. and Lewington, R. (2002). *Butterflies of Europe*. Princeton: Princeton
University Press.

Varpe, Ø. (2017). Life history adaptations to seasonality. *Integr. Comp. Biol.* **57**, 943–960.

Välimäki, P., Kivelä, S. M., Mäenpää, M. I. and Tammaru, T. (2013). Latitudinal clines
in alternative life histories in a geometrid moth. *J. Evol. Biol.* **26**, 118–129.

West-Eberhard, M. J. (2003). *Developmental plasticity and evolution*. New York, US:
Oxford Univ. Press.

Wigglesworth, W. B. (1972). *The Principles of Insect Physiology*. London: Chapman and
Hall.

Wiklund, C., Nylin, S. and Forsberg, J. (1991). Sex-related variation in growth rate as a
result of selection for large size and protandry in a bivoltine butterfly, *Pieris napi*.
Oikos **60**, 241–250.

Williams, C. M., Henry, H. A. and Sinclair, B. J. (2014). Cold truths: how winter drives

responses of terrestrial organisms to climate change. *Biol. Rev.* **90**, 214–235.

Williams, C. M., Thomas, R. H., MacMillan, H. A., Marshall, K. E., and Sinclair, B. J.

(2011). Triacylglyceride measurement in small quantities of homogenised insect tissue:

comparisons and caveats. *J. Insect Physiol.* **57**, 1602–1613.

Zebe, E. (1953). Über den Respiratorischen Quotienten der Lepidoptera.

Naturwissenschaften **40**, 298.

770 **Table 1.** Statistically significant (risk level 0.1) model-averaged (full average) fixed effects

771 of linear mixed-effects models fitted with the maximum likelihood method explaining life-

772 history variation between alternative developmental pathways (diapause / direct

773 development), experimental groups (control / respirometry) and cohorts (first / second). The

774 models included random family-specific intercepts.

Trait	Term (relative importance)	Factor levels	Averaged estimate	Adjusted S.E.	z-value	P-value
Development time		Intercept	9.28	0.192	48.3	<0.0001
	Pathway (1)	direct	-1.80	0.149	12.1	<0.0001
Time until peak mass		Intercept	6.72	0.146	46.0	<0.0001
	Pathway (1)	direct	-1.11	0.150	7.38	<0.0001
Peak mass		Intercept	177	5.35	33.1	<0.0001
	Cohort (0.92)	second	15.0	6.99	2.14	0.032
	Group (0.99)	respirometry	-10.1	4.04	2.50	0.012
	Pathway (1)	direct	-12.1	4.26	2.84	0.0044
Pupal mass		Intercept	146	4.16	35.0	<0.0001
	Pathway (0.94)	direct	-6.22	3.51	1.78	0.075
	Sex (0.99)	male	9.04	3.37	2.68	0.0073
Growth rate IV[a]		Intercept	0.728	0.0274	26.6	<0.0001
	Pathway (0.98)	direct	0.0907	0.0313	2.90	0.0037
Growth rate V[b]		Intercept	0.589	0.0233	25.3	<0.0001
	Pathway (1)	direct	0.0855	0.0287	2.98	0.0029

775 [a] 'varIdent' variance function to take different variances in different families into account.

776 [b] 'varIdent' variance function to take different variances in females and males into account.

Table 2. Statistically significant (risk level 0.1; insignificant lower-level terms are presented for interactions and squared effects; at least an intercept is presented for each trait) model-averaged (full average) fixed effects of linear mixed-effects models fitted with the maximum likelihood method explaining variation in CO_2 production rate and respiratory quotient (RQ) between alternative developmental pathways (diapause / direct development) and experimental cohorts (first / second). CO_2 production rate was ln-transformed in both instars.

Trait	Term (relative importance)	Factor levels	Averaged estimate	Adjusted S.E.	z-value	P-value
CO_2 IV[a]		Intercept	-3.51	0.0316	111	<0.0001
	Centered ln(mass) (1)		0.950	0.0403	23.5	<0.0001
	[Centered ln(mass)]^2 (1)		-0.548	0.0624	8.78	<0.0001
	Frass (0.99)		0.0814	0.0261	3.12	0.0018
CO_2 V[b]		Intercept	-2.19	0.0324	67.6	<0.0001
	Centered ln(mass) (1)		0.788	0.0339	23.2	<0.0001
	[Centered ln(mass)]^2 (1)		-0.302	0.0479	6.30	<0.0001
	Frass (0.99)		0.0568	0.0170	3.34	0.00082
	Pathway (1)	direct	0.128	0.0343	3.72	0.00020
RQ IV[c]		Intercept	0.500	0.455	1.10	0.27
RQ V[d]		Intercept	0.689	0.0582	11.8	<0.0001
	Mass (0.98)		0.00295	0.00106	2.79	0.0053
	Mass^2 (0.98)		-1.19×10^{-5}	4.42×10^{-6}	2.69	0.0072
	Cohort (1)	second	0.186	0.0155	11.0	<0.0001

[a] Random individual-specific intercepts and slopes in relation to centered mass.

[b] Random individual-specific intercepts and slopes in relation to centered mass; 'varIdent' variance function to take into account higher residual variance in females than males.

[c] Random individual-specific intercepts; 'varIdent' variance function to take into account higher residual variance in females than males.

[d] Random individual-specific intercepts.

Table 3. Statistically significant (risk level 0.1; insignificant lower-level terms are presented

for interactions and squared effects) model-averaged (full average) fixed effects of linear

mixed-effects models fitted with the maximum likelihood method explaining variation in

body water and lipid content between alternative developmental pathways (diapause / direct

development), and developmental stages (instar V larva / pupa). The models included random

family-specific intercepts.

Trait	Term (relative importance)	Factor levels	Averaged estimate	Adjusted S.E.	z-value	P-value
Water content V[a]		Intercept	-1.47	0.245	6.01	<0.0001
	ln(fresh mass) (1)		1.67	0.113	14.7	<0.0001
	[ln(fresh mass)]^2 (1)		-0.0849	0.0130	6.55	<0.0001
Lipid content V[b]		Intercept	-4.94	1.64	3.02	0.0026
	ln(fresh mass) (0.97)		1.37	0.720	1.91	0.056
Water content: larva vs. pupa[c]		Intercept	0.784	0.00604	130	<0.0001
	Pathway (1)	direct	0.000393	0.00726	0.054	0.96
	Stage (1)	pupa	-0.0746	0.00642	11.6	<0.0001
	pathway × stage (1)	direct × pupa	0.0625	0.00805	7.76	<0.0001
Lipid content: larva vs. pupa[d]		Intercept	0.236	0.00646	36.7	<0.0001
	Stage (1)	pupa	-0.0289	0.00888	3.25	0.0012

[a] 'varExp' variance function to take into account increasing residual variance with increasing

fitted value.

[b] 'varExp' variance function to take into account decreasing residual variance with increasing

fitted value.

^c 'varIdent' variance function to take into account different residual variances in different

combinations of developmental pathway and developmental stage (larva / pupa).

^d 'varIdent' variance function to take into account lower residual variance in pupae than in

larvae.

 Figure legends

 Figure 1. Larval growth rates under diapause and direct development in instars IV

 (left) and V (right). Each box includes 50% of observations, the horizontal line within the

 box depicting the median value. The whiskers depict minimum and maximum of the

 distribution while the circles depict outliers.

 Figure 2. Growth trajectories of experimental larvae (both control and respirometry

 group larvae are included) through instars IV (dotted lines) and V (continuous lines).

 Black and red lines depict growth trajectories of individuals entering pupal diapause and

 direct development into adult, respectively. Open circles depict pupal masses two days after

 pupation. Closed circles below the growth trajectories indicate mean times of the attainment

 of peak larval mass and pupation in the diapause (black points) and direct development (red

 points), while horizontal whiskers illustrate their 95% confidence intervals (CI). Closed black

 and red circles and associated vertical whiskers at top right corner indicate mean peak larval

 masses (in ln-scale) and their 95% CIs under diapause and direct development, respectively.

 Figure 3. Individual-specific trajectories of changing whole-organism CO_2 production

 rate (lines) along with growth (increasing mass) under diapause (black lines) and direct

 development (red lines) in instars IV (a) and V (b). The bottom row illustrates allometric

 regression curves fitted to ln-transformed data (points) in instars IV (c) and V (d). The

 regression curves are based on model-averaged coefficients (see Table 2) fitted to data

 including centered ln(mass) and illustrate within-instar allometry of CO_2 production rate.

 There was no difference between diapause and direct development in instar V, so only a

 single regression curve is presented in (c). The dotted lines in the bottom panels are isometric

 reference lines with arbitrary intercepts.

 Figure 4. Respiratory quotient, RQ, in relation to body mass in instars IV (a) and V (b).

 Points depict observations. The grey curves in (b) are cohort-specific regression curves based

 on the model-averaged coefficients (see Table 2).

 Figure 5. Instar V water content (a) and lipid content (b) in relation to body size in

 diapausing (closed circles) and directly developing (open circles) larvae, as well as

 comparison of proportion of water in fresh body matter (c) and proportion of lipids in

875   **dry body matter (d) between pupae and similarly-sized larvae.** The unbroken curves in

876   (a) and (b) are regression curves based on model-averaged coefficients (see Table 3), whereas

877   the dashed grey lines are isometric reference lines with an arbitrary intercept chosen for the

878   purpose of illustration. In (c) and (d), the boxes and whiskers illustrate the distribution of data

879   similarly as in Fig. 1.