1

The conserved transcriptional regulator CdnL is required for metabolic homeostasis and morphogenesis in *Caulobacter*

5 Selamawit Abi Woldemeskel[1], Laura Alvarez[2], Allison K. Daitch[1], Rilee Zeinert[3], Anant
6 Bhargava[1], Diego Gonzalez[4], Justine Collier[4], Peter Chien[3], Felipe Cava[2], Erin D. Goley[1]
7
1: Department of Biological Chemistry, Johns Hopkins University School of Medicine,
Baltimore, MD, USA
2: Department of Molecular Biology, Umeå University, Umeå, Sweden
3: Department of Biochemistry and Molecular Biology, University of Massachusetts-Amherst,
MA, USA
4: Department of Fundamental Microbiology, Faculty of Biology and Medicine, University of
Lausanne, Switzerland
15
16

Abstract

Bacterial growth and division require regulated synthesis of the macromolecules used to expand

and replicate components of the cell. Transcription of housekeeping genes required for metabolic

homeostasis and cell proliferation is guided by the sigma factor σ^{70}. The conserved CarD-like

transcriptional regulator, CdnL, associates with promoter regions where σ^{70} localizes and

stabilizes the open promoter complex. However, the contributions of CdnL to metabolic

homeostasis and bacterial physiology are not well understood. Here, we show that *Caulobacter*

crescentus cells lacking CdnL have severe morphological and growth defects. Specifically,

Δ*cdnL* cells grow slowly in both rich and defined media, and are wider, more curved, and have

shorter stalks than WT cells. These defects arise from transcriptional downregulation of most

major classes of biosynthetic genes. Notably, we find that Δ*cdnL* cells are severely limited in

glutamate synthesis, rendering them auxotrophic for that amino acid. Moreover, Δ*cdnL* cells

produce low amounts of the cell wall precursor lipid II, and Δ*cdnL* is synthetic lethal with other

genetic perturbations that limit lipid II production. Δ*cdnL* cells also have aberrant localization of

MreB and CtpS, cytoskeletal proteins required for maintaining proper cell width and curvature.

Interestingly, the localization of CtpS is dependent on availability of CTP, which is predicted to

be low in Δ*cdnL* cells. Our findings implicate CdnL as a global regulator of genes required for

metabolic homeostasis that impacts morphogenesis through availability of lipid II and through

metabolite-mediated changes in localization of cytoskeletal regulators of cell shape.

Main (400 words)

In order to replicate, bacterial cells must synthesize enough macromolecules to double in

size using nutrients available in their environment. These macromolecules are used to build

structural components of the cell such as the membrane and cell wall, to synthesize the multitude

of enzymes required for essential biochemical processes, and to duplicate genetic material.

Transcriptional control of housekeeping genes encoding factors that carry out these essential

functions contributes to maintaining metabolic homeostasis to support growth and development.

Transcription initiation of housekeeping genes is achieved by housekeeping sigma factors that

direct the core RNA polymerase (RNAP) to their promoter regions[1], but may be co-regulated by

other factors, including the CarD-like transcriptional regulator CdnL. CdnL is broadly conserved

in bacteria and is best-characterized in *Myxococcus xanthus* and in *Mycobacteria*[2–6]. It localizes

to promoter regions where the housekeeping sigma factor resides, directly binds to the RNAP β

subunit, stabilizes the open promoter complex, and is required for transcription from rRNA

promoters[2,3,5]. *Mycobacterium tuberculosis* CarD (the CdnL homolog) is essential for growth in

culture, persistence in mice, and survival during genotoxic stress and starvation[2,4]. CdnL is also

essential in *M. xanthus*, where its depletion causes cell filamentation[6]. This last observation

implies that CdnL-dependent changes in gene expression are required to support proper

morphogenesis, at least in *M. xanthus*.

Bacterial morphology is maintained by the peptidoglycan (PG) cell wall, which must be

expanded and remodeled in a regulated manner to allow for growth and division. It lies just

outside the inner membrane and is a continuous, covalently-linked structure made of glycan

strands crosslinked together by peptide side chains[7]. In addition to giving the cell its stereotyped

shape, the cell wall provides physical integrity and prevents cell lysis due to turgor pressure.

Different shape features (e.g. cell length, width, or curvature) are specified by spatially and

temporally regulated synthesis and remodeling of the PG that is largely orchestrated by

cytoskeletal proteins[7,8]. Though the details of PG synthesis and remodeling and their regulation

by cytoskeletal proteins has been the subject of intense study for several decades, how PG

metabolism is regulated in coordination with nutrient availability and the metabolic status of the

cell is incompletely understood.

Here, we investigate how CdnL affects metabolism, growth, and morphogenesis in the α-

proteobacterium *Caulobacter crescentus* (hereafter *Caulobacter*). *Caulobacter* undergoes

stereotyped morphological transitions as it progresses through the cell cycle and during

adaptation to a variety of stresses, making it an ideal organism to study bacterial morphogenesis[9].

Although it is known that *Caulobacter* CdnL is not required for viability, the consequences of its

loss on *Caulobacter* physiology have not been previously characterized[10]. Here, we demonstrate

that CdnL regulates transcription of biosynthetic genes required for metabolic homeostasis that,

in turn, supports proper growth, PG metabolism, and morphology in *Caulobacter*.

Results

<u>ΔcdnL cells have growth and morphology defects</u>

We became interested in *Caulobacter* CdnL through a screen for spontaneous suppressors of a

dominant lethal mutant of the cell division protein FtsZ (called FtsZΔCTL) that causes lethal

defects in PG metabolism[11]. Specifically, one of the FtsZΔCTL suppressors we identified carried

a point mutation in *cdnL* (*CCNA_00690*) encoding an I42N missense mutation that disrupted

CdnL protein stability (Supplementary figure 1a). To examine the role of CdnL in *Caulobacter*

growth and morphogenesis, we deleted it in a clean genetic background and compared growth

and morphology to wild type (WT) cells. We found that Δ*cdnL* cells grew more slowly than WT

(Figure 1a) and had pleiotropic morphological defects in complex PYE medium (Figure 1b, c).

Quantitative analysis of shape differences between WT and Δ*cdnL* cells grown in PYE using

CellTool[12] revealed significant differences in two shape modes which approximately correspond

to cell curvature (shape mode 2) and cell width (shape mode 3) (Figure 1c). Specifically, we

observed that in PYE, Δ*cdnL* cells were wider and more curved than WT cells (Figure 1b, c, d).

In these shape modes, Δ*cdnL* cells were also more variable in their range of values than WT,

indicating less precise maintenance of shape in the mutant (Figure 1c). Additionally, we

observed phase-light "ghost" cells in Δ*cdnL* samples indicative of cell lysis and suggesting a lack

of integrity in the cell envelope (Figure 1b, arrows).

To further characterize the Δ*cdnL* growth and morphology phenotypes, we grew WT and

Δ*cdnL* cells in the defined media M2G or HIGG. Surprisingly, we found that M2G was unable to

support growth of cells lacking CdnL, but that Δ*cdnL* cells grew in HIGG, albeit more slowly

than WT (Figure 1a). A comparison of the components of M2G and HIGG media suggested that

the missing nutrient in M2G that is required for growth of Δ*cdnL* cells might be glutamate.

Consistent with this, addition of sodium glutamate to M2G (called M2GG with sodium

glutamate added) supported growth of ΔcdnL cells at a rate similar to that in HIGG. These data

indicate that ΔcdnL cells are unable to synthesize sufficient glutamate to support growth and are

now glutamate auxotrophs (Figure 1a). No other amino acids are included in either M2G or

HIGG, suggesting a specific requirement for exogenous glutamate.

Similar to our observations with PYE, we found that ΔcdnL cells grew more slowly than

WT in M2GG or HIGG and also had aberrant shape (Figure 1a, b). ΔcdnL cells were again

hypercurved and wider than WT cells, but unlike in PYE, ΔcdnL cells were also consistently

shorter than WT in each defined medium (Figure 1b, c, d). The polar stalk is a prominent

morphological feature of *Caulobacter* cells and, like other features of cell shape, stalk biogenesis

and elongation require PG synthesis[13]. Since we observed stereotyped shape changes in cells

lacking CdnL, we also compared stalk morphogenesis between WT and ΔcdnL cells. We found

that in phosphate replete HIGG media (containing 200 μM PO_4^-), ΔcdnL cells had short stalks

compared to WT cells (Figure 1e, f), suggesting that CdnL impacts all aspects of morphogenesis

in *Caulobacter*. Stalks elongate in response to phosphate starvation in *Caulobacter* through a

pathway that is distinct from developmentally regulated stalk morphogenesis. Interestingly,

ΔcdnL cells significantly elongated their stalks during phosphate starvation (Figure 1 e, f)

suggesting that ΔcdnL cells can still alter their morphology in response to phosphate starvation.

Genes involved in biosynthetic processes are downregulated in ΔcdnL

Having established that the transcriptional regulator CdnL plays a role in growth and

morphogenesis, we next sought to understand how it exerts these effects. To identify genes that

are misregulated in ΔcdnL that may be responsible for the growth and morphology defects that

we observe, we extracted mRNA from WT and Δ*cdnL* cells and performed RNA-seq. The results

from this analysis showed that approximately 30% of the transcriptome is misregulated in cells

lacking CdnL (Supplementary table 1). A closer look at genes that are downregulated in the

Δ*cdnL* clone that we used for RNA-seq analysis suggested that a 50 kbp region comprising 51

genes between CCNA_02773 and CCNA_02826 was highly downregulated in Δ*cdnL*. This

region is flanked by identical sequences that encode transposases CCNA_02772 and

CCNA_02828, suggesting that this region may be deleted in the Δ*cdnL* clone (EG1415) used in

our RNA-seq analysis. Indeed, deletion of this region was confirmed by PCR. However, we

found no differences in growth or morphology between the Δ*cdnL* clone with the 50 kbp deletion

(EG1415, Δ50kbp) and Δ*cdnL* clones containing this region (e.g. EG1447), indicating that this

region does not contribute significantly to the growth and shape defects that we observe

(Supplementary figure 1b, c). Previously, this region of the chromosome was shown to be readily

lost in a background lacking the methyltransferase *ccrM*[14]. Similar to our observations with

Δ*cdnL*, loss of this region in Δ*ccrM* did not yield any specific growth advantages. To ensure that

loss of this region did not impact our analysis of the transcriptional consequences specific to loss

of CdnL, we analyzed the transcriptome of an independent Δ*cdnL* clone that has the 50 kbp

region intact (JC784) by microarray analysis (Supplementary table 1). We found significant

overlap between our RNA-seq dataset using EG1415 and microarray dataset using JC784

(Supplementary figure 1d, e, f, g). We generally used the intersection of our two datasets (RNA-

seq of EG1415 and microarray of JC874) as a conservative and rigorous representation of the

consequences of deleting *cdnL* on the *Caulobacter* transcriptome.

To obtain an overview of how loss of CdnL affects the transcriptome we used DAVID[15]

functional annotation analysis to functionally categorize genes that had at least a two-fold change

in transcript abundance in Δ*cdnL* in both the RNA-seq and microarray datasets. Of the genes that

are downregulated in Δ*cdnL*, we found an overrepresentation of genes involved in biosynthetic

and bioenergetic pathways including amino acid, nucleotide, fatty acid, lipid, cell wall, cell

membrane, and central carbon metabolism (Supplementary table 2). Genes that are over 2-fold

upregulated in Δ*cdnL* clustered into categories involved in transcription, transport, and motility.

<u>Δ*cdnL* cells have low levels of the cell wall precursor and changes in cell wall crosslinking</u>

Since our transcriptome analysis revealed downregulation of pathways involved in

macromolecular biosynthesis, we hypothesized that the severe defects in growth and morphology

may arise from limited amounts of substrates available for proliferative processes. Consistent

with our observations that Δ*cdnL* cells are unable to grow without exogenous glutamate, we

found that *gdhA* and *gltB*, genes essential for glutamate biosynthesis from α-ketoglutarate or

glutamine, are over 4-fold downregulated in both datasets (Figure 2a). Additionally, our findings

from the transcriptomic analysis indicated that genes required for PG precursor (lipid II)

biosynthesis are highly downregulated (Figure 2a) leading us to postulate that PG material may

be limiting in Δ*cdnL* cells. Since PG synthesis underlies both cell shape and cell envelope

integrity, changes in abundance of lipid II might underlie the shape and cell lysis phenotypes

observed for Δ*cdnL* cells. We directly compared lipid II levels in WT and Δ*cdnL* cells and found

that Δ*cdnL* cells have a striking 45% reduction in lipid II levels as compared to WT (Figure 2b).

To assess if CdnL-mediated changes in transcription result in alterations to PG metabolic

activities in addition to lipid II synthesis, we performed muropeptide analysis on PG isolated

from WT and Δ*cdnL* cells to identify changes in PG chemistry. Consistent with low levels of

lipid II, Δ*cdnL* cells have 23% less PG polymer than WT (Figure 2c). Although overall PG

crosslinking is unaffected, Δ*cdnL* cells had significantly more dimers and fewer trimers (higher

order crosslinks) than WT (Figure 2d, Supplementary Figure 2b). The mutant also shows an

increased chain length, as the relative amount of anhydro muropeptides (glycan chain termini)

were significantly reduced in Δ*cdnL* cells (Figure 2e, Supplementary Figure 2b). Moreover,

Δ*cdnL* cells had more pentapeptides and glycine-containing muropepetides compared to WT

(Figure 2e, Supplementary Figure 2b). Thus, the shape and envelope integrity defects observed

in Δ*cdnL* cells may arise, at least in part, from a combination of low levels of lipid II and

significant changes in the chemical structure of the cell wall.

<u>Pathways impacting PG metabolism become essential in Δ*cdnL*</u>

To gain insight into which transcriptional changes in Δ*cdnL* may be most relevant to fitness*,* we

performed comparative transposon sequencing (Tn-Seq) on WT and Δ*cdnL* strains. Consistent

with morphological defects and low amounts of lipid II observed for Δ*cdnL* cells, we found

several normally non-essential genes involved in pathways that interact with PG synthesis that

could not be disrupted in the Δ*cdnL* background (Figure 3a, Supplementary table 3).

Specifically, we identified genes involved in glutamate metabolism, PG metabolism, PG

recycling, and capsule synthesis that could not be disrupted in the Δ*cdnL* background (Figure

3a,b, Supplementary table 3). Of the genes involved in glutamate metabolism, *gdhZ* and *murI*

become essential in Δ*cdnL*. Each of these proteins link glutamate metabolism to morphogenesis:

MurI by making D-Glu for lipid II synthesis and GdhZ via its effects on the morphogenetic

protein FtsZ[16,17].

Our Tn-Seq analysis also indicated that *ampG* and *ampD* (*CCNA_02650*), genes required

for PG recycling, become essential in Δ*cdnL* (Figure 3b, Supplementary table 3). AmpG is a

permease that allows transport of cell wall material from the periplasm to the cytoplasm, and

AmpD degrades transported cell wall material to provide building blocks for lipid II synthesis[18].

We attempted to delete *cdnL* in a Δ*ampG* backgroundf but were unable to obtain clones with

deletions in both *ampG* and *cdnL*, supporting the genetic interaction suggested from the Tn-Seq.

Since lipid II levels are low in Δ*cdnL*, the cell apparently relies on pathways that allow for

recycling of the existing cell wall material for survival.

In addition to PG metabolic genes, we found putative synthetic lethal interactions

between capsule (exopolysaccharide) biosynthetic genes and *cdnL* (Figure 3a,b, Supplementary

table 3). During capsule synthesis, saccharide precursors are synthesized on the lipid anchor

undecaprenyl phosphate (Und-P) in the cytoplasm, flipped to the periplasmic space, and

assembled into a polysaccharide that is exported to the outer surface of the cell[19]. Thus, PG

synthesis and capsule biosynthesis both use Und-P as a lipid carrier for synthesis and export into

the periplasmic space. Previous work in *Escherichia coli* and *Bacillus subtilis* has shown that

disruption of pathways that use Und-P can lead to Und-P sequestration and cause severe growth

and morphological defects by inhibiting lipid II synthesis[20–22]. Similarly, we propose that

disrupting the capsule biosynthesis pathway in *Caulobacter* may sequester Und-P in dead-end

intermediates, limiting lipid II synthesis. We attempted to delete each of the capsule genes

predicted by Tn-Seq to genetically interact with *cdnL* in Δ*cdnL* cells. We were unsuccessful with

the exception of *CCNA_00164*, which encodes a putative O-antigen polymerase ligase and which

had the weakest predicted genetic interaction with *cdnL* of the capsule genes (Figure 3a, b and

Supplementary table 3). Nevertheless, deletion of both *cdnL* and *CCNA_00164* resulted in cells

with morphological defects that are more severe than Δ*cdnL* alone (Figure 3d). Moreover, these

double mutant cells grew more slowly than Δ*cdnL* (Figure 3c). Deletion of *CCNA_00164* by

itself had no obvious effect on growth or morphology. Our data support the model that mutations

in the capsule biosynthesis pathway sequester Und-P and limit its availability for lipid II

synthesis, exacerbating the existing lipid II deficiency of Δ*cdnL* cells to a lethal level.

Collectively, our transcriptomics, biochemical analyses, and genetic interaction results indicate

that CdnL is required to maintain adequate availability of the PG precursor lipid II to support

normal growth and morphology.

<u>MreB is hyperfocused at midcell in Δ*cdnL* cells</u>

Limitation of lipid II levels has previously been shown to cause an increase in cell width in

Caulobacter[23,24]. However, the morphological defects that we observe in Δ*cdnL* cells are distinct

from shape changes that occur when lipid II levels are low, suggesting additional players in the

shape defects of Δ*cdnL* cells. To investigate this possibility, we explored other proteins

implicated in shape regulation in *Caulobacter*, with a focus on the polymerizing proteins MreB,

FtsZ, crescentin, and CTP synthase. Since CdnL is a transcriptional regulator, we first probed for

changes in levels of these factors and found no differences in protein or transcript levels between

Δ*cdnL* and WT cells (Supplementary table 2 and data not shown).

Since Δ*cdnL* cells had significant differences in width and curvature, we focused on the

cytoskeletal proteins that determine these shape features. In bacteria, the actin homolog and

ATPase MreB is required for rod shape, width maintenance, and cell body elongation[25,26].

Additionally, MreB influences curvature of *Caulobacter* cells through its effects on the

intermediate filament-like protein, crescentin[27]. To determine if the localization of MreB is

altered in Δ*cdnL*, we expressed *venus-mreB* from the *xylX* promoter and visualized MreB

localization in WT or Δ*cdnL* cells grown in PYE, M2G, and HIGG media (Figure 4 a,b,c). In

WT cells grown in PYE, MreB has a patchy localization along the length of the cell in newborn

swarmer (G1) cells, shifts to midcell in stalked (S) and early predivisional cells, and assumes a

patchy localization in late predivisional cells[25]. Interestingly, we found that the localization of

MreB at midcell is more tightly focused in Δ*cdnL* cells compared to WT in all three media tested

(Figure 4a,b,c). Moreover, we found that most WT cells localize MreB along the length of the

cell when grown in HIGG media (Figure 4c). However, the localization of MreB in HIGG shifts

to mostly at midcell when *cdnL* is deleted. To better visualize the localization of MreB across

populations and throughout the cell cycle, we created demographs. In cells that have MreB

localized at midcell, we found that the localization of MreB is more tightly focused with minimal

signal outside of the midcell region (Figure 4d,e,f). To quantitatively compare the differences in

MreB localization between WT and Δ*cdnL*, we fit the MreB signal at midcell to an eight term

Fourier series model and calculated the full width at half max (FWHM). We found that the

FWHM at midcell is smaller in Δ*cdnL* cells compared to WT, consistent with what we inferred

from the images and demographs (Figure 4g,h,i).

To further characterize the link between MreB localization and loss of *cdnL,* we deleted

cdnL in backgrounds that have mutations near the ATP binding pocket of MreB. These

mutations cause MreB to localize at the poles, along the length of the cell, or at midcell

throughout the cell cycle[27]. These aberrant localizations of MreB cause changes in cell shape but

do not affect growth rate (Supplementary figure 3a,b,c). Deletion of *cdnL* in each of these

backgrounds caused severe morphological and growth defects (Supplementary figure 3a,b). The

severity of growth and shape defects was worse in mutant backgrounds that do not localize MreB

to midcell (Supplementary figure 3). Interestingly, we found that MreB V324A, which has a

patchy localization in a WT background, becomes enriched at midcell when *cdnL* is deleted and

grows similarly to mutants that localize MreB to midcell (Supplementary figure 3b,c,d,e). These

data suggest that localization of MreB at midcell is important for fitness in Δ*cdnL* cells. Since

MreB is essential for specification and maintenance of width and curvature, the mislocalization

of MreB may contribute to the width and curvature defects that we observe in Δ*cdnL* cells.

Since we found that Δ*cdnL* cells have low lipid II levels compared to WT and lipid II has

been linked to localization of MreB in *B. subtilis*[28], we wanted to determine if inhibiting lipid II

synthesis in *Caulobacter* cells was sufficient to cause mislocalization of MreB similar to Δ*cdnL*.

To test this, we inhibited MurA with a low dose of fosfomycin and visualized the localization of

Venus-MreB over time. We observed an increase in FWHM of midcell-localized MreB and cell

width after 5.5 hours of fosfomycin treatment (Supplementary Figure 4a,b,c). The changes in

localization of MreB during fosfomycin treatment are opposite to what we observe during *cdnL*

deletion, suggesting other mechanisms by which MreB is enriched at midcell in Δ*cdnL* cells.

<u>Hypercurvature of Δ*cdnL* cells is dependent on crescentin and may involve CTP synthase</u>

In *Caulobacter*, crescentin drives the patterning of PG synthesis required for cell curvature[8,29].

To determine if hypercurvature of Δ*cdnL* cells is dependent on crescentin, we deleted *cdnL* in a

Δ*creS* background. We observed that Δ*cdnL*Δ*creS* cells are straight, indicating that

hypercurvature is dependent on crescentin (Figure 5a,b). In addition to crescentin, curvature is

influenced by the metabolic enzyme CTP synthase (CtpS), which forms filaments near crescentin

when CTP is abundant[30,31]. CtpS filaments negatively regulate curvature in a crescentin-

dependent manner[31]. Since we observed that pyrimidine biosynthetic genes are downregulated

from our RNA-Seq analysis (Figure 5c), we hypothesized that the hypercurvature of Δ*cdnL* may

arise due to fewer CtpS filaments caused by low levels of CTP in Δ*cdnL*. To gain support for this

hypothesis, we expressed *mcherry-ctpS* from the xylose promoter in WT or Δ*cdnL* cells and

visualized its localization (Supplementary figure 5). Demograph analysis of mCherry-CtpS

localization indicated that mCherry-CtpS forms less robust filaments and has more dispersed

signal in Δ*cdnL* cells grown in PYE or M2GG compared to WT (Figure 5d, Supplementary

figure 5). Our data support the hypothesis that low levels of CTP in Δ*cdnL* cells lead to

formation of fewer CtpS filaments than in WT, which consequently yields the hypercurvature we

observe.

288
289

Discussion

In this work, we unveil a new role for the conserved global transcriptional regulator

CdnL in mediating transcription of biosynthetic genes required for metabolic homeostasis in

Caulobacter. We show that in Δ*cdnL* cells, transcription of many classes of biosynthetic genes is

downregulated, with physiologically critical defects in glutamate and lipid II metabolism in

particular. These changes in metabolic pools affect growth and morphology directly by limiting

macromolecules required for envelope expansion and indirectly by influencing morphogenetic

proteins. Downregulation of genes involved in biosynthetic and bioenergetics pathways in Δ*cdnL*

cells and the reliance of Δ*cdnL* cells on exogenous glutamate, which is a central node in multiple

biosynthetic pathways, suggests that CdnL is a global regulator of genes required for central

metabolism and macromolecular synthesis[32,33].

Previously, CdnL has been shown to directly bind to RNAP and stabilize the open

promoter complex required for transcription of rRNA promoters in *Mycobacteria* and *M.*

xanthus[3,5]. Similarly, *Caulobacter* CdnL has been shown to bind RNAP and regulate

transcription from an rRNA promoter[10]. Here, we demonstrate that loss of CdnL has global

effects on various aspects of bacterial metabolism. Interestingly, however, transcriptomic

changes that occur during loss of CdnL in *Caulobacter* are the opposite of what is observed in

Mycobacterium smegmatis. CarD depletion in *M. smegmatis* leads to upregulation of genes

encoding the translation machinery and enzymes in metabolic pathways, such as amino acid

biosynthesis, and to downregulation of genes involved in degradation of amino acids, fatty acid

metabolism and those coding for membrane proteins[2]. Although loss of CdnL generally has

opposite effects on expression of metabolic genes in these two divergent bacteria, CdnL-

mediated effects on the transcriptome impact proliferative and metabolic processes and are

thereby essential for proper growth in *Caulobacter* and viability in *Mycobacteria*.

In *Caulobacter*, loss of CdnL causes aberrant morphogenesis, at least in part, by limiting

production of lipid II. The effects of *cdnL* deletion are in some ways reminiscent of loss of the

master regulator and RNA chaperone Hfq in *Caulobacter*[23]. Deletion of *hfq* alters the levels of

several metabolites, including glutamate and components of the TCA cycle. Importantly, this

includes accumulation of α-oxoglutarate which, in turn, causes reduced synthesis of lipid II and

pleiotropic cell shape defects[23]. Though both Hfq and CdnL are required to maintain sufficiently

high levels of lipid II to support growth and stereotyped cell shape, CdnL has a distinct shape

phenotype, likely mediated by additional impacts on MreB and CtpS.

In recent years, a growing number of factors have been described that link the metabolic

status of the cell to morphology through cytoskeletal proteins [7,33,34]. The best-studied examples

of this employ moonlighting functions of metabolic enzymes to couple metabolism to

morphology. For example, the glucosyltransferases UgtP and OpgH in *B. subtilis* and *E. coli,*

respectively, alter FtsZ dynamics and affect cell size in a UDP-glucose dependent manner[35,36].

Similarly, in *Caulobacter*, GdhZ and KidO alter FtsZ assembly in a glutamate-dependent

manner, coupling metabolism to cell division[16,37]. In the case of CdnL, its impact on the

localization of the cytoskeletal proteins MreB and CtpS appears to be metabolite-mediated rather

than protein-mediated. Polymerization of CtpS into filaments that negatively regulate cell

curvature is dependent on CTP, which is predicted to be present in low amounts in Δ*cdnL* cells.

It is unclear how MreB localization is regulated in *Caulobacter*, though it appears to be linked to

its ATP-dependent polymerization cycle[27] and may involve cell wall precursors other than lipid

II. Thus, we propose that the disruption of metabolic homeostasis that arises from loss of CdnL

impacts morphology through metabolite-mediated effects on cytoskeletal proteins in addition to

reduced levels of lipid II.

Interestingly, depletion of CdnL in *M. xanthus* results in cell filamentation, consistent

with a conserved impact on morphogenesis[6]. The effects, if any, of CarD depletion in

Mycobacteria on cell morphology have not been reported, nor have the effects of CdnL depletion

on the *M. xanthus* transcriptome or metabolome. Future work in diverse organisms will reveal if

the role of CdnL as a global regulator of both morphogenesis and metabolism is conserved

across bacteria.

343
344

Methods

Bacterial strains, plasmids, growth conditions

Caulobacter crescentus NA1000 strains were grown in peptone yeast extract medium (PYE),

M2G[38], M2GG (M2G with 0.15% sodium glutamate), or Hutner base imidazole-buffered glucose

glutamate medium (HIGG)[39]. Xylose was used at 0.3% for induction experiments. EG1415 and

JC874 were used for RNA-Seq and microarray experiments, respectively. All other experiments

with Δ*cdnL* were performed using EG1447 (clean, in-frame deletion of *cdnL*) or EG1898 (*cdnL*

coding sequence replaced with *aacC1,* conferring gentamicin resistance) or derivatives thereof.

JC874, EG1447, and EG1898 contain the 50 kbp region that is deleted in EG1415. Growth

curves were performed in a Tecan Infinite M200 Pro plate reader, 100 µL culture volume, with

the OD_{600} of each strain monitored in biological triplicates every 30 min with intermittent

shaking. Antibiotics for growing *Caulobacter* and cloning and purifying plasmids from *E. coli*

were used at concentrations in liquid (solid) media as described in Woldemeskel *et al* 2017[40]. *C.*

crescentus: kanamycin 5 (25) µg/mL, gentamicin 1 (5) µg/mL. *E. coli:* ampicillin 50 (100)

µg/mL, kanamycin 30 (50) µg/mL, gentamicin 15 (20) µg/mL, and spectinomycin 50 (50)

µg/mL.

Phase contrast, ensemble fluorescence microscopy and image analysis

Cells in exponential phase of growth were spotted on 1% agarose pads and imaged using a Nikon

Eclipse Ti inverted microscope equipped with a Nikon Plan Fluor 100X (NA1.30) oil Ph3

objective and Photometrics CoolSNAP HQ[2] cooled CCD camera. Chroma filter cubes ET-EYFP

and ET-dsRED were used for Venus and mCherry, respectively. Images were processed using

Adobe Photoshop and demographs were made using Oufti[41]. Principal component analysis to

identify shape variations were performed using CellTool[12] after phase contrast images were

converted to binary masks in ImageJ[42] and edited in Adobe Photoshop to remove overlapping

cells. Full-width at half-maximum (FWHM) calculations of midcell MreB signal from Oufti[41]

output were performed using a custom MATLAB script which fit the normalized signal output

from Oufti[41] into an eighth term Fourier series model. Cells for which the fit had an adjusted R

square value greater than 0.85 were chosen for FWHM calculation. Cells that have midcell

localized MreB were selected for FWHM determination based on cell length bins guided by

demographs.

Immunoblotting to assess CdnL levels

His_6-SUMO-CdnL was overproduced in Rosetta (DE3) pLysS *E.coli* cells from pEG1129 by

inducing for 3 hr at 30 ^{o}C with 0.5 mM IPTG at OD_{600} of 0.5. Cells were harvested by

centrifugation, resuspended in lysis buffer (50 mM Tris-HCl, pH 8.0, 300 mM KCl, 20 mM

imidazole, and 10% glycerol), lysed with 1 mg/mL lysozyme, sonicated and centrifuged at 4 ^{o}C

for 30 min at 15000 x g. Supernatant was loaded onto a HisTrap FF 1 mL column (GE Life

Sciences) and eluted with lysis buffer with 300 mM imidazole, cleaved with His_6-Ulp1 (SUMO

protease) at 1:500 (protease: fusion) molar ratio overnight while dialyzing into lysis buffer.

Sample was reloaded onto HisTrap FF 1mL and flow through collected to separate His_6-SUMO

and His_6-Ulp1 from CdnL. CdnL fraction was dialyzed into PBS prior to antibody production.

Cells in log phase were isolated and lysed in SDS-PAGE loading buffer by boiling for 5 min.

Standard procedures were used for SDS-PAGE and transfer to nitrocellulose membrane. CdnL

antisera was generated by immunizing a rabbit with CdnL purified as above (Pocono Rabbit

Farm & Laboratory). Specificity was determined using cell lysates with deleted or overexpressed

cdnL. CdnL antisera was used at 1:5000.

RNA-Seq preparation

Cultures of three independent colonies of WT (EG865) and Δ*cdnL* (EG1415, Δ50kbp) cells were

grown in PYE and harvested at OD_{600} of 0.45 and total RNA was extracted using PureLink RNA

Mini Kit from Thermo Fisher using the protocol provided with the kit. Briefly, cells were

harvested, lysed, and homogenized using lysozyme, SDS, provided Lysis buffer, and

homogenizer. Nucleic acids were extracted with ethanol and loaded onto the provided spin

cartridge. DNA-free total RNA was extracted using the on-column PureLink DNAse Treatment

protocol. PureLink DNAse mixture was directly added on to the spin cartridge membrane,

incubated for 15 min and washed using provided wash buffers and RNA was eluted with

RNAse-free water. Finally, RNA quality was quantified using a bioanalyzer and rRNA was

removed using the Ribo-Zero rRNA Removal Kit (Gram-Negative Bacteria). RNA-Seq libraries

were prepared using the Illumina TruSeq stranded RNA kit and sequenced on an Illumina HiSeq

2500. Data analysis was performed using the Illumina's CASAVA 1.8.2, Bowtie2 (v 2.2.5), and

Bioconductor's DESeq package. rRNA clean up, quality control analysis, library prep,

sequencing and analysis was performed by the Johns Hopkins University School of Medicine

Next Generation Sequencing Center.

Microarray preparations

Biological triplicates of WT (JC450) and Δ*cdnL* (JC784) cultures grown in PYE were harvested

at OD_{660} of 0.3. RNA was extracted with Trizol, purified using Ambion PureLink RNA mini-kit

according to manufacturer's protocol. Double stranded cDNA was prepared with Roche Double

Stranded cDNA synthesis kit. Microarray analysis was performed on Nimblegen custom chips

following the manufacturer's protocol for labeling, hybridization on 4-plex chips, and washing.

Scanning was performed with Agilent microarray scanner at 3 μm and normalization was

performed with Nimblescan implemented RMA algorithm (3 x 4-plex chips in total, all samples

from the same culture conditions). The statistical analysis of significance was performed with R

using the Significance of Analysis of Microarrays (SAM) method implemented in the library

siggenes.[43]

DAVID analysis and functional clustering

Genes with a two-fold change in transcript levels in both RNA-Seq and microarray analysis were

functionally categorized using DAVID[15] with a medium classification stringency and default

parameters.

Transposon library preparation, sequencing, and analysis

Two Tn-Seq libraries each were generated for WT (EG865) and Δ*cdnL* (EG1447). 1L PYE

cultures were harvested at OD_{600} of 0.4-0.6, washed 5 times with 10% glycerol, and

electroporated with the Ez-Tn5 <Kan-2> transposome (Epicentre). Cells were recovered for 90

minutes at 30 $^{\circ}$C with shaking, and plated on PYE-Kan plates. WT libraries were grown for 5

and 6 days and Δ*cdnL* libraries were grown for 6 and 10 days. Colonies were scraped off plates,

combined, resuspended to form a homogeneous solution in PYE, and flash frozen in 20%

glycerol. Genomic DNA for each library was extracted from an aliquot using the DNeasy Blood

and Tissue Kit (Qiagen). Libraries were then prepared for Illumina Next-Generation sequencing

through sequential PCR reactions. The first PCR round used arbitrary hexamer primers with a

Tn5 specific primer going outward. The second round used indexing primers with unique

identifiers to filter artifacts arising from PCR duplicates. Indexed libraries were pooled and

sequenced at the University of Massachusetts Amherst Genomics Core Facility on the NextSeq

550 (Illumina).

Sequencing reads were first demultiplexed by index, each library was concatenated and clipped

of the molecular modifier added in the second PCR using Je[44]:

java -jar /je_1.2/je_1.2_bundle.jar clip F1=compiled.gz LEN=6

Reads were then mapped back to the *Caulobacter crescentus* NA1000 genome (NCBI Reference

Sequence: NC_011916.1) using BWA[45] and sorted using Samtools[46]:

bwa mem -t2 clipped.gz | samtools sort -@2 - > sorted.bam

Duplicates were removed using Je[44] and indexed with Samtools[46] using the following command:

java -jar /je_1.2/je_1.2_bundle.jar markdupes I=sorted.bam O=marked.bam M=METRICS.txt

MM=0 REMOVE_DUPLICATES=TRUE

samtools index marked.bam

The 5' insertion site of each transposon were converted into .wig files comprising of counts per

position and visualized using Integrative Genomics Viewer (IGV)[47,48]. Specific hits for each

library were determined with coverage and insertion frequency using a bedfile containing all

open reading frames from NC_011916.1 and the outer 20% of each removed to yield a clean and

thorough insertion profile. This was determined using BEDTools[49,50] and the following

commands:

bedtools genomecov -5 -bg marked.bam > marked.bed

bedtools map -a NA1000.txt -b marked.bed -c 4 > output.txt

Library comparisons were performed using the edgeR package in the Bioconductor suite using a

quasi-likelihood F-test (glmQLFit) to determine the false discovery rate adjusted p-values

reported here.

Lipid II extraction and analysis

Precursor extraction was performed as described previously and performed in triplicates[28].

Briefly, 500 mL of WT and $\Delta cdnL$ were grown in PYE to OD_{600} of 0.45. Cells were harvested,

resuspended in 5 mL PBS and poured into 50 mL flasks containing 20 mL $CHCl_3$:Methanol

(1:2). The mixture was stirred for 1 h at room temperature and centrifuged for 10 min at 4000 x g

at 4 ºC. The supernatant was transferred to clean 250 mL flasks containing 12 mL $CHCl_3$ and 9

mL PBS, stirred for 1 h at room temperature and centrifuged for 10 min at 4000 x g at 4 ºC. The

interface fraction (between the top aqueous and bottom organic layers) was collected and

vacuum dried. To remove lipid tail, samples were resuspended in 100 μL DMSO and 800 μL

H_2O, 100 μL ammonium acetate 100 mM pH 4.2 were added. This mixture was boiled for 30

min, dried in vacuum and resuspended in 300 μL H_2O. Samples were analyzed by UPLC

chromatography coupled to MS/MS analysis, using a Xevo G2- XS QTof system (Waters

Corporation, USA). The separation method used is identical to the one used for muropeptide

separation explained below. A library of compounds was used to target the identification of

peptidoglycan precursors and possible intermediates, although only lipid II (lipid II-penta) was

detected in these samples. Lipid II amounts were calculated based on the integration of the peaks

(total area), normalized to the culture OD.

Peptidoglycan (PG) purification and analysis

PG samples were analyzed as described previously[51,52]. Briefly, 50 mL of WT and Δ*cdnL* cells

were grown to an OD_{600} of 0.5 in PYE, harvested, and boiled in 5% SDS for 2 h. Sacculi were

repeatedly washed with MilliQ water by ultracentrifugation (110,000 rpm, 10 min, 20 ºC) until

total removal of the detergent and finally treated with muramidase (100 μg/mL) for 16 hours at

37 ºC. Muramidase digestion was stopped by boiling and coagulated proteins were removed by

centrifugation (10 min, 14,000 rpm). For sample reduction, the pH of the supernatants was

adjusted to pH 8.5-9.0 with sodium borate buffer and sodium borohydride was added to a final

concentration of 10 mg/mL. After incubating for 30 min at room temperature, pH was adjusted

to 3.5 with orthophosphoric acid.

UPLC analyses of muropeptides were performed on a Waters UPLC system (Waters

Corporation, USA) equipped with an ACQUITY UPLC BEH C18 Column, 130Å, 1.7 μm, 2.1

mm X 150 mm (Waters, USA) and a dual wavelength absorbance detector. Elution of

muropeptides was detected at 204 nm. Muropeptides were separated at 45 °C using a linear

gradient from buffer A (formic acid 0.1% in water) to buffer B (formic acid 0.1% in acetonitrile)

in an 18-minute run, with a 0.25 mL/min flow. Relative total PG amounts were calculated by

comparison of the total intensities of the chromatograms (total area) from three biological

replicates normalized to the same OD_{600} and extracted with the same volumes. Muropeptide

identity was confirmed by MS/MS analysis, using a Xevo G2-XS QTof system (Waters

Corporation, USA). Quantification of muropeptides was based on their relative abundances

(relative area of the corresponding peak) normalized to their molar ratio.

References

1. Österberg, S., Peso-Santos, T. del & Shingler, V. Regulation of Alternative Sigma Factor Use. *Annu. Rev. Microbiol.* **65**, 37–55 (2011).

2. Stallings, C. L. *et al.* CarD Is an Essential Regulator of rRNA Transcription Required for Mycobacterium tuberculosis Persistence. *Cell* **138**, 146–159 (2009).

3. Srivastava, D. B. *et al.* Structure and function of CarD, an essential mycobacterial transcription factor. *Proc. Natl. Acad. Sci.* **110**, 12619–12624 (2013).

4. Weiss, L. A. *et al.* Interaction of CarD with RNA Polymerase Mediates Mycobacterium tuberculosis Viability, Rifampin Resistance, and Pathogenesis. *J. Bacteriol.* **194**, 5621–5631 (2012).

5. Gallego-García, A. *et al.* Structural Insights into RNA Polymerase Recognition and Essential Function of Myxococcus xanthus CdnL. *PLOS ONE* **9**, e108946 (2014).

6. García-Moreno, D. *et al.* CdnL, a member of the large CarD-like family of bacterial proteins, is vital for Myxococcus xanthus and differs functionally from the global transcriptional regulator CarD. *Nucleic Acids Res.* **38**, 4586–4598 (2010).

7. Typas, A., Banzhaf, M., Gross, C. A. & Vollmer, W. From the regulation of peptidoglycan synthesis to bacterial growth and morphology. *Nat. Rev. Microbiol.* **10**, 123–136 (2012).

8. Cabeen, M. T. *et al.* Bacterial cell curvature through mechanical control of cell growth. *EMBO J.* **28**, 1208–1219 (2009).

9. Woldemeskel, S. A. & Goley, E. D. Shapeshifting to Survive: Shape Determination and Regulation in Caulobacter crescentus. *Trends Microbiol.* **25**, 673–687 (2017).

10. Gallego-García, A. *et al. Caulobacter crescentus* CdnL is a non-essential RNA polymerase-

binding protein whose depletion impairs normal growth and rRNA transcription. *Sci. Rep.* **7**,

43240 (2017).

11. Sundararajan, K. *et al.* The bacterial tubulin FtsZ requires its intrinsically disordered linker

to direct robust cell wall construction. *Nat. Commun.* **6**, 7281 (2015).

12. Pincus, Z. & Theriot, J. A. Comparison of quantitative methods for cell-shape analysis. *J.*

Microsc. **227**, 140–156 (2007).

13. Seitz, L. C. & Brun, Y. V. Genetic Analysis of Mecillinam-Resistant Mutants ofCaulobacter

crescentus Deficient in Stalk Biosynthesis. *J. Bacteriol.* **180**, 5235–5239 (1998).

14. Gonzalez, D. & Collier, J. Genomic Adaptations to the Loss of a Conserved Bacterial DNA

Methyltransferase. *mBio* **6**, e00952-15 (2015).

15. Dennis, G. *et al.* DAVID: Database for Annotation, Visualization, and Integrated Discovery.

Genome Biol. **4**, P3 (2003).

16. Beaufay, F. *et al.* A NAD-dependent glutamate dehydrogenase coordinates metabolism with

cell division in Caulobacter crescentus. *EMBO J.* **34**, 1786–1800 (2015).

17. Doublet, P., Heijenoort, J. van, Bohin, J. P. & Mengin-Lecreulx, D. The murI gene of

Escherichia coli is an essential gene that encodes a glutamate racemase activity. *J. Bacteriol.*

175, 2970–2979 (1993).

18. Park, J. T. Why does Escherichia coli recycle its cell wall peptides? *Mol. Microbiol.* **17**, 421–

426 (1995).

19. Ardissone, S. *et al.* Cell cycle constraints on capsulation and bacteriophage susceptibility.

eLife **3**, (2014).

20. Jorgenson, M. A., Kannan, S., Laubacher, M. E. & Young, K. D. Dead-end intermediates in

the enterobacterial common antigen pathway induce morphological defects in Escherichia

coli by competing for undecaprenyl phosphate. *Mol. Microbiol.* **100**, 1–14 (2016).

21. Jorgenson, M. A. & Young, K. D. Interrupting Biosynthesis of O Antigen or the

Lipopolysaccharide Core Produces Morphological Defects in Escherichia coli by

Sequestering Undecaprenyl Phosphate. *J. Bacteriol.* **198**, 3070–3079 (2016).

22. D'Elia, M. A. *et al.* Probing Teichoic Acid Genetics with Bioactive Molecules Reveals New

Interactions among Diverse Processes in Bacterial Cell Wall Biogenesis. *Chem. Biol.* **16**,

548–556 (2009).

23. Irnov, I. *et al.* Crosstalk between the tricarboxylic acid cycle and peptidoglycan synthesis in

Caulobacter crescentus through the homeostatic control of α-ketoglutarate. *PLOS Genet.* **13**,

e1006978 (2017).

24. Harris, L. K. & Theriot, J. A. Relative Rates of Surface and Volume Synthesis Set Bacterial

Cell Size. *Cell* **165**, 1479–1492 (2016).

25. Figge, R. M., Divakaruni, A. V. & Gober, J. W. MreB, the cell shape-determining bacterial

actin homologue, co-ordinates cell wall morphogenesis in Caulobacter crescentus. *Mol.*

Microbiol. **51**, 1321–1332 (2004).

26. Takacs, C. N. *et al.* MreB Drives De Novo Rod Morphogenesis in Caulobacter crescentus

via Remodeling of the Cell Wall. *J. Bacteriol.* **192**, 1671–1684 (2010).

27. Dye, N. A., Pincus, Z., Fisher, I. C., Shapiro, L. & Theriot, J. A. Mutations in the nucleotide

binding pocket of MreB can alter cell curvature and polar morphology in Caulobacter. *Mol.*

Microbiol. **81**, 368–394 (2011).

28. Qiao, Y. *et al.* Lipid II overproduction allows direct assay of transpeptidase inhibition by β-

lactams. *Nat. Chem. Biol.* **13**, 793–798 (2017).

29. Ausmees, N., Kuhn, J. R. & Jacobs-Wagner, C. The Bacterial Cytoskeleton: An Intermediate

Filament-Like Function in Cell Shape. *Cell* **115**, 705–713 (2003).

30. Barry, R. M. *et al.* Large-scale filament formation inhibits the activity of CTP synthetase.

eLife **3**, (2014).

31. Ingerson-Mahar, M., Briegel, A., Werner, J. N., Jensen, G. J. & Gitai, Z. The metabolic

enzyme CTP synthase forms cytoskeletal filaments. *Nat. Cell Biol.* **12**, 739–746 (2010).

32. Commichau, F. M., Forchhammer, K. & Stülke, J. Regulatory links between carbon and

nitrogen metabolism. *Curr. Opin. Microbiol.* **9**, 167–172 (2006).

33. Sperber, A. M. & Herman, J. K. Metabolism Shapes the Cell. *J. Bacteriol.* **199**, e00039-17

(2017).

34. Cabeen, M. T. & Jacobs-Wagner, C. Bacterial cell shape. *Nat. Rev. Microbiol.* **3**, 601–610

(2005).

35. Hill, N. S., Buske, P. J., Shi, Y. & Levin, P. A. A Moonlighting Enzyme Links Escherichia

coli Cell Size with Central Metabolism. *PLOS Genet.* **9**, e1003663 (2013).

36. Weart, R. B. *et al.* A metabolic sensor governing cell size in bacteria. *Cell* **130**, 335–347

(2007).

37. Radhakrishnan, S. K., Pritchard, S. & Viollier, P. H. Coupling Prokaryotic Cell Fate and

Division Control with a Bifunctional and Oscillating Oxidoreductase Homolog. *Dev. Cell* **18**,

90–101 (2010).

38. Marks, M. E. *et al.* The Genetic Basis of Laboratory Adaptation in Caulobacter crescentus. *J.*

Bacteriol. **192**, 3678–3688 (2010).

39. Poindexter, J. S. Selection for Nonbuoyant Morphological Mutants of Caulobacter

crescentus. **135**, 5 (1978).

40. Woldemeskel, S. A., McQuillen, R., Hessel, A. M., Xiao, J. & Goley, E. D. A conserved

coiled-coil protein pair focuses the cytokinetic Z-ring in Caulobacter crescentus. *Mol.*

Microbiol. **105**, 721–740 (2017).

41. Paintdakhi, A. *et al.* Oufti: An integrated software package for high-accuracy, high-

throughput quantitative microscopy analysis. *Mol. Microbiol.* **99**, 767–777 (2016).

42. Schneider, C. A., Rasband, W. S. & Eliceiri, K. W. NIH Image to ImageJ: 25 years of image

analysis. *Nat. Methods* **9**, 671–675 (2012).

43. Schwender, H., Krause, A. & Ickstadt, K. Comparison of the Empirical Bayes and the

Significance Analysis of Microarrays. 26

44. Girardot, C., Scholtalbers, J., Sauer, S., Su, S.-Y. & Furlong, E. E. M. Je, a versatile suite to

handle multiplexed NGS libraries with unique molecular identifiers. *BMC Bioinformatics* **17**,

419 (2016).

45. Li, H. & Durbin, R. Fast and accurate long-read alignment with Burrows–Wheeler transform.

Bioinformatics **26**, 589–595 (2010).

46. Li, H. *et al.* The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–

2079 (2009).

47. Robinson, J. T. *et al.* Integrative Genomics Viewer. *Nat. Biotechnol.* **29**, 24–26 (2011).

48. Thorvaldsdóttir, H., Robinson, J. T. & Mesirov, J. P. Integrative Genomics Viewer (IGV):

high-performance genomics data visualization and exploration. *Brief. Bioinform.* **14**, 178–

192 (2013).

49. McCarthy, D. J., Chen, Y. & Smyth, G. K. Differential expression analysis of multifactor

RNA-Seq experiments with respect to biological variation. *Nucleic Acids Res.* **40**, 4288–

4297 (2012).

50. Robinson, M. D., McCarthy, D. J. & Smyth, G. K. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. *Bioinformatics* **26**, 139–140

(2010).

51. Alvarez, L., Hernandez, S. B., de Pedro, M. A. & Cava, F. Ultra-Sensitive, High-Resolution

Liquid Chromatography Methods for the High-Throughput Quantitative Analysis of

Bacterial Cell Wall Chemistry and Structure. in *Bacterial Cell Wall Homeostasis: Methods*

and Protocols (ed. Hong, H.-J.) 11–27 (Springer New York, 2016). doi:10.1007/978-1-4939-

3676-2_2

52. Desmarais, S. M., Pedro, M. A. D., Cava, F. & Huang, K. C. Peptidoglycan at its peaks: how

chromatographic analyses can reveal bacterial cell wall structure and assembly. *Mol.*

Microbiol. **89**, 1–13 (2013).

Acknowledgements

We thank members of the Goley lab for helpful discussions and input. We thank Kousik

Sundararajan for identifying the initial mutant of CdnL that suppresses the FtsZΔCTL

phenotype. We would like to thank Julie Theriot, Lucy Shapiro, Christine Jacobs-Wagner, Regis

Hallez, and Zemer Gitai for providing materials and strains used in this project. We thank the

Johns Hopkins Genetic Resources Core Facility for RNA-Seq services and analysis and Zach

Pincus for guidance in using CellTool. This work was funded by the NIH through

R01GM108640 (to EDG), R01GM111706 (to PC), T32GM08515 (training grant support of RZ),

and T32GM007445 (training grant support of AKD) and Johns Hopkins Discovery Fund grant to

EDG. JC is funded by Swiss National Science Foundation (SNSF) grants 31003A_140758 and

31003A_173075. Research in the Cava lab is supported by MIMS, the Knut and Alice

Wallenberg Foundation (KAW), the Swedish Research Council, and the Kempe Foundation.

Author contributions

SAW and EDG conceived the study. SAW, LA, RZ, DG, JC, PC, FC, and EDG designed the

experiments. SAW performed the majority of experiments and analysis, except as noted. LA

performed muropeptide analysis and lipid II quantification. SAW, AKD, RZ, and PC performed

and/or analyzed Tn-Seq experiments. AB wrote custom software to analyze MreB distribution.

DG performed and analyzed microarray experiments. EDG purified CdnL, generated strains, and

performed CellTool analysis of shape. JC, PC, FC, and EDG supervised the research. SAW and

EDG wrote the manuscript with support and editing from all authors.

Figure Legends

Figure 1: Cells lacking CdnL have pleiotropic morphological and growth defects. a. Growth

of WT (EG865) and Δ*cdnL* (EG1447) cells in the indicated media. **b.** Phase contrast images of

WT and Δ*cdnL* cells grown in the indicated media. Arrows indicate ghost cells. Bar = 2 μm. **c.**

PCA of cell shape for WT and Δ*cdnL* shown in **b.** Scatter plots of shape modes 1, 2, and 3

approximately correspond to length, curvature and width. Contours on the left of scatter plots

indicate mean cell shape and standard deviations (s.d) from the mean. **d.** Mean shapes of WT and

Δ*cdnL* cells grown in the indicated media. **e.** Membrane staining of EG865 (WT) and EG1447

(Δ*cdnL*) cells with FM4-64 after growth in HIGG media with and without phosphate. **f.** Stalk

lengths measured using ImageJ's[42] line tool. Mean and SEM included for plots in **c.** and **f.**

Statistical analysis in **c.** and **f.** performed using one-way ANOVA with Bonferroni's Multiple

Comparison Test, n = 500 for each in **c.** n = 150, 127, 168, 155, 162, 172 from left to right in **e.**

*** = P < 0.001, * = P < 0.05, ns = not significant.

Figure 2: CdnL impacts cell wall precursor synthesis and cell wall metabolism. a. Heatmap

showing changes of the indicated transcripts involved in lipid II synthesis and cell wall

remodeling in WT (EG865) and Δ*cdnL* (EG1415) using RNA-Seq. **b. & c.** Relative amount of

lipid II and peptidoglycan in WT (EG865) and Δ*cdnL* (EG1447) cells. **d. & e.** Indicated

muropeptide species from WT (EG865) and Δ*cdnL* (EG1447). X-axis shown in **e.** are (1-6

anhydro) N-acetyl muramic acid (anhydro), Gly containing muropeptides (Gly),and

pentapeptides (penta). Values in **b., c., d. & e.** are from three independent cultures and mean and

SEM are indicated. Statistical analysis performed using unpaired t test. *** = P < 0.001, ** = P <

0.01, * = P < 0.05.

Figure 3: Mutations in cell wall recycling and capsule biosynthetic genes are synthetic

lethal with Δ*cdnL*. a. Volcano plot representation of Tn-Seq analysis. Negative $\log_{10}$ of the false

discovery rate (-log(FDR)) is plotted against $\log_2$ of the fold change in number of unique

transposon insertions in each gene in WT (EG865) vs Δ*cdnL* (EG1447). Genes that become

essential in the Δ*cdnL* background and are discussed in this study are color-coded. **b.** Transposon

insertion profile for selected genes from **a. c. & d.** Growth curves and phase contrast images for

WT (EG865), Δ*cdnL* (EG1447), Δ*CCNA_00164* (EG2604) and Δ*cdnL* Δ*CCNA_00164*

(EG2804). Bar = 2 μm.

Figure 4: MreB localization is aberrant in Δ*cdnL* cells. a., b. & c. Merged phase contrast

(blue) and fluorescence (yellow) images of WT (EG1795) and Δ*cdnL* cells (EG2465) expressing

venus-mreB for 2 hrs from the *xylX* promoter in the indicated media. Bar = 2 μm. **d., e. & f.**

Demographs of images shown in **a., b. &c.** n = 850 cells for each. **g., h. & i.** Scatter plots of full

width at half maximum (FWHM) values of midcell-localized MreB. Cell length range used for

FWHM calculation of midcell-localized MreB was determined from demographs shown in **d.-f.**.

For PYE, WT and Δ*cdnL*, cells 2 – 3 μm long have MreB localized at midcell and were selected

for FWHM calculations. For M2GG, WT and Δ*cdnL* cells between lengths of 3 – 3.75 μm and 2

– 3 μm, respectively, were selected for FWHM calculations. For HIGG, WT and Δ*cdnL* cells

between lengths of 3 – 3.5 μm and 2 – 3 μm, respectively, were selected for FWHM

calculations. Statistical analysis performed using unpaired t test. *** = P < 0.001. n = 140 each

for PYE and HIGG and n = 383 each for M2GG.

Figure 5: Hypercurvature of Δ*cdnL* cells is dependent on crescentin and may involve CTP

synthase. a. Phase contrast images of WT (EG865), Δ*creS* (EG1903), Δ*cdnL::gentR* (EG1898)

and Δ*creS* Δ*cdnL::gentR* (EG1912). **b.** Scatter plot depicting shape mode 2 (~curvature) from

PCA of cell shape for cells shown in **a**. Mean and SEM values are indicated. n = 447, 615, 734,

574 from left to right. Contours on the left of scatter plots indicate mean cell shape and standard

deviations (s.d) from the mean. Statistical analysis performed using one-way ANOVA with

Bonferroni's Multiple Comparison Test. *** = P < 0.001, ns = not significant. **c.** Heatmap

showing changes of indicated transcripts involved in pyrimidine biosynthesis in WT (EG865)

and Δ*cdnL* (EG1415) using RNA-Seq. **d.** Demographs of images from Supplementary Figure 4

of WT (EG1902) and Δ*cdnL* (EG1958) cells expressing *mCherry – ctpS* from the *xylX* locus

grown in indicated media. n = 400 cells for each.

Supplementary Figures Legends

Supplementary figure 1: RNA-seq and microarray analysis using strains EG1415 and

JC784 have significant overlap. a. Immunoblot against whole cell lysates of the indicated

strains probed with CdnL antisera. **b.** Phase contrast images and **c.** growth curves of WT

(EG865), Δ*cdnL* (EG1447) and Δ*cdnL; Δ50kb* (EG1415). Bar = 2 μm. **d.** Pie chart comparing

RNA-Seq data using EG1415 and microarray data using JC784. A 2-fold change in transcript

level was used as the cut-off for up- or downregulation. **e.** Pearson's correlation between

microarray (y – axis) and RNA-Seq (x – axis) data. Pearson r = 0.6331; significant correlation

with alpha = 0.05. **f & g.** Zoomed in scatter plots depicting correlation between microarray (y –

axis) and RNA-Seq (x – axis) for ≥2-fold upregulated genes **(f)** and ≥2-fold down regulated **(g)**

genes in both data sets.

Supplementary figure 2: Muropeptide composition is altered in Δ*cdnL* cells. a.

Representative chromatograms of the muramidase-digested sacculi of WT and Δ*cdnL* cells. **b.**

Relative molar abundance (%) of monomers, dimers, trimers, percentage of crosslinkage

(proportion of crosslinked peptide side chains), muropeptides with a residue of (1-6 anhydro) N-

acetyl muramic acid (Anhydro), Gly containing muropeptides (Gly muropep.), and

pentapeptides. **c.** Muropeptide relative molar abundance (%). GlcNAc: *N*-Acetyl glucosamine.

MurNAc: *N*-Acetyl muramic acid. Ala: Alanine. Glu: Glutamic acid. mDAP: meso-

diaminopimelic acid. Gly: Glycine. Statistical analysis performed using t-test analysis. * = P <

0.05 and > 10% variation compared to WT.

Supplementary figure 3: *mreB* mutants are synthetic sick with Δ*cdnL*. a. Phase contrast

images of WT and Δ*cdnL* cells expressing indicated *mreB* mutants at the *mreB* locus. Bar = 2

μm. **b.** Growth curves of strains shown in **(a)** grown in PYE. **c.** Summary of the localization of

MreB mutants from Dye et al, 2011.[27] **d.** Localization of indicated MreB mutants by expressing

mutant version of *venus-mreB* from *xylX* locus. **e.** Demographs of MreB mutants shown in **(d).**

Supplementary figure 4: Depletion of lipid II by treating cells with fosfomycin causes an

increase in cell width and FWHM of MreB localization. a. Merged images of cells expressing

venus-mreB from the *xylX* promoter and treated with fosfomycin for indicated amount of time

before imaging. Blue: phase contrast, yellow: Venus-MreB. Bar = 2 μm. **b.** Maximum cell width

of cells shown in **(a).** n = 500 each, *** = $P < 0.001$, ns = not significant using one-way ANOVA

with Bonferroni's Multiple Comparison Test. **c.** Full width at half maximum (FWHM) values of

Venus-MreB in cells between 2.5 – 3 μm long and which have MreB localized at midcell. n =

245, *** = $P < 0.001$, ns = not significant using one-way ANOVA with Bonferroni's Multiple

Comparison Test.

Supplementary figure 5. CTP synthase localization is weak in Δ*cdnL* cells compared to WT.

Phase contrast, fluorescent and merged images of WT (EG1902) and Δ*cdnL* (EG1958) cells

expressing *mCherry – ctpS* from the *xylX* locus grown in indicated media. Bar = 2 μm.

Supplementary table 1: RNA-Seq and microarray data for EG1415 and JC784, respectively.

Gene (CCNA), annotation, fold change in Δ*cdnL* over wild-type, $\log_2$ of fold change, and

statistical significance (1 = significant, 0 = not significant) are presented for each dataset. The 50

kb deletion in EG1415 is highlighted.

Supplementary table 2: DAVID[15] functional analysis of the set of genes great than or equal to

2-fold up-regulated or 2-fold down-regulated in both RNAseq and microarray datasets from

supplementary table 1. Functional annotation clusters identified, their enrichment scores, cluster

names, and the gene numbers for that cluster are indicated.

Supplementary table 3: Tn-Seq data for EG1447 (ΔcdnL) and EG865 (WT). The number of

unique transposon insertions in each gene in wild-type compared to Δ*cdnL* was used to

determine the $\log_2$ fold-change in number of unique transposon insertions (logFC), log counts

per million reads (logCPM), PValue, false-discovery rate (FDR), and negative $\log_{10}$ of the FDR

for each gene in WT versus Δ*cdnL*.

Supplementary table 4: Strains and plasmids used in this study.

Figure 1: Cells lacking CdnL have pleiotropic morphological and growth defects. a. Growth of WT (EG865) and ΔcdnL (EG1447) cells in the indicated media. b. Phase contrast images of WT and ΔcdnL cells grown in the indicated media. Arrows indicate ghost cells. Bar = 2 μm. c. PCA of cell shape for WT and ΔcdnL shown in b. Scatter plots of shape modes 1, 2, and 3 approximately correspond to length, curvature and width. Contours on the left of scatter plots indicate mean cell shape and standard deviations (s.d) from the mean. d. Mean shapes of WT and ΔcdnL cells grown in the indicated media. e. Membrane staining of EG865 (WT) and EG1447 (ΔcdnL) cells with FM4-64 after growth in HIGG media with and without phosphate. f. Stalk lengths measured using ImageJ's42 line tool. Mean and SEM included for plots in c. and f. Statistical analysis in c. and f. performed using one-way ANOVA with Bonferroni's Multiple Comparison Test, n = 500 for each in c. n = 150, 127, 168, 155, 162, 172 from left to right in e. *** = P < 0.001, * = P < 0.05, ns = not significant.

Figure 2: CdnL impacts cell wall precursor synthesis and cell wall metabolism. a. Heatmap showing changes of the indicated transcripts involved in lipid II synthesis and cell wall remodeling in WT (EG865) and ∆cdnL (EG1415) using RNA-Seq. b. & c. Relative amount of lipid II and peptidoglycan in WT (EG865) and ∆cdnL (EG1447) cells. d. & e. Indicated muropeptide species from WT (EG865) and ∆cdnL (EG1447). X-axis shown in e. are (1-6 anhydro) N-acetyl muramic acid (anhydro), Gly containing muropeptides (Gly),and pentapeptides (penta). Values in b., c., d. & e. are from three independent cultures and mean and SEM are indicated. Statistical analysis performed using unpaired t test. *** = P < 0.001, ** = P < 0.01, * = P < 0.05.

Figure 3: Mutations in cell wall recycling and capsule biosynthetic genes are synthetic lethal with ΔcdnL. a. Volcano plot representation of Tn-Seq analysis. Negative log10 of the false discovery rate (-log(FDR)) is plotted against log2 of the fold change in number of unique transposon insertions in each gene in WT (EG865) vs ΔcdnL (EG1447). Genes that become essential in the ΔcdnL background and are discussed in this study are color-coded. b. Transposon insertion profile for selected genes from a. c. & d. Growth curves and phase contrast images for WT (EG865), ΔcdnL (EG1447), ΔCCNA_00164 (EG2604) and ΔcdnL ΔCCNA_00164 (EG2804). Bar = 2 μm.

Figure 4: MreB localization is aberrant in ΔcdnL cells. a., b. & c. Merged phase contrast (blue) and fluorescence (yellow) images of WT (EG1795) and ΔcdnL cells (EG2465) expressing venus-mreB for 2 hrs from the xylX promoter in the indicated media. Bar = 2 µm. d., e. & f. Demographs of images shown in a., b. &c. n = 850 cells for each. g., h. & i. Scatter plots of full width at half maximum (FWHM) values of midcell-localized MreB. Cell length range used for FWHM calculation of midcell-localized MreB was determined from demographs shown in d.-f.. For PYE, WT and ΔcdnL, cells 2 – 3 µm long have MreB localized at midcell and were selected for FWHM calculations. For M2GG, WT and ΔcdnL cells between lengths of 3 – 3.75 µm and 2 – 3 µm, respectively, were selected for FWHM calculations. For HIGG, WT and ΔcdnL cells between lengths of 3 – 3.5 µm and 2 – 3 µm, respectively, were selected for FWHM calculations. Statistical analysis performed using unpaired t test. *** = P < 0.001. n = 140 each for PYE and HIGG and n = 383 each for M2GG.

Figure 5: Hypercurvature of ΔcdnL cells is dependent on crescentin and may involve CTP synthase. a. Phase contrast images of WT (EG865), ΔcreS (EG1903), ΔcdnL::gentR (EG1898) and ΔcreS ΔcdnL::gentR (EG1912). b. Scatter plot depicting shape mode 2 (~curvature) from PCA of cell shape for cells shown in a. Mean and SEM values are indicated. n = 447, 615, 734, 574 from left to right. Contours on the left of scatter plots indicate mean cell shape and standard deviations (s.d) from the mean. Statistical analysis performed using one-way ANOVA with Bonferroni's Multiple Comparison Test. *** = P < 0.001, ns = not significant. c. Heatmap showing changes of indicated transcripts involved in pyrimidine biosynthesis in WT (EG865) and ΔcdnL (EG1415) using RNA-Seq. d. Demographs of images from Supplementary Figure 4 of WT (EG1902) and ΔcdnL (EG1958) cells expressing mCherry – ctpS from the xylX locus grown in indicated media. n = 400 cells for each.

Supplementary figure 1: RNA-seq and microarray analysis using strains EG1415 and JC784 have significant overlap. a. Immunoblot against whole cell lysates of the indicated strains probed with CdnL antisera. b. Phase contrast images and c. growth curves of WT (EG865), ΔcdnL (EG1447) and ΔcdnL; Δ50kb (EG1415). Bar = 2 μm. d. Pie chart comparing RNA-Seq data using EG1415 and microarray data using JC784. A 2-fold change in transcript level was used as the cut-off for up- or downregulation. e. Pearson's correlation between microarray (y – axis) and RNA-Seq (x – axis) data. Pearson r = 0.6331; significant correlation with alpha = 0.05. f & g. Zoomed in scatter plots depicting correlation between microarray (y – axis) and RNA-Seq (x – axis) for ≥2-fold upregulated genes (f) and ≥2-fold down regulated (g) genes in both data sets.

a.

b.

Parameter	WT	ΔcdnL
Monomers	60.66	60.23
Dimers	26.23	32.22*
Trimers	13.10	7.55*
Crosslink	52.44	47.33
Anhydro	17.51	13.74*
Gly muropep.	21.66	43.54*
Pentapeptides	49.39	59.69*

c.

Peak	Muropeptide	Structure	WT	ΔcdnL
1	M3	GlcNAc-MurNAc-L-Ala-D-Glu-mDAP	0.60	1.41 *
2	M4^G	GlcNAc-MurNAc-L-Ala-D-Glu-mDAP-Gly	0.98	1.89 *
3	M5^G	GlcNAc-MurNAc-L-Ala-D-Glu-mDAP-D-Ala-Gly	10.69	20.39 *
4	M4	GlcNAc-MurNAc-L-Ala-D-Glu-mDAP-D-Ala	27.40	16.88 *
5	M2	GlcNAc-MurNAc-L-Ala-D-Glu	1.82	4.37 *
6	M44^G	M4-M4^G (DD-crosslink, dissaccharide lost)	0.00	1.54 *
7	M5	GlcNAc-MurNAc-L-Ala-D-Glu-mDAP-D-Ala-D-Ala	17.22	9.94 *
8	D44^G	M4-M4^G (DD-crosslink)	0.24	0.64 *
9	D43	M4-M3 (DD-crosslink)	0.16	0.75 *
10	D45^G	M4-M5^G (DD-crosslink)	5.18	12.50 *
11	D44	M4-M4 (DD-crosslink)	8.80	7.07 *
12	M5$^{G\ Anh}$	GlcNAc-(1-6anhydro)MurNAc-L-Ala-D-Glu-mDAP-D-Ala-Gly	0.75	2.37 *
13	D45	M4-M5 (DD-crosslink)	6.59	5.88
14	M4^{Anh}	GlcNAc-(1-6anhydro)MurNAc-L-Ala-D-Glu-mDAP-D-Ala	0.87	0.93
15	T445^G	M4-M4-M5^G (DD-crosslink)	0.63	1.35 *
16	M5^{Anh}	GlcNAc-(1-6anhydro)MurNAc-L-Ala-D-Glu-mDAP-D-Ala-D-Ala	0.33	0.51 *
17	T444	M4-M4-M4 (DD-crosslink)	1.27	0.98*
18	T445	M4-M4-M5 (DD-crosslink)	0.89	0.68 *
19	D45$^{G\ Anh}$	M4-M5$^{G\ Anh}$ (DD-crosslink)	0.92	1.57 *
20	D44^{Anh}	M4-M4^{Anh} (DD-crosslink)	2.41	1.57 *
21	D45^{Anh}	M4-M5^{Anh} (DD-crosslink)	1.93	2.25 *
22	T445$^{G\ Anh}$	M4-M4-M5$^{G\ Anh}$ (DD-crosslink)	1.77	0.73 *
23	T444^{Anh}	M4-M4-M4^{Anh} (DD-crosslink)	3.04	1.57 *
24	T445^{Anh}	M4-M4-M5^{Anh} (DD-crosslink)	1.99	0.76 *
25	T445$^{G\ Anh,\ Anh}$	M4-M4^{Anh}-M5$^{G\ Anh}$ (DD-crosslink)	0.50	0.57
26	T444$^{Anh,\ Anh}$	M4-M4^{Anh}-M4^{Anh} (DD-crosslink)	3.02	0.72 *
27	T445$^{Anh,\ Anh}$	M4-M4^{Anh}-M5^{Anh} (DD-crosslink)	0.00	0.20 *

Supplementary figure 2: Muropeptide composition is altered in ΔcdnL cells. a. Representative chromatograms of the muramidase-digested sacculi of WT and ΔcdnL cells. b. Relative molar abundance (%) of monomers, dimers, trimers, percentage of crosslinkage (proportion of cross-linked peptide side chains), muropeptides with a residue of (1-6 anhydro) N-acetyl muramic acid (Anhydro), Gly containing muropeptides (Gly muropep.), and pentapeptides. c. Muropeptide relative molar abundance (%). GlcNAc: N-Acetyl glucosamine. MurNAc: N-Acetyl muramic acid. Ala: Alanine. Glu: Glutamic acid. mDAP: meso-diaminopimelic acid. Gly: Glycine. Statistical analysis performed using t-test analysis. * = P < 0.05 and > 10% variation compared to WT.

Supplementary figure 3: mreB mutants are synthetic sick with ΔcdnL. a. Phase contrast images of WT and ΔcdnL cells expressing indicated mreB mutants at the mreB locus. Bar = 2 µm. b. Growth curves of strains shown in (a) grown in PYE. c. Summary of the localization of MreB mutants from Dye et al, 2011.27 d. Localization of indicated MreB mutants by expressing mutant version of venus-mreB from xylX locus. e. Demographs of MreB mutants shown in (d).

Supplementary figure 4: Depletion of lipid II by treating cells with fosfomycin causes an increase in cell width and FWHM of MreB localization. a. Merged images of cells expressing venus-mreB from the xylX promoter and treated with fosfomycin for indicated amount of time before imaging. Blue: phase contrast, yellow: Venus-MreB. Bar = 2 µm. b. Maximum cell width of cells shown in (a). n = 500 each, *** = P < 0.001, ns = not significant using one-way ANOVA with Bonferroni's Multiple Comparison Test. c. Full width at half maximum (FWHM) values of Venus-MreB in cells between 2.5 – 3 µm long and which have MreB localized at mid-cell. n = 245, *** = P < 0.001, ns = not significant using one-way ANOVA with Bonferroni's Multiple Comparison Test.

Supplementary figure 5. CTP synthase localization is weak in ∆cdnL cells compared to WT. Phase contrast, fluorescent and merged images of WT (EG1902) and ∆cdnL (EG1958) cells expressing mCherry – ctpS from the xylX locus grown in indicated media. Bar = 2 µm.