Glutarate metabolism in *Pseudomonas putida* is regulated by two distinct glutarate sensing

transcription factors

Mitchell G. Thompson[1,2,3], Pablo Cruz-Morales[1,2,4], Rohith N. Krishna[1,2,5],Jacquelyn M. Blake-

Hedges[1,2,5], Matthew R. Incha[1,2,3], Jay D. Keasling[1,2,6,7,8]

[1]Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.

[2]Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley,

CA 94720, USA.

[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720,

USA

[4]Centro de Biotecnologia FEMSA, Instituto Tecnologico y de Estudios superiores de Monterrey,

Mexico

[5]Department of Chemistry, University of California, Berkeley, CA 94720, USA

[6]Joint Program in Bioengineering, University of California, Berkeley/San Francisco, CA 94720,

USA

[7]Department of Chemical and Biomolecular Engineering, University of California, Berkeley, CA

94720, USA

[8]The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark,

Denmark

ABSTRACT

Transcription factor based biosensors can be leveraged to screen thousands of genetics designs for optimal production in engineered microbes. In this study we characterize two glutarate sensing transcription factors (CsiR and GcdR) from *Pseudomonas putida*. The genomic contexts of CsiR homologs were analyzed and DNA binding sites were bioinformatically predicted. Both CsiR and GcdR were purified and shown to bind upstream of their coding sequencing *in vitro*. CsiR was shown to dissociate from DNA *in vitro* when exogenous glutarate was added confirming it acts as a genetic repressor. Both transcription factors were then engineered into plasmid based biosensors and their respective sensing performance features calculated. Both sensor plasmids were then reintroduced into *P. putida* and were evaluated for their ability to sense flux through glutarate when grown on various lysine metabolites as sole carbon sources. These findings provide information describing glutarate flux in *P. putida*, and potentially useful tools for future metabolic engineering and synthetic biology efforts.

INTRODUCTION

A rate limiting step in the design, build, test cycle is often the test phase, wherein hundreds or thousands of genetic designs need to be evaluated for their productivity [1,2]. Though recent advances in analytical chemistry have dramatically increased sample throughput [3], transcription factor based biosensors still offer multiple advantages over traditional chromatographic and mass-spectrometry based detection methods. One of the most attractive benefits is the ability to rapidly screen constructs for the production of the target compounds via either plate based, or flow-cytometry based assays [1–3]. Additionally, biosensors may offer unmatched sensitivity towards specific ligands, with some sensors having picomolar affinity [3].

The evolution of diverse microbial metabolism has provided researchers with a wide array of

ligands capable of being sensed, ranging from complex natural products [4,5], to small central

metabolites [1,6].

Work in our lab has developed biosensors to detect multiple lactams [6], which can be

utilized to synthesize economically important polymers such as nylon [7]. These polymers also

require diacids, such as adipic acid, for their polymerization. The L-lysine metabolism of

Pseudomonas putida has been leveraged to produce the omega-amino acid 5-aminovaleric acid

(5AVA), as well as the diacid glutarate [7,8]. Recently it has been found that glutarate

metabolism in *P. putida* can be degraded either by a ketogenic route to acetyl-coA or through a

glucogenic route to succinate [8] (Figure 1). Work in our lab has also shown this metabolism,

and demonstrated that both glutarate catabolic pathways are highly upregulated in the presence

of glutarate [9]. While work in *Pseudomonas aeruginosa* has characterized a homolog of the

regulator of the ketogenic pathway (GcdR) [10], and work in *Escherichia coli* has preliminarily

characterized the glucogenic pathway regulator (CsiR), no work has directly investigated the

regulation of the *P. putida* L-lysine catabolic pathways.

In this work we sought to better characterize the two putative local regulators of glutarate

catabolism in *P. putida*, *csiR* and *gcdR*. First, we compared the genomic context of *csiR*

homolongs across bacteria, to bioinformatically predict a conserved DNA binding site. We then

biochemically, and genetically characterized both regulators. Finally we introduced RFP

transcriptional fusions of the promoter for both catabolic pathways into *P. putida* and evaluated

their induction upon the addition of various lysine metabolites.

RESULTS

<u>Genomic Contexts of CsiR and GcdR Homologs and Prediction of *P. putida* binding sites</u>

Work in *Pseudomonas aeruginosa* has characterized the GcdR regulation of ketogenic

glutarate metabolism, and shown that the binding site is conserved across multiple bacterial

species [10]. While binding sites for CsiR in *E. coli* have been identified, whether there is a

conserved binding site for homologs across bacterial species has yet to be investigated [11]. In

order to identify conserved binding sites of CsiR homologs, 19 selected genomes that contained

neighboring CsiD and CsiR homologs had their syntenic genomic contexts compared. Genes

encoding *csiR* were found in two distinct genomic contexts, either transcribed divergently from

csiD as found in *P. putida*, or transcribed as the last gene in the *csi* operon as in *E. coli* (Figure

2).   The genomic regions upstream of the *csiD* homolog were extracted and MEME was used to

identify a conserved CsiR binding motif. A consensus A(A/G)AAATCTAGA(C/T)ATTTT motif

was identified upstream of each *csiD* homolog. Previously, footprinting assays in *E. coli* revealed

two primary and two secondary bindings sites with the sequences $TTGTN_5TTTT$ and

$ATGTN_5TTTT$ respectively [11]. Our consensus motif agrees closely with the secondary binding

site identified, and highly suggests that the binding site of CsiR is conserved across the bacteria

where homologs are present.

Biochemical characterization of CsiR and GcdR

To determine if the *P. putida* CsiR also acts as a regulator and to identify its putative

binding sites, we biochemically characterized this protein using electrophoretic mobility shift

assays (EMSAs). The CsiR protein was purified and incubated with DNA probes consisting of

the intergenic region between *csiR* and *csiD*. The assay showed multiple binding sites in this

intergenic region, as four distinct bands appeared. These results appear to confirm the finding in

E. coli where four binding sites of the *E. coli* CsiR homolog were observed [11]. CsiR

demonstrated a high affinity for the DNA probe, with a calculated K_d of approximately 30 nM

(Figure 3a), which is similar to the CsiR/DNA K_d of the *E. coli* CsiR homolog which was

determined to be 10nM [11]. GcdR also bound to its cognate probe, with a single distinct shift,

with an estimated K_d of approximately 62.5 nM (Figure 3a) [10]. As CsiR is a GntR family

transcriptional regulator, many of which act as repressors [12], we evaluated whether glutarate

would decrease the DNA binding affinity of CsiR. EMSA assays were repeated in the presence

of increasing glutarate concentrations. Analysis by gel electrophoresis revealed increasing

quantities of free probe as glutarate concentrations increased (Figure 3b). These results confirm

that the *P. putida* CsiR is a glutarate-responsive repressor of the *csiDlhgO* operon.

Development of two glutarate biosensor vectors

In order to evaluate both CsiR and GcdR as biosensors we cloned both the regulator and

intergenic region of both the ketogenic and glucogenic glutarate catabolism pathways upstream

of RFP on a broad host range plasmid (Figure 4a). *E. coli* harboring either plasmid were grown

for 24 hours in LB supplemented with concentrations of glutarate ranging from 50 mM to 6.1

uM, after which both OD600 and RFP fluorescence were measured. At concentrations above

6.25 mM glutarate proved to be toxic to cells, and RFP measurements from these concentrations

were not used to calculate biosensor performance features (Figure 4b). The CsiR reporter vector

had significantly higher background than the GcdR vector, indicating leakier expression (Figure

4b). Normalized RFP expression data was then fit to the Hill equation to derive biosensor

performance (Figure 4c). Both sensors showed similar linear ranges (0.05 - 3.13 mM for CsiR,

and 0.05 - 6.25 mM for GcdR), however the K_m for CsiR was lower than GcdR (0.65 mM versus

2.24 mM). The GcdR reporter displayed greater sensitivity than the CsiR vector (Hill coefficient

of 1.09 versus 0.75), in addition to a greater maximum expression of RFP (65000 versus 42000).

Responsiveness of Glutarate Biosensors to Lysine metabolites in *P. putida*

To evaluate the ability of both reporter vectors to monitor glutarate flux *in vivo*, both

plasmids were introduced into wild-type *P. putida* KT2440 and grown on L-lysine and D-lysine

metabolites as sole carbon sources. Both strains were grown on 10 mM Glucose, L-lysine, D-

lysine, 5-aminovalerate, or 2-aminoadipatefor 48 hours with OD_{600} and RFP fluorescence being

measured continuously (Figures 5a and 5b). Neither reporter was induced when the bacterium

was grown on glucose (Figure 5b). The GcdR reporter was strongly induced by 5-aminovaleate,

and to a less extent 2-aminoadipate, D-lysine, and L-lysine (Figure 5b). Conversely, the strain

harboring the CsiR reporter plasmid only displayed r induction of RFP above background only

when grown on 5-aminovalerate. While 5-aminovalerate was able to induce RFP induction in the

strain harboring the CsiR reporter, induction was extremely limited compared to induction of

RFP from the GcdR reporter vector (Figure 5b).

DISCUSSION

Recent advances in high-throughput functional genomics have allowed researchers to

rapidly identify novel metabolic pathways, and in turn infer the function of novel transcription

factors [13,14]. Rigorous characterization of these regulators is a critical step to developing

novel parts for synthetic biology, as well as useful tools for metabolic engineering. The

discovery of two distinct pathways for glutarate catabolism within *P. putida*, regulated by

independent transcription factors, presents an interesting opportunity to compare and contrast the

relative sensing properties of each system.

The *P. aeruginosa* GcdR was demonstrated to be a positive regulator of the *gcdHG*

operon, with a binding site that was conserved in *P. putida* KT2440 among other *Pseudomonas*

strains [10]. While other strains of *P. aeruginosa* have *csiD* homologs, the type strain PA01 does

not. However, *csiDlghO* pathways have been rigorously characterized in both *P. putida* and *E.*

coli, though only in *E. coli* has the CsiR homolog been characterized [8,9,11]. Our bioinformatic

approach to predict the conserved binding site of CsiR largely agrees with the empirically

determined binding site identified in *E. coli*, and our EMSA results also confirm the existence of

four distinct CsiR binding sites. The dissociation of CsiR from the DNA probe with the addition

of glutarate *in vitro* verifies that CsiR is a negative regulator of CsiD, which is expected as many

GntR-type regulator tend to act as repressors. The existence of a primary and secondary series of

binding sites suggests a canonical autoregulatory negative feedback loop, found in an estimated

~60-70% of bacterial transcription factors [15]. Further work will need to be done to characterize

any such autoregulation.

Plasmid-based biosensors of both the CsiR and GcdR systems displayed low uM limits of

detection in *E. coli*. While the CsiR biosensor displayed greater sensitivity to glutarate at lower

inducer concentrations, its maximum output was considerably less than the GcdR system and

saturated at low mM concentrations of glutarate. Furthermore the CsiR system showed

substantially higher background RFP than the GcdR based system. While the CsiR system may

be useful for detecting low concentrations of glutarate, further engineering will need to be done

to increase its utility. GcdR displayed a larger linear range of detection, and a greater maximal

output that the CsiR system, in addition to a relatively low level of background expression.

These properties may make the GcdR system more attractive as a glutarate biosensor.

Furthermore, given that glutarate is relatively inexpensive, the GcdR system may make an

excellent inducible expression system. Further work will need to be conducted to evaluate the

utility of the GcdR expression system in other bacteria.

In addition to their utility as biosensors for metabolic engineering, these sensors are a

valuable tool in studying the carbon flux in the native host *P. putida*. Lysine metabolism in *P.*

putida is isomer specific, with each isomer being degraded by a separate catabolic pathway [9].

While cross-feeding between the pathways has been proposed previously [16], recent work by

our group has proposed a molecular mechanism for metabolite exchange between the D- and L-

catabolic pathways [9]. The exchange relies on the 2-oxoacid promiscuity of the non-heme Fe(II)

oxidase CsiD which normally catalyzes the hydroxylation of glutarate using 2-ketoglutarate as a

cosubstrate to yield 2-hydroxyglutarate and succinate. CsiD can also use 2-oxoadipate, a D-

lysine catabolic intermediate, as a 2-oxoacid cosubstrate to yield 2-hydroxyglutarate and

glutarate as products. The glutarate from this reaction could then proceed down the L-lysine

catabolic pathway. When 2-aminoadipate was fed into *P. putida* harboring the GcdR reporter,

fluorescence was observed in stationary phase. As 2-aminoadipate immediately precedes 2-

oxoadipate in the D-lysine catabolic pathway, these results support the hypothesis that CsiD

could act as a bridge between the two catabolic pathways.

There has been substantial interest in developing microbes to produce glutarate, with

strains of *E. coli*, *P. putida*, and *Corynebacterium glutamicum* all engineered to produce high

titers [8,17,18]. Further engineering of the GcdR system may be able to extend the linear range

to higher concentrations furthering its utility as a tool to achieve even higher titers of glutarate.

Though glutarate is a valuable commodity chemical, the C6 diacid adipate is used in much

greater quantities primarily as a monomer used to make nylons [19]. Recent work has readily

demonstrated the means to evolve transcription factors to sense non-native ligands [4]. Such

methods could be applied to either CsiR or GcdR in order to expand their utility in sensing other

industrially important diacids.

METHODS

<u>Media, chemicals, and culture conditions</u>

General *E. coli* cultures were grown in Luria-Bertani (LB) Miller medium (BD

Biosciences, USA) at 37 °C while *P. putida* was grown at 30 °C. When indicated, *P. putida* and

E. coli were grown on modified MOPS minimal medium [20]. Cultures were supplemented with

kanamycin (50 mg/L, Sigma Aldrich, USA), gentamicin (30 mg/L, Fisher Scientific, USA), or

carbenicillin (100mg/L, Sigma Aldrich, USA), when indicated. All other compounds were

purchased through Sigma Aldrich (Sigma Aldrich, USA) .

Strains and plasmids

All bacterial strains and plasmids used in this work are listed in Table 1. All strains and

plasmids created in this work are available through the public instance of the JBEI registry.

(https://public-registry.jbei.org/folders/390). All plasmids were designed using Device Editor

and Vector Editor software, while all primers used for the construction of plasmids were

designed using j5 software [21–23]. Plasmids were assembled via Gibson Assembly using

standard protocols [24], or Golden Gate Assembly using standard protocols [25]. Plasmids were

routinely isolated using the Qiaprep Spin Miniprep kit (Qiagen, USA), and all primers were

purchased from Integrated DNA Technologies (IDT, Coralville, IA).

Expression and purification of proteins

Proteins were purified as described previously [26]. Briefly, a 500 mL cultures of *E. coli*

BL21 (DE3) harboring expression plasmids were grown on LB media at 37 °C to an OD of 0.6

then induced with 1mM Isopropyl β-D-1-thiogalactopyranoside Cells were allowed to express

for 18 hours at 30 °C before being harvested via centrifugation. Cell pellets were stored at -80

°C until purification. Cell pellets were then resuspended in lysis buffer (50 mM sodium

phosphate, 300 mM sodium chloride, 10 mM imidazole, 8% glycerol, pH 7.5) and sonicated to

lyse cells. Insolubles were pelleted via centrifugation (30 minutes at 40,000xg). The supernatant

was applied to a fritted column containing Ni-NTA resin (Qiagen, USA) which had been pre-

equilibrated with several column volumes of lysis buffer. The resin was washed with lysis buffer

containing 50 mM imidazole, then the protein was eluted using a step-wise gradient of lysis

buffer containing increasing imidazole concentrations (100 mM, 200 mM, and 400 mM).

Fractions were collected and analyzed via SDS-PAGE. Purified proteins were concentrated using

Spin-X UF 20 (10 kDa MWCO) spin concentrators (Corning, Inc.). Concentrated protein was

stored at 4 ºC until *in vitro* analysis.

<u>Plate based growth and fluorescence assays</u>

Growth studies of bacterial strains were conducted a microplate reader kinetic assays.

Overnight cultures were inoculated into 10 mL of LB medium from single colonies, and grown

at 30 °C. These cultures were then washed twice with MOPS minimal media without any added

carbon and diluted 1:100 into 500 uL of MOPS medium with 10 mM of a carbon source in 48-

well plates (Falcon, 353072). Plates were sealed with a gas-permeable microplate adhesive film

(VWR, USA), and then optical density and fluorescence were monitored for 48 hours in an

Biotek Synergy 4 plate reader (BioTek, USA) at 30 °C with fast continuous shaking. Optical

density was measured at 600 nm, while fluorescence was measured using an excitation

wavelength of 561 nm and an emission wavelength of 605 nm with a manually set gain of 100.

<u>Transcriptional fusion fluorescence assays</u>

To measure RFP production from *E. coli* reporter plasmids, fluorescence measurements

were obtained from single time points of cells grown in deep-well 96-well plates as described

previously [27]. Briefly Cells were grown in 500 μL of LB medium supplemented with

kanamycin and a range of glutarate concentrations from 5 mM to 0 mM. Plates were sealed with

AeraSeal film (Excel Scientific, AC1201-02) and grown for 22 hours at 30 °C on a 200 rpm

shaker rack. After incubation, 100 μL from each well was aliquoted into a black, clear-bottom

96-well plate and fluorescence was measured with an Infinite F200 Pro (Tecan Life Sciences,

San Jose, CA) plate reader. Optical density was measured at 600 nm, while fluorescence was

measured using an excitation wavelength of 575 nm and an emission wavelength of 620 nm with

a manually set gain of 85.

<u>Electrophoretic mobility shift assays</u>

237   Electrophoretic mobility shift assays were performed as previously described [5]. 6-

Carboxyfluorescein labelled PCR products to generate probes for CsiR were generated from the

intergenic region between PP_2908 and PP_2909 using primers csiRprobeFOR 5'-6-

FAM/AGTTCGATCTGCGTAAAG-3' and csiRprobeREV 5'-CCCGCTGAATGCTGAGTT-

3'), while probes for GcdR were generated from the intergenic region between PP_0157 and

PP_0158 with primers gcdRprobeFOR 5'-6-FAM/CGGGTCGATCCAGTTGAAA-3' and

gcdRprobeREV 5'-GCATGTACGTCAACCTCACT-3'. Primers were purchased from IDT

Technologies (IDT, Coralville, IA). PCR product was then purified with a QIAquick PCR

Purification Kit (Qiagen, USA), and the amount of DNA was quantified via a NanoDrop 2000C

(There Fisher Scientific). Binding reactions were conducted with 10 ng of labelled probe, which

was added to 10 mM Tris–HCl (pH 8.0), 25 mM KCl, 2.5 mM MgCl2, 1.0 mM DTT and 2 ug

salmon sperm DNA in 20 uL reactions. CsiR was added to reactions in concentrations ranging

from 250 nM to ~4 nM in addition to no CsiR controls, and then allowed to incubate at 22 °C for

20 minutes. Reactions were then loaded into 10% native polyacrylamide gels buffered with 0.5x

TBE. After electrophoresis gels were imaged on a Alpha Innotech MultiImage III (Alpha

Innotech). Pixel intensity to estimate K_d was done in Adobe Photoshop.

<u>Bionformatic Analysis and Analysis of Biosensor Parameters</u>

Biosensing parameters were calculated as previously described [1]. For the phylogenetic

reconstructions, the best amino acid substitution model was selected using ModelFinder as

implemented on IQ-tree [28] phylogenetic trees were constructed using IQ-tree, nodes were

supported with 10,000 bootstrap replicates. The final tree figures were edited using FigTree

v1.4.3 (http://tree.bio.ed.ac.uk/software/figtree/). Orthologous syntenic regions of CsiR were

identified with CORASON-BGC [29] and manually colored and annotated. DNA-binding sites

were predicted with MEME [30].

Acknowledgements

Python code to analyze biosensor performance characteristics was partially written by Dr. Zak

Costello. The authors would like to thank the UC Berkeley SMART program for providing

support for R.K to conduct summer research. This work was part of the DOE Joint BioEnergy

Institute (https://www.jbei.org) supported by the U.S. Department of Energy, Office of Science,

Office of Biological and Environmental Research. The views and opinions of the authors

expressed herein do not necessarily state or reflect those of the United States Government or any

agency thereof. Neither the United States Government nor any agency thereof, nor any of their

employees, makes any warranty, expressed or implied, or assumes any legal liability or

responsibility for the accuracy, completeness, or usefulness of any information, apparatus,

product, or process disclosed, or represents that its use would not infringe privately owned rights.

Contributions

Conceptualization, M.G.T.; Methodology, M.G.T., J.M.B, P.C.M.; Investigation, M.G.T., J.M.B,

R.N.K, P.C.M, M.R.I..; Writing – Original Draft, M.G.T, M.R.I..; Writing – Review and Editing,

All authors.; Resources and supervision, J.D.K.

Competing Interests

J.D.K. has financial interests in Amyris, Lygos, Constructive Biology, Demetrix, Napigen and

Maple Bio.

Figure Legends

Figure 1: Organization of the two routes of glutarate metabolism in *P. putida*.

Figure 2: Genomic contexts of CsiR homologs and predicted binding regions: Phylogeny of

assorted gammaproteobacteria and the location of their *csiD* operons. MEME analysis of the

intergenic regions upstream of the *csiD* operon resulted in the sequence motifs depicted below

the tree.

Figure 3: Biochemical characterization of CsiR and GcdR. (A) EMSA of either purified CsiR

(left) or GcdR (right) incubated with fluorescent probe of intergenic region between *csiR* and

csiD. Concentrations of CsiR protein range from 0nM to 250nM. (B) EMSA showing 100nM

CsiR and fluorescent probe incubated with glutarate from concentrations of 0 mM to 5 mM. Far

left lane shows free probe with no CsiR.

Figure 4: Developing glutarate responsive reporter vectors for *E. coli*. (A) Plasmid designs for

glutarate responsive vectors (B) Fluorescence measured from either a CsiR (left) or GcdR (right)

transcriptional reporter in *E. coli*. Bars show fluorescence measured in arbitrary units normalized

to OD_{600} in response to mM concentrations of glutarate. Error bars show 95% CI, n=4. (C)

Fluorescence data fit to the Hill equation to derive biosensor performance characteristics. Table

below fitted data summarizes relevant biosensor features.

Sensor	RFPmax	Km (mM)	Hill Coefficient	Linear Range (mM)
CsiR	42000	0.65	0.75	0.05-3.13
GcdR	65000	2.24	1.09	0.05-6.25

Figure 5: Growth and fluorescence of PcsiD and PgcdH reporter in P. putida KT2440 (A)

Growth of strains harboring gcdR reporter (left), or csiR reporter (right) on MOPS minimal

media supplemented with 10mM of glucose, L-lysine, D-lysine, 5-aminovalerate, or 2-

aminoadipate as a sole carbon source. Shaded region represents 95% CI, n=3. (B) RFP

fluorescence of strains harboring gcdR reporter (left), or csiR reporter (right) on MOPS minimal

media supplemented with 10mM of glucose, L-lysine, D-lysine, 5-aminovalerate, or 2-

aminoadipate as a sole carbon source. Shaded region represents 95% CI, n=3.

Table 1.

Strain	JBEI Part ID	Reference
E. coli DH10B		[31]
E. coli BL21(DE3)		Novagen
P. putida KT2440		ATCC 47054
Plasmids		

pET21b		Novagen
pBADT		[32]
pBADT-gcdR-PgcdH::RFP	JPUB_010960	This work
pBADT-csiR-PcsiD::RFP	JPUB_010962	This work
pET21b-CsiR_Pput	JPUB_010964	This work
pET21b-GcdR_Pput	JPUB_010966	This work

Bibliography

1. Dietrich JA, Shis DL, Alikhani A, Keasling JD. Transcription factor-based screens and synthetic selections for microbial small-molecule biosynthesis. ACS Synth Biol. 2013;2: 47–58. doi:10.1021/sb300091d

2. Dietrich JA, McKee AE, Keasling JD. High-throughput metabolic engineering: advances in small-molecule screening and selection. Annu Rev Biochem. 2010;79: 563–590. doi:10.1146/annurev-biochem-062608-095938

3. Petzold CJ, Chan LJG, Nhan M, Adams PD. Analytics for metabolic engineering. Front Bioeng Biotechnol. 2015;3: 135. doi:10.3389/fbioe.2015.00135

4. Kasey CM, Zerrad M, Li Y, Cropp TA, Williams GJ. Development of Transcription Factor-Based Designer Macrolide Biosensors for Metabolic Engineering and Synthetic Biology. ACS Synth Biol. 2018;7: 227–239. doi:10.1021/acssynbio.7b00287

5. Lei C, Wang J, Liu Y, Liu X, Zhao G, Wang J. A feedback regulatory model for RifQ-mediated repression of rifamycin export in Amycolatopsis mediterranei. Microb Cell Fact. 2018;17: 14. doi:10.1186/s12934-018-0863-5

6. Zhang J, Barajas JF, Burdu M, Ruegg TL, Dias B, Keasling JD. Development of a Transcription Factor-Based Lactam Biosensor. ACS Synth Biol. 2017;6: 439–445. doi:10.1021/acssynbio.6b00136

7. Zhang J, Barajas JF, Burdu M, Wang G, Baidoo EE, Keasling JD. Application of an Acyl-

CoA Ligase from Streptomyces aizunensis for Lactam Biosynthesis. ACS Synth Biol.
2017;6: 884–890. doi:10.1021/acssynbio.6b00372

8.   Zhang M, Gao C, Guo X, Guo S, Kang Z, Xiao D, et al. Increased glutarate production by
blocking the glutaryl-CoA dehydrogenation pathway and a catabolic pathway involving L-
2-hydroxyglutarate. Nat Commun. 2018;9: 2114. doi:10.1038/s41467-018-04513-0

9.   Thompson MG, Blake-Hedges JM, Cruz-Morales P, Barajas JF, Curran SC, Harris NC, et
al. Massively parallel fitness profiling reveals multiple novel enzymes in Pseudomonas
putida lysine metabolism. BioRxiv. 2018; doi:10.1101/450254

10.   Madhuri Indurthi S, Chou H-T, Lu C-D. Molecular characterization of lysR-lysXE, gcdR-
gcdHG and amaR-amaAB operons for lysine export and catabolism: a comprehensive
lysine catabolic network in Pseudomonas aeruginosa PAO1. Microbiology (Reading,
Engl). 2016;162: 876–888. doi:10.1099/mic.0.000277

11.   Knorr S, Sinn M, Galetskiy D, Williams RM, Wang C, Müller N, et al. Widespread
bacterial lysine degradation proceeding via glutarate and L-2-hydroxyglutarate. Nat
Commun. 2018;9: 5071. doi:10.1038/s41467-018-07563-6

12.   Jain D. Allosteric control of transcription in GntR family of transcription regulators: A
structural overview. IUBMB Life. 2015;67: 556–563. doi:10.1002/iub.1401

13.   Price MN, Wetmore KM, Waters RJ, Callaghan M, Ray J, Liu H, et al. Mutant phenotypes
for thousands of bacterial genes of unknown function. Nature. 2018;557: 503–509.
doi:10.1038/s41586-018-0124-0

14.   Sagawa S, Price MN, Deutschbauer AM, Arkin AP. Validating regulatory predictions
from diverse bacteria with mutant fitness data. PLoS ONE. 2017;12: e0178258.
doi:10.1371/journal.pone.0178258

15.   Suvorova IA, Korostelev YD, Gelfand MS. GntR Family of Bacterial Transcription
Factors and Their DNA Binding Motifs: Structure, Positioning and Co-Evolution. PLoS
ONE. 2015;10: e0132618. doi:10.1371/journal.pone.0132618

16.   Revelles O, Espinosa-Urgel M, Molin S, Ramos JL. The davDT operon of Pseudomonas
putida, involved in lysine catabolism, is induced in response to the pathway intermediate
delta-aminovaleric acid. J Bacteriol. 2004;186: 3439–3446. doi:10.1128/JB.186.11.3439-
3446.2004

17.   Park SJ, Kim EY, Noh W, Park HM, Oh YH, Lee SH, et al. Metabolic engineering of
Escherichia coli for the production of 5-aminovalerate and glutarate as C5 platform
chemicals. Metab Eng. 2013;16: 42–47. doi:10.1016/j.ymben.2012.11.011

18.   Rohles CM, Gießelmann G, Kohlstedt M, Wittmann C, Becker J. Systems metabolic
engineering of Corynebacterium glutamicum for the production of the carbon-5 platform
chemicals 5-aminovalerate and glutarate. Microb Cell Fact. 2016;15: 154.

369   doi:10.1186/s12934-016-0553-0

370 19. Hagen A, Poust S, Rond T de, Fortman JL, Katz L, Petzold CJ, et al. Engineering a
371   polyketide synthase for in vitro production of adipic acid. ACS Synth Biol. 2016;5: 21–27.
372   doi:10.1021/acssynbio.5b00153

373 20. LaBauve AE, Wargo MJ. Growth and laboratory maintenance of Pseudomonas
374   aeruginosa. Curr Protoc Microbiol. 2012;Chapter 6: Unit 6E.1.
375   doi:10.1002/9780471729259.mc06e01s25

376 21. Ham TS, Dmytriv Z, Plahar H, Chen J, Hillson NJ, Keasling JD. Design, implementation
377   and practice of JBEI-ICE: an open source biological part registry platform and tools.
378   Nucleic Acids Res. 2012;40: e141. doi:10.1093/nar/gks531

379 22. Chen J, Densmore D, Ham TS, Keasling JD, Hillson NJ. DeviceEditor visual biological
380   CAD canvas. J Biol Eng. 2012;6: 1. doi:10.1186/1754-1611-6-1

381 23. Hillson NJ, Rosengarten RD, Keasling JD. j5 DNA assembly design automation software.
382   ACS Synth Biol. 2012;1: 14–21. doi:10.1021/sb2000116

383 24. Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. Enzymatic
384   assembly of DNA molecules up to several hundred kilobases. Nat Methods. 2009;6: 343–
385   345. doi:10.1038/nmeth.1318

386 25. Engler C, Kandzia R, Marillonnet S. A one pot, one step, precision cloning method with
387   high throughput capability. PLoS ONE. 2008;3: e3647. doi:10.1371/journal.pone.0003647

388 26. Jackson DR, Tu SS, Nguyen M, Barajas JF, Schaub AJ, Krug D, et al. Structural Insights
389   into Anthranilate Priming during Type II Polyketide Biosynthesis. ACS Chem Biol.
390   2016;11: 95–103. doi:10.1021/acschembio.5b00500

391 27. Thompson MG, Sedaghatian N, Barajas JF, Wehrs M, Bailey CB, Kaplan N, et al.
392   Isolation and characterization of novel mutations in the pSC101 origin that increase copy
393   number. Sci Rep. 2018;8: 1590. doi:10.1038/s41598-018-20016-w

394 28. Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. ModelFinder:
395   fast model selection for accurate phylogenetic estimates. Nat Methods. 2017;14: 587–589.
396   doi:10.1038/nmeth.4285

397 29. Cruz-Morales P, Ramos-Aboites HE, Licona-Cassani C, Selem-Mójica N, Mejía-Ponce
398   PM, Souza-Saldívar V, et al. Actinobacteria phylogenomics, selective isolation from an
399   iron oligotrophic environment and siderophore functional characterization, unveil new
400   desferrioxamine traits. FEMS Microbiol Ecol. 2017;93. doi:10.1093/femsec/fix086

401 30. Bailey TL, Williams N, Misleh C, Li WW. MEME: discovering and analyzing DNA and
402   protein sequence motifs. Nucleic Acids Res. 2006;34: W369-73. doi:10.1093/nar/gkl198

403 31. Grant SG, Jessee J, Bloom FR, Hanahan D. Differential plasmid rescue from transgenic
404   mouse DNAs into Escherichia coli methylation-restriction mutants. Proc Natl Acad Sci

405   USA. 1990;87: 4645–4649.

406   32.   Bi C, Su P, Müller J, Yeh Y-C, Chhabra SR, Beller HR, et al. Development of a broad-
407   host synthetic biology toolbox for Ralstonia eutropha and its application to engineering
408   hydrocarbon biofuel production. Microb Cell Fact. 2013;12: 107. doi:10.1186/1475-2859-
409   12-107

Glucogenic Catabolism

Ketogenic Catabolism

Alpha-ketoglutarate (2KG)

Glutarate

PP_2910

PP_2909

PP_0159

Glutaryl-CoA

Alpha-hydroxyglutarate (2HG)

PP_0158

Crotonyl-CoA

Succinate

TCA Cycle

Acetyl-CoA

Stenotrophomonas rhizophila Simmons01

Pseudomonas putida KT2440

Pseudomonas aeruginosa strain W60856

Halomonas xianhensis

Virgibacillus halodenitrificans S2

Escherichia fergusonii ECD227

Escherichia albertii B156

Escherichia coli B29595

Citrobacter braakii

Shimwellia blattae DSM 4481

Klebsiella quasipneumoniae Col-Kpn6

Klebsiella aerogenes MGH174

CsiR, transcriptional repressor of CsiD

Carbon starvation induced protein CsiD

L-2-hydroxyglutarate oxidase (EC 1.1.3.15)

Succinate-semialdehyde dehydrogenase [NAD(P)+] (EC 1.2.1.16)

Gamma-aminobutyrate:alpha-ketoglutarate aminotransferase (EC 2.6.1.19)

Gamma-aminobutyrate (GABA) permease

Tricarboxylate transport transcriptional regulator TctD

Tricarboxylate transport sensor protein TctE

Cys regulon transcriptional activator CysB

Others

Conserved CsiR binding site

100 99
100

0.07

A R A A A T G T A G A Y A Y T T Y

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17

A

CsiR Concentration   GcdR Concentration

B

Glutarate Concentration

Sensor	RFPmax	Km (mM)	Hill Coefficient	Linear Range (mM)
CsiR	42000	0.65	0.75	0.05-3.13
GcdR	65000	2.24	1.09	0.05-6.25