Robust characterization of two distinct glutarate sensing transcription factors of *Pseudomonas*

putida L-lysine metabolism

Mitchell G. Thompson[1,2,3], Zak Costello[1,2,4], Niklas F. C. Hummel[1,7], Pablo Cruz-Morales[1,2,5],

Jacquelyn M. Blake-Hedges[1,2,6], Rohith N. Krishna[1,2,6], Will Skyrud[6], Allison N. Pearson[1,2],

Matthew R. Incha[1,2,3], Patrick M. Shih[1,7], Hector Garcia-Martin[1,2,4,8], Jay D. Keasling[1,2,9,10,11,12,*]

[1]Joint BioEnergy Institute, 5885 Hollis Street, Emeryville, CA 94608, USA.

[2]Biological Systems & Engineering Division, Lawrence Berkeley National Laboratory, Berkeley,

CA 94720, USA.

[3]Department of Plant and Microbial Biology, University of California, Berkeley, CA 94720,

USA

[4]DOE Agile BioFoundry, Emeryville, CA, USA

[5]Centro de Biotecnologia FEMSA, Instituto Tecnologico y de Estudios superiores de Monterrey,

Mexico

[6]Department of Chemistry, University of California, Berkeley, CA 94720, USA

[7]Department of Plant Biology, University of California, Davis, Davis, CA, United States

[8]BCAM, Basque Center for Applied Mathematics, Bilbao, Spain

[9]Joint Program in Bioengineering, University of California, Berkeley/San Francisco, CA 94720,

USA

[10]Department of Chemical and Biomolecular Engineering, University of California, Berkeley,

CA 94720, USA

[11]The Novo Nordisk Foundation Center for Biosustainability, Technical University of Denmark,

Denmark

[12]Center for Synthetic Biochemistry, Institute for Synthetic Biology, Shenzhen Institutes for
Advanced Technologies, Shenzhen, China

*To whom correspondence may be addressed. Email: JDKeasling@lbl.gov

ABSTRACT

A significant bottleneck in synthetic biology involves screening large genetically encoded libraries for desirable phenotypes such as chemical production. However, transcription factor-based biosensors can be leveraged to screen thousands of genetic designs for optimal chemical production in engineered microbes. In this study we characterize two glutarate sensing transcription factors (CsiR and GcdR) from *Pseudomonas putida*. The genomic contexts of *csiR* homologs were analyzed and their DNA binding sites were bioinformatically predicted. Both CsiR and GcdR were purified and shown to bind upstream of their coding sequencing *in vitro*. CsiR was shown to dissociate from DNA *in vitro* when exogenous glutarate was added, confirming that it acts as a genetic repressor. Both transcription factors and cognate promoters were then cloned into broad host range vectors to create two glutarate biosensors. Their respective sensing performance features were characterized, and more sensitive derivatives of the GcdR biosensor were created by manipulating the expression of the transcription factor. Sensor vectors were then reintroduced into *P. putida* and evaluated for their ability to respond to glutarate and various lysine metabolites. Additionally, we developed a novel mathematical approach to describe the usable range of detection for genetically encoded biosensors, which may be broadly useful in future efforts to better characterize biosensor performance.

KEYWORDS

Biosensor, Transcription Factor, Pseudmonas putida, Glutarate, Monte Carlo Markov Chain

INTRODUCTION

A rate limiting step in the design-build-test-learn cycle is often the test phase, wherein hundreds or thousands of genetic designs need to be evaluated for their productivity [1,2]. Though recent advances in analytical chemistry have dramatically increased sample throughput [3], transcription factor-based biosensors still offer multiple advantages over traditional chromatographic and mass-spectrometry based detection methods [4,5]. One of the most attractive benefits is the ability to rapidly screen constructs for the production of the target compounds via either plate-based or flow-cytometry-based assays [1–3], which increases throughput by orders of magnitude compared to mass-spectrometry based methods. Additionally, biosensors may offer unmatched sensitivity towards specific ligands, with some sensors having picomolar affinity [3]. The evolution of diverse microbial metabolism has provided researchers with the ability to sense a wide array of ligands, ranging from complex natural products [6,7] to small central metabolites [1,8].

Diacids, polyamines, and lactams are petrochemical derivatives used to produce various polyester and nylon fibers [9,10]. In an effort to make production of these chemicals sustainable, many groups have developed engineered microbes to synthesize these precursors [11–14]. The L-lysine metabolism of *Pseudomonas putida* has been leveraged both in the native host and heterologously to produce valerolactam [15] the diacid glutarate [16,17]. Recently, this utility has inspired much work to uncover missing steps in the lysine catabolism of *P. putida*. These

missing steps included the discovery that glutarate is not only catabolized through the previously

known coA-dependent route to acetyl-coA, but is also catabolized through a coA-independent

route to succinate [16,18] (Figure 1). Recent work has also demonstrated that both glutarate

catabolic pathways are highly upregulated in the presence of glutarate [18]. The *Pseudomonas*

aeruginosa homolog of the ketogenic pathway regulator (GcdR) [19]and the *Escherichia coli*

homolog of the glucogenic pathway regulator (CsiR) have both been characterized. Furthermore

a rigorous investigation of the P. putida homologs has now been reported [20]..

Figure 1: A) The known lysine catabolism of *P. putida*. Dashed box shows the two known pathways of glutarate catabolism in *P. putida*. Highlighted in red is the CoA-independent route of glutarate catabolism, in blue the CoA-dependent route. Enzymes in the metabolism are DavB - L-lysine monooxygenase, DavA - 5-aminopentanamidase, DavT - 5-aminovalerate aminotransferase, DavD - glutaric semialdehyde dehydrogenase, CsiD - glutarate hydroxylase, LghO - L-2-hydroxyglutarate oxidase, GcdG glutaryl-coA transferase, GcdH - glutaryl-CoA dehydrogenase, Alr - alanine racemase, AmaD - D-lysine oxidase, DpkA - Δ1-piperideine-2-carboxylate reductase, AmaB - pipecolate oxidase, AmaA - L-aminoadipate-semialdehyde dehydrogenase, PP_4108 - L-2-aminoadipate aminotransferase, HglS - D-2-hydroxyglutarate synthase, PP_4493 - D-2-hydroxyglutarate dehydrogenase. B) Operonic structure of the two routes of glutarate metabolism in *P. putida*.

In this work we sought to also characterize the two putative local regulators of glutarate catabolism in *P. putida*, *csiR* and *gcdR*. First, we compared the genomic context of *csiR* homologs across bacteria to bioinformatically predict a conserved DNA binding site. We then biochemically and genetically characterized both regulators. Secondly, we developed a novel mathematical approach to rigorously determine the detection ranges for genetically encoded biosensors that can be used to systematically compare biosensors. . Finally, we introduced RFP transcriptional-fusions of the promoter for both catabolic pathways into *P. putida* and evaluated their induction upon the addition of various lysine metabolites.

RESULTS

Genomic contexts of *csiR* and *gcdR* homologs and prediction of *P. putida* binding sites

Work in *Pseudomonas aeruginosa* has characterized the GcdR regulation of ketogenic

glutarate metabolism, and shown that the binding site is conserved across multiple bacterial

species [19]. While binding sites for CsiR in *E. coli* have been identified, it was yet to be

investigated whether there is a conserved binding site for homologs across bacterial species [21]. In

order to identify conserved binding sites of CsiR homologs, we compared the syntenic genomic

contexts of 12 selected genomes that contained neighboring *csiD* and *csiR* homologs. Genes

encoding *csiR* were found in two distinct genomic contexts, either transcribed divergently from

csiD as found in *P. putida*, or transcribed as the last gene in the *csiD* operon as in *E. coli* (Figure

2).   The genomic regions upstream of the *csiD* homolog were extracted and Multiple EM for

Motif Elicitation (MEME) was used to identify a conserved CsiR binding motif [22]. A consensus

A(A/G)AAATCTAGA(C/T)ATTTT motif was identified upstream of each *csiD* homolog.

Previously, footprinting assays in *E. coli* BW25113 revealed two primary and two secondary

bindings sites with the sequences $TTGTN_5TTTT$ and $ATGTN_5TTTT$ respectively [21]. While this

manuscript was under review Zhang et al. demonstrated via DNase I footprinting that CsiR does

indeed binding at two locations upstream of *csiD*, including the conserved

A(A/G)AAATCTAGA(C/T)ATTTT motif [20]. Our consensus motif agrees closely with the

secondary binding site identified, and highly suggests that the binding site of CsiR is conserved

across the bacteria where homologs are present.

Figure 2: Genomic contexts of *csiR* homologs and predicted binding regions: Phylogeny of assorted gammaproteobacteria and the location of their *csiD* operons. MEME analysis of the intergenic regions upstream of the *csiD* operon resulted in the sequence motifs depicted below the tree.

Biochemical characterization of CsiR and GcdR

To determine if the *P. putida* CsiR also acts as a regulator and to identify its putative binding sites, we biochemically characterized this protein using electrophoretic mobility shift assays (EMSAs). The CsiR protein was purified (Figure S1) and incubated with DNA probes consisting of the intergenic region between *csiR* and *csiD*. The assay showed multiple binding

sites in this intergenic region, as four distinct bands appeared. These results appear to confirm

previous work in *E. coli in which* four binding sites of the *E. coli* CsiR homolog were observed

[21]. CsiR had a high affinity for the DNA probe, with a calculated K_d of approximately 30 nM

(Figure 3Aa), which is similar to the 10 nM CsiR/DNA K_d of the *E. coli* CsiR homolog, and

nearly identical to the results of Zhang et al. which showed complete probe shift at 6-fold molar

excess CsiR [20,21]. Purified GcdR (Figure S1) also bound to its cognate probe with a single

distinct shift and had an an estimated K_d of approximately 62.5 nM (Figure 3A) [19]. Again, these

results were consistent with Zhang et al., who showed near complete probe shifts at 8-fold molar

excess GcdR [20]. As CsiR is a GntR family transcriptional regulator, many of which act as

repressors [23], we evaluated whether glutarate would decrease the DNA binding affinity of CsiR.

EMSA assays were repeated in the presence of increasing glutarate concentrations. Analysis by

gel electrophoresis revealed increasing quantities of free probe as glutarate concentrations

increased (Figure 3B). These results suggest that the *P. putida* CsiR is a glutarate-responsive

repressor of the *csiDlhgO* operon[21]. Zhang et al. also showed that in addition to dissociating from

the DNA probe at high concentrations of glutarate, CsiR was also responsive to 2-

hydroxyglutarate (2HG) [20].

Figure 3: Biochemical characterization of CsiR and GcdR. (A) EMSA of either purified

CsiR (left) or GcdR (right) incubated with 10 nM fluorescent probe of intergenic region

between *csiR* and *csiD*. Concentrations of CsiR protein range from 0 nM to 250 nM. FP

denotes free probe (B) EMSA showing 100 nM CsiR and 10 nM fluorescent probe

incubated with glutarate from concentrations of 0 mM to 5 mM. Far left lane shows free

probe with no CsiR. FP denotes free probe.

Development of two glutarate biosensor vectors

In order to evaluate CsiR and GcdR as biosensors, we cloned both the regulator and

intergenic region of both the coA-independent and coA-dependent glutarate catabolism pathways

upstream of RFP on the broad host range vector pBADT. *E. coli* DH10B harboring either vector

were grown for 24 hours in EZ-Rich medium supplemented with concentrations of glutarate

ranging from 0.00015 mM to 2.5 mM, after which both OD600 and RFP fluorescence were

measured. At 2.5 mM glutarate, the GcdR and CsiR vectors demonstrated a maximal induction

over background of 55.5 and 1.5 times over uninduced cells, respectively; however, the CsiR

system showed considerable background signal (Table 1). Normalized RFP expression for each

sensor was fitted to the Hill equation to derive biosensor performance. The GcdR system was

found to have a Hill coefficient of 1.33, a K_d of 0.32 mM, and a maximum predicted normalized

RFP expression of 5403, while the CsiR system was shown to have a Hill coefficient of 1.61, a

K_d of 0.016 mM, and a maximum predicted normalized RFP expression of 3223 (Figure 4A). To

test for the ability of CsiR or GcdR to sense other diacids, *E. coli* harboring either vector were

grown in LB medium with 5 mM to 0.00015 mM of either succinate, adipate, or pimelate. While

pimelate and succinate were unable to induce either system, 2.5 mM adipate induced the CsiR

biosensor ~0.2x, and induced the GcdR system ~4x over background (Figure S2).

Given that GcdR showed a greater dynamic and substrate range with no response to 2-

HG, we tested whether the performance characteristics of these GcdR system could be altered.

The promoter region 50 bp upstream sequence of *gcdR* in the GcdR-sensing vector was replaced

by three different previously characterized constitutive promoters from the Anderson collection

(J23101, J23110, and J23113), representing a high (1791 RFP AU), medium (844 RFP AU) , and

low (21 RFP AU) strengths [24]. All of the engineered GcdR systems showed reduced limits of

detection and increased sensitivity to glutarate compared to natively regulated GcdR, with each

new vector showing a decreased K_d (J23113 : 0.008 mM, J23110 : 0.01 mM, J23101 : 0.01 mM)

(Figure 4B).All engineered GcdR vectors showed ~4x the maximal expression compared to the

native system (Figure 4B), but also had nearly 30x greater basal RFP expression (Table 1).

While all three engineered GcdR vectors performed similarly, the limit of detection of J23110

was 5x less than the other two vectors, likely due to the lower basal expression of the vector

(Table 1).

Figure 4: Development of glutarate responsive reporter vectors for *E. coli*. (A)

Fluorescence data fit to the Hill equation to derive biosensor performance characteristics

for native CsiR and GcdR systems. Points show individual experimental measurements.

187   Shaded area represents (+/-) one standard deviation, n=4. (B) Fluorescence data fit to the

188   Hill equation to derive biosensor performance characteristics for the engineered GcdR

189   systems. Points show individual experimental measurements. Relative promoter strength is

190   shown to the right. Shaded area represents (+/-) one standard deviation, n=4.

191   Table 1: Biosensor performance parameters with standard deviations. Sensitivity: Hill

192   coefficient of fitted data. Kd: Concentration of ligand to achieve half predicted maximal

193   RFP expression. Max (Normalized RFP): Predicted maximal RFP expression after OD

194   normalization and subtraction of uninduced expression. Dynamic range: Minimal and

195   maximal experimental range of OD normalized RFP. Induction: Ratio of maximal

196   experimental induction over basal expression. Inducer Range: Experimentally determined

197   range of glutarate that can detected with biosensor.

Reporter	Sensitivity	Kd (mM)	Max (Normalized RFP)	Dynamic Range (RFP/OD600)	Induction	Inducer Range (mM)
CsiR-Native	1.61 (+/- 0.16)	0.016 (+/- 0.002)	3223 (+/- 72)	5704-8357	1.5	0.005-0.0195
GcdR-Native	1.33 (+/- 0.03)	0.32 (+/- 0.006)	5353 (+/- 38.5)	97-5403	55.5	0.01-2.5
GcdR-J23113	1.64 (+/- 0.09)	0.008 (+/- 0.0003)	20896 (+/- 184)	3272-24027	7.3	0.001-0.0195
GcdR-J23101	1.74 (+/- 0.07)	0.01 (+/- 0.0003)	18881 (+/- 131)	2428-20099	8.3	0.001-0.0195

GcdR-J23110	1.42 (+/- 0.09)	0.01 (+/- 0.0004)	20113 (+/- 214)	2113-20690	9.8	0.0002-0.0195

Developing metrics to quantify biosensor performance

In previously published work the mathematical basis for determining linear range or limit

of detection has often been obscure and non-systematic [1,8,25]. Furthermore, a linear range may not

be able to adequately capture the performance characteristics of a biosensor over the range of

ligand concentrations where the sensor can still resolve differences. To address these

deficiencies, we sought to develop a mathematical method for evaluating the sensing

performances of genetically encoded transcription factors fitted to the Hill equation. Our

approach uses a probabilistic model to relate inducer concentration and corresponding

fluorescence measurements fit to the Hill equation assuming: (1) fluorescence measurements at a

particular concentration are normally distributed, (2) the variance of fluorescence measurements

is roughly constant over the range of measured values, and (3) the relationship between ligand

concentrations and fluorescence can be well modeled using the Hill function. This model allows

us to estimate the concentration of ligand compatible with our observed fluorescence data given

the variance of the data as determined via Markov Chain Monte Carlo (MCMC) sampling [26]. A

detailed methodological description can be found in the supplemental Jupyter notebook, which

can be used to analyze other biosensor data.

By applying MCMC sampling to the model of our native GcdR biosensor, we can readily

produce the probability density functions (i.e. the probability that the ligand produces the

observed fluorescent response) of specific ligand concentrations (Figure 5A). At glutarate

concentrations of 0.25 mM, 0.68 mM, and 1.125 mM associated fluorescence values can be

resolved from one another (no overlap), however the biosensor is less able to resolve ligand

concentrations between 1.5 mM and 2 mM (Figure 5A). When we apply MCMC sampling to the

GcdR sensor being driven by the J23101 promoter, we observe that this system possesses the

resolution to distinguish between 0.004 mM and 0.011 mM, but is less able to distinguish

between higher concentrations (Figure 5B). A biosensor's resolution window, defined as the

width of the 95% prediction interval of inducer concentrations derived from a set of fluorescence

measurements, can then be expressed as a continuous function across a range of ligand

concentrations for a given biosensor (Figure 5C). Below concentrations of ~0.01 mM glutarate

the J23101 GcdR biosensor has greater resolution, while at higher concentrations the native

GcdR sensor system has greater resolution(Figure 5C). Another important aspect of our approach

is that it allows for the resolution window to be calculated as a function of the number of

replicates in a biosensor experiment. If either variance decreases or sample size increases, the

resolution of a biosensor also increases. By simulating sample sizes of 1 through 100 via MCMC

sampling, the theoretical resolution of the native GcdR dramatically increases (Figure 4D). This

"power" analysis may serve as a guide for experimental design when a certain biosensor

resolution is required for a given application. We believe this approach may be generally useful

to any dataset derived from fluorescent transcription factor based biosensors. To demonstrate this

we also applied our MCMC methodology to two well characterized BglBrick vectors, pBbBSk-

rfp and pBbBEk-rfp, which express RFP from arabinose-inducible vectors from SC101 or ColE1

origins respectively (Figure S3). Fluorescence data from each vector (Figure S3A), was fit to the

Hill question (Figure S3B), and demonstrated that both vectors had the highest resolution at an

arabinose concentration of ~0.1% w/v (Figure S3C).

Figure 5: Development of analytics for biosensor performance. Probability density

functions of fluorescence values compatible with concentrations of glutarate for selected

ligand concentrations modelled to the Hill equation of the native GcdR system (A) or the

GcdR J23101 system (B). (C) Resolution of native GcdR or GcdR J23101 biosensor systems

over a select range of glutarate concentrations. (D) Theoretical resolution of the GcdR

native biosensor with differing number of replicates.

<u>Responsiveness of glutarate biosensors to lysine metabolites in *P. putida*</u>

To assess the ability of these vectors to function in *P. putida* both plasmids were

introduced into either wild type *P. putida* or a strain with both known pathways of glutarate

catabolism deleted (Δ*csiD*Δ*gcdH* - referred to as Δglutarate). The resulting strains were grown

in MOPS minimal medium supplemented with 10 mM glucose and glutarate ranging from 5 mM

to 0.01 mM for 24 hours. Both vectors responded to increased concentrations of exogenously

applied glutarate, though the GcdR vector had ~10x greater fluorescence than the CsiR vector

(Figure 6A). The Δglutarate strain showed increased levels of RFP induction using both the CsiR

and GcdR systems, suggesting both vectors are able to sense increased levels of glutarate (Figure

6A). To further examine the ability of these vectors to probe *P. putida* lysine metabolism, both

were also introduced into a Δ*davT* strain, which is unable to metabolize 5-aminovalerate to

glutarate semialdehyde (Figure 1A) and therefore precludes glutarate production. When wild

type, Δ*davT*, and Δglutarate strains harboring either the CsiR or GcdR systems were grown on

minimal medium supplemented with 10 mM glucose and 10 mM 5-aminovalerate, both vectors

in the Δ*davT* strain showed decreased fluorescence compared to wild type, while vectors in the

Δglutarate strains showed increased fluorescence (Figure 6B). Measurement of intracellular 5-

aminovalerate showed significant pools of the metabolite in the Δ*davT* strains (~1500

uM/OD600) with no detectable 5-aminovalerate in the other genetic backgrounds (Figure 6C).

These results highly suggest that both GcdR and CsiR are insensitive to 5-aminovalerate, an

essential feature of these sensors if they are to be used in organisms that derive glutarate from a

5-aminovalerate precursor.

To evaluate the ability of both reporter vectors to monitor the catabolism of other lysine

metabolites, wild-type *P. putida* harboring either GcdR or CsiR were grown in minimal media

supplemented with either 10 mM glucose, L-lysine, D-lysine, 5-aminovalerate, or 2-

aminoadipate for 48 hours with OD_{600} and RFP fluorescence being measured continuously

(Figure S4A and Figure S4B). Neither vector was induced when the bacterium was grown on

glucose (Figure S4). The GcdR vector was strongly induced when grown on 5-aminovalerate,

and to a lesser extent 2-aminoadipate, D-lysine, and L-lysine (Figure S4A). Conversely, the

strain harboring the CsiR vector only displayed induction of RFP above background when grown

on 5-aminovalerate (Figure S4B). While 5-aminovalerate was able to induce RFP induction in

the strain harboring the CsiR vector, induction was extremely limited compared to induction of

RFP from the GcdR vector (Figure S4).

Figure 6: Performance of CsiR and GcdR biosensors in *P. putida*. (A) RFP expression of either wild type *P. putida* or *P. putida* with the ability to metabolize glutarate knocked out, measured with either the CsiR or GcdR biosensor under different external glutarate concentrations. Error bars represent 95% CI, n=4 **(B)** RFP expression of wild type, Δ*davT*, or Δglutarate strains of *P. putida* harboring either the CsiR or GcdR biosensor when grown on 10 mM glucose and 10 mM 5-aminovalerate (5AVA). Error bars represent 95% CI, n=3. **(C)** Intracellular concentration of 5-aminovalerate of wild type, Δ*davT*, or Δglutarate strains of *P. putida* harboring either the CsiR or GcdR biosensor when grown on 10 mM glucose and 10 mM 5-aminovalerate (5AVA). Error bars represent 95% CI, n=3.

DISCUSSION

Recent advances in high-throughput functional genomics have allowed researchers to rapidly identify novel metabolic pathways, and in turn infer the function of novel transcription factors [27,28]. Rigorous characterization of these regulators is a critical step to developing novel parts for synthetic biology as well as useful tools for metabolic engineering. The discovery of two distinct pathways for glutarate catabolism within *P. putida*, regulated by independent transcription factors, presents an interesting opportunity to compare and contrast the relative sensing properties of each system.

CsiR and GcdR homologs have now been characterized thoroughly in multiple bacteria, and have demonstrated that CsiR acts as a repressor of csiD whereas GcdR acts as a positive regulator of gcdHG [19–21]. Our bioinformatic analyses suggest that the binding site of CsiR is highly conserved amongst bacteria that possess the regulator (Figure 2). While our CsiR biosensor was shown to have a lower limit of detection of glutarate than the unengineered GcdR biosensor (Table 1), Zhang et al. demonstrated that CsiR is also responsive to 2-HG [20]. Given

that the GcdR biosensor can readily be engineered to have lower limits of detections, it is likely

the better choice to detect glutarate via a genetically encoded transcription factor.

Multiple works have demonstrated that engineering changes to the expression of a

transcription factor, the responsiveness of the transcriptional output can also be changed [31,32]. .

By engineering the promoter of the *gcdR* transcription factor, the sensitivity of all the vectors

were increased from ~1.3 to ~1.7, while the K_d was lowered from 0.32 mM to ~0.01 mM (Figure

4B). This sensitivity to glutarate may make this vector useful in prototyping novel routes to

biological production of the C5 diacid. However at high concentrations of adipate (>1.25 mM)

GcdR is also activated, which may confound results (Figure S2).

While our preliminary work with exogenously applied ligand is promising, further work

remains to be done to evaluate the ability of these sensors to detect flux of glutarate in living

cells. Multiple recent publications have shown that glutarate metabolism is widespread in

bacteria, and evaluating the ability of the CsiR and GcdR sensors to measure this flux will

require careful experimentation with C^{13} labelled substrates. Another possible confounding

factor in utilizing these vectors is the presence of CsiR or GcdR binding sites in the host

organism. The overexpression of either transcription factor may misregulate host metabolism.

This hypothesis is supported by the differences in growth observed between GcdR and CsiR

biosensor containing *P. putida* when grown on lysine metabolites (Figure S4).

The overarching goal of synthetic biology is to apply engineering principles to biological

systems so that outcomes of genetic manipulation can become more predictable and repeatable

[33,34]. While there has been a large body of work devoted to the characterization and application

of biosensors, there has been conspicuously few attempts to rigorously describe at which ligand

concentrations a biosensor is useful. Often the analysis merely states limit of detection, and a

'linear range' with little in terms of a mathematical justification. Here we present an alternative

metric that allows for the calculation of ligand resolution across the entire range of detection for

a biosensor. By leveraging a MCMC approach to predicting ligand ranges compatible with

fluorescence values, researchers can more precisely describe a biosensors performance and

identify whether a given biosensing system is potentially useful for a given engineering task. The

MCMC approach also allows for the simulation of an increasing number of replicates, which

could inform the researcher of the replicates that may be required in an experimental design to

achieve a desired level of resolution. We hope that this initial work to better characterize

biosensor performance inspires other groups to develop even more sophisticated methods of

analysis.

In addition to their utility as biosensors for metabolic engineering, these sensors may be a

valuable tool in studying the carbon utilization in the native host *P. putida*. Work conducted here

demonstrates the ability of both CsiR and GcdR sensors to distinguish between glutarate

accumulating and mutants blocked in their ability to metabolize 5-aminovalerate to glutarate

(Figure 6). Lysine metabolism in *P. putida* is isomer specific, with each isomer being degraded

by a separate catabolic pathway [18]. While cross-feeding between the pathways has been proposed

previously [35], recent work by our group has proposed a molecular mechanism for metabolite

exchange between the D- and L- catabolic pathways [18]. The exchange relies on the 2-oxoacid

promiscuity of the non-heme Fe(II) oxidase CsiD, which normally catalyzes the hydroxylation of

glutarate using 2-ketoglutarate as a cosubstrate to yield 2-hydroxyglutarate and succinate. CsiD

can also use 2-oxoadipate, a D-lysine catabolic intermediate, as a 2-oxoacid cosubstrate to yield

2-hydroxyglutarate and glutarate as products. The glutarate from this reaction could then proceed

down the L-lysine catabolic pathway. When 2-aminoadipate was fed into *P. putida* harboring the

GcdR vector, fluorescence was observed in stationary phase. As 2-aminoadipate immediately

precedes 2-oxoadipate in the D-lysine catabolic pathway, these results support the hypothesis

that CsiD could act as a bridge between the two catabolic pathways. There has been substantial

interest in developing microbes to produce glutarate, with strains of *E. coli*, *P. putida*, and

Corynebacterium glutamicum all engineered to produce high titers [16,36,37]. Further engineering of

the GcdR system may be able to extend the resolvable range to higher concentrations, furthering

its utility as a tool to achieve even higher titers of glutarate. Though glutarate is a valuable

commodity chemical, the C6 diacid adipate is used in much greater quantities primarily as a

monomer used to make nylons 33. This slight sensitivity of GcdR toward adipic acid is

especially interesting, as recent work has demonstrated the effectiveness of evolving

transcription factors to sense non-native ligands [6]. Such methods could be applied to GcdR in

order to expand its utility in sensing other industrially important diacids.

METHODS

<u>Media, chemicals, and culture conditions</u>

E. coli cultures were grown in Luria-Bertani (LB) Miller medium (BD Biosciences,

USA) at 37 °C while *P. putida* was grown at 30 °C. When indicated, *P. putida* and *E. coli* were

grown on modified MOPS minimal medium [39]. Cultures were supplemented with kanamycin

(50 mg/L, Sigma Aldrich, USA), gentamicin (30 mg/L, Fisher Scientific, USA), or carbenicillin

(100 mg/L, Sigma Aldrich, USA), when indicated. All other compounds were purchased through

Sigma Aldrich (Sigma Aldrich, USA) .

<u>Strains and plasmids</u>

All bacterial strains and plasmids used in this work are listed in Table 3. All strains and

plasmids created in this work are available through the public instance of the JBEI registry.

(https://public-registry.jbei.org/folders/390). All plasmids were designed using DeviceEditor and

VectorEditor software, while all primers used for the construction of plasmids were designed

using j5 software [40–42]. Plasmids were assembled via Gibson Assembly using standard protocols

[43], or Golden Gate Assembly using standard protocols [44]. Plasmids were routinely isolated using

the Qiaprep Spin Miniprep kit (Qiagen, USA), and all primers were purchased from Integrated

DNA Technologies (IDT, Coralville, IA). Construction of *P. putida* deletion mutants was

performed as described previously [18].

Expression and purification of proteins

Proteins were purified as described previously [45]. Briefly, 500 mL cultures of *E. coli*

BL21 (DE3) harboring expression plasmids were grown in LB medium at 37 $^{\circ}$C to an OD of 0.6

then induced with 1mM isopropyl β-D-1-thiogalactopyranoside. Cells were allowed to express

for 18 hours at 30 $^{\circ}$C before being harvested via centrifugation. Cell pellets were stored at -80 $^{\circ}$C

until purification. Cell pellets were then resuspended in lysis buffer (50 mM sodium phosphate,

300 mM sodium chloride, 10 mM imidazole, 8% glycerol, pH 7.5) and sonicated to lyse cells.

Insolubles were pelleted via centrifugation (30 minutes at 40,000 x g). The supernatant was

applied to a fritted column containing Ni-NTA resin (Qiagen, USA) which had been pre-

equilibrated with several column volumes of lysis buffer. The resin was washed with lysis buffer

containing 50 mM imidazole, then the protein was eluted using a stepwise gradient of lysis

buffer containing increasing imidazole concentrations (100 mM, 200 mM, and 400 mM).

Fractions were collected and analyzed via SDS-PAGE. Purified proteins were concentrated using

Spin-X UF 20 (10 kDa MWCO) spin concentrators (Corning, Inc.). Concentrated protein was

stored at 4 $^{\circ}$C until *in vitro* analysis.

<u>Plate based growth and fluorescence assays in P. putida</u>

Growth studies of bacterial strains were conducted via microplate reader kinetic assays.

Overnight cultures were inoculated into 10 mL of LB medium from single colonies, and grown

at 30 °C. These cultures were then washed twice with MOPS minimal medium without any

added carbon and diluted 1:100 into 500 uL of MOPS medium with 10 mM of a carbon source in

48-well plates (Falcon, 353072). Plates were sealed with a gas-permeable microplate adhesive

film (VWR, USA), and then optical density and fluorescence were monitored for 48 hours in a

Biotek Synergy 4 plate reader (BioTek, USA) at 30 °C with fast continuous shaking. Optical

density was measured at 600 nm, while fluorescence was measured using an excitation

wavelength of 485 nm and an emission wavelength of 620 nm with a manually set gain of 100.

<u>Transcriptional fusion fluorescence assays</u>

To measure RFP production in *E. coli,* fluorescence measurements were obtained from

single time points of cells grown in deep-well 96-well plates as described previously with minor

changes[46]. Briefly, cells were grown in 500 µL of EZ-Rich medium supplemented with

kanamycin and a range of glutarate concentrations from 5 mM to 0 mM. Plates were sealed with

AeraSeal film (Excel Scientific, AC1201-02) and grown for 22 hours at 37 °C on a 200 rpm

shaker rack. After incubation, 100 µL from each well was aliquoted into a black, clear-bottom

96-well plate and fluorescence was measured with a Tecan Infinite F200 plate reader (Tecan,

USA). Optical density was measured at 600 nm, while fluorescence was measured using an

excitation wavelength of 535 nm and an emission wavelength of 620 nm with a manually set

gain of 60. Endpoint fluorescence assays in P. putida were carried out in LB media, and

measured by the same method.

Electrophoretic mobility shift assays

Electrophoretic mobility shift assays were performed as previously described [7]. 6-

Carboxyfluorescein labelled PCR products for CsiR probes were generated from the intergenic

region between PP_2908 and PP_2909 using primers csiRprobeFOR 5'-6-

FAM/AGTTCGATCTGCGTAAAG-3' and csiRprobeREV 5'-CCCGCTGAATGCTGAGTT-

3'), while probes for GcdR were generated from the intergenic region between PP_0157 and

PP_0158 with primers gcdRprobeFOR 5'-6-FAM/CGGGTCGATCCAGTTGAAA-3' and

gcdRprobeREV 5'-GCATGTACGTCAACCTCACT-3'. Primers were purchased from IDT

Technologies (IDT, Coralville, IA). PCR product was then purified with a QIAquick PCR

Purification Kit (Qiagen, USA), and the amount of DNA was quantified via a NanoDrop 2000C

(Thermo Fisher Scientific). Binding reactions were conducted with 10 ng of labelled probe,

which was added to 10 mM Tris–HCl (pH 8.0), 25 mM KCl, 2.5 mM MgCl2, 1.0 mM DTT and

2 ug salmon sperm DNA in 20 uL reactions. CsiR was added to reactions in concentrations

ranging from 250 nM to ~4 nM, in addition to a control reaction without CsiR, and then allowed

to incubate at 22 $^{\circ}$C for 20 minutes. Reactions were then loaded into 10% native polyacrylamide

gels buffered with 0.5x TBE. Afterwards, electrophoresis gels were imaged on an Alpha

Innotech MultiImage III (Alpha Innotech). Adobe Photoshop was used to average pixel intensity

over the entire band on EMSA gels in order to estimate the K_d.

Measurement of 5-aminovalerate

To measure intracellular concentrations of 5-aminovalerate, cells were quenched as

previously described [47]. LC/MS analysis was performed on an Agilent 6120 single quadrupole

LC/MS equipped with a Waters Atlantis Hilic 5 µM Silica column (4.6 x 150 mm). A linear

gradient of 100-30% 90:10 CH3CN:H20 with 10 mM ammonium formate and 0.1% formic acid

(v/v) over 20 min in 90:10 H2O:CH3CN with 10 mM ammonium formate and 0.1% formic acid

(v/v) at a flow rate of 1.0 mL/min was used. Extracted ion chromatograms were integrated and

peak area was used to construct a standard curve using an authentic 5-aminovalerate standard.

Concentrations of 5-aminovalerate within samples were interpolated from this curve.

<u>Analysis of biosensor parameters</u>

A model relating inducer concentrations and fluorescence measurements to characterize

the performance of a biosensor was generated under the following assumptions 1) the

relationship between analyte concentrations and fluorescence can be well modeled using the Hill

equation 2) fluorescence measurements at a particular concentration are normally distributed 3)

the variance of fluorescence measurements is roughly constant over the range of measured

values.

Under these assumptions we can phrase the following probabilistic model via equation 1

(Figure 7). Using the probabilistic model which captures our constraints on the problem the log

likelihood function is expressed as equation 2 (Figure 7).The log likelihood is used to express the

maximum likelihood estimation (MLE) problem as equation 3 (Figure 7),which when solved

results in the optimal parameters of the model given the characterization data. In order to

estimate the distribution of ligand concentrations that are compatible with experimental

fluorescence data, MCMC sampling was used to solve the MLE problem equation 4 (Figure 7).

We determined biosensor resolution by solving the above maximum likelihood estimation

problem iteratively over the range of observed fluorescences during the biosensor

characterization process. This can determine the relationship between an inducer concentration

estimate and the estimated standard deviation. The standard deviation of the estimate of inducer

concentration can be interpreted as the resolution window. In the case of this work, two standard

deviations is considered the resolution window of the sensor, as 95% of the compatible inducer

concentration estimates fall within the interval.

To determine minimal and maximal levels of ligand detection of a given biosensor, the

minimal detection limit was defined as the minimal concentration of inducer that resulted in a

OD600 normalized RFP value significantly (t-test pval <0.05) greater than that of uninduced

cultures, while the maximal detection limit was defined as the greatest concentration of inducer

that resulted in a OD600 normalized RFP value significantly (t-test pval <0.05) less than that of

cultures induced with the highest concentration of ligand experimentally tested (2.5 mM).

A comprehensive methodological description of calculating biosensor performance

parameters, as well as usable Jupyter notebooks can be found at

https://github.com/JBEI/biosensor_characterization_public.

Equation 1

$$P(F|C, \theta, \alpha) = \mathcal{N}(h_\theta(C), \sigma)$$

Equation 2

$$\ell(\theta, \sigma|D) = \sum_{i=1}^{N} \log P(F = f_i|C = c_i, \sigma, \theta).$$

Equation 3

$$\hat{\theta}, \hat{\sigma} = \arg \max_{\theta, \sigma} \ell(\theta, \sigma|D),$$

Equation 4

$$\hat{c} = \arg \max_{c} \sum_{i=1}^{N} \log P(F = f_i|C = c, \hat{\theta}, \hat{\sigma})$$

Figure 7: Equations used in this study for MCMC analysis of biosensor data.

<u>Bioinformatic analysis</u>

For the phylogenetic reconstructions, the best amino acid substitution model was selected

using ModelFinder as implemented on IQ-tree [48] , phylogenetic trees were constructed using IQ-

tree, and nodes were supported with 10,000 bootstrap replicates. The final tree figures were

edited using FigTree v1.4.3 (http://tree.bio.ed.ac.uk/software/figtree/). Orthologous syntenic

regions of CsiR were identified with CORASON-BGC [49] and manually colored and annotated.

DNA-binding sites were predicted with MEME [22].

SUPPORTING INFORMATION

The Supporting Information is available free of charge on the ACS Publications website.

S1.pdf contains supplementary figures.

Acknowledgements

The authors would like to thank Tyler Backman for assistance creating Git repositories. The

authors would like to thank the UC Berkeley SMART program for providing support for R.K to

conduct summer research. This work was part of the DOE Joint BioEnergy Institute

(https://www.jbei.org) supported by the U. S. Department of Energy, Office of Science, Office of

Biological and Environmental Research, and was part of the Agile BioFoundry

(http://agilebiofoundry.org) supported by the U.S. Department of Energy, Energy Efficiency and

Renewable Energy, Bioenergy Technologies Office, through contract DE-AC02-05CH11231

between Lawrence Berkeley National Laboratory and the U.S. Department of Energy. H.G.M.

was also supported by the Basque Government through the BERC 2018-2021 program and by

Spanish Ministry of Economy and Competitiveness MINECO: BCAM Severo Ochoa excellence

513   **Contributions**

517   **Competing Interests**

519   **Table 2. Strains and Plasmids used in this work**

Strain	JBEI Part ID	Reference
E. coli DH10B		50
E. coli BL21(DE3)		Novagen
E.coli S17		ATCC 47055
P. putida KT2440		ATCC 47054
P. putida Δ*davT*	JPUB_013544	This work
P. putida Δ*gcdH*Δ*csiD*		18
Plasmids		
pET21b		Novagen
pBADT		51
pMQ30		52
pBADT-gcdR-PgcdH::RFP	JPUB_010960	This work
pBADT-csiR-PcsiD::RFP	JPUB_010962	This work
pBADT-gcdR-J23101	JPUB_013546	This work
pBADT-gcdR-J23110	JPUB_013548	This work
pBADT-gcdR-J23113	JPUB_013550	This work
pMQ30 davT	JPUB_013544	This work
pET21b-CsiR_Pput	JPUB_010964	This work

pET21b-GcdR_Pput	JPUB_010966	This work
pBbS8k-rfp		[53]
pBbE8k-rfp		[53]

Bibliography

(1) Dietrich, J. A., Shis, D. L., Alikhani, A., and Keasling, J. D. (2013) Transcription factor-
based screens and synthetic selections for microbial small-molecule biosynthesis. *ACS Synth.*
Biol. 2, 47–58.
(2) Dietrich, J. A., McKee, A. E., and Keasling, J. D. (2010) High-throughput metabolic
engineering: advances in small-molecule screening and selection. *Annu. Rev. Biochem. 79*, 563–
590.
(3) Petzold, C. J., Chan, L. J. G., Nhan, M., and Adams, P. D. (2015) Analytics for metabolic
engineering. *Front. Bioeng. Biotechnol. 3*, 135.
(4) D'Ambrosio, V., and Jensen, M. K. (2017) Lighting up yeast cell factories by transcription
factor-based biosensors. *FEMS Yeast Res. 17*.
(5) Cheng, F., Tang, X.-L., and Kardashliev, T. (2018) Transcription Factor-Based Biosensors in
High-Throughput Screening: Advances and Applications. *Biotechnol. J. 13*, e1700648.
(6) Kasey, C. M., Zerrad, M., Li, Y., Cropp, T. A., and Williams, G. J. (2018) Development of
Transcription Factor-Based Designer Macrolide Biosensors for Metabolic Engineering and
Synthetic Biology. *ACS Synth. Biol. 7*, 227–239.
(7) Lei, C., Wang, J., Liu, Y., Liu, X., Zhao, G., and Wang, J. (2018) A feedback regulatory
model for RifQ-mediated repression of rifamycin export in Amycolatopsis mediterranei. *Microb.*
Cell Fact. 17, 14.
(8) Zhang, J., Barajas, J. F., Burdu, M., Ruegg, T. L., Dias, B., and Keasling, J. D. (2017)
Development of a Transcription Factor-Based Lactam Biosensor. *ACS Synth. Biol. 6*, 439–445.
(9) Gilbert, M. (2017) Aliphatic Polyamides, in *Brydson's Plastics Materials*, pp 487–511.
Elsevier.
(10) Chae, T. U., Ahn, J. H., Ko, Y.-S., Kim, J. W., Lee, J. A., Lee, E. H., and Lee, S. Y. (2019)
Metabolic engineering for the production of dicarboxylic acids and diamines. *Metab. Eng.*
(11) Lee, S. Y., Kim, H. U., Chae, T. U., Cho, J. S., Kim, J. W., Shin, J. H., Kim, D. I., Ko, Y.-
S., Jang, W. D., and Jang, Y.-S. (2019) A comprehensive metabolic map for production of bio-
based chemicals. *Nat. Catal. 2*, 18–33.
(12) Li, Z., Shen, Y.-P., Jiang, X.-L., Feng, L.-S., and Liu, J.-Z. (2018) Metabolic evolution and

a comparative omics analysis of Corynebacterium glutamicum for putrescine production. *J. Ind.*
Microbiol. Biotechnol. 45, 123–139.
(13) Li, M., Li, D., Huang, Y., Liu, M., Wang, H., Tang, Q., and Lu, F. (2014) Improving the
secretion of cadaverine in Corynebacterium glutamicum by cadaverine-lysine antiporter. *J. Ind.*
Microbiol. Biotechnol. 41, 701–709.
(14) Picataggio, S., Rohrer, T., Deanda, K., Lanning, D., Reynolds, R., Mielenz, J., and Eirich,
L. D. (1992) Metabolic engineering of candida tropicalis for the production of long–chain
dicarboxylic acids. *Nat. Biotechnol. 10*, 894–898.
(15) Thompson, M., Valencia, L. E., Blake-Hedges, J., Cruz-Morales, P., Velasquez, A.,
Pearson, A., Sermeno, L., Sharpless, W., Benites, V., Chen, Y., Baidoo, E., Petzold, C. J.,
Deutschbauer, A., and Keasling, J. D. (2019) Host engineering for improved valerolactam
production in Pseudomonas putida. *BioRxiv*.
(16) Zhang, M., Gao, C., Guo, X., Guo, S., Kang, Z., Xiao, D., Yan, J., Tao, F., Zhang, W.,
Dong, W., Liu, P., Yang, C., Ma, C., and Xu, P. (2018) Increased glutarate production by
blocking the glutaryl-CoA dehydrogenation pathway and a catabolic pathway involving L-2-
hydroxyglutarate. *Nat. Commun. 9*, 2114.
(17) Zhang, J., Barajas, J. F., Burdu, M., Wang, G., Baidoo, E. E., and Keasling, J. D. (2017)
Application of an Acyl-CoA Ligase from Streptomyces aizunensis for Lactam Biosynthesis. *ACS*
Synth. Biol. 6, 884–890.
(18) Thompson, M. G., Blake-Hedges, J. M., Cruz-Morales, P., Barajas, J. F., Curran, S. C.,
Eiben, C. B., Harris, N. C., Benites, V. T., Gin, J. W., Sharpless, W. A., Twigg, F. F., Skyrud,
W., Krishna, R. N., Pereira, J. H., Baidoo, E. E. K., Petzold, C. J., Adams, P. D., Arkin, A. P.,
Deutschbauer, A. M., and Keasling, J. D. (2019) Massively Parallel Fitness Profiling Reveals
Multiple Novel Enzymes in Pseudomonas putida Lysine Metabolism. *MBio 10*.
(19) Madhuri Indurthi, S., Chou, H.-T., and Lu, C.-D. (2016) Molecular characterization of lysR-
lysXE, gcdR-gcdHG and amaR-amaAB operons for lysine export and catabolism: a
comprehensive lysine catabolic network in Pseudomonas aeruginosa PAO1. *Microbiology*
(Reading, Engl) 162, 876–888.
(20) Zhang, M., Kang, Z., Guo, X., Guo, S., Xiao, D., Liu, Y., Ma, C., Gao, C., and Xu, P.
(2019) Regulation of Glutarate Catabolism by GntR Family Regulator CsiR and LysR Family
Regulator GcdR in Pseudomonas putida KT2440. *MBio 10*.
(21) Knorr, S., Sinn, M., Galetskiy, D., Williams, R. M., Wang, C., Müller, N., Mayans, O.,
Schleheck, D., and Hartig, J. S. (2018) Widespread bacterial lysine degradation proceeding via
glutarate and L-2-hydroxyglutarate. *Nat. Commun. 9*, 5071.
(22) Bailey, T. L., Williams, N., Misleh, C., and Li, W. W. (2006) MEME: discovering and
analyzing DNA and protein sequence motifs. *Nucleic Acids Res. 34*, W369-73.
(23) Jain, D. (2015) Allosteric control of transcription in GntR family of transcription regulators:
A structural overview. *IUBMB Life 67*, 556–563.
(24) Anderson, J. C., Dueber, J. E., Leguia, M., Wu, G. C., Goler, J. A., Arkin, A. P., and
Keasling, J. D. (2010) BglBricks: A flexible standard for biological part assembly. *J. Biol. Eng.*

4, 1.

(25) Siedler, S., Stahlhut, S. G., Malla, S., Maury, J., and Neves, A. R. (2014) Novel biosensors
based on flavonoid-responsive transcriptional regulators introduced into Escherichia coli. *Metab.*
Eng. 21, 2–8.

(26) Hastings, W. K. (1970) Monte Carlo sampling methods using Markov chains and their
applications. *Biometrika 57*, 97–109.

(27) Price, M. N., Wetmore, K. M., Waters, R. J., Callaghan, M., Ray, J., Liu, H., Kuehl, J. V.,
Melnyk, R. A., Lamson, J. S., Suh, Y., Carlson, H. K., Esquivel, Z., Sadeeshkumar, H.,
Chakraborty, R., Zane, G. M., Rubin, B. E., Wall, J. D., Visel, A., Bristow, J., Blow, M. J., and
Deutschbauer, A. M. (2018) Mutant phenotypes for thousands of bacterial genes of unknown
function. *Nature 557*, 503–509.

(28) Sagawa, S., Price, M. N., Deutschbauer, A. M., and Arkin, A. P. (2017) Validating
regulatory predictions from diverse bacteria with mutant fitness data. *PLoS ONE 12*, e0178258.

(29) Nitta, K. R., Jolma, A., Yin, Y., Morgunova, E., Kivioja, T., Akhtar, J., Hens, K., Toivonen,
J., Deplancke, B., Furlong, E. E. M., and Taipale, J. (2015) Conservation of transcription factor
binding specificities across 600 million years of bilateria evolution. *elife 4*.

(30) Suvorova, I. A., Korostelev, Y. D., and Gelfand, M. S. (2015) GntR Family of Bacterial
Transcription Factors and Their DNA Binding Motifs: Structure, Positioning and Co-Evolution.
PLoS ONE 10, e0132618.

(31) Mannan, A. A., Liu, D., Zhang, F., and Oyarzún, D. A. (2017) Fundamental Design
Principles for Transcription-Factor-Based Metabolite Biosensors. *ACS Synth. Biol. 6*, 1851–
1859.

(32) Ang, J., Harris, E., Hussey, B. J., Kil, R., and McMillen, D. R. (2013) Tuning response
curves for synthetic biology. *ACS Synth. Biol. 2*, 547–567.

(33) Way, J. C., Collins, J. J., Keasling, J. D., and Silver, P. A. (2014) Integrating biological
redesign: where synthetic biology came from and where it needs to go. *Cell 157*, 151–161.

(34) Wehrs, M., Tanjore, D., Eng, T., Lievense, J., Pray, T. R., and Mukhopadhyay, A. (2019)
Engineering Robust Production Microbes for Large-Scale Cultivation. *Trends Microbiol. 27*,
524–537.

(35) Revelles, O., Espinosa-Urgel, M., Molin, S., and Ramos, J. L. (2004) The davDT operon of
Pseudomonas putida, involved in lysine catabolism, is induced in response to the pathway
intermediate delta-aminovaleric acid. *J. Bacteriol. 186*, 3439–3446.

(36) Park, S. J., Kim, E. Y., Noh, W., Park, H. M., Oh, Y. H., Lee, S. H., Song, B. K., Jegal, J.,
and Lee, S. Y. (2013) Metabolic engineering of Escherichia coli for the production of 5-
aminovalerate and glutarate as C5 platform chemicals. *Metab. Eng. 16*, 42–47.

(37) Rohles, C. M., Gießelmann, G., Kohlstedt, M., Wittmann, C., and Becker, J. (2016) Systems
metabolic engineering of Corynebacterium glutamicum for the production of the carbon-5
platform chemicals 5-aminovalerate and glutarate. *Microb. Cell Fact. 15*, 154.

(38) Hagen, A., Poust, S., Rond, T. de, Fortman, J. L., Katz, L., Petzold, C. J., and Keasling, J.
D. (2016) Engineering a polyketide synthase for in vitro production of adipic acid. *ACS Synth.*

Biol. 5, 21–27.

(39) LaBauve, A. E., and Wargo, M. J. (2012) Growth and laboratory maintenance of Pseudomonas aeruginosa. *Curr. Protoc. Microbiol. Chapter 6*, Unit 6E.1.

(40) Ham, T. S., Dmytriv, Z., Plahar, H., Chen, J., Hillson, N. J., and Keasling, J. D. (2012) Design, implementation and practice of JBEI-ICE: an open source biological part registry platform and tools. *Nucleic Acids Res. 40*, e141.

(41) Chen, J., Densmore, D., Ham, T. S., Keasling, J. D., and Hillson, N. J. (2012) DeviceEditor visual biological CAD canvas. *J. Biol. Eng. 6*, 1.

(42) Hillson, N. J., Rosengarten, R. D., and Keasling, J. D. (2012) j5 DNA assembly design automation software. *ACS Synth. Biol. 1*, 14–21.

(43) Gibson, D. G., Young, L., Chuang, R.-Y., Venter, J. C., Hutchison, C. A., and Smith, H. O. (2009) Enzymatic assembly of DNA molecules up to several hundred kilobases. *Nat. Methods 6*, 343–345.

(44) Engler, C., Kandzia, R., and Marillonnet, S. (2008) A one pot, one step, precision cloning method with high throughput capability. *PLoS ONE 3*, e3647.

(45) Jackson, D. R., Tu, S. S., Nguyen, M., Barajas, J. F., Schaub, A. J., Krug, D., Pistorius, D., Luo, R., Müller, R., and Tsai, S.-C. (2016) Structural Insights into Anthranilate Priming during Type II Polyketide Biosynthesis. *ACS Chem. Biol. 11*, 95–103.

(46) Thompson, M. G., Sedaghatian, N., Barajas, J. F., Wehrs, M., Bailey, C. B., Kaplan, N., Hillson, N. J., Mukhopadhyay, A., and Keasling, J. D. (2018) Isolation and characterization of novel mutations in the pSC101 origin that increase copy number. *Sci. Rep. 8*, 1590.

(47) George, K. W., Thompson, M. G., Kim, J., Baidoo, E. E. K., Wang, G., Benites, V. T., Petzold, C. J., Chan, L. J. G., Yilmaz, S., Turhanen, P., Adams, P. D., Keasling, J. D., and Lee, T. S. (2018) Integrated analysis of isopentenyl pyrophosphate (IPP) toxicity in isoprenoid-producing Escherichia coli. *Metab. Eng. 47*, 60–72.

(48) Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A., and Jermiin, L. S. (2017) ModelFinder: fast model selection for accurate phylogenetic estimates. *Nat. Methods 14*, 587–589.

(49) Cruz-Morales, P., Ramos-Aboites, H. E., Licona-Cassani, C., Selem-Mójica, N., Mejía-Ponce, P. M., Souza-Saldívar, V., and Barona-Gómez, F. (2017) Actinobacteria phylogenomics, selective isolation from an iron oligotrophic environment and siderophore functional characterization, unveil new desferrioxamine traits. *FEMS Microbiol. Ecol. 93*.

(50) Grant, S. G., Jessee, J., Bloom, F. R., and Hanahan, D. (1990) Differential plasmid rescue from transgenic mouse DNAs into Escherichia coli methylation-restriction mutants. *Proc Natl Acad Sci USA 87*, 4645–4649.

(51) Bi, C., Su, P., Müller, J., Yeh, Y.-C., Chhabra, S. R., Beller, H. R., Singer, S. W., and Hillson, N. J. (2013) Development of a broad-host synthetic biology toolbox for Ralstonia eutropha and its application to engineering hydrocarbon biofuel production. *Microb. Cell Fact. 12*, 107.

(52) Shanks, R. M. Q., Kadouri, D. E., MacEachran, D. P., and O'Toole, G. A. (2009) New yeast

671 recombineering tools for bacteria. *Plasmid 62*, 88–97.

672 (53) Lee, T. S., Krupa, R. A., Zhang, F., Hajimorad, M., Holtz, W. J., Prasad, N., Lee, S. K., and

673 Keasling, J. D. (2011) BglBrick vectors and datasheets: A synthetic biology platform for gene

674 expression. *J. Biol. Eng. 5*, 12.

For Table of Contents Use Only

Robust characterization of two distinct glutarate sensing transcription factors of *Pseudomonas*

putida L-lysine metabolism

Mitchell G. Thompson[1,2,3], Zak Costello[1,2,4], Niklas F. C. Hummel[1,7], Pablo Cruz-Morales[1,2,5],

Jacquelyn M. Blake-Hedges[1,2,6],Rohith N. Krishna[1,2,6], Will Skyrud[6], Allison N. Pearson[1,2],

Matthew R. Incha[1,2,3], Patrick M. Shih[1,7], Hector Garcia-Martin[1,2,4,8], Jay D. Keasling[1,2,9,10,11,12]