1 Bottleneck revisited: increased adaptive variation despite reduced overall
2 genetic diversity in a rapidly adapting invader
3

4 Daniel Selechnik [1,2*], Mark F. Richardson [3,4], Richard Shine [1], Jayna DeVore [1], Simon
5 Ducatez [1], and Lee A. Rollins [2]
6
[1] School of Life and Environmental Sciences (SOLES), University of Sydney 2006
[2] Evolution & Ecology Research Centre, School of Biological, Earth and Environmental
Sciences, University of New South Wales, Sydney, NSW 2052
[3] Deakin Genomics Centre, School of Life and Environmental Sciences, Deakin University,
Locked Bag 20000, Geelong, VIC, Australia 3216
[4] Centre for Integrative Ecology, School of Life and Environmental Sciences, Deakin
University, Locked Bag 20000, Geelong, VIC, Australia 3216
[*] Correspondence: Dan Selechnik, Phone: +61 452 304 787; E-mail:
danselechnik@gmail.com

16
17

Abstract

Invasive species often exhibit rapid evolution following introduction, despite genetic

bottlenecks that may result from small numbers of founders; however, some invasions may

not fit this "genetic paradox." The invasive cane toad (*Rhinella marina*) displays high

phenotypic variation across its environmentally heterogeneous, introduced Australian range.

Here, we used three genome-wide datasets to characterize population structure and genetic

diversity in invasive toads: 1) RNA-Seq data from spleen sampled in the native range (French

Guiana), and introduced populations from Hawai'i (the source of the Australian introduction)

and Australia; 2) RNA-Seq data from brains sampled more extensively in Hawai'i and

Australia; and 3) previously published RADSeq data from transects across Australia. We

found that toads form three genetic clusters: (1) native range, (2) Hawai'i and long-

established areas near introduction sites in Australia, and (3) more recently established

northern Australian sites. In addition to strong divergence between native and invasive

populations, we find evidence for a reduction in genetic diversity following introduction.

However, we do not see this reduction in loci putatively under selection, suggesting that

genetic diversity may have been maintained at ecologically relevant traits, or that mutation

rates were high enough to maintain adaptive potential. Nonetheless, cane toads encounter

novel environmental challenges in Australia and appear to respond to selection across

environmental breaks; the transition between genetic clusters occurs at a point along the

invasion transect where temperature rises and rainfall decreases. We identify loci known to

be involved in resistance to heat and dehydration that show evidence of selection in

Australian toads. Despite well-known predictions regarding genetic drift and spatial sorting

during invasion, this study highlights that natural selection occurs rapidly and plays a vital

role in shaping the structure of invasive populations.

Introduction

The genetic paradox of invasion (Allendorf, 2003) describes a phenomenon that challenges widespread evidence of the relationship between genetic diversity and adaptive potential. High genetic diversity within a population is beneficial because it likely underlies phenotypic variation, allowing the population to respond to selection imposed by environmental change (Frankham, 2005; Reed & Frankham, 2003). Furthermore, a greater number of alleles confers an increased frequency of heterozygosity, which is often associated with population fitness (Reed & Frankham, 2003). Small or isolated populations with low genetic diversity have been shown to suffer declines due to inbreeding depression and the associated reduction of individual fitness (Blomqvist, Pauliny, Larsson, & Flodin, 2010; Madsen, Shine, Olsson, & Wittzell, 1999; Westemeier et al., 1998). Conservation efforts to salvage such populations by introducing individuals from allopatric populations (thereby introducing new alleles; "genetic rescue") have been successful, suggesting that the maintenance of genetic diversity can be crucial for population viability (Madsen et al., 1999; Westemeier et al., 1998).

Despite the fact that invasive populations are thought to undergo genetic bottlenecks due to the translocation of a small number of founders from their native range to an introduced range (Allendorf, 2003; Barrett & Kohn, 1991), invasive species are also characterized by their ability to establish and spread in their introduced ranges. Invasion success is commonly linked to rapid evolution, including adaptation to novel environmental conditions over short timescales (Franks & Munshi-South, 2014; Gao, Li, & Zhan, 2018; Leydet, Grupstra, Coma, Ribes, & Hellberg, 2018; Whitney & Gabler, 2008). Additionally, some invaders exhibit novel phenotypic traits that enhance invasive potential, such as increased growth and dispersal rates (White, Perkins, Heckel, & Searle, 2013; Whitney & Gabler, 2008). There are many examples of evolutionary change during invasion without high levels of genetic diversity (L.A. Rollins, Moles, & Lam, 2013).

Although low genetic diversity may limit the ability of an invasive population to

respond to natural selection, rapid evolution can also occur through non-adaptive processes:

(1) genetic drift may occur on range edges, reducing genetic diversity across an introduced

range (L. A. Rollins, Woolnough, Wilton, Sinclair, & Sherwin, 2009); (2) due to spatial

sorting, the invasion front may be inhabited exclusively by the individuals with the highest

dispersal rates (even if they are not the fittest) because they have arrived first and can only

breed with each other (Shine, Brown, & Phillips, 2011), resulting in a geographic separation

of phenotypes (C.M. Hudson, G. P. Brown, & R. Shine, 2016; Shine et al., 2011); (3)

admixture or hybridization may occur between individuals from different introductions or

sources (Mader, Castro, Bonatto, & de Freitas, 2016).

It has recently been suggested that the genetic paradox of invasion may be rare; some

invasive populations do not suffer a reduction in genetic diversity during introduction, and

others do not face novel adaptive challenges in their introduced ranges (A. Estoup et al.,

2016). Furthermore, some invasive systems demonstrate a 'spurious' paradox due to

inadequate estimation of genetic diversity (i.e. too few markers), or to maintenance of genetic

diversity only at ecologically relevant traits, or to a reduction in genetic diversity resulting

from natural selection rather than from genetic bottlenecks (A. Estoup et al., 2016). These

ideas can be tested with genome-wide data (Wellband, Pettitt-Wade, Fisk, & Heath, 2018;

Willoughby, Harder, Tennessen, Scribner, & Christie, 2018).

The Australian cane toad (*Rhinella marina*) is an invader that has exhibited rapid

evolution despite apparently low genetic diversity (L. A. Rollins, Richardson, & Shine,

2015). Cane toads were serially translocated from French Guiana and Guyana to Martinique

and Barbados, to Jamaica, to Puerto Rico, to Hawai'i before 101 founders were introduced

from the Hawai'ian island of Oahu to Queensland (QLD), Australia in 1935 (Turvey, 2009).

Toads have since spread westward through the Northern Territory (NT) into Western

Australia (WA; Fig 1). Prior to reaching Australia, toads were exposed to tropical

environments with relatively high mean annual rainfall (3200 mm in French Guiana, 1200-

2000 mm in Hawai'i) and high temperatures (26°C in French Guiana, 22-24°C in Hawaii)

(Hijmans, 2015). However, the Australian range is heterogeneous in several environmental

factors; on average, sites in QLD are more similar to the native range with respect to aridity,

receiving more annual rainfall than sites in the NT and WA (2000-3000 mm in QLD, 400-

1000 mm in NT and WA; Fig S1A), and have lower annual mean temperatures (21-24°C in

QLD, 24-27°C in NT and WA; Fig S1B) (Bureau of Meteorology, 2018). Heritable

phenotypic differences in behavior (J. Gruber, Brown, Whiting, & Shine, 2017), thermal

performance (G. K. Kosmala, Brown, Christian, Hudson, & Shine, 2018), morphology (C. M.

Hudson, G. P. Brown, & R. Shine, 2016; Hudson, Brown, Stuart, & Shine, 2018), and

immune function (Brown, Phillips, Dubey, & Shine, 2015c) have been documented in cane

toads at different localities across the Australian range. Nonetheless, surveys have reported

low levels of genetic diversity, based on microsatellites (Leblois, Rousset, Tikel, Moritz, &

Estoup, 2000) and MHC (Lillie, Shine, & Belov, 2014), and only one mitochondrial

haplotype in the *ND3* gene of 31 individuals sequenced from Hawai'i and Australia (Slade &

Moritz, 1998).

Here, we assessed genetic diversity and population structure using single nucleotide

polymorphisms (SNPs) identified in RNA-Seq data from samples originating from French

Guiana (the "native range"), Hawai'i (the "source"), QLD (the "range core"), NT

("intermediate" areas), and WA (the "invasion front") to test whether the genetic paradox is

evident in the cane toad invasion, and to investigate the evolutionary dynamics during

introduction. We predicted high divergence between native and invasive populations due to a

combination of genetic drift, selection, and spatial sorting, but little genetic differentiation

within invasion phases due to a putative lack of standing genetic diversity. Because of the

increase in aridity at intermediate areas and the invasion front, we predicted that loci involved

in thermal tolerance would be under selection, which may underlie the success that toads

have had dispersing through these areas.

RNA-Seq data provide genome-wide information, but are limited to transcribed

regions of the genome (Davey & Blaxter, 2011). RNA-Seq data may therefore contain

variants that exhibit strong signals for both demography and selection. Conversely, reduced

representation sequencing approaches such as restriction site-associated DNA sequencing

(RADSeq) provide genome-wide information on a subset of all coding and non-coding

sequences (only SNPs near restriction enzyme cleavage sites are detected); because non-

coding sequences make up most of the genome, SNPs detected using this technique are more

likely to represent neutral variation (Wang, Gerstein, & Snyder, 2009). To focus on how

selection may shape genomic diversity in invasive toads, we compared the results from our

RNA-Seq data to those from a publicly available RADSeq dataset (Trumbo et al., 2016) that

we reanalyzed.

Materials and Methods

Sample collection, RNA extraction, and sequencing

We collected samples from French Guiana, the Hawai'ian Islands and across Australia (Fig 1;

Table S1). We excised whole spleen tissue (sample sizes in Table S1A) and whole brain

tissue (sample sizes in Table S1B) from female toads immediately after euthanasia, and

preserved samples in RNAlater (QIAGEN, USA) at -20ºC for a maximum of one month,

after which they were drained and transferred to a -80°C freezer for long-term storage.

We conducted RNA extractions using the RNeasy Lipid Tissue Mini Kit (QIAGEN, USA)

following the manufacturer's instructions, with an additional genomic DNA removal step

using on-column RNase-free DNase treatment (QIAGEN, USA). We quantified the total

RNA extracted using a Qubit RNA HS assay on a Qubit 3.0 fluorometer (Life Technologies,

USA). Extracts were stored at -80∘C until sequencing (Macrogen Inc., ROK). mRNA

libraries were constructed using the TruSeq mRNA v2 sample kit (Illumina Inc., USA),

which included a 300bp selection step. In total, we sequenced 118 individually barcoded

samples (46 spleen and 72 brain) across 5 lanes of Illumina HiSeq 2500. Capture of mRNA

was performed using the oligo dT method, and size selection parameter choices were made

according to the HiSeq2500 manufacturer's protocol. Overall, this generated 709 million

(spleen) and 1.74 billion (brain) paired-end 2 x 125-bp reads. Raw sequence reads are

available as FASTQ files in the NCBI short read archive (SRA) under the BioProject

Accession PRJNA510261 (spleen data from French Guiana and Hawai'i), PRJNA395127

(spleen data from Australia), and PRJNA479937 (all brain data).

170
171

Data pre-processing and alignment

First, we examined per base raw sequence read quality (Phred scores) and GC content, and

checked for the presence of adapter sequences for each sample using FastQC v0.11.5 (S.

Andrews, 2010). We then processed raw reads (FASTQ format) from each sample with

Trimmomatic v0.35 (Bolger, Lohse, & Usadel, 2014), using the following parameters:

ILLUMINACLIP:TruSeq3-PE.fa:2:30:10:4 SLIDINGWINDOW:5:30 AVGQUAL:30

MINLEN:36. This removed any adaptor sequences, trimmed any set of five contiguous bases

with an average Phred score below 30, and removed any read with an average Phred score

below 30 or sequence length below 36 bp.

As a reference, we used the annotated *R. marina* transcriptome (Richardson et al.,

2018), which was constructed from brain, spleen, muscle, liver, ovary, testes, and tadpole

tissues. We conducted per sample alignments of reads (FASTQ files) to the reference using

STAR v2.5.0a (Dobin et al., 2013) in basic two-pass mode with default parameters, a

runRNGseed of 777, and specifying binary alignment map (BAM) alignment outputs. As

STAR-generated BAM files lack read groups (identifiers for reads that specify the individual

that they come from and the platform that was used to sequence them), we added them to our

BAM files using the AddOrReplaceReadGroups tool in Picard Tools (Institute, 2018). To

avoid making incorrect variant calls, we removed duplicate reads (reads that map to the same

position of the same gene in the same individual) using the MarkDuplicates tool in Picard

Tools. To split reads containing an N (region of the reference that is skipped in a read) into

individual exon segments, we used the SplitNCigarReads tool in the Genome Analysis

Toolkit (GATK) v3.8.0 (McKenna et al., 2010).

Variant calling and filtering

To call SNPs and insertion-deletions (indels), we used the HaplotypeCaller tool in GATK

(McKenna et al., 2010) on our alignment (BAM) files. This tool works by identifying regions

of the reference that show evidence of variation (called 'active regions'). Variants required a

minimum phred-scaled confidence of 20 to be called (marked as passing the quality filters)

and emitted (reported in the output), using the stand_call_conf and stand_emit_conf options.

201   To improve accuracy of SNP-calling, we performed this in 'GVCF' mode (which

generates one intermediate genomic variant call format file per sample). By using this mode,

we avoided missing SNPs at loci that match the reference in some but not all individuals. We

then used the GenotypeGVCFs tool to merge the GVCF files, re-calculate genotype

likelihoods at each SNP locus across all individuals, and re-genotype and re-annotate all SNP

loci. The results were written to one merged VCF file. Although we initially genotyped all

spleens and brains together, we discovered during downstream analyses that there was an

effect of tissue type on population assignment – even from the same individuals, and using

only SNPs from transcripts expressed in both tissues, spleen and brain samples were assigned

to separate populations). Thus, we genotyped spleens and brains separately, resulting in two

merged VCF files, and subsequently kept these separate for all downstream analyses. We

retained both datasets because each provides a unique benefit: the spleen dataset includes

native range samples, but the brain dataset has more extensive sampling of the invasive

populations.

215   Rather than following a random distribution, some SNPs are clustered. Clustered

SNPs may not be independent, and are thus filtered. We used the VariantFiltration tool to

identify and filter 'clusters' (sets of 3 SNPs that appear within a window of 35 bases) in each

of our merged VCF files. We also used this tool to filter variants with QualByDepth (QD;

variant confidence divided by the unfiltered depth of non-reference samples) less than 2.0,

depth of coverage (DP) less than 20.0, and allele frequency less than 0.05. We then subset our

VCF files to include only the variants that passed the filters we set in the VariantFiltration

step using the SelectVariants tool. This resulted in 803,489 SNPs from spleen data and

818,536 SNPs from brain data. We used bcftools (Li et al., 2009) to further filter the VCF

files, only retaining biallelic SNPs. We examined the results of filtering for minimum minor

allele frequency (min MAF) thresholds of 0.01 or 0.05, and several missing data tolerance

(MDT) thresholds (the maximum percentage of individuals in the dataset in which a genotype

for a locus can be absent without that locus being filtered out). Our population structure

results were consistent across min MAF and MDT thresholds, so we ultimately chose to filter

our data at min MAF = 0.05 and MDT = 0% (no missing data tolerated) because some

downstream analyses cannot handle missing data. These filtering steps reduced the number of

SNPs to 65,195 in spleen data and 35,842 in brain data (Table S1A-B).

Inference of population structure

We used PLINK (Purcell et al., 2007) to convert our VCF files to the Browser Extensible

Data (BED) format, which is readable to fastStructure. We then used fastStructure (Raj,

Stephens, & Pritchard, 2014) to infer population structure using a variational Bayesian

framework for calculating posterior distributions, and to identify the number of genetic

clusters in our dataset (K) using heuristic scores (Raj et al., 2014). We ran the structure.py ten

times each (K= 1 to 10). We then took the resulting meanQ files from fastStructure and

plotted them using the pophelper package (Francis, 2016) in R (Team, 2016).

In addition to using fastStructure for population assignment, we also performed a

Redundancy Analysis (RDA) using the vegan package (Oksanen et al., 2018) in R, which fits

genetic and environmental data to a multivariate linear regression, and then performs a

principal component analysis (PCA) on the fitted values. RDA visualizes both population

structure and the effects that environmental variables may have in shaping it. To do this, we

downloaded climatic data on French Guiana, Hawai'i, and Australia from the Bioclim

database (Hijmans, Cameron, Parra, Jones, & Jarvis, 2005) using the raster package

(Hijmans, 2015). Because these areas vary in aridity, we downloaded data on rainfall and

temperature; these data are averages of annual statistics over the period of 1970 to 2000.

Specifically, we downloaded: annual mean temperature, maximum temperature of the

warmest month, minimum temperature of the coldest month, annual precipitation,

precipitation of the wettest quarter, and precipitation of the driest quarter. We then used the

vcfR package (Knaus & Grünwald, 2017) to convert our VCF files to the GENLIGHT

format, which is readable to the vegan package.

Identification of candidate loci under selection

Loci that are under natural selection may have abnormally high F_{ST} values, causing them to

be outliers among all other loci in the transcriptome. Bayescan v2.1 (Foll & Gaggiotti, 2008)

detects loci with outlier F_{ST}s using the multinomial-Dirichlet model. We used PGDSpider

(Lischer & Excoffier, 2012) to convert our VCF files to the BAYESCAN format, and then

ran Bayescan to detect loci with outlier pairwise F_{ST} values between our three genetic clusters

in spleen data (native range toads versus source and core toads versus intermediate and

frontal toads) and our two genetic clusters in brain data (source and core toads versus

intermediate and frontal toads).

Loci under natural selection also may have allele frequencies that are associated with

environmental variables. We used the climatic data that we downloaded from the Bioclim

database (Hijmans et al., 2005) using the raster package (Hijmans, 2015) on rainfall during

the driest quarter and maximum temperature in the warmest month. We then used the lfmm

v2.0 package (Frichot & Francois, 2015) in R to perform a latent factor mixed model

(LFMM) to test the association between the allele frequencies of every locus and these two

environmental variables. We applied a Benjamini-Hochberg correction to all p-values from

the LFMM.

Because these scans reveal thousands of loci putatively under selection, and because

outlier tests have high false positive rates (Whitlock & Lotterhos, 2015), we took a more

conservative approach by cross-matching our list of F_{ST} outliers with each of our lists of

environmentally-associated loci. For individual investigation, we only looked at loci with

both an outlier F_{ST} value and an association with a putatively significant environmental

variable.

Evaluation of genetic differentiation and diversity

To quantify levels of genetic differentiation and diversity, we computed basic statistics in the

hierfstat (Goudet, 2005) and diveRsity (Keenan, McGinnity, Cross, Crozier, & Prodohl,

2013) packages in R (Team, 2016). We first used PGDSpider v2.1.1.3 (Lischer & Excoffier,

2012) to convert our VCF files to the FSTAT and GENEPOP formats, which are readable to

hierfstat and diveRsity, respectively. We then used these packages to calculate measures of

genetic differentiation, including global F_{ST} and pairwise F_{ST} (by genetic cluster, invasion

phase, and collection site), and genetic diversity, such as expected heterozygosity (He) and

rarefied allelic richness (AR). We also computed Shannon's Information Index (SI) with the

dartR package (B. Gruber, Unmack, Berry, & Georges, 2018). After calculating these

measures of diversity across all loci (N=65,195 from spleen data, N=35,842 from brain data),

we calculated the same measures in loci putatively under selection: those with outlier F_{ST}

values (N=648 from spleen data, N=203 from brain data) and those associated with

environmental variables (N=4179 from spleen data, N=530 from brain data). In spleen data,

this allowed us to investigate the hypothesis that genetic diversity is maintained at

ecologically relevant traits even if genome-wide diversity is lost. In brain data, this allowed

us to examine the effect of natural selection on genetic diversity within Hawai'i and

Australia. We used Kruskal-Wallis tests to assess the significance of the differences in

genetic diversity.

Annotation of SNPs

To visualize the types of genomic regions in which our SNPs lie, we used SnpEff (Cingolani

et al., 2012). SnpEff analyzes the information in a VCF file, annotates the SNPs, and

estimates their effects. We calculated the relative proportions of each type of SNP for both

tissue types using the full dataset of SNPs, and our two selection candidate datasets

(containing F_{ST} outlier loci associated with an environmental variable). We then used the stats

v3.5.0 R package to perform z-tests on the differences in proportions of SNP types between

the full and selection candidate datasets. We expected differences in the proportions of

certain types of SNPs, such as a lower proportion of synonymous variants in the selection

candidates because synonymous mutations do not result in a codon change (and thus are

unlikely to change protein function, i.e. phenotype).

Isolation by distance

To examine the effects of geographic distance on genetic distance across the Australian

range, we performed a Mantel test using ade4 v1.7-5 (Thioulouse & Dray, 2007). Because we

were focused on isolation by distance through range expansion after introduction (to

Australia), samples from Hawai'i and French Guiana were excluded; and because we

sampled brain tissue in a greater number of sites within Australia (as opposed to spleen

tissue; Table S1A-B), we only performed the Mantel test on brain data. For the geographic

distance matrix, we used the dist function in R (Team, 2016) to calculate the Euclidean

distances in geographic space between collection sites based on their coordinates. For the

genetic distance matrix, we used the collection site-based pairwise F_{ST} values generated from

hierfstat. We performed four Mantel tests, using: (1) all loci, (2) loci with outlier F_{ST} values,

(3) loci with outlier F_{ST} values and with an environmental association (temperature or

rainfall), (4) loci with outlier F_{ST} values but without an environmental association. A

significant result across all loci may suggest that the prominent driver of genetic structure is

genetic drift. However, clinal variation in allele frequencies may also result from spatial

sorting (particularly in loci underlying dispersal ability), or from selection associated with

environmental clines; distinguishing these evolutionary processes is difficult. Thus, we

performed the additional tests to attempt to separate the effects of genetic drift and spatial

sorting. Loci with outlier F_{ST} values are candidates for natural selection and spatial sorting;

thus, a significant result from only these loci would less likely be caused by genetic drift.

Outlier loci can be further separated into those with an environmental association, which are

likely candidates for natural selection driven by environmental variables, and those without

an environmental association, which are possible candidates for spatial sorting. If spatial

sorting is indeed influencing genetic differentiation, then we may expect to see different

patterns between these two groups of loci (e.g. linearity only in the outlier loci without an

environmental association).

Acquisition, processing, and analysis of RADSeq dataset

In addition to our own RNA-Seq dataset, we downloaded a publicly available RADSeq

dataset (Trumbo et al., 2016) to evaluate population structure in Australian cane toads. Unlike

RNA-Seq data, which consists of coding sequences only (Wang et al., 2009), RADSeq data

include SNPs from across the entire genome, but only around sites that are cleaved by a

selected set of restriction enzymes (Davey & Blaxter, 2011). This method allows us to

contrast the results of assessing loci potentially under 'tighter' selection (e.g., codon bias)

with those primarily constituting neutral variation (RADSeq). Individuals in the RADSeq

dataset were collected from QLD (range core towards intermediate areas, N=179) and the

border of NT/WA (intermediate areas towards the invasion front, N=441; Fig 1). However,

samples were lacking from a large part of the range between those regions; thus, identifying

geographic points of differentiation between genetic clusters may be unlikely using this

dataset. Furthermore, although there were many collection sites within these regions,

metadata with coordinates (or even location names) were unavailable. We downloaded raw

100-bp reads from NCBI short read archive (SRA) under the BioProject Accession

PRJNA328156. We then converted SRA files to the FASTQ format using the fastq-dump tool

in the SRA toolkit.

We used Stacks v2.0 (Catchen, Hohenlohe, Bassham, Amores, & Cresko, 2013) for

all RADSeq data processing. First, we used the process_radtags program to remove low

quality reads from the FASTQ files. Next, we used the denovo_map pipeline to perform a *de*

novo assembly for each sample, align matching DNA regions across samples (called

'stacks'), and call SNPs using a maximum likelihood framework. We allowed a maximum of

five base mismatches between stacks within an individual and three base mismatches

between stacks between individuals. This resulted in 499,623 SNPs. We then filtered the

results using the populations program, using one random SNP per locus and four

combinations of parameter choices (min MAF=0.05, MDT = 0.5; min MAF = 0.05, MDT =

0.2; min MAF = 0.01, MDT = 0.5; min MAF = 0.01, MDT = 0.2). Structure results were

consistent across min MAF values, but not MDT values: two distinct populations were only

detected using the more stringent MDT choice. Because population structure in this dataset is

relatively low, and because it is appropriate to use more stringent filtering choices when

analyzing datasets with low population structure (Selechnik et al., unpublished; Chapter 2),

we ultimately chose to filter this dataset with min MAF = 0.05 and MDT = 0.2. Furthermore,

using these filtering choices produced results that were most consistent with those reported

by Trumbo et al. (2016) and with those of our RNA-Seq dataset. Filtering reduced the

number of SNPs to 8,296. The results were written to a VCF file and a STRUCTURE file

(which could be read into fastStructure).

We ran fastStructure using the same methodology as we did for our RNA-Seq dataset.

In addition to the model-based fastStructure, we also performed a discriminant analysis of

principal components (DAPC) on this dataset using adegenet (Jombart & Ahmed, 2011).

DAPC is a multivariate approach that identifies the number of genetic clusters using K-means

of principal components and a Bayesian framework. We also converted the VCF to the

FSTAT and GENEPOP formats and computed the same basic statistics for the RADSeq

dataset as we did for our RNA-Seq datasets using hierfstat and diveRsity. Because collection

site coordinate metadata were unavailable for the RADSeq dataset, we did not perform a

Mantel test using these data.

Results

Population structure across the Hawai'ian-Australian invasion

We found three genetic clusters using spleen data (Fig 2A): French Guiana (native range)

formed its own genetic cluster, Hawai'i (source) clustered with QLD (core), and NT

(intermediate) clustered with WA (invasion front). In our brain dataset, population

differentiation seemed to align with environmental barriers (Fig 2B-C): Hawai'i (source)

clustered with coastal QLD (core), whereas inland QLD/NT (intermediate) clustered with

WA (invasion front). Inference of substructure within these two genetic clusters revealed that

source and core toads may further differentiate into two separate groups (Fig 2D); however,

intermediate and frontal toads remain one genetic cluster (Fig 2E).

RDA was mostly consistent with fastStructure for both spleen (Fig 3A) and brain (Fig

3B) data, except for the differentiation of source toads and core toads as two separate genetic

clusters. It also suggested that: (1) Source toads may have diverged from toads of other

clusters due to selection from rainfall during the driest quarter. (2) Core toads may have

diverged from toads of other clusters due to selection from rainfall during the wettest quarter,

mean annual rainfall, and minimum temperature during the coolest month. (3) Intermediate

and frontal toads may have diverged from toads of other clusters due to selection from

maximum temperature during the hottest month and mean annual temperature.

Tests for selection

Out of 65,195 SNPs from spleen data, 648 were identified as F_{ST} outliers, with a mean global

F_{ST} of 0.89. Mean pairwise F_{ST} values across outlier loci were 0.93 between the native and

source/core populations, 0.97 between the native and intermediate/front populations, and 0.34

between the source/core and intermediate/front populations. Also from the spleen dataset,

3088 SNPs were associated with maximum temperature during the warmest month and 1813

were associated with rainfall during the driest quarter. Of these loci, 772 were associated with

both environmental variables. We focused on SNPs that were detected both as F_{ST} outliers

and environmental correlates: there were 64 outlier SNPs associated with maximum

temperature and 47 outlier SNPs associated with rainfall in the driest quarter (including 18

outliers associated with both environmental variables).

Most of the 64 outlier SNPs associated with maximum temperature during the hottest

month (Table S2A) were in transcripts with functions such as metabolism, transcription

regulation, and immune function. There were also two SNPs in *HSP4*, a gene encoding heat

shock protein (HSP) 4 (70 kDa), which is involved in thermal tolerance (on exposure to heat

stress) and protein folding and unfolding (Consortium, 2017). Most of the 47 outlier SNPs

associated with rainfall during the driest quarter (Table S3A) were in transcripts with

functions such as cell signaling and immune function.

Out of 35,842 SNPs from brain data, 203 were identified as F_{ST} outliers, with a mean

global F_{ST} of 0.34. Mean pairwise F_{ST} was 0.50 between the source/core and

intermediate/front populations. Also from the brain dataset, 345 SNPs were associated with

maximum temperature during the warmest month and 194 were associated with rainfall

during the driest quarter. Nine of these loci were associated with both environmental

variables. Of SNPs that were detected both as F_{ST} outliers and environmental correlates, there

were 38 outlier SNPs associated with maximum temperature and 33 outlier SNPs associated

with rainfall in the driest quarter; no outliers were associated with both environmental

variables.

Most of the 38 outlier SNPs associated with maximum temperature during the hottest

month (Table S2B) were in genes involved in cell signaling, mitochondrial processes

(metabolism), or gene expression (Consortium, 2017). Similarly to the spleen data, there was

one SNP from brain data in *HSP4*. Another SNP was in *ARNT2*, a gene encoding aryl

hydrocarbon receptor nuclear translocator 2, a transcription factor involved in the

hypothalamic pituitary adrenal (HPA) axis and visual and renal function (Consortium, 2017).

Of the 33 outlier SNPs from brain data correlated with rainfall during the driest quarter

(Table S3B), 19 were in *MAGED2*, a gene encoding melanoma-associated antigen D2, which

is involved in renal sodium ion absorption (Consortium, 2017). Four were in *STXBP1*, a gene

encoding syntaxin-binding protein 1, which is involved in platelet aggregation and vesicle

fusion (Consortium, 2017). The rest were in genes generally involved in gene expression and

cell signaling (Consortium, 2017).

Evaluation of genetic differentiation, diversity, and isolation by distance

Our spleen data show strong divergence between the native and source/core populations

(pairwise $F_{ST} = 0.29$), and even stronger divergences between native and intermediate/frontal

populations (pairwise $F_{ST} = 0.32$). Differentiation between the source/core and

intermediate/frontal populations was much lower (pairwise $F_{ST} = 0.08$). Overall divergence

among natives and invaders was moderately high (global $F_{ST} = 0.18$)

In brain data, genetic differentiation between the source/core and intermediate/frontal

populations also was low (pairwise $F_{ST} = 0.07$). Differentiation between source and core

toads was lower when calculated with brain data as compared to spleen (pairwise $F_{ST} = 0.04$).

Generally, pairwise F_{ST} by invasion phase (Table S4A) and collection site (Table S4B)

calculated from brain data revealed that toads from areas that are more closely linked by

invasion history are less strongly differentiated. Because the brain data did not include native

range samples, overall divergence was estimated to be much lower than it was in the spleen

data (global $F_{ST} = 0.04$).

Diversity statistics from spleen data suggest that the native range population is only

slightly more genetically diverse than the source/core population (native: He = 0.27, AR =

1.74, SI = 0.40; source/core: He = 0.26, AR = 1.74, SI = 0.40; p = 5E-4 for He) but both are

more diverse than the intermediate/frontal population (He = 0.23, AR = 1.68, SI = 0.36; $p <$

2E-16 for all measures). However, we also estimated these measures by region (Table 1)

rather than by genetic cluster to detect changes in genetic diversity at a finer scale; these

calculations reveal a more obvious decline in genetic diversity from the native population to

all invasive collection sites.

Despite this reduction in genome-wide genetic diversity, subsets of loci putatively

under selection in the spleen dataset showed no loss of diversity during introduction (Table

1). Here there was an increase in diversity in the source/core population as compared to

native samples (native: He = 0.03, AR = 1.25, SI = 0.06; source/core: He = 0.09, AR = 1.37,

SI = 0.14; $p <$ 2E-16 for all measures). At these same loci, diversity was lowest in the

intermediate/front population (He = 0.02, AR = 1.14, SI = 0.04). These trends are also shown

in environmentally-associated loci (native: He = 0.17, AR = 1.53, SI = 0.25; source/core: He

= 0.37, AR = 1.99, SI = 0.55; intermediate/front: He = 0.23, AR = 1.81, SI = 0.35).

Genome-wide diversity estimates using brain data show similar patterns to spleen data

(source/core: He = 0.34, AR = 2.00, SI = 0.51; intermediate/front: He = 0.31, AR = 1.97, SI =

0.47; $p <$ 2E-16 for comparisons of all measures; Table 2). This reduction is greater in loci

with outlier F_{ST} values (source/core: He = 0.38, AR = 2.00, SI = 0.55; intermediate/front: He

= 0.15, AR = 1.70, SI = 0.24; $p <$ 2E-16 for comparisons of all measures) and in

environmentally-associated loci (source/core: He = 0.35, AR = 2.00, SI = 0.53;

intermediate/front: He = 0.08, AR = 1.57, SI = 0.15; $p <$ 2E-16 for comparisons of all

measures).

Geographic distance in collection sites and pairwise F_{ST} values by collection site (as

estimated with brain data) were significantly associated across all four of our Mantel tests

using: (1) all loci (p = 1E-3, r = 0.76; Fig 4A); (2) all loci with outlier F_{ST} values (p = 1E-3, r

= 0.75; Fig 4B); (3) only loci with outlier F_{ST} values and an environmental association (p =

1E-3, r = 0.62; Fig 4C); (4) only loci with outlier F_{ST} values but without an environmental

association (p = 1E-3, r = 0.77; Fig 4D).

SNP Annotations

Annotation of SNPs revealed that most of our loci were either missense, synonymous, or

found in the 3' untranslated region (UTR). A small percentage were found in the 5' UTR, and

the remaining loci involved the loss or gain of stop or start codons (Figs S2-S3). In the spleen

data, the temperature-associated outlier F_{ST} subset had a significantly higher proportion of

synonymous variants (p = 8E-3) than the full set of SNPs. The outlier loci within *HSP4* were

both synonymous variants. However, there were no significant differences in proportions

between the rainfall-associated outlier F_{ST} subset and the full set of SNPs.

In the brain data, there were no significant differences in proportions of SNP variant

types between the temperature-associated outlier F_{ST} subset and the full set. The locus within

HSP4 was a 3' UTR variant, and the locus within *ARNT2* was synonymous. The rainfall-

associated outlier F_{ST} subset had a significantly higher proportion of 3' UTR variants (p =

5E-3) and lower proportion of missense variants (p = 0.04) than did the full set. Eighteen of

the nineteen loci within *MAGED2* were 3' UTR variants (the last was missense), and two of

the four loci within *STXBP1* were missense (the other two were synonymous).

Population structure in Australian toads from the RADSeq experiment

Analysis of the RADSeq dataset through fastStructure and DAPC also showed two genetic

clusters within Australia (Fig 5A-B): QLD (core towards intermediate areas) in the first

group, and the border of NT/WA (intermediate areas towards the invasion front) in the

second group. There was little differentiation between these two groups (pairwise $F_{ST} = 0.09$)

or overall (global $F_{ST} = 0.04$). We found no evidence of significant substructure within either

of the two genetic clusters. The two groups were equal in allelic richness (AR = 2.00 for both

groups), but expected heterozygosity was higher in core toads than in intermediate/frontal

toads (core: He = 0.50; intermediate/front: He = 0.33).

Discussion

We predicted that genetic diversity would be significantly lower in invasive populations than

in native populations due to genetic bottlenecks occurring during serial introductions of cane

toads to Hawai'i and Australia. We also predicted strong divergence between native range

and invasive populations due to a combination of genetic drift, selection and spatial sorting.

In terms of population structure, we predicted low genetic structure within invasive

populations in Hawai'i and Australia resulting from putatively low levels of standing genetic

variation. Our analyses show evidence of a reduction in genetic diversity from native to

invasive populations across all loci. We found three genetic clusters: (1) native range toads,

(2) Hawai'ian source toads and eastern Australian range core toads, and (3) all toads from

further west in more recently colonized areas (intermediate and invasion front).

Differentiation is much higher between the native population and the invasive populations

than between the invasive populations. Loci identified as F_{ST} outliers had values greater than

0.9 between native and invasive populations, indicating that many of these loci are likely to

be fixed for different alleles in these populations. Genetic structure around the aridity barrier

suggest that natural selection is occurring in the western part of the Australian range.

However, the presence of isolation by distance across all loci also supports gradual genetic

drift across invasion phases; that this relationship is strengthened in the subset of F_{ST} outliers

without environmental associations may indicate the presence of spatial sorting. It seems

likely that all these evolutionary forces have shaped Australian invasive populations.

535   Evidence that invasive cane toads fit the genetic paradox paradigm is equivocal. We

did identify a reduction in overall genetic diversity, both in allelic richness and evenness (He

and SI). Previous studies on mtDNA (Slade & Moritz, 1998) and microsatellite (A. Estoup,

Wilson, Sullivan, Cornuet, & Moritz, 2001) data have also suggested losses in genetic

diversity from the native range to Hawai'i and Australia. Despite these findings, we did not

find this reduction in the subset of loci putatively under selection (loci with outlier F_{ST} values

or associations with environmental variables). This supports the prediction that some

invasions do not represent a genetic paradox, despite overall loss of genetic diversity and the

presence of novel adaptive challenges, because genetic diversity at ecologically relevant traits

is maintained by balancing selection (A. Estoup et al., 2016). However, diversity of this

subset of loci is not only maintained in the source/core population, it is higher than in the

native population. This may reflect admixture from multiple genetically-distinct

introductions; the Hawai'ian introduction was sourced from the Puerto Rican population,

which included introductions from French Guiana and Guyana (Turvey, 2009). Alternatively,

this may represent a genuine paradox; *de novo* mutation rates may have been high enough to

restore or even enhance adaptive potential in the source/core population (A. Estoup et al.,

2016). Although these toads have been in Hawai'i and Australia for less than 100 years, they

were separated from native toads in the early 1800s when toads were first introduced to

Caribbean islands. From the source/core population to the intermediate frontal population,

however, diversity of loci putatively under selection is highly reduced; we believe this is

because of directional selection, as outlined below.

Our two invasive genetic clusters (source/core and intermediate/front) diverge at the

transition from coastal to inland areas within Australia, in an area where temperatures

become higher and rainfall becomes scarcer, particularly during the dry season (Bureau of

Meteorology, 2018). This correlation suggests that these environmental variables may drive

population structure and that toads may be adapting to local conditions. Furthermore, RDA

indicates selection from temperature and rainfall may drive divergence between invasive

populations. Heterogeneous climates across introduced or expanding ranges have previously

been linked to genetic differentiation, possibly due to climatically-imposed selection (Leydet

et al., 2018; Weber & Schmid, 1998).

Diversity estimates from our brain data suggest a reduction in diversity from the

source/core population to the intermediate/frontal population in the full set of loci, but a

much greater reduction in diversity in the subset of loci putatively under selection. The

source (Hawai'i) and core (QLD) areas represent earlier phases of the invasion and

experience similar environmental conditions to those of the native range, whereas

intermediate (NT) and invasion front (WA) areas are more arid. Thus, the greater depletion in

diversity at loci putatively under selection across the Australian invasion may be due to the

effects of directional selection from the harsher climate at more recently colonized areas of

the Australian range (C. A. Andrews, 2010; A. Estoup et al., 2016).

Among our loci associated with temperature are several within a gene encoding

protein HSP 4 (70 kDa). HSPs protect protein folding during increased temperatures and

provide cells with enhanced thermal tolerance (Kiang & Tsokos, 1998), and expression of

HSP genes has been shown to underlie adaptive responses to environmental stress (Chen,

Feder, & Kang, 2018). Changes in HSP levels in response to thermal environment

manipulation vary between native and invasive cane toads, as well as between populations of

toads within Australia (G. K. Kosmala, Brown, & Shine, 2018). In our study, the SNPs

within *HSP4* included two synonymous variants (spleen data) and a 3' UTR variant (brain

data); 3' UTR variants are known to alter levels of protein functioning through differential

expression of genes (Skeeles, Fleming, Mahler, & Toland, 2013). In the case of HSPs,

adaptation may not necessarily mean a change in function, but rather a change in the

expression levels of proteins involved in the function, which could be caused by a 3' UTR

variant.

Aryl hydrocarbon receptor nuclear translocator 2 (from the *ARNT2* gene, also

associated with temperature), is a transcription factor that specifically acts on genes involved

in the HPA axis and visual and renal function. The HPA axis is stimulated by high

temperatures (Malmkvist et al., 2009), and the secretion of corticosterone can be modulated

adaptively in response to thermal change (Telemeco & Addis, 2014). Elevation of

corticosterone secretion in toads increases the rate of desiccation, thus making it maladaptive

in arid environments (Jessop, Letnic, Webb, & Dempster, 2013). Corticosterone levels in

Brazilian and Australian toads in response to manipulation of thermal environment display

similar patterns to those of HSPs (G. K. Kosmala, Brown, & Shine, 2018). Selection on loci

potentially affecting corticosterone secretion by affecting the HPA is consistent with these

results, although the SNP we investigated in *ARNT2* is synonymous.

More than half (19 of 33) of the loci under selection associated with rainfall (in brain

data) were found in *MAGED2* (encodes melanoma-associated antigen D2). This protein

regulates the expression and localization of two co-transporters that facilitate renal sodium

ion absorption, preventing excess loss of water and solutes via reabsorption through the

kidneys (Greger, 2000). Similarly to the SNP within *HSP4*, 18 of 19 SNPs within *MAGED2*

were 3' UTR variants, which can affect gene expression (Skeeles et al., 2013). The high

number of SNPs found in the 3' UTR of *MAGED2* is likely responsible for the difference in

proportions of variant types between the full dataset and the rainfall-associated outlier

dataset. Four SNPs (including missence variants) were found in *STXBP1* (encodes syntaxin-

binding protein 1, which is involved in vesicle fusion and blood clotting). Dehydration is

known to increase blood clotting rate (El-Sabban, Fahim, Al Homsi, & Singh, 1996), and

excessive blood clotting can be harmful to the heart, brain, and limbs (PubmedHealth, 2014).

Evolved changes in the rates of renal sodium absorption and blood clotting may allow toads

from intermediate and invasion front areas to survive in drier environments. The remaining

SNPs associated with rainfall or temperature were in genes generally involved in gene

expression or signal transduction; these genes may facilitate expression of (or signaling to)

the genes we have discussed, or may have independent functions in cellular maintenance.

Invasive species have been shown to rapidly adapt phenotypically to heterogeneous

climatic conditions, enabling range expansion (Colautti & Barrett, 2013). Some invaders

tolerate higher temperatures (Braby & Somero, 2006) and water loss (Godoy, Lemos-Filho,

& Valladares, 2011) better than do related indigenous taxa (Zerebecki & Sorte, 2011). There

is evidence of this in cane toads: under dehydrating conditions, wild-caught individuals from

Australia exhibit better locomotor performance than do conspecifics from Hawai'i and the

native range (G. Kosmala, Christian, Brown, & Shine, 2017), and within Australia, toads

from semi-arid areas (i.e. NT) exhibit better locomotor performance than do conspecifics

from wetter areas (i.e. coastal QLD) (G. Kosmala et al., 2017; Tingley, Greenlees, & Shine,

2012). This trait is heritable; captive-bred toads with parental origins from hotter areas

(northwestern Australia) outperform those with parental origins from cooler areas

(northeastern Australia) at high (but not low) temperatures in a common-garden setting (G.

K. Kosmala, Brown, Christian, et al., 2018). Coupled with the previously shown heritable

phenotypic patterns, our results suggest that intermediate and frontal toads in Australia may

be evolving enhanced thermal tolerance, thereby facilitating their continued westward range

expansion (Szucs et al., 2017). However, manipulative experiments are required to directly

test this hypothesis.

632   Although our data suggest that invasive toads may be responding to localized natural

selection, our data also support the presence of other evolutionary forces. We found evidence

of isolation by distance (IBD) in Australia using the full SNP dataset, which is likely to be

caused by genetic drift. However, loci under strong selection are less likely to show evidence

of IBD. Unsurprisingly, when we plot genetic vs. geographic distance in Australia, using loci

putatively associated with environmental selection, we see a clear discontinuity in these data

(Fig 4C). Other evolutionary forces may be more difficult to detect in these data. Spatial

sorting is the separation of individuals within a range-expanding population along phases of

their expansion based on their dispersive capabilities, and was first characterized in

Australian cane toads (Shine et al., 2011). Spatial sorting is expected to result in clinal

variation in dispersal-related allele frequencies along the range, so we might expect that the

relationship between genetic and geographic distance using loci underlying dispersal-related

characteristics would show a similar pattern to that of genetic drift. In this study, when we

examine this relationship only using loci identified as under selection in this introduction, but

not associated with environmental selection, there is a gradual decay of the relationship

between genetic and geographic distance (Fig 4D). Although we acknowledge that these loci

may be influenced by drift to some degree, it seems plausible that because they were

identified as F_{ST} outliers, they are likely to be influenced by other evolutionary forces,

including spatial sorting.

651   Our population structure analyses using the RADSeq dataset (Trumbo et al., 2016)

were similar to those from our RNA-Seq dataset; both showed two genetic clusters within

Australia. However, unlike the RNA-Seq results (which show a divide occurring in inland

QLD, where temperature increases and rainfall decreases), the RADSeq results cluster all

QLD toads in the first group, and NT/WA toads in the second group. In the RADSeq dataset,

the spatial location of the transition between the two groups is difficult to pinpoint due to the

absence of intermediate sampling sites. The difference in group divisions between the two

datasets may be because selection from environmental variables causes differentiation

between toads from coastal and inland QLD (as seen in the RNA-Seq data), whereas

demographic processes such as gene flow may lessen differentiation between them (as seen

in the RADSeq data). Alternatively, toads from coastal and inland QLD may cluster together

in the RADSeq data because the sampling sites in that study were south of those in our study,

and the environmental conditions of coastal and inland sites are more similar in that area

(Hijmans et al., 2005). Estimates of diversity from the RADSeq dataset were slightly higher

than those from the RNA-Seq dataset; this may be because adaptive variation is expected to

be lower than neutral variation due to evolutionary constraints (C. A. Andrews, 2010).

667   In conclusion, it appears that serial introductions have led to a reduction in genetic

diversity of invasive cane toads. Nonetheless, cane toads do indeed face novel adaptive

challenges in their introduced ranges, and are able to respond genetically to natural selection,

particularly in the harsh environmental conditions of northwestern Australia. The ability of

toads to adapt to these conditions may reflect the maintenance of diversity at ecologically

relevant loci, or it may reflect sufficiently high mutation rates to bolster adaptive potential;

thus, it remains unclear whether the toad invasion represents a true genetic paradox. We have

found evidence of natural selection, genetic drift, and spatial sorting in the Australian

invasion, but the impacts of these forces require further investigation to tease apart.

Manipulative experiments investigating the candidates that we have identified may be

especially informative. Additionally, genetic changes may not be the only mechanisms by

which adaptation occurs; studies on plasticity and epigenetic changes may also be useful for

uncovering the mysteries of rapid evolution.

Acknowledgements

This work was supported by the Australian Research Council (FL120100074, DE150101393)

and the Equity Trustees Charitable Foundation (Holsworth Wildlife Research Endowment).

We thank Olivier Francois and Natalie Hofmeister for their input during discussions about

our analyses. We thank Cam Hudson, Joachim Ehlenz, Serena Lam, and Chris Jolly for their

assistance with sample collection. We thank Andrea West for her assistance during RNA

extractions. All procedures involving live animals were approved by the University of

Sydney Animal Care and Ethics Committee.

References

Allendorf, F. W. (2003). Introduction: Population Biology, Evolution, and Control of Invasive Species. *Conservation Biology, 17*(1), 24 - 30.

Andrews, C. A. (2010). Natural Selection, Genetic Drift, and Gene Flow Do Not Act in Isolation in Natural Populations. *Nature Education Knowledge, 3*(10), 5.

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data. Retrieved from http://www.bioinformatics.babraham.ac.uk/projects/fastqc

Barrett, S. C. H., & Kohn, J. R. (1991). *Genetics and conservation of rare plants* D. A. Falk & K. E. Holsinger (Eds.), *Genetic and evolutionary consequences of small population size in plants: implications for conservation* (pp. 3 - 30).

Blomqvist, D., Pauliny, A., Larsson, M., & Flodin, L. A. (2010). Trapped in the extinction vortex? Strong genetic effects in a declining vertebrate population. *BMC Evolutionary Biology, 10*, 33.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics, 30*(15), 2114-2120. doi:10.1093/bioinformatics/btu170

Braby, C. E., & Somero, G. N. (2006). Following the heart: temperature and salinity effects on heart rate in native and invasive species of blue mussels (genus Mytilus). *Journal of Experimental Biology, 209*, 2554 - 2566.

Brown, G. P., Phillips, B. L., Dubey, S., & Shine, R. (2015c). Invader immunology: invasion history alters immune system function in cane toads (Rhinella marina) in tropical Australia. *Ecol Lett, 18*(1), 57-65. doi:10.1111/ele.12390

Bureau of Meteorology, A. G. (2018). Climate Data Online. Retrieved from http://www.bom.gov.au/

Catchen, J., Hohenlohe, P. A., Bassham, S., Amores, A., & Cresko, W. A. (2013). Stacks: an analysis tool set for population genomics. *Mol Ecol, 22*(11), 3124-3140. doi:10.1111/mec.12354

Chen, B., Feder, M. E., & Kang, L. (2018). Evolution of heat-shock protein expression underlying adaptive responses to environmental stress. *Mol Ecol, 27*(15), 3040-3054. doi:10.1111/mec.14769

Cingolani, P., Platts, A., Wang, L. L., Coon, M., Nguyen, T., Wang, L., . . . Ruden, D. M.
(2012). A program for annotating and predicting the effects of single nucleotide
polymorphisms, SnpEff. *Fly (Austin), 6*(2), 80-92.
Colautti, R. I., & Barrett, S. C. H. (2013). Rapid Adaptation to Climate Facilitates Range
Expansion of an Invasive Plant. *Science, 342*(6156), 364 - 366.
Consortium, T. U. (2017). UniProt: the universal protein knowledgebase. *Nucleic Acids Res.,*
45, D158-D169.
Davey, J. W., & Blaxter, M. L. (2011). RADSeq: next-generation population genetics.
Briefings in Functional Genomics, 9(5-6), 416-423.
Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., . . . Gingeras, T. R.
(2013). STAR: ultrafast universal RNA-seq aligner. *Bioinformatics, 29*(1), 15-21.
doi:10.1093/bioinformatics/bts635
El-Sabban, F., Fahim, M. A., Al Homsi, M. F., & Singh, S. (1996). Dehydration accelerates
in vivo platelet aggregation in pial arterioles of lead-treated mice. *J. therm. Biol.,*
20(6), 469-476.
Estoup, A., Ravigné, V., Hufbauer, R., Vitalis, R., Gautier, M., & Facon, B. (2016). Is There
a Genetic Paradox of Biological Invasion? *Annual Review of Ecology, Evolution, and*
Systematics, 47, 51-72.
Estoup, A., Wilson, I. J., Sullivan, C., Cornuet, J. M., & Moritz, C. (2001). Inferring
Population History From Microsatellite and Enzyme Data in Serially Introduced Cane
Toads, Bufo marinus. *Genetics, 159*, 1671-1687.
Foll, M., & Gaggiotti, O. (2008). A Genome-Scan Method to Identify Selected Loci
Appropriate for Both Dominant and Codominant Markers: A Bayesian Perspective.
Genetics, 180, 977-993.
Francis, R. M. (2016). pophelper: an R package and web app to analyse and visualize
population structure *Molecular Ecology Resources, 17*(1), 27-32.
Frankham, R. (2005). Genetics and extinction. *Biological Conservation, 126*(2), 131-140.
doi:10.1016/j.biocon.2005.05.002
Franks, S. J., & Munshi-South, J. (2014). Go forth, evolve and prosper: the genetic basis of
adaptive evolution in an invasive species. *Molecular Ecology, 23*, 2137-2140.
Frichot, E., & Francois, O. (2015). LEA: An R package for landscape and ecological
association studies. *Methods in Ecology and Evolution, 6*(8), 925-929.
Gao, Y., Li, S., & Zhan, A. (2018). Genome-wide single nucleotide polymorphisms (SNPs)
for a model invasive ascidian Botryllus schlosseri. *Genetica, 146*(2), 227-234.
doi:10.1007/s10709-018-0015-1
Godoy, O., Lemos-Filho, J. P., & Valladares, F. (2011). Invasive species can handle higher
leaf temperature under water stress than Mediterranean natives. *Environmental and*
Experimental Botany, 71(2), 207 - 214.
Goudet, J. (2005). HIERFSTAT, a package for R to compute and test hierarchical F-statistics.
Molecular Ecology Notes, 5, 184 - 186. doi:10.1111/j.1471-8278
Greger, R. (2000). Physiology of renal sodium transport. *Am J Med Sci., 319*(1), 51-62.
Gruber, B., Unmack, P. J., Berry, O. F., & Georges, A. (2018). dartr: An r package to
facilitate analysis of SNP data generated from reduced representation genome
sequencing. *Molecular Ecology Resources, 18*(3), 691-699.
Gruber, J., Brown, G. P., Whiting, M. J., & Shine, R. (2017). Is the behavioural divergence
between range-core and range-edge populations of cane toads (Rhinella marina) due
to evolutionary change or developmental plasticity? *Royal Society Open Science, 4.*
doi:10.1098/rsos.180197
Hijmans, R. J. (2015). raster: Geographic data analysis and modeling.

Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., & Jarvis, A. (2005). Very high
769   resolution interpolated climate surfaces for global land areas. *International Journal of*
770   *Climatology, 25*, 1965–1978.
771   Hudson, C. M., Brown, G. P., & Shine, R. (2016). It is lonely at the front: contrasting
772   evolutionary trajectories in male and female invaders. *R Soc Open Sci, 3*(12), 160687.
773   doi:10.1098/rsos.160687
774   Hudson, C. M., Brown, G. P., & Shine, R. (2016). It is lonely at the front: contrasting
775   evolutionary trajectories in male and female invaders. . *Royal Society Open Science,*
776   *3*, 160687.
777   Hudson, C. M., Brown, G. P., Stuart, K., & Shine, R. (2018). Sexual and geographical
778   divergence in head widths of invasive cane toads, Rhinella marina (Anura:
779   Bufonidae), is driven by both rapid evolution and plasticity. *Biological Journal of the*
780   *Linnean Society, 124*, 188-199.
781   Institute, B. (2018). Picard Tools. http://broadinstitute.github.io/picard. Retrieved from
782   http://broadinstitute.github.io/picard.
783   Jessop, T. S., Letnic, M., Webb, J. K., & Dempster, T. (2013). Adrenocortical stress
784   responses influence an invasive vertebrate's fitness in an extreme environment. *Proc*
785   *Biol Sci, 280*(1768), 20131444. doi:10.1098/rspb.2013.1444
786   Jombart, T., & Ahmed, I. (2011). adegenet 1.3-1: new tools for the analysis of genome-wide
787   SNP data. *Bioinformatics, 27*(21), 3070–3071.
788   Keenan, K., McGinnity, P., Cross, T. F., Crozier, W. W., & Prodohl, P. (2013).
789   diveRsity:AnR package for the estimation and exploration of popula tion genetics
790   parameters and their associated errors. *Methods in Ecology and Evolution, 4*, 782-788.
791   Kiang, J. G., & Tsokos, G. C. (1998). Heat Shock Protein 70 kDa: Molecular Biology,
792   Biochemistry, and Physiology. *Pharmacology & Therapeutics, 80*(2), 183 - 201.
793   Knaus, B. J., & Grünwald, N. J. (2017). vcfr: a package to manipulate and visualize variant
794   call format data in R. *Molecular Ecology Resources, 17*(1), 44-53.
795   Kosmala, G., Christian, K., Brown, G. P., & Shine, R. (2017). Locomotor performance of
796   cane toads differs between native-range and invasive populations. *The Royal Society*
797   *Open Science, 4*, 170517. doi:10.1098/rsos.170517 10.6084/m9
798   Kosmala, G. K., Brown, G. P., Christian, K. A., Hudson, C. M., & Shine, R. (2018). The
799   thermal dependency of locomotor performance evolves rapidly within an invasive
800   species *Ecology and Evolution, 8*(9), 4403-4408.
801   Kosmala, G. K., Brown, G. P., & Shine, R. (2018). Kicking back Down Under: Invasive
802   populations of cane toads (Rhinella marina) down-regulate their responses to thermal
803   and hydric stress in a climatically harsh environment. *Submitted*.
804   Leblois, R., Rousset, F., Tikel, D., Moritz, C., & Estoup, A. (2000). Absence of evidence for
805   isolation by distance in an expanding cane toad (Bufo marinus) population: an
806   individual-based analysis of microsatellite genotypes. *Molecular Ecology, 9*, 1905 -
807   1909.
808   Leydet, K. P., Grupstra, C. G. B., Coma, R., Ribes, M., & Hellberg, M. E. (2018). Host-
809   targeted RAD-Seq reveals genetic changes in the coral Oculina patagonica associated
810   with range expansion along the Spanish Mediterranean coast. *Mol Ecol, 27*(11), 2529-
811   2543. doi:10.1111/mec.14702
812   Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., . . . Genome Project
813   Data Processing, S. (2009). The Sequence Alignment/Map format and SAMtools.
814   *Bioinformatics, 25*(16), 2078-2079. doi:10.1093/bioinformatics/btp352
815   Lillie, M., Shine, R., & Belov, K. (2014). Characterisation of major histocompatibility
816   complex class I in the Australian cane toad, Rhinella marina. *PLOS ONE, 9*(8),
817   e102824. doi:10.1371/journal.pone.0102824

Lischer, H. E., & Excoffier, L. (2012). PGDSpider: an automated data conversion tool for
connecting population genetics and genomics programs. *Bioinformatics, 28*(2), 298-
299. doi:10.1093/bioinformatics/btr642

Mader, G., Castro, L., Bonatto, S. L., & de Freitas, L. (2016). Multiple introductions and
gene flow in subtropical South American populations of the fireweed, Senecio
madagascariensis(Asteraceae). *Genet Mol Biol, 39*(1), 135-144.

Madsen, T., Shine, R., Olsson, M., & Wittzell, H. (1999). Restoration of an inbred adder
population. *Nature, 402*, 34 - 35.

Malmkvist, J., Damgaard, B. M., Pedersen, L. J., Jorgensen, E., Thodberg, K., Chaloupkova,
H., & Bruckmaier, R. M. (2009). Effects of thermal environment on hypothalamic-
pituitary-adrenal axis hormones, oxytocin, and behavioral activity in periparturient
sows *Journal of Animal Science, 87*(9), 2796–2805.

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., . . .
DePristo, M. A. (2010). The Genome Analysis Toolkit: a MapReduce framework for
analyzing next-generation DNA sequencing data. *Genome Research, 20*, 1297-1303.

Oksanen, J., Blanchet, F. G., Friendly, M., Kindt, R., Legendre, P., McGlinn, D., . . . Wagner,
H. (2018). vegan: Community Ecology Package. R package.

PubmedHealth. (2014). Excessive Blood Clotting.   Retrieved from
https://www.ncbi.nlm.nih.gov/pubmedhealth/PMH0062998/

Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., . . .
Sham, P. C. (2007). PLINK: a toolset for whole-genome association and population-
based linkage analysis. *American Journal of Human Genetics, 81*.

Raj, A., Stephens, M., & Pritchard, J. K. (2014). fastSTRUCTURE: Variational Inference of
Population Structure in Large SNP Data Sets. *Genetics, 197*(2), 573 - 589.
doi:10.1534/genetics.114.164350/-/DC1

Reed, D. H., & Frankham, R. (2003). Correlation between Fitness and Genetic Diversity.
Conservation Biology, 17(1), 230 - 237.

Richardson, M. F., Sequeira, F., Selechnik, D., Carneiro, M., Vallinoto, M., Reid, J. G., . . .
Rollins, L. A. (2018). Improving amphibian genomic resources: a multi-tissue
reference transcriptome of an iconic invader. *GigaScience, 7*(1), 1-7.

Rollins, L. A., Moles, A. T., & Lam, S. (2013). High genetic diversity is not essential for
successful introduction. *Ecology and Evolution, 3*, 4501 - 4517.

Rollins, L. A., Richardson, M. F., & Shine, R. (2015). A genetic perspective on rapid
evolution in cane toads (Rhinella marina). *Mol Ecol, 24*(9), 2264-2276.
doi:10.1111/mec.13184

Rollins, L. A., Woolnough, A. P., Wilton, A. N., Sinclair, R., & Sherwin, W. B. (2009).
Invasive species can't cover their tracks: using microsatellites to assist management of
starling (Sturnus vulgaris) populations in Western Australia. *Mol Ecol, 18*(8), 1560-
1573. doi:10.1111/j.1365-294X.2009.04132.x

Shine, R., Brown, G. P., & Phillips, B. L. (2011). An evolutionary process that assembles
phenotypes through space rather than through time. *Proceedings of the National
Academy of Sciences of the United States of America, 108*, 5708 - 5711.

Skeeles, L. E., Fleming, J. L., Mahler, K. L., & Toland, A. E. (2013). The impact of 3'UTR
variants on differential expression of candidate cancer susceptibility genes. *PLOS
ONE, 8*(3), e58609.

Slade, R. W., & Moritz, C. (1998). Phylogeography of Bufo marinus from its natural and
introduced ranges. *Proceedings of the Royal Society of London. Series B: Biological
Sciences, 265*, 769.

Szucs, M., Vahsen, M. L., Melbourne, B. A., Hoover, C., Weiss-Lehman, C., & Hufbauer, R.
A. (2017). Rapid adaptive evolution in novel environments acts as an architect of

868   population range expansion. *Proc Natl Acad Sci U S A, 114*(51), 13501-13506.
869   doi:10.1073/pnas.1712934114

870 Team, R. C. (2016). R: A language and environment for statistical computing. Vienna,
871   Austria: R Foundation for Statistical Computing. Retrieved from https://www.r-
872   project.org/

873 Telemeco, R. S., & Addis, E. A. (2014). Temperature has species-specific effects on
874   corticosterone in alligator lizards. *Gen. Comp. Endocrinol., 206*, 184-192.

875 Thioulouse, J., & Dray, S. (2007). Interactive Multivariate Data Analysis in R with the ade4
876   and ade4TkGUI Packages. *Journal of Statistical Software, 22*(5), 1 - 14.

877 Tingley, R., Greenlees, M. J., & Shine, R. (2012). Hydric balance and locomotor
878   performance of an anuran (Rhinella marina) invading the Australian arid zone. *Oikos,*
879   *121*, 1959-1965.

880 Trumbo, D. R., Epstein, B., Hohenlohe, P. A., Alford, R. A., Schwarzkopf, L., & Storfer, A.
881   (2016). Mixed population genomics support for the central marginal hypothesis across
882   the invasive range of the cane toad (Rhinella marina) in Australia. *Molecular Ecology,*
883   *25*(17), 4161–4176 doi:10.1111/mec.13754

884 Turvey, N. (2009). A Toad's Tale. *Hot Topics from the Tropics, 1*, 1 - 10.

885 Wang, Z., Gerstein, M., & Snyder, M. (2009). RNA-Seq: a revolutionary tool for
886   transcriptomics. *Nature Reviews Genetics, 10*(57-63).

887 Weber, E., & Schmid, B. (1998). Latitudinal population differentiation in two species of
888   Solidago (Asteraceae) introduced into Europe. *American Journal of Botany, 85*(9),
889   1110-1121.

890 Wellband, K. W., Pettitt-Wade, H., Fisk, A. T., & Heath, D. D. (2018). Standing genetic
891   diversity and selection at functional gene loci are associated with differential invasion
892   success in two non-native fish species. *Mol Ecol, 27*(7), 1572-1585.
893   doi:10.1111/mec.14557

894 Westemeier, R. L., Brawn, J. D., Simpson, S. A., Esker, T. L., Jansen, R. W., Walk, J. W., . .
895   . Paige, K. N. (1998). Tracking the long-term decline and recovery of an isolated
896   population. *Science, 282*, 1695-1698.

897 White, T. A., Perkins, S. E., Heckel, G., & Searle, J. B. (2013). Adaptive evolution during an
898   ongoing range expansion: the invasive bank vole (Myodes glareolus) in Ireland. *Mol*
899   *Ecol, 22*(11), 2971-2985. doi:10.1111/mec.12343

900 Whitlock, M. C., & Lotterhos, K. E. (2015). Reliable Detection of Loci Responsible for Local
901   Adaptation: Inference of a Null Model through Trimming the Distribution of FST.
902   *The American Naturalist, 186*(S1), 24-36.

903 Whitney, K. D., & Gabler, C. A. (2008). Rapid evolution in introduced species, 'invasive
904   traits' and recipient communities: challenges for predicting invasive potential.
905   *Diversity and Distributions, 14*, 569-580. doi:10.1111/j

906 Willoughby, J. R., Harder, A. M., Tennessen, J. A., Scribner, K. T., & Christie, M. R. (2018).
907   Rapid genetic adaptation to a novel environment despite a genome-wide reduction in
908   genetic diversity. *Mol Ecol, 27*(20), 4041-4051. doi:10.1111/mec.14726

909 Zerebecki, R. A., & Sorte, C. J. B. (2011). Temperature Tolerance and Stress Proteins as
910   Mechanisms of Invasive Species Success. *PLOS ONE, 6*(4), e14806.

917 **Figures:**

918
Fig 1. Map of two invaded territories (Australia and Hawai'i) and the native range (French

Guiana) of the cane toad (*Rhinella marina*). The shaded region represents the toad's current

Australian range, which is continuing to expand. Black diamonds in French Guiana, Hawai'i,

and Australia indicate collection sites for our RNA-Seq experiment. White circles in

Australia indicate collection sites for the RADSeq experiment from which we downloaded

data.

Fig 2. (A) Population structure of cane toads from French Guiana, Hawai'i, and Australia, as
inferred by SNPs from spleen RNA-Seq data. (B) Rainfall during the driest quarter in each
location across collection sites for invasive cane toads in French Guiana, Hawaii, and
Australia. A similar environmental barrier occurs in temperature (Fig S1). (C) Population
structure of invasive cane toads from Hawai'i and Australia, as inferred by SNPs from brain
RNA-Seq data. (D) Substructure within toads from Hawai'i (HI) and Queensland (QLD) as
inferred by SNPs from brain RNA-Seq data. (E) Substructure within toads from Northern
Territory (NT) and Western Australia (WA) as inferred by SNPs from brain RNA-Seq data.

Fig 3. (A) Relationship between environmental variables and population structure of cane

toads from French Guiana, Hawai'i, and Australia, as inferred by Redundancy Analysis

(RDA) on SNPs from spleen RNA-Seq data. Environmental data include: annual mean

temperature (Mean Temp), maximum temperature of the warmest month (Max Temp),

minimum temperature of the coldest month (Min Temp), annual precipitation (Mean Rain),

precipitation of the wettest quarter (Wet Rain), and precipitation of the driest quarter (Dry

Rain). (B) Relationship between environmental variables and population structure of invasive

cane toads from Hawai'i and Australia, as inferred by RDA on SNPs from brain RNA-Seq

data.

952   Table 1. Expected heterozygosity (He), allelic richness (AR), and Shannon's Information

953   Index (SI) estimated using SNPs from spleen RNA-Seq data on native and invasive

954   populations of cane toads (*Rhinella marina*) at regions across French Guiana, Hawai'i, and

955   Australia. QLD = Queensland, NT = Northern Territory, WA = Western Australia.

Region	All loci (He, AR, SI)	Outlier F_{ST} loci (He, AR, SI)	Environmentally associated loci (He, AR, SI)
French Guiana (native, N=8)	0.27, 1.74, 0.40	0.03, 1.25, 0.06	0.17, 1.53, 0.25
Hawai'i (source, N=10)	0.25, 1.72, 0.37	0.08, 1.26, 0.13	0.34, 1.90, 0.51
QLD (core, N=10)	0.25, 1.73, 0.38	0.08, 1.29, 0.13	0.34, 1.91, 0.50
NT (intermediate, N=8)	0.23, 1.69, 0.35	0.02, 1.07, 0.03	0.23, 1.73, 0.36
WA (front, N=10)	0.23, 1.68, 0.34	0.02, 1.11, 0.04	0.21, 1.68, 0.33

956
957
958   Table 2. Expected heterozygosity (He), allelic richness (AR), and Shannon's Information

959   Index (SI) estimated using SNPs from brain RNA-Seq data on native and invasive

960   populations of cane toads (*Rhinella marina*) at regions across French Guiana, Hawai'i, and

961   Australia. QLD = Queensland, NT = Northern Territory, WA = Western Australia.

Region	All loci (He, AR, SI)	Outlier F_{ST} loci (He, AR, SI)	Environmentally associated loci (He, AR, SI)
Hawai'i (source, N=18)	0.33, 1.97, 0.49	0.35, 1.87, 0.51	0.41, 2.00, 0.60
QLD (core, N=18)	0.33, 1.98, 0.50	0.39, 2.00, 0.56	0.24, 1.92, 0.38
NT (intermediate, N=18)	0.31, 1.95, 0.47	0.17, 1.66, 0.28	0.09, 1.52, 0.16
WA (front, N=18)	0.29, 1.93, 0.45	0.12, 1.49, 0.19	0.07, 1.46, 0.12

970
Fig 4. Relationship between geographic distance (km) and genetic distance (pairwise F_{ST})

between cane toads collected across their invasive Australian range (Fig 1) as inferred by

brain RNA-Seq data. (A) represents the full dataset of loci (p = 1E-3, r = 0.76). (B) represents

all F_{ST} outliers (p = 1E-3, r = 0.75). (C) represents only F_{ST} outliers with an association with

either rainfall or temperature (p = 1E-3, r = 0.62). (D) represents only F_{ST} outliers without an

association with either rainfall or temperature (p = 1E-3, r = 0.77).

980
981 Fig 5. (A) Population structure of cane toads across their invasive range in Australia, as

982 inferred from running fastStructure on SNPs from RADSeq data. Because coordinate

983 metadata for collection sites were unavailable, the samples could not be arranged from east to

984 west along the Australian range. (B) Population structure of cane toads across their invasive

985 range in Australia, as inferred from DAPC on SNPs from RADSeq data.