TITLE PAGE

TITLE

Mouse navigation strategies for odor source localization

RUNNING TITLE

Odor source localization by mice

AUTHOR NAMES AND AFFILIATIONS

Liu, Annie[1,4]*, Papale, Andrew E[1]*, Hengenius, James[2]*, Patel, Khusbu[3], Ermentrout, Bard[2,4], and

Urban Nathaniel N[1,4]

* These three authors contributed equally to this manuscript

1. Department of Neurobiology, University of Pittsburgh School of Medicine, Pittsburgh, PA

2. Department of Mathematics, University of Pittsburgh, Pittsburgh, PA

3. University of Pittsburgh, Pittsburgh, PA

4. Center for the Neural Basis of Cognition, Pittsburgh, PA

AUTHOR CONTRIBUTIONS

JH, AL, BE, and NNU designed the behavioral experiments. AEP, JH, BE, and NNU designed the

simulations. AL and KP conducted the behavioral experiments. AEP, JH, and AL analyzed the

data. AEP and JH wrote the MATLAB code. AEP calculated the statistics. AEP, JH, AL, BE, and

NNU wrote the paper.

CORRESPONDING AUTHOR

Andrew E. Papale

papalea@pitt.edu

KEYWORDS

Olfaction, navigation, casting, binaral-olfaction, binaral-sniffing, serial-sniffing

SUMMARY STATEMENT

Use of a naturalistic odor-source localizing task uncovers key strategies underlying successful mouse navigation. Concentration-dependent models successfully recapitulate mouse behavior and reveal important behavioral components.

ABSTRACT

Navigating an odor landscape is a critical behavior for the survival of many species, including mice. An ethologically relevant mouse behavior is locating food using information about odor concentration, such as the relatively laminar gradient established by diffusion-convection near surfaces. To approximate this behavior, we use an open field odor-based spot-finding task indoors with little wind, examining navigation strategies as mice search for and approach an odor source. After mice were trained to navigate to odor sources paired with food reward, we observed behavioral changes consistent with localization 10-45cm away from the source. These behaviors included both orientation towards the source, decreased velocity, and increased exploration time. We also found that the amplitude of 'casting,' which we broadly define as lateral back and forth movement of the olfactory sensors, increased with proximity to the source. Based on these observations, we created a concentration-sensitive model to simulate mouse behavior. This model provided evidence for a binaral-sniffing strategy (inter-nostril comparison of concentration in a single sniff) and a serial-sniffing strategy (sampling concentration, moving in space, and then sampling again). Serial-sniffing may be accomplished at farther distances by moving the body and at closer distances by moving the head (casting). Together, these results elucidate key components of behavioral strategies for odor-based navigation.

INTRODUCTION

Mice, insects, and other animals navigate odor trails and locate odor sources with high fidelity, a remarkable feat given the complexity and variability of odor stimuli. Prior studies of olfactory navigation have focused on the mechanisms of navigating through odor plumes in air (Gire et al., 2016; Vickers, 2000) or water (Koehl et al., 2001), or of scent-tracking on the ground, specifically following odor trails (Khan et al., 2012; Means et al., 1992; Porter et al., 2007; Thesen et al., 1993; Wallace et al., 2002). However, different strategies may emerge for search under conditions near surfaces where intermittency (the fraction of samples above detection threshold) is high (Connor et al., 2018). In this study, we examine the odor-driven behaviors involved in the search for a discrete

odor object under low wind conditions where gradient information is preserved, and develop a

model of olfactory navigation to describe multiple strategies used for effective odor localization.

One general navigation strategy preserved across species and environmental conditions involves

the use of lateral back-and-forth changes in orientation of the olfactory sensors during active odor-

sampling (Baker et al., 2018). Moths navigating odor plumes use this strategy to reacquire an odor

plume, a behavior termed "zig-zagging" or "casting" depending on the magnitude and angle of

displacement (Carde & Willis, 2008; Kennedy, 1983; Kuenen & Carde, 1994; Marsh et al., 1978).

Flies (Budick & Dickinson, 2006) and cockroaches (Willis et al., 2008) use a similar casting strategy

during plume navigation. Lateral back-and-forth behaviors are also often observed in animals

following odor trails. Rats employ head oscillations that sweep back and forth across a trail as they

track it (Khan et al., 2012), a strategy also used by humans (Porter et al., 2007) and ants (Draft et

al., 2018; Hangartner, 1969) following trails. In this study, we use the term 'casting' to describe the

amplitude of the movement of the olfactory sensors during navigation toward an odor source

(Baker et al., 2018; Zhao et al., 2017).

Successful navigation of a complex odor environment likely involves different behavioral strategies

at different times. Knowledge of the odor environment, characteristics of specific odor cues, and

general task context may each influence odor-guided behavior. Animals demonstrate strategy-

switching between distinct behaviors as a consequence of both changes in odor stimulus

properties and task context (Gire et al., 2016; Pierce-Shimomura et al., 1999; Thesen et al., 1993;

Wesson et al., 2008). Environmental familiarity and visual cues also influence decision-making

strategy during odor source localization. Given visual cues to navigate, exploratory casting based

on odor cues disappears on tasks with a small number of potential goal locations; instead, mice

begin directly navigating to known odor source locations in straight-line trajectories (Bhattacharyya

& Bhalla, 2015; Gire et al., 2016). This resembles the transition from exploration to exploitation

behavior observed during successive search trials (Hills et al., 2015; Sutton & Barto, 1998). Thus,

both behavioral context and the availability of orienting stimuli are important for strategy choice in

odor source localization.

Here, we use an open field odor-based spot-finding task to examine strategies of olfactory search

for an odor source deposited on the surface under air flow conditions typical of an indoor setting.

By randomizing odor source location, performing experiments in the dark, and using a discrete

target odorant object, we prevent mice from switching to methods of systematic search based on

memory, visual, or proprioceptive cues (Gire et al., 2016). We find that multiple dimensions of

behavior are varied as mice approach the odor source, including velocity and casting, suggesting

that these behaviors are associated with odor detection or localization. We then created agent-

based models of olfactory search and use simulations to interrogate the roles of binaral-sniffing,

variable velocity, and variable casting on effective odor localization. Both behavioral and simulation

data suggest that multiple behavioral strategies are important for successful odor navigation to a

deposited odor source under typical indoor air flow conditions in mice.

MATERIALS AND METHODS

Ethical approval. All experiments were completed in compliance with the guidelines established by

the Institutional Animal Care and Use Committee of the University of Pittsburgh.

Animals. Four male and one female adult M72-IRES-ChR2-YFP knock-in mice (The Jackson

Laboratory) (N=5) were used for this study, with training begun after age P47. Average age at

training start was P72. All mice were housed singly with behavioral enrichment (running wheel and

house) and received unlimited access to water in their home cage.

Behavioral arena. A large, open field behavioral arena was used for data collection, as described in

Jones and Urban, 2018 (Figure 1A, adapted with permission from Jones and Urban, 2018). The

arena is a custom built 36" x 45" transparent plexiglass and acrylic surface without walls mounted 4

feet above the floor in an aluminum frame. Rows of infrared light-emitting diodes (LEDs) were

mounted along each of the four table surface edges. Data were collected using a camera mounted

below the table surface (1280x1024 resolution, 11.2pixels/cm, 50 frames s^{-1}, Flea3, Point Grey

Imaging). Trials were run in the dark and LEDs from equipment in the room were blacked out with

electrical tape. Remaining light from computer monitors was red-shifted so as to not be visible to

the mice.

Odorized crayon. We created odorized wax crayons similar to what was described in Jones and

Urban, 2018. Briefly, 4.9 g of a Crayola crayon, 0.1 g of Crayola chalk, and 0.1 g of odorant

consisting of one of three different concentrations (0.1%, 1%, and 2% by weight) of methyl

salicylate (Sigma-Aldrich) diluted in mineral oil (Fisher Chemical Paraffin Oil, 0121-1) were mixed

together, and the mixture was poured into a silicon or rubber crayon mold. After ten minutes, the

mixture cooled and hardened into an odorized crayon.

Task: Prior to each trial, the table was wiped clean with 200 proof ethanol using a paper towel.

After the table dried, a spot ~1cm in diameter was drawn in a pseudorandom location in the central

area of the table (Figure 1B). Based on recent spot placements, the experimenter attempted to

place the spot at a different location each subsequent trial. For baited trials, a small piece of

chocolate or peanut (~0.1 g) was placed in the middle of the spot, while for unbaited trials, the spot

was left empty. No more than 2 unbaited trials were run on a given day to prevent extinction of

motivated behavior. With the lights off, a mouse was placed onto the table at a pseudorandom

starting location along the right side of the table and allowed to freely explore the arena for

approximately 3 minutes. Over the course of 5 months, 133-197 trials were collected per animal, 3-

5 trials per day, 3-5 days per week, at a time between 9 am and 2 pm on a normal light-dark cycle.

Food deprivation and task training. To motivate mice to perform the task, they were food deprived

to 78±07% of original body weight. Animals received between 1.8-2.2 g of their normal chow per

day and were given free access to water. Animals were trained to associate a spot of odor (<1cm

diameter) with food reward, either a small sliver of peanut or milk chocolate. Training consisted of

two stages over the course of 7-10 trials: 1) habituation (2-3 days); and, 2) odor-reward association

(5-10 days). During habituation, mice were placed on the table for one 15 minute long trial on two

consecutive days and allowed to explore the arena with no food reward present. During initial food-

odor association, plexiglass dividers were used to decrease the arena size to 25% of the full arena.

A baited odor spot was placed on the table. Once animals found the baited spot in the small arena

during at least 66% of the day's trials, the arena size was increased to 50% the size of the full

arena, followed by an increase to the full 45" x 36" arena once the spot was again found during

66% of the day's baited spot trials. Unbaited spot finding trials were introduced only after the

animals successfully performed the baited spot task in the full arena. Training ended and data

collection began upon successful performance of unbaited spot trials, which occurred following 6-

13 days of training (18-39 trials). All reported data are from the full sized arena, but contain both

baited and unbaited trials. Baited and unbaited trials were analyzed separately, but were found to

be qualitatively similar and so are grouped together.

Video Processing. 44 trials were removed from analysis for technical reasons, leaving N=915 trials

analyzed. Trials were removed if 1) the video file was corrupted; or, 2) tracking the mouse using

Optimouse (see below) was judged by the experimenter to be unsuccessful before data analysis

began. This likely occurred due to unintended variation in table illumination on different trials. Of

these 915 trials, 54 had odor spot placement within 4 cm of the edge, and were removed from

further analysis due to potential obstruction of mouse trajectories near the spots. Finally, 5 trials

were removed because the mouse found the spot within 1 s, indicating that it was initially placed on

the arena too close to the odor spot (most trials where this occurred the mouse started within 10

cm of the odor spot). This left N=853 trials analyzed here. Raw avi formatted files were processed

using the Optimouse open source program (Ben-Shaul, 2017) to extract body position and nose

position. Extracted position files were then post-processed with in-house software written in Matlab

(www.mathworks.com). Position frames were then passed through a 5^{th} order median filter to

suppress jitter in the position data and remove high-frequency noise. Position data was excluded if

1) Optimouse failed to detect a position, 2) the $\log_{10}$ of mean mouse brightness was less than 0.5

times the mean for that trial, 3) if mean velocity was greater than 100 cm s^{-1}, 4) if the mouse length

was greater than 8 cm, 5) if the mouse position 'jumped' by more than 35 pixels (3.13 cm) between

consecutive frames, and 6) if the mouse position was 4 cm or closer from the edge of the arena.

Note that the edges of the arena were metal fasteners and the mouse was often partially obscured

from below when it walked around the edges. Exclusion criteria 6) accounted for the majority (N=

6E6; 84%) of excluded frames, as mice spent a good deal of time at the edges of the arena.

Finally, a nearest neighbor filter was used to remove times when the mouse was stationary as the

computation orientation of the mouse was unstable here. For the majority of analyses, we focused

on either the time before a spot was found, or a matched sample of times for unsuccessful trials.

Behavioral Analysis. A successful spot trial was scored when the mouse's nose came within 1.5

cm of the spot, based on a sharp increase in the survival function on minimum distance to the spot

(Figure 2A).

The fraction of the arena explored was calculated by binning the arena into approximately equal-

area parts of 9-10,000 partitions and, then, the percentage of time exploring the center of the arena

(excluding 4 cm around the edges) was calculated.

Analysis of percent exploration time entailed defining 100 circular rings around the odor spots at
intervals of 1 cm. The number of samples in each ring was then normalized by the total number of
samples in the trial to (% time). For trials when the mouse failed to find the odor source, the entire
session was used.
199
200 To compute the probability of successfully finding an odor source as a function of distance, the
201 occupancy was divided into successful trials (hits) and unsuccessful trials (misses). The fraction of
202 hits was divided by the total number of trials (hits + misses) and averaged across mice.
203
204 The orientation relative to the odor spot was calculated by computing the difference in the four-
205 quadrant arctangent of the vectors from body-to-nose and body-to-spot (Figure 3E).
206
207 Derivatives were computed using the adaptive windowing Janabi-Sharifi algorithm (Janabi-Sharifi
208 et al., 2000) available online (Manurung, 2016), but modified to handle missing data points (NaNs)
209 and to allow for post-smoothing. Parameters used were a window size of 50 (1 s) and an error term
210 of 0.2, with post smoothing of 0.1s.
211
Casting behavior was quantified by computing the numerical derivative of the arctangent of the
nose velocity, taking the absolute value, and then dividing by the linear velocity of the nose
(|dPhi|/nV). This is a modification of the zIdPhi measurement (Papale et al., 2012). |dPhi|/nV is
equivalent to the signed curvature formula with units of m^{-1} and is therefore an appropriate
measure of tortuosity. The algorithm was extensively tested on variable-frequency sine waves to
verify that segments of high curvature were well-separated from segments of high linearity across a
wide range of frequencies. Then, for position data from mice, the $\log_{10}$ of the absolute value of
|dPhi|/nV was taken to generate a roughly normal distribution. An example of the |dPhi|/nV casting
measure (Figure 1C) for one analyzed behavioral trial shows 'elbows' and angled segments with
high curvature (black arrow) well-separated from straight segments of high linearity.
222
223 An alternate measure of casting used was the angle of the nose relative to the midline. For this
224 measure, the angle from the nose to the midline was determined by calculating a vector for the
225 mouse's current heading and computing the shortest distance from the detected nose point to that
226 line, then taking the arccosine of that angle.
227

Anemometer Measurements. A hot wire anemometer (TSI Velocicalc 9535A) was placed at 9

different positions across the table (Figure 1D). 10 measurements were taken with the sensor

aligned to the long-axis or short-axis of the table at each position and the respective averages were

taken.

PID Measurements. A cotton swab dipped in 99% Methyl Salicylate liquid was placed at the center

of the table (Figure 1D, gold star) at approximately 0.5cm from the surface. Concentration was

measured for 30s at distances from 0.5cm to 15cm using a Photoionization Detector (PID)

(miniPid, Aurora Scientific). Raw measurements had noise peaks at 60Hz and 72Hz, broad

spectrum high frequency noise, and substantial baseline drift. To remove the noise peaks, two 2nd

order Butterworth IIR notch filters were used. To remove high frequency noise, a (0.2s backward,

0.02s forward) moving mean filter was used. To remove baseline drift, a baseline estimation and

denoising function was used (Duval, 2018). The denoised traces (Figure 1E) show large signal

fluctuations and, generally, decreasing mean output with distance. Intermittency was calculated as

the mean number of points greater than a threshold of 0.005.

Shuffled Spot Control. As a control, spots were randomly shuffled and paired with tracking data

from one session. Shuffled spots had to be at least 5cm away from the original spot. Analyses

were repeated with the shuffled control data.

Model Specification. Based on observations of mouse behavior, we developed an agent-based

model that navigated a virtual odor environment. This agent made temporally discrete sniff-to-sniff

comparisons of odor concentration as it moved through virtual space, altering its heading toward

higher concentrations and away from lower concentrations. The agent consisted of a body with

coordinates (x, y) and moved through space along a heading θ at a velocity v_c:

253
$$x_{t+\Delta t} = x_t + v_c \, \Delta t \, cos \, \theta \qquad\qquad (n1)$$

254
$$y_{t+\Delta t} = y_t + v_c \, \Delta t \, sin \, \theta \qquad\qquad (n2)$$

where Δt=0.1 s represents a constant the inter-sniff interval. The agent's nose was located at a

distance ℓ=5 cm from the body at coordinates (x_n, y_n). The nose was further divided into left and

right nares, (x_{nL}, y_{nL}) and (x_{nR}, y_{nR}), separated by an inter-nares distance, d_{nares}=1.8 mm. The two

nares are separated by an angle 2γ. The nose was capable of limited movement independent of

the body heading, with an angular deflection ϕ relative to the forward heading. Nares positions

were given as:

$$x_{nL/nR} = x + \sqrt{l^2 + \left(\frac{d_{nares}}{2}\right)^2} \cos(\theta + \phi \pm \gamma) \qquad \text{(n3)}$$

$$y_{nL/nR} = y + \sqrt{l^2 + \left(\frac{d_{nar}}{2}\right)^2} \sin(\theta + \phi \pm \gamma) \qquad \text{(n4)}$$

with nose deflection bounded within $\pm \phi_{max}$ as dictated by mouse anatomy. The agent geometry is

shown superimposed on a mouse body (Figure S2A).

The independent motion of the nose is what allows the agent to sample concentrations to the left

and right of its current heading and adjust its heading toward higher concentrations (Figure 5C).

The distribution of observed mouse nose deflections was approximately Gaussian (Figure S2C);

therefore, we modeled the angular movement of the nose as an Ornstein-Uhlenbeck (OU) process,

a mean-reverting correlated random walk with a Gaussian stationary distribution. At each sniff, the

nose deflection was updated according to:

$$\phi_t = \phi_{t-\Delta t} - \frac{\phi_{t-\Delta t}}{\tau}\Delta t + dW_t\sqrt{\frac{\Delta t}{\tau}} + \tanh\big(k_{binaral}(c_L - c_R)\big)\Delta t \qquad \text{(n5)}$$

$$dW_t \sim N(0, \sigma_c^2) \qquad \text{(n6)}$$

$$\sigma_c = \sigma_{min} + (\sigma_{max} - \sigma_{min})\frac{C_t}{k + C_t} \qquad \text{(n7)}$$

where τ is the characteristic time constant of the process, σ_c is the concentration-dependent

standard deviation of the nose deflection, dW_t is a normally distributed random variable with mean

of zero and standard deviation σ_c. To this OU process, we add binaral bias in the form of the

hyperbolic tangent of the left-right concentration difference; when $C_L > C_R$, the nose is biased to the

left and *vice versa*. Here, $k_{binaral}$ controls the response of the nose to binaral inputs. The

concentration value C_t is the average concentration across both nares at time t:

$$C_t = \frac{(c_L + c_R)}{2} \qquad \text{(n8)}$$

Here, c_L is the concentration at the left naris and c_R is the concentration at the right naris.

We observed that mouse speed increases with distance from the odor source (see Figure 3F).

Because the agent in our model does not have information about its distance from the odor source,

instead we used a concentration-dependent modulation of speed. Specifically, we use a sigmoidal

function with a quartic coefficient to approximate concentration-dependence of speed:

$$v_c = v_{max}\left(1 - \frac{C_t^4}{k + C_t^4}\right) \qquad \text{(n9)}$$

where v_{max} is the maximum velocity and $k=\frac{1}{2}$ concentration units. Note that this is the same k that

appears in eq n7.

Model Odor Environment. The agent must iteratively sample concentrations in an odor

environment to navigate in the direction of higher concentration. To simulate the ambient

concentration, *C(x,y),* we use a noisy bivariate exponential distribution normalized to a value of one

at the source:

295
$$C(x,y) = e^{-\left(\frac{\sqrt{(x-x_{sp})^2+(y-y_{sp})^2}}{2\,\sigma_{sp}^2}\right)}(1+\eta) \tag{n10}$$

296
$$\eta \sim U(-k_n, k_n) \tag{n11}$$

Here, $(x_{sp},\ y_{sp})$ is the location of the source, σ_{sp} controls the width of the odor distribution around the

source (model results were obtained with σ_{sp}=20 cm), and η is multiplicative uniform noise added

to each point in a spatial grid to approximate the effect of turbulent flow, bounded at k_n=0.5.

To represent the table, we generate a static smooth odor landscape grid with a resolution of 1 mm

according to eq n10 (Figure 5B, step 1). Next, we then apply noise according to eq n11 (Figure 5B,

step 2). We then simulate the PID intermittency in the odor signal by randomly setting grid values

to zero (Figure 5B step,3). The probability that a grid value is zeroed, P_{int}, increases with distance

from the source according to

306
$$P_{int} = 1 - e^{k_{int}\sqrt{(x-x_{sp})^2+(y-y_{sp})^2}} \tag{n12}$$

where k_{int} is the decay parameter set to -0.002 cm^{-1}. Finally, we apply a Gaussian filter with a

standard deviation of 4 mm to introduce local spatial correlations into our landscape (Figure 5B,

step 4).

To reflect the apparent lack of mouse orientation toward the source beyond 30 cm, (Figure 3D), the

models have a concentration detection threshold of 0.25 concentration units. This represents a

distance of approximately 30cm from the source.

A distribution of simulated spot positions was used for simulations, matched to data (Figure 1B),

but due to edge effects reducing overall model performance, spots closer than 10cm to the walls

were not used.

Concentration-Sensitive Model (CSM). The agent moves along forward heading θ at velocity v_c,

iteratively sniffing the concentration at its nose coordinates (Eqs n1-n4). At each sniff, the agent's

nose randomly samples to the left or right of the forward heading depending on the state of the

random nose deflection process (Eqs n5-n7). It compares the current concentration sample, C_t, to

the sample from the previous sniff, $C_{t-\Delta t}$. If C_t is greater than $C_{t-\Delta t}$, the agent sets its heading to the

direction of the nose, $\theta_{t+\Delta t} := \theta_t + \phi_t$. If C_t is less than or equal to $C_{t-\Delta t}$, the agent turns away from the

direction of the nose, $\theta_{t+\Delta t} := \theta_t - \phi_t$. This algorithm is effective at reproducing key parameters of

mouse odor source localization (Figure 5):

$$\theta_{t+\Delta t} = \theta_t + \phi_t \; if \; C_t > C_{t-\Delta t} \; else \; \theta_t - \phi_t \; if \; C_t \leq C_{t-\Delta t} \tag{n13}$$

Correlated Random Walk (CRW) Controls. To test model behavior in the absence of odor-

sensitivity, and to serve as a control in case mice were randomly finding the spots, concentration-

dependent turning was removed from the model and replaced by a random turning decision. Eq

n13 is replaced by:

$$\theta_{t+\Delta} = \theta_t + \phi_t \; if \; z > 0.5 \; else \; \theta_t - \phi_t; \; z\sim U(0,1) \tag{n14}$$

To our surprise, CRW control models had a similar success rate to mice when run for 180 s. After

observing mice on videos; however, we determined they were not exploring the arena for the entire

180 s trial. On successful trials, before finding the odor spots, the average time mice actually spent

exploring the center of the arena was only ~30 s. The CRW controls continuously explored the

center of the arena as their decision rule at table boundaries was to bounce off like a billiard ball.

Therefore, we ran all simulations for 30s to match average mouse exploration time in the center of

the arena.

Reduced Model Controls. To evaluate the effects of different hypothesized features of mouse

navigation, we constructed 8 model variant combinations by leaving out binaral-sniffing,

concentration-dependent velocity, and/or concentration-dependent casting amplitude. We then

evaluated CSM model performance in the absence of these features. To eliminate binaral-sniffing

from the model, $k_{binaral}$ in eq n5 was set to zero. To remove concentration-dependent velocity from

the model, eq n9 was modified so that v_c was equal to the constant v_{max}. To remove concentration-

dependent casting amplitude from the model, eq n7 was modified so that σ_c was set equal to the

constant σ_{min}. The same 8 model variants were created for the CRW control, and results were

averaged together to obtain a control distribution (Figure 6, black lines; Figure 7A, grey bar).

RESULTS

An open field odor-based spot-finding task was developed.

To examine the behaviors relevant to odor source localization, we designed a task that minimized non-olfactory cues. Five mice were trained to find a <1cm diameter spot of odorized wax within a rectangular behavioral arena of approximately 1m^2 in the dark (Figure 1A, and Methods). Odor sources (methyl salicylate with(out) bait) for each of the trials (N=855) across all mice (N=5) were pseudo randomly distributed within the center of the testing table (Figure 1B). An example trajectory shows the nose position of the mouse starting at the right-hand side of the arena and moving leftward before successfully finding the spot (red circle; Figure 1C). In this figure, the color indicates a measurement of curvature (|dPhi|/nV; see Methods) used to detect casting amplitude. Additional example trajectories are displayed in Figure S1, with color indicating relative time (in seconds).

Wind Conditions in the Room. We did not generate any airflow during these experiments but of course air flow occurs in every typical environment and air movement likely dominates the effects of odor diffusion in the vast majority of environments. The room in which these behavioral data were acquired had a ventilation input vent at roughly the center of the room and the primary outflow in the room was underneath the door. Airflow measurements were taken on the behavior table (height 1.135m above the floor) at 9 positions and 2 orthogonal orientations. Wind speed was measured between 1.9ft/min – 9.6ft/min along the short axis and 9.2ft/min – 35.1ft/min along the long axis of the table (Figure 1D). Wind direction was estimated by taking the four-quadrant arctangent (green arrows).

Photoionization Detector (PID) Measurements on Behavior Table. A cotton swab dipped in 99% methyl salicylate was placed at the center of the table (Figure 1D, gold star). The tip of a PID was placed at distances from 0.5cm to 15cm away from the odor source. After denoising (see Methods) PID traces were highly variable but mean output decreased reliably with distance (Figure 1E). Three aspects of the PID data were characterized. The mean variance increased for distances less than 2.5cm and then decreased (Figure 1F, left panel). The increase at small distances could be due to saturation of the PID at high concentrations. The mean output also decreased with distance from the source from about 2.5cm (Figure 1F, center panel).

Intermittency was measured as the fraction of samples above baseline, and decreased with

distance from the source (Figure 1F, right panel).

Mice learn the open field odor-based spot-finding task. To determine an appropriate distance

threshold for achieving a successful trial and an approximate minimum concentration-detection

threshold, we examined the survival function with respect to distance from the spot (Figure 2A).

Typically, a survival function is used to determine the probability that a variable of interest survives

past some time point; here, we use it to describe the probability of a mouse reaching a given

distance away from the odor spot. For mice, there was a sharp increase in survival at about ~1.5

cm, with a more gradual increase toward 100% at about 90 cm from the odor spot (red line). Thus,

mice on all trials successfully came within ~90 cm of the spot. In contrast, the survival function

from shuffled data rose smoothly, crossing the mouse data at about 30cm. We therefore set the

distance threshold for finding a spot to be 1.5cm and estimated the minimum concentration-

detection threshold to be 30cm for later modeling purposes.

Given a threshold of 1.5cm, mice were successful in 64% (N=532/826) of trials (Figure 2B). In

comparison, shuffled controls were successful a mere 1.67% (N=1,524/91,446). Mice had a

significantly higher success rate than shuffled controls (Welch's t-test; p=6.22E-5, t-stat=14.69,

df=4). A logistic regression on successful versus unsuccessful trials for mouse data revealed

several highly predictive explanatory variables (Table S1). Baited trials had higher success rates

(p=4.68E-12, β=2.51), potentially indicating higher motivation during these trials or a contribution of

the odor of food to the localization. Greater arena exploration also predicted success (~45 cm^2

bins; p=1.65E-15, β=15.51; see Figure 3A). Slower mice (p=6.89E-06, β=-0.29) were more

successful, though they had faster average nose velocity (p=1.07E-04, β=0.27) and higher casting

(integrated $\log_{10}$[|dphi|/nV] within 30cm; p=2.56E-3, β=1.4), suggesting that slower forward body

velocity and higher casting amplitude may be features of successful search strategies. Successful

completion of this spot-finding task is thus dependent on features of individual trajectories.

We conclude that mice find odor spots above chance levels established by the shuffled control,

and predict that further evaluation of specific trajectory features will provide useful information

about odor localization strategy.

Mice efficiently navigate to the odor source in comparison to shuffled controls.

Next, we compared navigation efficiencies of mice to data from shuffled controls by examining time

to spot divided by initial distance, nose displacement divided by initial distance, and the probability

of finding the spot as a function of distance. Time to spot was divided by initial distance to spot,

with the average for mice (0.51 s/cm) significantly lower than shuffled controls (1.71 s/cm; Figure

2C; Welch's t-test; p=7.25E-18, t-stat=8.72, df=1700). Then, nose displacement was divided by

initial distance to spot, with the average for mice (4.75x) significantly lower than shuffled controls

(17.78x; Figure 2D; Welch's t-test; p=1.2E-19, t-stat=-9.11, df=1573). To determine the probability

of finding the odor source at various distances away, successful samples (spot found) were divided

by total samples (spot found + not found) in concentric annuli centered at the spot (Figure 2E). For

mice, this resulted in a fairly linear distribution (red line), while for shuffled controls the distribution

was highly nonlinear (cyan line). We conclude that mice are more efficient at navigating to the

odor source than shuffled controls.

Mouse behavior varies systematically as a function of distance from the odor source.

As suggested by the logistic regression analysis of mouse data, exploration percentage, velocity,

and casting amplitude are important for odor source localization. Next we tested whether mice

reliably change behavior depending on proximity to the odor source. We conducted four analyses

to test this hypothesis: 1) looking at the percentage of the arena explored, 2) computing the

percentage of time exploring at a given distance on a given trial; and separately, 3) examining

mouse velocity as a function of distance from the odor source; and 4) analyzing mouse heading

relative to the odor source.

Using a bin size of ~45 cm^2, mice explored an average of 33±17% (mean±s.d.) of the table before

finding the spot. This estimate varied as a function of bin size and successful spot-finding (Figure

3A; two-way ANOVA interaction term; p<1E-100, F=83.95, df=97). Mice also explored

progressively less of the table with increasing trial (~45 cm^2 bins; ANOVA; p=2.65E-51, F=57.16,

df=5), suggesting refinement of learning throughout the experiment.

We then computed the percentage of time mice occupied different distances from the odor source,

normalized by the area of the annulus (Figure 3B). This revealed a decreasing function with

distance on successful trials (red line) compared to a flat distribution with distance on unsuccessful

trials (blue line). These lines diverged at a distance of 44cm (two sample Kolmogorov-Smirnov

test; p<1E-10), suggesting that mice are exploring more within 44cm of the odor source on

successful trials.

Next, we examined mouse velocity as a function of distance from the odor source (Figure 3C). As

mice approached the odor source across all trials, nose velocity slowed down (ANOVA; p=2.72E-

91, F=18.73, df=28), as did body velocity (ANOVA; p=1.02E-160, F=31.12, df=28).

We next examined mouse heading relative to the odor source for successful trials (Figure 3D). If a

mouse detects odor cues that provide insight into the location of the odor source, then the mouse's

nose should orient towards the source. At each position sample, the difference in angle between

body-to-spot and body-to-nose was computed (Figure 3E). We found that orientation relative to the

spot depended on distance from the odor spot (Watson-Williams Test; p<1E-100, F=877.94,

df=48), and has a median value not significantly different from zero (i.e. oriented directly at the

spot) within 9.54cm (p>1E-3, circular test for equal medians with Bonferroni correction).

In summary, we found that mice vary their search strategy and orient towards the spot as they

approach it. Mice also slowed down as they approach the spot. Together, these behavioral

changes potentially represent a speed-accuracy trade-off during the final stage of navigation

towards an odor object, roughly within ~ 10-45cm.

Casting increases near the odor spot and is modulated by velocity

Casting is seen in diverse odor navigation circumstances ranging from plume navigation by moths

to trail tracking by dogs; thus, mice may also cast during the open field odor-based spot-finding

task. To measure casting amplitude, we calculated the $\log_{10}$ of the absolute value of the angular

velocity of the nose divided by the linear velocity of the nose (Figure 1C; and see Methods). This is

an appropriate measure for unsigned path curvature with units of (m^{-1}). We found that curvature

increased as mice approached the odor source (ANOVA; p=2.2E-117, F=23.4, df=28) with an

interaction between distance and successful versus unsuccessful trials (two-way ANOVA;

p=0.0041, F=1.85, df=28).

We used a stepwise Generalized Linear Model (GLM) to determine the relative effect of velocity

and distance on the curvature measure of casting. The first term to include in the stepwise GLM

was velocity (p<1E-100, F=9636, Deviance=1500) and the second was distance (p=7.1E-41,

F=1481, Deviance=1481). No nonlinear terms were added to the model.

As curvature explicitly depended on inverse nose velocity (see Methods and results from stepwise

GLM), we computed the angle between the head and body centerline as an alternate measure of

casting (Figure S3B). Using this alternate measure of nose deflection, we again found that casting

increases as mice approach the odor source (ANOVA; p=2.18E-56, F=12.82, df=28), with an

interaction between distance and successful versus unsuccessful trials (two-way ANOVA; p=0.02,

F=1.61, df=28).

To assess whether animals began using non-odor cues to navigate towards the odor spots on later

trials, we performed one-way ANOVAs on curvature versus binned trial number (N=6). We found

an effect of session on curvature (ANOVA; p=1.42E-11, F=12.34, df=5), suggesting that after 60-70

trials, a shift in strategy occurred which led to systematically less casting. Further analysis

revealed that this effect was consistent with systematically increased nose velocity on later trials

(ANOVA; p=9.16E-7, F=7.36, df=5).

We also examined casting amplitude as a function of approach vector. To compute approach

vectors, the angle of the nose with respect to the body was taken and positions were centered with

respect to the spot, then binned every 2.3cm. The length of the vector was then normalized by the

standard deviation of the angles at that point. The vector colormap indicates the $\log_{10}(|dphi|/nV)$ at

that point. Note that on most trials mice approached the spot from the northeast direction (the

'downwind' direction, see Figure 1D). It was in this direction, at points closest to the source, where

casting increased (Figure 4B).

Overall, we found that casting amplitude of mice increases at distances close to the spot,

suggesting that it may play a role in the successful navigation to a discrete odor object by mice.

This change in casting is potentially correlated with velocity, which also decreases significantly

during the approach to the spot.

A concentration-sensitive model (CSM) was developed to capture key aspects of mouse behavior discovered on the spot-finding task.

The core of the CSM model is the temporal comparison of sniff-to-sniff concentration. At each discrete "sniff," it samples the odor at the position of its nose and compares the current odor concentration with the previously sampled concentration. Part of the algorithm averages across the two nares (Eq n8) while one component (binaral component) compares the difference between nares (right-most term in Eq n5). A typical simulated nose trajectory showing a successful capture displays inter-sniff differences in odor concentration in arbitrary units (Figure 5C). The CSM model used here had parameters: Δt=0.1s, ℓ=5 cm, v_{max}=25cm/s, k=0.5 concentration units, τ=0.1 s, σ_{min}=0.2 radians, σ_{max}=0.3 radians, and $k_{binaral}$=200 The value of τ chosen gave a good balance of sharp turning nose trajectories with smooth nose trajectories. Values an order of magnitude lower resulted in tight spirals of the nose, while values an order of magnitude higher resulted in loss of maneuverability of the model. v_{max} was chosen as it was the approximate average velocity of mice. Because simulated odor concentrations were scaled to a maximum of one, k=0.5 was chosen as the half-maximal value for behavioral response. Changing k could be used to 'tune' the model's performance without qualitatively affecting its trajectories. Values of σ_{min}, $\sigma_{max,}$ and $k_{binaral}$ were chosen so that the model's nose curvature remained qualitatively "mouselike" in character. *I.e.*, the model trajectories did not appear overly tortuous or linear when compared by eye to the mouse trajectories.

To model the observed decrease in velocity close to the odor source, we implemented a concentration-dependent velocity term (Eq n9). In this term, the maximum velocity, v_{max} decreases with increasing concentration according to a sigmoidal function. This function was chosen as it was roughly linear in the center of the concentration range and saturated at high concentrations. To model the increase in curvature with approach to the spot, a concentration-dependent widening of the normal distribution was implemented (Eq n7). These equations were linked by a common parameter k that controlled their rates of change as a function of concentration.

As a control, we created a Correlated Random Walk (CRW) model that turned independently of concentration (Eq n14), aligning its heading with its nose 50% of the time and away from its nose 50% of the time, resulting in left and right turns in equal proportion. Models using odor cues to navigate should outperform the CRW.

CSM simulations have similar general properties to mice.

From our data, we formulated a hypothesis that mice are detecting concentration gradients in order to navigate to the target. While some prior work has suggested that animals can navigate to sources without concentration gradients (Balkovsky & Shraiman, 2002; Vergassola et al., 2007), which may be necessary in highly turbulent environments, odor concentration gradients will develop over time through convection-diffusion and are a likely candidate for rapid olfactory navigation to targets under conditions of low turbulence. We refined our hypothesis to suggest that mice are 1) moving rapidly where the slope of the gradient is low and detecting changes at different physical locations (a versus b); versus, 2) moving slowly where the slope of the gradient is high and detecting changes by sweeping their heads back and forth (A versus B), which we interpret as casting (Figure 5A).

Initially, we had used a simple stimulus model of multiplicative random noise added to a smooth gradient. However, after examining the fluctuations of concentration near the table on a PID, we decided to implement a more complex odor stimulus model. This included 1) generation of a smooth concentration gradient; 2) addition of multiplicative noise which increased near the source; 3) setting pixels equal to zero at a proportion dictated by an intermittency fit such that the farther away from the source the higher proportion of zeros; and, 4) the smoothing of the resulting stimulus by a two-dimensional local Gaussian filter (Figure 5B).

The CSM model with 4-step stimulus generation generally resembled mouse behavior (Figure 5C). Here, concentration difference between consecutive samples is plotted so that the turning away from high-to-low and turning towards low-to-high concentrations is visible. The simulated odor spot is indicated by the red circle.

Spots positions for the CSM and CRW were taken from the mouse data, with spots less than 10cm from the edges being removed. The success rate of the CSM was 94%, significantly greater than the 19% for CRW controls (Figure 5D; Welch's t-test; p=2.16E-22, t-stat=59.45, df=17). The CSM therefore successfully used information about concentration to navigate within 1.5cm of the odor source.

The nose displacement over initial distance ratio for the CSM (~5.6x) was significantly less than the

CRW (~13x) (Figure 5E; Welch's t-test; p=3.5E-11, t-stat=-34-41, df=9), indicating that the CSM

found the odor spots more effectively than the CRW. The survival function of CSM minimum

distance to the spot was shifted to left and up compared to the CRW control (Figure 5F). The

probability of finding the spot measure for the CSM was also consistently higher than the CRW

control (Figure 5G).

Examining the structure of CSM trajectories suggests that they approximate mouse behavior.

The orientation relative to the spot for the CSM was qualitatively similar to mice (Figure 6A,

compare with Figure 3D). The CSM occupancy increased with respect to the CRW occupancy as

agents approached the spot (Figure 6B; two-sample Kolmogorov-Smirnov test; p<1E-10 at

<26cm). This was similar to the occupancy observed in mice on successful trials (Figure 3B).

Mean nose velocity of the CSM decreased as agents approached the spot (Figure 6C), as

expected, given eq n9. On successful trials, the CSM explored *less* than it did on unsuccessful

trials (Figure 6D). This contrasts with the behavior observed in mice (Figure 3A), suggesting that

there are unexplored differences between the CSM and mice. CSM casting increased as agents

approach the spot (Figure 6E), as expected, given eq n7. This approximates the effect observed in

mice (Figure 4A).

Simulations allow separation of component strategies for odor-driven navigation.

Simulations allowed for further dissection of the odor-driven factors that impact behavior and

successful odor source localization such as decrease in velocity close to the source, increase in

casting close to the source, and binaral sniffing. We created reduced CSMs to examine the relative

contributions of three hypothesized behavioral components: binaral-sniffing, concentration-

dependent velocity, and concentration-dependent casting amplitude. These model variants are

denoted (-V) for the model lacking concentration-dependent velocity, (-B) for the model lacking

binaral olfaction, (-C) for the model lacking concentration-dependent casting, (-VC), (-VB), (-SC) for

models lacking multiple components, and (-VBC) for the model lacking all three components. While

all reduced model variants outperformed CRW controls (Figure 7A), removal of any of these

behavioral components decreased model success. The removal of concentration-dependent

velocity led to the largest change in success rate (-V, Δ30%), followed by concentration-dependent

casting amplitude (-C, Δ8%), and binaral-sniffing (-B, Δ3%). The change in success rate was

significant for removal of concentration-dependent velocity (Welch's t-test; p<1E-100, t-stat=169,

df=1472), concentration-dependent casting (Welch's t-test; p<1E-100, t-stat=58.3, df=1759), and

binaral sniffing (Welch's t-test, p=2.65E-100, t-stat=22.6, df=1927). The effect size depended on

the number of bootstrapped samples (out of 100,000) and groups in each sample. For this

analysis, we chose 1000 groups with 10,000 samples in each group, selected without replacement.

These behavioral components of olfactory search are synergistic; simultaneous removal of two or

more of them leads to a greater reduction in success rate than removal of individual components.

Model variant (-VC), in which both velocity decrease and casting increase are removed, resulted in

a significant Δ12% change in model success relative to (-V) alone (Welch's t-test; p<1E-100, t-

stat=-55.9, df=1997). All groups performed significantly better than the CRW Control, even with all

three additional components of the model removed (-VBC; Welch's t-test; p<1E-100, t-stat=67.08,

df=1997). This is logical since (-VBC) still makes decisions based on inter-sniff concentration

differences, while the CRW Control makes random decisions.

Significance of model parameters was also established by shuffling 12 parameters of the model

50k times, and comparing the effect of each parameter using a logistic regression (Table S2). We

also included two dependent variables from the model trajectories that we thought might impact

success rate, the mean casting ($\log_{10}$(|dPhi|/nV) within 30cm, and the mean nose velocity within

30cm. Note that for this analysis, we split k into separate parameters for casting ($k = k_c$ in eq n7)

and velocity ($k = k_v$ in eq n9). Likewise, we labeled and varied the power parameters (n_c = 1 in eq

n7) and (n_v = 4 in eq n9) from 0 - 5. We also added a coefficient in front of the *tanh* binaral term in

eq n5 ($k_{binaral2}$), and where $k_{binaral}$ in eq n5 is relabeled as $k_{binaral1}$. All parameters and both

dependent variables significantly impacted success rate, except for the parameters τ, σ_{min}, and k_c.

To more rigorously test the effects of different levels of noise, the noise level k_n in eq n11 (also see

step 2 in Figure 5B) was varied from 0 to 1 and the Gaussian smoothing (4mm in reported CSM

variant; step 4 in Figure 5B) was varied from 0mm to 10mm. Success rate rapidly dropped off as

k_n approached 1 or smoothing approached 0mm (Figure 7B).

To more rigorously test the effects of the 2-parameter binaral term, we re-ran the 6 reduced model

variants 100,000 times each with a coefficient of $k_{binaral2}$ =3 in front of the *tanh* term in eq n5. Again,

we chose 1000 groups with 10,000 samples in each group, selected without replacement. This

resulted in smaller effects of removing (-V) and (-C), but a larger effect of removing (-B), suggesting

that strong binaral-sniffing may be able to compensate for removal of variable velocity and casting

(Figure 7C).

In summary, reduced models revealed that individual components of the overall strategy worked

together to maximize task performance, with concentration-dependent velocity and binaral-sniffing

having the greatest impact on successful task performance.

DISCUSSION

Here, we described the navigation strategies of mice during an open field odor-based spot-finding

task. Over the course of 7-10 trials, mice could be trained to perform this task on a 45" x 36" open

field arena. Performance of mice exceeded that of shuffled controls, indicating that mice were

using odor cues to localize the odor target. Olfactory-guided behavior changed as a function of

distance; specifically, 1) nose velocity decreased close to the source, 2) casting increased close to

the source, and, 3) occupancy increased close to the source. Mice demonstrate reliable spot-

finding within 10-45cm of the odor source. The Concentration Sensitive Model (CSM) was built to

capture these observations by incorporating concentration-dependent velocity and casting terms,

as well as a postulated binaral comparison term. Removing components of the CSM demonstrated

that each had a significant impact on performance.

There is a potential speed-accuracy trade-off in search strategy close to the odor spot.

As mice approach an odor source, they slow down and increase their exploration time in a smaller

annulus around the spot. This is similar to observations in fruit flies following a plume upwind to an

odor source (Budick & Dickinson, 2006). We suggest this phenomenon may be a type of speed-

accuracy trade-off (Heitz, 2014) in olfactory search behavior. Specifically, when concentration

reaches some perceptual threshold, mice may "switch" into a local search strategy designed to

increase accuracy at the expense of speed to target. In this local search strategy, the mouse

spends more time in regions of high concentration where it is more likely to hit its target. In

agreement with this hypothesis, elimination of concentration-dependent velocity decrease led to

less success in reduced models (Figure 7A). An alternative explanation is that as the mouse

approaches the odor source, it may reduce its running speed to avoid 'overshooting' the odor

source.

Binaral-sniffing facilitates search near the odor source.

Use of binaral-cues for navigation has been observed in moles (Catania, 2013), snakes (Schwenk,

1994), and insects (Martin, 1965; Steck et al., 2010) suggesting it may be an evolutionarily

conserved strategy. In mammals, neurons in the anterior olfactory nucleus may play a key role in

integration of bilateral stimuli that are key to binaral-sniffing (Kikuta et al., 2010) as this is one of

the earliest locations in which information from both left and right naris can be integrated. We

suggest that binaral-sniffing cues may play a pivotal role in fine discrimination of odor identity

(Uchida & Mainen, 2003), as opposed to gross determination of odor source localization in space.

Once a trail's direction, or the rough direction to a spot is determined, binaral-sniffing cues may be

useful for determining the direction to cast in order to continue following the direction of the trail

(Khan et al., 2012), or steer towards the odor spot. Binaral-sniffing allowed the CSM to

preferentially cast in the direction of the odor source only when the odor gradient was large (i.e.,

near the source). Removal of binaral-sniffing led to a statistically significant decrease in CSM

success (Figure 7C), an observation compatible with the model developed by Catania, 2013,

where binaral-sniffing becomes a relevant strategy on steep concentration gradients. Depending

on the strength of the $k_{binaral}$ term (Eq n5), binaral-sniffing may help bias increased lateral search

towards the direction of the source, allowing the nose to be "pulled" toward the source in relatively

smooth arc-like trajectories.

Future research should examine if the physical distance between nostrils is correlated with the

'effective' nares separation of air drawn into the nasal cavity, or if the effective nares separation is

determined by physical parameters such as inhalation volume, flux, and gradient. For example,

inhalation in rats is largely from the lateral direction, rather than the front of the animal (Wilson &

Sullivan, 1999). Pressure differences generated during sniffing inhalation at different magnitudes

and frequencies (slow deep breath versus rapid sniffing) may also draw in relatively different

concentrations of molecules from different distances away from the nares.

Odor source localization may rely on cognitive mapping

Navigation to an odor source with serial-sniffing is probably intimately tied to navigation through

space using the cognitive map (Jacobs, 2012). Specifically, the distance traveled between the first

to the second sniff in the concentration comparison helps determine an updated heading towards

the source or along the trail. In support of this view, hippocampal place cells may change their

firing patterns (called remapping) to match the relocation of an orienting odor source in a circular

arena (Zhang & Manahan-Vaughan, 2015). Thus, serial-sniffing is potentially more relevant to

gross odor localization than binaral-sniffing. For example, to determine trail direction using binaral-

sniffing most efficiently, an animal would have to be oriented roughly perpendicular to the trail so

that its nares are sensing maximally different concentrations. In contrast, studies with dogs

repeatedly find that serial-sniffing of 2-5 footprints is required to determine a trail's direction

(Hepper & Wells, 2005; Thesen et al., 1993). Similarly, initial odor source seeking may utilize

serial-sniffing to quickly map larger spatial areas, including eliminating subspaces where no odor is

detected (Vergassola et al., 2007).

Casting is related to olfactory-guided navigation to a discrete, static odor source.

Casting has been observed at frequencies of ~4 Hz in moths flying upwind during pheromone-

tracking (Charlton et al., 1993), and similar head-scanning behaviors have been observed in

Drosophila larvae, rats, moles, and humans (Baker et al., 2018). In moths, casting has been

reported to occur *outside* of an odor plume when a moth is attempting to re-acquire the odor

(Kennedy, 1983) and is characterized by loop-like flight trajectories (Kuenen & Carde, 1994). In

contrast, here we find that mouse casting amplitude increases during approach to the odor source.

The moth data suggest that casting increases whenever there are intermittent gaps in the odor

signal, as they cast to re-acquire the odor plume. If this were true for mice performing this task, we

would expect casting to be higher farther away from the odor source where the signal is sparse.

The nature of this task, where mice seek a discrete odor object on a near-bed surface rather than

an odor plume in the air, may explain these differences in behavior. While we did not find that

addition of concentration-dependent casting improves performance (Figure 5B), we did not prove

that this was true for all parameter sets in the CSM model. Therefore, a set of untested parameters

may explain the lack of improvement with addition of concentration-dependent casting.

Complexities to address in future studies include decision-rules and individual variability in strategy

utilization.

Our data suggest that mice may use multiple search strategies during an open field odor-based

spot-finding task depending on their distance from the source. We speculate that the relative

contribution of strategies may also be variable between individuals, as has been observed during

fruit fly navigation during odor presentation (Tao et al., 2019). This may mean a specific

instantiation of our model (with one set of parameters) captures behavior for one individual mouse,

but fails to capture behavior for some other subset of mice. For example, we found that the k

parameter could be used to 'tune' aspects of the model's trajectories which may better fit the data

for a subpopulation of mice. A much larger cohort of animals than was used here would be

necessary to evaluate this idea; however, the general principle is that similar-looking ethologically-

relevant behavior can emerge from different model permutations.

A limitation of the current model is that it incorporates an 'all or nothing' decision at each sniff. It

may be more appropriate to use a probabilistic decision rule to combine binaral- and serial-sniffing

into a unified output. Also, a model where pausing emerges during epochs of uncertainty in

sampling of the odor concentration gradient may be advantageous. Additionally, the model is

currently limited to detection of one odorant—it would be useful to distinguish different features of

diverse odorants. Finally, stimulus models with 'zeros' distributed throughout the simulated odor

landscape produced high failure rates. Developing a model that is robust to this type of

intermittency would be a logical next step.

We show that the CSM is capable of using serial concentration sampling of gradients to locate

odors, even when the odor landscape has many local minima and maxima surrounding the odor

source (e.g., Figure 5B, step 4). While the CSM is capable of finding the odor source when the

odor is consistently above threshold, it does not attempt to capture mouse behavior in the

environmental regime where the odor concentration is below the animal's detection threshold.

Rather than attempting to model mouse behavior in this subthreshold condition, the CSM engages

in a correlated random walk that may bring it in contact with suprathreshold odor concentrations.

The mouse may be engaging in more complex navigation behavior in this dilute regime, including

mapping of the odor environment. Models that attempt to navigate complex odor landscapes

without gradient ascent have their own drawbacks. The model of Balkovsky and Shairaman, 2002

requires an additional wind cue - anemotaxis - to function properly. The model of Vergasolla, *et* al,

2007 builds a cognitive map of odor source location likelihood; it utilizes both sparse odor detection

events and frequent subthreshold observations to update its map of source location likelihood.

In summary, we discover that mice vary their search strategy, including casting, during active

navigation toward a discrete odor object, modulating features of behavior as a function of distance

from the target. A concentration-sensitive model successfully reproduces mouse behavior, while

allowing for the analysis of the relative contributions of binaral-sniffing, concentration-dependent

velocity, and concentration-dependent casting amplitude. Future work is needed to experimentally

dissect how these variables impact mouse navigation and whether additional parameters are

important in successful navigation to a static odor source.

ACKNOWLEDGEMENTS

We thank Gregory LaRocca for technical help, and members of the Urban laboratory as well as Yu

Chen, Robert Kass, Nour Riman, Erin Connor, and John Crimaldi for helpful discussions. We

thank Kathy Nagel and Jonathan Victor for helpful comments on an earlier draft of the manuscript.

A special thanks to Peter Jones for designing and building the behavior table used in this

experiment. Thanks to Chris D. Rand from Aurora Scientific for technical help with the PID.

COMPETING INTERESTS

The authors declare no competing financial interests.

FUNDING

This work was supported by the National Science Foundation [1555862 to N.N.U., 1555916 to

B.E.]; the National Institutes of Health, National Institute on Deafness and Other Communication

Disorders [F30DC015161 to A.L.], and the Pennsylvania Department of Health's Commonwealth

Universal Research Enhancement Program [N.N.U.].

FIGURE CAPTIONS

Figure 1. Experimental setup and methods. A) The behavioral arena is a 45" x 36" table made

from three layers of plexiglass, with an infrared (IR) camera placed underneath to view the mice

from below. Odor spots were placed pseudo randomly on the table by the experimenter before

starting the trial, either unbaited or baited with a food reward. (Adapted with permission from Jones

and Urban, 2018.) **B)** The pseudo random odor spot distribution for all trials (N=853). **C)** An

example trajectory where the mouse finds the odor spot, showing nose tracking with the casting

measure (unsigned curvature) on the color axis. The mouse is placed on the right side of the table

and begins to hone in on the odor spot (red circle). The casting measure clearly indicates points of

high curvature (yellow) from points of high linearity (blue). **D)** Anemometer measurements at 9

positions across the behavioral arena. Average wind speed in the <x> and <y> direction are

indicated by the text at each point, with estimated wind velocity vectors indicated by the green

arrows. **E)** Photoionization Detector (PID) readings from 0.5cm to 15cm away from an odor source

placed at the center of the table (gold star in Figure 1D). Measurements were taken to the right of

the star, in the 'downwind' direction. **F)** PID analysis suggests that mean signal variance, mean

signal amplitude, and intermittency decrease starting at about 2.5cm from the source out to about

15cm.

Figure 2. Mice and the CSM efficiently locate the odor source. A) The survival function shows

the fraction of closest distances to the spot as a function of distance from the spot. At about 1.5cm

away from the spot, the closest distance starts to drop off for mice, suggesting this would be a

good threshold distance for counting a trial as successful. In comparison, there is no sharp

distinction in the shuffled control data. Instead, shuffled minimum distances tend to increase

smoothly as a function of increasing distance. At about 30cm away from the odor spot, the

minimum distances are indistinguishable between mice and shuffled controls. **B)** Mice are more

successful at finding the odor spots than shuffled controls, suggesting that they are using odor

cues to guide their search. **C)** The time to spot is divided by the initial distance to the spot, and

mice find the spot more quickly than shuffled controls. **D)** The nose displacement was divided by

the initial distance to the spot, and the relative distance traveled by the nose of mice is less than

that of the shuffled controls. **E)** The number of samples in 100 concentric rings around the spot

were counted up. The number of samples in each ring on successful trials was divided by the total

number of samples to give a measure of probability of finding the spot as a function of distance.

This measure decreased roughly linearly for mice but highly nonlinearly for shuffled controls. This

suggests mice are efficiently navigating toward the spot.

Figure 3. Mouse behavior varies systematically as mice approach the odor source. A) The

center of the arena was broken up into bins of different area and the percentage of those bins

visited by mice was counted up and plotted versus the bin area. In general, this 'exploration

percentage' increased with increasing bin size, while successful trials entailed a higher exploration

percentage than unsuccessful trials. **B)** The number of samples in 100 concentric rings around the

spot was tallied, and normalized by both the total number of samples in a given trial and the area of

the ring. This gave a measure of occupancy with units of %/cm^2. As mice approach the spot on

successful trials, their occupancy increases. In contrast, the occupancy on unsuccessful trials

remains flat as a function of distance. **C)** Mean nose velocity decreases as mice approach the

spot on successful trials. In contrast, mean nose velocity remains flat as a function of distance on

unsuccessful trials. **D)** The orientation of the mouse relative to the odor spot was computed and

normalized to 100% at each distance from the spot. For this measure, orientation of 0° indicates

that the mouse's nose is pointing directly at the odor spot and an orientation of 180° means that it's

nose is pointing away from the odor spot. Orientation tends to converge at 0° as mice approach

the odor spot. **E)** Diagram demonstrating the computation of orientation relative to the spot. The

angle β between the body and spot (red circle) is subtracted from the angle α between the nose

and the body.

Figure 4. Analysis of casting amplitude using curvature. A) Unsigned curvature of the nose

was computed as a measure of casting amplitude (see Methods) and averaged across successful

trials (red line=mean±s.e.m.). Casting increased as mice approached the odor spot (ANOVA;

p=1.21E-64, F=42, df=8), and was significantly higher on successful trials within ~12 cm compared

to unsuccessful trials (Welch's t-test, p=7.2E-5, t-stat=-4.26, df=61, Bonferroni correction). **B)** The

approach vector for successful trials are the vector components of the circular mean of the nose-to-

body orientation after centering position data at the odor spots. Vectors are normalized by the

circular standard deviation. Color indicates the curvature measure for casting at that point. Most

trajectories approach the spot from the northeast.

Figure 5. Development of concentration-sensitive model. A) Our general hypothesis in

constructing the model was that lower concentration differences (ΔC_{ab}) far from the source would

be detectable by gross movement of the body (right panel). In contrast, higher concentration

differences (ΔC_{AB}) would be detectable by fine movement of the head (left panel). In this

illustration, the wavy green line represents the concentration of an odorant. The blue shading

indicates averaging concentration over an inhalation. The dashed lines indicate the mean

concentration for each sniff. **B)** The odor concentration profile was constructed in four steps: 1) A

smooth concentration gradient was computed; 2) multiplicative random noise was added to the

gradient; 3) 'Intermittency' was modeled by setting a proportion of pixels equal to zero. This

proportion increased with distance from the source. Finally, 4) Gaussian smoothing was

implemented to simulate averaging of the signal over a sniff. **C)** An example trajectory from the

concentration-sensitive model (CSM) that successfully found the odor spot (red circle).

Concentration difference (A.U.) is indicated. **D)** The success rate of the CSM was above 75%,

while the CRW control model had a success rate of below 25%. **E)** The nose displacement divided

by the initial distance to the spot was lower for the CSM than for the CRW control, indicating more

direct travel to the odor spot using concentration to guide the model. **F)** The survival function of

approach distances shows that nearly all CSM trajectories approach within 30cm of the source,

while CRW control trajectories are more spread out across the entire table. **G)** The probability of

finding the odor spot was computed for the models as in Figure 2E. With this measure, the CSM

was more successful at finding odor spots at all distances from the spot compared to CRW

controls.

Figure 6. Detailed analysis of CSM trajectories. A) The orientation of the CSM model relative

to the odor spot resembled that of mice (compare Figure 3D). **B)** The occupancy of the CSM

resembled that of mice (magenta line; compare red line Figure 3B), and significantly diverged from

the CRW control (gray line) at around 20cm from the odor spot. **C)** The mean nose velocity of the

CSM (magenta line) decreased as the model approached the odor spot, consistent with what was

observed with mice (compare red line Figure 3C). The nose velocity of the CSM was different from

the CRW control (gray line). **D)** In contrast to mice, successful trials of the CSM (magenta line)

showed *less* exploration than unsuccessful trials (gray line) (compare red line to blue line in Figure

3A). **E)** Casting increased as the CSM approached the spot (magenta line) consistent with

observations in mice (compare red line Figure 4A). Casting in the CSM increased with respect to

the CRW control (gray line) at a distance of 7cm.

Figure 7. Analysis of reduced models and parameter sweeps. A) The full model (CSM)

included three components: 1) a concentration-dependent velocity term (V), 2) a concentration-

dependent casting term (C); 3) a binaral-olfaction term (S). Removal of any of these components

(-V), (-B), or (-C) may, in principle, reduce model performance. Removal of concentration-

dependent velocity (-V) had the largest effect on performance. Removal of concentration-

dependent casting (-C) had a smaller effect on performance. Removal of binaral-olfaction (-B) had

the smallest effect on performance. Removal of both concentration-dependent velocity and casting

(-VC) had a multiplicative effect on performance. Removal of all three components (-VBC) was still

significantly better than CRW controls. **B)** Comparison of noise effects on CSM success rate

shows that the CSM model is robust for a wide array of different noise conditions. **C)** Because (-B)

had the smallest effect on performance, we wished to investigate its role further. We added a

coefficient in front of the hyperbolic tangent term in eq n5 and set it equal to 3. With this additional

term, success rate reduction from removing the binaral term increased. However, success

reduction from removing (-V) or (-C) terms decreased. This suggests that binaral sniffing can

compensate somewhat for lack of velocity or casting changes.

SUPPLEMENTAL CAPTIONS

Figure S1. Example Traces. Successful trials (left and center panels) have the time to spot

indicated by the text. The relative time of each position sample is indicated by the color.

Unsuccessful trials (right panel) have the total trial time indicated by the text. Color represents the

relative time of each position sample. Spot positions are indicated by red circles.

Figure S2. Detailed model geometry. A) Model specification of the body position (x,y), and the

body angle θ superimposed on a diagram of the mouse. The nose position (x_n, y_n) is the mean

position between the left and right nares. **B)** Nose angle ϕ (gray dashed line) is the angle between

θ and nose position. Nose angle ϕ is constrained by a maximum angle ϕ_{max} . The head-body

length ℓ is the distance between the body position and the nose position. Nose position is sub-

divided into two nostrils (left indicated by red, right indicated by green), separated by a distance

d_{nares}. The angle from the center of the nose to the edge of the nostril is γ. **C)** The distribution of

mouse nose deflection angles ϕ (blue bars) is approximately Gaussian (dashed red line; a*exp(-

((x-b)/c)^2); a=0.04, b=0.06, c=1.92).

Figure S3. Additional Casting Analyses. A) Curvature as measured by $\log_{10}(|dPhi|/nV)$ depends

on nose velocity. The colormap indicates the percentage of samples in each bin. **B)** Averaged

across successful trials, nose deflection angle ϕ increases as the distance to the spot decreases

(line=mean±s.e.m; ANOVA; p=2.18E-56, F=12.82, df=28).

Table S1. Logistic regression analysis on spot-finding success in mice. Significant variables in bold.

Table S2. Logistic regression analysis of CSM parameters on spot-finding success. Significant parameters are in bold.

ABBREVIATIONS

CSM – concentration-sensitive model

CRW – correlated random walk

OU – Ornstein-Uhlenbeck

Δt – change in time (s)

ℓ – nose-to-body length parameter (cm)

v_{max} – maximum velocity parameter (cm/s)

k – concentration-dependent parameter

σ_{min} – concentration-dependent parameter

σ_{max} – concentration-dependent parameter

$k_{binaral}$ – binaral-olfaction parameter

d_{nares} – inter-nostril parameter (cm)

ANOVA – Analysis Of VAriance

LED – light emitting diode

|dPhi|/nV – curvature

x' – derivative (of x)

x_n – nose coordinate

θ – heading of model body (degrees)

φ – heading of model nose relative to body (degrees)

C_t – concentration magnitude (A.U.)

η_t – multiplicative uniform noise

(-C) – CSM model without concentration-dependent casting, $\sigma = \sigma_{min}$

(-B) – CSM model without binaral-sniffing component, $k_{binaral}=0$

(-V) – CSM model without concentration-dependent velocity, $v=v_{max}$

IR – infrared

REFERENCES

Baker, K. L., Dickinson, M., Findley, T. M., Gire, D. H., Louis, M., Suver, M. P., . . . Smear, M. C. (2018). Algorithms for Olfactory Search across Species. *Journal of Neuroscience, 38*(44), 9383-9389. doi:10.1523/JNEUROSCI.1668-18.2018

Balkovsky, E., & Shraiman, B. I. (2002). Olfactory search at high Reynolds number. *Proceedings of the National Academy of Sciences of the United States of America, 99*(20), 12589-12593. doi:10.1073/pnas.192393499

Ben-Shaul, Y. (2017). OptiMouse: a comprehensive open source program for reliable detection and analysis of mouse body and nose positions. *BMC Biol, 15*(1), 41. doi:10.1186/s12915-017-0377-3

Bhattacharyya, U., & Bhalla, U. S. (2015). Robust and Rapid Air-Borne Odor Tracking without Casting. *eNeuro, 2*(6). doi:10.1523/ENEURO.0102-15.2015

Budick, S. A., & Dickinson, M. H. (2006). Free-flight responses of Drosophila melanogaster to attractive odors. *Journal of Experimental Biology, 209*(Pt 15), 3001-3017. doi:10.1242/jeb.02305

Carde, R. T., & Willis, M. A. (2008). Navigational strategies used by insects to find distant, wind-borne sources of odor. *Journal of Chemical Ecology, 34*(7), 854-866. doi:10.1007/s10886-008-9484-5

Catania, K. C. (2013). Stereo and serial sniffing guide navigation to an odour source in a mammal. *Nat Commun, 4*, 1441. doi:10.1038/ncomms2444

Charlton, R. E., Kanno, H., Collins, R. D., & Carde, R. T. (1993). Influence of Pheromone Concentration and Ambient-Temperature on Flight of the Gypsy-Moth, Lymantria-Dispar (L), in a Sustained-Flight Wind-Tunnel. *Physiological Entomology, 18*(4), 349-362.

Connor, E. G., McHugh, M. K., & Crimaldi, J. P. (2018). Quantification of airborne odor plumes using planar laser-induced fluorescence. *Experiments in Fluids, 59*(9). doi:ARTN 13710.1007/s00348-018-2591-3

Draft, R. W., McGill, M. R., Kapoor, V., & Murthy, V. N. (2018). Carpenter ants use diverse antennae sampling strategies to track odor trails. *Journal of Experimental Biology, 221*(22). doi:UNSP jeb18512410.1242/jeb.185124

Duval, L. (2018). BEADS: Baseline Estimation And Denoising with Sparsity, version 1.7.0.1. https://www.mathworks.com/matlabcentral/fileexchange/49974-beads-baseline-estimation-and-denoising-with-sparsity

Gire, D. H., Kapoor, V., Arrighi-Allisan, A., Seminara, A., & Murthy, V. N. (2016). Mice Develop Efficient Strategies for Foraging and Navigation Using Complex Natural Stimuli. *Curr Biol, 26*(10), 1261-1273. doi:10.1016/j.cub.2016.03.040

Hangartner, W. (1969). Structure and variability of the individual odor trail in Solenopsis geminata Fabr. (Hymenoptera Formicidae). *Z. Vgl. Physiol., 62*, 111-120.

Heitz, R. P. (2014). The speed-accuracy tradeoff: methodology, and behavior. *Frontiers in Neuroscience, 8*. doi:ARTN 15010.3389/fnins.2014.00150

Hepper, P. G., & Wells, D. L. (2005). How many footsteps do dogs need to determine the direction of an odour trail? *Chemical Senses, 30*(4), 291-298. doi:10.1093/chemse/bji023

1027 Hills, T. T., Todd, P. M., Lazer, D., Redish, A. D., Couzin, I. D., & Grp, C. S. R. (2015). Exploration versus
1028   exploitation in space, mind, and society. *Trends in Cognitive Sciences, 19*(1), 46-54.
1029   doi:10.1016/j.tics.2014.10.004
1030
1031 Jacobs, L. F. (2012). From chemotaxis to the cognitive map: The function of olfaction. *Proceedings of the*
1032   *National Academy of Sciences of the United States of America, 109*, 10693-10700.
1033   doi:10.1073/pnas.1201880109
1034
1035 Janabi-Sharifi, F., Hayward, V., & Chen, C. S. J. (2000). Discrete-time adaptive windowing for velocity
1036   estimation. *Ieee Transactions on Control Systems Technology, 8*(6), 1003-1009. doi:Doi
1037   10.1109/87.880606
1038
Kennedy, J. S. (1983). Zigzagging and Casting as a Programmed Response to Wind-Borne Odor - a
Review. *Physiological Entomology, 8*(2), 109-120. doi:DOI 10.1111/j.1365-3032.1983.tb00340.x
Khan, A. G., Sarangi, M., & Bhalla, U. S. (2012). Rats track odour trails accurately using a multi-layered
strategy with near-optimal sampling. *Nature Communications, 3*. doi:ARTN
70310.1038/ncomms1712
Kikuta, S., Sato, K., Kashiwadani, H., Tsunoda, K., Yamasoba, T., & Mori, K. (2010). From the Cover:
Neurons in the anterior olfactory nucleus pars externa detect right or left localization of odor sources.
Proc Natl Acad Sci U S A, 107(27), 12363-12368. doi:10.1073/pnas.1003999107
1047
1048 Koehl, M. A., Koseff, J. R., Crimaldi, J. P., McCay, M. G., Cooper, T., Wiley, M. B., & Moore, P. A. (2001).
1049   Lobster sniffing: antennule design and hydrodynamic filtering of information in an odor plume.
1050   *Science, 294*(5548), 1948-1951. doi:10.1126/science.1063724
1051
1052 Kuenen, L. P. S., & Carde, R. T. (1994). Strategies for Recontacting a Lost Pheromone Plume - Casting and
1053   Upwind Flight in the Male Gypsy-Moth. *Physiological Entomology, 19*(1), 15-29. doi:DOI
1054   10.1111/j.1365-3032.1994.tb01069.x
1055
1056 Manurung, A. (2016). Numerical methods for velocity estimation.
1057   https://www.mathworks.com/matlabcentral/fileexchange/52381-numerical-methods-for-velocity-
1058   estimation
1059
1060 Marsh, D., Kennedy, J. S., & Ludlow, A. R. (1978). An analysis of anemotactic zigzagging flight in male
1061   moths stimulated by pheromone. *Physiol. Entomol., 3*, 221-240.
1062
1063 Martin, H. (1965). Osmotropotaxis in Honey-Bee. *Nature, 208*(5005), 59-&. doi:DOI 10.1038/208059a0
1064
1065 Means, L. W., Alexander, S. R., & Oneal, M. F. (1992). Those Cheating Rats - Male and Female Rats Use
1066   Odor Trails in a Water-Escape Working Memory Task. *Behavioral and Neural Biology, 58*(2), 144-
1067   151. doi:Doi 10.1016/0163-1047(92)90387-J
1068
1069 Papale, A. E., Stott, J. J., Powell, N. J., Regier, P. S., & Redish, A. D. (2012). Interactions between
1070   deliberation and delay-discounting in rats. *Cognitive Affective & Behavioral Neuroscience, 12*(3),
1071   513-526. doi:10.3758/s13415-012-0097-7
1072
1073 Pierce-Shimomura, J. T., Morse, T. M., & Lockery, S. R. (1999). The fundamental role of pirouettes in
1074   Caenorhabditis elegans chemotaxis. *Journal of Neuroscience, 19*(21), 9557-9569.
1075
Porter, J., Craven, B., Khan, R. M., Chang, S. J., Kang, I., Judkewitz, B., . . . Sobel, N. (2007). Mechanisms
of scent-tracking in humans. *Nature Neuroscience, 10*(1), 27-29. doi:10.1038/nn1819

Schwenk, K. (1994). Why snakes have forked tongues. *Science, 263*(5153), 1573-1577.
doi:10.1126/science.263.5153.1573

Steck, K., Knaden, M., & Hansson, B. S. (2010). Do desert ants smell the scenery in stereo? *Animal
Behaviour, 79*(4), 939-945. doi:10.1016/j.anbehav.2010.01.011

Sutton, R. S., & Barto, A. G. (1998). *Reinforcement learning : an introduction*. Cambridge, Mass.: MIT
Press.

Tao, L. Y., Ozarkar, S., Beck, J. M., & Bhandawat, V. (2019). Statistical structure of locomotion and its
modulation by odors. *Elife, 8*. doi:ARTN e4123510.7554/eLife.41235

Thesen, A., Steen, J. B., & Doving, K. B. (1993). Behavior of Dogs during Olfactory Tracking. *Journal of
Experimental Biology, 180*, 247-251.

Uchida, N., & Mainen, Z. F. (2003). Speed and accuracy of olfactory discrimination in the rat. *Nature
Neuroscience, 6*(11), 1224-1229. doi:10.1038/nn1142

Vergassola, M., Villermaux, E., & Shraiman, B. I. (2007). 'Infotaxis' as a strategy for searching without
gradients. *Nature, 445*(7126), 406-409. doi:10.1038/nature05464

Vickers, N. J. (2000). Mechanisms of animal navigation in odor plumes. *Biological Bulletin, 198*(2), 203-212.
doi:Doi 10.2307/1542524

Wallace, D. G., Gorny, B., & Whishaw, I. Q. (2002). Rats can track odors, other rats, and themselves:
implications for the study of spatial behavior. *Behavioural Brain Research, 131*(1-2), 185-192. doi:Pii
S0166-4328(01)00384-9Doi 10.1016/S0166-4328(01)00384-9

Wesson, D. W., Donahou, T. N., Johnson, M. O., & Wachowiak, M. (2008). Sniffing behavior of mice during
performance in odor-guided tasks. *Chemical Senses, 33*(7), 581-596. doi:10.1093/chemse/bjn029

Willis, M. A., Avondet, J. L., & Finnell, A. S. (2008). Effects of altering flow and odor information on plume
tracking behavior in walking cockroaches, Periplaneta americana (L.). *Journal of Experimental
Biology, 211*(Pt 14), 2317-2326. doi:10.1242/jeb.016006

Wilson, D. A., & Sullivan, R. M. (1999). Respiratory airflow pattern at the rat's snout and an hypothesis
regarding its role in olfaction. *Physiology & Behavior, 66*(1), 41-44. doi:Doi 10.1016/S0031-
9384(98)00269-8

Zhang, S. J., & Manahan-Vaughan, D. (2015). Spatial Olfactory Learning Contributes to Place Field
Formation in the Hippocampus. *Cerebral Cortex, 25*(2), 423-432. doi:10.1093/cercor/bht239

Zhao, W. Q., Gong, C. X., Ouyang, Z. H., Wang, P. F., Wang, J., Zhou, P. P., . . . Gong, Z. F. (2017). Turns
with multiple and single head cast mediate Drosophila larval light avoidance. *Plos One, 12*(7).
doi:ARTN e018119310.1371/journal.pone.0181193

1121

1122

1123

1124

1125

1126

1127

FIGURE 1

1129

1130

1131
1132
1133
1134
1135 **FIGURE 2**
1136

A

C   D

E

1137
1138
1139

1140

1141

1142

FIGURE 3

1143

1144

A

B

C

D

E

1145

1146

1147

1148

1149

1150

1151

1152 **FIGURE 4**

1153

1154

1155

1156

1157

1158

1159

1160

FIGURE 5

1166

1167

FIGURE 6

1169

1170

1171

1172

1173

FIGURE 7

1176

1177 **SUPPLEMENTAL FIGURE 1**

1188

1189 **SUPPLEMENTAL FIGURE 2**

1190

A

B

C

1191

1192

1193

1194

SUPPLEMENTAL FIGURE 3

1196

1198

1199

1200 **SUPPLEMENTAL TABLE 1**

Variable	p value	β value
Experimenter	4.20E-01	-0.43
Baited	**4.68E-12**	**2.51**
Concentration	4.93E-03	0.56
Order of Trial	0.08	0.23
Trial	0.98	4.47E-03
Mouse	0.11	**0.16**
Initial Distance	**6.54E-08**	**4.53E-02**
Initial Orientation	0.79	3.26E-04
Mean Body Velocity	**6.89E-06**	**-0.29**
Mean Nose Velocity	**1.07E-04**	**0.27**
Displacement Ratio	**4.19E-19**	**-0.38**
Mean Curvature	**2.56E-03**	**1.4**
Exploration %	**1.65E-15**	**15.51**
Mean Nose Acceleration	0.01	-0.05
Time Stopped	0.22	-0.02

1201

1202

1203

1204

1205

1206

1207

1208

1209

1210

1211

1212

1213

1214

1215

1216

1217

 SUPPLEMENTAL TABLE 2

Variable	p value	β value	Range
d_{nares}	**1.00E-50**	**0.54**	**0.01 - 1cm**
k_c	4.84E-03	-0.1	0 - 1
$k_{binaral1}$	**6.86E-26**	**1.27E-03**	**0.01 - 300**
$k_{binaral2}$	**7.33E-106**	**0.16**	**0 - 5**
k_v	**2.63E-47**	**0.62**	**0 - 1**
ℓ (neck length)	**1.35E-191**	**0.26**	**0.01 - 10cm**
Mean Curvature (dist < 30cm)	**6.13E-09**	**0.6**	**Dependent Variable**
Mean Nose Velocity (dist < 30cm)	**2.30E-19**	**-0.016**	**Dependent Variable**
n_c	**4.67E-09**	**-0.043**	**0 - 5**
n_v	**4.76E-160**	**0.22**	**0 - 5**
σ_{max}	**1.11E-39**	**0.77**	**0.01 - 1**
σ_{min}	2.50E-01	0.074	0.01 - 1
τ	6.40E-01	-0.02	0.01 - 1
V_{max}	**9.26E-54**	**0.03**	**20 - 40cm/s**

1219