Bacterial microbiota composition of *Ixodes ricinus* ticks: the role of environmental variation,

tick characteristics and microbial interactions

Tuomas Aivelo[1,2,*], Anna Norberg[2], Barbara Tschirren[3]

1: Department of Evolutionary Biology and Environmental Studies, University of Zürich

2: Organismal and Evolutionary Biology research program, University of Helsinki

3: Centre for Ecology and Conservation, University of Exeter, Treliever Road, Penryn, TR10 9FE,

United Kingdom

* Corresponding author

Address: Viikinkaari 1 (PL 56), 00014, Helsingin yliopisto, Finland

Email: tuomas.aivelo@helsinki.fi

Tel.: +358 40 828 6466

Running title: Species distribution modelling of tick microbiota

Competing interests: The authors declare no competing interests.

Abstract

Ecological factors, host characteristics and / or interactions among microbes may all shape the occurrence of microbes and the structure of microbial communities within organisms. In the past, disentangling these factors and determining their relative importance in shaping within-host microbiota communities has been hampered by analytical limitations to account for (dis)similar environmental preferences ('environmental filtering'). Here we used a joint species distribution modelling (JSDM) approach to characterize the bacterial microbiome of one of the most important disease vectors in Europe, the sheep tick *Ixodes ricinus,* along ecological gradients in the Swiss Alps. Although our study captured extensive environmental variation along elevational clines, the explanatory power of such large-scale ecological factors was comparably weak, suggesting that tick-specific traits and behaviours, microhabitat and -climate experienced by ticks, and interactions among microbes play an important role in shaping tick microbial communities. Indeed, when accounting for shared environmental preferences, evidence for significant patterns of positive or negative co-occurrence among microbes was found, which is indicative of competition or facilitation processes. Facilitation occurred primarily among human pathogens, leading to co-infection within ticks, whereas competition was observed between the tick endosymbiont *Spiroplasma* and human pathogens. These findings highlight the important role of small-scale ecological variation and microbe-microbe interactions in shaping tick microbial communities and the dynamics of tick-borne disease.

Keywords: disease ecology, tick-borne pathogens, species distribution modelling, community composition, Borrelia burgdorferi, Lyme disease, facilitation, endosymbionts

Introduction

Microbial communities within organisms consist of symbionts, commensals, mutualists and pathogens that co-occur simultaneously and potentially influence each other (Petney and Andrews 1998, Rigaud et al. 2010, Sofonea et al. 2015). These microbial communities may be shaped by a range of factors and processes, including the environment, host and microbe genetics and the occurrence and abundance of other microbial species (Adair and Douglas 2017). For example, certain microbial species might tolerate only specific biotic or abiotic conditions, which makes it more likely that species with similar requirements co-occur within a host ('environmental filtering', Dallas and Presley, (2014)). Similarly, the host's immune system can influence colonization success of microbes (Hawley and Altizer 2011), with cross-immunity preventing the colonization of different microbes with similar antigenic properties (Durand et al. 2015). Finally, direct interactions among microbes might affect colonization, or replication success after colonization, through competition or facilitation processes. Competition may occur when different microbes use the same, limited resources within a host (Lello et al. 2004), whereas facilitation may occur directly through the production of public goods (West and Buckling 2003) or indirectly through the modification of the host's physiology (Abraham et al. 2017) or immune defense (Rodríguez et al. 1999).

Ixodes ricinus is the most common tick species in Europe and an important vector for a range of human and wildlife pathogens (Medlock et al. 2013). Its distribution and abundance are strongly influenced by environmental conditions, in particular temperature and humidity (Cortinas et al. 2002, Gatewood et al. 2009). Previous studies on the bacterial community composition of *I. ricinus* ticks have found approximately 100 OTUs (operational taxonomic units; Stackebrandt and Goebel 1994) per tick (Greay et al. 2018). Most of them are environmental and free-living bacteria but also several endosymbionts and human or wildlife pathogens were found, including *Borrelia* (Mannelli et al. 2012), *Rickettsia* (Venclikova et al. 2014), *Anaplasma* (Jahfari et al. 2014) and *Candidatus* Neoehrlichia (Kawahara et al. 2004).

Differences in the bacterial community structure and composition of ticks across habitats (Estrada-

Peña et al. 2018), geographical sites (Carpi et al. 2011), and tick life stages and sexes (Carpi et al.

2011, Vayssier-Taussat et al. 2013) have been documented. Large-scale biotic or abiotic factors

such as vegetation structure, elevation, temperature or rainfall may influence tick microbial

communities directly, or indirectly through effects on tick physiology or activity patterns (van

Treuren et al. 2015) or via influencing the distribution and abundance of tick hosts species

(Randolph et al. 1999, MacDonald et al. 2017). Small-scale and/or tick-specific effects on microbial

communities may be explained by microhabitat or microclimatic conditions experienced by

individual ticks (Gern et al. 2008), individual tick behavior or genetics (Hawlena et al. 2013), direct

interactions among microbes (Moutailler et al. 2016) or parallel acquisition from a host during a

bloodmeal (Andersson et al. 2014, Belli et al. 2017, Swei and Kwan 2017).

Currently, the relative importance of these factors in shaping tick microbial communities is not well

understood, which hampers progress in our understanding of the processes shaping microbial

communities in nature and predicting the occurrence of specific microbes (e.g., human pathogens).

Elevational gradients are excellently suited to quantify the importance of large-scale ecological

variation in shaping tick bacterial microbiota because they cover a large range of environmental

conditions within a small geographical area. Furthermore, including replicated transects along

gradients allow us to quantify the robustness of ecological associations within sites and along

elevational clines on tick microbial communities.

Ixodes ticks are commonly found to be co-infected with several human or wildlife pathogens

(Andersson et al. 2013, Michelet et al. 2014, Diuk-Wasser et al. 2016, Moutailler et al. 2016).

Currently, it is unknown whether these co-infection patterns are caused by similar environmental

preferences of pathogens, parallel acquisition from host communities or direct microbe-microbe

interactions within ticks. Yet, previous studies suggest that the latter process, (i.e., facilitation and

competition processes among microbes) may play a role in shaping the microbial communities

(Haine 2008, Bonnet et al. 2017). For example, it has been found that pathogenic *Rickettsia* species

prevent co-infection with other *Rickettsia* species in *Dermacentor variabilis* ticks (Macaluso et al.

2002), whereas the presence of *Francisella* sp. endosymbionts increases the colonization success of

pathogenic *Francisella novicida* in *D. andersoni* ticks (Gall et al. 2016). Facilitation has also been

suggested to promote co-infection with different *Borrelia afzelii* strains in *Ixodes ricinus* ticks

(Andersson et al. 2013). Most strikingly, dysbiosis in *I. scapularis* ticks (i.e., ticks with low

microbiotal diversity) leads to a defective peritrophic matrix which decreases the colonization

success of *B. burgdorferi* s.s., suggesting that the pathogen requires the presence of an intact

microbiota to be able to invade ticks (Narasimhan et al. 2014). Thus, the microbial community may

have a crucial impact on vector competence of ticks and thereby on disease dynamics.

Yet, as outlined above, co-occurrence of microbes can be due to environmental filtering or direct

microbial interactions, and distinguishing between these processes is non-trivial. Indeed, previous

studies that have documented pathogen co-occurrence in ticks have not accounted for potential

confounding variables such as shared ecological requirements, and are thus limited in their ability to

differentiate between accidental co-occurrences due to shared environmental niches, and co-

occurrence shaped by facilitation or competition among microbes.

To address these gaps, we exploited the substantial environmental heterogeneity along replicated

elevational gradients in the Swiss Alps to quantify the relative importance of environmental factors,

tick characteristics and direct microbial interactions in influencing the structure of bacterial

communities in *I. ricinus* ticks in general, and the (co-)occurrence of human or wildlife pathogens

in particular, using a combination of 16S sequencing and joint species distribution modelling

(JSDM) (Ovaskainen et al. 2015, Warton et al. 2015). By taking shared environmental preferences

into account, JSDMs allows to identify residual co-occurrence patterns among microbes that can

result from unaccounted environmental effects or direct microbial interactions.

Specifically, we ask (i) how do large-scale abiotic factors and small scale tick-level variables affect

tick microbiota composition, (ii) which large-scale abiotic and small-scale tick-level variables

predict pathogen occurrence, and (iii) are there patterns of non-random microbial co-occurrence that

cannot be explained by environmental responses, implying unmeasured variables, such as microbial

interactions.

Materials and methods

Tick sampling

Questing *Ixodes ricinus* ticks were collected at three locations in the Swiss Alps (Kanton

Graubünden). At each location, one site at low (630 - 732 m above sea level, masl), one at medium

(1 094 – 1 138 masl) and one at high (1 454 – 1 673 masl) elevation were identified (Fig. 1, Table 1,

N = 9 sampling sites). At each site, questing ticks were sampled thrice, once in June, once in July,

and once in August 2014 by dragging a white blanket (1 m x 1 m) over the ground vegetation as

described previously (Lemoine et al. 2018). Ticks were collected from the blanket and stored in

95% ethanol. Tick species, sex and life stage were verified by morphological features following

(Hillyard 1996) using a stereomicroscope.

Environmental variables

For each sampling site, we compiled information on large-scale, site-level ecological variables by

obtaining data on elevation, slope and aspect using DHM25, land use data from swissTLM3D (both

from Federal Office of Topography swisstopo) and data on temperature and precipitation from

Landscape Dynamics (Swiss Federal Research Institute for Water, Snow and Landscape Research

WSL and Federal Office of Meteorology and Climatology MeteoSwiss, (Thornton et al. 1997)).

Data on *I. ricinus* abundance and the abundance of a key tick host, the bank vole (*Myodes*

glareolus), as well as the ratio of bank vole to other rodents at our sampling sites were obtained

from (Cornetti et al. 2018). Details on the different variables and a justification why they were

included to characterise large-scale ecological conditions is provided in the Supplementary

Material.

Tick DNA isolation and quantification of neutral genetic diversity

The number of analysed ticks per site is presented in Table 1. Although we aimed to include similar

numbers of *I. ricinus* ticks from each sampling site and month, variation in the number of ticks per

site (Table 1) was unavoidable because of variation in tick abundance across sites (Lemoine et al.

2018). To avoid environmental contamination, we performed DNA isolation and amplifications in a

laminar flow cabinet. Each tick was washed thrice with sterile water before sterilizing it with 3%

hydrogen peroxide. Ticks were then cut in half with a sterilized blade to facilitate DNA isolation.

DNA was extracted using DNeasy Blood & Tissue kit (Qiagen; Hilden, Germany).

Host genetics may affect pathogen and endosymbiont colonisation and replication success (Archie

and Ezenwa 2011). In order to quantify individual and population-level genetic diversity, we

genotyped ticks at 11 microsatellite markers in two multiplexed amplifications (see Supplementary

Material for details). Not all markers were successfully amplified in all samples, but none of the

samples contained more than two failed markers. We used package *poppr* (Kamvar et al. 2014) in R

3.4.1 (Team 2013) to test for linkage disequilibrium and deviation from Hardy-Weinberg

equilibrium. Individual observed heterozygosity was determined for each tick as a proportion of

heterozygous markers to all successfully amplified markers. Expected population level

heterozygosity was determined with *poppr*. The former was used as a tick-level explanatory

variable (together with tick sex and life stage), the latter was used as a site-level explanatory

variable.

Tick microbiota sequencing

We quantified tick bacterial community composition by sequencing the hypervariable V3-V4

region of the 16S rRNA (16S) gene. Negative controls (extraction reagent blank, N=2 and PCR

controls, N=3) were processed alongside the tick samples. Sequencing libraries were prepared

following the Earth Microbiome 16S Illumina Amplicon protocol, using the primers 515FB and

806RB (Carey et al. 2013) (see Supplementary Material for details). Samples and negative controls

were randomized across two plates. The libraries were sequenced on Illumina MiSeq at the

Functional Genomic Center Zurich with a target length of 250 bp following the manufacturer's

protocol. The obtained sequence data were analyzed following the *mothur* pipeline with MiSeq

standard operation procedures (Kozich et al. 2013). Sequences have been deposited to the Sequence

Read Archive under BioProject PRJNA506875. The complete metadata of the samples and their

matching sequence accession numbers have been submitted to FigShare (doi:

10.6084/m9.figshare.7380767).

A special focus of our analysis was on tick endosymbionts and tick-borne human or wildlife

pathogens (Table 2), which are obligate residents. Identification of endosymbionts and pathogens is

described in Supplementary Material.

Joint species distribution modelling of microbiota composition

Only samples with > 500 amplicons and OTUs which were present in at least two samples were

included in the analyses (Table 1). As the most common OTU, the intra-mitochondrial

endosymbiont *Candidatus* Midichloria (Lo et al. 2006), was present in all samples, it was not

included in the modelling. For the occurrence matrix, an OTU was determined to be present in a

tick if >5 amplicons were identified in a sample (Aivelo and Norberg 2017).

We used a JSDM framework called Hierarchical Modelling of Species Communities (HMSC,

Ovaskainen et al. 2017a) to examine how environmental variables correlate with human or wildlife

pathogen and tick endosymbiont occurrence in ticks, and whether there are non-random residual

associations among different OTUs and/or oligotypes, implying direct facilitation or competition

effects among microbes. This approach combines information on environmental covariates,

bacterial species traits, spatiotemporal context and sampling design to explain the presence or

absence of OTUs (Fig. S1). The associations among OTUs are captured with the latent part of the

framework, modelling the residual variance after accounting for the effects of the environment with

latent variables. The estimates for these latent variables can be then translated into residual

correlations among response variables, i.e. OTUs and/or oligotypes. These correlations thus reflect

(dis)associations which cannot be explained by shared responses to the environment.

We compiled occurrence matrices for OTUs for each individual tick as a response variable. For

each sampling unit, i.e. a row in our response variable matrix, we included information on the

identity of the sampling unit (tick ID), its location, sampling site and month, describing the study

design. To reach a better resolution within specific OTUs, we analyzed known human or wildlife

pathogens, tick endosymbionts and their close relatives within the 100 most common OTUs with

oligotyping pipeline (Eren et al. 2014). Oligotyping uses all the sequences, which form an OTU,

and performs Shannon Entropy Analysis to regroup sequences based on within-OTU variation. This

results in higher-resolution grouping than OTUs as the different oligotypes might differ only by a

single nucleotide (Eren et al. 2014). We used the standard operation procedures of the oligotyping

pipeline software (Eren et al. 2013).

Including a large number of explanatory variables in statistical models is inherently challenging. To

reduce the number of variables, while maintaining their information value, we used two variable

sets in the model: a) a set of full-effect explanatory variables, and b) explanatory variables under

variable selection (Ovaskainen et al. 2017a). The full-effect variable set included an intercept, two

tick-level variables (tick sex or life stage and individual heterozygosity) and two site-level variables

(tick abundance and elevation of the sampling site). Additionally, we included information whether

208 a specific OTU is an endosymbiont and/or a human or wildlife pathogen (Abrego et al. 2016). This

209 allowed us to test if endosymbionts and/or human or wildlife pathogens respond differentially to

210 environmental conditions than other OTUs. The set of explanatory variables under variable

211 selection included additional information on the environmental conditions of the sites (namely the

212 number of days above 7 °C during the year, monthly precipitation, mean monthly temperature,

213 forest coverage, slope, aspect, bank vole abundance, the proportion of voles to other rodents and

214 expected tick heterozygosity) (Table S1). We considered all parameter estimates, including

215 associations among bacterial OTUs, having strong statistical support and thus being "significant" if

216 the 90% central credible interval of the parameter did not overlap with zero (following Aivelo and

217 Norberg 2017; see Supplementary Material for additional model details).

218 Although JSDM is a powerful approach to model community structure, it has a number of

219 limitations. First, it assumes that interactions among microbes are similar across environments (but

220 see Tikhonov et al. 2017). This is not necessarily the case as both abiotic and biotic factors may

221 shape microbial interactions (Elliot et al. 2002). Second, the model assumes that the explanatory

222 variables affect the microbial community composition (or rather, the presence or absence of

223 individual OTUs), but not vice versa. However, this is a valid assumption for most environmental

224 (e.g. elevation and temperature) and tick-related variables (e.g. tick sex, life stage) included in our

225 models. Thirdly, covariation among explanatory variables poses a problem to any correlative

226 modelling approach. Our model is built on two distinct variable sets to aid in handling such

227 covariation: the full variable set includes elevation, whereas the variables with the strongest

228 covariation (i.e., temperature and precipitation) are included in the variable selection set.

230 **Results**

231 Ixodes ricinus *microbiota composition*

We 16S sequenced the bacterial community of 92 *Ixodes ricinus* ticks which resulted in 13 214 477

amplicons. No amplification was observed in the negative controls (i.e., their sequencing did not

result in any sequences). After contig assembly and quality control 1 656 287 sequences were

retained. Most of the discarded amplicons were either shorter than 250 bp or chimeras. There was a

median of 1 562 quality-controlled amplicons per sample, with an interquartile range of 6 319. 82

samples with more than 500 amplicons per sample, a plateauing accumulation curve and a Good's

coverage estimator ≥0.95 were included in the subsequent analyses. In total, 5 181 bacterial OTUs

were identified. The median number of OTUs when rarified to 500 amplicons per sample was 89

OTUs, with a 95% confidence interval of 78.3 - 98.5 OTUs.

Six OTUs were present in at least 90% of the samples: *Ca.* Midichloria (Otu0001), *Sphingomonas*

(Otu0002, 0006 and 0007*)*, *Pseudomonas* (Otu0011) and *Delftia* (Otu0012). Together, they

represented 50.2 % of all amplicons. We used oligotyping to further divide OTU0031 '*Rickettsia*'

into two oligotypes labelled as '*R. helvetica*' and '*R. monacensis*', and OTU0086 '*Borrelia*' into

four oligotypes labelled as '*B. afzelii*', '*B. valaisian*a' and '*B. garinii*' and '*B. miyamotoi*'. After

excluding rare OTUs, 635 OTUs and oligotypes were used in subsequent analyses, including 14

endosymbionts and / or human or wildlife pathogens (Table 2).

Tick microbiota variance partitioning

Variance partitioning revealed that most of the variation in tick microbiota composition explained

by our model related to tick ID: for the hundred most common OTUs, tick ID accounted for 64.1%

of the variation explained by the model. Fixed effects (e.g., tick life stage, elevation; see Table S1)

accounted for 12.5% (tick-level: 7.3%, site-level: 5.2%) and spatial and temporal random effects

(i.e., location, site and month) explained 23.3% (Fig. 2). This suggests that there is extensive tick-

level variation which cannot be accounted for by tick-specific characteristics included in our model

(i.e., sex, life stage, genetic diversity) or site-level environmental factors. The pattern differed

slightly for endosymbionts and human or wildlife pathogens: while tick ID was still the most

important variable explaining 49.9%, fixed effects explained 31.8% (tick-level: 17.5%, site-level:

14.3%) and spatial and temporal random effects explained 18.3% of the total variation explained by

the model, when averaged over all pathogens and endosymbionts (Fig. 2).

Tick-specific and environmental factors related to OTU occurrence

The occurrence of tick endosymbionts and human or wildlife pathogens was strongly associated

with specific explanatory variables, yet associations were typically microbe-specific rather than

universal (Table 3). The two most important variables explaining the presence or absence of tick

endosymbionts and human or wildlife pathogens were tick sex and elevation of the sampling site:

adult female ticks were less likely to harbour the endosymbionts *Spiroplasma*, *Rickettsiella*,

Lariskella and *Rickettsia* spp. (Table 3), and ticks at higher elevations had higher probability to

harbour *R. helvetica* and *R. monacensis*, but were less probable to harbour *B. garinii* (Table 3).

Topography was also a significant predictor of tick endosymbionts and human or wildlife pathogen

occurrence, with ticks from sites facing northwards having a higher probability of harbouring

Spiroplasma and *B. afzelii*, and ticks from sites on steeper slopes having a higher probability of

harbouring *Rickettsia sp*. (Table 3). Furthermore, a higher tick density was associated with a higher

probability of *Rickettsiella* and *Ca.* Neoehrlichia occurrence (Table 3). Relationships between tick

life stage, mean temperature, the number of days > 7 °C or forest cover and the occurrence of

specific OTUs were not strongly statistically supported.

The effect sizes of strongly statistically supported associations varied substantially (Fig. S3a-i). For

example, threefold increase in vole abundance corresponded to less than one percentage point

decrease of *R. monacensis* prevalence (Fig. S3b), whereas a threefold increase in tick abundance

corresponded to a threefold increase in *Neoehrlichia* prevalence from 8% to 27% (Fig. S3e).

Patterns of microbial association within ticks

Numerous bacterial OTUs were either significantly more or less likely to co-occur within a tick

than expected by chance after accounting for shared environmental preferences (Fig. 3a; Table S5).

At the level of the individual tick, the occurrence of the tick endosymbiont *Spiroplasma* was

negatively associated with the occurrence of the endosymbiont *Lariskella* and several tick-borne

pathogens, namely *Rickettsia* sp., *Ca*. Neoehrlichia and *B. miyamotoi* (Fig. 3a). Associations among

pathogens, if they occurred, were all positive (Fig. 3a), suggesting that ticks are more likely to be

co-infected with several human or wildlife pathogens simultaneously than expected by chance or

based on shared environmental preferences. *Borrelia* oligotypes showed positive co-occurrence

patterns among each other, except for *B. miyamotoi*, which was not associated with other *Borrelia*

sp., but negatively with *Spiroplasma* and positively with *Lariskella*. At the level of the sampling

site, significant associations were sparser. *Spiroplasma* was more likely to co-occur with *Lariskella*

and *Rickettsiella* across sites, whereas the latter two were less likely to co-occur across sites than

expected by chance after accounting for shared environmental preferences (Fig. 3b).

Discussion

We used a JSDM framework to quantify the relative importance of large scale, site-level

environmental variables, tick-level characteristics and interactions among microbes in shaping tick

microbiota composition along elevational gradients in the Swiss Alps. We show that although our

study captured extensive environmental variation, with sampling sites spanning across an

elevational gradient from 630 – 1 580 masl, and a large number of ecological variables was

considered in our models, the explanatory power of such large-scale ecological factors was

comparably weak. In contrast, individual tick ID explained over 60% of the variation in microbiota

composition. This substantial microbiota variation across individual ticks may be partly explained

by some of the bacteria present in ticks being non-resident (i.e., bacteria that were accidentally

obtained from the environment). Indeed, there has been a debate whether ticks have a stable

microbiota (Ross et al. 2017), mirroring the wider debate on how common resident microbiota is in

arthropod hosts (Hammer et al. 2017).

However, also for endosymbionts and human or wildlife pathogens, which are obligate resident,

tick ID accounted for half of the variation in occurrence, suggesting that microhabitat or -climatic

conditions experienced by individual ticks, tick-specific traits and behaviors not included in our

models, as well as microbial interactions within ticks such as facilitation and competition (Abraham

et al. 2017, Gurfield et al. 2017), play a crucial role in shaping microbiota composition and the

occurrence of endosymbionts and human or wildlife pathogens in *I. ricinus*. Focusing on such

small-scale variables, rather than large-scale climatic or environmental factors as is usually done

when modelling tick-borne pathogen prevalence (Norman et al. 2016, Rosà et al. 2018), is thus

likely a more fruitful approach to advance our understanding of microbiota composition of natural

populations as well as (tick-borne) disease dynamics.

Co-occurrence of human or wildlife pathogens in ticks has been documented previously, both in *I.*

ricinus (Lommano et al. 2012, Michelet et al. 2014) and other tick species (Gurfield et al. 2017,

Laaksonen et al. 2018). Yet, previous studies did not control for shared environmental preferences

(i.e., environmental filtering), which limited their ability to disentangle shared responses to the

environment from direct microbe-microbe interactions. Our study revealed that when accounting for

shared environmental preferences, associations among human or wildlife pathogens were often

pronounced and mostly positive. These positive associations may result from direct facilitation

among microbes or parallel colonization from co-infected tick hosts.

Within ticks, the strong positive associations among the Lyme disease-causing *Borrelia* genospecies

(*B. afzelii*, *B. garinii* and *B. valaisiana*) were particularly striking. This positive co-occurrence is

surprising because *B. garinii* and *B. valaisiana* are bird specialists (Hanincova et al. 2003a,

Comstedt et al. 2011), whereas *B. afzelii* is a rodent specialist (Hanincova et al. 2003b). Thus, the

parallel colonization from co-infected tick hosts cannot explain this pattern. Rather the positive co-

occurrence is indicative of facilitation processes among *Borrelia* genospecies, as has been

suggested previously (Andersson et al. 2013). Such facilitation, and the resulting co-infection of

ticks with several *Borrelia* genospecies has implications for the severity, diagnosis, treatment and

control of Lyme disease. Finally, the co-occurrence of these different *Borrelia* genospecies suggests

that *I. ricinus* feeds on multiple, phylogenetically diverse host species during its life cycle and does

not show pathogen-mediated host specialization as has been suggested previously (McCoy et al.

2005, 2013).

Although associations among microbes were mostly positive, there were negative associations

between the tick endosymbiont *Spiroplasma* and several human or wildlife pathogens, which may

be explained by competition. The most common infection route for *Spiroplasma* is maternal (i.e.,

vertical) transmission (Herren and Lemaitre 2011), indicating that horizontal or environmental

transfer plays a minor role in its transmission. Protective effects of *Spiroplasma* have been

previously described in *Drosophila* spp., where *Spiroplasma* is associated with a decreased

probability of nematode and parasitoids infections (Xie et al. 2010, Jaenike et al. 2013). Although

the exact mechanisms mediating *Spiroplasma*-induced competition effects are currently unknown,

this finding may stimulate further research into the potential of tick endosymbionts to manage tick-

borne pathogens.

In contrast to the numerous positive or negative associations among microbes at the tick-level, little

statistical support for positive or negative microbial co-occurrence was found at the site-level, with

the exception of the associations among three endosymbionts. Interestingly, the pattern of co-

occurrence of *Spiroplasma* and *Lariskella* at the site-level was opposite from what was observed at

the tick-level. It suggests that *Spiroplasma* and *Lariskella* are more likely to co-occur at the same

sites but not within the same tick. Co-occurrence at the site-level is most likely due to an

environmental variable not included in our model, for which the three OTUs had similar responses.

Importantly, the substantial reduction of microbe-microbe associations at the site level highlights

relevant environmental variation across sites was captured in our models, strengthening our

conclusion that the patterns of positive or negative co-occurrence among microbes observed at the

tick-level can be due to direct microbial interactions.

Despite the large among-tick variation in microbiota composition, we identified a range

environmental variables that significantly predicted the occurrence of specific tick endosymbionts

and human or wildlife pathogens. However, the predictor variables as well as their effect were

typically OTU-specific rather than universal. For example, *B. garinii* was *less* likely to occur at

higher elevations, whereas *R. helvetica* and *R. monascensis* were *more* likely to occur at higher

elevations. Generally, the environmental factors shaping *Rickettsia* spp. distribution are poorly

understood, as is their range of host species (Halos et al. 2010, Eremeeva and Dasch 2015). Yet, it

has previously been found that spotted fever incidence in humans, caused by *R. ricketsii*, is highest

in areas or regions, where ticks are less common (Atkinson et al. 2013). This is in line with our

findings and suggests that *Rickettsia* spp. are more likely to colonize ticks living under suboptimal

conditions (e.g. at range edges).

The finding that *B. garinii* is less likely to occur at higher elevations is in line with previous

observations (Jouda et al. 2004a, Cornetti et al. 2018) and may be explained by changes in

vegetation structure and associated changes in host communities (Halos et al. 2010), in particular

changes in the diversity and/or abundance of birds, the natural hosts of *B. garinii* (Comstedt et al.

2011). In contrast, the occurrence of the mammal specialist *B. afzelii* was not related to elevation,

potentially because elevational clines in mammal diversity and/or abundance are less pronounced

(McCain 2005). Indeed, we did not observe an association between elevation and bank vole

abundance across our study sites (ANOVA: $F_{1,8}=0.357$, $p = 0.57$, $R^2 = 0.05$).

Interestingly, temperature and precipitation, which vary strongly across elevational gradients

(average temperature and precipitation: high elevation sites: 11.8 °C and 17.8 mm per month; in

low sites: 16.5 °C and 12.1 mm per month), were not significant predictors of the occurrence of

endosymbionts or human or wildlife pathogens, with the exception of precipitation correlating

positively with the probability of *Rickettsiella* occurrence. This may be partly explained by the

temperature and precipitation measures included in our models not fully capturing the microclimatic

variation across sites and along elevational clines. Indeed, slope and aspect, which are important

determinants of the topography, and thus microclimate (Bennie et al. 2008), were significant

predictors of pathogen and endosymbiont occurrence. The probability of *Rickettsia* sp. occurrence

was higher on steeper slopes. Furthermore, the probability of occurrence was higher on north-facing

slopes for *B. afzelii* and *Spiroplasma* and higher on south-facing slopes for *Rickettsiella* (see also

(Stuen et al. 2013)). Microclimatic conditions may affect microbial occurrence directly, or

indirectly via affecting tick behavior or host community composition (Swei et al. 2011, Lawson et

al. 2014). Furthermore, topography can affect population connectivity and dispersal in

metapopulation networks (Swei and Kwan 2017).

Previous work has found that tick abundance is a strong predictor of *Borrelia* spp. prevalence,

potentially because larger tick populations facilitate co-feeding transmission (Jouda et al. 2004b).

No relationship between *Borrelia* spp occurrence and tick abundance was observed in our study.

However, both *Ca.* Neoehrlichia and *Rickettsiella* were more common at sites where ticks were

more abundant, suggesting that co-feeding transmission may also play a role in the life cycle of

these microbes.

Finally, differences in host competence can lead to dilution effects and thus affect the prevalence of

tick-borne pathogens (Keesing et al. 2006). Whereas for some tick-borne pathogens the vertebrate

hosts are known or suspected (e.g. small mammals for *B. afzelii* (Hanincova et al. 2003b) and *Ca.*

Neoehrlichia (Jahfari et al. 2012), birds for *B. garinii* and *B. valaisiana* (Hanincova et al. 2003a),

both for *Anaplasma* (Keesing et al. 2012) and *R. helvetica* (Sprong et al. 2009)), for others the host

species range is less well understood (e.g. *B. miyamotoi* (Wagemakers et al. 2015)). The bank vole

is a common tick host at our study sites and their abundance was a significant negative predictor of

R. monacensis and *R. helvetica* occurrence. Interestingly, bank voles are not known hosts for either

(Burri et al. 2014). Most likely, the relation is thus indirect, explained by an unmeasured biotic or

abiotic variable that correlates with bank vole abundance. No evidence was found that the

proportion of bank voles to other rodents affects the prevalence of tick-borne pathogens.

In conclusion, our study demonstrates that a JSDM framework can contribute to a better

understanding of the factors shaping bacterial communities in natural populations as well as

patterns of co-occurrence among microbes. Overall, our study highlights the role of small-scale,

tick-level characteristics rather than large-scale ecological variation in shaping microbial

communities of *I. ricinus*. We identified a number of ecological variables that predict the

occurrence of specific tick endosymbionts and human or wildlife pathogens with strong statistical

support, but these effects were typically microbe-specific rather than universal. This highlights that

environmental change can have different, even opposite effects on different human pathogens, and

thus disease risk. Furthermore, by accounting for shared environmental preferences, our approach

identified patterns of microbial co-occurrence that are consistent with microbe-microbe interactions,

which result in pathogen co-infections within ticks, as well as competition between *Spiroplasma*

and a number of human or wildlife pathogens. The latter opens up new and exciting avenues for the

control and management of tick-borne diseases in regions with high human disease incidence.

Acknowledgements

This study was funded by Finnish Cultural Foundation Postdoc Pool grant (to TA), the Stiftung für wissenschaftliche Forschung an der Universität Zürich (17_027), the Swiss National Science Foundation (PP00P3_128386 and PP00P3_157455), the University of Zurich Research Priority Program "Evolution in Action: from Genomes to Ecosystems", the Faculty of Science of the University of Zurich, and the Baugarten Stiftung (all to BT). We thank Mélissa Lemoine for providing ticks, the numerous people who helped collecting ticks in the field, Glauco Camenisch, Elisa Granato, Jennifer Morger and Alessia Pennachia for help with laboratory work, Lucy Poveda for help with MiSeq sequencing, Frédéric Guillaume for providing access to IT infrastructure, Janine Bolliger and Dirk Schmatz from Swiss Federal Institute for Forest, Snow and Landscape Research WSL for providing spatial data and for help with spatial analyses, and Otso Ovaskainen for discussions on the modelling.

References

Abraham, N. M. et al. 2017. Pathogen-mediated manipulation of arthropod microbiota to promote infection. - Proc. Natl. Acad. Sci. 114: E781–E790.

Abrego, N. et al. 2016. Measuring and predicting the influence of traits on the assembly processes of wood-inhabiting fungi. - J. Ecol. in press.

Adair, K. L. and Douglas, A. E. 2017. Making a microbiome: the many determinants of host-associated microbial community composition. - Curr. Opin. Microbiol. 35: 23–29.

Aivelo, T. and Norberg, A. 2017. Parasite-microbiota interactions potentially affect intestinal communities in wild mammals. - J. Anim. Ecol. 87: 438–447.

Andersson, M. et al. 2013. Multiple-strain infections of Borrelia afzelii: a role for within-host interactions in the maintenance of antigenic diversity? - Am. Nat. 181: 545–54.

Andersson, M. et al. 2014. Infection dynamics of the tick-borne pathogen "Candidatus Neoehrlichia mikurensis" and

coinfections with Borrelia afzelii in bank voles in Southern Sweden. - Appl. Environ. Microbiol. 80: 1645–9.

Archie, E. a and Ezenwa, V. O. 2011. Population genetic structure and history of a generalist parasite infecting multiple sympatric host species. - Int. J. Parasitol. 41: 89–98.

Atkinson, S. F. et al. 2013. Modelling spatial concordance between Rocky Mountain spotted fever disease incidence and habitat probability of its vector Dermacentor variabilis (American dog tick). - Geospat. Health 7: 91–100.

Belli, A. et al. 2017. Ticks infected via co-feeding transmission can transmit Lyme borreliosis to vertebrate hosts. - Sci. Rep. 7: 1–13.

Bennie, J. et al. 2008. Slope, aspect and climate□: Spatially explicit and implicit models of topographic microclimate in chalk grassland. - Ecol. Modell. 6: 47–59.

Bonnet, S. I. et al. 2017. The tick microbiome: Why non-pathogenic microorganisms matter in tick biology and pathogen transmission. - Front. Cell. Infect. Microbiol. 7: 236.

Burri, C. et al. 2014. Are Apodemus spp. mice and Myodes glareolus reservoirs for Borrelia miyamotoi, Candidatus Neoehrlichia mikurensis, Rickettsia helvetica, R. monacensis and Anaplasma phagocytophilum? - Ticks Tick. Borne. Dis. 5: 245–251.

Carey, H. V et al. 2013. Seasonal restructuring of the ground squirrel gut microbiota over the annual hibernation cycle. - Am. J. Physiol. Regul. Integr. Comp. Physiol. 304: R33-42.

Carpi, G. et al. 2011. Metagenomic profile of the bacterial communities associated with Ixodes ricinus ticks. - PLoS One 6: e25604.

Comstedt, P. et al. 2011. Global ecology and epidemiology of Borrelia garinii spirochetes. - Infect. Ecol. Epidemiol. 1: 9545.

Cornetti, L. et al. 2016. Higher genetic diversity on mountain tops: The role of historical and contemporary processes in shaping genetic variation in the bank vole. - Biol. J. Linn. Soc. 118: 233–244.

Cornetti, L. et al. 2018. Small-scale spatial variation in infection risk shapes the evolution of a *Borrelia* resistance gene in wild rodents. - Mol. Ecol.: 3515–3524.

Cortinas, M. R. et al. 2002. Detection, characterization, and prediction of tick-borne disease foci. - Int. J. Med. Microbiol. 291: 11–20.

Dallas, T. and Presley, S. J. 2014. Relative importance of host environment, transmission potential and host phylogeny

to the structure of parasite metacommunities. - Oikos 123: 866–874.

Diuk-Wasser, M. A. et al. 2016. Coinfection by Ixodes Tick-Borne Pathogens: Ecological, Epidemiological, and

Clinical Consequences. - Trends Parasitol. 32: 30–42.

Durand, J. et al. 2015. Cross-immunity and community structure of a multiple-strain pathogen in the tick vector. - Appl.

Environ. Microbiol. in press.

Elliot, S. L. et al. 2002. Host-pathogen interactions in a varying environment: temperature, behavioural fever and

fitness. - Proc. R. Soc. B Biol. Sci. 269: 1599–607.

Eremeeva, M. E. and Dasch, G. A. 2015. Challenges Posed by Tick-Borne Rickettsiae: Eco-Epidemiology and Public

Health Implications. - Front. Public Heal. 3: 1–17.

Eren, A. M. et al. 2013. Oligotyping: Differentiating between closely related microbial taxa using 16S rRNA gene data.

- Methods Ecol. Evol. 4: 1111–1119.

Eren, A. M. et al. 2014. Oligotyping analysis of the human oral microbiome. - Proc. Natl. Acad. Sci. 111: E2875–

E2884.

Estrada-Peña, A. et al. 2018. High Throughput Sequencing and Network Analysis Disentangle the Microbial

Communities of Ticks and Hosts Within and Between Ecosystems. - Front. Cell. Infect. Microbiol. 8: 236.

Gall, C. A. et al. 2016. The bacterial microbiome of Dermacentor andersoni ticks influences pathogen susceptibility. -

ISME J. 10: 1846–1855.

Gatewood, A. G. et al. 2009. Climate and tick seasonality are predictors of Borrelia burgdorferi genotype distribution. -

Appl. Environ. Microbiol. 75: 2476–83.

Gern, L. et al. 2008. Influence of some climatic factors on Ixodes ricinus ticks studied along altitudinal gradients in two

geographic regions in Switzerland. - Int. J. Med. Microbiol. 298: 55–59.

Greay, T. L. et al. 2018. Recent insights into the tick microbiome gained through next-generation sequencing. -

Parasites and Vectors 11: 12.

Gurfield, N. et al. 2017. Endosymbiont interference and microbial diversity of the Pacific coast tick, Dermacentor

occidentalis, in San Diego County, California. - PeerJ 5: e3202.

Haine, E. R. 2008. Symbiont-mediated protection. - Proc. R. Soc. B Biol. Sci. 275: 353–361.

Halos, L. et al. 2010. Ecological factors characterizing the prevalence of bacterial tick-borne pathogens in ixodes ricinus

ticks in pastures and woodlands. - Appl. Environ. Microbiol. 76: 4413–4420.

Hammer, T. J. et al. 2017. Caterpillars lack a resident gut microbiome. - Proc. Natl. Acad. Sci. 114: 9641–9646.

Hanincova, K. et al. 2003a. Songbirds in Slovakia. - Appl. Environ. Microbiol. 69: 2825–2830.

Hanincova, K. et al. 2003b. Association of Borrelia afzelii with rodents in Europe. - Parasitology 126: 11–20.

Hawlena, H. et al. 2013. The arthropod, but not the vertebrate host or its environment, dictates bacterial community

composition of fleas and ticks. - ISME J. 7: 221–3.

Hawley, D. M. and Altizer, S. M. 2011. Disease ecology meets ecological immunology: understanding the links

between organismal immunity and infection dynamics in natural populations. - Funct. Ecol. 25: 48–60.

Herren, J. K. and Lemaitre, B. 2011. Spiroplasma and host immunity: Activation of humoral immune responses

increases endosymbiont load and susceptibility to certain Gram-negative bacterial pathogens in Drosophila

melanogaster. - Cell. Microbiol. 13: 1385–1396.

Hillyard, P. D. 1996. Ticks of north-west Europe. - Field Studies Council.

Jaenike, J. et al. 2013. Adaptation via symbiosis: recent spread of a Drosophila defensive symbiont. - Science (80-.).

329: 212–215.

Jahfari, S. et al. 2012. Prevalence of Neoehrlichia mikurensis in ticks and rodents from North-west Europe. - Parasit.

Vectors 5: 74.

Jahfari, S. et al. 2014. High seroprevalence of Borrelia miyamotoi antibodies in forestry workers and individuals

suspected of human granulocytic anaplasmosis in the Netherlands. - New Microbes New Infect. 2: 144–149.

Jouda, F. et al. 2004a. Ixodes ricinus density, and distribution and prevalence of Borrelia burgdorferi sensu lato

infection along an altitudinal gradient. - J. Med. Entomol. 41: 162–169.

Jouda, F. et al. 2004b. Density of questing Ixodes ricinus nymphs and adults infected by Borrelia burgdorferi sensu lato

in Switzerland: spatiotemporal pattern at a regional scale. - Vector-borne Zoonotic Dis. 4: 23–32.

Kamvar, Z. N. et al. 2014. *Poppr*: an R package for genetic analysis of populations with clonal, partially clonal,

and/or sexual reproduction. - PeerJ 2: e281.

Kawahara, M. et al. 2004. Ultrastructure and phylogenetic analysis of "Candidatus Neoehrlichia mikurensis" in the

family Anaplasmataceae, isolated from wild rats and found in Ixodes ovatus ticks. - Int. J. Syst. Evol. Microbiol.

54: 1837–1843.

Keesing, F. et al. 2006. Effects of species diversity on disease risk. - Ecol. Lett. 9: 485–98.

Keesing, F. et al. 2012. Reservoir competence of vertebrate hosts for Anaplasma phagocytophilum. - Emerg. Infect.

Dis. 18: 2013–2016.

Kozich, J. J. et al. 2013. Development of a dual-index sequencing strategy and curation pipeline for analyzing amplicon

sequence data on the MiSeq Illumina sequencing platform. - Appl. Environ. Microbiol. 79: 5112–20.

Laaksonen, M. et al. 2018. Tick-borne pathogens in Finland: comparison of Ixodes ricinus and I. persulcatus in

sympatric and parapatric areas. - Parasit. Vectors 11: 556.

Lawson, C. R. et al. 2014. Topographic microclimates drive microhabitat associations at the range margin of a butterfly.

- Ecography (Cop.). 37: 732–740.

Lello, J. et al. 2004. Competition and mutualism among the gut helminths of a mammalian host. - Nature 428: 20–24.

Lemoine, M. et al. 2018. Does Borrelia burgdorferi sensu lato facilitate the colonisation of marginal habitats by Ixodes

ricinus? A correlative study in the Swiss Alps. - bioRxiv in press.

Lo, N. et al. 2006. Widespread distribution and high prevalence of an alpha-proteobacterial symbiont in the tick Ixodes

ricinus. - Environ. Microbiol. 8: 1280–1287.

Lommano, E. et al. 2012. Infections and coinfections of questing Ixodes ricinus ticks by emerging zoonotic pathogens

in Western Switzerland. - Appl. Environ. Microbiol. 78: 4606–4612.

Macaluso, K. R. et al. 2002. Rickettsial Infection in *Dermacentor variabilis* (Acari: Ixodidae) Inhibits Transovarial

Transmission of a Second *Rickettsia*. - J. Med. Entomol. 39: 809–813.

MacDonald, A. J. et al. 2017. Lyme disease risk in southern California□: abiotic and environmental drivers of Ixodes

pacificus (Acari: Ixodidae) density and infection prevalence with Borrelia burgdorferi. - Parasit. Vectors: 7.

Mannelli, A. et al. 2012. Ecology of Borrelia burgdorferi sensu lato in Europe: transmission dynamics in multi-host

systems, influence of molecular processes and effects of climate change. - FEMS Microbiol. Rev. 36: 837–61.

McCain, C. M. 2005. Elevational gradients in diversity of small mammals. - Ecology 86: 366–372.

McCoy, K. D. et al. 2005. Recurrent evolution of host-specialized races in a globally distributed parasite. - Proc. R. Soc.

B Biol. Sci. 272: 2389–2395.

McCoy, K. D. et al. 2013. Host specialization in ticks and transmission of tick-borne diseases: a review. - Front. Cell.

Infect. Microbiol. 3: 57.

Medlock, J. M. et al. 2013. Driving forces for changes in geographical distribution of *Ixodes ricinus* ticks in Europe. -

Parasit. Vectors 6: 1.

Michelet, L. et al. 2014. High-throughput screening of tick-borne pathogens in Europe. - Front. Cell. Infect. Microbiol.

4: 103.

Moutailler, S. et al. 2016. Co-infection of ticks: the rule rather than the exception. - PLoS Negl. Trop. Dis. 10:

e0004539.

Narasimhan, S. et al. 2014. Gut microbiota of the tick vector Ixodes scapularis modulate colonization of the Lyme

disease spirochete. - Cell Host Microbe 15: 58–71.

Ogden, N. H. et al. 2007. Vector seasonality, host infection dynamics and fitness of pathogens transmitted by the tick

Ixodes scapularis. - Parasitology 134: 209–27.

Ovaskainen, O. et al. 2015. Using latent variable models to identify large networks of species-to-species associations at

different spatial scales. - Methods Ecol. Evol. 7: 549–555.

Ovaskainen, O. et al. 2017a. How are species interactions structured in species-rich communities？ A new method for

analysing time-series data. - Proc. R. Soc. B Biol. Sci. 284: 20170768.

Ovaskainen, O. et al. 2017b. How to make more out of community data? A conceptual framework and its

implementation as models and software. - Ecol. Lett. 20: 561–576.

Petney, T. N. and Andrews, R. H. 1998. Multiparasite communities in animals and humans: frequency , structure and

pathogenic significance. - Int. J. Parasitol. 28: 377–393.

Randolph, S. E. et al. 1999. Incidence from coincidence: Patterns of tick infestations on rodents facilitate transmission

of tick-borne encephalitis virus. - Parasitology 118: 177–186.

Rigaud, T. et al. 2010. Parasite and host assemblages: embracing the reality will improve our knowledge of parasite

transmission and virulence. - Proc. R. Soc. B Biol. Sci. 277: 3693–3702.

Rodríguez, M. et al. 1999. Susceptibility to Trypanosoma cruzi is modified by a previous non-related infection. -

Parasite Immunol. 21: 177–185.

Ross, B. D. et al. 2017. Ixodes scapularis does not harbor a stable midgut microbiome. - bioRxiv: 198267.

Shapira, M. 2016. Gut microbiotas and host evolution: Scaling up symbiosis. - Trends Ecol. Evol. 31: 539–549.

Sofonea, M. T. et al. 2015. From within-host interactions to epidemiological competition: A general model for multiple

infections. - Philos. Trans. R. Soc. B Biol. Sci. 370: 20140303.

Sprong, H. et al. 2009. Ixodes ricinus ticks are reservoir hosts for Rickettsia helvetica and potentially carry flea-borne

Rickettsia species. - Parasites and Vectors 2: 41.

Stackebrandt, E. and Goebel, B. M. 1994. Taxonomic note: A place for DNA-DNA reassociation and 16S rRNA

sequence analysis in the present species definition in bacteriology. - Int. J. Syst. Evol. Microbiol. 44: 846–849.

States, S. L. et al. 2017. Co-feeding transmission facilitates strain coexistence in Borrelia burgdorferi, the Lyme disease

agent. - Epidemics 19: 33–42.

Stuen, S. et al. 2013. Anaplasma phagocytophilum—a widespread multi-host pathogen with highly adaptive strategies. -

Front. Cell. Infect. Microbiol. 3: 31.

Swei, A. and Kwan, J. Y. 2017. Tick microbiome and pathogen acquisition altered by host blood meal. - ISME J. 11:

813–816.

Swei, A. et al. 2011. Influence of abiotic and environmental factors on the density and infection prevalence of Ixodes

pacificus (Acari: Ixodidae) with Borrelia burgdorferi. - J. Med. Entomol. 48: 20–28.

Team, R. C. 2013. R: A language and environment for statistical computing. in press.

Thornton, P. E. et al. 1997. Generating surfaces of daily meteorological variables over large regions of complex terrain.

- J. Hydrol. 190: 214–251.

Tikhonov, G. et al. 2017. Using joint species distribution models for evaluating how species-to-species associations

depend on the environmental context. - Methods Ecol. Evol. 8: 443–452.

van Treuren, W. et al. 2015. Variation in the microbiota of Ixodes ticks with regard to geography, species, and sex. -

Appl. Environ. Microbiol. 81: 6200–6209.

Vayssier-Taussat, M. et al. 2013. Next generation sequencing uncovers unexpected bacterial pathogens in ticks in

western Europe. - PLoS One 8: e81439.

Venclikova, K. et al. 2014. Rickettsiae in questing Ixodes ricinus ticks in the Czech Republic. - Ticks Tick. Borne. Dis.

5: 135–138.

Wagemakers, A. et al. 2015. Borrelia miyamotoi: A widespread tick-borne relapsing fever spirochete. - Trends

Parasitol. 31: 260–269.

Warton, D. I. et al. 2015. So many variables: joint modeling in community ecology. - Trends Ecol. Evol. 30: 766–779.

West, S. A. and Buckling, A. 2003. Cooperation, virulence and siderophore production in bacterial parasites. - Proc. R.

Soc. B Biol. Sci. 270: 37–44.

Xie, J. et al. 2010. Spiroplasma bacteria enhance survival of Drosophila hydei attacked by the parasitic wasp

Leptopilina heterotoma. - PLoS One 5: e12149.

Table 1: Tick sampling sites in the Swiss Alps.

Location	Site	Coordinates		Elevation	Succesfully sequenced *Ixodes ricinus* ticks		
		North	East	(masl)	nymphs	males	females
1	Sagogn	46.783	9.233	693	0	9	15
	Flims	46.827	9.280	1138	3	5	3
	Ruschein	46.795	9.169	1454	0	1	1
2	Rodels	46.760	9.425	630	2	5	4
	Tomils	46.772	9.453	1144	3	6	4
	Feldis	46.789	9.453	1673	1	1	0
3	Passug	46.840	9.538	732	0	5	6
	Castiel	46.826	9.569	1094	0	3	3
	Praden	46.817	9.589	1582	1	0	1

621   Table 2: Common tick endosymbionts and/or putative human pathogens observed in *I. ricinus* ticks. See

622   Supplementary Materials for information on OTU assignment.

OTU	Label	Human pathogen / tick endosymbiont	Occurrence (% of analyzed ticks)
Otu0001	*Midichloria*	endosymbiont	100
Otu0003	*Spiroplasma*	endosymbiont	41
Otu0005	*Rickettsiella*	endosymbiont	63
Otu0021	*Lariskella*	endosymbiont	49
Otu0031	*Rickettsia helvetica*	both	16
	R. monacensis	both	6
Otu0067	*Rickettsia sp.*	both	25
Otu0076	*Anaplasma*	both	33
Otu0086	*Candidatus* Neoehrlichia	both	22
Otu0088	*Borrelia afzelii*	pathogen	9
	B. miyamotoi	pathogen	10
	B. garinii	pathogen	6
	B. valaisiana	pathogen	2

625   Table 3. Associations between tick-specific and environmental variables and the occurrence of

626   endosymbionts and human pathogens in *I. ricinus* ticks. A positive sign indicates that higher variable values

627   are associated with a higher probability of OTU occurrence. A higher aspect value means that a site is facing

628   northwards. Only associations with strong statistical support (based on the 90% central credible interval)

629   are presented.

		Full variable set					Variable selection set									
		Tick sex (Female)	Tick life stage (Nymph)	Tick abundance	Tick heterozygosity	Elevation	Tick population expected heterozygosity	Number of days > 7C°	Precipitation	Mean temperature	Forest cover	Slope	Aspect	Vole abundance	Vole/ other rodents ratio
Otu0003	*Spiroplasma*	−											+		−
Otu0005	*Rickettsiella*	−		+	−		−		+				−		
Otu0022	*Lariskella*	−			−										
Otu0031	*Rickettsia helvetica*					+							−		
	R. monacensis					+	−						−		
Otu0067	*Rickettsia sp.*	−										+			
Otu0076	*Anaplasma*														
Otu0086	*Ca.* Neoehrlichia			+											
Otu0088	*Borrelia afzelii*												+		
	B. miyamotoi														
	B. garinii					−									
	B. valaisiana														

     Figure legends

Figure 1: Location of tick sampling sites in the Swiss Alps. Different colours of circles (i.e., black,

grey, white) represent the different locations. Rivers and motorway are shown in black. Modified

from (Cornetti et al. 2016).

Figure 2: Tick microbial community variance partitioning for different fixed and random effects.

The first three columns represent tick endosymbionts, the next three columns are OTUs which are

both tick endosymbionts and human pathogens and the subsequent six columns represent human

pathogens. The other columns represent the 88 most common OTUs found in *I. ricinus,* ordered by

amplicon frequency. Month, sampling site, location and tick ID were included in the model as

random effects, whereas fixed effects were divided into environmental (elevation, temperature,

precipitation, forest coverage, slope, aspect, vole abundance and vole-to-other-rodents ratio) and

tick-specific variables (life stage or sex, individual heterozygosity, abundance, expected population

heterozygosity). See raw data in Figshare for information on OTU labels.

Figure 3: Residual association patterns among endosymbionts and human pathogens within ticks

on a) individual tick-level and b) on site-level after accounting for shared environmental preference.

Red lines represent positive associations and blue lines negative associations. Only associations

with strong statistical support (i.e., based on the 90% central credible interval) are presented. Darker

colors indicate stronger associations.