1   Nanoparticle-mediated transformation expands horizon of organism engineering
2
3

4   Katherine E. French*[1]
5
6   [1] Department of Plant and Microbiology, University of California Berkeley, 111 Koshland Hall, Berkeley,
7   CA
8
9   Correspondence: katherine.french@berkeley.edu

10
11

12   **Our ability to engineer organisms will unlock new discoveries in how cellular processes work and enable new**
13   **advances in biotechnology. However, many organisms are recalcitrant to these methods. Here, we describe a**
14   **new nanoparticle-based method of transformation which can be used to transform bacteria, plants, algae and**
15   **diatoms.**

16
17
18
Our ability to engineer organisms will unlock new discoveries in how cellular processes work and enable

new advances in biotechnology (Khalil and Collins 2010). This process is known as 'transformation' and

relies on electric currents, heat shock, or chemicals to create tiny pores in organism cell membranes that

allow foreign genetic material to enter the host. However, many organisms are recalcitrant to these

methods (Brenner et al. 2008). Recent advances in nanoparticle technology have illustrated that quantum

dots can deliver siRNA (typically of 20-22 bp) and other molecules to human and mice cells, but these

methods do not work on bacteria or organisms with cell walls (Kim et al. 2017; Pierrat et al. 2015). Here,

we describe a new carbon-nanoparticle based transformation system. Nanoparticle composition,

polymer coating, pH, and plasmid DNA (p-DNA) concentration were key factors in transformation success.

Pilot experiments with eukaryotic, prokaryotic, and difficult-to-transform species suggest any organism

that can take-up the nanoparticles we designed can be transformed using this method.

We prepared 15 different nanoparticles to assess the following properties (1) organism uptake; (2) cellular

localization; (3) transformation ability; and (4) cellular toxicity. We replicated two nanoparticle types from

previous studies; the rest of the nanoparticles are unique to this study. For nanoparticles with desired

optical and functional properties, we experimented with different polymer coatings (bPEI and PEG),

polymer concentrations (0.0125-0.05 g/mL), plasmid concentrations (1 ng-1 ug/ul), and nanoparticle-

plasmid incubation time (10 min. to 24 h). The particles best suited for general (non-organelle)

transformation were made of citric acid or glucose, passivated with low concentrations of bPEI (0.01-0.02

g/mL) and plasmid (1-4ng/ul), and with a short nanoparticle-plasmid conjugation time (15-20 min.).

Particles made with threonine localized to chloroplasts and those made with arginine, lysine, valine and

serine localized to mitochondria (**SI Table 1**). Nanoparticles made with specific amino acid combinations

may mimic the role of terminal sequences in guiding proteins targeted to specific cellular compartments

(Carrie et al. 2009; de Castro et al. 1996; Lodish et al. 2000).

Cellular uptake studies were carried out in the following organisms: plants/mosses (*Arabidopsis thaliana,*

Nicotiana benthamiana, Physcomitrella patens and *Selaginella kraussiana*), protists (*Physarum*

polycephalum), bacteria (*Pseudomonas flourescens*), cyanobacteria (*Synecoccochus*), diatoms

(*Thalassiosira oceanica*), algae (*Penium margaritaceum*, *Spyrogira* sp.), fungi (*Trichoderma reesei*), and

coccolithophores (*Pleurochrysis certerae, Gephyrocapsa oceanica*). This is the first study to show algae,

diatoms and coccolithophores will take up nanoparticles.   Nanoparticles entered cells in two ways: (1)

through membrane pores or (2) invagination of cell membranes. For example, in diatoms nanoparticles

appear to enter the cell through external pores in the frustule; in algae such as *Penium*, in-folding of the

cell membrane around nanoparticle aggregates was observed and these vesicles seemed to transport

nanoparticles in and out of the cell (**Fig. 1 A and B**). The latter correlates with Giraldo et al. 2014, who

proposed that single-walled carbon nanotubes may enter the double-membranes of chloroplasts via

diffusion, becoming coated with glycerolipids in the process. Experiments with vascular plants,

filamentous fungi, and *P. polycephalum* indicate nanoparticles can be transported long distances (either

within the vascular system or within vesicles) (**SI Fig. 1**). Based on material composition, nanoparticles

localized to six places: the cell membrane, cytoplasm, vacuole, nucleus, mitochondria, and chloroplasts.

In general, carbon-based nanoparticles are non-toxic to cells. This is in contrast to nanoparticles made out

of heavy metals (e.g. 'quantum dots') which can kill cells and may cause environmental harm (Hardman

2006; Valizadeh et al. 2012). Study organisms did show signs of stress when exposed to nanoparticles

with high bPEI (0.05-0.1 g/mL) and plasmid concentrations (300 ng-1 ug/ul) or with low pH (< pH 4). Signs

of stress included retraction of the cytoplasm from the cell wall in vascular plants; cell rupture; and loss

of innate chloroplast structure (specifically in algae). Once nanoparticles were optimized to decrease cell

toxicity, organisms could survive on media (agar gel plates or liquid growth media) containing

nanoparticles for over 30 days.

The following organisms were successfully transformed using glucose and citric acid based nanoparticles

coated with bPEI and conjugated with a Venus NLS plasmid, a MCherry NLS plasmid, or a GFP HDEL

plasmid: *A. thaliana*, *N. benthamiana*, *S. kraussiana, P. margaritaceum* and *T. oceanica* (**Fig. 1 A, C-F**; **SI**

Fig. 2). Organisms were exposed to the following treatments: control (no treatment), bPEI, bPEI + pDNA,

CND, CND + pDNA. No transformation was seen using plasmids bound to bPEI, a transfection method

commonly used in mammalian cells (Kim et al. 2017; Pierrat et al. 2015), and exposure to bPEI alone

resulted in cell death for plants and algae. Successful transformation using plasmids bound to CNDs was

indicated by fluorescence of the nuclear envelope and the endoplasmic reticulum. To further confirm

transformation, we transformed *P. margaritaceum* with a 35S ubi-GUS plasmid. Non-transformed cells

appear clear or yellowish; transformed cells appear blue (**Fig. 1 E**). This system can be used for transient

and stable expression of foreign genes in both chassis organisms and difficult to transform species. For

example, the alga *P. margaritaceum* showed nuclear YFP and MCherry labelling eight days after

transfection, suggesting the plasmid DNA was inserted into the hosts' genomes. Other images show cells

dividing, with both cells exhibiting nuclear fluorescence, suggesting the DNA is inherited (**Fig. 1 F**).

Carbon nanoparticle-based transformation presents a promising new method of altering the genetic

make-up of cells. This is the first study to show the transport and expression of large plasmids (>10 kbp)

into recalcitrant organisms using carbon nanoparticles. Future studies should look at the metabolic impact

of CND exposure, the degradation of CNDs in cells and the environment, plasmid size limitations which

may affect transformation, and whether organelle-targeting nanoparticles can be used to genetically

modify chloroplasts and mitochondria. Further development of this technology will lead to new

discoveries across the life sciences and commercial gains in the growing biotech sector.

91
92

93
Fig. 1. **Organism uptake of carbon nanoparticles and transformation**. A. Expression on VNLS in *T. oceanica*. The
image shows dorsal and lateral views of several diatoms in different stages of life. B. Uptake of CNDs in *P.*
margaritaceum. Arrows point to vesicular structures filled with nanoparticles. C. Transformation of *N. benthamiana*
with VNLS. D. Expression of VNLS in *S. kraussiana* 16 hours after exposure to CND-pDNA complexes; boxes show

magnified cells with fluorescent nuclei. E. GUS-staining of *P. margaritaceum*. F. Dividing *P. margaritaceum* eight days
after exposure to CND-pDNA complexes. N= nucleus. Confocal images (A, B, C, F) taken using excitation
405/488/543nm. Images are composed of superimposed z-sections. Images D and E were taken with a Leica light
microscope using a YFP filter and an LED light respectively.
102
103 Author contributions
104
105 KEF designed the nanoparticles, performed the nanoparticle-uptake and transformation experiments,
106 imaged the experiments, and wrote the manuscript.
107
108 Declaration of interests
109
110 The authors declare no competing interests.
111
112 **References**
113

114 Brenner, K., L. You and F. Arnold. 2008. Engineering microbial consortia: a new frontier in synthetic
115 biology. *Trends in Biotechnology* 26.9: 483-489.
116
117 Carrie, C., E. Giraud and J. Whelan. 2009. Protein transport in organelles: Dual targeting of proteins to
118 mitochondria and chloroplasts. *FEBS Journal* 276: 1187–1195.
119
120 de Castro Silva Filho, M., F. Chaumont, S. Leterme, M. Boutry. 1996. Mitochondrial and chloroplast
121 targeting sequences in tandem modify protein import specificity in plant organelles. *Plant Molecular*
122 *Biology* 30: 769-780.
123
124 Giraldo, J.P., M. P. Landry, S. M. Faltermeier, T. P. McNicholas, N. M. Iverson, A. A. Boghossian, N. F.
125 Reuel, A. J. Hilmer, F. Sen, J. A. Brew and M. S. Strano. 2014. Plant nanobionics approach to augment
126 photosynthesis and biochemical sensing. Nature Materials 13: 400-410. DOI: 10.1038/NMAT3890.
127
128 Hardman, R. 2006. A Toxicologic Review of Quantum Dots: Toxicity Depends on Physicochemical and
129 Environmental Factors. *Environmental Health Perspectives* 114.2: 165–172.
130
131 Kim, S., Y. Choi, G. Park, C. Won, Y. Park, Y. Lee. 2017. Highly efficient gene silencing and bioimaging
132 based on fluorescent carbon dots *in vitro* and *in vivo. NanoResearch* 10.2: 503–519.
133
134 Khalil, A. and J. Collins. 2010. Synthetic biology: applications come of age. *Nature Reviews Genetics* 11:
135 367-379.
136
137 *Molecular Cell Biology*. 4th edition. Lodish ,H., A. Berk, S.L. Zipursky, et al. New York: W. H. Freeman.
138 2000.
139
140 Pierrat, P., R. Wang, D. Kereselidze, M. Lux, P. Didier, A. Kichler, F. Pons, L. Lebeau. 2015. Efficient in
141 vitro and in vivo pulmonary delivery of nucleic acid by carbon dot-based nanocarriers. *Biomaterials* 51:
142 290e302.
143
144 Valizadeh, A., H. Mikaeili, M. Samiei, S. M. Farkhani, N. Zarghami, M. Kouhi, A. Akbarzadeh, S. Davaran.
145 2012. Quantum dots: synthesis, bioapplications, and toxicity. *Nanoscale Research Letters* 7.1: 480.